Title: Identification of erythroid cell positive blood transcriptome phenotypes associated with severe respiratory syncytial virus infection

Authors: Darawan Rinchai[1], Matthew B Altman[2,3], Oceane Konza[4], Signe Hässler[4,5], Federica Martina[4], Mohammed Toufiq[1], Mathieu Garand[1], Basirudeen Kabeer[1], Karolina Palucka[6], Asuncion Mejias[6], Octavio Ramilo[6], Davide Bedognetti[1], Encarnita Mariotti-Ferrandiz[5], David Klatzmann[4,5] and Damien Chaussabel[1]

Affiliations:
[1] Sidra Medicine, Doha, Qatar
[2] Benaroya Research Institute, Seattle, WA, USA
[3] University of Washington, Seattle, WA, USA
[4] AP-HP, Hôpital Pitié-Salpêtrière, Biotherapy (CIC-BTi) and Inflammation-Immunopathology-Biotherapy Department (i2B), Paris, France
[5] Sorbonne Université, INSERM, Immunology-Immunopathology-Immunotherapy (i3), Paris, France
[6] Jackson Laboratory for Genomic Medicine, Farmington, CT, United States
[7] Division of Infectious Diseases, Nationwide Children's Hospital, Columbus, OH, United States

ABSTRACT

Biomarkers to assess the severity of acute respiratory syncytial virus (RSV) infection

are needed. We conducted a meta-analysis of 490 unique profiles from six public RSV

blood transcriptome datasets. A repertoire of 382 well-characterized transcriptional

modules was used to define dominant host responses to RSV infection. The

consolidated RSV cohort was stratified according to four traits: "interferon response"

(IFN), "neutrophil-driven inflammation" (Infl), "cell cycle" (CC), and "erythrocytes" (Ery).

Eight prevalent blood transcriptome phenotypes were thus identified. Among those

three Ery+ phenotypes comprised higher proportions of patients requiring intensive

care. We posit that the erythrocyte module is linked to an overabundance of

immunosuppressive erythroid cells that might underlie progression to severe RSV

infection. These findings outline potential priority areas for biomarker development and

investigations into the immune biology of RSV infection. The approach that was

employed here will also permit to delineate prevalent blood transcriptome phenotypes in

other settings.

39
INTRODUCTION

Respiratory syncytial virus (RSV) infection is the leading cause of hospitalization and

the second cause of infant mortality worldwide (1). There are well-characterized

populations at risk for severe disease, but most infants who develop a severe RSV

infection have no underlying health conditions (2,3). The mechanisms underlying RSV

morbidity are poorly understood, but studies suggest that immature or under-developed

lungs and/or a dysregulated immune response might have a role (4).

Several groups of researchers, including us, have undertaken blood

transcriptome profiling studies of patients with RSV infection (5–13). This approach

involves measuring the abundance of blood leukocyte transcripts on a genome-wide

scale (14,15). Whole blood comprises a heterogeneous mix of leukocyte populations;

thus, changes in transcript abundance might be attributable to either gene expression

regulation or relative changes in cell abundance. Regardless, blood transcriptome

profiling remains one of the most straightforward approaches to implement in clinical

settings and on a large scale (15). Among the RSV blood transcriptome studies, several

aimed to identify factors associated with severe disease. For example, we reported an

increase in abundance of neutrophil, inflammation and erythrocyte genes in severe

pediatric cases (7). Brand et al. pinpointed that an increase in abundance of transcripts

coding for Olfactomedin 4, a factor involved in inflammatory responses, is strongly

associated with disease severity (10). More recently, Do et al. linked RSV disease

severity with follicular T helper cell development and BCL6-dependant inflammation (9).

Building consensus around biomarker signatures and finding a path to clinical

utility may involve performing meta-analyses that regroup datasets derived from multiple

independent studies (16). Work that permitted the development of novel diagnostic

products for sepsis provides a good example (17,18).   An obvious benefit of

consolidating data from multiple studies is that it permits to achieve larger sample sizes.

Arguably the heterogeneity of the patient populations and clinical settings could also

help improve the robustness of the resulting biomarker signatures (16). Challenges

include the presence of important technical variability between studies, such as in the

sampling methods, the profiling platform used or data pre-processing. Another potential

limitation is the varying depth and lack of harmonization of sample or subject

information available from one study to another.

Such meta-analyses tend to focus on the identification of consensus biomarker

signatures: here, the deliverable is a set of differentially expressed genes or predictors

of a given clinical outcome. In the present work we endeavored to identify discrete

molecular traits (e.g. "interferon = IFN", "inflammation = Inf", "Erythrocytes" = Ery")

underlying inter-individual differences among patients with an RSV infection. We used

such traits to define blood transcriptome phenotypes and to stratify patient cohorts on

the basis of their individual status of each trait: increased, decreased or unchanged vs

the uninfected comparators (e.g. of a given phenotype being IFN+ Infl0 Ery-). The next

key step was to assess the clinical relevance of such a classification, for instance in

terms of differences in the degrees of RSV disease severity. We finally endeavored to

investigate the biological basis of the inter-individual variation being measured, in

particular for the traits showing the highest degree of association with severe

presentations.

A fixed repertoire of transcriptional modules formed the basis for this work. This

repertoire consists of a collection of co-expressed gene sets. Co-expression was

determined in a collection of reference datasets encompassing 16 distinct

immunological states (19) (see methods section). This 382-module repertoire is "fixed",

in the sense that it serves as a reusable framework for analysis and interpretation of

transcriptome data. As such, transcriptional modules are not re-formed every time a

new dataset is analyzed. Using transcriptional modules is a key aspect of our approach.

Indeed, it is the drastic reduction in the number of variables that permits the selection of

molecular traits that underpin patient phenotyping and cohort stratification. The fact that

this repertoire is fixed is also important, as it permits considerable functional annotations

of these transcriptional modules. This annotation can in turn prove critical in unravelling

the biological significance of the patient phenotypes being defined.

In summary, we present here a meta-analysis of six RSV blood transcriptome

datasets that include 490 unique subject profiles. Specifically, we aimed to: 1) measure

inter-individual variability and molecularly stratify RSV patients; 2) identify the

associations between patient molecular phenotypes, clinical parameters and outcomes;

and 3) identify and interpret the immunobiological processes associated with each

molecular phenotype.

RESULTS

A collection of RSV blood transcriptome datasets can be assembled from earlier submissions to public repositories.

Several researchers investigating the host responses to RSV infection have made their blood transcriptome datasets public. We consolidated the datasets contributed by six independent studies (5–7,20–22) and performed meta-analyses to delineate distinct blood transcriptome phenotypes among RSV subjects. A criterion for including studies in this meta-analysis was the availability of uninfected controls. This point is important because control groups serve as a common denominator between studies and provide the basis for data normalization. Thus, public datasets for which such controls were unavailable could not be included in this meta-analysis.

As each study had different goals and designs, it was first important to identify the key differences, so that they are accounted for when interpreting the meta-analysis results. Information about the six studies is summarized in **Table 1**. The most notable outlier in this collection was the dataset from Liu et al. [GSE73072 (21)] as it consisted of samples from adult subjects collected before and after experimental exposure to RSV. All other studies comprised pediatric subjects with community-acquired RSV infection and a separate group of uninfected controls. Among the latter, the study by Mejias et al. addressed the question of disease severity most directly [GSE38900 (7)], while the work of de Steenhuijsen Piters [GSE77087 (6)] examined the effects of of microbiome composition on the disease course and blood transcriptome signatures. Rodriguez-Fernandez et al. examined the influence of RSV genotypes on blood transcriptional signatures [GSE103842 (5)]]. The study from McDonald et al. focused on

identifying pathways involved in disease pathogenesis [GSE80179 (22)]. In the study by

Herberg et al. [GSE42026 (20)] the RSV dataset was mostly used as a comparator in a

study focusing on responses to H1N1 influenza. The latter two studies were conducted

in Europe, while all others were conducted in the United States. Finally, in terms of

technical variables, samples from the adult exposure study were run using Affymetrix

GeneChips, while the others were run on Illumina BeadArrays. The sample types were

otherwise homogenous across all studies and consisted of RNA stabilized whole blood.

The studies used one of two popular commercial sample collection tubes for this type of

application: Paxgene blood RNA tubes (3 studies) or Tempus tubes (3 studies).

Altogether the consolidated dataset collection that was constituted for this meta-

analysis encompassed 490 profiles, of which 319 were from subjects with an RSV

infection. We hypothesized that this expanded sample size should permit us to define

blood transcriptome phenotypes and stratify patient cohorts more effectively than each

individual study could.

Comparing the changes in transcript abundance across module aggregates

identifies a consensus RSV signature.

Extracting meaningful biological information from large-scale datasets is a

notable challenge. Meta-analyses are particularly compounded by the high degree of

technical variation existing between independent studies. To at least partly address

such challenges, we used a fixed repertoire of transcriptional modules (19) as a

framework for data analysis and interpretation. In brief, this repertoire comprises 382

modules, each formed by a set of genes grouped together based on patterns of co-

expression across a reference collection of 16 blood transcriptome datasets. This

collection was comprised of 985 unique transcriptome profiles and spanned 16 different

immunologically relevant pathological or physiological states. A higher degree of

organization was further achieved by organizing, in turn, the modules into 38

"aggregates" (designated A1-A38). This grouping was based on similarities in patterns

of transcript abundance, determined this time at the module-level and across the 16

reference datasets. Each aggregate comprised between 1 and 42 modules (27 of the 28

aggregates comprised 2 modules or more).

Here, we mapped the changes in transcript abundance for each dataset against

this modular framework. From a practical perspective, this means determining for a

given module the percentage of its constitutive transcripts that are significantly changed.

This procedure is repeated in turn for each module and across each one of the six

datasets comprised in our collection. This approach made it possible to assess, as a

first step, the degree of consistency in the RSV response signatures across the

datasets. To facilitate interpretation, we represented the changes at the least granular

level by showing on a heatmap the abundance profiles for each of the six RSV datasets

(columns) across 27 module aggregates (rows) (Figure 1A). From this highly reduced

set of variables, we could pinpoint the most conserved molecular signatures across the

six datasets. These included seven aggregates showing consistent increases in

transcript abundance (observed in at least 5/6 datasets: A26, A27, A28, A33, A35, A37,

A38) and two aggregates showing consistent decreases (A1, A3). Changes were also

observed for another set of modules but in only 3/6 RSV datasets (A15, A16, A29, A30,

A34, A36). Technical or biological parameters (Table 1) did not yield an obvious

explanation for the differences between these two groups of studies. The amplitude of

the changes in other aggregates was minimal. Taken together, this step permitted the

mapping of transcriptional changes measured across different RSV data using the same

transcriptional module framework. This was useful in relating changes observed

between the studies and pinpointing signatures that appear to be most robustly

associated with RSV infection.

Changes in transcript abundance can be mapped to a fixed transcriptional

module repertoire to facilitate functional interpretation.

To functionally interpret the conserved signatures observed across RSV datasets, a

more granular level of information is needed. We thus examined transcriptomic changes

at the level of the modules forming the 27 aggregates mentioned above. We

represented the changes in transcript abundance as grid plots for each of the RSV

datasets (**Figure 1B**, **Supplementary Figure 1**). A first vertical reading of the grid

across the rows provides a sense of the changes at the aggregate level already

summarized in the heatmap that was presented earlier (**Figure 1A**). A second

horizontal reading across the columns provides a sense of the changes occurring at the

module level within each of the aggregates.

Because the positions on the grid are fixed, it is possible to overlay other

information, such as functional annotations (color-coded grid in **Figure 1B**). We found

that some of the conserved signatures that were increased during RSV infection

comprised modules preferentially associated with interferon responses (A28),

inflammation (A33, A35), erythrocytes (A37, A38) and cell cycle (A27), while those that

were decreased were associated with lymphocytic responses (A1, A3). Some of these responses are further interpreted below, and all details are accessible via interactive web presentations for modules constituting each aggregate (web links are listed in **Table 2**). The presentations include reports from functional profiling analyses carried out using different tools. Heatmaps representing the patterns of abundance for transcripts constituting each module across reference datasets are also available. Furthermore, a dedicated web application was develop in support of the work presented here and permits users to access the fingerprint grid plots presented here and generate other types of plots which are presented throughout this manuscript. This resource can be accessed via this link: https://drinchai.shinyapps.io/RSV_Meta_Module_analysis/. A video demonstration can be accessed here: https://youtu.be/htNSMreM8es.

In summary, we mapped the conserved RSV signatures identified to a well-annotated modular framework. This mapping made it relatively straightforward to assign each signature to a predetermined functional category. The added granularity and available online resources make it possible to further dissect those signatures in subsequent analyses, as exemplified below.

Blood transcriptional signature present a high level of inter-individual among cohorts of RSV patients

Blood transcriptome profiling provides a means to measure inter-individual differences with a high degree of resolution. Understanding the biological and clinical significance of this variability is important but requires the study of large patient cohorts. The collective re-analysis of the six datasets assembled here provides a unique opportunity to

investigate inter-individual differences among patients with RSV infection at the

molecular level.

The approach that we used next to map changes in transcript abundance for

individual patients against the repertoire of transcriptional modules is very similar to that

described above for groups of patients. We expressed the changes for each individual

RSV patient as a percentage of constitutive genes for which the abundance was

increased or decreased compared to the respective control group (see methods for

details). As an illustration, we generated a heatmap (**Figure 2**) of the results obtained

for subjects comprising the de Steenhuijsen Piters dataset (**Figure 1B**). The patterns in

the changes in abundance are only shown for the modules constituting the nine

aggregates deemed to be conserved across the collection of RSV datasets (highlighted

in **Figure 1**). We generated similar plots for each of the remaining datasets

(**Supplementary File 1**). There are also accessible and customizable via our web

application (https://drinchai.shinyapps.io/RSV_Meta_Module_analysis/).

From the heatmaps, we observed that inter-individual variability exists even for

signatures that at the group level were well-conserved across the six datasets (**Figure**

1A). In reality, only a minority of patients in this illustrative dataset (11/81) matched the

prototypical pattern defined above based on conserved changes observed for nine

module aggregates (i.e. A1- A3- A27+ A28+ A33+ A35+ A37+ A38+; **Figure 2**). The

degree of inter-individual variability differed from one module aggregate to another. For

example, in the modules forming aggregates A1 (Lymphocytic) or A33 (Inflammation),

changes in abundance only varied in amplitude without, for the most part, showing an

inversion of trends. In other modules, inverted trends were much more common, as

exemplified by A37 (Erythrocytes).

Taken together, examining changes in transcript abundance at the level of

individuals revealed a significant degree of heterogeneity among cohorts of RSV

patients. This paradigm was also true for signatures deemed to be conserved when

carrying out comparisons at the group level.

Distinct blood transcriptome phenotypes are identified among a consolidated

cohort of RSV patients

The fact that the consensus disease signature defined earlier was not reflected at the

level of individual patients highlighted the need to characterize distinct RSV blood

transcriptome phenotypes. For this, we used the combined set of patients from the six

public transcriptome datasets. First, we generated PCA plots to evaluate the sources of

variance among this composite set of samples (**Figure 3A**). The results indicated the

absence of study bias when abundance measures where normalized to the respective

control group and reduced to module-level summarized values. This finding was largely

confirmed when representing inter-individual differences on a tSNE plot (**Figure 3B**)

(23). One dataset did show partial separation from the others, but this only concerned a

minority of subjects and could also be attributable to biological sources. The same shift

was observed on the PCA plot but only along PC3, which accounts for only 9% of the

overall variance.

As a first step, we sought to identify signatures with the highest degree of inter-

patient variability, assuming that these would be best able to discriminate patients

according to phenotypes. We identified these signatures at the least granular, module-

aggregate level, thus starting from a set of only 38 variables (**Supplementary Figure**

2). We selected the following four aggregates: 1) A27, comprising five modules

functionally associated with the "cell cycle"; 2) A28, comprising six modules functionally

associated with the "interferon response"; 3) A35, comprising 21 modules predominantly

associated with "inflammation"; and 4) A37, comprising 11 modules predominantly

associated with "erythrocytes". We discarded other aggregates that exhibited a similar

degree of inter-patient variability due to likely redundancy with the four selected

aggregates. These aggregates included A33, which like A35 is also associated with

"inflammation", and A38, which like A37 is associated with "erythrocytes".

Next, we assigned the status for each aggregate signature in a given individual

using the corresponding percentage of increased or decreased transcripts: if the value

was >15% the aggregate was considered to be increased (noted as +), if ≤15% it was

considered to be decreased (noted as -), else it was considered not changed (noted as

0). For example, Infl0/IFN+/CC+/Ery-. We then generated the distribution of subjects

constituting the combined RSV cohorts across all 81 possible Infl/IFN/CC/Ery

phenotypes (**Figure 4**). We found that a small subset of phenotypes comprised a higher

number of patients than others (>10 per phenotype). These phenotypes were all

positive for the interferon "trait" (IFN+), positive or showed no changes for the

"inflammation" and "cell cycle" traits (Infl+/0 or CC+/0) and exhibited any erythrocytes

trait status (Ery+/0/-). Phenotypes where the interferon status was unchanged or

decreased were comparatively less prevalent (five patients per group at most), and so

were phenotypes where interferon was increased but inflammation or cell cycle
phenotypes were decreased.

Overall, we have shown here that a principled approach using a small subset of
highly reduced variables can identify a discrete number of interpretable RSV blood
transcriptome phenotypes. It should be noted, however, that alternate classifications
could be obtained by modifying the selection criteria. This would be the case if, for
instance, more emphasis was put on biological significance than inter-patient variability.

296 A subset of RSV blood transcriptome phenotypes is associated with severe
297 disease

An obvious next question is whether the stratification of RSV patients according to blood
transcriptome phenotypes, such as those described above, have any clinical relevance.
The extent of phenotypic information made available alongside the blood transcriptome
datasets varied significantly from study to study. Notably, pertinent information reflecting
disease severity (e.g. respiratory rate, transcutaneous O_2 saturation) were lacking for
many patients. As a result, we had to use a relatively crude metric of disease severity
that relied on the type of care the patient required; i.e. whether they were outpatients,
inpatients cared for in the ward or were admitted to the pediatric intensive care unit
(PICU).

We first visualized the patterns of transcript abundance at the module level for
individuals belonging to the seven most prevalent phenotypes (**Figure 5**). This heatmap
verified that the phenotypic categories presented a high degree of homogeneity. Upon
overlaying the phenotypic information on this plot, we gained a first indication of a

possible association between age and disease severity. Specifically, it was possible to

discern a trend towards a younger age among Ery+ subjects in comparison to Ery- or

Ery0 subjects. Importantly, the different studies also seemed to be well represented in

each of the phenotypes, indicative of their underpinnings by biological rather than

technical factors.

We then looked at the relative proportion of severe patients for each high-prevalence

phenotype and the contributions by the different studies (**Figure 6A**). Four of the eight

phenotypes were Ery+, of which three comprised a proportion of PICU patients that was

on average 5.6 times higher than the four Ery- and Ery0 phenotypes. The Ery+ patients

included the quadruple positive IFN+/Infl+/CC+/Ery+ phenotype with 32% of PICU

patients, while its IFN+/Infl+/CC+/Ery- counterpart had 12% of patients. We found no

severe patients in the IFN+/Infl+/CC0/Ery0 group. Furthermore, subjects with Ery+

phenotypes were significantly younger than subjects with Ery- phenotypes (p <0.001,

Figure 6B).

We next endeavored to determine whether the presence of the Ery trait was

associated with heightened severity, regardless of age. For this we examined the

distribution of severe cases across the same phenotypes but focusing on infants <4

months old (**Figure 6A)**. Again, the severe cases were distributed preferentially among

Ery+ phenotypes [Ery+ = 29 PICU) cases, Ery-/0 = 4 PICU cases; of note the

IFN+/Infl+/CC+/Ery- phenotype comprised only two patients, one of which was a PICU

case].

Finally, we investigated the associations between each of the four traits used for

RSV patient phenotyping and stratification, and disease severity (**Figure 7**). Here we

found that the abundance of transcripts forming the A37/erythrocytes cells aggregate were significantly increased in patients cared in the PICU compared to the ward (Ery trait; p<0.01). We made a similar finding for the A35/"inflammation" aggregate, although to a lesser degree (Infl trait ; p<0.05). We found no significant differences were found for the A27/cell cycle or A28/interferon aggregates. Associations can be explored for various aggregates as a function of age via our web application (https://drinchai.shinyapps.io/RSV_Meta_Module_analysis/).

Taken together, these findings suggest that our RSV stratification system might be clinically relevant. This conclusion is illustrated by the fact that a high proportion of severe subjects was observed among most phenotypes positive for the Ery trait. This finding might be particularly relevant in infants <4 months–of-age who would otherwise carry a similar risk of developing severe RSV disease when taking age into consideration.

The RSV "Erythrocyte" signature is shared with melanoma patients and liver transplant recipients

Beyond the question of clinical relevance of these RSV blood transcriptome phenotypes, we next sought to understand their biological significance. For this we relied on several resources. First, we used a web application providing access to a reference collection of module fingerprints for the 16 pathological or physiological states (19) (accessible via: https://drinchai.shinyapps.io/dc_gen3_module_analysis/#; demonstration video: https://youtu.be/y__7xKJo5e4).

We generated fingerprint grid plots to compare the changes in transcript

abundance in acute influenza and RSV infections (**Figure 8A**). Acute influenza infection

highly resembles RSV infection in terms of clinical presentation, especially in infants. As

could be expected, the fingerprints of both of these respiratory infections featured a

potent interferon signature (modules in aggregate A28; i.e. the IFN trait defined earlier).

The modules associated with inflammation (comprised in aggregate A35; Infl trait) were

also generally increased in both diseases. However, one of the most marked

differences between the influenza and RSV fingerprints concerned the erythrocyte

signature (aggregate A37; Ery trait), which was consistently increased in RSV but was

unchanged in influenza.

Another fingerprint dominated by an interferon signature was that of systemic

lupus erythematosus (SLE), but as was the case for influenza, it did not comprise an

"A37/erythrocyte" signature. Among the fingerprints of other reference datasets, those

of *Staphylococcus aureus* infection, liver transplantation and metastatic melanoma also

showed an elevation in abundance for transcripts constituting modules belonging to the

A37 aggregate (**Figure 8A**). The *S. aureus* infection fingerprint showed the highest

degree of alteration overall, with widespread changes in transcript levels occurring in

most aggregates. This finding contrasts with the fingerprint for metastatic melanoma,

which for the most part was quiescent, except for a marked increase in the abundance

of genes constituting the A37 modules. The signature observed in liver transplant

recipients receiving maintenance chemotherapy was more perturbed than that of

melanoma patients, but it was likewise characterized by an increase in the abundance

of transcripts constituting the "A37/erythrocyte" modules.

Next, we used the same web application to examine module abundance profiles

specifically for the IFN, Infl, CC and Ery traits across all 16 reference datasets (**Figure**

8B). For the IFN trait (A28), RSV clustered among the diseases showing an

intermediate level of response, along with liver transplant recipients, patients with

systemic onset juvenile idiopathic arthritis (SoJIA), *S. aureus* infection (pediatric) or

sepsis caused by various pathogens (adults). Influenza was clustered among diseases

showing the highest IFN responses, including other infections such as tuberculosis or

HIV, as well as systemic lupus erythematosus (SLE) (**Supplementary Figure 3**). For

the Infl trait (A35), RSV clustered again with diseases showing an intermediate

response level, and predictably lower than those measured not only in SoJIA, sepsis,

and *S.* aureus, but also influenza infection. For the CC trait (A27), the RSV and HIV

cohorts formed a cluster with the highest increase in abundance. Finally, for the Ery trait

(A37) the RSV cohort was one of only three diseases in the high abundance cluster,

along with the melanoma and liver transplant cohorts. We observed increases to a

lesser extent in diseases characterized by overt systemic inflammation, such as sepsis

or SoJIA, as well as in pregnant women. This trait tended to be decreased in other viral

illnesses, such as influenza and HIV infection.

Overall, this contextual interpretation of the dominant traits comprising the RSV

signature identified some peculiarities. Notably, the interferon response that, while

robust, seemed to be somewhat muted when compared to other viral infections. More

strikingly was the atypical overall elevation in the abundance levels associated with the

erythrocyte signature. The extent of the observed change was only found in melanoma

patients and liver transplant recipients: in these two cohorts, the erythrocyte signature

dominated the overall changes observed in the blood transcriptome. Notably, both

patient populations relate to states characterized by marked immunosuppression, driven

by the disease in the first case and pharmacological treatment aiming at maintaining

graft tolerance in the other.

Expression of transcripts constituting the A37/"erythrocyte" modular signature is

restricted to a population of fetal GlyA+ erythroid cells

In our final analyses, we focused our interpretations on the erythrocyte signature (A37).

Although we had observed an association with RSV disease severity, we did not

ascertain causality. Based on functional profiling results that were run using multiple

approaches, we attributed the erythrocyte annotation to 8/11 modules in aggregate A37

(interactive   reports   available   via: https://prezi.com/view/YyQs4WiXSNf0YXE79lfS/).

Examining the abundance patterns for the transcripts comprising the A37 modules in

reference datasets consisting in isolated leukocyte cell populations provided further

insight (**Figure 9;** with additional heatmaps accessible interactively via the weblink

provided above). In one such reference dataset contributed by Novershtern et al. (24),

the expression of A37 transcripts was narrowly restricted to populations of glycophorin

A-positive (GlyA+) fetal erythroid cells. This pattern was irrespective of CD71 marker

expression. However, genes comprising A37 modules were not expressed in CD71+ but

GlyA- cells. We observed similar expression patterns for modules constituting

aggregates A36 and A38, which both comprised one module functionally associated

with the erythrocyte annotation (**Supplementary Figure 4**, **Supplementary Figure 5**)

Erythroid precursors of fetal origin can circulate in the blood of neonates for up to

3-4 weeks following birth. Immunosuppressive properties have been attributed to these

populations (25); for example, this cell population confers susceptibility to Listeria

infection in neonates (26). However, a possible role for these circulating erythroid cells

in the context of RSV infection has not been investigated to date. Others have also

described the presence of an erythroid cell population with potent immunosuppressive

properties associated with anemia in adults with late stage cancer (27). This finding is

consistent with our observation of a prominent A37/erythrocyte signature in the

melanoma patients included in our study.

Taken together, these observations support the notion that an increase in A37

transcript abundance is associated with the presence of circulating erythroid cells.

These cells might possess immunosuppressive properties and conceivably contribute to

the worsening of RSV infection. While these findings might be particularly relevant to

young infants, we also observed increases in adult subjects exposed experimentally to

the virus (GSE73072: **Figure 1A, Supplementary Figure 1**).

DISCUSSION

This work has built on earlier studies investigating host responses to RSV via blood

transcriptome profiling. The approach we adopted did not focus on identifying sets of

classifiers or predictors. Rather, we primarily documented the inter-individual

differences among this large consolidated set of patients.

Relying on highly reduced dimensions made it possible to define dominant blood

transcriptome phenotypes among this aggregated RSV patient cohort. The four "traits"

or signatures that were retained included: interferon (A28 aggregate / IFN trait),

inflammation (A35 / Infl), the cell cycle (A27 / CC) and erythrocytes (A37 / Ery). Out of

319 RSV subjects, 199 were distributed in just eight phenotypes out of a possible 81.

These dominant phenotypes were all positive for the interferon trait and positive or

unchanged for the inflammation and cell cycle traits. The erythrocyte trait status ranged

from being increased, unchanged and decreased.

From the standpoint of clinical significance, the phenotypes positive for both the

interferon and erythrocyte traits were generally associated with a higher proportion of

severe subjects. However, the levels of increase in abundance of interferon-inducible

transcripts did not correlate with disease severity. Rather, erythrocyte transcripts

showed the strongest association with infection severity. This finding confirmed the

association previously described by Mejias et al., which was based on the analysis of

one of the datasets comprised in this collection (28). We also observed an association,

although to a lesser degree, between the level of increase of transcripts forming an

inflammation signature (aggregate A35) and infection severity. Again, this finding is in

line with previous data (9,10). Follow-up studies in large patient cohorts are now

warranted to validate such classification scheme. Then, the number of traits needed for

this classification to be clinically relevant must be determined. Thus far, our analysis

suggests that the erythrocyte trait might constitute a valuable risk indicator if testing

focuses on a narrowly defined age group (e.g. <4 months-of-age).

Our meta-analysis of available RSV blood transcriptome datasets also yielded

several insights relevant to the immune biology of this disease. The interferon signature

(A28 / "IFN" trait) is a hallmark of RSV infection and is observed in a wide range of other

viral and bacterial infections, as well as autoimmune diseases, as illustrated in the set of

16 blood transcriptome datasets used in our interpretation. Our previous work has

suggested that subsets of modules constituting this aggregate are preferentially induced

by type I interferons [M8.1, M10.1 (19)]. Consistently, we also observed changes for

these modules in patients with RSV infection (**supplementary Figure 3**). Others have

also described robust interferon responses as measured via blood transcriptome

profiling, in RSV patients (11). The role of type I interferons *per se* is not widely

reported, but a recent study describes a dependency on interferon alpha and beta for

developing antibody-mediated responses (29). Several reports, however, have identified

suppressed interferon gamma responses in the context of RSV infection, especially

when compared to the response observed during influenza virus infection (30–33). This

finding is also consistent with our observation of a somewhat muted interferon response

compared to what we measured in response to not only influenza, but also to HIV or TB

infection, as well as in patients with SLE.

We propose that the inflammation signature (A35 / "Infl" trait) is associated with

"neutrophil-driven" inflammation, given the preferential expression of its constitutive

transcript in neutrophils and induction in patients with sepsis. This information is

derived, in part, from a dataset contributed by Linsley et al. that comprised RNAseq

profiles of isolated leukocyte fractions (34). The patterns of transcript abundance are

available for the modules constituting aggregate A35 via an interactive web presentation

(https://prezi.com/view/7Q20FyW6Hrs5NjMaTUyW/). This aggregate was the focus of a

recent re-analysis we conducted in which an increase in abundance of A35 transcripts

was a dominant feature of the psoriasis blood transcriptome fingerprint (35). We

hypothesized in turn that this inflammatory response might be driven more specifically

by interleukin-17 (IL17). Indeed, several researchers have found a role for IL17 in the

context of RSV infection (36,37) and in one instance specifically indicating the

involvement of neutrophils in IL17-mediated antiviral responses (38).

We posit that the cell cycle signature (A27 / "CC" trait) is associated with the

expansion of plasmablasts, which are responsible for antibody production. Indeed,

modules in this aggregate comprised an overabundance of genes involved in the cell

cycle, such as cyclins. One of the modules also comprises several genes expressed by

plasmablasts (M12.15: CD38, IGJ, TNFRSF17) (39–41) that are markedly expressed

between 7 and 14 days after the administration of trivalent influenza or pneumococcal

vaccines (42). In the context of this study, changes in abundance of these markers was

confirmed by flow cytometry to be correlated with the presence of these antibody

producing cells. These levels also correlated with antibody titers measured four weeks

post-vaccination. These cell populations are also expanded during the course of RSV

infection. Habibi et al. reported a peak 10 days after experimental exposure to RSV and

an correlation with the levels of neutralizing antibody developed by the individuals (43).

Consistent with our earlier findings (28), the erythrocyte signature (A37 / "Ery"

trait) was most strongly associated with severity. However, such an association has not

been reported in other studies comprising the consolidated dataset collection. We

putatively link this signature with the presence of circulating erythroid cell precursors,

based on the restriction of A37 transcripts in a reference transcriptome dataset to fetal

erythroid cells (**Figure 9**). Erythroid precursors would normally be found in the bone

marrow, but cells originating from the fetal liver do persist in infants in the circulation for

a few weeks after birth (44). In adults, extramedullary erythropoiesis is observed in the

spleen and liver, and occurs under various circumstances, including anemia,

pregnancy, severe infection or chronic stress (45,46). We hypothesize that circulating

erythroid cells (CECs) associated with this signature might have immunosuppressive

functions during an RSV infection. Thus, CEC-mediated immunosuppression would in

turn drive a worsening of disease and severity in patients with RSV infection. This

assertion is supported by various lines of evidence. First, Elahi et al. have described a

wide range of mechanisms conferring immunosuppressive properties to this cell

population, including via soluble factors (such as arginase, TGFbeta, reactive oxygen

species) or cell surface receptors (such as PD1/PDL-1 and VISTA) (25). Second, we

also observed a marked increase in A37 transcript abundance in metastatic melanoma

and liver transplant recipients under maintenance therapy. Both of these states are

characterized by marked immunosuppression and were categorized in the high

abundance profile cluster for A37. Of the 14 other cohorts in this reference collection

only RSV was present in this same cluster. Others have recently described

immunosuppression exerted by CECs in patients with late stage cancer: these cells

were found to be at least in part responsible for the impaired T-cell responses observed

in this patient population (27). Third, the RSV literature provides indications of this virus'

ability to subvert the immune response (4). While the possibility of an involvement of

CECs in this immune modulation of the response to RSV is novel, the contribution of the

hyporesponsiveness of the neonatal immune system as an underlying factor to

progression to severe RSV infection has been clearly outlined (47). A key question

could thus center on the possible contribution of CECs to the reduced competence of

the neonatal immune system. Indeed, results obtained by Elahi et al. in animal models

indicate that CEC depletion can restore neonatal immune responsiveness and confer

resistance to Listeria infection (26).

The synthesis that we conducted here builds upon and extends earlier findings. It

also identifies new avenues of investigations. Notably, it points to the potential

relevance of blood transcriptional phenotyping for stratification of patients with RSV

infection. More specifically, a signature putatively attributed to immunosuppressive

erythroid cells was found to be associated with clinical severity, even in homogenously

younger patients. Clinical relevance of such candidate biomarker signature would need

to be assessed next. Further investigation of circulating erythroid cells population in the

context of RSV infection are also warranted by these findings. One of the central

questions being whether these cells merely accompany clinical worsening of the

disease or constitute one of its drivers. More generally, this work also highlights the

need for follow on large-scale blood transcriptome profiling studies of responses to RSV

patients, especially over multiple time points. Coordination and cooperation between the

groups that may engage in such endeavors would also prove beneficial for generating

large inter-operable blood transcriptome dataset collections.

ACKNOWLEDGEMENTS

Medicine is a member of the Qatar Foundation for Education, Science and Community

Development. We would like to acknowledge Insight Editing London for editing the

manuscript prior to submission.

CONTRIBUTIONS

Conceptualization: DR, DC, DB, EM, DK, MA. Data curation and validation: DR, MT,

MG, BK, OK, SA, FM. Visualization: DR, DC, OK, SA, FM. Analysis and interpretation:

DR, OK, SH, FM, AM, OR, EM, DK, DC. Writing of the first draft: DR, DC. Funding

acquisition: DK, DC. Methodology development: DR, DC. Writing – review & editing:

DR, MA, MT, MG, BK, OK, SH, FM, AM, OR, DB, EM, DK, DC. The contributor's roles

listed above follow the Contributor Roles Taxonomy (CRediT) managed by The

Consortia Advancing Standards in Research Administration Information (CASRAI)

(https://casrai.org/credit/).

DECLARATIONS OF INTERESTS

The authors declare no competing interests.

FIGURE LEGENDS

Figure 1: Modular repertoire changes in patients with RSV infection vs uninfected

controls. A. Fingerprint heatmap comparing the module aggregate-level patterns of

transcript abundance across six RSV datasets. The summarized module aggregate-

level values on this heatmap are arranged in rows and the datasets in columns. The

datasets are grouped via hierarchical clustering, according to similarities in patterns of

transcript abundance across module aggregates. B. Fingerprint grid for GSE77087.

Modules are assigned a fixed position on the grid, with each row corresponding to a

"module aggregate" constituted of modules following similar patterns of change in

transcript abundance. The numbers of constitutive modules for each aggregate range

from two (A16) to 42 (A2). Aggregates comprising a single module are not represented

on this map (A9-A14; A19-A23). The percentage of constitutive transcripts for a given

module showing an increase in abundance in RSV patients over controls is indicated by

a red spot. The percentage of constitutive transcripts showing a decrease in abundance

for a given module is indicated by a blue spot. The color key at the bottom indicates the

functions that have been associated with some of the modules on the grid.

Figure 2. Heatmap representation of changes in abundance of transcriptional

modules across RSV infected individuals. This heatmap was generated for the

GSE77087 dataset that is also represented as a fingerprint grid plot in Figure 1B. The

modules comprised in aggregates identified as being conserved (indicated by the

colored triangles in Figure 1A) are arranged as rows and the RSV subjects comprised in

this dataset are arranged as columns. The colored spots represent the percentage of

transcripts within each module deemed to be differentially expressed (up = red, down =

blue). The modules are arranged based on similarities in abundance patterns via

hierarchical clustering within each aggregate. A general function is attributed to some of

the aggregates, as indicated by the colored symbols and key.

Figure 3: Individual subjects PCA plots and tSNE plots. The modular fingerprint

profiles of individual subjects from each of the six different studies were combined and

PCA and tSNE plots were generated. The total number of subjects was 490 (319

subjects with RSV infection and 171 controls). **A.** 2D PCA plots. The subjects are color-

coded according to grouping information (Groups) and dataset membership (Studies).

B. tSNE plots. In both types of plot, each sample is represented by a dot. The proximity

between the dots is an indication of the modular transcriptional profile similarity.

Figure 4: Stratification of RSV patients according to blood transcriptome

phenotypes. Phenotypes were defined according to four different "traits". Each of the

319 RSV patients comprising the consolidated cohort used in this meta-analysis were

assigned to a phenotype according to their status for each of the four traits: positive

(red), negative (blue) or unchanged (white). This determination was made in reference

to each respective control baseline. The bar graph shows the number of patients

assigned to each of the phenotypes. The gray line indicates the threshold used to select

the phenotypes considered to be the most abundant (> 10 subjects).

Figure 5: The association of dominant RSV blood transcriptional phenotypes with

clinical and demographic attributes. Heatmaps were generated for the seven most

prevalent phenotypes identified on the distribution plot in Figure 4. The subjects (rows)

were first arranged according to their phenotype, and then arranged within each

phenotype according to similarities in abundance patters. Modules constituting the four

aggregates selected for the definition of molecular phenotypes are shown as columns.

The respective traits for module aggregates A28, A35, A27 and A37 are IFN

(interferon), Infl (inflammation), CC (cell cycle) and Ery (Erythrocytes). The status for

each phenotype is indicated by a red, blue or white spot (increase, decrease or no

change, respectively). The concentric circle plots (right) indicate the distribution of

patients constituting each phenotype according to age, study membership and RSV

severity status.

Figure 6: The association of dominant RSV blood transcriptional phenotypes with

disease severity as a function of age. A. The relative frequencies of patients in PICU

care across dominant IFN/Infl/CC/Ery phenotypes. Bottom panel: the red or blue spots

define the status for four traits corresponding to the eight most prevalent RSV

phenotypes identified in Figure 4 (>10 subjects). Middle panel: the relative frequency of

subjects cared for in the hospital ward, PICU, outpatients, and experimentally exposed

subjects. This information was not available for all studies. Top panel: the relative

contribution of the different datasets selected for this meta-analysis. The combination of

tables and graphs on the left is for all subjects. Information for a subset of subjects <4

months-of-age is shown on the right. B. Ages according to Ery trait status for all patients

and a subset aged <4 months old. The box plots represent the age in months of

individuals comprising the consolidated RSV cohort used throughout this study. Patients

were categorized according to their Ery trait status. The plot on the right shows the

same information but of a subset of patients <4 months old. Dots are color-coded

according to severity status (Red = PICU, Blue = ward, Yellow = outpatient).   * p< 0.05,

** p< 0.01, *** p< 0.001.

Figure 7: Association of blood transcriptomic traits with RSV disease severity in infants < 4 month of age. Box plots (left) show levels of transcript abundance measured for individual subjects for a given aggregate. This value represents the percentage of transcripts constituting the aggregate that are increased or decreased for an individual compared to the median of the uninfected control group (+100% = all transcripts are increased; -100% = all transcripts are decreased). Individuals are grouped according to health status: uninfected controls, inpatients cared for in the hospital ward or inpatients cared for in the PICU. * p< 0.05, ** p< 0.01, *** p< 0.001.

The "Signature survival" curves (right) represent the relative frequency of subjects (y-axis) for whom the percentage response falls at or above a given threshold (x-axis). The percentage response was calculated in the same manner as described for the box plot. Thus, all subjects would have a percentage of response falling between -100% and +100%, as indicated by the curves showing the frequency values of 1 at x = -100%. As the range narrows, the frequency decreases; in most cases a very small proportion of patients have a percentage of response values falling between +90% and +100% (at x = +90%). The separation of the curves is an indication of the differences in the distribution of percentage responses between groups (pale blue = control, dark blue = ward, red = PICU).

Figure 8: Contextual interpretation of RSV blood transcriptome fingerprints A. Fingerprint grid plots displaying changes in the levels of transcript abundance in

patients with RSV infection, Influenza infection, Systemic Lupus Erythematosus, stage

IV melanoma or in liver transplant recipients. The visualization scheme is similar to the

one described for the fingerprint grid plot in Figure 1B. The four traits used to

molecularly stratify RSV patients are highlighted. Because the position of the modules

on the grid is fixed, the color key from Figure 1 can also be used for functional

interpretation of the modules from the other rows. The datasets on which the fingerprint

maps are based are publicly available under GEO accession ID GSE100150. B.

Heatmaps displaying the changes in transcript abundance for modules belonging to four

aggregates (columns) across 16 reference datasets. As for the grid plots, an increase

and decrease in the abundance of transcripts constituting these modules are shown by

a red or blue spot, respectively. The rows (datasets from each disease cohort) and

columns (modules) were arranged by hierarchical clustering based on similarities in

patterns of transcript abundance. All the plots can be generated and exported via a web

application: https://drinchai.shinyapps.io/dc_gen3_module_analysis/# video

demonstration https://youtu.be/y__7xKJo5e4.

Figure 9: Expression levels of A37 genes across cell populations isolated from

human peripheral blood and cord blood. The abundance levels of transcripts

comprised in the 11 modules constituting A37 (columns) across blood-cell populations

(rows). The dataset is publicly available under GEO accession ID GSE24759 (24). The

populations are separated based on whether they were isolated from adult venous

blood (top) or from neonate cord blood (bottom). Distinct erythroid cell populations

isolated on the basis of cell surface expression of CD34, CD71 and GlyA antigens are

also shown.

Supplementary Figure 1: Fingerprint grid maps of modular repertoire changes in

six independent RSV blood transcriptome datasets. The visualization scheme is

similar to the one described for the fingerprint grid map in Figure 1B. Because the

position of the modules on the grid is fixed, the color key from Figure 1 can also be used

for functional interpretation. The maps were generated from multiple independent RSV

blood transcriptome datasets that are available in the NCBI Gene Expression Omnibus

(5, 6, 20–22, 28).

Supplementary Figure 2: Inter-individual variance of summarized module-level

abundance values. The variance in the percentage of the transcript response across

319 RSV patients was calculated across 382 modules. The box plots display those

values with each module being grouped in its respective module aggregate, from A1 to

A38.

Supplementary Figure 3: Patterns of abundance of six A28 modules across

several disease and physiological states. Each column on the heatmap corresponds

to one of six modules constituting the A28 aggregate. Each row corresponds to one of

16 reference datasets. Red spots on the heatmap indicate an increase in abundance of

the transcripts constituting a given module for a given dataset. A blue spot indicates a

decrease in abundance. No color indicates no change. Disease or physiological states

were arranged based on similarity in patterns of aggregate activity. The circle (top left)

is a representation of the six modules constituting aggregate 28, and of the transcripts

constituting each of the modules. Some genes on the Illumina BeadArrays can map to

multiple probes, which explains the few instances where the same gene can be found in

different modules. The smaller circles (right) represent changes in transcript abundance

of individual transcripts for two of the disease clusters. The first cluster (red arrow)

comprises four diseases with the highest degree of increase in transcript abundance.

The second cluster (orange arrow) comprises five diseases with an intermediate level of

increase in abundance. The insert (bottom) shows two circles representing changes in

abundance in patients with hepatitis C treated with IFNα (51), and patients with multiple

sclerosis treated with IFNβ (52). The corresponding NCBI GEO accession IDs are

indicated for each.

Supplementary Figure 4: The expression levels of A36 genes across the cell

populations isolated from human peripheral blood and cord blood. The abundance

levels of transcripts comprised in the 11 modules constituting aggregate A36 (columns)

across blood cell populations (rows). The dataset is publicly available under GEO

accession ID GSE24759 (24). The populations are separated based on whether they

were isolated from adult venous blood (top) or from neonate cord blood (bottom).

Distinct erythroid cell populations isolated on the basis of cell surface expression of

CD34, CD71 and GlyA antigens are shown.

Supplementary Figure 5: The expression levels of A38 genes across the cell

populations isolated from human peripheral blood and cord blood. The abundance

levels of transcripts comprised in the 11 modules constituting aggregate A38 (columns)

across blood cell populations (rows). The dataset is publicly available under GEO

accession ID GSE24759 (24). The populations are separated based on whether they

were isolated from adult venous blood (top) or from neonate cord blood (bottom).

Distinct erythroid cell populations isolated on the basis of cell surface expression of

CD34, CD71 and GlyA antigens are shown.

758 **TABLES**
759
760 **Table 1: Description of the public RSV blood transcriptome datasets.**

GEO ID	Reference	Title	N Subjects	Age demogra-phics	Sample type	Platform	Cluster # (Figure 1A)
GSE42026	Herberg et al. / 23901082 / (20)	Transcriptomic profiling in childhood H1N1/09 influenza reveals reduced expression of protein synthesis genes	22 RSV & 33 controls	Pediatric	Whole blood (PaxGene)	Illumina HumanHT-12 V3.0	1
GSE38900	Mejias et al. / 24265599 / (28)	Genome-wide analysis of whole blood transcriptional response to RSV, Influenza and Rhinovirus (LRTI) in children	107 RSV & 31 controls	Pediatric	Whole blood (Tempus)	Illumina Human HT-12 V3.0]	1
GSE80179	McDonald et al. /27822537 / (22)	A simple screening approach to prioritize genes for functional analysis identifies a role for IRF7 in the control of RSV disease.	27 RSV & 52 controls	Pediatric	Whole blood (PaxGene)	Illumina HumanHT-12 V4.0	1
GSE73072	Liu et al. / 26801061 / (21)	Host gene expression signatures of H1N1,	20 subjects 2 time points	Adult	Whole blood (PaxGene)	Affymetrix GeneChip Human	2

		H3N2, HRV, RSV virus infection in adults				Genome U133A 2.0 Array	
GSE77087	de Steenhuijsen Piters et al. / 27135599 / (6)	Nasopharyngeal microbiota, host transcriptome and disease severity in children with RSV infection	81 RSV & 23 controls	Pediatric	Whole blood (Tempus)	Illumina HumanHT-12 V4.0	2
GSE103842	Rodriguez-Fernandez et al. / 29045741 / (5)	RSV Genotypes and Disease Severity in Young Children Hospitalized with Bronchiolitis	62 RSV & 12 controls	Pediatric	Whole blood (Tempus)	Illumina HumanHT-12 V4.0	2

762
763   RSV: respiratory syncytial virus, LRTI: lower respiratory tract infection
774   **Table 2: Links to module aggregates annotation pages**

Aggregate	Function	Links
A1	Lymphocytic	https://prezi.com/view/sxap39tKxkmCNTTNIlVO/
A2	TBD	https://prezi.com/view/96GWajx5mZjuRS4B6gjA/
A3	TBD	https://prezi.com/view/OWFVI51FND0WWwNgsgJZ/
A4	TBD	https://prezi.com/view/2Zbq8ZDYbO4hbUd4r2KF/
A5	Lymphocytic	https://prezi.com/view/62tgA5E6roOlk5DRNvS1/
A6	Lymphocytic	https://prezi.com/view/Uks2Nd4lvizNNFVPtBEy/
A7	TBD	https://prezi.com/view/kKfergNj0SkLXyFtm0Dg/
A8	TBD	https://prezi.com/view/Y4uk1RPJyNcSndJYnFX6/
A9	TBD	https://prezi.com/view/jgYehQ9QhyADAttEsdoI/
A15	TBD	https://prezi.com/view/jgYehQ9QhyADAttEsdoI/
A16	TBD	https://prezi.com/view/SKzHeA0XYdLYvy2sY8gP/
A17	TBD	https://prezi.com/view/FS7sE1Vqew5g8EKOM1AM/
A18	TBD	https://prezi.com/view/aZMLflMNVrV7JnVaIILm/
A24	Oxidative phosphorylation	https://prezi.com/view/eiXvf2LNBLFRgrtaeCuM/
A25	TBD	https://prezi.com/view/pwyojaU62Z7GT102ZYwM/
A26	TBD	https://prezi.com/view/9CErpW3NwpN2HgRS3Hzf/
A27	Cell cycle	https://prezi.com/view/GgIiA0K9kSFHbpVj2I85/

A28	Interferon	https://prezi.com/view/E34MhxE5uKoZLWZ3KXjG/
A29	TBD	https://prezi.com/view/W4TShTd32dEJx0XPOF1U/
A30	TBD	https://prezi.com/view/kl7VHoJTWug0sn7TgXut/
A31	TBD	https://prezi.com/view/GqtUO22JJlSf16zMJKbB/
A32	TBD	https://prezi.com/view/qlbG9VFzegOndQKD8aiy/
A33	Inflammation	https://prezi.com/view/VBqKqHuLWCra3OJOIZRR/
A34	TBD	https://prezi.com/view/HcSgIEGP3TJjTSpaPCxv/
A35	Inflammation	https://prezi.com/view/7Q20FyW6Hrs5NjMaTUyW/
A36	Erythroid	https://prezi.com/view/M7dnztl2h61gXrKFQeB2/
A37	Erythroid	https://prezi.com/view/YyQs4WiXSNf0YXE79lfS/
A38	Erythroid	https://prezi.com/view/0KUlPICKUZGeUjb206R5/

776
777

METHODS

Selection of public blood transcriptome datasets

Datasets deposited in the NCBI Gene Expression Omnibus, GEO, were used in this meta-analysis. Accession IDs along with descriptive information and references can be found in **Table 1**. A reference dataset, which consisted of transcriptome profiles derived from adult blood cell populations and cord blood was also used to support the functional interpretation of our findings. This dataset was contributed to the GEO by Novershtern et al. with accession ID GSE24759 (24).

Module repertoire construction

The construction of a transcriptional module repertoire for blood transcriptome analyses has been described previously (48,49). The version that was used in this study is the

third one developed by our group and is the object of a separate publication (available

on a pre-print server (19)). Briefly, the approach consists of identifying sets of co-

expressed transcripts for a given biological system (in this case blood) and across a

wide range of disease or physiological states (perturbations of steady state). In this

case, co-expression was determined based on patterns of co-clustering observed for all

gene pairs across a collection of 16 reference datasets. These datasets encompassed

viral and bacterial infectious diseases (HIV, influenza, RSV, Melioidosis,

Staphylococcus aureus, Tuberculosis) as well as several inflammatory or autoimmune

diseases (systemic lupus erythematosus, multiple sclerosis, chronic obstructive

pulmonary disease, Kawasaki disease, juvenile dermatomyosistis, systemic onset

juvenile idiopathic arthritis), B-cell deficiency, liver transplantation, stage IV melanoma

and pregnancy. The overall collection comprised 985 blood transcriptome profiles. A

weighted co-expression network was built on the basis of co-clustering patterns that

were obtained. Here, the weight of the nodes connecting a gene pair being based on

the number of times co-clustering was observed, thus ranging from a weight of 1 (where

co-clustering occurs in one of 16 datasets) to a weight of 16 (where co-clustering occurs

in all 16 datasets). Next, this network was mined using a graph theory algorithm

(identification of cliques and paracliques) to define a subset of densely connected gene

sets that constituted our module repertoire. Overall, 382 modules were identified via this

process, encompassing 14,168 transcripts. A supplemental file including the definition

of this module repertoire along with the functional annotations is available from a

companion publication.

Constitution of module aggregates

To maintain the number of variables within a manageable number and to facilitate data interpretation, a second tier of clustering was performed to group the modules into "aggregates". This was achieved by segregating the set of 382 modules according to the patterns of transcript abundance across the 16 reference datasets that were used for module construction. This segregation resulted in the formation of 38 aggregates, each comprising between one and 42 modules. The second level of granularity that was thus obtained was used to define distinct RSV blood transcriptome phenotypes and as a basis for the fingerprint grid plot representation (see **Figure 1** and **Figure 7**). With such grids, the first vertical reading of the fingerprint grid provides an overview of the changes in transcript abundance observed among module aggregates, while the horizontal reading provides the changes observed within an aggregate and across modules.

Module repertoire analysis workflow

The modular analysis was performed using 14,168 transcripts. The fold change was computed using gene expression data prior to log2 transformation. For group comparisons, a paired t-test was performed on the log2-transformed data [Fold change (FC) cut off = 1.5; FDR cut off = 0.1]. For individual-patients analyses, each sample was compared to the mean value of control samples in each dataset. The cut off comprised an absolute FC >1.5 and a difference in gene expression level >10. The results for each module analysis are reported as the percentage of its constitutive transcripts for which the abundance was increased or decreased. Because gene sets are selected based on

the co-expression observed in blood, the changes in abundance within a given module

tend to be coordinated and the dominant trend is therefore selected (the greater value

of the percentage increased vs percentage decreased). A module was considered to be

"responsive" when the proportion of differentially expressed transcripts (as defined

above) was >15%.

Data visualization

The results were visualized in a fingerprint format, either as a grid plot (group level,

Figure 1) or as a heatmap (individual level, **Figure 5**), using the same illustrative RSV

dataset. For each module, the percentage of increased transcripts is represented by a

red spot and the percentage of decreased transcripts is represented by a blue spot. The

largest of the two values was retained for visualization. In the grid format (**Figure 1**), the

position of each module is fixed. A row of modules corresponds to a "module

aggregate", which as described above, is a set of modules following a similar pattern of

activity across the 16 input datasets corresponding to different disease or physiological

states. A few "aggregates" comprised only a single module and thus are not shown on

the grid. The fingerprint grid plots were generated using "ComplexHeatmap" (50).

For the heatmaps (**Figure 2, Figure 5, Figure 7B**), each row corresponds to a module

and each column to a sample. The columns and rows are arranged based on

similarities in the patterns of module activity. Filters can be applied to remove modules

that show only low levels of activity across the samples or to retain only the modules

associated with functional annotations.

REFERENCES

1.   Shi T, McAllister DA, O'Brien KL, Simoes EAF, Madhi SA, Gessner BD, et al. Global, regional,
and national disease burden estimates of acute lower respiratory infections due to
respiratory syncytial virus in young children in 2015: a systematic review and modelling
study. Lancet. 2017 Sep 2;390(10098):946–58.

2.   Hall CB, Weinberg GA, Iwane MK, Blumkin AK, Edwards KM, Staat MA, et al. The burden of
respiratory syncytial virus infection in young children. N Engl J Med. 2009 Feb
5;360(6):588–98.

3.   García CG, Bhore R, Soriano-Fallas A, Trost M, Chason R, Ramilo O, et al. Risk factors in
children hospitalized with RSV bronchiolitis versus non-RSV bronchiolitis. Pediatrics. 2010
Dec;126(6):e1453-1460.

4.   Ascough S, Paterson S, Chiu C. Induction and Subversion of Human Protective Immunity:
Contrasting Influenza and Respiratory Syncytial Virus. Front Immunol. 2018;9:323.

5.   Rodriguez-Fernandez R, Tapia LI, Yang C-F, Torres JP, Chavez-Bueno S, Garcia C, et al.
Respiratory Syncytial Virus Genotypes, Host Immune Profiles, and Disease Severity in
Young Children Hospitalized With Bronchiolitis. J Infect Dis. 2017 27;217(1):24–34.

6.   de Steenhuijsen Piters WAA, Heinonen S, Hasrat R, Bunsow E, Smith B, Suarez-Arrabal M-
C, et al. Nasopharyngeal Microbiota, Host Transcriptome, and Disease Severity in Children
with Respiratory Syncytial Virus Infection. Am J Respir Crit Care Med. 2016
01;194(9):1104–15.

7.   Mejias A, Dimo B, Suarez NM, Garcia C, Suarez-Arrabal MC, Jartti T, et al. Whole blood
gene expression profiles to assess pathogenesis and disease severity in infants with
respiratory syncytial virus infection. PLoS Med. 2013 Nov;10(11):e1001549.

8.   McClain MT, Park LP, Nicholson B, Veldman T, Zaas AK, Turner R, et al. Longitudinal
analysis of leukocyte differentials in peripheral blood of patients with acute respiratory
viral infections. J Clin Virol. 2013 Dec;58(4):689–95.

9.   Do LAH, Pellet J, van Doorn HR, Tran AT, Nguyen BH, Tran TTL, et al. Host Transcription
Profile in Nasal Epithelium and Whole Blood of Hospitalized Children Under 2 Years of Age
With Respiratory Syncytial Virus Infection. J Infect Dis. 2017 27;217(1):134–46.

10.   Brand HK, Ahout IML, de Ridder D, van Diepen A, Li Y, Zaalberg M, et al. Olfactomedin 4
Serves as a Marker for Disease Severity in Pediatric Respiratory Syncytial Virus (RSV)
Infection. PLoS ONE. 2015;10(7):e0131927.

11.   Bucasas KL, Mian AI, Demmler-Harrison GJ, Caviness AC, Piedra PA, Franco LM, et al.
Global gene expression profiling in infants with acute respiratory syncytial virus

broncholitis demonstrates systemic activation of interferon signaling networks. Pediatr
Infect Dis J. 2013 Feb;32(2):e68-76.

12. Fjaerli H-O, Bukholm G, Krog A, Skjaeret C, Holden M, Nakstad B. Whole blood gene
expression in infants with respiratory syncytial virus bronchiolitis. BMC Infect Dis. 2006
Dec 13;6:175.

13. Tsalik EL, Henao R, Nichols M, Burke T, Ko ER, McClain MT, et al. Host gene expression
classifiers diagnose acute respiratory illness etiology. Sci Transl Med. 2016 Jan
20;8(322):322ra11.

14. Banchereau R, Cepika A-M, Banchereau J, Pascual V. Understanding Human Autoimmunity
and Autoinflammation Through Transcriptomics. Annu Rev Immunol. 2017 26;35:337–70.

15. Chaussabel D. Assessment of immune status using blood transcriptomics and potential
implications for global health. Semin Immunol. 2015 Feb;27(1):58–66.

16. Sweeney TE, Haynes WA, Vallania F, Ioannidis JP, Khatri P. Methods to increase
reproducibility in differential gene expression via meta-analysis. Nucleic Acids Res. 2017
09;45(1):e1.

17. Sweeney TE, Perumal TM, Henao R, Nichols M, Howrylak JA, Choi AM, et al. A community
approach to mortality prediction in sepsis via gene expression analysis. Nat Commun. 2018
15;9(1):694.

18. Schaack D, Siegler BH, Tamulyte S, Weigand MA, Uhle F. The immunosuppressive face of
sepsis early on intensive care unit-A large-scale microarray meta-analysis. PLoS ONE.
2018;13(6):e0198555.

19. Altman MC, Rinchai D, Baldwin N, Whalen E, Garand M, Kabeer BA, et al. A Novel
Repertoire of Blood Transcriptome Modules Based on Co-expression Patterns Across
Sixteen Disease and Physiological States. bioRxiv. 2019 Jan 23;525709.

20. Herberg JA, Kaforou M, Gormley S, Sumner ER, Patel S, Jones KDJ, et al. Transcriptomic
profiling in childhood H1N1/09 influenza reveals reduced expression of protein synthesis
genes. J Infect Dis. 2013 Nov 15;208(10):1664–8.

21. Liu T-Y, Burke T, Park LP, Woods CW, Zaas AK, Ginsburg GS, et al. An individualized
predictor of health and disease using paired reference and target samples. BMC
Bioinformatics. 2016 Jan 22;17:47.

22. McDonald JU, Kaforou M, Clare S, Hale C, Ivanova M, Huntley D, et al. A Simple Screening
Approach To Prioritize Genes for Functional Analysis Identifies a Role for Interferon
Regulatory Factor 7 in the Control of Respiratory Syncytial Virus Disease. mSystems. 2016
Jun;1(3).

23.   Maaten L van der, Hinton G. Visualizing Data using t-SNE. Journal of Machine Learning
Research. 2008;9(Nov):2579–605.

24.   Novershtern N, Subramanian A, Lawton LN, Mak RH, Haining WN, McConkey ME, et al.
Densely interconnected transcriptional circuits control cell states in human hematopoiesis.
Cell. 2011 Jan 21;144(2):296–309.

25.   Elahi S. Neglected Cells: Immunomodulatory Roles of CD71+ Erythroid Cells. Trends
Immunol. 2019 Mar;40(3):181–5.

26.   Elahi S, Ertelt JM, Kinder JM, Jiang TT, Zhang X, Xin L, et al. Immunosuppressive CD71+
erythroid cells compromise neonatal host defence against infection. Nature. 2013 Dec
5;504(7478):158–62.

27.   Zhao L, He R, Long H, Guo B, Jia Q, Qin D, et al. Late-stage tumors induce anemia and
immunosuppressive extramedullary erythroid progenitor cells. Nat Med.
2018;24(10):1536–44.

28.   Mejias A, Dimo B, Suarez NM, Garcia C, Suarez-Arrabal MC, Jartti T, et al. Whole blood
gene expression profiles to assess pathogenesis and disease severity in infants with
respiratory syncytial virus infection. PLoS Med. 2013 Nov;10(11):e1001549.

29.   Jans J, Pettengill M, Kim D, van der Made C, de Groot R, Henriet S, et al. Human newborn B
cells mount an interferon-α/β receptor-dependent humoral response to respiratory
syncytial virus. J Allergy Clin Immunol. 2017;139(6):1997-2000.e4.

30.   Roberts NJ, Hiscott J, Signs DJ. The limited role of the human interferon system response
to respiratory syncytial virus challenge: analysis and comparison to influenza virus
challenge. Microb Pathog. 1992 Jun;12(6):409–14.

31.   Aberle JH, Aberle SW, Dworzak MN, Mandl CW, Rebhandl W, Vollnhofer G, et al. Reduced
interferon-gamma expression in peripheral blood mononuclear cells of infants with severe
respiratory syncytial virus disease. Am J Respir Crit Care Med. 1999 Oct;160(4):1263–8.

32.   Aberle JH, Aberle SW, Rebhandl W, Pracher E, Kundi M, Popow-Kraupp T. Decreased
interferon-gamma response in respiratory syncytial virus compared to other respiratory
viral infections in infants. Clin Exp Immunol. 2004 Jul;137(1):146–50.

33.   Schauer U, Hoffjan S, Rothoeft T, Bartz H, Konig S, Fuchs E, et al. Severe respiratory
syncytial virus infections and reduced interferon-gamma generation in vitro. Clin Exp
Immunol. 2004 Oct;138(1):102–9.

34.   Linsley PS, Speake C, Whalen E, Chaussabel D. Copy number loss of the interferon gene
cluster in melanomas is linked to reduced T cell infiltrate and poor patient prognosis. PLoS
ONE. 2014;9(10):e109760.

35.   Rawat A, Rinchai D, Toufiq M, Marr A, Kino T, Garand M, et al. A neutrophil-driven
inflammatory signature characterizes the blood cell transcriptome fingerprints of Psoriasis
and Kawasaki Disease. bioRxiv. 2020 Feb 25;2020.02.24.962621.

36.   Bera MM, Lu B, Martin TR, Cui S, Rhein LM, Gerard C, et al. Th17 cytokines are critical for
respiratory syncytial virus-associated airway hyperreponsiveness through regulation by
complement C3a and tachykinins. J Immunol. 2011 Oct 15;187(8):4245–55.

37.   Mebratu YA, Tesfaigzi Y. IL-17 Plays a Role in Respiratory Syncytial Virus-induced Lung
Inflammation and Emphysema in Elastase and LPS-injured Mice. Am J Respir Cell Mol Biol.
2018;58(6):717–26.

38.   Zhang G, Zhou KF, Lu ZH. Interleukin-17 enhances the removal of respiratory syncytial
virus in mice by promoting neutrophil migration and reducing interferon-gamma
expression. Genet Mol Res. 2016 Mar 24;15(1).

39.   Cole S, Walsh A, Yin X, Wechalekar MD, Smith MD, Proudman SM, et al. Integrative
analysis reveals CD38 as a therapeutic target for plasma cell-rich pre-disease and
established rheumatoid arthritis and systemic lupus erythematosus. Arthritis Res Ther.
2018 02;20(1):85.

40.   Lin P, Owens R, Tricot G, Wilson CS. Flow cytometric immunophenotypic analysis of 306
cases of multiple myeloma. Am J Clin Pathol. 2004 Apr;121(4):482–8.

41.   Owczarczyk K, Lal P, Abbas AR, Wolslegel K, Holweg CTJ, Dummer W, et al. A plasmablast
biomarker for nonresponse to antibody therapy to CD20 in rheumatoid arthritis. Sci Transl
Med. 2011 Sep 21;3(101):101ra92.

42.   Obermoser G, Presnell S, Domico K, Xu H, Wang Y, Anguiano E, et al. Systems scale
interactive exploration reveals quantitative and qualitative differences in response to
influenza and pneumococcal vaccines. Immunity. 2013 Apr 18;38(4):831–44.

43.   Habibi MS, Jozwik A, Makris S, Dunning J, Paras A, DeVincenzo JP, et al. Impaired Antibody-
mediated Protection and Defective IgA B-Cell Memory in Experimental Infection of Adults
with Respiratory Syncytial Virus. Am J Respir Crit Care Med. 2015 May 1;191(9):1040–9.

44.   Delyea C, Bozorgmehr N, Koleva P, Dunsmore G, Shahbaz S, Huang V, et al. CD71+
Erythroid Suppressor Cells Promote Fetomaternal Tolerance through Arginase-2 and PDL-
1. J Immunol. 2018 15;200(12):4044–58.

45.   Alamo IG, Kannan KB, Loftus TJ, Ramos H, Efron PA, Mohr AM. Severe trauma and chronic
stress activates extramedullary erythropoiesis. J Trauma Acute Care Surg. 2017;83(1):144–
50.

46. Jackson A, Nanton MR, O'Donnell H, Akue AD, McSorley SJ. Innate immune activation
during Salmonella infection initiates extramedullary erythropoiesis and splenomegaly. J
Immunol. 2010 Nov 15;185(10):6198–204.

47. Verhoeven D. Influence of Immunological Maturity on Respiratory Syncytial Virus-Induced
Morbidity in Young Children. Viral Immunol. 2019;32(2):76–83.

48. Chaussabel D, Baldwin N. Democratizing systems immunology with modular
transcriptional repertoire analyses. Nat Rev Immunol. 2014;14(4):271–80.

49. Chaussabel D, Quinn C, Shen J, Patel P, Glaser C, Baldwin N, et al. A modular analysis
framework for blood genomics studies: application to systemic lupus erythematosus.
Immunity. 2008 Jul 18;29(1):150–64.

50. Gu Z, Eils R, Schlesner M. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. Bioinformatics. 2016 15;32(18):2847–9.

51. Taylor MW, Tsukahara T, McClintick JN, Edenberg HJ, Kwo P. Cyclic changes in gene
expression induced by Peg-interferon alfa-2b plus ribavirin in peripheral blood monocytes
(PBMC) of hepatitis C patients during the first 10 weeks of treatment. J Transl Med. 2008
Nov 5;6:66.

52. Malhotra S, Bustamante MF, Pérez-Miralles F, Rio J, Ruiz de Villa MC, Vegas E, et al. Search
for specific biomarkers of IFNβ bioactivity in patients with multiple sclerosis. PLoS ONE.
2011;6(8):e23634.

1015

Figure 1

GSE77087: de Steenhuijsen Piters et al.

RSV (81)

% Response

Figure 2

IFN — Interferon

Infl — Inflammation

Ery — Erythrocytes

CC — Cell cycle

Lyc — Lymphocytic

A. PCA

Groups
- Control
- RSV

Studies
- GSE103842
- GSE38900
- GSE42026
- GSE73072
- GSE77087
- GSE80179

B.

tSNE

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

A37 (11 modules)

z scores
3
2
1
0
-1

naive B-cells
mature B-cell able to switch
mature B-cells
mature B-cell class switched
plasmacytoid DC
myeloid DC
mature NK cell CD56- CD16+ CD3-
mature NK cell CD56+ CD16+ CD3-
mature NK cell CD56- CD16- CD3-
NKT
CD8+ Effector Memory RA
naive CD8+ T-cell
CD8+ Effector Memory
CD8+ Central Memory
naive CD4+ T-cell
CD4+ Effector Memory
CD4+ Central Memory

ADULT BLOOD

basophils
common myeloid progenitor
eosinophil
erythroid CD34+ CD71+ GlyA-   } erythroid GlyA-
erythroid CD34- CD71+ GlyA-
erythroid CD34- CD71+ GlyA+
erythroid CD34- CD71lo GlyA+   } erythroid GlyA+
erythroid CD34- CD71- GlyA+
granulocyte/monocyte progenitor
colony forming unit - granulocyte
neutrophilic metamyelocyte
neutrophil
hematopoietic SC CD133+ CD34dim
hematopoietic SC CD133+ CD34dim
colony forming unit megakaryocitic
megakaryocyte
megakaryocyte / erythroid progenitor
colony forming unit monocyte
monocyte
early B-cell
pro B-cell

CORD BLOOD

GSE24759 samples

A37 Genes

Figure 9

RSV

OVER UNDER

% significant transcripts

cutoff p<0.05

GSE42026: Herberg et al.

GSE38900: Mejias et al.

GSE73072: Liu et al.

GSE77087: de Steenhuijsen Piters et al.

GSE80179: McDonald et al.

GSE103842: Rodriguez-Fernandez et al.

Supplementary Figure 1

Supplementary Figure 3

Supplementary Figure 4

A38   x 4 modules

z scores
3
2
1
0
-1

naive B-cells
mature B-cell able to switch
mature B-cells
mature B-cell class switched
plasmacytoid DC
myeloid DC
mature NK cell CD56- CD16+ CD3-
mature NK cell CD56+ CD16+ CD3-
mature NK cell CD56- CD16- CD3-
NKT
CD8+ Effector Memory RA
naive CD8+ T-cell
CD8+ Effector Memory
CD8+ Central Memory
naive CD4+ T-cell
CD4+ Effector Memory
CD4+ Central Memory

ADULT BLOOD

basophils
common myeloid progenitor
eosinophil
erythroid CD34+ CD71+ GlyA-
erythroid CD34- CD71+ GlyA-

erythroid GlyA-

erythroid CD34- CD71+ GlyA+
erythroid CD34- CD71lo GlyA+
erythroid CD34- CD71- GlyA+

erythroid GlyA+

granulocyte/monocyte progenitor
colony forming unit - granulocyte
neutrophilic metamyelocyte
neutrophil
hematopoietic SC CD133+ CD34dim
hematopoietic SC CD133+ CD34dim
colony forming unit megakaryocitic
megakaryocyte
megakaryocyte / erythroid progenitor
colony forming unit monocyte
monocyte
early B-cell
pro B-cell

CORD BLOOD

Supplementary Figure 5