1

Paternally-acting canonical RNA-directed DNA methylation pathway genes

sensitize Arabidopsis endosperm to paternal genome dosage

P.R. V. Satyaki[1], Mary Gehring[1,2]*

[1]Whitehead Institute for Biomedical Research, [2]Department of Biology, Massachusetts
Institute of Technology, Cambridge MA 02142

* author for correspondence: mgehring@wi.mit.edu

Short Title: Small RNA pathways control paternal dosage response

Abstract

Seed development is sensitive to parental dosage, with excess maternal or paternal genomes creating reciprocal phenotypes. Paternal genomic excess frequently results in extensive endosperm proliferation without cellularization and seed abortion. We previously showed that loss of the RNA Pol IV gene *nrpd1* in tetraploid fathers represses seed abortion in paternal excess crosses. Here we show genetically that RNA-directed DNA methylation (RdDM) pathway activity in the paternal parent is sufficient to determine the viability of paternal excess seeds. We compared transcriptomes, DNA methylation, and small RNAs from endosperm of balanced crosses (diploid x diploid) and lethal (diploid x tetraploid) and viable paternal excess (diploid x tetraploid *nrpd1*). Endosperm from both lethal and viable paternal excess seeds share widespread transcriptional and DNA methylation changes at genes and TEs. Interploidy seed abortion is thus unlikely to be caused by either transposable element or imprinted gene mis-regulation, and its repression by loss of paternal RdDM is associated with only modest gene expression changes. Finally, using allele-specific transcription data, we present evidence for a transcriptional buffering system that increases expression of maternal alleles and represses paternal alleles in response to excess paternal genomic dosage. These findings prompt reconsideration of models for dosage sensitivity in endosperm.

Introduction

The endosperm of flowering plants is an essential tissue for seed viability. It is most commonly triploid, formed when the diploid central cell is fertilized by a haploid sperm. In many flowering plants, endosperm proceeds through the early phases of proliferative development as a syncytium before differentiating into three sub-types: micropylar, peripheral and chalazal endosperm (Li and Berger, 2012). Cellularization is a critical step in endosperm development, after which cell division slows and eventually ceases (Hehenberger et al., 2012; Li and Berger, 2012). In some species, like Arabidopsis, the endosperm is almost completely degraded at seed maturation as nutrients are assimilated and stored in the embryo (Li and Berger, 2012), whereas in other species, like grasses, it is persistent and mobilized later during seed germination. The endosperm serves several important functions: it mediates resource transfer from the mother to the growing embryo or germinating seedling (Li and Berger, 2012), signals to the maternal integuments to promote their proliferation and allow accommodation of the growing offspring (Figueiredo et al., 2016), is required for embryonic development, and influences seed dormancy and germination (Fiume and Fletcher, 2012; Piskurewicz et al., 2016; Yan et al., 2014). Through these activities, endosperm influences seed size.

Balance between the maternal and paternal genomes in endosperm is important for normal endosperm development and, consequently, seed development and viability. Violations of the 2:1 maternal:paternal (m:p) genome ratio in endosperm lead to developmental defects, although species vary in the extent to which they tolerate violations of this ratio (Cooper, 1951; Esen and Soost, 1973; Håkansson and Ellerström, 1950; Milbocker and Sink, 1969; Muntzing, 1936; Povilus et al., 2018; Scott et al., 1998, 1998; Stoute et al., 2012). In Arabidopsis, increased maternal genome dosage (maternal excess) leads to premature cellularization of the endosperm and the formation of smaller seeds (Scott et al., 1998). By contrast, increased paternal genome dosage (paternal excess) in crosses between diploid mothers and hexaploid fathers leads to a failure of endosperm cellularization, prolonged cell proliferation, and seed abortion (Scott et al., 1998). However, there is intra-specific variation in the levels of

seed abortion observed in paternal excess crosses. For example, whereas tetraploid

Col-0 induces seed abortion when pollinating diploid mothers, tetraploid C24 and Cvi do

not (Dilkes et al., 2008; Lu et al., 2012; Piskurewicz et al., 2016; Scott et al., 1998).

What determines the threshold between seed lethality and viability in conditions

of paternal genomic excess? Mutants that repress seed abortion in interploidy crosses,

as well data from incompatible interspecific crosses, which share features with

interploidy crosses, have provided some answers to this question. Components in the

endosperm, in the gametophytes, or in the parental sporophytes, have been proposed

to be responsible for endosperm dosage sensitivity in interploid crosses. Some of the

elements that determine the critical threshold might be linked to interactions between

the endosperm and maternal genotype. In one model, the imbalance is between the

paternal dose and the female gametophyte (Birchler, 2014; von Wangenheim and

Peterson, 2004). A second model involves interactions between the paternal excess

endosperm and the diploid maternal integuments, which develop into the seed coat

(Muntzing, 1936). The ability of maternal sporophytic mutations in the flavonoid pathway

to repress paternal excess seed abortion is consistent with this model (Dilkes et al.,

2008; Doughty et al., 2014). The levels of *AGLs*, the small RNA *siRNA854*, chromatin

regulators, cell wall genes, and defense response genes, all presumed to be acting in

the endosperm, have been linked to paternal excess seed abortion (Burkart-Waco et al.,

2013; Jiang et al., 2017; Kradolfer et al., 2013; Walia et al., 2009; Wang et al., 2018;

Wolff et al., 2015).

Multiple models suggest that endosperm ploidy incompatibilities are caused by

epigenetic abnormalities. The epigenetic mark DNA methylation is established and

partially maintained through the RNA-directed DNA methylation (RdDM) pathway.

During this process, short non-coding RNAs generated by RNA Pol IV are converted

into double-stranded RNAs by RDR2. These are then subsequently diced by DCL3 into

24 nt small RNAs (sRNAs), loaded into an ARGONAUTE complex (usually AGO4 or

AGO6), and then interact with a non-coding RNA transcribed by another polymerase,

RNA Pol V, to direct DNA methylation by the *de novo* methyltransferase DRM2. In

normal triploid endosperm, maternal chromosomes are DNA hypomethylated relative to

paternal chromosomes due to the activity of the DNA demethylase *DME* (Gehring et al.,
2009; Hsieh et al., 2009; Ibarra et al., 2012). DNA methylation represses expression of
some endosperm genes and promotes the expression of others (Satyaki and Gehring,
2017). It has been proposed that a cause of interploid seed abortion is increased
expression of transposable elements (TEs) due to epigenetic alterations (Martienssen,
2010). Under this model, a maternally deposited sRNA dose is insufficient to silence the
doubled number of paternally inherited TEs. Another model is influenced by the parental
conflict theory (Haig and Westboy, 1991). The theory argues that in a polyandrous
system in which the mother provisions resources for her offspring from various fathers,
genes restricting resource allocation to any one seed and favoring equitable distribution
of resources across all the progeny become predominantly maternal in expression
(maternally expressed genes or MEGs). On the other hand, genes promoting resource
allocation and larger seed production are predominantly paternally expressed
(paternally expressed genes or PEGs). This type of allele-specific expression is an
epigenetic phenomenon referred to as gene imprinting. According to these ideas, an
excess dose of paternal chromosomes leads to over-expression of PEGs (genes that
promote endosperm proliferation) eventually leading to seed abortion. Consistent with
this model, PEGs were reported to be over-expressed in paternal excess seeds
(Kradolfer et al., 2013; Schatlowski et al., 2014) and loss-of-function mutations in
multiple PEGs repress seed abortion in conditions of paternal genomic excess (Huang
et al., 2017; Jiang et al., 2017; Kradolfer et al., 2013; Wolff et al., 2015).

        We recently proposed another model based on discoveries about the function of
NRPD1, which codes for the largest subunit of RNA Pol IV, in normal triploid endosperm
(Erdmann et al., 2017). Expression of 24 nt sRNAs in *Arabidopsis thaliana* endosperm
is paternally biased and concentrated in pericentromeric heterochromatin, but a subset
of 24 nt sRNAs are expressed predominantly from the maternal alleles of genes
(Erdmann et al., 2017). We showed that paternal loss of *NRPD1* in 2m:1p triploid
endosperm increased the maternal fraction of the transcriptome, suggesting that wild-
type *NRPD1* represses maternal genome dosage, probably via the production of 24 nt
small RNAs. We also discovered that paternally inherited mutations in *NRPD1* repress
seed abortion caused by excess paternal genomes (i.e. 2N Col x 4N Col), leading to the

hypothesis that the increased maternal fraction of the transcriptome in *nrpd1* mutant
endosperm compensates for increased paternal genomic dosage. In this model, the
loss of RNA Pol IV-dependent sRNA production in the endosperm is essential for
viability (Erdmann et al., 2017).

Finally, the "easiRNA" model argues that *NRPD1* and *RDR6* act together in a
non-canonical pathway in the male gametophyte to produce easiRNAs whose
concentration scales with ploidy (Borges et al., 2018; Martinez et al., 2018). easiRNAs
are defined as 21-22 nt sRNAs produced from the processing of epigenetically activated
TE mRNAs (Creasey et al., 2014). It is proposed that easiRNAs are transmitted during
fertilization from sperm to the central cell, where they inhibit RdDM in the developing
endosperm (Borges et al., 2018; Martinez et al., 2018). In this model, easiRNAs are lost
from sperm when the paternal copy of *NRPD1* is mutated. The loss of paternal
easiRNAs allows the restoration of a functional RdDM pathway in endosperm using
maternal copies of *NRPD1*. The restored RdDM pathway was proposed to repress
excess paternal dosage and restore seed viability.

To test among these models and further understand the nature of intolerance or
tolerance to paternal genomic excess, we tested the genetic and molecular
contributions to interploidy seed abortion and repression. We created tetraploid mutants
for members of the canonical and non-canonical small RNA pathways to test the
genetic requirements for seed abortion in paternal excess crosses. We also profiled
sRNAs, DNA methylation, and gene expression in balanced (2N x 2N), lethal paternal
excess (2N x 4N), and viable paternal excess (2N x 4N *nrpd1*) endosperm. We found
that genes of the canonical RdDM pathway were necessary in the male parent only for
seed abortion induced by paternal genomic excess. Through transcriptomic profiling we
found that there were extensive gene and transposable element expression changes
associated with lethal paternal genomic excess but that only a small fraction were
ameliorated in viable paternal excess endosperm. Our analysis also revealed the
signatures of a potential buffering system that attempts to rebalance transcription in
conditions of paternal excess by repressing paternal alleles and activating maternal
alleles. These observations indicate that endosperm development can tolerate

surprisingly large variations in gene expression at most genes and suggests that

relatively small gene expression changes contribute to the critical dosage that

determines seed viability under conditions of paternal genomic excess.

Results

Paternal loss of canonical RdDM pathway genes is sufficient for suppression of interploid seed abortion

In crosses between diploid females and tetraploid males, tetraploid *nrpd1* mutant

fathers repress seed abortion, whereas diploid *nrpd1* mothers have no effect (Erdmann

et al., 2017; Martinez et al., 2018). We tested if other members of the canonical RdDM

pathway behaved genetically in the same manner. Using colchicine-induced tetraploid

mutants of *shh1, rdr2, dcl3, nrpe1,* and *drm2,* we examined seeds obtained from

crosses between 2N wild-type (Col) mothers and 4N (Col) fathers that were either wild-

type or mutant for one of the RdDM pathway components (Figure 1). Mature seeds

were scored as normal, aborted, or abnormal and multiple independent crosses were

analyzed. Typically, seed abortion in each set of crosses varied by between 20 and

30%, regardless of whether parents were wild-type or mutant (Figure 1A). For example,

in wild-type paternal excess crosses (2N Col x 4N Col) seed abortion ranged from

57.8%-100%, with mean seed abortion at 81.1%. Therefore, we considered only those

mutants capable of substantially enhancing seed viability across multiple crosses as

being true repressors of seed abortion.

All paternally inherited RdDM pathway mutations, except *shh1*, substantially

repressed seed abortion and promoted the production of fully developed seeds capable

of germination (Figure 1, Supplemental Figure 1). SHH1 is required for RNA Pol IV

recruitment at a subset of its target sites (Law et al., 2013), suggesting that SHH1-

independent activity of RNA Pol IV is important for seed abortion. Mutations in *nrpe1*

(encoding the largest subunit of RNA Pol V) and *drm2* mirrored *nrpd1* in their

substantial repression of seed abortion; mean seed abortion among examined seeds

was 18.9% for *nrpd1,* 21.4% for *nrpe1* and 11.3% for *drm2* (Figure 1A, Supplemental

Figure 1: Loss of paternal RdDM genes but not *RDR6* represses seed abortion in paternal excess crosses.
a) Each circle represents seed abortion rate in one cross and represents multiple siliques from a single infloroscence.
b) Each circle represents the percent of seeds that failed to germinate from each scored collection of seed. Failure to germinate was defined as the inability to produce either a radicle or a hypocotyl. Bars show median and interquartile range. * at bottom represents statistically significant difference (p<0.05) in comparisons between indicated cross and cross between wild-type (WT) diploid (2N) Col-0 mothers and wild-type tetraploid (4N) Col-0 fathers. * at top represents statistically significant differences (p<0.05) between crosses indicated. Statistical significance calculated by Wilcoxon test.

Supplemental Figure 1: Loss of paternal RdDM genes but not *RDR6* and *CMT3* represses seed abortion in paternal excess crosses. a,b,f,g) Histograms depicting percentage of seed collected that was either aborted, abnormal or normal. **c)** Histograms depicting the germination status of collected seed. Seeds grouped under "normal" produced both true leaves and roots. Abnormally germinating seeds germinated but failed to produce a radicle or had abnormal cotyledons. Seeds that failed to produce either a radicle or a hypocytl are grouped under "did not germinate". **d)** Loss of CMT3 does not repress paternal excess lethality. Wilcox test did not show statistically significant differences between crosses in which the tetraploid father is either wild-type or *cmt3*. **e)** Interploidy crosses with L*er* mothers are similar to crosses with Col-0 mothers. Wilcox test showed statistically significant differences between crosses. For **d** and **e**, circles represent a single cross and includes multiple siliques from a single inflorescence. Bars represent median and interquartile range. * at bottom represents statistically significant difference in seed abortion or germination (p<0.05) in comparisons between indicated cross and cross between wild-type diploid mothers and wild-type tetraploid fathers. Statistical significance calculated by Chi-Squared test. N represents number of seeds/seedlings counted in each cross. WT = Col-0. This supplement supports Figure 1.

Figure 1). *rdr2* and *dcl3* mutations repressed seed abortion to a lesser extent (Figure 1,

Supplemental Figure 1). The previously described redundancy between *DCL3* and

other *DCL* paralogs (Gasciolli et al., 2005; Stroud et al., 2013) likely explains the lower

repression of seed abortion by loss of *DCL3*, but it remains unclear as to why loss of

RDR2 does not repress seed abortion to the same extent as that observed upon loss of

NRPD1.

For all mutants the seed abortion ratios were reflected in the percentage of seeds

that germinated on plates (Figure 1B, Supplemental Figure 1). Tetraploid *drm2* and

nrpd1 mutants also suppressed interploidy seed abortion when crossed with wild-type

diploid L*er* mothers (Supplemental Figure 1E,G) (Erdmann et al., 2017), indicating the

effect was not specific to Col mothers. RNA Pol IV has also been proposed to function

in partnership with RDR6 in a non-canonical RdDM pathway (Martinez et al., 2018). We

therefore tested if tetraploid fathers mutant for *rdr6* could suppress paternal excess

seed abortion. A paternally inherited *rdr6* mutation did not repress seed abortion and did

not act additively with *rdr2* (Figure 1A, Supplemental Figure 1B). We also tested

whether mutations in the CHG methyltransferase *CMT3* could suppress interploidy seed

abortion when inherited through the male – no effect was observed (Supplemental

Figure 1D,F).

In contrast to repression of seed abortion by paternally inherited loss-of-function

mutations in RdDM pathway genes, most loss-of-function mutations inherited through

the diploid mother did not repress seed abortion in paternal excess crosses with wild-

type tetraploid fathers (Figure 1, Supplemental Figure 1A,C). An exception was

maternal loss of *DRM2*, which resulted in a statistically significant repression in seed

abortion, although the magnitude of the repression was small (mean percentage of

aborted seed was 66.9% for *drm2* compared to 81.1% for WT) and did not phenocopy

the extensive reduction in seed abortion observed upon the loss of paternal RdDM

pathway components (Figure 1, Supplemental Figure 1A,C). It was previously reported

that suppression of paternal excess seed abortion also occurs when both parents are

mutant for *NRPD1* (Erdmann et al., 2017; Martinez et al., 2018). We observed the same

effect for all other genes tested: *rdr2*, *dcl3*, *nrpe1*, and *drm2* (Figure 1, Supplemental

Figure 1A,C). In sum, these genetic results indicate 1) that paternal loss of the

canonical RdDM pathway is sufficient to suppress seed abortion caused by paternal

genomic excess and 2) that a genetically complete canonical RdDM pathway is not

required in endosperm itself for suppression of seed abortion.

230 Massive, shared gene mis-regulation in lethal and viable paternal excess

231 endosperm

To determine what genes or processes were associated with interploidy seed

lethality and its genetic suppression, we examined the transcriptional effects of doubling

paternal dosage in the endosperm. We performed mRNA-seq to high depth on

developing endosperm from three biological replicates of balanced crosses (L*er* x Col),

lethal paternal excess (L*er* x 4N Col), and viable paternal excess (L*er* x 4N Col *nrpd1*)

(Figure 2, Supplemental Table 1). Surprisingly, comparisons of the three transcriptomes

by principal component analysis indicated that lethal and viable paternal excess

endosperm were more similar to each other than to balanced endosperm (Figure 2A).

About a third of the transcriptome was significantly mis-expressed (2-fold or greater

difference at q < 0.05) in lethal paternal-excess endosperm compared to balanced

endosperm: 4,054 genes were more highly expressed and 3,855 genes decreased in

expression (Figure 2B, Supplemental Dataset 1). GO analyses (Supplemental Dataset

2) showed that genes with lower transcript abundance in lethal paternal excess relative

to balanced endosperm were enriched for those encoding light harvesting proteins,

proteins in glucose and starch metabolism, hormone responses, response to abiotic

stimuli, and cell wall organization. Whereas the predicted consequences of many of

these changes remain unclear, the decreased expression of cell wall genes is

consistent with the failure in endosperm cellularization that has been previously

described in lethal paternal excess crosses (Wolff et al., 2015). GO analyses of genes

with increased expression identified enrichment for those encoding proteins involved in

protein deneddylation, ribosome biogenesis, DNA replication, chromosome segregation,

the cell-cycle. The increased expression of these genes is consistent with increased cell

proliferation observed in paternal excess endosperm (Scott et al., 1998; Tiwari et al.,

2010). Compared to balanced endosperm, viable paternal excess endosperm also

Figure 2: Lethal and viable paternal excess endosperm are transcriptionally more similar to each other than to balanced endosperm. a) PCA plot of read counts for genes from biological replicate mRNA-Seq samples.
b) Plot of number of genes differentially expressed in comparisons of balanced endosperm with both lethal (purple) and viable (yellow) ♂ excess endosperm. Only 614 genes were differentially expressed between viable and lethal ♂ excess endosperm (gray). **c)** Correction value in viable paternal excess endosperm for each gene that was called as being significantly differentially expressed in comparisons of balanced and lethal ♂ excess endosperm. The value was calculated as % Correction= 100-(((log$_2$ (Viable/Balanced)/log$_2$ (Lethal/Balanced))*100).
A value of 100% indicates that the gene, which was mis-regulated in lethal ♂ excess, was not differentially expressed in viable ♂ excess relative to balanced endosperm. A value of 0% represents similar mis-regulation in both lethal and viable ♂ excess relative to balanced endosperm. Fold change values and significance for fold-change for **b** and **c** were calculated using CUFFDIFF. **d)** Lethal paternal excess endosperm was enriched for chalazal endosperm gene expression; viable paternal excess endosperm showed both chalazal and peripheral markers. Tissue enrichment for each biological replicate is shown.

Supplemental Table 1: Details of sequencing libraires

RNA-Seq

Genotype	Mapped reads	Mapping efficiency
Lethal paternal excess replicate 1	123133930	91.60%
Lethal paternal excess replicate 2	116724380	92.30%
Lethal paternal excess replicate 3	105331054	89.30%
Viable paternal excess replicate 1	88644757	92%
Viable paternal excess replicate 2	99177965	92.70%
Viable paternal excess replicate 3	93705685	81.70%

small RNA sequencing

Genotype	Reads remaining after processing	Mapped reads	Mapping efficiency
Lethal paternal excess replicate 1	4636843	3777633	81.47%
Lethal paternal excess replicate 2	15358892	13311163	86.67%
Viable paternal excess replicate 1	19840772	17098180	86.18%
Viable paternal excess replicate 2	20129312	17518163	87.03%

Methylome

Genotype	Mapped read pairs (unique only)	Mapping efficiency*	Reads remaining after deduplication by bismark
Balanced replicate 1	9495700	41.70%	7096231
Balanced replicate 2	14406400	44%	10176085
Lethal paternal excess replicate 1	10075611	49.60%	6850969
Lethal paternal excess replicate 2	14320244	49.50%	9505864
Viable paternal excess replicate 1	20027548	46.80%	13152774
Viable paternal excess replicate 2	12051462	44.30%	8589294

Note that biological replicates for the methylomes were merged for analysis

* Mapping efficiency refers to % of uniquely mapping paired end reads

Supplemental Figure 2: Analysis of gene expression differences in ♂ excess endosperm derived using the *osd1* mutation (data from Martinez et al., 2018). a) PCA plot shows that the transcriptomes of L*er* x Col *osd1* (lethal paternal excess) and L*er* x Col *osd1 nrpd1* (viable paternal excess) are more similar to each other than to L*er* x Col (balanced).
b) Frequency distribution of all gene expression changes between balanced, lethal ♂ excess and viable ♂ excess. Fold change was calculated using Cuffdiff. All genes whose output FC was greater than 1 or less than -1 shown.
c) Correction value was calculated for genes that are statistically significantly different in comparisons of balanced and lethal ♂ excess endosperm. The value, calculated as % Correction= 100-(($\log_2$ (viable/balanced))/($\log_2$ (lethal/balanced))*100). Value of 100% indicates that a gene misregulated in lethal ♂ excess is not misregulated in viable ♂ excess. 0% represents similar misregulation in both lethal and viable ♂ excess. Fold change values and significance were calculated using Cuffdiff. Boxplot here is a Tukey plot. d) Both L*er* x Col *osd1* and L*er* x Col *osd1 nrpd1* endosperm are enriched for chalazal endosperm marker gene expression. Analysis was performed using the tissue enrichment test (Schon and Nodine, 2017) on gene expression measured by the HT-Seq count command. Tissue enrichment per biological replicate shown. This supplement supports Figure 2.

showed a similar quantitative change in gene expression: 3150 genes increased in

expression and 2845 decreased (Figure 2B, Supplemental Dataset 1). Many of same

genes were mis-regulated in viable and lethal paternal excess endosperm compared to

balanced endosperm. Compared to lethal paternal excess, only 188 genes were

reduced in viable paternal excess endosperm and 426 genes were more highly

expressed (Figure 2B, Supplemental Dataset 1). To further test if the viability of paternal

excess seeds was predicated on the transcriptome coming nearer to the dosage of

balanced endosperm, we examined the extent to which gene expression was

"corrected" in viable paternal excess endosperm (Figure 2C). Most genes with

increased expression in lethal paternal excess endosperm were not corrected, but a

small subset of genes with decreased expression in lethal paternal excess were

moderately corrected in viable paternal excess (Figure 2C).

We analyzed the transcriptomic data using a tissue enrichment tool to assess if

the abundances of markers enriched in chalazal, peripheral, and micropylar endosperm

were altered in paternal excess endosperm (Schon and Nodine, 2017). Paternal excess

endosperm adopted a gene expression program characteristic of chalazal endosperm

(Figure 2D), consistent with phenotypic analysis of lethal paternal excess seeds

(Martinez et al., 2018; Scott et al., 1998; Wolff et al., 2015). Viable paternal excess

endosperm transcriptomes also showed increased chalazal endosperm marker gene

expression, but there was also slightly elevated peripheral endosperm marker gene

expression relative to lethal paternal excess endosperm. These results suggest that

lethal and viable paternal excess endosperm differ similarly from balanced endosperm

in gene expression and subsequent developmental programs.

We also reanalyzed published endosperm transcriptome data from lethal and

viable paternal excess endosperm generated using *osd1* and *osd1 nrpd1* mutations

(Martinez et al., 2018) (Supplemental Figure 2). Consistent with our data, we found that

both L*er* x Col *osd1* (lethal paternal excess) and L*er* x Col *osd1 nrpd1* (viable paternal

excess) showed elevated expression of chalazal markers (Supplemental Figure 2D).

Endosperm from L*er* x Col *osd1 nrpd1* also showed extensive genic mis-regulation, but

the percentage of the transcriptome that was mis-regulated was lower than in our viable

paternal excess endosperm datasets (Figure 2B, Supplemental Figure 2B). Additionally,

the extent of gene expression correction in viable paternal excess endosperm was also

lower in our datasets compared to theirs (Figure 2C, Supplemental Figure 2C). These

discrepancies could stem from multiple differences between our experiments. Our

endosperm data also had more replicates, higher mappable read depth, and lower

levels of seed coat contamination (Supplemental Figure 2, Supplemental Table 1,

(Martinez et al., 2018)). We created paternal excess endosperm using tetraploid fathers

while Martinez et al. created paternal excess via the use of the *osd1* mutation, which

causes omission of the second meiotic division and thus generates diploid pollen. There

might be biological differences, as yet unclear, between diploid sperm produced from a

diploid parent compared to diploid sperm produced from a tetraploid parent. Also, the

osd1 mutation was backcrossed into Col from another accession (d'Erfurth et al., 2009;

Martinez et al., 2018) and interploidy seed abortion is sensitive to the genetic

background of both parents (Lu et al., 2012; Piskurewicz et al., 2016; Scott et al., 1998).

Mis-regulation of imprinted genes characterizes both lethal and viable paternal

excess endosperm

Imprinted gene mis-regulation in the endosperm has been suggested as a culprit

for the developmental catastrophe of paternal excess crosses (Gutierrez-Marcos et al.,

2003; Haig and Westboy, 1991). We found that imprinted genes are disproportionately

more likely to show increased expression than all genes in the genome in conditions of

lethal paternal excess (N-1 chi-square test, *p= 3x10^{-4} for MEGs and p<10^{-4} for PEGs)

(Figure 3). Of 43 previously identified Col-L*er* PEGs and 130 Col-L*er* MEGs (Pignatta et

al., 2014), 25 PEGs (58%) and 27 MEGs (19%) displayed at least a two-fold increase in

transcript abundance in lethal paternal excess endosperm (Figure 3A). The abundance

of two PEGs and 18 MEGs decreased. The vast majority of these imprinted genes

remained mis-regulated in viable paternal excess endosperm (Figure 3B,C). Only two

MEGs were differentially expressed in comparisons of viable and lethal paternal excess

endosperm – *JLO* transcript abundance increased 2.6-fold and *GSR1* transcript

abundance decreased 2.4-fold (Figure 3C, Supplemental Dataset 1). Similarly, only one

PEG, AT4G20800, had lower transcript abundance in viable paternal excess

Figure 3: **Imprinted genes and transposons are mis-regulated in both lethal and viable ♂ excess endosperm. a-c)** Expresson of Col-L*er* imprinted genes in endosperm. FPKM is normalized expression; statistical significance of difference in abundance calculated by Cuffdiff. q<0.05 represented by black circles. q>0.05 represented by gray circles. **d-f)** Expression from transposable elements is elevated in lethal and viable ♂ excess endosperm. RNA-Seq reads were mapped to consensus sequences from Repbase. Black circles represent TEs with significant differences in transcript abundance according to DEGseq. Gray circles represent TEs without significant differences in transcript abundance.

Supplemental Figure 3: Suppression of paternal excess lethality by loss of *NRPD1* does not require normalization of genes implicated in interploid seed abortion. Expression level of genes shown to repress seed abortion in *osd1*-based paternal excess crosses is not repressed by loss of *NRPD1*. Fold change and statistical significance (represented by *) calculated by Cuffdiff. This supplement supports Figure 3.

endosperm compared to lethal paternal excess endosperm (Figure 3C, Supplemental

Dataset 1). These results suggest that mis-regulation of multiple imprinted genes in

endosperm is unlikely to be the cause of endosperm dysfunction and seed lethality in

interploidy crosses.

It was previously demonstrated that mutations in a subset of PEGs or their

interactors can suppress paternal excess seed abortion when inherited paternally (Wolff

et al., 2015; Huang et al., 2017; Kradolfer et al., 2013; Jiang et al., 2017). We found that

whereas several of these genes indeed have increased transcript abundance in lethal

paternal excess relative to balanced endosperm, their expression remains altered in

viable paternal excess (Supplemental Figure 3). This observation suggests that seed

viability brought about by loss of paternal *nrpd1* is independent of gene expression

normalization in endosperm of genes whose paternal loss also represses seed abortion.

In summary, these gene expression data indicate that the expression of only a

small number of genes distinguishes lethal paternal excess endosperm from viable

paternal excess endosperm at this stage of development and suggest that correction of

gene expression to levels observed in balanced endosperm, even for imprinted genes,

is not necessary for paternal excess seed viability.

A subset of TEs are mis-regulated in lethal and viable paternal excess seeds

Increased expression of transposable elements (TEs) in paternal excess

endosperm has been speculated as a cause of seed abortion (Castillo and Moyle, 2012;

Martienssen, 2010). We compared TE expression levels in endosperm by mapping

mRNA-seq reads to 375 consensus TE sequences from REPBASE (Figure 3D-F) (Bao

et al., 2015). In lethal paternal excess endosperm, 73 families showed a statistically

significant change in transcript abundance and 40 increased by 5-fold or more, including

the transpositionally active family *ONSEN/ATCOPIA78* (Figure 3D, Supplemental

Dataset 3). The TE families *ATLINE1-10A* and *ATGP7* exhibited the largest increases,

at least 52-fold (Figure 3D). Additionally, the transcript abundance of 26 TE families

decreased in paternal excess crosses. *TAG1*, a transpositionally active TE that is

present only in the maternally inherited L*er* genome (Tsay et al., 1993), was the most

repressed, nearly 56-fold (Figure 3D). Like for genes, viable paternal excess

endosperm displayed a TE expression profile similar to lethal paternal excess

endosperm (Figure 3E,F, Supplemental Dataset 3). Only 14 TE families exhibited

decreased transcript abundance in viable paternal excess endosperm relative to lethal

endosperm and 19 TE families exhibited increased transcript abundance (Figure 3F,

Supplemental Dataset 3). The observation that lethal and viable paternal excess

endosperm exhibit TE mis-regulation to a similar extent argues that TE mis-regulation

by itself is unlikely to be the cause of seed abortion induced by paternal genomic

excess.

The RdDM pathway is attenuated in paternal excess endosperm

A number of proteins that establish or maintain CG, CHG and CHH methylation

were differentially expressed among balanced, lethal, and viable paternal excess

endosperm (Figure 4, Supplemental Dataset 1). The differential expression of genes

encoding members of the RdDM pathway is particularly noteworthy. With the exception

of *NRPD1*, the expression of genes encoding subunits of RNA Pol IV and V, along with

RDR2, *AGO4*, and *DRM2*, was significantly decreased in lethal paternal excess

endosperm (Figure 4A, Supplemental Dataset 1). Consistent with these findings, the

expression of the 5-methylcytosine DNA glycosylase *ROS1*, whose expression is

directly promoted by RdDM (Williams et al., 2015), was also reduced (Figure 4A). The

expression of most of these genes increased slightly in viable paternal excess

endosperm compared to lethal endosperm, but only increases in *RDR2* and *ROS1* were

significant (Figure 4A).

We tested if sRNA production was impacted in paternal excess endosperm. We

sequenced endosperm small RNAs from two replicates of lethal and viable paternal

excess endosperm (Supplemental Table 1) and compared them to previously published

sRNA data from balanced endosperm (Erdmann et al., 2017). Previous evaluations of

small RNAs in paternal excess crosses have been performed on whole seeds, not

endosperm, but noted reduced 24 nt sRNAs in lethal paternal excess seeds (Lu et al.,

2012; Martinez et al., 2018). We assessed the overall functionality of small RNA

production in the endosperm by examining small RNA size profiles. In lethal paternal

excess endosperm, 24 nt sRNAs represented the most abundant size class, but their

Figure 4: Canonical RdDM pathway function in endosperm is affected by paternal excess. a) Genes encoding RdDM components are down-regulated in lethal and viable ♂ excess endosperm. *ROS1* expression is a read-out of RdDM activity and reflects differential activity of RdDM. * represents statistically significantly different gene expression. **b)** Small RNA production is impacted in paternal excess endosperm. Size profiles of sRNA reads mapped to TAIR10 genome for three replicates of balanced endosperm and two replicates each of lethal and viable paternal excess endosperm. **c)** CHH methylation losses in paternal excess endosperm. Venn diagram shows intersections of CHH DMRs obtained from comparisons of balanced, lethal, and viable paternal excess endosperm. Lethal and viable paternal excess endosperm share a significant proportion of regions that are hypomethylated relative to balanced endosperm. A smaller subset of DMRs lose more methylation in lethal relative to viable. **d)** Loss of CHH methylation is associated with loss of 24 nt sRNAs. Venn diagrams show the relationship between changes in sRNA abundance and CHH methylation levels. A subset of CHH DMRs are associated with loss of sRNAs. A smaller subset is associated with gains in 24 nt sRNAs. Significance of overlaps was calculated using the Fisher test option from Bedtools.

Supplemental Figure 4. Small RNA in balanced, lethal and viable paternal excess endosperm. a) Genic sRNAs are reduced in lethal paternal excess. Size profiles of sRNA reads overlapping genes for three replicates of balanced endosperm, two replicates of lethal paternal excess and two replicates of viable paternal excess endosperm. Sum of all reads overlapping genes was used as denominator to determine proportion of small RNA at each size. **b)** TE insertion sRNAs are reduced in lethal paternal excess. Analysis as in a). Frequency distribution of significant changes in sRNA abundance at **c)** genes and **d)** transposable element insertions. Fold-change and significance calculated using DESeq2. Read counts for TAIR10 TEs generated using bedtools. This supplement supports Figure 4.

proportion was reduced compared to balanced endosperm (Figure 4B). In viable

paternal excess endosperm, the proportion of 24 nt sRNAs was comparable to that of

balanced endosperm (Figure 4B). Genic and TE-associated sRNAs in lethal and viable

paternal excess showed a pattern similar to that of bulk sRNAs (Supplemental Figure

4A,B). These observations suggest that the capacity to produce 24 nt sRNAs is

relatively normal in viable paternal excess endosperm, presumably because of restored

RDR2 expression (Figure 4A,B). We also note that the proportion of 21 nt sRNAs

increased in both lethal and viable paternal excess endosperm (Figure 4B,

Supplemental Figure 4). While the pathways involved in the biogenesis of these 21 nt

sRNAs remain unclear, they could represent the activity of post-transcriptional gene

silencing (mRNA cleavage) pathways on transcripts that are mis-regulated in both lethal

and viable paternal excess endosperm (Borges and Martienssen, 2015; Marí-Ordóñez

et al., 2013)

A hallmark of the activity of the 24 nt sRNA pathway is CHH methylation. We

performed whole genome bisulfite sequencing to profile the methylomes of balanced,

lethal, and viable paternal excess endosperm (Supplemental Table 1). To identify

regions with altered CHH methylation, we divided the genome into 300 bp windows with

200 bp overlaps and identified those windows with a 10% or greater CHH methylation

difference between genotypes (i.e. 20% vs 30% methylation.) Windows with significant

differences were merged into differentially methylated regions. CHH methylation was

slightly higher in viable paternal excess endosperm relative to lethal paternal excess

endosperm, but in both genotypes it was drastically reduced relative to balanced

endosperm on both maternal and paternal alleles (Figure 4C, Supplemental Datasets 4

& 5). Regions with reduced CHH methylation significantly overlapped with regions with

lowered sRNA levels in lethal paternal excess endosperm (Figure 4D).

Paternal excess differentially affects the expression of maternally and paternally

inherited alleles

Total gene expression levels are derived from the contribution of maternally and

paternally inherited alleles. Based on the effect of *nrpd1* mutations on allele-specific

expression in balanced endosperm, we previously proposed that 4N *nrpd1* might

suppress paternal excess lethality by increasing transcriptional dosage from maternal

alleles for many genes (Erdmann et al., 2017). To test this hypothesis, we evaluated

allele-specific expression in balanced, lethal, and viable paternal excess endosperm for

genes with at least 50 allele-specific reads in both balanced and paternal excess

crosses (Figure 5) (Supplemental Dataset 9). Balanced endosperm has a 2:1 m:p ratio

and paternal excess endosperm has a 2:2 ratio of maternal:paternal genomes. Mirroring

the genomic ratios, the median paternal fraction for genes was 33.8% in balanced

endosperm, 50.1% in lethal paternal excess endosperm, and 50.6% in viable paternal

excess (Figure 5A). To assess potential allelic mis-regulation of individual genes, we

normalized transcriptomic fractions to genomic fractions by calculating the deviation

from paternal genomic contributions for each gene (% paternal-33% for balanced

crosses and % paternal-50% for paternal excess). Contrary to our expectation, there

was a small but detectable *decrease* in paternal contribution relative to genomic

expectation for specific genes in lethal paternal excess endosperm (*two-sided*

D'Agostino's K^2 test, skew = -0.58, *p< 2.2e-16* ; Figure 5B, purple line, Supplemental

Dataset 9). A similar bias was observed in viable paternal excess endosperm (*two-sided*

D'Agostino's K^2 test, *skew = -0.53, p <2.2e-16;* Figure 5B, yellow line, Supplemental

Dataset 9*)*. Thus, in viable relative to lethal paternal excess endosperm there were few

changes in overall allelic bias (*two-sided D'Agostino's K^2* test, *skew = 0.0443*, *p =*

0.05887; Figure 5B, gray line, Supplemental Dataset 9).

To explore the contribution of allelic mis-regulation to genic mis-regulation, we

examined shifts in paternal deviation from genomic expectation for genes with

increased or decreased expression in lethal paternal excess endosperm compared to

balanced endosperm (Figure 5C-D, Supplemental Figure 5A). Genes with decreased

expression were more heavily influenced by loss of paternal allele contributions (*two-*

sided D'Agostino's K^2 test, skew for down-regulated genes =-0.4506, p-value= 1.243e-

16; Figure 5C*)*. 12.1% of genes with sufficient allele-specific reads showed at least 20%

decrease in paternal bias, indicating increased repression of paternal alleles, whereas

2.3% showed at least a 20% increase in paternal bias, indicating increased repression

of maternal alleles (Figure 5C). Genes with increased expression in lethal paternal

excess endosperm were more impacted by increases in maternal allele contributions

a

b

Lethal ♂ excess - Balanced
Viable ♂ excess - Balanced
Lethal - Viable ♂ excess

c

Genes with decreased expression in lethal paternal excess

232
Repressed ♂
alleles

44
Repressed ♀
alleles

% Paternal deviation from
genomic expectation

WOX8

Decreased
♀ allele

AT2G01580

Decreased
♂ allele

d

Genes with increased expression in lethal paternal excess

119
Activated ♀
alleles

66
Activated ♂
alleles

% Paternal deviation from
genomic expectation

SDG21

Increased
♀ allele

HEXO3

Increased
♂ allele

e

All sRNAs

Genes

TEs

Balanced
Viable ♂ excess
Lethal ♂ excess

Figure 5: Allelic contributions to expression differences between balanced and lethal paternal excess endosperm. a) Paternal fraction in genic transcripts. Boxplot represents all genes with at least 50 allele specific reads in indicated genotype. Number of genes in balanced = 10,388, lethal ♂ excess =11,709, viable ♂ excess =11,430. For balanced endosperm, a total of five replicate libraries were analyzed (Erdmann et al., 2017). **b)** Paternal excess endosperm transcriptome is maternally skewed. Frequency distribution plot of % paternal deviation for each gene. Paternal deviation was calculated for genes with at least 50 allele specific reads in each pair of genotypes being compared. Number of genes in lethal ♂ excess - balanced = 9,550, viable ♂ excess - balanced = 9,639, viable ♂ excess - lethal ♂ excess = 10,919. **c)** Decreased paternal allele contribution at a larger proportion of genes with decreased expression in lethal paternal excess compared to balanced. N = 1910 genes. Impacts of allele-specific changes on gene expression at two examples, *WOX8* and AT2G01580, are shown. **d)** Increased maternal allele activation at a larger proportion of genes with increased expression in lethal paternal excess. N= 1608 genes. Impacts of allele specific changes on gene expression at two examples, *SDG21* and *HEXO3*, are shown. Genes analyzed for **c** and **d** were detected as being significantly different in gene expression by Cuffdoff. A gene with an allelic shift of at least 20% was considered to have a modified allelic balance. Boxplots represent median values for paternal deviation from genomic expectation. For gene-specific histograms, maternal allele is red and paternal allele is blue. **e)** sRNA populations are increasingly maternally biased in paternal excess endosperm. The paternal fraction of sRNA populations from viable paternal excess endosperm is intermediate between lethal paternal excess and balanced endosperm. Data from three replicates of balanced endosperm and two replicates each of lethal and viable paternal excess endosperm are plotted. sRNA reads mapped to TAIR10 genome were first split based on size and then into Col and L*er* reads and reads using SNPs. % of paternal reads, and deviation from genomic expectation at each size was calculated.

Supplemental Figure 5: Allelic contributions to gene expression differences. a) Contribution of allelic mis-regulation to differential gene expression in lethal ♂ excess. Scatter plot of relationship between change in paternal deviation and change in gene expression between balanced and lethal paternal excess. **b)** Contribution of allelic mis-regulation to differential gene expression between lethal and viable ♂ excess. Scatter plot of relationship between change in paternal deviation and change in gene expression between balanced and lethal paternal excess. In **a** and **b**, black circles represent genes called as being significantly different by Cuffdiff; gray cricles represent genes that are not significantly different. **c)** Allele-specific changes in genes upregulated in viable relative to lethal paternal excess (N=189). **d)** Allele specific changes among genes downregulated in viable paternal excess relative to lethal paternal excess. (N=83) In **a-d**, genes with more than 50 allele specific reads were considered for analysis. In **c** and **d**, genes with at least a 20% shift in allelic balanced between viable and lethal ♂ excess were considered to have a modified allelic contribution. This supplement supports Figure 5.

Supplemental Figure 6: Allelic bias of some imprinted genes is lost in lethal paternal excess.
Changes in paternal deviation for differentially expressed MEGs and PEGs. Maximal deviation for a MEG in balanced endosperm and paternal excess endosperm is -33 and -50 respectively. Maximal deviation for PEGs in balanced endosperm and paternal excess endosperm is 67 and 50 respectively. Our data did not show imprinting in balanced endosperm at a small subset of imprinted loci previously shown to be imprinted in balanced endosperm. This supplement supports Figure 5.

(*skew for up-regulated genes= -0.52084, p-value= 6.61e-16*; Figure 5D). 7.4% showed

at least a 20% decrease in paternal deviation, indicating increased maternal allele

expression, and 4.1% showed at least a 20% increase in paternal bias, indicating

increased paternal allele expression (Figure 5D).

Comparison of allelic contributions between viable and lethal endosperm

revealed large differences at a very limited number of genes (Supplemental Figure 5B-

D). Among genes with increased transcript abundance in viable compared to lethal

paternal excess endosperm, increased paternal allele contributions were observed at 9

loci (Supplemental Figure 5B,C, Supplemental Dataset 9). Increased maternal allele

contributions were observed at another 9 loci (Supplemental Figure 5B,C, Supplemental

Dataset 9). Among genes with decreased abundance, only two genes showed strong

evidence for a repression of paternal allele expression and two genes had evidence for

repression of maternal allele expression (Supplemental Figure 5B,D).

We also tested if the increased expression of imprinted genes in paternal excess

endosperm was related to a breakdown of imprinting (Supplemental Figure 6). In

comparisons of balanced endosperm and lethal paternal excess, loss-of-imprinting was

apparent among MEGs. Among 27 MEGs with increased expression in lethal paternal

excess endosperm, paternal deviation increased for 8 genes, which became bi-

allelically expressed (Supplemental Figure 6). However, strongly maternally biased

expression was maintained at 11 MEGs (Supplemental Figure 6). Of 25 PEGs with

increased total expression, 13 had reduced paternal bias (Supplemental Figure 6).

However, most of these changes were small, with the exception of AT4G20800, which

was bi-allelically expressed (Supplemental Figure 6). These observations indicate that

loss-of-imprinting contributes to increases in imprinted gene expression (Figure 3) at

only a subset of imprinted genes.

The allele-specific changes in gene expression (Figure 5A-D) led us to test if

sRNAs change in an allele-specific manner. The sRNA pool in balanced endosperm is

overall paternally biased (Erdmann et al., 2017) (Figure 5E). For most sRNA size

classes, lethal paternal excess endosperm was more maternally-biased than expected

from the ratio of maternal and paternal genomes (Figure 5E). For example, in lethal

paternal excess, 24 nt sRNAs were 13.8% more maternal than expected given the 2:2

ratio of maternal to paternal genomes. This indicates, counterintuitively, that extra

paternal genomes are associated with decreased small RNAs from paternal genomes

(or increased sRNAs from maternal genomes). sRNAs in viable paternal excess

endosperm are slightly more paternal than lethal paternal excess endosperm, although

to a lesser extent than is observed in balanced endosperm (Figure 5E), consistent with

increased functionality of the RdDM pathway in viable paternal excess endosperm

(Figure 4A).

In summary, from these analyses we conclude that increasing paternal genome

copy number can impact the expression of only maternal or only paternal alleles for a

subset of loci. Thus, maternally and paternally inherited alleles contribute unequally to

genome-wide transcriptome mis-regulation. However, overall, we did not find evidence

for our hypothesis that a shift toward maternal allele expression is associated with

repression of paternal excess seed abortion (Erdmann et al., 2017), as *both* lethal and

viable paternal excess endosperm have a higher fraction of maternal transcripts than

expected from parental genomic ratios.

DISCUSSION

Impact on models for interploidy seed abortion

We have shown that an extra copy of the paternal genome induces regulatory

changes at both maternal and paternal alleles and drives massive changes in gene

expression in the endosperm, varying little between viable and inviable seeds (Figures

2-5). Our data allow evaluation of a number of proposals regarding the transcriptional

changes at genes, TEs, or sRNAs that could result in interploid seed lethality. Inspired

by models of transposon-induced dysgenesis in *Drosophila* leading to atrophied ovaries

(reviewed in Kelleher, 2016), it was speculated that an imbalance between sRNAs

deposited by a diploid mother and the TE load in a tetraploid father could lead to TE

expression, which would then trigger seed abortion (Castillo and Moyle, 2012;

Martienssen, 2010). Our finding that TEs are similarly mis-regulated in lethal and viable

paternal excess endosperm (Figure 3) leads us to conclude that TE expression levels

are unlikely to be a determinant of interploidy seed viability. While it is perhaps

surprising that viable seeds have high TE transcript abundance, it is not unprecedented.

Arabidopsis mutants like *met1*, which have high levels of TE expression (Oberlin et al.,

2017; Zilberman et al., 2007), also produce viable seed (Xiao et al., 2006).

Transcription of imprinted genes is potentially a strong candidate for the critical

gene dosage difference separating lethality and viability (Haig and Westboy, 1991;

Lafon-Placette and Köhler, 2015; Wolff et al., 2015). But, like TEs, the similar

expression levels of imprinted genes between lethal and viable interploid seeds (Figure

3) suggest their differential expression is not determinant for viability. We did identify

614 differentially expressed genes between lethal and viable interploidy seeds (Figure

2B,C, Supplemental Dataset 1). These 614 differentially expressed genes include

several genes encoding proteins involved in phytohormone signaling and

developmental regulators – all potentially good candidates for determining seed viability

(Supplemental Dataset 1). It is possible that the expression of multiple loci of modest

effect or partial renormalization of expression patterns of multiple loci constitute the

dosage threshold between lethal and viable seeds. Based on the present data, we are

unable to distinguish if any of the differentially expressed genes or partially corrected

genes are causal for the effect on seed viability, or whether they represent a

consequence of endosperm development associated with seed viability.

Several distinct hypotheses revolving around sRNAs have been postulated to

explain how loss of *NRPD1* in tetraploid fathers, but not diploid mothers, represses seed

abortion in paternal excess crosses (Erdmann et al., 2017; Martinez et al., 2018). The

easiRNA model for interploid seed abortion is based on two key conclusions from the

recent data of Martinez et al (Martinez et al., 2018). First, a non-canonical RNA Pol IV-

RDR6 pathway functions in pollen to make easiRNAs, which scale with paternal

genome dosage. Second, the transfer of increased easiRNAs into the endosperm from

diploid pollen inhibits RdDM; it is then the absence of these easiRNAs in *nrpd1* mutant

pollen that permits the RdDM pathway to function in paternal excess endosperm. Our

data do not support this model. We failed to obtain a significant increase in viable

paternal excess seeds with 4N *rdr6-15* pollen (Figure 1A, Supplemental Figure 1B),

suggesting that any sRNAs exclusively dependent on *RDR6* are not linked to paternal

excess seed abortion. Consistent with this result, paternally produced siRNA854, which

is dependent on *DCL2/4* and *RDR6*, is required for viability of paternal excess seeds

(McCue et al., 2012; Wang et al., 2018). By contrast, we found that multiple members of

the canonical RdDM pathway restore paternal excess seed viability (Figure 1,

Supplemental Figure 1A,C). Unlike the reported relatively low levels of viable seed

produced by mutations in non-RdDM genes such as *RDR6, DCL2, DCL4*, *AGO2* and

miRNA845 (approximately 20-30% of the seeds are viable) (Borges et al., 2018;

Martinez et al., 2018), loss of *NRPE1* and *DRM2* phenocopy *NRPD1* in the large

magnitude of repression of seed abortion (approximately 80% of the seeds are viable)

(Figure 1, Supplemental Figure 1). The cause of the discrepancy among the genetic

results between our studies remains unclear, but may be linked to the use of *osd1*

mutation to create conditions of paternal excess. Similar discrepancies have been

observed elsewhere; *osd1 pkr2* mutants did not repress paternal excess seed abortion

whereas 4N *pkr2* showed reduced seed abortion (Huang et al., 2017; Wolff et al., 2015).

Finally, the easiRNA model also suggests that the RdDM pathway's recovery in the

viable paternal excess endosperm is essential for seed viability. However, loss of both

paternal and maternal copies of *NRPD1, RDR2, DCL3, NRPE1*, and *DRM2* in paternal

excess crosses did not abolish the seed viability obtained upon loss of the paternal copy

alone (Figure 1, Supplemental Figure 1) (Erdmann et al., 2017; Martinez et al., 2018).

This indicates that the activity of a genetically complete canonical RdDM pathway in the

endosperm is not necessary for suppression of paternal excess seed abortion.

Another model posits that RNA Pol IV sRNA levels are determined by maternal

dosage and that these small RNAs normally repress *AGL*s and trigger endosperm

cellularization (Lu et al., 2012). This model argues that in paternal excess endosperm

siRNA levels are disproportionate to target transcript levels of *AGL* genes, which in turn

leads to prolonged endosperm proliferation and cellularization. This model makes a key

testable prediction – the requirement for a functional RdDM pathway in the endosperm.

However, we find that RdDM in endosperm is not essential for seed viability in

conditions of paternal excess (Figure 1, Supplemental Figure 1).

We previously showed that maintaining a 2:1 ratio of maternal to paternal allele

transcripts in balanced endosperm was dependent on *NRPD1*; in its absence the

expression of many genes shifted more maternal (Erdmann et al., 2017). Thus, we

suggested that the loss of *NRPD1* repressed paternal excess seed abortion by raising

the maternal fraction of the transcriptome and balancing excess paternal gene dosage

(Erdmann et al., 2017). A central tenet of our model was that the reduction in the RNA

Pol IV sRNA pathway in endosperm (caused by loss of paternal *NRPD1*) was essential

for seed viability. However, we find that the RdDM pathway is in fact *more* functional in

viable paternal excess endosperm than in lethal paternal excess endosperm (Figure 4A-

B, Figure 5E, Supplemental Figure 4). Additionally, the transcriptome in lethal paternal

excess is already maternally biased relative to genomic expectation. This maternal bias

is not increased in viable paternal excess endosperm, as would be predicted under our

previous model (Figure 4).

What, then, might be the mechanism by which suppression of seed lethality

occurs? Although our current data cannot address whether mutations in all canonical

RdDM genes suppress paternal excess seed lethality in the same manner, given what

is known about the RdDM pathway it is likely that differences in DNA methylation, in

either in the paternal sporophyte or the male gametophyte (or both), play a key role. It

has been shown in diploid plants that loss of *NRPD1* or *DRM2* in the father can impact

gene expression in the endosperm of balanced crosses (Erdmann et al., 2017; Vu et al.,

2013) and that *DRM2* establishes methylation patterns at a subset of genomic sites in

the male germline (Walker et al., 2018). Some studies suggest that tetraploidization

itself induces DNA methylation changes (Baubec et al., 2010; Mittelsten Scheid et al.,

2003; Yu et al., 2010; Zhang et al., 2015). In one scenario, DNA methylation changes in

tetraploid pollen could ultimately direct changes in gene expression in the endosperm,

which would not occur in endosperm fathered by RdDM mutant tetraploids. An

alternative possibility is that loss of RdDM in tetraploids could lead to an altered

methylation state not found in either wild-type diploid or tetraploid males. This ectopic

paternal methylation state could then dictate a gene expression pattern that represses

seed abortion.

Evidence for buffering mechanisms in endosperm

Our results suggest that transcriptional buffering, operational in both lethal and
viable paternal excess endosperm, is one feature that could contribute to the ability of
the seed to withstand the addition of an entire extra paternal genome. In transcriptional
buffering, expression per copy of a gene can be increased or decreased based on the
number of copies of the gene relative to that of the rest of the genome (Birchler and
Veitia, 2012; Zhang et al., 2010). We observed the signatures of buffering in paternal
excess transcriptomes (Figure 5). Decreased paternal transcript levels contributed
predominantly to genes with decreased expression in paternal excess, whereas
increases in the levels of transcripts from maternal alleles contributed predominantly to
genes with increased expression (Figure 7C-D). The outcome is that some parts of the
transcriptome are more maternal than expected from parental genomic dosage,
constituting a push back against excess paternal genomes. While the mechanistic basis
of this buffering system remains unclear, it could be linked at least in part to the
reduction in the 24 nt small RNA pathway in endosperm (Figure 4). RNA Pol IV function
in balanced endosperm has been previously shown to be required for repression of the
maternal transcriptome (Erdmann et al., 2017). The increase in the maternal fraction of
the transcriptome for specific genes in paternal excess endosperm (Figure 5B) is
therefore consistent with the reduced expression of genes encoding subunits of RNA
Pol IV and other downstream members of the RdDM pathway (Figure 4A). It is
important to note that the increase in maternal fraction can be observed in both lethal
and viable paternal excess endosperm and is thus unlinked to the repression of seed
abortion brought about by paternal loss of RdDM. Other buffering mechanisms might
also play important roles and could act through post-transcriptional pathways (Wang et
al., 2018), protein degradation pathways, or via signaling pathways that have been
shown to influence paternal excess crosses (Batista et al., 2019; Doughty et al., 2014).
One striking observation is that even wild-type paternal excess crosses produce about
20% viable seed (Figure 1, Supplemental Figure 1), suggesting that seed lethality and
viability exist on a continuum. Our data further suggest that the slim margin that
separates seed life and death is likely determined by modest changes in the
endosperm, which are ultimately instructed by paternal imprints laid down by RdDM .

Methods

Arabidopsis strains, tetraploid production, and tissue collection

Mutants used in this study were: *nrpd1a-4*, *rdr2-1*, *dcl3-1*, *nrpd1b-11*, *drm2-2*, *rdr6-15* (CS879578), *rdr2-1 rdr6-15* (CS66111), and *cmt3-11t* (CS16392) and were obtained from ABRC. Tetraploids were generated by applying 0.25% colchicine in 0.2% Silwet to the apices of 2-3 week old diploid plants. Progeny from treated plants were screened by flow cytometry to identify tetraploids. For crosses, wild-type diploid Col-0 or L*er* buds were emasculated and pollination was carried out two days later with diploid or tetraploid pollen in Col-0 strain background. To assess seed abortion, siliques were harvested after drying and seeds examined under a dissecting microscope. To assess germination rates, seeds were sterilized in 2% PPM (Plant Cell Technologies) for three days at 4°C and then plated on 0.5X MS/Phytagar media. Endosperm from approximately one hundred seeds at 7 DAP from at least three siliques were dissected away from embryo and seed coat and pooled for each replicate as previously described (Gehring et al., 2009; 2011). Replicates were collected from a different sets of crosses with different individuals as parents. For lethal paternal excess seeds, endosperm was collected only from seeds with arrested embryonic growth. For viable paternal excess seeds, endosperm was collected only from seeds where embryonic growth was progressing normally.

mRNA-Seq library preparation and transcriptome analyses

Long RNA was isolated as previously described (Erdmann et al, 2017) and mRNA-seq libraries were constructed at the Whitehead Institute Genome Technology Core using the SMARTerUltra-lowPOLYA-V4 kit. Libraries were sequenced on a 40 base, single read cycle. Sequence data were filtered for quality with " *trim_galore -q 25 --phred64 -- fastqc --length 20 --stringency 5*". Filtered reads were aligned to the TAIR10 genome using "*tophat -i 30 -I 3000 --solexa1.3-quals -p 5 -g 5 --segment-mismatches 1 -- segment-length 18 --b2-very-sensitive*" (Kim et al., 2013). Tophat version 2.1.1 with Bowtie version 2.3.4 was used. We used Cuffdiff (version 2.2.1) with default settings

and the ARAPORT11 genome annotation to calculate changes in gene expression and

their statistical significance. Genes with q-value <0.05 were considered to be

significantly different. We used a custom script (assign_to_allele.py;

https://github.com/clp90/imprinting_analysis/tree/master/helper_scripts) and SNPs

between Col-0 and L*er* to identify allele specific reads. Allele specific reads per gene

were assessed using "*htseq-count -s no –m union*" and the ARAPORT11 annotation.

To assess TE transcript levels, reads were aligned to the REPBASE consensus

sequence (Jurka et al., 2005) using "*bowtie -v 2 -m 3 --best --strata -p 5 --phred64"* and

reads mapping to each TE family were summed. *META1, ATHILA6A* and *ATGP8* were

precluded from further analyses because of mapping artifacts. Differential abundance of

TE mapping reads were calculated using Fisher's exact test (p<0.05) option with

multiple-test correction in DEGseq (Wang et al., 2010).

Tissue enrichment test

Expression per gene was measured using HT-Seq count and analyses were performed

using the seed tissue enrichment test (Schon and Nodine, 2017). The time point was

set to bent cotyledon.

Small RNA-Seq library preparation and analyses

RNA was isolated from manually dissected endosperm using the RNAqueous micro kit

(Ambion). Small RNA was obtained as previously described (Erdmann et al, 2017).

Libraries were built using the NEXTflex sRNA-seq kit v3 (Bioo Scientific). Final library

amplification was carried out for 24 cycles; resultant libraries were size selected (135-

160 bp) with a pippin prep. 40 base single read sequencing was carried out an Illumina

HiSeq 2500. Sequencing reads were trimmed for quality with "*fastq_quality_trimmer -v -*

t 20 -l 25". Reads were further filtered and adapter containing reads were retained using

"*cutadapt -a TGGAATTCTCGGGTGCCAAGG --trimmed-only --quality-base 64 -m 24 -*

M 40 --max-n 0.5". The reads from libraries produced by the NEXTflex library prep kit

include four random nucleotides at both 5' and 3' ends of the reads. Taking advantage

of these tags, we were able to remove PCR duplicates using "*prinseq-lite –fastq <infile>*

-out_format 3 –out_good <outfile> -derep 1". Reads were aligned to TAIR10 with

"*bowtie -v 1 --best -5 4 -3 4 --sam --phred64-quals or --phred33-quals"* . We used a

custom script (assign_to_allele.py;

<u>https://github.com/clp90/imprinting_analysis/tree/master/helper_scripts</u>) to identify

allele-specific reads. Regions with differential sRNA levels were identified by counting

reads in 300 bp windows with 200 bp overlaps. DESeq2 (Love et al., 2014) was used to

identify windows with differential abundance of sRNAs (adjusted *p-value* less than or

equal to 0.05 and log$_2$ fold change =+/-1). Overlapping windows with significant

increases or decreases in sRNA were merged to obtain regions with differences in

sRNA. To identify genes with differences in 21 and 24 nt sRNAs, mapped reads were

separated based on size. 21 and 24 nt sRNAs mapping to genes were counted using

the ARAPORT11 annotation and "*htseq-count -s no –m union*". To identify TE insertions

with differences in 21 and 24 nt sRNAs, we first identified TE insertions that did not

overlap with genes and used "*coverageBed -counts*" command from Bedtools suite

(Quinlan and Hall, 2010) to count reads mapping to TE insertions. We identified genes

and TEs with differential abundance in 21 and 24 nt sRNA using DESeq2.

Methylome library preparation and analyses

Genomic DNA was isolated from manually dissected endosperm using the QiaAMP

DNA microkit (QIAGEN 56304); dissected tissue was incubated in ATL buffer and

proteinase K at 56°C overnight on a shaking incubator. 80-100 ng of DNA obtained from

these protocols was subjected to bisulfite treatment using Methylcode Bisulfite

Conversion kit (Invitrogen). Bisulfite libraries were constructed from these materials

using Illumina's Truseq DNA methylation kit. Bisulfite libraries were sequenced on

Illumina's HISeq 2500 in a paired end configuration. Reads were filtered for quality with

"*trim_galore --phred64 --fastqc --stringency 5 --length 15 --paired --clip_R1 2 --clip_R2*

2". The reads were then aligned to TAIR10 using *"bismark -N 1 -L 20"*. Duplicate reads

were removed using Bismark version 0.19. Bismark methylation extractor and custom

scripts previously described (Pignatta et al., 2014, 2015) were used to obtain per base

methylation. Briefly, differences in methylation were calculated genome-wide for 300 bp

sliding windows with 200 bp overlaps. To be included, windows had at least 3

overlapping cytosines in both genotypes with a read depth of at least 6 reads per

cytosine. Windows called as significantly different were at least 10% different between

tested genotypes for CHH methylation, 20% for CHG methylation and 30% for CG

methylation. Significance of difference was calculated by F.E.T with a Bonferroni-

Hochberg correction (p<0.01). Overlapping windows with significant differences in DNA

methylation were merged to obtained differentially methylated regions.

Data Access

712 Whole genome bisulfite sequencing data, small RNA sequencing and mRNA

713 sequencing data are deposited in NCBI GEO under accession GSE126932.

Figure Legends

Figure 1: **Loss of paternal RdDM genes but not *RDR6* represses seed abortion in**

paternal excess crosses. a) Each circle represents seed abortion rate in one

cross and represents multiple siliques from a single inflorescence. **b)** Each circle

represents the percent of seeds that failed to germinate from each scored collection of

seed. Failure to germinate was defined as the inability to produce either a radicle or a

hypocotyl. Bars show median and interquartile range. * at bottom represents statistically

significant difference (p<0.05) in comparisons between indicated cross and cross

between wild-type (WT) diploid (2N) Col-0 mothers and wild-type tetraploid (4N) Col-0

fathers. * at top represents statistically significant differences (p<0.05) between crosses

indicated. Statistical significance calculated by Wilcoxon test.

Figure 2: Lethal and viable paternal excess endosperm are transcriptionally more

similar to each other than to balanced endosperm. a) PCA plot of read counts for

genes from biological replicate mRNA-Seq samples. **b)** Plot of number of genes

differentially expressed in comparisons of balanced endosperm with both lethal (purple)

and viable (yellow) ♂ excess endosperm. Only 614 genes were differentially expressed

between viable and lethal ♂ excess endosperm (gray). **c)** Correction value in viable

paternal excess endosperm for each gene that was called as being significantly

differentially expressed in comparisons of balanced and lethal ♂ excess endosperm.

The value was calculated as % Correction= 100-((($\log_2$ (viable/balanced)/$\log_2$

(lethal/balanced))*100). A value of 100% indicates that the gene, which was mis-

regulated in lethal ♂ excess, was not differentially expressed in viable ♂ excess relative

to balanced endosperm. A value of 0% represents similar mis-regulation in both lethal

and viable ♂ excess relative to balanced endosperm. Fold change values and

significance for fold-change for b and c were calculated using Cuffdiff. Boxplot is a

Tukey plot. **d)** Lethal paternal excess endosperm was enriched for chalazal endosperm

gene expression; viable paternal excess endosperm showed both chalazal and

peripheral markers. Tissue enrichment for each biological replicate is shown.

Figure 3: Imprinted genes and transposons are mis-regulated in both lethal and

viable ♂ excess endosperm. a-c) Expression of Col-L*er* imprinted genes in

endosperm. FPKM is normalized expression; statistical significance of difference in

abundance calculated by Cuffdiff. q<0.05 represented by black circles. q>0.05

represented by gray circles. d-f) Expression from transposable elements is elevated in

lethal and viable ♂ excess endosperm. RNA-Seq reads were mapped to consensus

sequences from Repbase. Black circles represent TEs with significant differences in

transcript abundances according to DEGseq. Gray circles represent TEs without

significant differences in transcript abundances.

Figure 4: Canonical RdDM pathway function in endosperm is affected by paternal

excess. a) Genes encoding RdDM components are down-regulated in lethal and viable

♂ excess endosperm. ROS1 expression is a read-out of RdDM activity and reflects

differential activity of RdDM. * represents statistically significantly different gene

expression. **b)** Small RNA production is impacted in paternal excess endosperm. Size

profiles of sRNA reads mapped to TAIR10 genome for three replicates of balanced

endosperm and two replicates each of lethal and viable paternal excess endosperm. **c)**

CHH methylation losses in paternal excess endosperm. Venn diagram shows

intersections of CHH DMRs obtained from comparisons of balanced, lethal, and viable

paternal excess endosperm. Lethal and viable paternal excess endosperm share a

significant proportion of regions that are hypomethylated relative to balanced

endosperm. A smaller subset of DMRs lose more methylation in lethal relative to viable.

d) Loss of CHH methylation is associated with loss of 24 nt sRNAs. Venn diagrams

show the relationship between changes in sRNA abundance and CHH methylation

levels. A subset of CHH DMRs are associated with loss of sRNAs. A smaller subset is

associated with gains in 24 nt sRNAs. Significance of overlaps was calculated using the

Fisher test option from Bedtools.

Figure 5: Allelic contributions to expression differences between balanced and

lethal paternal excess endosperm. a) Paternal fraction in genic transcripts. Boxplot

represents all genes with at least 50 allele specific reads in indicated genotype. Number

of genes in balanced = 10,388, lethal ♂ excess =11,709, viable ♂ excess =11,430. For

balanced endosperm, a total of five replicate libraries were analyzed (Erdmann et al.,

2017). **b)** Paternal excess endosperm transcriptome is maternally skewed. Frequency

distribution plot of % paternal deviation for each gene. Paternal deviation was calculated

for genes with at least 50 allele specific reads in each pair of genotypes being

compared. Number of genes in lethal ♂ excess - balanced = 9,550, viable ♂ excess -

balanced = 9,639, viable ♂ excess - lethal ♂ excess = 10,919. **c)** Decreased paternal

allele contribution at a larger proportion of genes with decreased expression in lethal

paternal excess compared to balanced. N = 1910 genes. Impacts of allele-specific

changes on gene expression at two examples, *WOX8* and *AT2G01580*, are shown. **d)**

Increased maternal allele activation at a larger proportion of genes with increased

expression in lethal paternal excess. N= 1608 genes. Impacts of allele specific changes

on gene expression at two examples, *SDG21* and *HEXO3*, are shown. Genes analyzed

for **c** and **d** were detected as being significantly different in gene expression by Cuffdiff.

A gene with an allelic shift of at least 20% was considered to have a modified allelic

balance. Boxplots represent median values for paternal deviation from genomic

expectation. For gene-specific histograms, maternal allele is red and paternal allele is

blue. **e)** sRNA populations are increasingly maternally biased in paternal excess

endosperm. The paternal fraction of sRNA populations from viable paternal excess

endosperm is intermediate between lethal paternal excess and balanced endosperm.

Data from three replicates of balanced endosperm and two replicates each of lethal and

viable paternal excess endosperm are plotted here. sRNA reads mapped to TAIR10

genome were first split based on size and then into Col and L*er* reads and reads using

SNPs. % of paternal reads, and deviation from genomic expectation at each size was

calculated.

Acknowledgements

We thank Dr. Daniela Pignatta for deriving the 4N Col-0 line. This research was supported by NSF MCB CAREER Award 1453459 to M.G.

Author Contributions

P.R.V.S and M.G. conceived of and designed the study, P.R.V.S. performed experiments and analyzed data, and P.R.V.S. and M.G. wrote the manuscript.

References

Bao, W., Kojima, K.K., and Kohany, O. (2015). Repbase Update, a database of repetitive elements in eukaryotic genomes. Mob. DNA *6*, 11.

Batista, R.A., Figueiredo, D.D., Santos-González, J., and Köhler, C. (2019). Auxin regulates endosperm cellularization in Arabidopsis. Genes Dev. 10.1101/gad.316554.118

Baubec, T., Dinh, H.Q., Pecinka, A., Rakic, B., Rozhon, W., Wohlrab, B., von Haeseler, A., and Scheid, O.M. (2010). Cooperation of Multiple Chromatin Modifications Can Generate Unanticipated Stability of Epigenetic States in Arabidopsis. Plant Cell *22*, 34–47.

Birchler, J.A. (2014). Interploidy hybridization barrier of endosperm as a dosage interaction. Front. Plant Sci. *5*.

Birchler, J.A., and Veitia, R.A. (2012). Gene balance hypothesis: Connecting issues of dosage sensitivity across biological disciplines. Proc. Natl. Acad. Sci. *109*, 14746–14753.

Borges, F., and Martienssen, R.A. (2015). The expanding world of small RNAs in plants. Nat. Rev. Mol. Cell Biol. *16*, 727–741.

Borges, F., Parent, J.-S., Ex, F. van, Wolff, P., Martínez, G., Köhler, C., and Martienssen, R.A. (2018). Transposon-derived small RNAs triggered by miR845 mediate genome dosage response in Arabidopsis. Nat. Genet. *50*, 186.

Burkart-Waco, D., Ngo, K., Dilkes, B., Josefsson, C., and Comai, L. (2013). Early Disruption of Maternal–Zygotic Interaction and Activation of Defense-Like Responses in Arabidopsis Interspecific Crosses. Plant Cell *25*, 2037–2055.

Castillo, D.M., and Moyle, L.C. (2012). Evolutionary Implications of Mechanistic
Models of TE-Mediated Hybrid Incompatibility. Int J Evol Biol 2012, 698198

Cooper, D.C. (1951). Caryopsis Development Following Matings Between Diploid and
Tetraploid Strains of Zea Mays. Am. J. Bot. *38*, 702–708.

Creasey, K.M., Zhai, J., Borges, F., Van Ex, F., Regulski, M., Meyers, B.C., and
Martienssen, R.A. (2014). miRNAs trigger widespread epigenetically activated siRNAs from
transposons in *Arabidopsis*. Nature *508*, 411–415.

Dilkes, B.P., Spielman, M., Weizbauer, R., Watson, B., Burkart-Waco, D., Scott, R.J.,
and Comai, L. (2008). The Maternally Expressed WRKY Transcription Factor TTG2 Controls
Lethality in Interploidy Crosses of Arabidopsis. PLOS Biol. *6*, e308.

Doughty, J., Aljabri, M., and Scott, R.J. (2014). Flavonoids and the regulation of seed
size in Arabidopsis. Biochem. Soc. Trans. *42*, 364–369.

Erdmann, R.M., Satyaki, P.R.V., Klosinska, M., and Gehring, M. (2017). A Small RNA
Pathway Mediates Allelic Dosage in Endosperm. Cell Rep. *21*, 3364–3372.

d'Erfurth, I., Jolivet, S., Froger, N., Catrice, O., Novatchkova, M., and Mercier, R.
(2009). Turning Meiosis into Mitosis. PLOS Biol. *7*, e1000124.

Esen, A., and Soost, R.K. (1973). Seed Development in Citrus with Special Reference to
2x X 4x Crosses. Am. J. Bot. *60*, 448–462.

Figueiredo, D.D., Batista, R.A., Roszak, P.J., Hennig, L., and Köhler, C. (2016). Auxin
production in the endosperm drives seed coat development in Arabidopsis. ELife *5*, e20542.

Fiume, E., and Fletcher, J.C. (2012). Regulation of Arabidopsis Embryo and Endosperm
Development by the Polypeptide Signaling Molecule CLE8. Plant Cell *24*, 1000–1012.

Gasciolli, V., Mallory, A.C., Bartel, D.P., and Vaucheret, H. (2005). Partially redundant
functions of Arabidopsis DICER-like enzymes and a role for DCL4 in producing trans-acting
siRNAs. Curr. Biol. *15*, 1494–1500.

Gehring, M., Bubb, K.L., and Henikoff, S. (2009). Extensive Demethylation of
Repetitive Elements During Seed Development Underlies Gene Imprinting. Science *324*, 1447–
1451.

Gehring, M., Missirian, V., and Henikoff, S. (2011). Genomic analysis of parent-of-
origin allelic expression in Arabidopsis thaliana seeds. PloS One *6*, e23687.

Gutierrez-Marcos, J.F., Pennington, P.D., Costa, L.M., and Dickinson, H.G. (2003).
Imprinting in the endosperm: a possible role in preventing wide hybridization. Philos. Trans. R.
Soc. Lond. B Biol. Sci. *358*, 1105–1111.

Haig, and Westboy (1991). Genomic imprinting in endosperm: its effect on seed
development in crosses between species, and its implications for the evolution of apomixis.
Philos. Trans. R. Soc. Lond. B. Biol. Sci. *333*, 1–13.

Håkansson, A., and Ellerström, S. (1950). SEED DEVELOPMENT AFTER
RECIPROCAL CROSSES BETWEEN DIPLOID AND TETRAPLOID RYE. Hereditas *36*,
256–296.

Hehenberger, E., Kradolfer, D., and Köhler, C. (2012). Endosperm cellularization defines
an important developmental transition for embryo development. Development *139*, 2031–2039.

Hsieh, T.-F., Ibarra, C.A., Silva, P., Zemach, A., Eshed-Williams, L., Fischer, R.L., and
Zilberman, D. (2009). Genome-Wide Demethylation of Arabidopsis Endosperm. Science *324*,
1451–1454.

Huang, F., Zhu, Q.-H., Zhu, A., Wu, X., Xie, L., Wu, X., Helliwell, C., Chaudhury, A.,
Finnegan, E.J., and Luo, M. (2017). Mutants in the imprinted PICKLE RELATED 2 gene
suppress seed abortion of fertilization independent seed class mutants and paternal excess
interploidy crosses in Arabidopsis. Plant J. *90*, 383–395.

Ibarra, C.A., Feng, X., Schoft, V.K., Hsieh, T.-F., Uzawa, R., Rodrigues, J.A., Zemach,
A., Chumak, N., Machlicova, A., Nishimura, T., et al. (2012). Active DNA Demethylation in
Plant Companion Cells Reinforces Transposon Methylation in Gametes. Science *337*, 1360–
1364.

Jiang, H., Moreno-Romero, J., Santos-González, J., Jaeger, G.D., Gevaert, K., Slijke,
E.V.D., and Köhler, C. (2017). Ectopic application of the repressive histone modification
H3K9me2 establishes post-zygotic reproductive isolation in Arabidopsis thaliana. Genes Dev.
31: 1272-1287

Jurka, J., Kapitonov, V.V., Pavlicek, A., Klonowski, P., Kohany, O., and Walichiewicz,
J. (2005). Repbase Update, a database of eukaryotic repetitive elements. Cytogenet. Genome
Res. *110*, 462–467.

Kelleher, E.S. (2016). Reexamining the P-Element Invasion of Drosophila melanogaster
Through the Lens of piRNA Silencing. Genetics *203*, 1513–1531.

Kim, D., Pertea, G., Trapnell, C., Pimentel, H., Kelley, R., and Salzberg, S.L. (2013).
TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene
fusions. Genome Biol. *14*, R36.

Kradolfer, D., Wolff, P., Jiang, H., Siretskiy, A., and Köhler, C. (2013). An imprinted
gene underlies postzygotic reproductive isolation in Arabidopsis thaliana. Dev. Cell *26*, 525–
535.

Lafon-Placette, C., and Köhler, C. (2015). Epigenetic mechanisms of postzygotic
reproductive isolation in plants. Curr. Opin. Plant Biol. *23*, 39–44.

Law, J.A., Du, J., Hale, C.J., Feng, S., Krajewski, K., Palanca, A.M.S., Strahl, B.D.,
Patel, D.J., and Jacobsen, S.E. (2013). Polymerase IV occupancy at RNA-directed DNA
methylation sites requires SHH1. Nature *498*, 385–389.

Li, J., and Berger, F. (2012). Endosperm: food for humankind and fodder for scientific
discoveries. New Phytol. *195*, 290–305.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550.

Lu, J., Zhang, C., Baulcombe, D.C., and Chen, Z.J. (2012). Maternal siRNAs as
regulators of parental genome imbalance and gene expression in endosperm of Arabidopsis
seeds. Proc. Natl. Acad. Sci. U. S. A. *109*, 5529–5534.

Marí-Ordóñez, A., Marchais, A., Etcheverry, M., Martin, A., Colot, V., and Voinnet, O.
(2013). Reconstructing *de novo* silencing of an active plant retrotransposon. Nat. Genet. *45*,
1029–1039.

Martienssen, R.A. (2010). Heterochromatin, small RNA and post-fertilization dysgenesis
in allopolyploid and interploid hybrids of Arabidopsis. New Phytol. *186*, 46–53.

Martinez, G., Wolff, P., Wang, Z., Moreno-Romero, J., Santos-González, J., Conze, L.L.,
DeFraia, C., Slotkin, R.K., and Köhler, C. (2018). Paternal easiRNAs regulate parental genome
dosage in Arabidopsis. Nat. Genet. *50*, 193.

McCue, A.D., Nuthikattu, S., Reeder, S.H., and Slotkin, R.K. (2012). Gene Expression
and Stress Response Mediated by the Epigenetic Regulation of a Transposable Element Small
RNA. PLOS Genet. *8*, e1002474.

Milbocker, D., and Sink, K. (1969). Embryology of diploid × diploid and diploid ×
tetraploid crosses in poinsettia. Can. J. Genet. Cytol. *11*, 598–601.

Mittelsten Scheid, O., Afsar, K., and Paszkowski, J. (2003). Formation of stable
epialleles and their paramutation-like interaction in tetraploid *Arabidopsis thaliana*. Nat. Genet.
34, 450–454.

Muntzing, A. (1936). The Evolutionary Significance of Autopolyploidy. Hereditas 21:
363-378

Oberlin, S., Sarazin, A., Chevalier, C., Voinnet, O., and Marí-Ordóñez, A. (2017). A
genome-wide transcriptome and translatome analysis of Arabidopsis transposons identifies a
unique and conserved genome expression strategy for Ty1/Copia retroelements. Genome Res.
27, 1549–1562.

Pignatta, D., Erdmann, R.M., Scheer, E., Picard, C.L., Bell, G.W., and Gehring, M.
(2014). Natural epigenetic polymorphisms lead to intraspecific variation in Arabidopsis gene
imprinting. ELife *3*, e03198.

Pignatta, D., Bell, G., and Gehring, M. (2015). Whole Genome Bisulfite Sequencing and
DNA Methylation Analysis from Plant Tissue. BIO-Protoc. *5*.

Piskurewicz, U., Iwasaki, M., Susaki, D., Megies, C., Kinoshita, T., and Lopez-Molina,
L. (2016). Dormancy-specific imprinting underlies maternal inheritance of seed dormancy in
Arabidopsis thaliana. ELife *5*, e19573.

Povilus, R.A., Diggle, P.K., and Friedman, W.E. (2018). Evidence for parent-of-origin
effects and interparental conflict in seeds of an ancient flowering plant lineage. Proc. Biol. Sci.
285.

Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for
comparing genomic features. Bioinformatics *26*, 841–842.

Satyaki, P.R.V., and Gehring, M. (2017). DNA methylation and imprinting in plants:
machinery and mechanisms. Crit. Rev. Biochem. Mol. Biol. *52*, 163–175.

Schatlowski, N., Wolff, P., Santos-González, J., Schoft, V., Siretskiy, A., Scott, R.,
Tamaru, H., and Köhler, C. (2014). Hypomethylated Pollen Bypasses the Interploidy
Hybridization Barrier in Arabidopsis. Plant Cell *26*, 3556–3568.

Schon, M.A., and Nodine, M.D. (2017). Widespread Contamination of Arabidopsis
Embryo and Endosperm Transcriptome Data Sets. Plant Cell *29*, 608–617.

Scott, R.J., Spielman, M., Bailey, J., and Dickinson, H.G. (1998). Parent-of-origin effects
on seed development in Arabidopsis thaliana. Development *125*, 3329–3341.

Stoute, A.I., Varenko, V., King, G.J., Scott, R.J., and Kurup, S. (2012). Parental genome
imbalance in Brassica oleracea causes asymmetric triploid block: Parent-of-origin effects in
Brassica oleracea. Plant J. 71: 503-516.

Stroud, H., Greenberg, M.V.C., Feng, S., Bernatavichute, Y.V., and Jacobsen, S.E.
(2013). Comprehensive Analysis of Silencing Mutants Reveals Complex Regulation of the
Arabidopsis Methylome. Cell *152*, 352–364.

Tiwari, S., Spielman, M., Schulz, R., Oakey, R.J., Kelsey, G., Salazar, A., Zhang, K.,
Pennell, R., and Scott, R.J. (2010). Transcriptional profiles underlying parent-of-origin effects in
seeds of Arabidopsis thaliana. BMC Plant Biol. *10*, 72.

Tsay, Y.F., Frank, M.J., Page, T., Dean, C., and Crawford, N.M. (1993). Identification of
a mobile endogenous transposon in Arabidopsis thaliana. Science *260*, 342–344.

Vu, T.M., Nakamura, M., Calarco, J.P., Susaki, D., Lim, P.Q., Kinoshita, T.,
Higashiyama, T., Martienssen, R.A., and Berger, F. (2013). RNA-directed DNA methylation
regulates parental genomic imprinting at several loci in Arabidopsis. Development *140*, 2953–
2960.

Walia, H., Josefsson, C., Dilkes, B., Kirkbride, R., Harada, J., and Comai, L. (2009).
Dosage-Dependent Deregulation of an AGAMOUS-LIKE Gene Cluster Contributes to
Interspecific Incompatibility. Curr. Biol. *19*, 1128–1132.

Walker, J., Gao, H., Zhang, J., Aldridge, B., Vickers, M., Higgins, J.D., and Feng, X.
(2018). Sexual-lineage-specific DNA methylation regulates meiosis in Arabidopsis. Nat. Genet.
50, 130.

Wang, G., Jiang, H., Del Toro de León, G., Martinez, G., and Köhler, C. (2018).
Sequestration of a Transposon-Derived siRNA by a Target Mimic Imprinted Gene Induces
Postzygotic Reproductive Isolation in Arabidopsis. Dev. Cell *46*, 696-705.

Wang, L., Feng, Z., Wang, X., Wang, X., and Zhang, X. (2010). DEGseq: an R package
for identifying differentially expressed genes from RNA-seq data. Bioinforma. Oxf. Engl. *26*,
136–138.

von Wangenheim, K.-H., and Peterson, H.-P. (2004). Aberrant endosperm development
in interploidy crosses reveals a timer of differentiation. Dev. Biol. *270*, 277–289.

Williams, B.P., Pignatta, D., Henikoff, S., and Gehring, M. (2015). Methylation-Sensitive
Expression of a DNA Demethylase Gene Serves As an Epigenetic Rheostat. PLOS Genet. *11*,
e1005142.

Wolff, P., Jiang, H., Wang, G., Santos-González, J., and Köhler, C. (2015). Paternally
expressed imprinted genes establish postzygotic hybridization barriers in Arabidopsis thaliana.
ELife *4:e10074*.

Xiao, W., Brown, R.C., Lemmon, B.E., Harada, J.J., Goldberg, R.B., and Fischer, R.L.
(2006). Regulation of Seed Size by Hypomethylation of Maternal and Paternal Genomes. Plant
Physiol. *142*, 1160–1168.

Yan, D., Duermeyer, L., Leoveanu, C., and Nambara, E. (2014). The Functions of the
Endosperm During Seed Germination. Plant Cell Physiol. *55*, 1521–1533.

Yu, Z., Haberer, G., Matthes, M., Rattei, T., Mayer, K.F.X., Gierl, A., and Torres-Ruiz,
R.A. (2010). Impact of natural genetic variation on the transcriptome of autotetraploid
Arabidopsis thaliana. Proc. Natl. Acad. Sci. *107*, 17809–17814.

Zhang, J., Liu, Y., Xia, E.-H., Yao, Q.-Y., Liu, X.-D., and Gao, L.-Z. (2015).
Autotetraploid rice methylome analysis reveals methylation variation of transposable elements
and their effects on gene expression. Proc. Natl. Acad. Sci. U. S. A. *112*, E7022-7029.

Zhang, Y., Malone, J.H., Powell, S.K., Periwal, V., Spana, E., MacAlpine, D.M., and
Oliver, B. (2010). Expression in Aneuploid Drosophila S2 Cells. PLOS Biol. *8*, e1000320.

Zilberman, D., Gehring, M., Tran, R.K., Ballinger, T., and Henikoff, S. (2007). Genome-
wide analysis of *Arabidopsis thaliana* DNA methylation uncovers an interdependence between
methylation and transcription. Nat. Genet. *39*, 61–69.

1000

Supplemental Data Files

Supplemental Figure 1: Paternal loss of RdDM genes but not *RDR6* and *CMT3* represses seed abortion in paternal excess crosses.

Supplemental Figure 2: Analysis of gene expression differences in paternal excess endosperm derived using the *osd1* mutation.

Supplemental Figure 3: Rescue by loss of *NRPD1* does not require normalization of genes implicated in interploid seed abortion.

Supplemental Figure 4: Small RNA in balanced, lethal and viable paternal excess endosperm.

Supplemental figure 5: Allelic contributions to gene expression differences.

Supplemental Figure 6: Allelic bias of some imprinted genes is lost in lethal paternal excess.

Supplemental Table 1: Details of sequencing libraries.

Supplemental Dataset 1: Cuffdiff output comparing gene expression in balanced, lethal and viable paternal excess endosperm.

Supplemental Dataset 2: Gene ontology analysis of genes that are differentially expressed between balanced, lethal and viable paternal excess endosperm.

Supplemental Dataset 3: DEGseq output comparing transposon transcript levels between balanced, lethal and viable paternal excess endosperm.

Supplemental Dataset 4: Differentially methylated regions in the CHH context in comparisons between balanced, lethal and viable paternal excess endosperm.

Supplemental Dataset 5: Differentially methylated region in the CHG context in comparisons between balanced, lethal and viable paternal excess endosperm.

Supplemental Dataset 6: Differentially methylated region in the CG context in comparisons between balanced, lethal and viable paternal excess endosperm.

Supplemental Dataset 7: DESeq output for differential abundance of 21 and 24 nt

genic sRNA between balanced, lethal and viable paternal excess endosperm.

Supplemental Dataset 8: DESeq output for differential abundance of 21 and 24 nt TE

sRNA between balanced, lethal and viable paternal excess endosperm.

Supplemental Dataset 9: Allele-specific mRNA read counts and differences in allelic

contributions between balanced, lethal and viable paternal excess endosperm.

1032