The TRAPP complex mediates secretion arrest induced by stress granule assembly

Francesca Zappa[1], Cathal Wilson[1], Giuseppe Di Tullio[1], Michele Santoro[1], Piero Pucci[2],

Maria Monti[2], Davide D'Amico[1], Sandra Pisonero Vaquero[1], Rossella De Cegli[1],

Alessia Romano[1], Moin A. Saleem[3], Elena Polishchuk[1], Mario Failli[1], Laura Giaquinto[1],

Maria Antonietta De Matteis[1, 2]

1 Telethon Institute of Genetics and Medicine, Pozzuoli (Naples), Italy

2 Federico II University, Naples, Italy

3 Bristol Renal, Bristol Medical School, University of Bristol, UK

Correspondence to: f.zappa@tigem.it, dematteis@tigem.it

The TRAnsport Protein Particle (TRAPP) complex controls multiple steps along the secretory, endocytic and autophagic pathways and is thus strategically positioned to mediate the adaptation of membrane trafficking to diverse environmental conditions including acute stress. We identified TRAPP as a key component of a branch of the integrated stress response that impinges on the secretory pathway. We found that TRAPP associates with, and drives the recruitment of the inner components of the COPII coat to, SGs. The sequestration to SGs of COPII, required for the ER export of newly synthesized proteins, and of TRAPP, the exchange factor for Rab1, a GTPase controlling the architecture of the Golgi complex, leads to arrest of the secretory activity and to the disorganization of the Golgi complex. Interestingly, the relocation of TRAPP and COPII to SGs only occurs in actively proliferating cells and it does it in a CDK1/2 dependent manner. We show that CDK1/2 activity controls the membrane-cytosol cycle of COPII at ER exit sites (ERES) and that CDK1/2 inhibition/depletion prevents the relocation of COPII and TRAPP to SGs by stabilizing them at the ERES. Importantly, TRAPP is not just a passive constituent of SGs but controls their maturation. This includes the acquisition of signaling components, such as RACK1 and Raptor, whose sequestration to SGs is needed to mediate the pro-survival response. SGs that assemble in TRAPP-depleted cells are no longer able to sequester RACK1 and Raptor thus rendering the cells more prone to undergo apoptosis upon stress exposure.

Altogether, our findings reveal a new property of the integrated stress response that is acquired through the recruitment of TRAPP to SGs: a high grade of plasticity that tailors the secretory pathway to stress according to different cell growth states.

Introduction

The TRAPP (TRAnsport Protein Particle) complex is a conserved multimolecular complex intervening in multiple segments of membrane trafficking along the secretory, the endocytic and the autophagy pathways (Kim et al., 2016).

Originally identified in yeast as a tethering factor acting in ER-to-Golgi trafficking, it was subsequently discovered to act as a GEF for Ypt1 and Ypt31/32 in yeast and for Rab1 and possibly Rab11 in mammals (Westlake et al., 2011; Yamasaki et al., 2009; Zou et al., 2012). The TRAPP complex has a modular composition and is present as two forms in mammals: TRAPPII and TRAPPIII which share a common heptameric core (TRAPPC1, TRAPPC2, TRAPPC2L, TRAPPC3, TRAPPC4, TRAPPC5, TRAPPC6) and additional subunits specific for TRAPPII (TRAPPC9/ TRAPPC10) or for TRAPPIII (TRAPPC8, TRAPPC11, TRAPPC12, TRAPPC13) (Sacher et al., 2018). TRAPPII has been implicated in late Golgi trafficking while TRAPPIII has a conserved role in the early secretory pathway (ER-to-Golgi) and in autophagy (Scrivens et al., 2011; Yamasaki et al., 2009). Recent lines of evidence have expanded the range of activities of the TRAPP complex by showing that it takes part in a cell survival response triggered by agents that disrupt the Golgi complex (Ramírez-Peinado et al., 2017) and can drive the assembly of lipid droplets in response to lipid load (Li et al., 2017).

The importance of the TRAPP complex in humans is testified by the deleterious consequences caused by mutations in genes encoding distinct TRAPP subunits. Mutations in TRAPPC2L, TRAPPC6A, TRAPPC6B, TRAPPC9 and TRAPPC12 cause neurodevelopmental disorders leading to intellectual disability and dysmorphic syndromes (Harripaul et al., 2018; Khattak and Mir, 2014; Milev et al., 2018, 2017, p. 12; Mohamoud et al., 2018), mutations in TRAPPC11 lead to ataxia (Koehler et al., 2017), and mutations in TRAPPC2 lead to the spondyloepiphyseal dysplasia tarda (SEDT) (Gedeon et al., 1999).

SEDT is characterized by short stature, platyspondyly, barrel chest and premature osteoarthritis that manifest in late childhood/prepubertal age. We have shown that pathogenic mutations or deletion of TRAPPC2 alter the ER export of procollagen (both type I and type II) and that TRAPPC2 interacts with the procollagen escort protein TANGO1 and regulates the cycle of the GTPase Sar1 at the ER exit sites

71	(ERES) (Venditti et al., 2012)). The Sar1 cycle in turn drives the cycle of the COPII coat
72	complex, which mediates the formation of carriers containing neo-synthesized cargo
73	to be transported to the Golgi complex. While this role of TRAPPC2 in the ER export
74	of PC might explain the altered ECM deposition observed in patient's cartilage
75	(Venditti et al., 2012; Tiller et al., 2001), it leaves the late onset of the disease signs
76	unexplained. We hypothesized that the latter could be due to an inability to
77	maintain long-term cartilage tissue homeostasis, possibly due to an impaired
78	capacity of chondrocytes to face the physiological stresses that underlie and guide
79	their development and growth. These include mechanical stress that can induce
80	oxidative stress leading to apoptosis (Henrotin et al., 2003; Zuscik et al., 2008).
81	Here, by analyzing the cell response to different stresses, we show that the TRAPPC2
82	and the entire TRAPP complex, are components of stress granules (SGs), membrane-
83	less organelles that assemble in response to stress (Protter and Parker, 2016).
84	The recruitment of TRAPP to SGs has multiple impacts on the stress response as it
85	induces the sequestration of Sec23/Sec24 (the inner layer of the COPII complex)
86	onto SGs thus inhibiting trafficking along the early secretory pathway, leads to the
87	inactivation of the small GTPase Rab1 with a consequent disorganization of the Golgi
88	complex, and is required for the recruitment of signalling proteins, such as Raptor
89	and RACK1, to SGs thus contributing to the anti-apoptotic role of SGs. Interestingly,
90	TRAPP and COPII recruitment to SGs only occurs in actively proliferating cells and is
91	under control of cyclin-dependent-kinases (CDK 1 and 2).
92	The TRAPP complex thus emerges as a key element in conferring an unanticipated
93	plasticity to SGs, which adapt their composition and function to cell growth activity,
94	reflecting the ability of cells to adjust their stress response to their proliferation state
95	and energy demands.
96	
97	
98	**Results**
99	
100	**TRAPP redistributes to SGs in response to different stress stimuli**
101	To investigate the involvement of TRAPPC2 in the stress response, we exposed
102	chondrocytes (Figure 1A) or HeLa cells (Figure 1B) to sodium arsenite (SA) treatment

that induces oxidative stress. We have shown previously that TRAPPC2 associate with ERES under steady state conditions (Venditti et al., 2012). SA treatment led to dissociation of TRAPPC2 from ERES and a complete relocation to roundish structures. Oxidative stress is known to lead to the formation of stress granules (SGs; Anderson and Kedersha, 2002), therefore, we considered the possibility that these TRAPPC2-positive structures might be SGs. Indeed, co-labeling with an anti-eIF3 antibody, a canonical SG marker (Aulas et al., 2017), showed the co-localization of TRAPPC2 with eIF3 after SA treatment (Figure 1A,B). Other TRAPP components, such as TRAPPC1 and TRAPPC3, exhibited the same response to SA treatment (Figure 1C). TRAPP recruitment to SGs also occurs in response to heat shock (Figure 1—figure supplement 1A), and in different cell lines (Figure 1—figure supplement 1B,C). In addition, TRAPP elements associate to SGs in yeast cells exposed to heat stress (Figure 1—figure supplement 2, Figure 1—supplement video 1 and 2) thus suggesting that this is a conserved process in evolution. Of note, the association of TRAPP with SGs is fully reversible after stress removal (Figure 1—figure supplement 3).

We studied the kinetics of TRAPPC2 redistribution to SGs. The appearance of eIF3-positive SGs occurred 7 min after SA treatment (Figure 1D), while TRAPPC2 re-localization to SGs began 15 min after exposure to stress and gradually increased over time, with massive recruitment occurring 60 min after treatment, thus lagging behind the initial assembly of SGs (Figure 1D).

We observed that oxidative stress did not affect the overall integrity of the TRAPP complexes (both TRAPPII and TRAPPIII), as analyzed by gel filtration. However, in stressed cells TRAPPII and TRAPPIII components were also found in higher molecular weight fractions (Figure 1E) containing the SG component TIAR-1 (Figure 1E), possibly reflecting the SG-associated TRAPP.

To investigate the mechanisms that could mediate the recruitment of TRAPP to SGs, we performed a proteomics analysis of the TRAPPC2 interactors (see Materials and methods). This analysis (Figure 1F, Table S1), confirmed known TRAPPC2 interactors (e.g. TRAPP complex components, Rab1, COPII and COPI) and revealed the presence of many RNA binding proteins (RBPs), including those with a central role in the assembly of SGs (Jain et al., 2016). This suggested that, as described for other

components of SGs, TRAPPC2 may be recruited to growing SGs by a piggyback

mechanism (Anderson and Kedersha, 2008), i.e. via the interaction with RBPs that

are components of SGs.

Altogether, these results led us to conclude that following stress the TRAPP complex,

through TRAPPC2 interaction with RBPs, becomes a SG component.

TRAPPC2 is required for recruiting COPII to SGs

The translocation of TRAPP to SGs prompted us to investigate whether other

components of membrane trafficking machineries behaved similarly. We screened

different coat complex components (COPI, COPII, clathrin adaptors, clathrin), and

other cytosolic proteins associated with the exocytic and endocytic pathways. Out of

the 15 proteins tested, only components of the inner layer of the COPII coat, Sec24

(Figure 2A, B) and Sec23 (Figure 2—figure supplement 1A), but none of the others

(Figure 2C, and Figure 2—figure supplement 1B-N, Figure 2—figure supplement 2)

associated with SGs. Notably, Sec24 was recruited to SGs with kinetics similar to

those of the TRAPP complex (Figure 2A).

Since TRAPP and COPII are known to interact and TRAPP is recruited to ERES in a

Sar1- and COPII-dependent manner (Lord et al., 2011; Venditti et al., 2012), we

asked whether COPII and TRAPP recruitment to SGs is interdependent. The depletion

of the COPII inner layer proteins Sec23 and Sec24 had no impact on the recruitment

of the TRAPP complex to SGs (Figure 2D,E, Figure 2—figure supplement 1A,B) while

the depletion of the entire TRAPP complex (by KD of TRAPPC3, which destabilizes the

entire complex) or of the TRAPPC2 subunit (by KD of TRAPPC2 that leaves unaltered

the rest of the TRAPP complex; Venditti et al., 2012) abrogated the recruitment of

COPII components to SGs (Figure 2F, Figure 2—figure supplement 1C). These results

indicated that the TRAPP complex, through its component TRAPPC2, drives the

recruitment of COPII to SGs.

It is worth mentioning that COPII components have been reported to associate, in

Drosophila S2 cells, with other membrane-less organelles: the "Sec bodies"

(Zacharogianni et al., 2014). However, Sec bodies are distinct from SGs since they do

not contain ribonucleoproteins and form in response to prolonged starvation, but

167 not to oxidative stress. In addition, the recruitment of COPII to Sec bodies in
168 Drosophila S2 cells depends of the ERES component Sec16. We found that in
169 mammalian cells Sec16 neither significantly associates with SGs (Figure 2—figure
170 supplement 2A,B) nor is it required for the recruitment of COPII to SGs (Figure 2—
171 figure supplement 2C). These results indicate that stress-induced COPII phase
172 separation diverges in Drosophila and mammalian cells for the inducing stresses
173 (starvation vs oxidative or heat stress), for the composition of the membrane-less
174 organelles (Sec bodies vs SGs) and for the underlying mechanisms (Sec16-dependent
175 vs TRAPP-dependent mechanism).

176 We then moved to investigate the nature of association of COPII to SGs since the
177 COPII coat cycles in a very dynamic fashion at ERES (D'Arcangelo et al., 2013). To this
178 end, we permeabilized living cells (Kapetanovich et al., 2005) at steady state and
179 upon stress induction. Treatment with digitonin forms pores at the plasma
180 membrane that allows the exit of the cytosolic content into the extracellular space.
181 Under these conditions, we found that the COPII pool recruited to SGs is less
182 dynamic than the pool cycling at the ERES, since COPII remained associated with SGs
183 in permeabilized cells exposed to stress while the COPII pool cycling at the ERES was
184 completely lost upon permeabilization of unstressed cells (Figure 3A). We then
185 assessed the possible interplay between the COPII fraction at ERES and the pool that
186 eventually relocates to SGs. We asked whether changes in the association of COPII at
187 the ERES could have an impact on the recruitment of COPII to SGs. To this end, we
188 stabilized COPII and TRAPP at ERES by expressing either a constitutively active GTP-
189 bound Sar1 mutant (Sar1-H79G) or decreasing the rate of GTP hydrolysis on Sar1 by
190 depleting Sec31, which is a co-GAP that potentiates by an order of magnitude the
191 Sar1 GAP activity of Sec23-24 (Bi et al., 2007). Under both conditions, COPII and
192 TRAPP were more tightly associated with ERES (Figure 3—figure supplement 1),
193 while their translocation to SGs upon exposure to stress was significantly reduced
194 (Figure 3B,C). Thus, COPII/TRAPP recruitment to SGs is commensurate with the rate
195 of their association at the ERES indicating that the cytosolic pool of COPII generated
196 through the fast cycling at ERES is the one available to be recruited to SGs upon
197 stress.

198

199 **The association of TRAPP is under the control of CDK1/2**

200 To dissect the regulation of TRAPP/COPII recruitment to SGs, we screened a library

201 of kinase inhibitors for their ability to affect the re-localization of COPII to SGs. HeLa

202 cells were pre-treated with kinase inhibitors for 150 min and then exposed to SA for

203 a further 30 min in the continuous presence of the inhibitors. Out of the 273

204 inhibitors tested, 194 had no effect on SG formation or on COPII recruitment, 13

205 were generally toxic (65% mortality) at the tested concentrations, one (AZD4547)

206 affected the formation of SGs, while the remaining 32 specifically inhibited (by at

207 least 35%) the association of Sec24C with SGs without affecting SG formation (Figure

208 4A). Enrichment analysis on this last group of inhibitors highlighted that CDK

209 inhibitors were the most represented class (Figure 4B). Western blot analysis

210 confirmed that CDK activity was impaired in cells treated with the CDK inhibitors

211 (Figure 4C). In addition, we observed that there is an increase in CDK activity in SA-

212 treated cells that was sensitive to CDK inhibitors (Figure 4C).

213 The CDK protein kinase family is composed of 21 members (Malumbres, 2014) and

214 available inhibitors have limited selectivity (Asghar et al., 2015). To identify which

215 specific CDK might be involved in Sec24C recruitment, we compared the potency of

216 selected CDK inhibitors (Flavopiridol hydrochloride, SNS-032, Dinaciclib, AT7519,

217 PHA-793887, ADZ5438, JNJ-7706621, PHA-767491, BMS-265246, PHA-848125,

218 Roscovitine, Palbociclib, BS-181HCl) in the Sec24C recruitment assay with their

219 ability to inhibit different CDK isoforms, as described by Selleckchem

220 (https://www.selleckchem.com/CDK.html). This analysis revealed that CDK1 and 2

221 were those commonly targeted by the inhibitors that most potently affected Sec24C

222 recruitment to SGs (Figure 4D), strongly suggesting that CDK1/2 are the most likely

223 CDKs controlling COPII recruitment to SGs. This was confirmed by the observation

224 that down-regulating the expression of CDK1 and CDK2 by siRNA inhibited COPII

225 recruitment to SGs (Figure 4E, Figure 4—figure supplement 1). The three best hits

226 (Flavopiridol, Dinaciclib, SNS032, Figure 4D,F) were assayed using a dose-response

227 analysis and were found to be effective at concentrations as low as 100 nM (Figure

228 4G). As a negative control, the specific CDK7 inhibitor (BS-181HCl) was ineffective,

229 even at the highest concentration tested (Figure 4G).

Since COPII recruitment to SGs is driven by TRAPP recruitment (Figure 2F), we

predicted that CDK1/2 inhibitors would also inhibit TRAPP recruitment. In line with

our expectations, TRAPP re-localization to SGs was reduced in cells treated with

CDK1/2 inhibitors (Figure 4H).

Altogether, these results indicate that a signaling pathway involving CDK1/2

regulates the redistribution of TRAPP and COPII from ERES to SGs and prompted us

to search for the process and possibly the substrates targeted by CDK1/2.

We excluded the possibility that CDK1/2 could directly target TRAPPC2 or Sec23/24

as none of them appeared to undergo CDK1/2 dependent phosphorylation (data not

shown). Thus, we hypothesized that CDK inhibition might indirectly inhibit the

recruitment of TRAPP/COPII to SGs by stabilizing their association with ER

membranes. Indeed, we found that CDK inhibitors stabilize TRAPP and COPII on ERES

membranes even in unstressed cells (Figure 5A). The analysis of the ERES associated

pool using a permeabilization assay confirmed that the CDK1/2 inhibitors strongly

stabilized COPII at the ERES, thus increasing the membrane associated pool and

decreasing the cytosolic one (Figure 5B,C).

To identify a possible target of CDK1/2 among the regulators of COPII cycle at the

ERES, we focused on Sec31 for three main considerations. First, Sec31 regulates Sar1

cycling at the ERES acting as co-GAP of Sec23-24 (Bi et al., 2007); second, we knew

that the regulation of Sar1 cycle by Sec31 is required for the relocation of

COPII/TRAPP from ERES to SGs (Figure 3C); finally, it has been shown that Sec31 is a

target of CDKs in yeast (Holt et al., 2009) and in *Trypanosoma brucei* (Hu et al., 2016).

In mammals, Sec31 activity is known to be modulated by post-translational

modifications and specifically by phosphorylation on multiple sites (Koreishi et al.,

2013) one of which (serine 799) is included in a consensus motif for CDK

phosphorylation. We tested whether Sec31 could be a substrate for CDK using an

antibody specific for phosphoSer CDK substrates. We found that Sec31 is indeed a

target for CDKs (Figure 5D), and that its CDK-dependent phosphorylation is increased

upon SA treatment and decreased by the CDK inhibitor Dinaciclib (Fig. 5D).

Altogether, the above results suggest that at least one of the mechanisms underlying

the regulation of TRAPP/COPII recruitment to SGs by CDK1/2 involves the ability of

CDK1/2 to control the rate of Sar1 and COPII cycling at the ERES. This control is likely

to be exerted through Sec31, which, acting as a co-GAP for Sar1, speeds up the COPII

cycle at the ERES, thus controlling the size of the COPII pool that can be mobilized to

SGs in response to stress.

The recruitment of TRAPP and COPII to SGs occurs in actively proliferating cells

The observation that the recruitment of TRAPP and COPII to SGs is under control of

CDK1/2 suggested that it could be linked to the cell proliferation. We explored this

possibility by decreasing the cell proliferation rate in diverse and independent ways.

First, cells were subjected to prolonged nutrient starvation (HBSS for 8 hr) and then

were exposed to SA for 30 min. Under these conditions, which inhibited cell

proliferation, TRAPP/COPII association with SGs was reduced (Figure 6A,B).

Second, HeLa cells were seeded at different confluency and then treated with SA for

30 min. Strikingly, the extent of confluency was inversely related to the extent of

TRAPP/COPII association with SGs and to the degree of proliferation (Figure 6C).

Third, we monitored the recruitment of TRAPP/COPII to SGs in podocyte cells

expressing a temperature sensitive LTA-SV40 that actively proliferate at the

permissive temperature of 33ºC but arrest proliferation and differentiate at the

restrictive temperature of 37ºC (Saleem et al., 2002) (Figure 6—figure supplement 1).

While TRAPP and COPII associate with SGs in undifferentiated podocytes in response

to stress, this association was reduced in differentiated podocytes (Figure 6D). Of

note, the proliferation to differentiation switch, as described (Saurus et al., 2016), is

also accompanied by a decline in CDK activity (Figure 6E,F). Interestingly, also the

extent of CDK-dependent phosphorylation of Sec31 is dampened in differentiated as

compared with proliferating podocytes (Figure 6G), thus reinforcing the correlation

between the phosphorylation of Sec31 and the recruitment of COPII and TRAPP to

SGs.

Altogether these data indicate that the association of COPII and TRAPP with SGs

occurs only in proliferating cells and requires active CDK1/2 signaling.

The TRAPP complex imposes SG size, composition, and function

293 The identification of TRAPP as a component of SGs led us to ask whether it could
294 play any role in the assembly and function of SGs themselves. Indeed, we observed
295 that depleting the entire TRAPP complex or only TRAPPC2 (through siRNA-mediated
296 depletion) decreased the size of SGs (Figure 2F and Figure 7A). Microinjection of a
297 TRAPPC3-blocking Ab (Yu et al., 2006; Venditti et al., 2012) had a similar effect
298 (Figure 7B). Moreover, a smaller size of SGs was also observed upon treatment with
299 CDK1/2 inhibitors, which prevents TRAPP translocation to SGs (Figure 4F,H). Under
300 all of the above conditions the size of the SGs recall that of "immature SGs"
301 (Wheeler et al., 2016).

302 In fact, the processes that lead to SG nucleation, i.e. shut-down of protein synthesis
303 or phosphorylation of eIF2alpha induced by oxidative stress, were not affected by
304 TRAPP depletion (Figure 7—figure supplement 1A,B). Instead, the successive
305 increase in size of the nucleated SGs (i.e. SG maturation) was impaired in TRAPP-
306 depleted cells, consistent with our observation that TRAPP is recruited after the
307 initial formation of SGs (Figure 1D) and occupies an outer position in the SGs (figure
308 7C). A feature of SG maturation is the acquisition of an array of signaling molecules
309 (which can be either hyperactivated or deactivated) to drive the pro-survival
310 program (Franzmann and Alberti, 2019). This is the case of the energy sensor
311 mTORC1 (through the sequestration of its partner Raptor) and the kinase RACK1
312 (Arimoto et al., 2008; Thedieck et al., 2013; Wippich et al., 2013). To evaluate
313 whether the absence of TRAPP could affect SG maturation and/or signaling function
314 by interfering with the recruitment of these proteins, we checked the localization of
315 Raptor (Figure 7D) and RACK1 (Figure 7E, Figure 7—figure supplement 2) in TRAPP-
316 and or TRAPPC2-depleted cells subjected to oxidative stress. We observed that the
317 amount of these signaling proteins sequestered to SGs was strongly reduced by
318 TRAPP depletion, thus indicating that the TRAPP complex is necessary to generate
319 "mature" SGs that can accomplish a fully functional stress response. Localization of
320 RACK1 and Raptor was also evaluated in a context of CDK inhibition. Again,
321 treatment with CDK inhibitors strongly reduced Raptor (Figure 7F) and RACK1 (Figure
322 7G) sequestration, further supporting that the TRAPP complex is needed for SG
323 maturation. Notably, the distribution of Raptor in response to oxidative stress was

324 also studied by transiently expressing a cMyc-tagged version of the protein in control

325 cells, in TRAPP KD and in cells treated with CDK inhibitors. While myc-Raptor readily

326 relocated to SGs in control cells exposed to stress, it failed to associate with SGs

327 under conditions that prevented TRAPP association with SGs (i.e. TRAPP depletion

328 and CDK inhibition; data not shown).

329 SGs are part of a pro-survival program that aims to save energy to cope with stress

330 insults. Interfering with this program by impairing SG formation sensitizes cells to

331 apoptosis (Takahashi et al., 2013). We thus assayed stress resistance in cells with an

332 impaired capability to recruit TRAPP to SGs, i.e. upon CDK inhibition. As shown in

333 Figure 7H, CDK inhibition significantly sensitized cells to cell death, suggesting that

334 the presence of TRAPP at SGs is required for their anti-apoptotic role and

335 highlighting the TRAPP complex as a novel player in the cell response to stress.

336

337 **The recruitment of TRAPP and COPII to SGs impairs ER export of newly synthesized**

338 **cargoes and induces the fragmentation of the Golgi complex**

339 We finally investigated the functional relevance of the recruitment of TRAPP/COPII

340 to SGs by analyzing its impact on the function and structure of the secretory pathway.

341 We first analysed the ER export of procollagen type I (PC-I), which is known to be

342 regulated by COPII and TRAPP (Gorur et al., 2017; Venditti et al., 2012). PC-I

343 transport can be synchronized by temperature, being unfolded and blocked in the ER

344 at 40°C while properly folded and assembled and able to be exported from the ER,

345 transported to the Golgi complex (GC) and secreted at 32°C (Mironov et al., 2001,

346 and Methods). Human fibroblasts were incubated at 40°C, left untreated or treated

347 with SA, which induces the recruitment of Sec24C to SGs (Figure 8A). PC-I was

348 blocked in the ER under both conditions (Figure 8A). Monitoring of PC-I transport

349 after the shift to 32°C showed that while PCI reaches the GC in untreated cells, it

350 remains blocked in the ER in SA-treated cells (Figure 8A).

351 We then investigated to what extent the inhibition of cargo export from the ER

352 induced by oxidative stress is dependent on COPII/TRAPP sequestering onto SGs. To

353 this end we compared the ER export of neosynthesised cargo in control cells and in

354 SA-treated cells either under conditions permissive for the formation of SGs (and of

the recruitment of COPII/TRAPP to SG) or under conditions that prevent SG formation. For the latter, we took advantage of the use of ISRIB, a compound that antagonises the effect of eIF2α phosphorylation on translation and impairs SG assembly (Sidrauski et al., 2015). In preliminary experiments, we checked the ability of ISRIB to impair SG formation under our experimental conditions and found that ISRIB completely abrogates SG assembly after 15 min SA treatment, delaying their appearance at later time points of treatment and thus significantly also impairing COPII sequestering to SGs (Figure 8—figure supplement 1). To monitor the synchronized export of neosynthesised cargo we followed a reporter cargo, VSVG-UVR8, that exits the ER in a UV light-inducible manner (Chen et al., 2013). After UV light induction, VSVG-UVR8 exited the ER and reached the Golgi in control cells, while its ER export was severely impaired in cells treated with SA under conditions permissive for SG formation where COPII was massively recruited to SGs (Figure 8B, D). Interestingly, the inhibition of ER export of VSVG-UVR8 was largely prevented when the cells were exposed to similar oxidative stress, but under conditions that prevented SG formation (i.e. when cells were pretreated with ISRIB (Figure 8B, D)).

We next tested whether COPII and TRAPP maintain their functionality once released from SGs upon stress removal (Figure 1—figure supplement 3). Cells were treated with SA for 30 min, the SA was washed out, and cells were left to recover for 180 min in the presence of cycloheximide to prevent *de novo* synthesis of TRAPP and COPII components. Under these conditions SGs were resolved, COPII returned to its native location (ERES/cytosol) and cells completely recovered their capability to transport cargo to the Golgi apparatus (Figure 8C,D). These data indicate that sequestration of COPII/TRAPP onto SGs halts ER-to-Golgi trafficking while removal of the stress releases COPII/TRAPP and allows trafficking to resume.

COPII and TRAPP not only control ER export but are also needed to maintain the organization of the GC. In particular, the TRAPP complex acts as GEF for Rab1, a GTPase with a key role in the organization and function of the GC (Tisdale et al., 1992; Wilson et al., 1994). We monitored Golgi complex morphology in response to SA by following the distribution of the early Golgi marker GM130. Strikingly, the GC starts to fragment after 30 min and completely redistributes throughout the

cytoplasm after 60 min of SA treatment (Figure 9A). This time window overlaps with the progressive recruitment of the TRAPP complex onto SGs (Figure 1D). To assess whether the Golgi fragmentation induced by oxidative stress was due to sequestration of TRAPP/COPII onto SGs, we analyzed the morphology of the GC in cells exposed to oxidative stress under conditions that prevent COPII/TRAPP recruitment to SGs, i.e. treatment with CDK inhibitors, highly confluent cells or cells treated with ISRIB. We found that CDK inhibitors prevented the fragmentation of the GC in cells exposed to oxidative stress (Figure 9B,C) and that oxidative stress had no impact on the organization of the Golgi complex in highly confluent cells (Figure 8C). Finally, ISRIB was effective in preventing the fragmentation of the Golgi complex induced by SA (Figure 8D).

These results indicated a causal link between TRAPP/COPII sequestration to SGs and Golgi fragmentation. To gain insight into the nature of the fragmentation of the Golgi complex, we performed ultrastructural analysis that revealed the presence of residual swollen cisternae with the loss of stacked structures and extensive vesiculation in SA-treated cells (Figure 9E), a situation that is reminiscent of the effect of Rab1 inactivation (Wilson et al., 1994). We hypothesized that the dismantling of the Golgi complex could have been due to impaired Rab1 activation as a consequence of TRAPP (the Rab1 GEF) sequestering to SGs. We monitored Rab1 activity by using an antibody that specifically recognizes the GTP-bound Rab1 and found, indeed, that the fraction of active Rab1 (i.e. the GTP-bound form) is progressively reduced in cells exposed to SA (Figure 9F). Overexpression of wt Rab1 reduced fragmentation/vesiculation of the GC induced by oxidative stress without affecting the capacity of TRAPP to migrate to SGs (Figure 9G, Figure 9—figure supplement 1), establishing a causal a link between Rab1 inactivation and Golgi fragmentation. It is worth mentioning that a further contribution to the disorganization of the Golgi complex upon oxidative stress may derive from the delocalization of the TGN-located PARP12 to SGs, which, however, affects mainly late Golgi compartments (Catara et al., 2017).

Discussion

417

418 We described a branch of the integrated stress response that is dependent on the

419 assembly of SGs and that impinges on the early secretory pathway. A key player of

420 this branch is TRAPP, a multimolecular complex that intervenes in multiple

421 membrane trafficking steps.

422

423 We showed that TRAPP is massively recruited to SGs in response to different acute

424 stress stimuli and in turn recruits components of the inner layer of the COPII

425 complex Sec23-Sec24, the protein complex that drives the export of newly

426 synthesized proteins from the ER, to SGs. The functional consequences of the

427 recruitment of TRAPP/COPII to SGs are double-edged since on the one hand they

428 impact on the function and organization of the secretory pathway while on the other

429 they impact on the composition and function of the SGs (Figure 10).

430 We have shown that the sequestering to SGs of these two complexes, one, COPII,

431 with a pivotal role in ER export and another, TRAPP, acting as a GEF for Rab1, a

432 master GTPase orchestrating the organization of the Golgi complex, induces a slow-

433 down of the ER-to-Golgi transport of newly synthesized proteins and the

434 disorganization of the structure of the Golgi complex. This halt in secretory activity,

435 an energetically costly process, can be seen as a way to limit energy expenditure

436 upon acute stress but also to prevent the ER export of newly synthesized proteins

437 that might have been damaged/misfolded by the acute stress before an adequate

438 UPR is installed and becomes operative. Importantly, these changes are completely

439 reversible since ER export resumes and proper Golgi morphology is restored upon

440 stress removal and relocation of the two complexes to their natural sites.

441 An intriguing feature of this branch of the integrated stress response that involves

442 the secretory pathway is its strict dependence on the proliferation status of the cells.

443 Indeed, the sequestration of TRAPP and COPII to SGs occurs only when the acute

444 stress affects actively proliferating cells. Cells with a slow/halted proliferation rate

445 still form SGs in response to acute stress stimuli, but do not delocalize TRAPP/COPII

446 from their steady state sites, i.e., the ERES, to SGs. We have used multiple strategies

447 to slow down (or halt) proliferation: nutrient deprivation, a high cell density, or a

448 switch from a proliferation to a differentiation program. Under all circumstances, we

observed a very tight correlation between the extent of recruitment of TRAPP/COPII

to SGs and the proliferation rate, indicating that acquisition of components of the

secretory machinery does not occur by default as a consequence of the assembly of

the SGs but is an active process that is subject to specific regulation.

We found that this regulation involves the activity of CDK1&2, since CDK1&2

inhibitors and/or CDK1&2 downregulation prevented the relocation of TRAPP/COPII

to SGs by stabilizing their association with the ERES. Indeed, the function of the

ERES and some of the components of the COPII coat are known to be controlled by

CDKs. A member of the CDK family, PCTAIRE, is recruited to ERES via its interaction

with Sec23 and controls ER export (Palmer et al., 2005), while Sec31, a component

and a regulator of Sar1/COPII cycle, is a substrate for CDKs in yeast and in

Trypanosoma, and, as we have shown here, also in mammalian cells (Figure 7C,D).

Further, the phosphorylation of Sec31 in response to growth factor and mitogenic

signals has also been reported (Dephoure et al. 2008; Olsen et al., 2006). We have

shown here that, consistent with the described role of Sec31 as a co-GAP for Sar1,

Sec31 function is required to control the COPII cycle at the ERES, since Sec31

depletion induces a tighter association of Sec23-24 with ERES and prevents their

translocation to SGs.

In fact, our results indicate that the extent of translocation of COPII/TRAPP to SGs is

directly proportional to their cycling rate at the ERES membranes. We have shown

that this rate is diminished by CDK inhibitors, but also, importantly, in starved cells as

compared to actively proliferating cells. These observations are in line with previous

reports showing that the cycling rate of ERES components is under control of growth

factors and growth factor-dependent signaling (Farhan et al., 2010; Tillmann et al.,

2015). Thus, the activity of the ERES, as measured by their number and by the rate of

cycling of COPII and other ERES components, is stimulated under conditions that

demand maximal output efficiency from the ER (i.e. during cell proliferation) as they

involve cell size increase and organelle expansion. If cells experience an acute stress

under these conditions of active growth, then the fast cycling COPII components are

susceptible to being temporarily sequestered to the SGs to instantly slow down ER

output and reduce energy consumption. By contrast, conditions that are not

permissive for cell proliferation, such as nutrient starvation, stabilize the association

of COPII/TRAPP with ERES membranes and prevent their translocation to SGs. Indeed, a profound remodeling of ERES is known to occur during nutrient starvation and coincides with the rerouting of COPII and TRAPP from their role in mediating the ER export of newly synthesized proteins to their role in autophagosome formation (Ge et al., 2017; Imai et al., 2016; Kim et al., 2016; Lamb et al., 2016; Ramírez-Peinado et al., 2017; van Leeuwen et al., 2018). The kinetics of the COPII cycling at the sites of phagosome formation in the ER have not been specifically explored. It is tempting to speculate that it might be slower than that at the "conventional" ERES under feeding conditions and that this more stable association of COPII with the ER during autophagosome formation may in turn prevent its sequestration to SGs upon stress exposure. Preventing the sequestration of TRAPP/COPII in starved cells exposed to stress would preserve the function of COPII and TRAPP in the autophagy process, a process that leads to nutrient recycling and energy production, two desirable events in cells exposed to stress.

Finally, we have shown that TRAPP is required for the proper maturation of SGs. Impeding the recruitment of TRAPP to SGs or depleting TRAPP does not impair the formation of SGs *per se* but hampers their maturation, as evaluated by their size (smaller SGs in the absence of TRAPP) and composition. In particular, key signaling components, such as RACK1 and Raptor, are no longer recruited to SGs in TRAPP-depleted cells. The mechanisms underlying the TRAPP-dependent maturation of SGs remain to be identified. Two possible non-exclusive scenarios are worthy of consideration, the engagement of specific protein-protein interactions between TRAPP components and signaling molecules or a central role of TRAPP, originally identified as a tethering complex, in mediating tethering/coalescence of smaller SGs into larger ones, which might be the only ones competent for recruiting signaling molecules. Whatever the underlying mechanisms, the impaired recruitment to SGs of RACK1 and Raptor in TRAPP-depleted cells leads to lower resistance to stress and higher tendency to undergo apoptosis, as the association of these signaling elements with SGs exerts an anti-apoptotic role (Arimoto et al., 2008; Thedieck et al., 2013; Wippich et al., 2013).

512 Our finding that TRAPP, and in particular TRAPPC2, acts as a mediator of the

513 secretory arrest in response to stress and as a driver of SG maturation elicits the

514 question as to whether and how these unsuspected properties of TRAPPC2 may be

515 relevant for the pathogenesis of SEDT, which is caused by mutations in TRAPPC2

516 (Gedeon et al., 1999). This, as well as the more general question on the physiological

517 relevance of assembling SGs in response to stress (Protter and Parker, 2016), remain

518 outstanding questions for further investigation. For now, we can only speculatively

519 propose that, given the role of TRAPPC2 in mediating secretory arrest and SG

520 maturation in actively proliferating cells, the main target cells in SEDT are likely to be

521 proliferative chondrocytes. Derived from chondrocyte progenitor cells, rapidly

522 proliferating chondrocytes are key components of the epiphyseal growth plate and

523 undergo terminal differentiation into postmitotic hypertrophic chondrocytes, which

524 eventually transdifferentiate into osteoblasts or undergo apoptosis, being replaced

525 by osteoblasts and osteoclasts. It is established that the rate of proliferation,

526 together with the height of columnar hypertrophic chondrocytes, are major

527 determinant of growth plate function and thus of bone length (Lui et al., 2018). We

528 hypothesize that TRAPPC2-defective proliferative chondrocytes are less resistant to

529 repetitive stresses, including mechanical and oxidative stresses (Henrotin et al.,

530 2003; Zuscik et al., 2008), and are thus more prone to undergo apoptosis. The

531 accelerated loss of proliferative chondrocytes would thus lead to a premature

532 growth plate senescence that could manifest as progressive delayed growth.

533

534 **Supplementary Materials**

535

536 **Materials and methods**

537 **Reagents and antibodies**

538 Primary antibodies used in this study were: mouse monoclonal antibody anti-G3BP
539 (BD Transduction Laboratories cat. no. 611126), rabbit polyclonal antibody anti-G3BP
540 (Bethil, cat. no. A302-033), mouse monoclonal antibody anti-TRAPPC5 (Abnova cat.
541 no. H00126003-A01), goat polyclonal antibody anti-eIF3 (Santa Cruz cat. no. sc-
542 16377), rabbit polyclonal antibody anti-Sec24C (Sigma-Aldrich cat. no. HPA040196),
543 mouse monoclonal anti-TIAR (BD Transduction Laboratories cat. no. 610352), mouse
544 monoclonal TRAPPC12 (Abcam cat. no. 88751), mouse monoclonal TRAPPC10 (Sanza
545 Cruz cat. no. SC-101259) mouse monoclonal anti-flag (Sigma-Aldrich cat. no. F1804),
546 rabbit polyclonal antibody anti-phosphor-Ser CDK substrates motif (Cell signaling cat.
547 no. 9477), mouse monoclonal anti-GM130 (BD Transduction Laboratories cat. no.
548 610823), human monoclonal antibody anti-Rab1-GTP (Adipogen cat. no. AG-27B-
549 0006), mouse monoclonal anti-RACK1 (Sanza Cruz cat. no. 17754), rabbit polyclonal
550 anti-Raptor (Cell signaling cat.no. 2280), rabbit polyclonal Sec16A (Bethil, cat.no.
551 A300-648A), sheep polyclonal anti-TGN46 (Serotec cat. no. AHP500), rabbit
552 polyclonal antibody anti-Sec31A (Sigma-Aldrich cat. no. HPA005457), mouse
553 monoclonal antibody AP-II (Pierce, cat. no. MA1-064), mouse monoclonal   adaptin
554 (Transduction cat. no. 618305), mouse monoclonal antibody anti-Retinoblastoma
555 (Cell Signaling cat. no. 9309), rabbit polyclonal antibody anti-phospho
556 Retinoblastoma (Cell Signaling cat. no. 9516), rabbit polyclonal antibody anti-eiF2α
557 (Cell Signaling cat. no. 9722), rabbit polyclonal antibody anti-phospho-eiF2α (ser51)
558 (Cell Signaling cat. no. 3597), mouse monoclonal antibody anti-puromycin (Millipore
559 cat.no. MABE343), rabbit polyclonal anti-ß-actin (Sigma-Aldrich cat. no. A2066),
560 mouse monoclonal antibody anti-GAPDH (Santa Cruz cat. no. sc-32233). Guinea pig
561 polyclonal antibody anti-Synaptopodin (Acris cat. no. AP33487SU-N). Polyclonal
562 antibodies against Arfaptin, p115, G-97, COPI, TRAPPC2 and TRAPPC3 were obtained
563 in our labs. Media, serum and reagents for tissue culture were purchased from
564 Thermo Fisher Scientific.

Sodium arsenite (cat. no. S7400), Puromycin (cat.no. A1113802) and cycloheximide

(cat.no. C7698) were purchased from Sigma-Aldrich, Flavopirodol hydrochloride (cat.

no. S2679), SNS-032 (BMS-387032) (cat. no. S1145) and Dinaciclib (SCH727965) (cat.

no. S2768) were purchased from Sellekchem.

Plasmid construction

TRAPPC3-GFP, Sec23A-GFP, Sar1GTP-GFP, Rab1-GFP, OCRL-GFP, Rab5-GFP and Rab7-

GFP were obtained as previously described (De Leo et al., 2016; Venditti et al., 2012;

Vicinanza et al., 2011). Construct encoding GFP-ARF1 has been described (Dubois et

al., 2005).

Cell culture, transfection, and RNA interference

HeLa cells were grown in high glucose (4,500 mg/L) DMEM supplemented with 10%

FCS. U2OS were grown in McCoy's supplemented with 20% FCS. Human fibroblasts

were grown in DMEM and M199 (1:4) supplemented with 10% FCS. Podocytes were

grown in RPMI-1640 supplemented with 10% Insulin-transferrin-Selenium (ITS) and

10% FCS.

For transfection of DNA plasmids, HeLa cells were transfected using either TransIT-

LT1 (Mirus Bio LLC, for BioID2 experiment) or JetPEI (Polyplus, for

immunofluorescence analysis) as transfection reagents, and the expression was

maintained for 16 hr before processing. Microinjection of TRAPPC3 antibody was

performed as described (Venditti et al., 2012).

siRNA sequences used in this study are listed in table S2. HeLa cells were treated for

72 hr with Oligofectamine (Life Technologies) for direct transfection.

Knock-down efficiency was evaluated by western blot for TRAPPC2, TRAPPC3 and

Sec31 (figure 2-Supplementary 3C and figure 3-Supplementary 1) and by q-PCR for

Sec16, Sec23 and sec24 isoforms, CDK1 and CDK2 (figure 2-Supplementary 2; figure

2-Supplementary 3A,B; figure 4 Supplementary 1).

Stress stimuli

Mammalian cells were treated with 300 µM SA for 30 min (unless otherwise stated)

in DMEM 10% FBS supplemented with 20mM Hepes. Temperature was maintained

597 at 37°C for the time of treatment in water bath. Heat treatment (44°C) was for 45

598 min in water bath.

599

600 **Immunofluorescence microscopy**

601 HeLa, human fibroblast, U2OS and rat chondrosarcoma cells were grown on

602 coverslips and fixed with 4% PFA for 10 min, washed three times with PBS, blocked

603 and permeabilized for 5 min in 0.1% Triton in PBS. Samples were washed three times

604 and blocked with blocking solution (0.05% saponin, 0.5% BSA, 50 mM NH_4Cl in PBS)

605 and incubated with primary antibodies diluted in blocking solution for 1 hr at

606 RT. Coverslips were washed with PBS and incubated with fluorochrome-conjugated

607 secondary antibodies (Alexa-Fluor-488, Alexa-Fluor-568 and Alexa-Fluor-633) for 1 hr

608 at RT. Fixed cells were mounted in Mowiol and imaged with Plan-Apochromat

609 63x/1.4 oil objective on a Zeiss LSM800 confocal system equipped with an ESID

610 detector and controlled by a Zen blue software. Fluorescence images presented are

611 representative of cells imaged in at least three independent experiments and were

612 processed with Fiji (ImageJ; National Institutes of Health) software. Sec24C and

613 TRAPPC2 quantification and Golgi particle analysis are described in the

614 corresponding figure legends. Differences among groups were performed using the

615 unpaired Student's *t* test calculated with the GraphPad Prism software. All data are

616 reported as mean ± s.e.m. as indicated in the figure legends. For Structured

617 Illumination Microscopy (SIM, Figure 7D, figure7-supplementary 1C), cells were

618 imaged with a Plan-Apochromat 63x/1.4 oil objective on a Zeiss LSM880 confocal

619 system. Image stacks of 0.88 μm thickness were acquired with 0.126 μm z-steps and

620 15 images (three angles and five phases per angle) per z-section. The field of view

621 (FOV) was ~ 75 x 75 μm at 0.065 μm/pixel, and a SIM grating of 34 μm was used.

622 Image stacks were processed and reconstructed with Zen software, using the SIM

623 tool in the Processing palette.

624

625
626 **Gel filtration**

The lysate used for gel filtration was prepared by cell cracker homogenizing 1.8×10^8

cells in 2 ml of buffer 20 mM HEPES, pH 7.2, 150 mM NaCl, 2 mM EDTA, 1 mM DTT

and Protease inhibitor (Roche). The sample was centrifuged at 100,000xg for 1 h. 10

mg of protein was concentrated in 350μL and loaded onto a Superose6 gel filtration

column (GE) and 400 μL of each fraction was collected. 50μL of fractions were

processed for SDS-PAGE analysis and proteins were detected by Western blot using

specific antibodies as described in Figure 1E.

Yeast Methods

The centromeric plasmid pUG23-Bet3-GFP (His selection) was described previously

(Mahfouz et al., 2012). A Pab1-mRFP expressing plasmid was constructed by PCR

amplification of the Pab1 gene plus 419 bp of the promoter region from yeast

genomic DNA and cloning into the vector backbone of the centromeric plasmid

pUG35 (Ura selection), removing the Met25 promoter and replacing GFP with mRFP,

using standard cloning techniques. Yeast cells (BY4743) were transformed with both

plasmids and grown in –Ura-His media with 2% Dextrose at 30°C to early log phase.

Stress granules were induced by incubating cells at 46°C for 10 min (Riback et al.,

2017) and observed immediately on an LSM800 microscope.

Immunoprecipitation

For the studies of TRAPPC2 interactors, HeLa cells were transfected with the

TRAPPC2-3XFLAG and 3XFLAG constructs. 16 hr post-transfection, cells were lysed

with lysis buffer (50 mM Tris-HCl pH 7.4, 100 mM NaCl, 1 mM EDTA, 0.5% Lauryl

Maltoside, protease and phosphatase inhibitors).

Cell extracts from control transfected HeLa cells or cells expressing TRAPPC2-3XFLAG

were immune-precipitated for 5 hr at 4°C using M2-flag agarose beads. Immune-

precipitates were analysed by Western blot or by LC-MS/MS (Central Proteomics

Facility, Sir William Dunn Pathology School, Oxford University).

HeLa cells or podocytes were used to analyse Sec31 phosphorylation status. IP of

HeLa cells: cells were plated in 15 mm plates. The day after, samples were treated

with Dinaciclib (10 μM in DMSO) for 150 min and with SA (300 μM, 10 min) alone or

in combination. Cells were then lysed with RIPA buffer (20mM Tris-HCl pH 7.5, 150

mM NaCl, 1 mM Na2EDTA, 1 mM EGTA, 1% NP-40, 1% sodium deoxycholate,

protease and phosphatase inhibitors). Cells extracts were immune-precipitated for

30 min at 4°C using the anti-Sec31A antibody. Finally, protein-A Sepharose beads

were added to the samples and left for 45 min at 4°C. Samples were analyzed by

Western blot using the antibodies indicated in the main text. IP of podocytes:

growing and differentiated podocytes were lysed and processed as described for

HeLa cells.

Functional annotation of the TRAPPC2 interactors

TRAPPC2-FLAG IP samples were compared with the proteins immunoprecipitated in

the control IP (Table S1): 427 proteins were specifically immunoprecipitated in

TRAPPC2-FLAG samples. Moreover, 113 were enriched in the TRAPPC2-FLAG IP

samples versus the control IP (ratio >2). Functional Annotation analysis (Dennis et al.,

2003; Huang et al., 2009) was used to group the putative TRAPPC2 protein partners

according to Molecular Function (MF). The DAVID online tool (DAVID Bioinformatics

Resources 6.7) was used restricting the output to all MF terms (MF_FAT). Of note,

we found that 251 out of the total 539 proteins were annotated as RNA-binding

proteins (GO:0003723) (Ashburner et al., 2000).

Digitonin Assay

HeLa cells were exposed to SA or treated with Dinaciclib (10 μM). After treatment,

cells were washed twice with permeabilization buffer (25mM HEPES, 125 mM

CH_3COOK, 2.5 mM $Mg(CH_3COO)_2$, 5 mM EGTA, 1mM DTT) and then permeabilized in

PB buffer supplemented with 30 μg/mL of digitonin plus the indicated antibody to

visualize permeabilized cells. To analyze Sec24C association with SGs, digitonin was

left for 6 min at room temperature. Cells were then washed three times with PB

buffer and left in PB buffer for 10 min at RT. Finally, samples were fixed with 4% PFA

and processed for IF. To analyse Sec24C association at ERES in the presence of

Dinaciclib, cells were fixed for 2 min after digitonin permeabilization.

High content screening

HeLa cells were plated in 384-well culture plates. The day after, a kinase inhibitor
library, purchased from Selleckchem, was dispensed using a liquid-handler system
(Hamilton) to give a final concentration of 10 μM. 150 minutes after drug
administration, cells were treated with sodium arsenite (300 μM) for 30 min at 37°C
in the presence of compounds. Finally, samples were fixed for 10 min at room
temperature by adding 1 volume of 4% PFA (paraformaldehyde in PBS) to the growth
medium and stained with the appropriate antibodies.

For image acquisition, at least 25 images per field were acquired per well of the 384-
well plate using confocal automated microscopy (Opera high content system; Perkin-
Elmer). A dedicated script was developed to perform the analysis of Sec24C
localization on the different images (Harmony and Acapella software; Perkin-Elmer).
The script calculated the co-localization value of Sec24C with the SG marker (G3BP).
The results were normalized using positive (mock cells exposed to SA) control
samples in the same plate.

P-values were calculated on the basis of mean values from three independent wells.
The data are represented as a percentage of Sec24C recruitment in the control cells
(100%) using Excel (Microsoft) and Prism software (GraphPad software).

Cell cycle analysis by flow cytometry

HeLa cells were harvested and resuspended in PBS. For the fixation, a 9-fold volume
of 70% ethanol was added and incubated a 4°C for at least 1hr. Next, cells were
centrifuged, washed in PBS and resuspended in PBS containing RNase A 0.1 mg/ml.
After incubation for 1hr at 37°C, propidium iodide was added to a final concentration
of 10 μg/ml and samples were analyzed in an Accuri C6 flow cytometer.

Cell proliferation analysis by High Content Imaging

Cell cycle analysis by high content imaging was performed using the Click-iT Plus EdU
Alexa Fluor 488 Imaging Kit (Life Technologies) according to the manufacturer's
instructions. Images were acquired with an automated confocal microscopy (Opera
System, Perkin-Elmer) and analyzed through Columbus Image Data Storage and
Analysis System (Perkin-Elmer). Nuclear intensity of EdU (5-ethynyl-2'-deoxyuridine,

a nucleoside analog of thymidine) in EdU-positive nuclei (S-phase cells) was used as a

measure of DNA replication rate.

Cell death assay

HeLa cells were pre-treated with CDK inhibitors (SNS-032, Flavopiridol and Dinaciclib)

for 150 min and exposed to SA (500 µM, 1 h). Subsequently, the stress stimulus was

washed out and cells left to recover for 16 h. To stain dead cells, a fluorophore dye

cocktail containing the cell-permeant nuclear dye Hoechst 33342 and the cell-

impermeant nuclear dye BOBO ™-3 that selectively labels dying cells (Invitrogen,

H3570, B3586), was added to the growth media and incubated for 45 min (37°C, 5%

CO_2). Cells were analyzed at automated microscopy (Operetta high content system;

Perkin-Elmer) using a customized script to calculate the ratio between BOBO ™-3

positive nuclei and total cells.

Puromycin Assay

The puromycin (PMY) assay was modified from David et al., JCB 2012 (David et al.,

2012). In brief, mock, TRAPPC2-KD or TRAPPC3-KD HeLa cells were exposed to SA

(500 µM, 30 min) in DMEM 10% FCS. Cells were washed three times in DMEM 1X

and incubated with 9 µM PMY in DMEM for 5 min at 37°C. Samples were lysed in

RIPA buffer and processed for Western blot analysis with the anti-puromycin

antibody.

Transport assay

VSVG-mEOS2-2XUVR8 was a gift from Matthew Kennedy (AddGene plasmid #49803).

HeLa cells were transfected with the plasmid for 16 h and treated with SA, CHX and

ISRIB for the indicated times. A UV-A lamp was used to illuminate samples (4 pulses,

15 sec each). After the light pulses, cells were left for 10 min at 37°C, then fixed with

a volume of 4% PFA and processed for immunofluorescence.

The PC-I transport assay was performed in human fibroblasts as previously described

(Venditti et al., 2012). For our purposes, cells were treated with SA (300 µM) for 120

min at 40°C and analyzed 10 min after the temperature switch (40 to 32°C). Cells

were then fixed and stained with appropriate antibodies.

752

753 **Electron microscopy**

754 EM samples were prepared as previously described (D'Angelo et al., 2007). Briefly,

755 cells were fixed by adding to the culture medium the same volume of a mixture of

756 PHEM buffer (10 mM EGTA, 2 mM MgCl$_2$, 60 mM PIPES, 25 mM HEPES, pH 6.9), 4%

757 paraformaldehyde, 2% glutaraldehyde for 2 h, and then stored in storage solution

758 (PHEM buffer, 0.5% paraformaldehyde) overnight. After washing with 0.15 M glycine

759 buffer in PBS, the cells were scraped and pelleted by centrifugation, embedded in

760 10% gelatin, cooled on ice, and cut into 0.5 mm blocks. The blocks were infused with

761 2.3 M sucrose and trimmed with an ultramicrotome (Leica Ultracut R) at −120°C

762 using a dry diamond knife.

763

764 **Quantitative real-time PCR**

765 Real-time quantitative PCR (qRT-PCR) was carried out with the LightCycler 480 SYBR

766 Green I mix (Roche) using the Light Cycler 480 II detection system (Roche) with the

767 following conditions: 95 °C, 10 min; (95 °C, 10 s; 60 °C, 10 s; 72 °C, 15 s) × 45 cycles.

768 For expression studies the qRT-PCR results were normalized against an internal

769 control (HPRT1).

770

771

References

Anderson P, Kedersha N. 2008. Stress granules: the Tao of RNA triage. *Trends Biochem Sci* 33:141–150. doi:10.1016/j.tibs.2007.12.003

Anderson P, Kedersha N. 2002. Stressful initiations. *J Cell Sci* 115:3227–3234.

Arimoto K, Fukuda H, Imajoh-Ohmi S, Saito H, Takekawa M. 2008. Formation of stress granules inhibits apoptosis by suppressing stress-responsive MAPK pathways. *Nat Cell Biol* 10:1324–1332. doi:10.1038/ncb1791

Asghar U, Witkiewicz AK, Turner NC, Knudsen ES. 2015. The history and future of targeting cyclin-dependent kinases in cancer therapy. *Nat Rev Drug Discov* 14:130–146. doi:10.1038/nrd4504

Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, Harris MA, Hill DP, Issel-Tarver L, Kasarskis A, Lewis S, Matese JC, Richardson JE, Ringwald M, Rubin GM, Sherlock G. 2000. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet* 25:25–29. doi:10.1038/75556

Aulas A, Fay MM, Szaflarski W, Kedersha N, Anderson P, Ivanov P. 2017. Methods to Classify Cytoplasmic Foci as Mammalian Stress Granules. *J Vis Exp JoVE*. doi:10.3791/55656

Ballock RT, O'keefe RJ. 2003. The Biology Of The Growth Plate. *J Bone Jt Surg-Am Vol* 85:715–726.

Bi X, Mancias JD, Goldberg J. 2007. Insights into COPII Coat Nucleation from the Structure of Sec23•Sar1 Complexed with the Active Fragment of Sec31. *Dev Cell* 13:635–645. doi:10.1016/j.devcel.2007.10.006

Catara G, Grimaldi G, Schembri L, Spano D, Turacchio G, Lo Monte M, Beccari AR, Valente C, Corda D. 2017. PARP1-produced poly-ADP-ribose causes the PARP12 translocation to stress granules and impairment of Golgi complex functions. *Sci Rep* 7:14035. doi:10.1038/s41598-017-14156-8

Chen D, Gibson ES, Kennedy MJ. 2013. A light-triggered protein secretion system. *J Cell Biol* 201:631–640. doi:10.1083/jcb.201210119

David A, Dolan BP, Hickman HD, Knowlton JJ, Clavarino G, Pierre P, Bennink JR, Yewdell JW. 2012. Nuclear translation visualized by ribosome-bound nascent chain puromycylation. *J Cell Biol* 197:45–57. doi:10.1083/jcb.201112145

D'Arcangelo JG, Stahmer KR, Miller EA. 2013. Vesicle-mediated export from the ER: COPII coat function and regulation. *Biochim Biophys Acta* 1833:2464–2472. doi:10.1016/j.bbamcr.2013.02.003

De Leo MG, Staiano L, Vicinanza M, Luciani A, Carissimo A, Mutarelli M, Di Campli A,

819 Polishchuk E, Di Tullio G, Morra V, Levtchenko E, Oltrabella F, Starborg T, Santoro M,
820 Di Bernardo D, Devuyst O, Lowe M, Medina DL, Ballabio A, De Matteis MA. 2016.
821 Autophagosome-lysosome fusion triggers a lysosomal response mediated by TLR9
822 and controlled by OCRL. *Nat Cell Biol* 18:839–850. doi:10.1038/ncb3386
823
824 Dennis G, Sherman BT, Hosack DA, Yang J, Gao W, Lane HC, Lempicki RA. 2003.
825 DAVID: Database for Annotation, Visualization, and Integrated Discovery. *Genome*
826 *Biol* 4:P3.
827
828 Dephoure N, Zhou C, Villén J, Beausoleil SA, Bakalarski CE, Elledge SJ, Gygi SP. 2008 A
829 quantitative atlas of mitotic phosphorylation. *Proc Natl Acad Sci U S A.*
830 105(31):10762-7
831
832 Dubois T, Paléotti O, Mironov AA, Fraisier V, Stradal TEB, De Matteis MA, Franco M,
833 Chavrier P. 2005. Golgi-localized GAP for Cdc42 functions downstream of ARF1 to
834 control Arp2/3 complex and F-actin dynamics. *Nat Cell Biol* 7:353–364.
835 doi:10.1038/ncb1244
836
837 Farhan H, Wendeler MW, Mitrovic S, Fava E, Silberberg Y, Sharan R, Zerial M, Hauri
838 H-P. 2010. MAPK signaling to the early secretory pathway revealed by
839 kinase/phosphatase functional screening. *J Cell Biol* 189:997–1011.
840 doi:10.1083/jcb.200912082
841
842 Franzmann TM, Alberti S. 2019. Protein Phase Separation as a Stress Survival
843 Strategy. *Cold Spring Harb Perspect Biol*. doi:10.1101/cshperspect.a034058
844
845 Ge L, Zhang M, Kenny SJ, Liu D, Maeda M, Saito K, Mathur A, Xu K, Schekman R. 2017.
846 Remodeling of ER-exit sites initiates a membrane supply pathway for
847 autophagosome biogenesis. *EMBO Rep* 18:1586–1603.
848 doi:10.15252/embr.201744559
849
850 Gedeon AK, Colley A, Jamieson R, Thompson EM, Rogers J, Sillence D, Tiller GE,
851 Mulley JC, Gécz J. 1999. Identification of the gene (SEDL) causing X-linked
852 spondyloepiphyseal dysplasia tarda. *Nat Genet* 22:400–404. doi:10.1038/11976
853
854 Gorur A, Yuan L, Kenny SJ, Baba S, Xu K, Schekman R. 2017. COPII-coated membranes
855 function as transport carriers of intracellular procollagen I. *J Cell Biol* 216:1745–1759.
856 doi:10.1083/jcb.201702135
857
Harripaul R, Vasli N, Mikhailov A, Rafiq MA, Mittal K, Windpassinger C, Sheikh TI,
Noor A, Mahmood H, Downey S, Johnson M, Vleuten K, Bell L, Ilyas M, Khan FS, Khan
V, Moradi M, Ayaz M, Naeem F, Heidari A, Ahmed I, Ghadami S, Agha Z, Zeinali S,
Qamar R, Mozhdehipanah H, John P, Mir A, Ansar M, French L, Ayub M, Vincent JB.
2018. Mapping autosomal recessive intellectual disability: combined microarray and
exome sequencing identifies 26 novel candidate genes in 192 consanguineous
families. *Mol Psychiatry* 23:973–984. doi:10.1038/mp.2017.60
865

866   Henrotin YE, Bruckner P, Pujol J-PL. 2003. The role of reactive oxygen species in
867   homeostasis and degradation of cartilage. *Osteoarthritis Cartilage* 11:747–755.
868
869   Holt LJ, Tuch BB, Villén J, Johnson AD, Gygi SP, Morgan DO. 2009. Global analysis of
870   Cdk1 substrate phosphorylation sites provides insights into evolution. *Science*
871   325:1682–1686. doi:10.1126/science.1172867
872
873   Hu H, Gourguechon S, Wang CC, Li Z. 2016. The G1 Cyclin-dependent Kinase CRK1 in
874   Trypanosoma brucei Regulates Anterograde Protein Transport by Phosphorylating
875   the COPII Subunit Sec31. *J Biol Chem* 291:15527–15539.
876   doi:10.1074/jbc.M116.715185
877
878   Huang DW, Sherman BT, Zheng X, Yang J, Imamichi T, Stephens R, Lempicki RA. 2009.
879   Extracting biological meaning from large gene lists with DAVID. *Curr Protoc*
880   *Bioinforma* Chapter 13:Unit 13.11. doi:10.1002/0471250953.bi1311s27
881
882   Imai K, Hao F, Fujita N, Tsuji Y, Oe Y, Araki Y, Hamasaki M, Noda T, Yoshimori T. 2016.
883   Atg9A trafficking through the recycling endosomes is required for autophagosome
884   formation. *J Cell Sci* 129:3781–3791. doi:10.1242/jcs.196196
885
886   Jain S, Wheeler JR, Walters RW, Agrawal A, Barsic A, Parker R. 2016. ATPase
887   modulated stress granules contain a diverse proteome and substructure. *Cell*
888   164:487–498. doi:10.1016/j.cell.2015.12.038
889
890   Kapetanovich L, Baughman C, Lee TH. 2005. Nm23H2 Facilitates Coat Protein
891   Complex II Assembly and Endoplasmic Reticulum Export in Mammalian Cells. *Mol*
892   *Biol Cell* 16:835–848. doi:10.1091/mbc.E04-09-0785
893
894   Khattak NA, Mir A. 2014. Computational analysis of TRAPPC9: candidate gene for
895   autosomal recessive non-syndromic mental retardation. *CNS Neurol Disord Drug*
896   *Targets* 13:699–711.
897
898   Kim JJ, Lipatova Z, Segev N. 2016. TRAPP Complexes in Secretion and Autophagy.
899   *Front Cell Dev Biol* 4:20. doi:10.3389/fcell.2016.00020
900
901   Koehler K, Milev MP, Prematilake K, Reschke F, Kutzner S, Jühlen R, Landgraf D, Utine
902   E, Hazan F, Diniz G, Schuelke M, Huebner A, Sacher M. 2017. A novel TRAPPC11
903   mutation in two Turkish families associated with cerebral atrophy, global retardation,
904   scoliosis, achalasia and alacrima. *J Med Genet* 54:176–185. doi:10.1136/jmedgenet-
905   2016-104108
906
907   Koreishi M, Yu S, Oda M, Honjo Y, Satoh A. 2013. CK2 Phosphorylates Sec31 and
908   Regulates ER-To-Golgi Trafficking. *PloS One* 8:e54382.
909   doi:10.1371/journal.pone.0054382
910
Lamb CA, Nühlen S, Judith D, Frith D, Snijders AP, Behrends C, Tooze SA. 2016.
TBC1D14 regulates autophagy via the TRAPP complex and ATG9 traffic. *EMBO J*

35:281–301. doi:10.15252/embj.201592695

Li C, Luo X, Zhao S, Siu GK, Liang Y, Chan HC, Satoh A, Yu SS. 2017. COPI-TRAPPII
activates Rab18 and regulates its lipid droplet association. *EMBO J* 36:441–457.
doi:10.15252/embj.201694866

Lord C, Bhandari D, Menon S, Ghassemian M, Nycz D, Hay J, Ghosh P, Ferro-Novick S.
2011. Sequential interactions with Sec23 control the direction of vesicle traffic.
Nature 473:181–186. doi:10.1038/nature09969

Lui JC, Jee YH, Garrison P, Iben JR, Yue S, Ad M, Nguyen Q, Kikani B, Wakabayashi Y,
Baron J. 2018. Differential aging of growth plate cartilage underlies differences in
bone length and thus helps determine skeletal proportions. *PLoS Biol* 16:e2005263.
doi:10.1371/journal.pbio.2005263

Malumbres M. 2014. Cyclin-dependent kinases. *Genome Biol* 15:122.
Milev MP, Graziano C, Karall D, Kuper WFE, Al-Deri N, Cordelli DM, Haack TB,
Danhauser K, Iuso A, Palombo F, Pippucci T, Prokisch H, Saint-Dic D, Seri M, Stanga D,
Cenacchi G, van Gassen KLI, Zschocke J, Fauth C, Mayr JA, Sacher M, van Hasselt PM.
2018. Bi-allelic mutations in TRAPPC2L result in a neurodevelopmental disorder and
have an impact on RAB11 in fibroblasts. *J Med Genet* 55:753–764.
doi:10.1136/jmedgenet-2018-105441

Milev MP, Grout ME, Saint-Dic D, Cheng Y-HH, Glass IA, Hale CJ, Hanna DS, Dorschner
MO, Prematilake K, Shaag A, Elpeleg O, Sacher M, Doherty D, Edvardson S. 2017.
Mutations in TRAPPC12 Manifest in Progressive Childhood Encephalopathy and Golgi
Dysfunction. *Am J Hum Genet* 101:291–299. doi:10.1016/j.ajhg.2017.07.006

Mironov AA, Beznoussenko GV, Nicoziani P, Martella O, Trucco A, Kweon H-S,
Giandomenico DD, Polishchuk RS, Fusella A, Lupetti P, Berger EG, Geerts WJC, Koster
AJ, Burger KNJ, Luini A. 2001. Small cargo proteins and large aggregates can traverse
the Golgi by a common mechanism without leaving the lumen of cisternae. *J Cell Biol*
155:1225–1238. doi:10.1083/jcb.200108073

Mohamoud HS, Ahmed S, Jelani M, Alrayes N, Childs K, Vadgama N, Almramhi MM,
Al-Aama JY, Goodbourn S, Nasir J. 2018. A missense mutation in TRAPPC6A leads to
build-up of the protein, in patients with a neurodevelopmental syndrome and
dysmorphic features. *Sci Rep* 8:2053. doi:10.1038/s41598-018-20658-w

Moujalled D, James JL, Yang S, Zhang K, Duncan C, Moujalled DM, Parker SJ,
Caragounis A, Lidgerwood G, Turner BJ, Atkin JD, Grubman A, Liddell JR,

Olsen JV[1], Blagoev B, Gnad F, Macek B, Kumar C, Mortensen P, Mann M 2006 Global,
in vivo, and site-specific phosphorylation dynamics in signaling networks
Cell. 2006;127(3):635-48

Proepper C, Boeckers TM, Kanninen KM, Blair I, Crouch PJ, White AR. 2015.

Phosphorylation of hnRNP K by cyclin-dependent kinase 2 controls cytosolic
accumulation of TDP-43. *Hum Mol Genet* 24:1655–1669. doi:10.1093/hmg/ddu578
962

963 Palmer KJ, Konkel JE, Stephens DJ. 2005. PCTAIRE protein kinases interact directly
964 with the COPII complex and modulate secretory cargo transport. *J Cell Sci* 118:3839–
965 3847. doi:10.1242/jcs.02496
966

967 Protter DSW, Parker R. 2016. Principles and Properties of Stress Granules. *Trends Cell*
968 *Biol* 26:668–679. doi:10.1016/j.tcb.2016.05.004
969

970 Ramírez-Peinado S, Ignashkova TI, van Raam BJ, Baumann J, Sennott EL, Gendarme
971 M, Lindemann RK, Starnbach MN, Reiling JH. 2017. TRAPPC13 modulates autophagy
972 and the response to Golgi stress. *J Cell Sci* 130:2251–2265. doi:10.1242/jcs.199521
973

974 Sacher M, Shahrzad N, Kamel H, Milev MP. 2018. TRAPPopathies: An emerging set of
975 disorders linked to variations in the genes encoding transport protein particle
976 (TRAPP)-associated proteins. *Traffic Cph Den*. doi:10.1111/tra.12615
977

978 Saleem MA, O'Hare MJ, Reiser J, Coward RJ, Inward CD, Farren T, Xing CY, Ni L,
979 Mathieson PW, Mundel P. 2002. A conditionally immortalized human podocyte cell
980 line demonstrating nephrin and podocin expression. *J Am Soc Nephrol JASN* 13:630–
981 638.
982

983 Saurus P, Kuusela S, Dumont V, Lehtonen E, Fogarty CL, Lassenius MI, Forsblom C,
984 Lehto M, Saleem MA, Groop P-H, Lehtonen S. 2016. Cyclin-dependent kinase 2
985 protects podocytes from apoptosis. *Sci Rep* 6:21664. doi:10.1038/srep21664
986

987 Scrivens PJ, Noueihed B, Shahrzad N, Hul S, Brunet S, Sacher M. 2011. C4orf41 and
988 TTC-15 are mammalian TRAPP components with a role at an early stage in ER-to-
989 Golgi trafficking. *Mol Biol Cell* 22:2083–2093. doi:10.1091/mbc.E10-11-0873
990

991 Sidrauski C, McGeachy AM, Ingolia NT, Walter P. 2015. The small molecule ISRIB
992 reverses the effects of eIF2α phosphorylation on translation and stress granule
993 assembly. *eLife* 4. doi:10.7554/eLife.05033
994

995 Takahashi M, Higuchi M, Matsuki H, Yoshita M, Ohsawa T, Oie M, Fujii M. 2013.
996 Stress Granules Inhibit Apoptosis by Reducing Reactive Oxygen Species Production.
997 *Mol Cell Biol* 33:815–829. doi:10.1128/MCB.00763-12
998

999 Thedieck K, Holzwarth B, Prentzell MT, Boehlke C, Kläsener K, Ruf S, Sonntag AG,
1000 Maerz L, Grellscheid S-N, Kremmer E, Nitschke R, Kuehn EW, Jonker JW, Groen AK,
1001 Reth M, Hall MN, Baumeister R. 2013. Inhibition of mTORC1 by Astrin and Stress
1002 Granules Prevents Apoptosis in Cancer Cells. *Cell* 154:859–874.
1003 doi:10.1016/j.cell.2013.07.031
1004

1005 Tiller GE, Hannig VL, Dozier D, Carrel L, Trevarthen KC, Wilcox WR, Mundlos S, Haines
1006 JL, Gedeon AK, Gecz J. 2001. A recurrent RNA-splicing mutation in the SEDL gene

1007 causes X-linked spondyloepiphyseal dysplasia tarda. *Am J Hum Genet* 68:1398–1407.
1008 doi:10.1086/320594
1009
1010 Tillmann KD, Reiterer V, Baschieri F, Hoffmann J, Millarte V, Hauser MA, Mazza A,
1011 Atias N, Legler DF, Sharan R, Weiss M, Farhan H. 2015. Regulation of Sec16 levels and
1012 dynamics links proliferation and secretion. *J Cell Sci* 128:670–682.
1013 doi:10.1242/jcs.157115
1014
1015 Tisdale EJ, Bourne JR, Khosravi-Far R, Der CJ, Balch WE. 1992. GTP-binding mutants
1016 of rab1 and rab2 are potent inhibitors of vesicular transport from the endoplasmic
1017 reticulum to the Golgi complex. *J Cell Biol* 119:749–761.
1018
1019 van Leeuwen W, van der Krift F, Rabouille C. 2018. Modulation of the secretory
1020 pathway by amino-acid starvation. *J Cell Biol* 217:2261–2271.
1021 doi:10.1083/jcb.201802003
1022
1023 Venditti R, Scanu T, Santoro M, Di Tullio G, Spaar A, Gaibisso R, Beznoussenko GV,
1024 Mironov AA, Mironov A, Zelante L, Piemontese MR, Notarangelo A, Malhotra V,
1025 Vertel BM, Wilson C, De Matteis MA. 2012. Sedlin controls the ER export of
1026 procollagen by regulating the Sar1 cycle. *Science* 337:1668–1672.
1027 doi:10.1126/science.1224947
1028
1029 Vicinanza M, Di Campli A, Polishchuk E, Santoro M, Di Tullio G, Godi A, Levtchenko E,
1030 De Leo MG, Polishchuk R, Sandoval L, Marzolo M-P, De Matteis MA. 2011. OCRL
1031 controls trafficking through early endosomes via PtdIns4,5P$_2$-dependent regulation
1032 of endosomal actin. *EMBO J* 30:4970–4985. doi:10.1038/emboj.2011.354
1033
1034 Westlake CJ, Baye LM, Nachury MV, Wright KJ, Ervin KE, Phu L, Chalouni C, Beck JS,
1035 Kirkpatrick DS, Slusarski DC, Sheffield VC, Scheller RH, Jackson PK. 2011. Primary cilia
1036 membrane assembly is initiated by Rab11 and transport protein particle II (TRAPPII)
1037 complex-dependent trafficking of Rabin8 to the centrosome. *Proc Natl Acad Sci U S A*
1038 108:2759–2764. doi:10.1073/pnas.1018823108
1039
1040 Wheeler JR, Matheny T, Jain S, Abrisch R, Parker R. n.d. Distinct stages in stress
1041 granule assembly and disassembly. *eLife* 5. doi:10.7554/eLife.18413
1042
1043 Wilson BS, Nuoffer C, Meinkoth JL, McCaffery M, Feramisco JR, Balch WE, Farquhar
1044 MG. 1994. A Rab1 mutant affecting guanine nucleotide exchange promotes
1045 disassembly of the Golgi apparatus. *J Cell Biol* 125:557–571.
1046
1047 Wippich F, Bodenmiller B, Trajkovska MG, Wanka S, Aebersold R, Pelkmans L. 2013.
1048 Dual specificity kinase DYRK3 couples stress granule condensation/dissolution to
1049 mTORC1 signaling. *Cell* 152:791–805. doi:10.1016/j.cell.2013.01.033
1050
Yamasaki A, Menon S, Yu S, Barrowman J, Meerloo T, Oorschot V, Klumperman J,
Satoh A, Ferro-Novick S. 2009. mTrs130 is a component of a mammalian TRAPPII
complex, a Rab1 GEF that binds to COPI-coated vesicles. *Mol Biol Cell* 20:4205–4215.

doi:10.1091/mbc.e09-05-0387

Yu S, Satoh A, Pypaert M, Mullen K, Hay JC, Ferro-Novick S. 2006. mBet3p is required
for homotypic COPII vesicle tethering in mammalian cells. *J Cell Biol* 174:359–368.
doi:10.1083/jcb.200603044

Zacharogianni M, Aguilera-Gomez A, Veenendaal T, Smout J, Rabouille C. n.d. A
stress assembly that confers cell viability by preserving ERES components during
amino-acid starvation. *eLife* 3. doi:10.7554/eLife.04132

Zou S, Liu Y, Zhang XQ, Chen Y, Ye M, Zhu X, Yang S, Lipatova Z, Liang Y, Segev N.
2012. Modular TRAPP complexes regulate intracellular protein trafficking through
multiple Ypt/Rab GTPases in Saccharomyces cerevisiae. *Genetics* 191:451–460.
doi:10.1534/genetics.112.139378

Zuscik MJ, Hilton MJ, Zhang X, Chen D, O'Keefe RJ. 2008. Regulation of
chondrogenesis and chondrocyte differentiation by stress. *J Clin Invest* 118:429–438.
doi:10.1172/JCI34174

Acknowledgements

We thank Andrea Ballabio, Carmine Settembre, Leandro Raul Soria, Maria Chiara
Masone, Rossella Venditti and Graciana Diez Roux for helpful discussion. MADM
acknowledges the support of Telethon (grant TGM11CB1), the Italian Association for
Cancer Research (AIRC, grant IG2013_14761), European Research Council Advanced
Investigator grant no. 670881 (SYSMET).

Author contributions

F.Z. and M.A.D.M. conceived the work. F.Z. planned and analyzed most of the
experiments, C.W. performed experiments on yeast, G.D.T. performed gel filtration
and Sec31-IP, M.S. provided technical support, P.P. and M.P. performed MS/MS
analysis, D.D. helped with cell culture, S.P.V. performed cell proliferation and flow
cytometry assays, R.D.C. and M.F. analyzed MS/MS data, A.R. helped in performing
cell death assay, M.A.S. provided podocytes cell line, E.P. performed E.M. analyses,
L.G. designed the script for high-content analysis, F.Z. and M.A.D.M. conceptualized
the work and strategy and wrote the manuscript.

1091 Conflict of interest

1092 The authors declare that they have no conflicts of interest with the contents of this
1093 article.

Figure 1. The TRAPP complex is recruited to stress granules. (A,B) Localization of endogenous TRAPPC2 under steady state conditions or after sodium arsenite (SA) treatment in (A) a rat chondrosarcoma cell line (300 µM, 1 hr) and (B) in HeLa cells (300 µM, 30 min). Fluorescence microscopy of fixed cells using antibodies against TRAPPC2, eIF3 (to label SGs), Sec31 (to label ERES). DAPI (blue). Small panels in (A) show details and merge of boxed areas. **(C)** Localization of endogenous TRAPPC1 and

of GFP-TRAPPC3 to eIF3-labeled SGs in HeLa cells after SA treatment. **(D)** Representative images of a time course analysis of TRAPPC2 redistribution to SGs. Cells were treated as in (B). Graph, quantification of TRAPPC2 localization at SGs over time (the ratio between TRAPPC2 (mean fluorescence intensity) in SG puncta and cytosolic TRAPPC2). Mean ± s.e.m. n = 50 cells per experiment, N = 3. Scale bar, 10 μm in (A-D) **(E)** Extracts from untreated and SA-treated HeLa cells were fractionated by size exclusion chromatography on a Superose6 column. Fractions were analyzed by Western blot using antibodies against TRAPPC2 and TRAPPC3 (components of both TRAPPII and TRAPPIII complexes), TRAPPC10 (a component of the TRAPPII complex), TRAPPC12 (a component of the TRAPPIII complex), and against the SG marker TIAR-1. TRAPP components co-fractionated with a high molecular weight complex (red dashed box) containing TIAR-1 in SA-treated cells. Arrows indicate protein standards: thyroglobulin (670 kDa), -globulin (158 kDa), ovalbumin(44 kDa) **F)** Schematic representation of MS/MS analysis of TRAPPC2 interactors. 251 out of 512 (46%) proteins are RNA-binding proteins, including several SG proteins (yellow ellipse).

Figure 1—figure supplement 1. Re-localization of TRAPP to SGs is independent of stress stimulus and cell type.

Figure 1—figure supplement 2. Stress-induced localization of Bet3 (TRAPPC3) to SGs in yeast cells.

Figure 1—supplement video 1. Time lapse fluorescent microscopy of Bet3 (TRAPPC3) and Pab1 at 30°C in yeast cells.

Figure 1—supplement video 2. Time lapse fluorescent microscopy of Bet3 (TRAPPC3) and Pab1 at 46°C in yeast cells.

Figure 1—figure supplement 3. Association of TRAPP with SGs is reversible after SG disassembly.

1129
Figure 2. TRAPPC2 is required for Sec23/24 relocalization to SGs. (A) HeLa cells,
untreated or treated with SA, were fixed and visualized by fluorescence microscopy
using anti-Sec24C Ab, anti-eIF3 Ab, and DAPI (blue). **(B)** Quantification of Sec24C
redistribution to SGs over time after SA treatment (the ratio between Sec24C (mean
fluorescence intensity) in SG puncta and cytosolic Sec24C). Mean ± s.e.m. n = 50 cells
per experiment, N = 3. **(C)** Cells treated as in **(A)** stained with an antibody recognizing
the coatomer I (COPI), which does not relocalize to SGs (see also Figure 2—figure
supplement 1A-N). **(D)** Representative images of TRAPPC2 localization in Sec23AB-
KD and Sec24ABCD-KD cells treated with SA. Cells were fixed and visualized by
fluorescence microscopy using anti-TRAPPC2 Ab, anti-G3BP Ab (to label SGs) and
DAPI (blue). **(E)** Quantification of TRAPPC2 redistribution to SGs after KD of the
indicated Sec23 and Sec24 combinations, calculated as the ratio between TRAPPC2
(mean intensity) in SG puncta and cytosolic TRAPPC2. Mean ± s.e.m., n = 40-60 cells
per experiment, N = 3. ns: not significant. **(F)** Representative images of Sec24C

localization at SGs (stained for G3BP) in TRAPPC3-KD and TRAPPC2-KD cells treated with SA. Depletion of TRAPP or TRAPPC2 reduces Sec24C recruitment. Graph, quantification of Sec24C at SGs, calculated as in (A). Mean ± s.e.m., n = 40-60 cells per experiment, N = 3. **** $P < 0.0001$.

Figure 2—figure supplement 1. Sec24C and Sec23, but not components of other coat complexes or other cytosolic proteins associated with the exocytic and endocytic pathways, associate with SGs.

Figure 2—figure supplement 2. Sec16 does not significantly associate with SGs nor is it required for the recruitment of COPII to SGs.

Figure 2—figure supplement 3. Evaluation of Sec23, Sec24, TRAPPC2 and TRAPPC3 knock down efficiency.

1155
1156 **Figure 3. Comparison and relationships between the ERES-associated and the SG-**
1157 **associated pools of TRAPP and COPII. (A)** TRAPP and COPII associate more stably
1158 with SGs than with ERES. The membrane association of Sec24C was evaluated in
1159 non-permeabilized or permeabilized cells with or without SA treatment, as indicated.
1160 G3BP was used as an SG marker. Left panel insets, G3BP staining in non-
1161 permeabilized cells. Dashed white lines show the outline of permeabilized SA-
1162 untreated cells. Blue, DAPI. **(B-C)** Stabilizing TRAPP and COPII at the ERES prevents

their relocalization to SGs. **(B)** HeLa cells overexpressing GFP-Sar1H79G were treated
with SA and immunostained for Sec24C, TRAPPC2, and eIF3, as indicated. Blue, DAPI.
Scale bar 10 μm. Graphs show quantification of Sec24C or TRAPPC2 at SGs in GFP-
Sar1H79G-expressing cells relative to non-transfected cells. Data are the ratio
between Sec24C or TRAPPC2 mean intensity in SG puncta and Sec24C or TRAPPC2
mean intensity at ERES, expressed as a percentage of the non-transfected cells.
Mean ± s.e.m. n = 60-70, three independent experiments ****p<0.0001. **(C)** Effect of
Sec31A depletion on TRAPPC2 and Sec24C recruitment to SGs. Cells were mock-
treated or KD for Sec31, treated with SA (300 μM, 30 min), and then immunostained
for TRAPPC2, Sec24C and eIF3 as indicated. Scale bar, 10 μm. Graph, quantification
of Sec24C and TRAPPC2 at SGs, calculated as in (B). Mean ± s.e.m. of TRAPPC2 and
Sec24C, one representative experiment; n=60-80. ** p<0.001, ****p<0.0001.
Figure 3—figure supplement 1. Effect of Sec31 depletion on Sec24C localization at
ERES.

Figure 4. The association of TRAPP with SGs is under control of CDK1/2. (A) Cells
were treated with 273 kinase inhibitors (10 μM) from the Sellekchem library and
with SA, and Sec24C localization in G3BP puncta (% of total G3BP area) was
calculated (see Materials and methods) and is reported as a percentage of the
control (cells treated with SA alone). Gray dashed line, mean of control; gray box,
control standard deviation (± 18.2%); dashed red line, threshold of positive hits. **(B)**
Enrichment Analysis of the positive hits; for each class of inhibitor, the enrichment

(in percentage) of compounds falling into positive hits and the enrichment (in percentage) in the total number of compounds was calculated and expressed as a ratio. **(C)** Evaluation of CDK kinase activity upon SA treatment. Western blot analysis of phosphoSer-CDK substrates in control, SA (300μM, 30 min), Dinaciclib (10 μM, 180 min) and SA+Dinaciclib (150 min Dinaciclib and 30 minutes SA). CDKs are hyperactivated upon oxidative stress and this activation is partially prevented upon CDK inhibition. Left, Western blot, right: Ponceau was used as loading control. Image, one representative experiments out of 3 independent replicates. **(D)** Drug specificity, from high (++++) to low (+). for the different CDKs as described by Selleckchem, and Sec24C recruitment to SGs as a percentage of control. **(E)** Analysis of Sec24C recruitment to SGs in control (MOCK), CDK1-KD, CDK2-KD and CDK1+2-KD HeLa cells. Data are expressed as percentage of Sec24C (mean intensity) in SG puncta normalized for cytosolic signal as a percentage of control conditions. Quantification of one representative experiment. N>100, mean ± s.e.m. ****$p<0.0001$. **(F)** Representative images of HeLa cells treated with the best three hits (Flavopiridol, SNS-032, Dinaciclib; 1μM 150 min) and then exposed to SA (300 μM, 30 min), followed by immunostaining for Sec24C and G3BP. Scale bar, 10μm. **(G)** HeLa cells were treated with Flavopiridol, SNS-032, or Dinaciclib for 150 min at the indicated concentrations, subsequently treated with SA (300 μM, 30 min), and then immunostained for Sec24C and G3BP and imaged by OPERA. BS-181, a specific CDK7 inhibitor, was used as negative control. Sec24C localization to SGs is expressed as a percentage of the control (cells treated with SA alone). Dashed red line, mean of control; red box, standard deviation of control (± 9.8%). **(H)** Cells were treated as in (G) and immunostained for TRAPPC2 and eIF3. Scale bar, 10 μm. The graph shows the quantification of TRAPPC2 localization in SG puncta (mean intensity). Data are mean ± s.e.m. expressed as a percentage of TRAPPC2 signal in SGs after Flavopiridol, SNS-032 or Dinaciclib treatment compared to the control (cells treated with SA alone). N=3, three independent experiments, n=60-80 cells per experiment. ****$p<0.0001$.

Figure 4—figure supplement 1. Evaluation of CDK1 and CDK2 knock down efficiency.

1216
Figure 5. CDK1/2 modulate the COPII cycle at ERES. (A) Immunofluorescence images
of Sec24C and TRAPPC2 in vehicle-treated and SNS-032-treated (1 μM, 3 h) cells.
Scale bar, 10 μm. Graphs show the quantification of Sec24C and TRAPPC2 in the peri-
Golgi area (mean intensity) normalized in SNS-032-treated cells relative to vehicle-
treated cells (set as 100%). Mean ± s.e.m. of three independent experiments.
p<0.05**, p<0.0001. **(B)** HeLa cells were treated with Dinaciclib (1 μM, 3 h) and
permeabilized or not with digitonin as described in Materials and methods. A GM130
antibody was added to the buffer of living cells to monitor permeabilization
efficiency. Upper panels, non-permeabilized (NP) control cells; middle panels,
permeabilized control cells; lower panels, permeabilized Dinaciclib-treated cells.
Scale bar, 10 μm. **(C)** Quantification of Sec24C membrane association after CDK
inhibitor treatment. Digitonin-permeabilized cells treated with vehicle (CTRL) or the
indicated CDK inhibitor (1 μM, 3 h) were immunostained for Sec24C. The mean
intensity of Sec24C in the perinuclear area normalized for the cytosolic Sec24C signal
in drug-treated cells is expressed as fold change compared to the control; n= 60-80;
mean ± s.e.m. of three independent experiments ****p<0.0001. **(D)** Cell lysates
from HeLa cells treated with SA (300 μM, 10 min), Dinaciclib (10 μM, 180 min), or
their combination were immunoprecipitated using a Sec31 Ab. Western blots of the
immunoprecipitates probed using the phosphor-Ser-CDK Ab (upper panel) or the
anti-Sec31 Ab (lower panel). Note that the anti-Sec31 Ab immunoprecipitates both
Sec31 isoforms

Figure 6. TRAPP and Sec23/24 migration to SGs depends on the proliferation state of cells. (A) HeLa cells were starved for 8 h with HBSS and subsequently exposed to SA (30 min, 300 µM), and immunostained for Sec24C or TRAPPC2 and eIF3. Scale bar 10 µm. **(B)** Analysis of the proliferation status of the cells after 8 h starvation in HBSS. Left, representative images of starved and non-starved cells using EdU incorporation (see Materials and methods) and, right, quantification of EdU incorporation in starved cells as a percentage of incorporation in control fed cells. **(C)** HeLa cells were

seeded at different confluency, treated with SA and stained for Sec24C and G3BP as an SG marker. Scale bar, 10 μm. Flow cytometry (FACS) analysis (right panels) was performed to evaluate the distribution of cell cycle phases in HeLa cell populations seeded at different confluency. The graph shows quantification of Sec24C mean intensity in SG puncta normalized for the cytosolic Sec24C at the indicated cell confluency. Mean ± s.e.m. of a representative experiment out of n=5 biological replicates. n= 50-80. ns, not significant; ****p<0.0001. **(D)** Growing and differentiated podocytes were treated with SA (300 μM, 30 min) and stained for TRAPPC2 or Sec24C and eIF3. Scale bar, 10μm. The graphs show quantification of TRAPPC2 and Sec24C (mean intensity) in SGs. Mean ± s.e.m. of three independents experiments ****p<0.0001. **(E)** CDK kinase activity in growing and differentiated podocytes. Western blot using a specific antibody recognizing phosphoSer-CDK substrates was used on total cell lysates. **(F)** Western blot analysis of phosphorylated retinoblastoma (p-RB) in growing (non-differentiated) versus differentiated podocytes. ß-actin was used as loading control. **(G)** Quantification of immunoprecipitated Sec31 from growing and differentiated podocytes using the phosphor-Ser-CDK Ab. Mean ± s.e.m. of three independents experiments. ***p<0.0007.

Figure 6—figure supplement 1. Differentiation of podocytes.

1268
Figure 7. The TRAPP complex controls SG composition and function. (A) Box plot
representing SG area (see Materials and methods) in mock, TRAPPC2- and TRAPPC3-
depleted cells treated with SA. Distribution of values from three independent
experiments. ****p<0.0001. **(B)** HeLa cells microinjected with control preimmune
IgG or a TRAPPC3-specific antibody (right panels in green) were treated with SA. The
TRAPPC3 Ab disrupts the Golgi (Yu et al., 2006), monitored using an anti-TGN46 Ab.
Anti-G3BP was used to stain SGs. The graph shows quantification of the SG area.

Mean ± s.e.m. three independent experiments; n>80. ****p<0.0001. **(C)** Structured Illumination Microscopy (SIM)-Super resolution (SR) images of endogenous TRAPPC2 and GFP-Sec23 localizing at SGs, stained for G3BP. Right, magnification of boxed area. **(D,E)** Localization of Raptor **(D)** and RACK1 **(E)** in mock, TRAPPC3-KD and TRAPPC2-KD HeLa cells treated with SA. G3BP was used to stain SGs. Scale bar, 10 μm. Each graph shows the quantification (mean intensity) of the respective protein in SG spots expressed as a percentage of the mock. Mean ± S.D. three independent replicates. *p<0.02; **p<0.009 in **(D)**, *p<0.05; ****p<0.0001 in **(E)**. **(F,G)** Localization of Raptor **(F)** and RACK1 **(G)** in untreated cells or cells pre-treated with the indicated CDK inhibitor (1 μM, 150 min) and then with SA (300 μM, 30 min). G3BP was used to stain SGs. Scale bar, 10 μm. Graphs show quantification of the localization of the respective protein with SGs, expressed as a percentage of the control. Mean ± s.e.m. of one representative experiment out of three independent replicates, n=60-80. ****p<0.0001. **(H)** Analysis of cell death after overnight recovery of HeLa cells treated or untreated with CDKi (SNS-032; Flavopiridol or Dinaciclib, 1 μM, 150 min) and then treated with SA (500 μM, 3 h). Images were automatically acquired by OPERETTA microscopy. Values indicate the percentage of the total number of nuclei (stained with DAPI) positive for BoBo-3 staining. Mean ± s.d. of one representative experiment out of three independent replicates. ns, not significant, ****p<0.0001.

Figure 7—figure supplement 1. TRAPP depletion does not affect protein translation inhibition caused by SA treatment.

1314 **Figure 8. The recruitment of TRAPP and COPII to SGs slows down ER export (A)** Left,
1315 control or SA-treated (500µM, 120 min) human fibroblasts (HFs) were incubated at
1316 40°C for 180 min and stained for eIF3, PCI, and Sec24C. Right, the cells were imaged
1317 10 min after shifting the temperature from 40°C to 32°C and stained for eIF3, PCI,
1318 and Giantin (to label the Golgi). Scale bar, 10 µm. **(B)** HeLa cells expressing VSV-G-
1319 UVR8 mEOS were left untreated or treated with SA alone or in combination with
1320 ISRIB (1 µM) for the indicated times and then pulsed with blue light. Images were

taken 10 minutes after the UV pulse and processed for staining. **(C)** Cells were
treated with SA for 30 min, the SA was washed out, and cells were left to recover for
180 min in the presence of cycloheximide, followed by a blue light pulse and
processing for staining as described in (B). **(D)** Quantification of VSV-G (mean
intensity) in the Golgi area to the total VSV-G per cell under the conditions described
in (B,C)**.** Data are expressed as percentage of the control. Mean ± s.e.m. of three
independent experiment. ****p<0.0001; ns, not significant.
Figure 8—figure supplement 1. ISRIB delays SG nucleation and inhibits Sec24C
relocalization to SGs.

1330

Figure 9. The recruitment of TRAPP induces the fragmentation of the GC in a Rab1-dependent manner. (A) AiryScan images of HeLa cells treated with SA for the indicated times and stained with GM130 (gray). Scale bar, 10 μm. The graph shows quantification of Golgi particles in SA-treated cells. Mean ± s.e.m. of three independent experiments. n=90-100. ****p<0.0001; ns, not significant. **(B)** Control cells and cells treated with Dinaciclib (CDKi) were treated with SA and stained for GM130 (in gray). Insets, eIF3. Blue, nuclear DAPI staining. The graph shows

quantification of Golgi particles in HeLa cells untreated or treated with SNS-032, Flavopiridol, or Dinaciclib for 150 min and treated with SA (300 µM, 30 min). Mean ± s.e.m. of three independent experiment. ****p<0.0001. **(C)** Quantification of Golgi particles in cells plated at different percentage of confluency. Mean ± s.e.m. of three independent experiment. ** p<0.001; *** p<0.0002; ns: not significant. **(D)** Quantification of Golgi objects in cells pre-treated with ISRIB (1 µM) for 3 h and exposed to SA (30 min) in the presence of ISRIB. *** p<0.002. **(E)** Electron microscopy images of a cell exposed to SA (300 µM, 30 min). GC: Golgi complex. Scale bar 200 nm. **(F)** AiryScan images of HeLa cells at steady state or exposed to SA for the indicated times. A Rab1-GTP specific antibody was used to monitor Rab1-GTP activity and GM130 to stain the Golgi complex. Insets, eIF3. The graph shows quantification of Rab1-GTP at the Golgi complex expressed as a percentage of the control. Mean ± s.e.m. of three independent experiments. *p<0.011, **p<0.002. **(G)** HeLa cells transfected or not with WT GFP-Rab1B were left untreated or treated with SA and stained for GM130 and G3BP to monitor SGs. Dashed white line, WT GFP-Rab1B transfected cells. The graph shows quantification of Golgi particles in the different conditions. NT: non-transfected. Mean ± s.e.m. of three independent experiment. ****p<0.0001; n=30-50.

Figure 9—figure supplement 1. TRAPPC2 migrates to SGs in Rab1 overexpressing cells.

Figure 10. Proposed model for secretion arrest mediated by TRAPP upon SG assembly in actively proliferating cells. Upon stress, the TRAPP complex relocalizes to SGs and drives the sequestering of Sec23/24, RACK1 and Raptor to SGs. This phenomenon leads to a block in ER-export and contributes to SG maturation.

1367
1368 **Figure 1—figure supplement 1.** Re-localization of TRAPP to SGs is independent of
1369 stress stimulus and cell type. (**A**) HeLa cells exposed to heat shock (44°C, 45 min), (**B**)
1370 human fibroblasts treated with SA (500 μM, 180 min), (**C**) U2OS cells treated with SA
1371 (300 μM, 60 min), stained for eIF3 and TRAPPC2.

1372

1373

1374 **Figure 1—figure supplement 2.** Stress-induced localization of Bet3 (TRAPPC3) to SGs
1375 in yeast cells. Cells co-expressing Bet3-GFP and the stress granule marker RFP-PAB1
1376 were exposed to heat shock (46°C, 10 min) and imaged.
1377 **Figure 1—supplement video 1.** Time-lapse fluorescent microscopy of Bet3-GFP
1378 (TRAPPC3) and RFP-Pab1 at 30°C in yeast cells. Images were taken every 8 sec for 64
1379 sec.
1380 **Figure 1—supplement video 2.** Time-lapse fluorescent microscopy of Bet3-GFP
1381 (TRAPPC3) and RFP-Pab1 at 46°C in yeast cells. Images were taken every 8 sec for 64
1382 sec.

1383

1384
1385 **Figure 1—figure supplement 3.** Association of TRAPP with SGs is reversible after SG

1386 disassembly. Cells were treated with SA for 30 min, the SA was washed out, and cells

1387 were left to recover for 120 min and processing for staining. Scale bar 10 μm.

1388

Figure 2—figure supplement 1. Sec24C and Sec23, but not components of other coat complexes or other cytosolic proteins associated with the exocytic and endocytic pathways, associate with SGs. (**A-L**) Cells at steady state or treated with SA were visualized by fluorescence microscopy using antibodies against the endogenous proteins or Flag-tagged proteins, or visualizing transfected GFP-tagged proteins, and co-stained for eIF3 or G3BP, as indicated. (**M**) Cells exposed to heat shock (44°C, 45 min) were stained for Sec24C or COPI, and G3BP. Blue, DAPI. (**N**) Schematic diagram showing the cellular location of the tested proteins that associate (red) or not (black) with SGs.

1402
Figure 2—figure supplement 2. Sec16 does not significantly associate with SGs nor is
it required for the recruitment of COPII to SGs. **(A)** Cells were treated with SA (300
µM) and co-stained for endogenous Sec16 and G3BP as an SG marker. **(B)** Cells
exposed to heat shock were processed as described in (A). **(C)** Representative images
of Sec24C localization in Sec16-KD cells treated with SA. Cells were fixed and
visualized by fluorescence microscopy using anti-Sec24C Ab, anti-G3BP Ab, and DAPI
(blue). Quantification of TRAPPC2 redistribution to SGs after KD calculated as the
ratio between Sec24C (mean intensity) in SG puncta and cytosolic Sec24C. Mean ±
s.e.m., n = 40-60 cells per experiment, N = 3. ns: not significant. **(D)** Sec16A and B
mRNA levels in Sec16A+B KD cells were evaluated by q-PCR and expressed as fold
change with respect to the MOCK. Dashed red line: MOCK.

1415
1416 **Figure 2—figure supplement 3. Evaluation of Sec23, Sec24, TRAPPC2 and TRAPPC3**
1417 **knock down efficiencies. (A)** Sec23A and Sec23B mRNA levels were evaluated by q-
1418 PCR and expressed as fold change with respect to the MOCK. Dashed red line:
1419 MOCK. **(B)** Sec24A, B, C and D mRNA levels were evaluated as described in (A). **(C)**
1420 Representative Western blot of HeLa cell extracts depleted for TRAPPC2 and
1421 TRAPPC3; GAPDH was used as loading control.
1422

1426 **Figure 3—figure supplement 1. (A)** Effect of Sec31 depletion on Sec24 localization at
1427 ERES. Cells were mock-treated or KD for Sec31 and then immunostained for Sec24C
1428 and cTAGE to stain ERES. Scale bar 10µm. The graph shows the mean intensity of
1429 Sec24C at ERES normalized for cytosolic signal as a percentage of control conditions.
1430 **(A)** Western blot analysis of Sec31A KD cells. ß-actin was used as loading control.
1431

1434   Figure 4—figure supplement 1 Evaluation of CDK1 and CDK2 knock down efficiency.
1435   CDK1 and CDK2 mRNA levels were evaluated by q-PCR and expressed as fold change
1436   with respect to the MOCK. Dashed red line: MOCK.

1438
1439

1440
1441 **Figure 6—figure supplement 1.** Differentiation of podocytes. Western blot of
1442 growing and differentiated podocytes. Synaptopodin was used as a differentiation
1443 marker, GAPDH as a loading control.
1444

1445

1446
1447 **Figure 7—figure supplement 1.** TRAPP depletion does not affect protein translation
1448 inhibition caused by SA treatment. **(A)** Puromycin assay (see Materials and methods)
1449 of MOCK, TRAPPC3 and TRAPPC2-KD HeLa cells at steady state and upon SA
1450 treatment. **(B)** Evaluation of phosphorylated eIF2a in TRAPPC3 and TRAPPC2
1451 depleted cells exposed to SA. ß-actin was used as a loading control. **(C)** Structured
1452 Illumination Microscopy (SIM)-Super resolution (SR) images of endogenous RACK1
1453 and G3BP. Scale bar 500 nm.
1454

Figure 8—figure supplement 1. ISRIB delays SG nucleation and prevents Sec24C relocalization to SGs. **(A)** HeLa cells were exposed to SA (200 μM) for the indicated time, alone or in combination with ISRIB and SGs were analyzed. Quantification of SG area (μm) of three independent experiment Mean ± s.e.m. **(B)** Graph, Quantification of Sec24C localization at SGs over time (the ratio between Sec24C (mean fluorescence intensity) in SG puncta and cytosolic) Sec24C. Mean ± SD. n = 60-80 cells per experiment, N = 3.

1466
1467 **Figure 9—figure supplement 1.** TRAPPC2 migrates to SGs in Rab1 overexpressing
1468 cells. HeLa cells overexpressing GFP-Rab1B-WT were treated with SA (300 µM 30
1469 min) and stained for TRAPPC2 (in red) and eIF3 (in gray). Dashed white line:
1470 Overexpressing cells.

1471

Table S1 TRAPPC2 interactors.

EEF1B2	P24534	28	24919	2	2	2	2	0.46	Elongation factor 1-beta OS=Homo sapiens GN=EEF1B2 PE=1 SV=3
EEF1D	E9PKY8	346	77150	8	7	7	6	0.44	Elongation factor 1-delta OS=Homo sapiens GN=EEF1D PE=1 SV=1
EEF1E1	H0YAL7	50	15444	2	1	2	1	0.35	Eukaryotic translation elongation factor 1 epsilon-1 (Fragment) OS=Homo sapiens GN=EEF1E1 PE=4 SV=1
EFTUD2	Q15029	84	106286	5	4	5	4	0.19	Isoform 2 of 116 kDa U5 small nuclear ribonucleoprotein component OS=Homo sapiens GN=EFTUD2
EHD2	Q9NZN4	58	46144	3	2	3	2	0.23	Isoform 2 of EH domain-containing protein 2 OS=Homo sapiens GN=EHD2
EIF2S2	P20042	292	38706	13	10	12	9	1.98	Eukaryotic translation initiation factor 2 subunit 2 OS=Homo sapiens GN=EIF2S2 PE=1 SV=2
EIF3G	O75821	446	35874	18	13	13	10	2.71	Eukaryotic translation initiation factor 3 subunit G OS=Homo sapiens GN=EIF3G PE=1 SV=2
EIF3J	O75822	181	29159	5	5	5	5	1.23	Eukaryotic translation initiation factor 3 subunit J OS=Homo sapiens GN=EIF3J PE=1 SV=2
EIF3K	Q9UBQ5	213	25329	9	6	8	5	1.52	Eukaryotic translation initiation factor 3 subunit K OS=Homo sapiens GN=EIF3K PE=1 SV=1
EIF4A2	Q14240	501	46688	12	10	8	7	1.03	Isoform 2 of Eukaryotic initiation factor 4A-II OS=Homo sapiens GN=EIF4A2
EIF4A3	P38919	241	47126	7	6	6	5	0.65	Eukaryotic initiation factor 4A-III OS=Homo sapiens GN=EIF4A3 PE=1 SV=4
EIF5A	I3L397	65	16237	3	2	3	2	0.78	Eukaryotic translation initiation factor 5A-1 (Fragment) OS=Homo sapiens GN=EIF5A PE=1 SV=2
EIF5B	O60841	432	139198	17	14	16	14	0.61	Eukaryotic translation initiation factor 5B OS=Homo sapiens GN=EIF5B PE=1 SV=4
ELMO2	H0YCM7	29	51873	3	1	3	1	0.10	Engulfment and cell motility protein 2 (Fragment) OS=Homo sapiens GN=ELMO2 PE=1 SV=1
EMG1	V9GYP5	115	26809	6	3	5	3	0.69	Uncharacterized protein (Fragment) OS=Homo sapiens PE=1 SV=1
EPRS	P07814	571	172080	20	20	17	17	0.64	Bifunctional glutamate/proline--tRNA ligase OS=Homo sapiens GN=EPRS PE=1 SV=5
ERAL1	J3QRV9	21	31877	1	1	1	1	0.16	GTPase Era, mitochondrial (Fragment) OS=Homo sapiens GN=ERAL1 PE=1 SV=2
ERLIN1	O75477	38	39072	1	1	1	1	0.13	Erlin-1 OS=Homo sapiens GN=ERLIN1 PE=1 SV=1
ESYT1	Q9BSJ8	16	124439	1	1	1	1	0.04	Isoform 2 of Extended synaptotagmin-1 OS=Homo sapiens GN=ESYT1
EWSR1	B0QYK0	267	65174	8	8	5	5	0.54	RNA-binding protein EWS OS=Homo sapiens GN=EWSR1 PE=1 SV=1
EXOC3	D6RBS9	15	53607	18	1	1	2	0.09	Exocyst complex component 3 OS=Homo sapiens GN=EXOC3 PE=1 SV=1
EXOSC1	B1AMU7	15	18231	3	1	3	1	0.29	Exosome complex component C1L4 (Fragment) OS=Homo sapiens GN=EXOSC1 PE=1 SV=2
EXOSC10	Q01780	24	98826	3	1	3	1	0.05	Isoform 2 of Exosome component 10 OS=Homo sapiens GN=EXOSC10
EYA3	B1APR7	18	45806	1	1	1	1	0.11	Eyes absent homolog 3 OS=Homo sapiens GN=EYA3 PE=1 SV=1
FABP5	Q01469	33	15497	1	1	1	1	0.35	Fatty acid-binding protein, epidermal OS=Homo sapiens GN=FABP5 PE=1 SV=3
FAM126A	H7C0W7	29	52801	2	1	2	1	0.09	Hyccin (Fragment) OS=Homo sapiens GN=FAM126A PE=4 SV=1
FAM83B	Q5T0W9	36	115185	3	1	3	1	0.04	Protein FAM83B OS=Homo sapiens GN=FAM83B PE=1 SV=1
FAM98A	Q8NCA5	165	55694	8	6	7	6	0.66	Isoform 2 of Protein FAM98A OS=Homo sapiens GN=FAM98A
FAM98B	Q52LJ0	49	45918	1	1	1	1	0.11	Isoform 2 of Protein FAM98B OS=Homo sapiens GN=FAM98B
FASN	P49327	48	275877	5	1	5	1	0.02	Fatty acid synthase OS=Homo sapiens GN=FASN PE=1 SV=3
FASTKD2	Q9NYY8	61	75343	1	1	1	1	0.06	Isoform 2 of FAST kinase domain-containing protein 2 OS=Homo sapiens GN=FASTKD2
FAU	E9PR30	42	10898	1	1	1	1	0.53	40S ribosomal protein S30 OS=Homo sapiens GN=FAU PE=1 SV=1
FMC1	Q96HJ9	144	54761	13	6	13	6	0.68	Isoform 2 of UPF0562 protein C7orf55 OS=Homo sapiens GN=C7orf55
FMR1	Q06787	146	67215	5	3	5	3	0.23	Isoform 1 of Fragile X mental retardation protein 1 OS=Homo sapiens GN=FMR1
FOXRED1	Q96CU9	98	30868	2	2	2	2	0.36	Isoform 2 of FAD-dependent oxidoreductase domain-containing protein 1 OS=Homo sapiens GN=FOXRED1
FSCN1	Q16658	37	55123	2	1	2	1	0.09	Fascin OS=Homo sapiens GN=FSCN1 PE=1 SV=3
FTSJ1	Q9UET6	22	36471	1	1	1	1	0.14	Isoform 2 of Putative tRNA (cytidine(32)/guanosine(34)-2'-O)methyltransferase OS=Homo sapiens GN=FTSJ1
FUBP3	Q96I24	48	61944	2	1	2	1	0.08	Far upstream element-binding protein 3 OS=Homo sapiens GN=FUBP3 PE=1 SV=2
FUS	P35637	432	53551	15	14	9	8	1.21	Isoform Short of RNA-binding protein FUS OS=Homo sapiens GN=FUS
FXR2	P51116	197	74520	9	6	9	6	0.46	Fragile X mental retardation syndrome-related protein 2 OS=Homo sapiens GN=FXR2 PE=1 SV=2
GANAB	F5H6X6	41	96441	1	1	1	1	0.05	Neutral alpha-glucosidase AB OS=Homo sapiens GN=GANAB PE=1 SV=1
GART	P22102	26	46631	1	1	1	1	0.11	Isoform Short of Trifunctional purine biosynthetic protein adenosine-3 OS=Homo sapiens GN=GART
GATAD2B	Q8WXI9	19	65562	2	1	2	1	0.07	Transcriptional repressor p66-beta OS=Homo sapiens GN=GATAD2B PE=1 SV=1
GDI2	P50395	63	51087	2	2	2	2	0.20	Rab GDP dissociation inhibitor beta OS=Homo sapiens GN=GDI2 PE=1 SV=2
GEMIN5	Q8TEQ6	48	170821	5	1	5	1	0.03	Gem-associated protein 5 OS=Homo sapiens GN=GEMIN5 PE=1 SV=3
GLUD2	P49448	44	61738	2	1	2	1	0.08	Glutamate dehydrogenase 2, mitochondrial OS=Homo sapiens GN=GLUD2 PE=1 SV=2
GNAS	Q5JWF2	106	110299	3	2	3	2	0.09	Isoform XLas-2 of Guanine nucleotide-binding protein G(s) subunit alpha isoforms XLas OS=Homo sapiens GN=GNAS
GNB3	F5H0S8	34	30024	2	1	2	1	0.17	Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-3 (Fragment) OS=Homo sapiens GN=GNB3 PE=4 SV=1
GNL3L	Q9NVN8	46	66216	1	1	1	1	0.07	Guanine nucleotide-binding protein-like 3-like protein OS=Homo sapiens GN=GNL3L PE=1 SV=1
GOLGA3	Q08378	110	155987	8	4	7	3	0.10	Isoform 2 of Golgin subfamily A member 3 OS=Homo sapiens GN=GOLGA3
GPATCH4	Q5T300	50	51125	3	1	3	1	0.10	Isoform 3 of G patch domain-containing protein 4 OS=Homo sapiens GN=GPATCH4
GRWD1	M0QX71	98	25632	2	2	2	2	0.44	Glutamate-rich WD repeat-containing protein 1 (Fragment) OS=Homo sapiens GN=GRWD1 PE=1 SV=1
GSTM3	P21266	30	26998	1	1	1	1	0.19	Glutathione S-transferase Mu 3 OS=Homo sapiens GN=GSTM3 PE=1 SV=3
GTPBP1	O00178	28	73035	3	1	3	1	0.07	GTP-binding protein 1 OS=Homo sapiens GN=GTPBP1 PE=1 SV=3
HADHB	B5MD38	31	38128	4	2	3	2	0.28	3-ketoacyl-CoA thiolase OS=Homo sapiens GN=HADHB PE=1 SV=1
HCK	H0Y3C5	20	59986	3	1	3	1	0.08	Tyrosine-protein kinase HCK OS=Homo sapiens GN=HCK PE=1 SV=1
HERC5	Q9UI4I	13	118203	1	1	1	1	0.04	E3 ISG15--protein ligase HERC5 OS=Homo sapiens GN=HERC5 PE=1 SV=2
HIST1H1D	P16402	27	22336	2	1	2	1	0.23	Histone H1.3 OS=Homo sapiens GN=HIST1H1D PE=1 SV=2
HIST1H2BK	O60814	71	13882	5	2	5	2	0.95	Histone H2B type 1-K OS=Homo sapiens GN=HIST1H2BK PE=1 SV=3
HLTF	Q14527	31	100882	2	1	2	1	0.05	Isoform 2 of Helicase-like transcription factor OS=Homo sapiens GN=HLTF
HNRNPA1	P09651	1099	34289	32	30	19	18	16.79	Isoform 2 of Heterogeneous nuclear ribonucleoprotein A1 OS=Homo sapiens GN=HNRNPA1
HNRNPAB	Q99729	219	36059	6	6	5	5	0.92	Isoform 2 of Heterogeneous nuclear ribonucleoprotein A/B OS=Homo sapiens GN=HNRNPAB
HNRNPDL	O14979	161	33739	8	7	7	6	1.31	Isoform 2 of Heterogeneous nuclear ribonucleoprotein D-like OS=Homo sapiens GN=HNRNPDL
HNRNPF	P52597	308	45985	15	12	10	7	1.05	Heterogeneous nuclear ribonucleoprotein F OS=Homo sapiens GN=HNRNPF PE=1 SV=3
HNRNPK	P61978	2753	48760	82	73	24	23	14.00	Isoform 3 of Heterogeneous nuclear ribonucleoprotein K OS=Homo sapiens GN=HNRNPK
HNRNPUL1	Q9BUJ2	448	84903	19	13	16	11	0.95	Isoform 3 of Heterogeneous nuclear ribonucleoprotein U-like protein 1 OS=Homo sapiens GN=HNRNPUL1
HRNR	Q86YZ3	176	283140	3	3	2	2	0.03	Hornerin OS=Homo sapiens GN=HRNR PE=1 SV=2
HSP90AA1	P07900	345	98670	16	13	14	12	0.86	Isoform 2 of Heat shock protein HSP 90-alpha OS=Homo sapiens GN=HSP90AA1
HSP90AB1	P08238	639	83554	26	23	22	20	2.28	Heat shock protein HSP 90-beta OS=Homo sapiens GN=HSP90AB1 PE=1 SV=4
HSP90B1	P14625	243	92696	6	5	5	4	0.23	Endoplasmin OS=Homo sapiens GN=HSP90B1 PE=1 SV=1
HSPB1	P04792	52	22826	4	4	3	3	0.85	Heat shock protein beta-1 OS=Homo sapiens GN=HSPB1 PE=1 SV=2
HSPD1	P10809	367	61187	10	9	10	9	1.00	60 kDa heat shock protein, mitochondrial OS=Homo sapiens GN=HSPD1 PE=1 SV=2
HTATSF1	Q5H918	57	27898	1	1	1	1	0.11	HIV Tat-specific factor 1 (Fragment) OS=Homo sapiens GN=HTATSF1 PE=1 SV=1
IDH1	O75874	33	46915	1	1	1	1	0.11	Isocitrate dehydrogenase [NADP] cytoplasmic OS=Homo sapiens GN=IDH1 PE=1 SV=2
IGF2BP1	Q9NZI8	468	63783	14	11	12	11	1.26	Insulin-like growth factor 2 mRNA-binding protein 1 OS=Homo sapiens GN=IGF2BP1 PE=1 SV=2
IGF2BP2	Q9Y6M1	219	61918	6	5	6	5	0.46	Isoform 2 of Insulin-like growth factor 2 mRNA-binding protein 2 OS=Homo sapiens GN=IGF2BP2
IGHV4OR15-8	A0A075B7B6	22	13105	1	1	1	1	0.43	Putative V-set and immunoglobulin domain-containing-like protein IGHV4OR15-8 (Fragment) OS=Homo sapiens GN=IGHV4OR15-8 PE=4 SV=1
IGKV2-30	P06310	4080	14811	88	77	3	3	1.5	Ig kappa chain V-II region RPMI 6410 OS=Homo sapiens PE=4 SV=1
IKBIP	Q70UQ0	122	43228	4	3	4	3	0.39	Isoform 2 of Inhibitor of nuclear factor kappa-B kinase-interacting protein OS=Homo sapiens GN=IKBIP
IL17RC	G3V177	34	5031	1	1	1	1	1.45	Interleukin 17 receptor C, isoform CRA_g OS=Homo sapiens GN=IL17RC PE=4 SV=1
ILF2	Q12905	156	43263	7	5	7	5	0.72	Interleukin enhancer-binding factor 2 OS=Homo sapiens GN=ILF2 PE=1 SV=2
ILF3	Q12906	416	76441	17	12	13	10	0.85	Isoform 2 of Interleukin enhancer-binding factor 3 OS=Homo sapiens GN=ILF3
IPO8	O15397	418	120945	11	9	11	9	0.42	Importin-8 OS=Homo sapiens GN=IPO8 PE=1 SV=2
ITPR1	E7EVP7	39	315354	7	1	6	1	0.02	Inositol 1,4,5-trisphosphate receptor type 1 OS=Homo sapiens GN=ITPR1 PE=1 SV=1
JUP	P14923	103	82434	3	2	3	2	0.12	Junction plakoglobin OS=Homo sapiens GN=JUP PE=1 SV=3
KARS	Q15046	192	71908	7	7	7	7	0.58	Isoform Mitochondrial of Lysine--tRNA ligase OS=Homo sapiens GN=KARS
KCTD17	H0Y231	59	20928	1	1	1	1	0.25	BTB/POZ domain-containing protein KCTD17 (Fragment) OS=Homo sapiens GN=KCTD17 PE=4 SV=1
KCTD2	Q14681	323	28794	8	8	5	5	1.66	BTB/POZ domain-containing protein KCTD2 OS=Homo sapiens GN=KCTD2 PE=1 SV=3
KIF2C	Q5JR89	71	32827	1	1	1	1	0.15	Kinesin-like protein KIF2C (Fragment) OS=Homo sapiens GN=KIF2C PE=1 SV=1
KLB	Q86Z14	35	120473	2	1	2	1	0.04	Beta-klotho OS=Homo sapiens GN=KLB PE=1 SV=1
KLHL6	Q9P2J3	23	70239	1	1	1	1	0.07	Kelch-like protein 6 OS=Homo sapiens GN=KLHL6 PE=1 SV=2
KPNB1	J3QR48	71	16502	2	1	2	1	0.33	Importin subunit beta-1 (Fragment) OS=Homo sapiens GN=KPNB1 PE=1 SV=2
KRAS	G3V4K2	20	4664	1	1	1	1	1.61	GTPase KRas OS=Homo sapiens GN=KRAS PE=4 SV=1
KRI1	Q8N9T8	42	83004	2	1	2	1	0.06	Protein KRI1 homolog OS=Homo sapiens GN=KRI1 PE=1 SV=1
KRT10	P13645	2747	59020	72	65	36	33	16.74	Keratin, type I cytoskeletal 10 OS=Homo sapiens GN=KRT10 PE=1 SV=6
KRT2	P35908	1146	65678	33	31	25	25	5.95	Keratin, type II cytoskeletal 2 epidermal OS=Homo sapiens GN=KRT2 PE=1 SV=2
KRT6A	P02538	644	60293	27	21	21	19	3.79	Keratin, type II cytoskeletal 6A OS=Homo sapiens GN=KRT6A PE=1 SV=3
KRT72	Q14CN4	107	56342	11	5	8	4	0.40	Isoform 2 of Keratin, type II cytoskeletal 72 OS=Homo sapiens GN=KRT72
KRT86	O43790	55	55120	6	3	5	3	0.29	Keratin, type II cuticular Hb6 OS=Homo sapiens GN=KRT86 PE=1 SV=1
KTN1	Q86UP2	108	149804	10	3	9	3	0.10	Isoform 2 of Kinectin OS=Homo sapiens GN=KTN1
LANCL1	F8WDS9	26	9334	1	1	1	1	0.64	LanC-like protein 1 OS=Homo sapiens GN=LANCL1 PE=1 SV=1
LAPTM4B	Q86VI4	29	25972	2	2	2	2	0.43	Isoform 2 of Lysosomal-associated transmembrane protein 4B OS=Homo sapiens GN=LAPTM4B
LARP4	Q71RC2	110	81877	5	2	5	2	0.12	Isoform 4 of La-related protein 4 OS=Homo sapiens GN=LARP4
LARP4B	Q92615	27	80902	2	2	2	2	0.12	La-related protein 4B OS=Homo sapiens GN=LARP4B PE=1 SV=3
LARS	B4DER1	140	132410	8	7	8	7	0.28	Leucine--tRNA ligase, cytoplasmic OS=Homo sapiens GN=LARS PE=1 SV=1
LDHB	P07195	226	36900	5	5	5	5	0.89	L-lactate dehydrogenase B chain OS=Homo sapiens GN=LDHB PE=1 SV=2
LGALS3BP	Q08380	29	66202	2	2	2	2	0.15	Galectin-3-binding protein OS=Homo sapiens GN=LGALS3BP PE=1 SV=1
LIFR	P42702	19	125090	1	1	1	1	0.04	Leukemia inhibitory factor receptor OS=Homo sapiens GN=LIFR PE=1 SV=1
LIMA1	Q9UHB6	569	85630	25	19	22	17	1.56	LIM domain and actin-binding protein 1 OS=Homo sapiens GN=LIMA1 PE=1 SV=1
LMNA	P02545	130	65153	5	3	4	3	0.24	Isoform C of Prelamin-A/C OS=Homo sapiens GN=LMNA
LMO7	J3KP06	889	191723	38	32	34	29	1.10	LIM domain only protein 7 OS=Homo sapiens GN=LMO7 PE=1 SV=1
LRMDA	M0R2H0	25	6258	3	1	2	1	1.07	Leucine-rich repeat-containing protein C10orf11 OS=Homo sapiens GN=C10orf11 PE=1 SV=1
LRPPRC	P42704	290	159003	9	6	9	6	0.20	Leucine-rich PPR motif-containing protein, mitochondrial OS=Homo sapiens GN=LRPPRC PE=1 SV=3
LRRK2	Q5S007	14	289568	9	1	8	1	0.02	Leucine-rich repeat serine/threonine-protein kinase 2 OS=Homo sapiens GN=LRRK2 PE=1 SV=2
LSM14A	I3L4Q1	18	11439	1	1	1	1	0.50	Protein LSM14 homolog A OS=Homo sapiens GN=LSM14A PE=1 SV=1
LUC7L	Q9NQ29	80	38781	5	3	5	3	0.44	Isoform 2 of Putative RNA-binding protein Luc7-like 1 OS=Homo sapiens GN=LUC7L
LUC7L3	J3KPP4	81	58698	2	1	2	1	0.08	Cisplatin resistance-associated overexpressed protein, isoform CRA_b OS=Homo sapiens GN=LUC7L3 PE=1 SV=1
MAGED2	Q5H907	71	55932	3	3	3	3	0.29	Melanoma antigen family D, 2, isoform CRA_d OS=Homo sapiens GN=MAGED2 PE=1 SV=1
MAP7	Q14244	39	87014	7	1	7	1	0.06	Isoform 7 of Ensconsin OS=Homo sapiens GN=MAP7
MAT2A	P31153	341	43975	11	10	9	8	1.62	S-adenosylmethionine synthase isoform type-2 OS=Homo sapiens GN=MAT2A PE=1 SV=1
MCAT	Q8IVS2	32	43390	1	1	1	1	0.11	Malonyl-CoA-acyl carrier protein transacylase, mitochondrial OS=Homo sapiens GN=MCAT PE=1 SV=2
MCRIP2	I3L172	50	7085	2	1	1	1	0.90	Protein FAM195A OS=Homo sapiens GN=FAM195A PE=1 SV=1

Gene name	UNIPROT id	Score	Mass	Num. of matches	Num. of significant matches	Num. of sequences	Num. of significant sequences	emPAI	Description
MARC1	Q5VT66	21	39999	2	1	2	1	0.13	Isoform 2 of Mitochondrial amidoxime-reducing component 1 OS=Homo sapiens GN=MARC1
AAAS	Q9NRG9	130	60392	6	3	6	3	0.26	Aladin OS=Homo sapiens GN=AAAS PE=1 SV=1
ABCF1	Q8NE71	249	92080	11	8	10	8	0.51	Isoform 2 of ATP-binding cassette sub-family F member 1 OS=Homo sapiens GN=ABCF1
AC093512.2	J3KPS3	104	40249	5	4	4	4	0.60	Aldolase A, fructose-bisphosphate, isoform CRA_b OS=Homo sapiens GN=ALDOA PE=1 SV=1
ACADVL	P49748	22	68414	1	1	1	1	0.07	Isoform 2 of Very long-chain specific acyl-CoA dehydrogenase, mitochondrial OS=Homo sapiens GN=ACADVL
ACLY	P53396	102	120608	3	1	3	1	0.04	Isoform 2 of ATP-citrate synthase OS=Homo sapiens GN=ACLY
ACOT9	H7C5Q2	26	30256	1	1	1	1	0.17	Acyl-coenzyme A thioesterase 9, mitochondrial (Fragment) OS=Homo sapiens GN=ACOT9 PE=1 SV=1
ACTBL2	Q562R1	394	42318	18	14	7	5	1.18	Beta-actin-like protein 2 OS=Homo sapiens GN=ACTBL2 PE=1 SV=2
ACTN1	H9KV75	172	95279	7	6	6	5	0.28	Alpha-actinin-1 OS=Homo sapiens GN=ACTN1 PE=1 SV=2
ACTN4	O43707	146	105245	7	7	6	6	0.31	Alpha-actinin-4 OS=Homo sapiens GN=ACTN4 PE=1 SV=2
ACTR1A	P61163	29	42701	3	1	3	1	0.12	Alpha-centractin OS=Homo sapiens GN=ACTR1A PE=1 SV=1
ADAR	H0YCK3	61	133761	3	1	3	1	0.04	Double-stranded RNA-specific adenosine deaminase (Fragment) OS=Homo sapiens GN=ADAR PE=1 SV=1
ADRM1	Q16186	22	42412	1	1	1	1	0.12	Proteasomal ubiquitin receptor ADRM1 OS=Homo sapiens GN=ADRM1 PE=1 SV=2
AGO2	Q9UKV8	96	94758	5	4	5	4	0.22	Isoform 2 of Protein argonaute-2 OS=Homo sapiens GN=AGO2
AHNAK	Q09666	68	629213	10	2	10	2	0.02	Neuroblast differentiation-associated protein AHNAK OS=Homo sapiens GN=AHNAK PE=1 SV=2
AIFM1	O95831	99	66539	10	4	8	4	0.33	Isoform 3 of Apoptosis-inducing factor 1, mitochondrial OS=Homo sapiens GN=AIFM1
AIMP1	Q12904	164	37358	5	5	5	5	0.88	Isoform 2 of Aminoacyl tRNA synthase complex-interacting multifunctional protein 1 OS=Homo sapiens GN=AIMP1
AIMP2	Q13155	57	35668	3	1	3	1	0.14	Aminoacyl tRNA synthase complex-interacting multifunctional protein 2 OS=Homo sapiens GN=AIMP2 PE=1 SV=2
AKAP8L	Q9ULX6	55	72060	2	1	2	1	0.07	A-kinase anchor protein 8-like OS=Homo sapiens GN=AKAP8L PE=1 SV=3
AKAP9	Q99996	19	454989	5	1	5	1	0.01	Isoform 2 of A-kinase anchor protein 9 OS=Homo sapiens GN=AKAP9
ALPG	P10696	344	57626	10	10	8	8	0.93	Alkaline phosphatase, placental-like OS=Homo sapiens GN=ALPP L2 PE=1 SV=4
ANKS6	Q68DC2	15	92575	2	1	2	1	0.05	Isoform 3 of Ankyrin repeat and SAM domain-containing protein 6 OS=Homo sapiens GN=ANKS6
ANXA1	P04083	33	38918	4	1	4	1	0.13	Annexin A1 OS=Homo sapiens GN=ANXA1 PE=1 SV=2
ANXA2	H0YM50	166	28582	8	6	7	5	1.28	Annexin (Fragment) OS=Homo sapiens GN=ANXA2 PE=1 SV=1
AP2A1	O95782	152	106378	9	5	9	5	0.25	Isoform 8 of AP-2 complex subunit alpha-1 OS=Homo sapiens GN=AP2A1
AP2B1	P63010	40	106537	5	2	5	2	0.09	Isoform 2 of AP-2 complex subunit beta OS=Homo sapiens GN=AP2B1
AP2M1	E9PFW3	35	52612	2	1	2	1	0.09	AP-2 complex subunit mu OS=Homo sapiens GN=AP2M1 PE=1 SV=1
AP2S1	M0QYZ2	23	19145	3	1	3	1	0.28	AP-2 complex subunit sigma OS=Homo sapiens GN=AP2S1 PE=1 SV=1
AP3S2	H0YLI7	19	23611	1	1	1	1	0.22	AP-3 complex subunit sigma-2 OS=Homo sapiens GN=AP3S2 PE=1 SV=1
ARF5	C9J1Z8	30	17153	2	1	2	1	0.31	ADP-ribosylation factor 5 (Fragment) OS=Homo sapiens GN=ARF5 PE=1 SV=1
ARHGDIA	J3KTF8	20	21561	2	1	2	1	0.24	Rho GDP-dissociation inhibitor 1 (Fragment) OS=Homo sapiens GN=ARHGDIA PE=1 SV=2
ARHGEF2	V9GYM8	103	117025	7	2	7	2	0.08	Rho guanine nucleotide exchange factor 2 OS=Homo sapiens GN=ARHGEF2 PE=1 SV=1
ASF1B	K7ES22	33	16102	1	1	1	1	0.34	Histone chaperone ASF1B OS=Homo sapiens GN=ASF1B PE=1 SV=1
ASPH	Q12797	226	83672	11	8	10	7	0.48	Isoform 10 of Aspartyl/asparaginyl beta-hydroxylase OS=Homo sapiens GN=ASPH
ASS1	P00966	39	46786	3	3	3	3	0.35	Argininosuccinate synthase OS=Homo sapiens GN=ASS1 PE=1 SV=2
ATAD3A	Q9NVI7	161	66291	12	9	11	8	0.77	Isoform 2 of ATPase family AAA domain-containing protein 3A OS=Homo sapiens GN=ATAD3A
ATP1A2	B1AKY9	109	112046	4	2	4	2	0.09	Sodium/potassium-transporting ATPase subunit alpha-2 OS=Homo sapiens GN=ATP1A2 PE=1 SV=1
ATP2A2	P16615	61	111103	3	2	3	2	0.09	Isoform 2 of Sarcoplasmic/endoplasmic reticulum calcium ATPase 2 OS=Homo sapiens GN=ATP2A2
ATP2C1	H0Y8X9	16	24342	1	1	1	1	0.21	Calcium-transporting ATPase type 2C member 1 (Fragment) OS=Homo sapiens GN=ATP2C1 PE=1 SV=1
ATP5F1A	P25705	354	59828	12	11	11	10	1.20	ATP synthase subunit alpha, mitochondrial OS=Homo sapiens GN=ATP5A1 PE=1 SV=1
ATP5F1B	P06576	500	56525	18	12	12	9	1.51	ATP synthase subunit beta, mitochondrial OS=Homo sapiens GN=ATP5B PE=1 SV=3
ATP5F1C	P36542	69	33032	3	3	3	3	0.53	ATP synthase subunit gamma, mitochondrial OS=Homo sapiens GN=ATP5C1 PE=1 SV=1
ATP5MD	Q96IX5	28	6510	1	1	1	1	1.00	Up-regulated during skeletal muscle growth protein 5 OS=Homo sapiens GN=USMG5 PE=1 SV=1
ATP5PB	Q5QNZ2	16	22318	1	1	1	1	0.23	ATP synthase F(0) complex subunit B1, mitochondrial OS=Homo sapiens GN=ATP5F1 PE=1 SV=1
ATP5PO	H7C086	112	8242	2	2	2	2	2.05	ATP synthase subunit O, mitochondrial (Fragment) OS=Homo sapiens GN=ATP5O PE=1 SV=1
ATXN2	Q99700	78	106610	4	2	4	2	0.09	Isoform 2 of Ataxin-2 OS=Homo sapiens GN=ATXN2
BASP1	P80723	28	17654	1	1	1	1	0.30	Isoform 2 of Brain acid soluble protein 1 OS=Homo sapiens GN=BASP1
BLM	P54132	29	160611	3	1	3	1	0.03	Bloom syndrome protein OS=Homo sapiens GN=BLM PE=1 SV=1
BRIX1	Q8TDN6	32	41660	2	1	2	1	0.12	Ribosome biogenesis protein BRX1 homolog OS=Homo sapiens GN=BRIX1 PE=1 SV=2
BST2	Q10589	66	18611	1	1	1	1	0.28	Isoform 2 of Bone marrow stromal antigen 2 OS=Homo sapiens GN=BST2
BYSL	Q13895	83	49798	4	2	4	2	0.21	Bystin OS=Homo sapiens GN=BYSL PE=1 SV=3
C1QBP	Q07021	137	31742	4	4	3	3	0.56	Complement component 1 Q subcomponent-binding protein, mitochondrial OS=Homo sapiens GN=C1QBP PE=1 SV=1
C7orf50	H7C0T1	30	18846	1	1	1	1	0.28	Uncharacterized protein C7orf50 (Fragment) OS=Homo sapiens GN=C7orf50 PE=1 SV=1
CALM2	H0Y7A7	123	20863	4	2	3	1	0.25	Calmodulin (Fragment) OS=Homo sapiens GN=CALM2 PE=1 SV=1
CALR	K7EJB9	63	28528	2	2	2	2	0.39	Calreticulin (Fragment) OS=Homo sapiens GN=CALR PE=1 SV=1
CAND1	Q86VP6	26	119182	2	1	2	1	0.04	Isoform 2 of Cullin-associated NEDD8-dissociated protein 1 OS=Homo sapiens GN=CAND1
CAPZB	B1AKB5	37	29703	2	2	2	2	0.37	F-actin-capping protein subunit beta OS=Homo sapiens GN=CAPZB PE=1 SV=1
CAVIN1	Q6NZI2	41	33342	2	2	1	1	0.33	Isoform 2 of Polymerase I and transcript release factor OS=Homo sapiens GN=PTRF
CCAR1	Q8IX12	127	131646	13	7	12	7	0.29	Isoform 2 of Cell division cycle and apoptosis regulator protein 1 OS=Homo sapiens GN=CCAR1
CCDC124	Q96CT7	34	25820	3	1	3	1	0.20	Coiled-coil domain-containing protein 124 OS=Homo sapiens GN=CCDC124 PE=1 SV=1
CCDC137	Q6PKO4	31	33268	2	1	2	1	0.15	Coiled-coil domain-containing protein 137 OS=Homo sapiens GN=CCDC137 PE=1 SV=1
CCT5	P48643	247	60089	10	9	9	8	1.03	T-complex protein 1 subunit epsilon OS=Homo sapiens GN=CCT5 PE=1 SV=1
CD55	B1AP13	108	50561	5	5	5	5	0.59	Complement decay-accelerating factor OS=Homo sapiens GN=CD55 PE=1 SV=1
CD59	E9PNW4	19	12604	1	1	1	1	0.44	CD59 glycoprotein OS=Homo sapiens GN=CD59 PE=1 SV=1
CDC5L	Q99459	77	92422	6	2	6	2	0.11	Cell division cycle 5-like protein OS=Homo sapiens GN=CDC5L PE=1 SV=2
CEP164	Q9UPV0	15	163958	5	1	5	1	0.02	Isoform 2 of Centrosomal protein of 164 kDa OS=Homo sapiens GN=CEP164
CHCHD3	F8WAR4	69	27832	2	2	2	2	0.40	Coiled-coil-helix-coiled-coil-helix domain-containing protein 3, mitochondrial OS=Homo sapiens GN=CHM45 PE=1 SV=1
CHERP	J3QK89	225	105324	11	9	9	7	0.43	Calcium homeostasis endoplasmic reticulum protein OS=Homo sapiens GN=CHERP PE=1 SV=1
CKAP5	Q14008	17	220041	2	1	2	1	0.02	Isoform 2 of Cytoskeleton-associated protein 5 OS=Homo sapiens GN=CKAP5
CKB	G3V4N7	64	24272	1	1	1	1	0.25	Creatine kinase B-type (Fragment) OS=Homo sapiens GN=CKB PE=1 SV=1
CKMT1B	F8WCN3	44	33782	1	1	1	1	0.15	Creatine kinase U-type, mitochondrial OS=Homo sapiens GN=CKMT1B PE=1 SV=1
CLIC1	O00299	1590	27248	69	58	13	13	8.42	Chloride intracellular channel protein 1 OS=Homo sapiens GN=CLIC1 PE=1 SV=4
CLIC5	Q9NZA1	79	28389	7	3	2	2	0.39	Isoform 1 of Chloride intracellular channel protein 5 OS=Homo sapiens GN=CLIC5
CLNS1A	P54105	959	26370	25	23	7	7	6.05	Methylosome subunit pICln OS=Homo sapiens GN=CLNS1A PE=1 SV=1
CLPX	O76031	130	69922	5	5	4	4	0.31	ATP-dependent Clp protease ATP-binding subunit clpX-like, mitochondrial OS=Homo sapiens GN=CLPX PE=1 SV=2
CLTC	Q00610	167	189538	6	4	6	4	0.11	Isoform 2 of Clathrin heavy chain 1 OS=Homo sapiens GN=CLTC
CMAS	Q8NFW8	14	49033	2	1	2	1	0.10	N-acylneuraminate cytidylyltransferase OS=Homo sapiens GN=CMAS PE=1 SV=2
CMBL	Q96DG6	982	28372	36	31	18	18	18.55	Carboxymethylenebutenolidase homolog OS=Homo sapiens GN=CMBL PE=1 SV=1
COPA	P53621	2978	140832	107	99	61	61	9.15	Isoform 2 of Coatomer subunit alpha OS=Homo sapiens GN=COPA
COPB1	P53618	117	108214	3	3	2	2	0.14	Coatomer subunit beta OS=Homo sapiens GN=COPB1 PE=1 SV=3
COPB2	P35606	1727	99839	50	46	35	32	4.50	Isoform 2 of Coatomer subunit beta' OS=Homo sapiens GN=COPB2
COPE	O14579	778	34688	23	22	12	12	8.98	Coatomer subunit epsilon OS=Homo sapiens GN=COPE PE=1 SV=3
COPG1	Q9Y678	66	98967	3	2	3	2	0.10	Coatomer subunit gamma-1 OS=Homo sapiens GN=COPG1 PE=1 SV=1
CPSF6	F8WJN3	249	52409	7	5	6	4	0.43	Cleavage and polyadenylation specificity factor subunit 6 OS=Homo sapiens GN=CPSF6 PE=1 SV=1
CPSF7	Q8N684	133	51179	7	4	7	4	0.45	Isoform 2 of Cleavage and polyadenylation specificity factor subunit 7 OS=Homo sapiens GN=CPSF7
CPT1A	P50416	123	88995	6	5	6	5	0.30	Carnitine O-palmitoyltransferase 1, liver isoform OS=Homo sapiens GN=CPT1A PE=1 SV=2
CROCC	Q5TZA2	65	228787	16	1	16	1	0.02	Rootletin OS=Homo sapiens GN=CROCC PE=1 SV=1
CSDE1	E9PLT0	49	75164	5	2	4	2	0.13	Cold shock domain-containing protein E1 OS=Homo sapiens GN=CSDE1 PE=1 SV=1
CSNK1A1	P48729	70	37771	3	2	3	2	0.28	Isoform 2 of Casein kinase I isoform alpha OS=Homo sapiens GN=CSNK1A1
CSNK1E	H0YS59	51	12971	2	2	2	2	1.04	Casein kinase I isoform epsilon (Fragment) OS=Homo sapiens GN=CSNK1E PE=1 SV=1
CTAG2	O75638	50	18453	1	1	1	1	0.29	Isoform LAGE-1A of Cancer/testis antigen 2 OS=Homo sapiens GN=CTAG2
CTTN	H7C314	32	25136	1	1	1	1	0.21	Src substrate cortactin (Fragment) OS=Homo sapiens GN=CTTN PE=1 SV=1
CWF19L1	Q69YN2	52	46078	1	1	1	1	0.11	Isoform 3 of CWF19-like protein 1 OS=Homo sapiens GN=CWF19L1
DAP3	P51398	167	41361	7	5	7	5	0.77	Isoform 2 of 28S ribosomal protein S29, mitochondrial OS=Homo sapiens GN=DAP3
DARS	H7BZ35	58	21708	5	4	5	4	1.36	Aspartate--tRNA ligase, cytoplasmic (Fragment) OS=Homo sapiens GN=DARS PE=1 SV=1
DBN1	Q16643	76	72068	4	2	4	2	0.14	Isoform 2 of Drebrin OS=Homo sapiens GN=DBN1
DCD	P81605	171	11391	4	4	3	3	2.37	Dermcidin OS=Homo sapiens GN=DCD PE=1 SV=2
DCTN4	Q9UJW0	16	54030	2	1	2	1	0.09	Isoform 3 of Dynactin subunit 4 OS=Homo sapiens GN=DCTN4
DDB1	Q16531	43	128142	4	2	4	2	0.08	DNA damage-binding protein 1 OS=Homo sapiens GN=DDB1 PE=1 SV=1
DDOST	P39656	38	46719	2	1	2	1	0.11	Isoform 2 of Dolichyl-diphosphooligosaccharide--protein glycosyltransferase 48 kDa subunit OS=Homo sapiens GN=DDOST
DDR2	Q16832	21	97586	2	1	2	1	0.05	Discoidin domain-containing receptor 2 OS=Homo sapiens GN=DDR2 PE=1 SV=2
DDX21	Q9NR30	43	80121	4	2	4	2	0.13	Isoform 2 of Nucleolar RNA helicase 2 OS=Homo sapiens GN=DDX21
DDX39B	Q13838	385	51103	13	13	10	10	1.52	Isoform 2 of Spliceosome RNA helicase DDX39B OS=Homo sapiens GN=DDX39B
DDX47	Q9H0S4	50	45312	3	2	3	2	0.11	Isoform 2 of Probable ATP-dependent RNA helicase DDX47 OS=Homo sapiens GN=DDX47
DECR1	E5RJG7	60	19429	2	1	1	1	0.27	2,4-dienoyl-CoA reductase, mitochondrial (Fragment) OS=Homo sapiens GN=DECR1 PE=1 SV=1
DERL1	B4E1G1	16	17079	1	1	1	1	0.31	Derlin-1 OS=Homo sapiens GN=DERL1 PE=1 SV=1
DHCR24	Q15392	57	55456	3	2	3	2	0.19	Isoform 2 of Delta(24)-sterol reductase OS=Homo sapiens GN=DHCR24
DHX30	Q7L2E3	579	137057	22	17	20	15	0.68	Isoform 2 of Putative ATP-dependent RNA helicase DHX30 OS=Homo sapiens GN=DHX30
DHX37	F5H3Y4	17	108432	2	1	1	1	0.04	Probable ATP-dependent RNA helicase DHX37 OS=Homo sapiens GN=DHX37 PE=1 SV=1
DHX9	Q08211	828	142181	26	26	24	20	0.95	ATP-dependent RNA helicase A OS=Homo sapiens GN=DHX9 PE=1 SV=4
DIMT1	D6RCL3	44	17173	3	1	2	1	0.31	Probable dimethyladenosine transferase OS=Homo sapiens GN=DIMT1 PE=1 SV=1
DLD	E9PEX6	24	52182	1	1	1	1	0.09	Dihydrolipoyl dehydrogenase, mitochondrial OS=Homo sapiens GN=DLD PE=1 SV=1
DNAJA1	P31689	161	37762	8	6	5	4	0.65	Isoform 2 of DnaJ homolog subfamily A member 1 OS=Homo sapiens GN=DNAJA1
DNAJA2	O60884	71	46344	3	2	3	2	0.23	DnaJ homolog subfamily A member 2 OS=Homo sapiens GN=DNAJA2 PE=1 SV=1
DNAJB11	H7C2Y5	57	19249	5	1	4	1	0.28	DnaJ homolog subfamily B member 11 (Fragment) OS=Homo sapiens GN=DNAJB11 PE=1 SV=2
DNAJC8	O75937	105	29823	3	3	3	3	0.60	DnaJ homolog subfamily C member 8 OS=Homo sapiens GN=DNAJC8 PE=1 SV=1
DNAJC9	Q8WXX5	29	30062	1	1	1	1	0.17	DnaJ homolog subfamily C member 9 OS=Homo sapiens GN=DNAJC9 PE=1 SV=1
DPM1	H0Y368	47	33537	4	1	4	1	0.15	Dolichol-phosphate mannosyltransferase subunit 1 (Fragment) OS=Homo sapiens GN=DPM1 PE=1 SV=1
DRG1	Q9Y295	106	40802	6	3	5	3	0.41	Developmentally-regulated GTP-binding protein 1 OS=Homo sapiens GN=DRG1 PE=1 SV=1
DYNC1H1	Q14204	31	534809	8	1	8	1	0.01	Cytoplasmic dynein 1 heavy chain 1 OS=Homo sapiens GN=DYNC1H1 PE=1 SV=5
EBNA1BP2	H7C2Q8	49	40830	4	2	3	2	0.26	EBNA1 binding protein 2, isoform CRA_c OS=Homo sapiens GN=EBNA1BP2 PE=1 SV=1

1474

Gene	Accession							Ratio	Description
PTBP2	Q9UKA9	90	57725	4	3	4	3	0.28	Isoform 2 of Polypyrimidine tract-binding protein 2 OS=Homo sapiens GN=PTBP2
PTCD3	Q96EY7	32	32128	1	1	1	1	0.16	Isoform 2 of Pentatricopeptide repeat domain-containing protein 3, mitochondrial OS=Homo sapiens GN=PTCD3
PTCHD3	Q3KNS1	17	87842	3	1	2	1	0.06	Patched domain-containing protein 3 OS=Homo sapiens GN=PTCHD3 PE=1 SV=3
PTS	Q03393	163	16489	8	8	5	5	6.19	6-pyruvoyl tetrahydrobiopterin synthase OS=Homo sapiens GN=PTS PE=1 SV=1
PUF60	H0YCP8	38	28260	3	1	2	1	0.18	Poly(U)-binding-splicing factor PUF60 (Fragment) OS=Homo sapiens GN=PUF60 PE=1 SV=1
PURA	Q00577	59	35003	3	1	3	1	0.14	Transcriptional activator protein Pur-alpha OS=Homo sapiens GN=PURA PE=1 SV=2
PURB	Q96QR8	30	33392	2	1	2	1	0.15	Transcriptional activator protein Pur-beta OS=Homo sapiens GN=PURB PE=1 SV=2
PWP1	B4DJV5	35	49464	2	1	2	1	0.10	Periodic tryptophan protein 1 homolog OS=Homo sapiens GN=PWP1 PE=1 SV=1
PYM1	Q9BRP8	147	22691	7	6	6	6	2.46	Isoform 2 of Partner of Y14 and mago OS=Homo sapiens GN=WIBG
RAB1A	P62820	76	22891	5	2	3	2	0.51	Ras-related protein Rab-1A OS=Homo sapiens GN=RAB1A PE=1 SV=3
RAB5C	P51148	60	27247	2	1	2	1	0.19	Isoform 2 of Ras-related protein Rab-5C OS=Homo sapiens GN=RAB5C
RAB9A	P51151	27	23108	1	1	1	1	0.22	Ras-related protein Rab-9A OS=Homo sapiens GN=RAB9A PE=1 SV=1
RACK1	P63244	167	35511	7	5	6	5	0.94	Guanine nucleotide-binding protein subunit beta-2-like 1 OS=Homo sapiens GN=GNB2L1 PE=1 SV=3
RANBP9	Q96S59	24	43607	1	1	1	1	0.11	Isoform 2 of Ran-binding protein 9 OS=Homo sapiens GN=RANBP9
RARS	P54136	222	76129	12	9	12	9	0.75	Arginine--tRNA ligase, cytoplasmic OS=Homo sapiens GN=RARS PE=1 SV=2
RAVER1	K7EKR9	54	10993	1	1	1	1	0.52	Ribonucleoprotein PTB-binding 1 (Fragment) OS=Homo sapiens GN=RAVER1 PE=1 SV=1
RBBP7	Q5JPO1	62	31883	4	2	2	2	0.34	Histone-binding protein RBBP7 (Fragment) OS=Homo sapiens GN=RBBP7 PE=1 SV=1
RBM39	H0Y4X3	64	36888	3	3	3	3	0.47	RNA-binding protein 39 (Fragment) OS=Homo sapiens GN=RBM39 PE=1 SV=2
RBM47	A0AV96	39	57340	1	1	1	1	0.09	Isoform 2 of RNA-binding protein 47 OS=Homo sapiens GN=RBM47
RBM6	P78332	98	129192	6	4	6	4	0.16	RNA-binding protein 6 OS=Homo sapiens GN=RBM6 PE=1 SV=5
RCL1	Q9Y2P8	23	41273	1	1	1	1	0.12	RNA 3'-terminal phosphate cyclase-like protein OS=Homo sapiens GN=RCL1 PE=1 SV=3
RFC1	P35251	65	128617	4	2	4	2	0.08	Isoform 2 of Replication factor C subunit 1 OS=Homo sapiens GN=RFC1
RFC2	P35250	40	39588	2	1	2	1	0.13	Replication factor C subunit 2 OS=Homo sapiens GN=RFC2 PE=1 SV=3
RFC3	P40938	51	35589	3	1	3	1	0.14	Isoform 2 of Replication factor C subunit 3 OS=Homo sapiens GN=RFC3
RFC4	P35249	78	40170	4	3	4	3	0.42	Replication factor C subunit 4 OS=Homo sapiens GN=RFC4 PE=1 SV=1
RFC5	P40937	32	36367	3	1	2	1	0.14	Isoform 2 of Replication factor C subunit 5 OS=Homo sapiens GN=RFC5
RHOC	Q5JR08	34	21794	1	1	1	1	0.24	Rho-related GTP-binding protein RhoC (Fragment) OS=Homo sapiens GN=RHOC PE=1 SV=2
ROCK1	Q13464	140	159102	13	7	12	7	0.23	Rho-associated protein kinase 1 OS=Homo sapiens GN=ROCK1 PE=1 SV=1
RPA1	I3L4R8	39	35378	2	1	2	1	0.14	Replication protein A 70 kDa DNA-binding subunit (Fragment) OS=Homo sapiens GN=RPA1 PE=1 SV=1
RPL10	X1WI28	540	23416	20	17	13	10	12.46	60S ribosomal protein L10 (Fragment) OS=Homo sapiens GN=RPL10 PE=1 SV=1
RPL15	P61313	191	24245	9	8	8	7	2.88	60S ribosomal protein L15 OS=Homo sapiens GN=RPL15 PE=1 SV=2
RPL23A	K7EJV9	260	19354	7	7	7	7	4.44	60S ribosomal protein L23a (Fragment) OS=Homo sapiens GN=RPL23A PE=1 SV=1
RPL26	J3KTJ8	159	11491	9	7	6	6	15.54	60S ribosomal protein L26 (Fragment) OS=Homo sapiens GN=RPL26 PE=4 SV=2
RPL32	D3YTB1	19	15721	1	1	1	1	0.34	60S ribosomal protein L32 (Fragment) OS=Homo sapiens GN=RPL32 PE=1 SV=1
RPL34	P49207	121	13513	5	5	5	5	4.55	60S ribosomal protein L34 OS=Homo sapiens GN=RPL34 PE=1 SV=1
RPL36A	J3KQN4	41	16653	4	1	4	1	0.32	60S ribosomal protein L36a OS=Homo sapiens GN=RPL36A PE=3 SV=1
RPL37A	C9J4Z3	99	7847	6	4	4	2	2.19	60S ribosomal protein L37a OS=Homo sapiens GN=RPL37A PE=1 SV=1
RPL6	Q02878	501	32765	13	13	11	11	3.85	60S ribosomal protein L6 OS=Homo sapiens GN=RPL6 PE=1 SV=1
RPL8	P62917	302	28235	12	10	10	8	3.47	60S ribosomal protein L8 OS=Homo sapiens GN=RPL8 PE=1 SV=2
RPLP2	P05387	134	11658	4	4	4	4	3.90	60S acidic ribosomal protein P2 OS=Homo sapiens GN=RPLP2 PE=1 SV=1
RPN2	Q5JYR7	90	37200	3	2	3	2	0.29	Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit 2 (Fragment) OS=Homo sapiens GN=RPN2 PE=1 SV=2
RPS16	M0R210	279	14524	11	10	8	8	11.91	40S ribosomal protein S16 OS=Homo sapiens GN=RPS16 PE=1 SV=1
RPS27	Q5T4L4	201	7523	7	7	4	4	19.62	40S ribosomal protein S27 OS=Homo sapiens GN=RPS27 PE=1 SV=1
RPS27A	P62979	302	18296	17	14	7	6	4.95	Ubiquitin-40S ribosomal protein S27a OS=Homo sapiens GN=RPS27A PE=1 SV=2
RPS27L	Q71UM5	90	9813	4	3	2	2	3.12	40S ribosomal protein S27-like OS=Homo sapiens GN=RPS27L PE=1 SV=1
RPS8	P62241	318	24475	12	10	8	8	3.64	40S ribosomal protein S8 OS=Homo sapiens GN=RPS8 PE=1 SV=2
RPUSD3	Q6P087	34	37264	2	1	2	1	0.13	Isoform 2 of RNA pseudouridylate synthase domain-containing protein 3 OS=Homo sapiens GN=RPUSD3
RRBP1	F8W7S5	75	84674	7	2	6	2	0.12	Ribosome-binding protein 1 OS=Homo sapiens GN=RRBP1 PE=1 SV=1
RRP1B	Q14684	60	82467	1	1	1	1	0.06	Isoform 2 of Ribosomal RNA processing protein 1 homolog B OS=Homo sapiens GN=RRP1B
RTRAF	Q9Y224	268	28165	7	7	5	5	1.30	UPF0568 protein C14orf166 OS=Homo sapiens GN=C14orf166 PE=1 SV=1
RUVBL1	Q9Y265	60	42499	2	2	2	2	0.25	Isoform 2 of RuvB-like 1 OS=Homo sapiens GN=RUVBL1
RUVBL2	M0R0Y3	131	38845	5	5	5	5	0.83	RuvB-like 2 OS=Homo sapiens GN=RUVBL2 PE=1 SV=1
SAFB2	Q14151	70	107921	5	2	5	2	0.09	Scaffold attachment factor B2 OS=Homo sapiens GN=SAFB2 PE=1 SV=1
SAMM50	Q9Y512	41	52342	2	1	2	1	0.09	Sorting and assembly machinery component 50 homolog OS=Homo sapiens GN=SAMMSO PE=1 SV=1
SAR1A	Q9NR31	20	17603	3	1	2	1	0.30	Isoform 2 of GTP-binding protein SAR1a OS=Homo sapiens GN=SAR1A
SAR1B	Q9Y6B6	37	22510	3	1	2	1	0.23	GTP-binding protein SAR1b OS=Homo sapiens GN=SAR1B PE=1 SV=1
SART1	O43290	77	90371	3	2	3	2	0.11	U4/U6.U5 tri-snRNP-associated protein 1 OS=Homo sapiens GN=SART1 PE=1 SV=1
SCYL2	Q6P3W7	98	104327	3	3	2	2	0.09	SCY1-like protein 2 OS=Homo sapiens GN=SCYL2 PE=1 SV=1
SERPINH1	E9PPV6	94	35758	4	3	4	3	0.48	Serpin H1 OS=Homo sapiens GN=SERPINH1 PE=1 SV=1
SF1	Q15637	44	68817	1	1	1	1	0.07	Isoform 2 of Splicing factor 1 OS=Homo sapiens GN=SF1
SF3A1	Q15459	452	88888	23	19	19	17	1.60	Splicing factor 3A subunit 1 OS=Homo sapiens GN=SF3A1 PE=1 SV=1
SF3A2	Q15428	79	49338	4	3	4	3	0.33	Splicing factor 3A subunit 2 OS=Homo sapiens GN=SF3A2 PE=1 SV=2
SF3A3	Q12874	304	59154	11	9	9	7	0.75	Splicing factor 3A subunit 3 OS=Homo sapiens GN=SF3A3 PE=1 SV=1
SF3B1	O75533	1022	146479	41	34	32	27	1.55	Splicing factor 3B subunit 1 OS=Homo sapiens GN=SF3B1 PE=1 SV=3
SF3B4	Q5SZ64	106	20733	2	2	2	2	0.57	Splicing factor 3B subunit 4 (Fragment) OS=Homo sapiens GN=SF3B4 PE=1 SV=1
SKP1	E5RJR5	65	18879	1	1	1	1	0.09	S-phase kinase-associated protein 1 OS=Homo sapiens GN=SKP1 PE=1 SV=1
SLC1A3	P43003	90	55284	2	1	2	1	0.09	Isoform 2 of Excitatory amino acid transporter 1 OS=Homo sapiens GN=SLC1A3
SLC25A1	B4DP62	54	23423	1	1	1	1	0.22	Solute carrier family 25 (Mitochondrial carrier citrate transporter), member 1, isoform CRA_b OS=Homo sapiens GN=SLC25A1 PE=1 SV=1
SLC25A13	Q9UJ50	18	74656	2	1	2	1	0.07	Isoform 2 of Calcium-binding mitochondrial carrier protein Aralar2 OS=Homo sapiens GN=SLC25A13
SLC25A5	P05141	88	33059	3	3	3	3	0.53	ADP/ATP translocase 2 OS=Homo sapiens GN=SLC25A5 PE=1 SV=7
SLC3A2	J3KPF3	171	68230	6	3	6	3	0.23	4F2 cell-surface antigen heavy chain OS=Homo sapiens GN=SLC3A2 PE=1 SV=1
SLC7A5	Q01650	66	55659	1	1	1	1	0.05	Large neutral amino acids transporter small subunit 1 OS=Homo sapiens GN=SLC7A5 PE=1 SV=2
SLIRP	Q9GZT3	18	12171	1	1	1	1	0.46	Isoform 2 of SRA stem-loop-interacting RNA-binding protein, mitochondrial OS=Homo sapiens GN=SLIRP
SND1	Q7KZF4	163	102618	12	6	11	6	0.32	Staphylococcal nuclease domain-containing protein 1 OS=Homo sapiens GN=SND1 PE=1 SV=1
SNRNP200	O75643	135	246006	17	4	13	4	0.08	U5 small nuclear ribonucleoprotein 200 kDa helicase OS=Homo sapiens GN=SNRNP200 PE=1 SV=2
SNRNP40	Q96DI7	18	45001	1	1	1	1	0.11	Isoform 2 of U5 small nuclear ribonucleoprotein 40 kDa protein OS=Homo sapiens GN=SNRNP40
SNRNP70	P08621	43	51583	4	2	4	2	0.20	U1 small nuclear ribonucleoprotein 70 kDa OS=Homo sapiens GN=SNRNP70 PE=1 SV=2
SNRPA1	P09661	288	28512	9	8	8	7	2.16	U2 small nuclear ribonucleoprotein A' OS=Homo sapiens GN=SNRPA1 PE=1 SV=2
SNRPB2	P08579	44	25470	2	1	2	1	0.20	U2 small nuclear ribonucleoprotein B'' OS=Homo sapiens GN=SNRPB2 PE=1 SV=1
SNRPD1	P62314	113	13273	3	3	2	2	1.01	Small nuclear ribonucleoprotein Sm D1 OS=Homo sapiens GN=SNRPD1 PE=1 SV=1
SNRPD2	P62316	169	13632	7	7	6	6	9.82	Small nuclear ribonucleoprotein Sm D2 OS=Homo sapiens GN=SNRPD2 PE=1 SV=1
SNRNPG15	A8MWD9	90	8595	6	3	4	2	4.01	Putative small nuclear ribonucleoprotein G-like protein 15 OS=Homo sapiens GN=SNRPGP15 PE=5 SV=2
SNRPN	J3QLE5	488	17706	15	15	7	7	9.64	Small nuclear ribonucleoprotein-associated protein N (Fragment) OS=Homo sapiens GN=SNRPN PE=4 SV=1
SORBS2	O94875	21	135708	2	1	2	1	0.04	Isoform 11 of Sorbin and SH3 domain-containing protein 2 OS=Homo sapiens GN=SORBS2
SPDEF	O95238	20	35920	2	1	2	1	0.14	Isoform 2 of SAM pointed domain-containing Ets transcription factor OS=Homo sapiens GN=SPDEF
SPINDOC	Q9BUA3	73	41297	7	3	6	3	0.41	Uncharacterized protein C11orf84 OS=Homo sapiens GN=C11orf84 PE=1 SV=1
SPTAN1	Q13813	2509	282906	83	74	71	63	1.92	Isoform 3 of Spectrin alpha chain, non-erythrocytic 1 OS=Homo sapiens GN=SPTAN1
SPTLC1	O15269	121	53281	2	2	2	2	0.19	Serine palmitoyltransferase 1 OS=Homo sapiens GN=SPTLC1 PE=1 SV=1
SPTLC3	H0Y733	18	13620	2	2	1	1	0.41	Serine palmitoyltransferase 3 (Fragment) OS=Homo sapiens GN=SPTLC3 PE=4 SV=1
SRP68	Q9UHB9	75	67775	2	1	2	1	0.07	Isoform 2 of Signal recognition particle subunit SRP68 OS=Homo sapiens GN=SRP68
SRPK2	H7CSL6	30	33605	2	1	1	1	0.15	SRSF protein kinase 2 (Fragment) OS=Homo sapiens GN=SRPK2 PE=1 SV=1
SRPRB	Q9Y5M8	98	29912	5	4	5	4	0.88	Signal recognition particle receptor subunit beta OS=Homo sapiens GN=SRPRB PE=1 SV=3
SRRT	H7C3A1	68	57154	1	1	1	1	0.09	Serrate RNA effector molecule homolog (Fragment) OS=Homo sapiens GN=SRRT PE=1 SV=1
SRSF1	J3KTL2	322	28426	9	9	7	7	2.18	Serine/arginine-rich-splicing factor 1 OS=Homo sapiens GN=SRSF1 PE=1 SV=1
SRSF3	P84103	104	14422	5	4	4	4	2.63	Isoform 2 of Serine/arginine-rich splicing factor 3 OS=Homo sapiens GN=SRSF3
SRSF4	Q08170	69	56759	4	3	4	3	0.28	Serine/arginine-rich splicing factor 4 OS=Homo sapiens GN=SRSF4 PE=1 SV=2
SRSF7	C9JAB2	133	27140	6	6	4	4	1.37	Serine/arginine-rich splicing factor 7 OS=Homo sapiens GN=SRSF7 PE=1 SV=1
SSR1	C9IZQ1	119	33866	2	2	1	1	0.15	Translocon-associated protein subunit alpha OS=Homo sapiens GN=SSR1 PE=1 SV=1
SSR4	P51571	125	19158	3	3	3	3	1.07	Translocon-associated protein subunit delta OS=Homo sapiens GN=SSR4 PE=1 SV=1
STAU2	Q9NUL3	37	43397	4	2	4	2	0.24	Isoform 6 of Double-stranded RNA-binding protein Staufen homolog 2 OS=Homo sapiens GN=STAU2
STOM	P27105	445	31882	13	10	11	9	3.37	Erythrocyte band 7 integral membrane protein OS=Homo sapiens GN=STOM PE=1 SV=3
STRAP	B4DNJ6	37	40095	3	2	3	2	0.26	Serine-threonine kinase receptor-associated protein OS=Homo sapiens GN=STRAP PE=1 SV=1
STX6	O43752	13	17480	2	1	2	1	0.31	Isoform 2 of Syntaxin-6 OS=Homo sapiens GN=STX6
SUB1	P53999	43	14386	1	1	1	1	0.38	Activated RNA polymerase II transcriptional coactivator p15 OS=Homo sapiens GN=SUB1 PE=1 SV=3
SVIL	O95425	39	202262	2	1	2	1	0.02	Isoform 12 of Supervillin OS=Homo sapiens GN=SVIL
SYTL2	Q9HCH5	28	84684	2	1	1	1	0.06	Isoform 12 of Synaptotagmin-like protein 2 OS=Homo sapiens GN=SYTL2
TAB2	Q9NYJ8	77	58323	3	2	3	2	0.18	Isoform 2 of TGF-beta-activated kinase 1 and MAP3K7-binding protein 2 OS=Homo sapiens GN=TAB2
TCP1	P17987	174	60819	10	6	9	5	0.59	T-complex protein 1 subunit alpha OS=Homo sapiens GN=TCP1 PE=1 SV=1
TECR	Q9NZ01	40	36410	2	1	2	1	0.14	Very-long-chain enoyl-CoA reductase OS=Homo sapiens GN=TECR PE=1 SV=1
TFRC	G3V0E5	80	76259	3	3	3	3	0.20	Transferrin receptor (P90, CD71), isoform CRA_c OS=Homo sapiens GN=TFRC PE=1 SV=1
TGFBR1	B4DYD6	29	49642	3	1	2	1	0.10	TGF-beta receptor type-1 OS=Homo sapiens GN=TGFBR1 PE=1 SV=1
TIMM50	M0R0D3	68	28432	3	1	3	1	0.18	Mitochondrial import inner membrane translocase subunit TIM50 (Fragment) OS=Homo sapiens GN=TIMM50 PE=1 SV=1
TJP2	Q9UDY2	68	131516	4	3	4	3	0.11	Isoform C1 of Tight junction protein ZO-2 OS=Homo sapiens GN=TJP2
TKT	B4E022	146	63410	7	5	6	4	0.35	Transketolase OS=Homo sapiens GN=TKT PE=1 SV=1
TMED10	G3V2K7	65	17008	1	1	1	1	0.32	Transmembrane emp24 domain-containing protein 10 OS=Homo sapiens GN=TMED10 PE=1 SV=1
TMED2	F5GX39	36	13679	1	1	1	1	0.41	Transmembrane emp24 domain-containing protein 2 OS=Homo sapiens GN=TMED2 PE=1 SV=1
TMEM214	H7BZ30	68	15615	2	1	2	1	0.35	Transmembrane protein 214 (Fragment) OS=Homo sapiens GN=TMEM214 PE=1 SV=1
TMEM59	Q9BKS4	85	36884	4	2	3	2	0.29	Transmembrane protein 59 OS=Homo sapiens GN=TMEM59 PE=1 SV=1
TMEM9	B1ALM5	16	21458	2	1	2	1	0.24	Transmembrane protein 9 OS=Homo sapiens GN=TMEM9 PE=1 SV=1
TMOD3	Q9NVL9	85	39741	3	3	3	3	0.43	Tropomodulin-3 OS=Homo sapiens GN=TMOD3 PE=1 SV=1
TNFRSF1A	F5H061	27	47081	2	1	2	1	0.11	Tumor necrosis factor receptor superfamily member 1A OS=Homo sapiens GN=TNFRSF1A PE=4 SV=1
TPI1	P60174	139	26938	6	6	6	6	1.84	Isoform 2 of Triosephosphate isomerase OS=Homo sapiens GN=TPI1

1475

Gene	Accession								Description
MCU	S4R468	60	19186	1	1	1	1	0.28	Calcium uniporter protein, mitochondrial (Fragment) OS=Homo sapiens GN=MCU PE=4 SV=1
MDH2	G3XAL0	58	24921	1	1	1	1	0.21	Malate dehydrogenase OS=Homo sapiens GN=MDH2 PE=1 SV=1
MGST2	Q99735	62	8524	1	1	1	1	0.71	Isoform 2 of Microsomal glutathione S-transferase 2 OS=Homo sapiens GN=MGST2
MGST3	Q5VV87	93	14427	1	1	1	1	0.38	Microsomal glutathione S-transferase 3 OS=Homo sapiens GN=MGST3 PE=1 SV=1
MISP	Q8IVT2	45	75482	4	1	2	1	0.06	Mitotic interactor and substrate of Plk1 OS=Homo sapiens GN=MISP PE=1 SV=1
MLEC	H0YG07	41	19893	2	1	2	1	0.27	Malectin (Fragment) OS=Homo sapiens GN=MLEC PE=1 SV=1
MOB1B	Q7L9L4	50	25654	1	1	1	1	0.20	Isoform 2 of MOB kinase activator 1B OS=Homo sapiens GN=MOB1B
MOGS	Q13724	111	92032	3	2	3	2	0.11	Mannosyl-oligosaccharide glucosidase OS=Homo sapiens GN=MOGS PE=1 SV=5
MORF4L2	Q15014	43	32345	2	1	2	1	0.16	Mortality factor 4-like protein 2 OS=Homo sapiens GN=MORF4L2 PE=1 SV=1
MRPL11	Q9Y3B7	58	20727	3	2	3	2	0.57	39S ribosomal protein L11, mitochondrial OS=Homo sapiens GN=MRPL11 PE=1 SV=1
MRPL14	Q6P1L8	38	16165	2	1	2	1	0.33	39S ribosomal protein L14, mitochondrial OS=Homo sapiens GN=MRPL14 PE=1 SV=1
MRPL22	J3KQY1	105	26686	4	4	4	4	1.02	39S ribosomal protein L22, mitochondrial OS=Homo sapiens GN=MRPL22 PE=1 SV=1
MRPL24	Q96A35	40	25013	1	1	1	1	0.21	39S ribosomal protein L24, mitochondrial OS=Homo sapiens GN=MRPL24 PE=1 SV=1
MRPL3	H0Y9G6	57	40898	5	2	5	2	0.26	39S ribosomal protein L3, mitochondrial (Fragment) OS=Homo sapiens GN=MRPL3 PE=1 SV=1
MRPL30	Q8TCC3	15	21939	1	1	1	1	0.24	Isoform 2 of 39S ribosomal protein L30, mitochondrial OS=Homo sapiens GN=MRPL30
MRPL37	S4R369	60	55659	4	1	4	1	0.09	39S ribosomal protein L37, mitochondrial OS=Homo sapiens GN=MRPL37 PE=1 SV=1
MRPL38	Q96DV4	65	44968	2	2	2	2	0.23	39S ribosomal protein L38, mitochondrial OS=Homo sapiens GN=MRPL38 PE=1 SV=1
MRPL4	K7ES61	151	33771	6	4	5	3	0.52	39S ribosomal protein L4, mitochondrial (Fragment) OS=Homo sapiens GN=MRPL4 PE=1 SV=1
MRPL40	Q9NQ50	15	24475	1	1	1	1	0.21	39S ribosomal protein L40, mitochondrial OS=Homo sapiens GN=MRPL40 PE=1 SV=1
MRPL44	Q9H9J2	127	37854	7	4	6	4	0.64	39S ribosomal protein L44, mitochondrial OS=Homo sapiens GN=MRPL44 PE=1 SV=1
MRPL45	Q9BRJ2	78	35613	3	2	3	2	0.30	39S ribosomal protein L45, mitochondrial OS=Homo sapiens GN=MRPL45 PE=1 SV=1
MRPL48	F5H8D0	45	13699	2	1	2	1	0.41	39S ribosomal protein L48, mitochondrial OS=Homo sapiens GN=MRPL48 PE=1 SV=1
MRPL49	H0YDP7	68	14731	3	1	3	1	0.37	39S ribosomal protein L49, mitochondrial (Fragment) OS=Homo sapiens GN=MRPL49 PE=1 SV=1
MRPL9	Q9BYD2	37	30395	2	1	2	1	0.17	39S ribosomal protein L9, mitochondrial OS=Homo sapiens GN=MRPL9 PE=1 SV=2
MRPS2	Q5T8A0	44	30745	1	1	1	1	0.17	28S ribosomal protein S2, mitochondrial OS=Homo sapiens GN=MRPS2 PE=1 SV=1
MRPS23	J3QLR8	56	17563	1	1	1	1	0.30	28S ribosomal protein S23, mitochondrial OS=Homo sapiens GN=MRPS23 PE=1 SV=1
MRPS27	G5EA06	37	41588	3	3	3	3	0.40	28S ribosomal protein S27, mitochondrial OS=Homo sapiens GN=MRPS27 PE=1 SV=1
MRPS28	H0YAT2	46	16063	2	1	2	1	0.34	28S ribosomal protein S28, mitochondrial (Fragment) OS=Homo sapiens GN=MRPS28 PE=1 SV=1
MRPS30	Q9NP92	23	50960	1	1	1	1	0.10	28S ribosomal protein S30, mitochondrial OS=Homo sapiens GN=MRPS30 PE=1 SV=2
MRPS34	C9JJ19	44	26373	2	1	2	1	0.19	28S ribosomal protein S34, mitochondrial OS=Homo sapiens GN=MRPS34 PE=1 SV=2
MRPS7	J3QKW2	70	20882	3	3	2	2	0.57	28S ribosomal protein S7, mitochondrial OS=Homo sapiens GN=MRPS7 PE=1 SV=1
MRPS9	P82933	131	46034	6	4	6	4	0.51	28S ribosomal protein S9, mitochondrial OS=Homo sapiens GN=MRPS9 PE=1 SV=2
MSI2	J3KTC1	62	15507	1	1	1	1	0.35	Protein turtle homolog B (Fragment) OS=Homo sapiens GN=IGSF9B PE=1 SV=2
MSN	P26038	23	67892	2	1	2	1	0.07	Moesin OS=Homo sapiens GN=MSN PE=1 SV=3
MTDH	Q86UE4	254	63856	12	9	10	9	0.95	Protein LYRIC OS=Homo sapiens GN=MTDH PE=1 SV=2
MTERF3	Q96E29	29	48055	2	1	2	1	0.10	Transcription termination factor 3, mitochondrial OS=Homo sapiens GN=MTERF3 PE=1 SV=2
MTHFD1	F5H2F4	44	111342	3	1	3	1	0.04	C-1-tetrahydrofolate synthase, cytoplasmic OS=Homo sapiens GN=MTHFD1 PE=1 SV=1
MUM1	Q2TAK8	14	71715	2	1	1	1	0.07	Isoform 2 of PWWP domain-containing protein MUM1 OS=Homo sapiens GN=MUM1
MYBPB1A	I3L1L3	64	141124	3	2	3	2	0.07	Myb-binding protein 1A (Fragment) OS=Homo sapiens GN=MYBBP1A PE=1 SV=1
MYO1E	Q12965	62	127552	3	1	3	1	0.04	Unconventional myosin-Ie OS=Homo sapiens GN=MYO1E PE=1 SV=2
MYO5B	Q9ULV0	65	215135	4	2	4	2	0.04	Unconventional myosin-Vb OS=Homo sapiens GN=MYO5B PE=1 SV=3
NACA	E9PAV3	62	205979	3	1	2	1	0.02	Nascent polypeptide-associated complex subunit alpha, muscle-specific form OS=Homo sapiens GN=NACA PE=1 SV=1
NAP1L1	H0YH88	120	21179	5	5	4	4	2.01	Nucleosome assembly protein 1-like 1 (Fragment) OS=Homo sapiens GN=NAP1L1 PE=1 SV=1
NAT10	Q9H0A0	85	108084	3	1	3	1	0.04	Isoform 2 of N-acetyltransferase 10 OS=Homo sapiens GN=NAT10
NCBP1	Q09161	61	92864	3	2	3	2	0.11	Nuclear cap-binding protein subunit 1 OS=Homo sapiens GN=NCBP1 PE=1 SV=1
NCOA5	Q9HCD5	75	65725	5	2	5	2	0.15	Nuclear receptor coactivator 5 OS=Homo sapiens GN=NCOA5 PE=1 SV=2
NDC1	Q9BTX1	61	61768	4	2	4	2	0.17	Isoform 4 of Nucleoporin NDC1 OS=Homo sapiens GN=NDC1
NDUFA9	Q16795	56	42654	3	1	3	1	0.12	NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 9, mitochondrial OS=Homo sapiens GN=NDUFA9 PE=1 SV=2
NELFE	E9PD43	25	30046	3	1	3	1	0.17	Negative elongation factor E (Fragment) OS=Homo sapiens GN=NELFE PE=1 SV=1
NEMF	O60524	41	121085	3	1	3	1	0.04	Isoform 3 of Nuclear export mediator factor NEMF OS=Homo sapiens GN=NEMF
NES	P48681	105	177788	6	3	6	3	0.08	Nestin OS=Homo sapiens GN=NES PE=1 SV=3
NFATC4	Q14934	20	91153	4	1	3	1	0.05	Isoform 11 of Nuclear factor of activated T-cells, cytoplasmic 4 OS=Homo sapiens GN=NFATC4
NGRN	Q9NPE2	39	32502	2	1	2	1	0.16	Neugrin OS=Homo sapiens GN=NGRN PE=1 SV=2
NIPSNAP1	Q9BPW8	51	33460	2	2	2	2	0.32	Protein NipSnap homolog 1 OS=Homo sapiens GN=NIPSNAP1 PE=1 SV=1
NKRF	O15226	47	80130	3	1	3	1	0.06	Isoform 2 of NF-kappa-B-repressing factor OS=Homo sapiens GN=NKRF
NME1	P15531	118	19869	4	4	3	3	1.56	Isoform 2 of Nucleoside diphosphate kinase A OS=Homo sapiens GN=NME1
NME2P1	O60361	111	15690	5	4	4	3	2.29	Putative nucleoside diphosphate kinase OS=Homo sapiens GN=NME2P1 PE=5 SV=1
NOTCH2	Q04721	78	279082	4	3	4	3	0.05	Neurogenic locus notch homolog protein 2 OS=Homo sapiens GN=NOTCH2 PE=1 SV=3
NSMCE1	H3BR31	13	8492	2	1	1	1	0.71	Non-structural maintenance of chromosomes element 1 homolog OS=Homo sapiens GN=NSMCE1 PE=4 SV=1
NSUN5	Q96P11	24	51120	2	1	2	1	0.10	Isoform 2 of Probable 28S rRNA (cytosine-C[5])-methyltransferase OS=Homo sapiens GN=NSUN5
NT5C2	P49902	69	61858	2	2	2	2	0.16	Isoform 2 of Cytosolic purine 5'-nucleotidase OS=Homo sapiens GN=NT5C2
NUBP2	H3BNF0	36	26764	1	1	1	1	0.19	Cytosolic Fe-S cluster assembly factor NUBP2 OS=Homo sapiens GN=NUBP2 PE=1 SV=1
OCIAD1	D6RBN5	19	24409	2	1	2	1	0.21	OCIA domain-containing protein 1 OS=Homo sapiens GN=OCIAD1 PE=1 SV=1
OGT	O15294	60	116887	5	5	5	5	0.22	Isoform 1 of UDP-N-acetylglucosamine--peptide N-acetylglucosaminyltransferase 110 kDa subunit OS=Homo sapiens GN=OGT
P4HB	P07237	99	57480	6	4	5	4	0.39	Protein disulfide-isomerase OS=Homo sapiens GN=P4HB PE=1 SV=3
PABPC1	E7EQV3	1430	65878	45	39	30	28	8.92	Polyadenylate-binding protein 1 OS=Homo sapiens GN=PABPC1 PE=1 SV=1
PACSIN3	Q9UKS6	71	48799	3	1	3	1	0.10	Protein kinase C and casein kinase substrate in neurons protein 3 OS=Homo sapiens GN=PACSIN3 PE=1 SV=2
PAIP1	D6REB4	86	40225	2	2	2	2	0.26	Uncharacterized protein OS=Homo sapiens GN=PAIP1 PE=1 SV=1
PCBP2	F8VZX2	409	34121	13	12	9	8	2.46	Poly(rC)-binding protein 2 OS=Homo sapiens GN=PCBP2 PE=1 SV=1
PCNA	P12004	53	29092	3	1	3	1	0.18	Proliferating cell nuclear antigen OS=Homo sapiens GN=PCNA PE=1 SV=1
PDCD6	H0Y9X3	37	12277	2	1	2	1	0.46	Programmed cell death protein 6 (Fragment) OS=Homo sapiens GN=PDCD6 PE=1 SV=1
PDE4D	Q08499	24	24584	2	1	1	1	0.21	Isoform 12 of cAMP-specific 3',5'-cyclic phosphodiesterase 4D OS=Homo sapiens GN=PDE4D
PDIA3	P30101	53	57146	1	1	1	1	0.09	Protein disulfide-isomerase A3 OS=Homo sapiens GN=PDIA3 PE=1 SV=4
PDIA6	Q15084	33	54380	1	1	1	1	0.09	Isoform 2 of Protein disulfide-isomerase A6 OS=Homo sapiens GN=PDIA6
PEG10	Q86TG7	85	80692	3	2	3	2	0.12	Retrotransposon-derived protein PEG10 OS=Homo sapiens GN=PEG10 PE=1 SV=1
PFKFB2	B0FLL2	61	55199	4	2	4	2	0.19	6-phosphofructo-2-kinase/fructose-2,6-biphosphatase 2 transcript variant 3 OS=Homo sapiens GN=PFKFB2 PE=1 SV=1
PFN1	P07737	140	15216	4	4	4	4	2.40	Profilin-1 OS=Homo sapiens GN=PFN1 PE=1 SV=2
PGAM4	Q8N0Y7	29	28930	2	1	2	1	0.18	Probable phosphoglycerate mutase 4 OS=Homo sapiens GN=PGAM4 PE=2 SV=1
PHF5A	Q7RTV0	75	13138	3	2	3	2	1.03	PHD finger-like domain-containing protein 5A OS=Homo sapiens GN=PHF5A PE=1 SV=1
PI4KA	J3KN10	19	239244	4	1	4	1	0.02	Phosphatidylinositol 4-kinase alpha OS=Homo sapiens GN=PI4KA PE=1 SV=1
PKM	P14618	333	58470	12	9	11	9	1.07	Pyruvate kinase PKM OS=Homo sapiens GN=PKM PE=1 SV=4
PKP2	Q99959	59	97868	2	1	2	1	0.05	Plakophilin-2 OS=Homo sapiens GN=PKP2 PE=1 SV=2
PLEC	Q15149	####	533462	317	290	228	210	6.25	Plectin OS=Homo sapiens GN=PLEC PE=1 SV=3
PLIN4	Q96Q06	115	135204	6	5	6	5	0.19	Perilipin-4 OS=Homo sapiens GN=PLIN4 PE=2 SV=2
PLP2	Q04941	23	17022	1	1	1	1	0.32	Proteolipid protein 2 OS=Homo sapiens GN=PLP2 PE=1 SV=1
POLDIP3	Q9BY77	18	43096	2	1	1	1	0.12	Isoform 2 of Polymerase delta-interacting protein 3 OS=Homo sapiens GN=POLDIP3
POP1	Q99575	48	116346	7	4	7	4	0.18	Ribonucleases P/MRP protein subunit POP1 OS=Homo sapiens GN=POP1 PE=1 SV=2
PPFIBP1	Q86W92	23	97303	3	1	3	1	0.05	Isoform 3 of Liprin-beta-1 OS=Homo sapiens GN=PPFIBP1
PPIA	P62937	320	18229	10	10	8	8	9.05	Peptidyl-prolyl cis-trans isomerase A OS=Homo sapiens GN=PPIA PE=1 SV=2
PPIB	P23284	43	23785	2	1	2	1	0.22	Peptidyl-prolyl cis-trans isomerase B OS=Homo sapiens GN=PPIB PE=1 SV=2
PPP1CC	F8VYE8	26	35669	2	2	2	2	0.30	Serine/threonine-protein phosphatase OS=Homo sapiens GN=PPP1CC PE=1 SV=1
PPP2R1A	P30153	69	66065	7	2	7	2	0.15	Serine/threonine-protein phosphatase 2A 65 kDa regulatory subunit A alpha isoform OS=Homo sapiens GN=PPP2R1A PE=1 SV=4
PPP2R2A	E5RFR9	64	23448	2	2	2	2	0.49	Serine/threonine-protein phosphatase 2A 55 kDa regulatory subunit B alpha isoform (Fragment) OS=Homo sapiens GN=PPP2R2A PE=1 SV=1
PRDX2	P32119	103	22049	5	4	5	4	1.33	Peroxiredoxin-2 OS=Homo sapiens GN=PRDX2 PE=1 SV=5
PRDX4	A6NJJ0	65	21403	3	1	2	1	0.24	Peroxiredoxin-4 (Fragment) OS=Homo sapiens GN=PRDX4 PE=1 SV=1
PRKACA	P17612	21	40042	3	1	3	1	0.12	Isoform 2 of cAMP-dependent protein kinase catalytic subunit alpha OS=Homo sapiens GN=PRKACA
PRKAR1A	X6RAV4	23	15032	2	1	2	1	0.35	cAMP-dependent protein kinase type I-alpha regulatory subunit OS=Homo sapiens GN=PRKAR1A PE=1 SV=1
PRKDC	E7EUY0	499	470050	26	16	25	15	0.16	DNA-dependent protein kinase catalytic subunit OS=Homo sapiens GN=PRKDC PE=1 SV=1
PRKRA	O75569	17	33672	2	1	2	1	0.15	Isoform 2 of Interferon-inducible double-stranded RNA-dependent protein kinase activator A OS=Homo sapiens GN=PRKRA
PRMT1	H7C2I1	59	43061	5	3	5	3	0.39	Protein arginine N-methyltransferase 1 OS=Homo sapiens GN=PRMT1 PE=1 SV=1
PRPF19	Q9UMS4	130	55603	7	5	5	5	0.53	Pre-mRNA-processing factor 19 OS=Homo sapiens GN=PRPF19 PE=1 SV=1
PRPF31	Q8WWY3	300	55649	18	12	9	9	0.97	U4/U6 small nuclear ribonucleoprotein Prp31 OS=Homo sapiens GN=PRPF31 PE=1 SV=2
PRPF4	O43172	81	58969	2	1	2	1	0.08	Isoform 2 of U4/U6 small nuclear ribonucleoprotein Prp4 OS=Homo sapiens GN=PRPF4
PRPF8	Q6P2Q9	63	274738	6	2	4	2	0.04	Pre-mRNA-processing-splicing factor 8 OS=Homo sapiens GN=PRPF8 PE=1 SV=2
PRPS2	P11908	585	35431	15	12	9	7	1.53	Isoform 2 of Ribose-phosphate pyrophosphokinase 2 OS=Homo sapiens GN=PRPS2
PRPSAP1	B4DP31	632	31474	17	15	8	8	3.44	Phosphoribosyl pyrophosphate synthase-associated protein 1 OS=Homo sapiens GN=PRPSAP1 PE=1 SV=1
PRPSAP2	O60256	602	41299	18	16	10	9	2.51	Phosphoribosyl pyrophosphate synthase-associated protein 2 OS=Homo sapiens GN=PRPSAP2 PE=1 SV=1
PRRC2C	E7EPN9	127	309159	13	6	12	6	0.10	Isoform PRRC2C OS=Homo sapiens GN=PRRC2C PE=1 SV=1
PRSS56	P0CW18	48	65583	1	1	1	1	0.07	Serine protease 56 OS=Homo sapiens GN=PRSS56 PE=1 SV=1
PSMA1	F5GX11	26	26774	1	1	1	1	0.19	Proteasome subunit alpha type-1 OS=Homo sapiens GN=PSMA1 PE=1 SV=1
PSMB2	P49721	42	22993	1	1	1	1	0.23	Proteasome subunit beta type-2 OS=Homo sapiens GN=PSMB2 PE=1 SV=1
PSMB5	H0YJM8	48	13021	1	1	1	1	0.43	Proteasome subunit beta type-5 (Fragment) OS=Homo sapiens GN=PSMB5 PE=1 SV=1
PSMC3	R4GNH3	27	47608	3	1	3	1	0.10	Proteasome subunit OS=Homo sapiens GN=PSMC3 PE=1 SV=1
PSMC4	P43686	41	43595	4	1	4	1	0.11	Isoform 2 of 26S protease regulatory subunit 6B OS=Homo sapiens GN=PSMC4
PSMC5	P62195	147	44927	6	4	6	4	0.52	26S protease regulatory subunit 8 OS=Homo sapiens GN=PSMC5
PSMC6	H0YJC0	30	29019	2	1	2	1	0.17	26S protease regulatory subunit 10B (Fragment) OS=Homo sapiens GN=PSMC6 PE=1 SV=1
PSMD11	O00231	46	47790	3	2	3	2	0.22	Isoform 2 of 26S proteasome non-ATPase regulatory subunit 11 OS=Homo sapiens GN=PSMD11
PSMD13	J3KNQ3	16	40090	1	1	1	1	0.12	Isoform 2 of 26S proteasome non-ATPase regulatory subunit 13 OS=Homo sapiens GN=PSMD13 PE=1 SV=1
PSMD14	O00487	33	34726	1	1	1	1	0.14	26S proteasome non-ATPase regulatory subunit 14 OS=Homo sapiens GN=PSMD14 PE=1 SV=1
PSMD2	Q13200	105	82825	3	2	3	2	0.12	Isoform 2 of 26S proteasome non-ATPase regulatory subunit 2 OS=Homo sapiens GN=PSMD2
PSMD6	C9J7B7	16	16725	1	1	1	1	0.32	26S proteasome non-ATPase regulatory subunit 6 (Fragment) OS=Homo sapiens GN=PSMD6 PE=1 SV=1
PSMD7	P51665	41	37060	2	2	2	2	0.29	26S proteasome non-ATPase regulatory subunit 7 OS=Homo sapiens GN=PSMD7 PE=1 SV=2

1476

TPM1	B7Z596	19	31904	2	1	2	1	0.16	Tropomyosin alpha-1 chain OS=Homo sapiens GN=TPM1 PE=1 SV=1
TRA2B	P62995	105	22036	5	5	4	4	1.33	Isoform 3 of Transformer-2 protein homolog beta OS=Homo sapiens GN=TRA2B
TRAPPC1	**I3L1R8**	**40**	**5018**	**1**	**1**	**1**	**1**	**1.45**	**Trafficking protein particle complex subunit 1 OS=Homo sapiens GN=TRAPPC1 PE=1 SV=1**
TRAPPC11	**Q7Z392**	**97**	**125008**	**4**	**3**	**3**	**3**	**0.12**	**Isoform 3 of Trafficking protein particle complex subunit 11 OS=Homo sapiens GN=TRAPPC11**
TRAPPC12	**Q8WVT3**	**100**	**79781**	**4**	**3**	**4**	**3**	**0.19**	**Trafficking protein particle complex subunit 12 OS=Homo sapiens GN=TRAPPC12 PE=4 SV=3**
TRAPPC13	**A5PLN9**	**89**	**46589**	**3**	**3**	**2**	**2**	**0.22**	**Isoform 2 of Trafficking protein particle complex subunit 13 OS=Homo sapiens GN=TRAPPC13**
TRAPPC2	**P0DI81**	**547**	**20418**	**20**	**17**	**7**	**6**	**6.85**	**Isoform 3 of Trafficking protein particle complex subunit 2 OS=Homo sapiens GN=TRAPPC2**
TRAPPC2L	**H3BUK6**	**63**	**12667**	**4**	**2**	**3**	**2**	**1.08**	**Trafficking protein particle complex subunit 2-like protein OS=Homo sapiens GN=TRAPPC2L PE=1 SV=1**
TRAPPC3	**O43617**	**645**	**20432**	**16**	**16**	**10**	**10**	**14.60**	**Trafficking protein particle complex subunit 3 OS=Homo sapiens GN=TRAPPC3 PE=1 SV=1**
TRAPPC4	**E9PN70**	**62**	**29313**	**3**	**2**	**3**	**2**	**0.38**	**Trafficking protein particle complex subunit 4 OS=Homo sapiens GN=TRAPPC4 PE=1 SV=1**
TRAPPC5	**Q8IUR0**	**235**	**20884**	**10**	**9**	**8**	**7**	**3.80**	**Trafficking protein particle complex subunit 5 OS=Homo sapiens GN=TRAPPC5 SV=1**
TRAPPC8	**J3QQJ5**	**175**	**156369**	**10**	**7**	**8**	**6**	**0.20**	**Trafficking protein particle complex subunit 8 OS=Homo sapiens GN=TRAPPC8 PE=1 SV=1**
TRAPPC9	**Q96Q05**	**139**	**140775**	**12**	**6**	**12**	**6**	**0.22**	**Isoform 2 of Trafficking protein particle complex subunit 9 OS=Homo sapiens GN=TRAPPC9**
TRIM25	Q14258	196	72581	5	5	4	4	0.30	E3 ubiquitin/ISG15 ligase TRIM25 OS=Homo sapiens GN=TRIM25 PE=1 SV=2
TRIM28	M0R0K9	34	49031	1	1	1	1	0.10	Transcription intermediary factor 1-beta (Fragment) OS=Homo sapiens GN=TRIM28 PE=1 SV=1
TRIM29	Q14134	20	64488	1	1	1	1	0.08	Isoform Beta of Tripartite motif-containing protein 29 OS=Homo sapiens GN=TRIM29
TRMT1L	Q7Z2T5	14	82893	3	1	3	1	0.06	TRMT1-like protein OS=Homo sapiens GN=TRMT1L PE=1 SV=2
TROVE2	P10155	46	59129	1	1	1	1	0.08	Isoform 3 of 60 kDa SS-A/Ro ribonucleoprotein OS=Homo sapiens GN=TROVE2
TSR1	Q2NL82	48	92151	3	1	3	1	0.05	Pre-rRNA-processing protein TSR1 homolog OS=Homo sapiens GN=TSR1 PE=1 SV=1
TUBA1C	F5H5D3	1002	58606	29	28	21	21	4.42	Tubulin alpha-1C chain OS=Homo sapiens GN=TUBA1C PE=1 SV=1
TUBA4A	P68366	655	48982	24	23	18	18	4.66	Isoform 2 of Tubulin alpha-4A chain OS=Homo sapiens GN=TUBA4A
TUFM	P49411	111	49852	3	3	3	3	0.33	Elongation factor Tu, mitochondrial OS=Homo sapiens GN=TUFM PE=1 SV=2
TXN	P10599	42	9674	1	1	1	1	0.61	Isoform 2 of Thioredoxin OS=Homo sapiens GN=TXN
U2AF1	Q01081	42	28378	1	1	1	1	0.18	Isoform 2 of Splitting factor U2AF 35 kDa subunit OS=Homo sapiens GN=U2AF1
U2SURP	H7C4P4	50	10874	1	1	1	1	0.53	U2 snRNP-associated SURP motif-containing protein (Fragment) OS=Homo sapiens GN=U2SURP PE=1 SV=1
U2SURP	O15042	343	118618	18	12	15	10	0.49	Isoform 2 of U2 snRNP-associated SURP motif-containing protein OS=Homo sapiens GN=U2SURP
U3KQJ1	U3KQJ1	56	42136	2	1	2	1	0.12	Polymerase delta-interacting protein 2 OS=Homo sapiens GN=POLDIP2 PE=1 SV=1
UBTF	E9PKP7	27	87722	4	2	4	2	0.11	Nucleolar transcription factor 1 OS=Homo sapiens GN=UBTF PE=1 SV=1
UPF2	Q9HAU5	21	148972	3	1	3	1	0.03	Regulator of nonsense transcripts 2 OS=Homo sapiens GN=UPF2 PE=1 SV=1
USP10	Q14694	59	93281	2	1	2	1	0.05	Isoform 2 of Ubiquitin carboxyl-terminal hydrolase 10 OS=Homo sapiens GN=USP10
USP15	Q9Y4E8	109	110482	3	3	3	3	0.14	Isoform 2 of Ubiquitin carboxyl-terminal hydrolase 15 OS=Homo sapiens GN=USP15
VCP	P55072	546	89950	20	16	18	15	1.20	Transitional endoplasmic reticulum ATPase OS=Homo sapiens GN=VCP PE=1 SV=4
VDAC1	P21796	275	30868	7	7	7	7	1.90	Voltage-dependent anion-selective channel protein 1 OS=Homo sapiens GN=VDAC1 PE=1 SV=2
WDR1	D6RD66	42	27066	3	1	3	1	0.11	WD repeat-containing protein 1 (Fragment) OS=Homo sapiens GN=WDR1 PE=1 SV=1
WDR13	Q9H1Z4	13	43637	2	1	2	1	0.11	Isoform 2 of WD repeat-containing protein 13 OS=Homo sapiens GN=WDR13
WDR26	Q9H7D7	37	71385	5	2	4	2	0.14	Isoform 2 of WD repeat-containing protein 26 OS=Homo sapiens GN=WDR26
WDR77	H0Y711	1820	31706	47	37	10	9	2.79	Methylosome protein 50 (Fragment) OS=Homo sapiens GN=WDR77 PE=1 SV=1
WLS	H0YCG9	31	23225	1	1	1	1	0.22	Protein wntless homolog (Fragment) OS=Homo sapiens GN=WLS PE=1 SV=1
WWP1	Q9H0M0	41	92174	4	2	3	2	0.11	Isoform 3 of NEDD4-like E3 ubiquitin-protein ligase WWP1 OS=Homo sapiens GN=WWP1
XPC	Q01831	23	102639	2	1	2	1	0.05	Isoform 2 of DNA repair protein complementing XP-C cells OS=Homo sapiens GN=XPC
YBX3	P16989	301	31928	9	9	6	6	1.42	Isoform 2 of Y-box-binding protein 3 OS=Homo sapiens GN=YBX3
YES1	J3QRU1	30	61861	4	1	3	1	0.08	Tyrosine-protein kinase Yes OS=Homo sapiens GN=YES1 PE=1 SV=1
YME1L1	Q96TA2	79	80011	4	1	4	1	0.06	Isoform 2 of ATP-dependent zinc metalloprotease YME1L1 OS=Homo sapiens GN=YME1L1
YTHDC2	D6R9T8	28	15543	1	1	1	1	0.35	Probable ATP-dependent RNA helicase YTHDC2 (Fragment) OS=Homo sapiens GN=YTHDC2 PE=4 SV=1
YTHDF3	R4GN55	38	63772	1	1	1	1	0.08	YTH domain-containing family protein 3 OS=Homo sapiens GN=YTHDF3 PE=1 SV=1
YWHAB	P31946	114	27947	7	4	4	4	0.96	Isoform Short of 14-3-3 protein beta/alpha OS=Homo sapiens GN=YWHAB
YWHAE	P62258	161	26658	6	5	5	5	1.41	Isoform SV of 14-3-3 protein epsilon OS=Homo sapiens GN=YWHAE
YWHAG	P61981	82	28456	5	4	4	4	0.94	14-3-3 protein gamma OS=Homo sapiens GN=YWHAG PE=1 SV=2
YWHAZ	E7EX29	233	28190	10	6	7	6	1.71	14-3-3 protein zeta/delta (Fragment) OS=Homo sapiens GN=YWHAZ PE=1 SV=1
ZBTB42	B2RXF5	31	47602	1	1	1	1	0.10	Zinc finger and BTB domain-containing protein 42 OS=Homo sapiens GN=ZBTB42 PE=1 SV=1
ZRANB2	O95218	251	36753	8	7	8	7	1.45	Isoform 2 of Zinc finger Ran-binding domain-containing protein 2 OS=Homo sapiens GN=ZRANB2
ZYX	H0Y2Y8	79	58804	3	2	3	2	0.17	Zyxin (Fragment) OS=Homo sapiens GN=ZYX PE=1 SV=1

1477

1478   * TRAPP interactors are highlighted in bold

1479

1480   **Table S2. Small interfering RNA oligonucleotides used in the study.**

Gene name		siRNA sequence
TRAPPC2	#1	GGGCAUAUGAGGUUUAUUA
	#2	GAGCUUCUACUUUGUAAUU
	#3	ACAUGUGGCUAUCGAACAA
	#4	CAUAAGACAAGAAGAUGGA
TRAPPC3		GGGCAUCACUCCAAGCAUU
Sec23A	#1	GUUAUGCUGGUAUAUCUGA
	#2	GCAUAAUGCUCCAAUUCCU
Sec23B	#1	CACGUUACAUCAACACGGA
	#2	CACUAUGAGAUGCUUGCUA
Sec24A		GAGUCAGUGAGCCAAGGAU
Sec24B		GCCGAUCCUGUCGAACGUAUA
Sec24C		GGCUGCUGUGUAGAUCUCU
Sec24D		CUGUCUUACCCAGGAGGCU
Sec31A		GCUUCUCUUCCACUCAAUA
Sec16A		GGGCGCAAAGUGAGCUGCCAGAUUU
Sec16B		CCCUGCCUAGUUUCCAGGUGUUUAA
CDK1	#1	GAACUUCGUCAUCCAAAUA
	#2	GGUUAUAUCUCAUCUUUGA
	#3	GAAUCUUUACAGGACUAUA
CDK2	#1	CGGAGCUUGUUAUCGCAAA
	#2	GAGUCCCUGUUCGUACUUA
	#3	CAAGAUCUCAAGAAAUUCA

1481

1482   **Table S3. Primers for real time PCR used in this study**

Gene name		Primer sequence
Sec23A	(+)	CCAGATTTCCAATGCCAAGATAC
Sec23A	(-)	TGAGTCTGTGAAGGGTTGAC
Sec23B	(+)	GAATGTGAAAGGACCGTGTG
Sec23B	(-)	AGATGCCAAGTGTAGATGTAGG
Sec24A	(+)	GTTTATGGAGAACCTCACAGAAG
Sec24A	(-)	ACAAAGAGATACACTGGAGGC
Sec24B	(+)	GGGCTCCTGTACCTAACTTG
Sec24B	(-)	AGTAAAGCCTGTGTCTGTGG
Sec24C	(+)	CATCAAACCGCTGGCAAG
Sec24C	(-)	GGGACACATGTATGCTTTGC
Sec24D	(+)	GAGGACAAGTTTATGCCACC
Sec24D	(-)	TGTACAACGGATGAATCGAGG
Sec16A	(+)	AACCAACGAAGCAATCCAG
Sec16A	(-)	GCGGCAGGAGTAGATGAAC
Sec16B	(+)	GGTTTTAGAAAGACGCAGTGG
Sec16B	(-)	TGAGCGAGTAGGATGAGGAT
HPRT	(+)	TGCTGACCTGCTGGATTACA
HPRT	(-)	CCTGACCAAGGAAAGCAAAG
CDK1	(+)	AAGGAACAATTAAACTGGCTGAT
CDK1	(-)	TGACCCCAGCAATACTTCTG
CDK2	(+)	GCATTCCTCTTCCCCTCATC
CDK2	(-)	AGGTTTAAGGTCTCGGTGG

1483