# Spontaneously Arising *Streptococcus mutans* Variants With Reduced Susceptibility to Chlorhexidine Display Genetic Defects and Diminished Fitness

Justin R. Kaspar[1][#], Matthew J. Godwin[1], Irina M. Velsko[2,3], Vincent P. Richards[2], and Robert A. Burne[1]

[1]Department of Oral Biology, University of Florida, Gainesville, Florida
[2]Department of Biological Sciences, Clemson University, Clemson, South Carolina
[3]Max Planck Institute for the Science of Human History, Jena, Germany

**Running title:** characterization of chlorhexidine reduced suspectability variants in *Streptococcus mutans*
**Keywords:** dental caries, antimicrobial tolerance, chlorhexidine, biofilms, stress response, adaptation, microbial antagonism

[#] Corresponding author
Mailing address: Department of Oral Biology, University of Florida, College of Dentistry, P.O. Box 100424, Gainesville, FL 32610.
Phone: (352) 273-8848
E-mail: jkaspar@dental.ufl.edu

## ABSTRACT

Chlorhexidine (CHX) has been used to control dental caries, caused by acid-tolerant bacteria such as *Streptococcus mutans*, since the 1970s. Repeated CHX exposures in other bacterial species results in development of variants with reduced susceptibility that also become more resistant to other antimicrobials. It has not been tested if such variants arise when streptococci are exposed to CHX. Here, we passage *S. mutans* in increasing concentrations of CHX and isolate spontaneously-arising reduced susceptibility variants (RSVs) from separate lineages that have minimal inhibitory concentrations (MICs) that are up to 3-fold greater than the parental strain. The RSVs have increased growth rates at neutral pH and under acidic conditions in the presence of CHX, but accumulate less biomass in biofilms. RSVs display higher MICs for daptomycin and clindamycin, but increased sensitivity to dental-relevant antimicrobials triclosan and sodium fluoride. Plate-based assays for competition with health-associated oral streptococci reveal decreased bacteriocin production by the RSVs, increased sensitivity to hydrogen peroxide, and diminished competitive fitness in a human-derived *ex vivo* biofilm consortium. Whole genome sequencing identified common single nucleotide polymorphisms (SNPs) within a diacylglycerol kinase homolog and a glycolipid synthesis enzyme, which could alter accumulation of lipoteichoic acids and other envelope constituents; as well as a variety of mutations in other genes. Collectively, these findings confirm that *S. mutans* and likely other streptococci can develop tolerance to CHX, but that increased tolerance comes at a fitness cost, such that CHX-induced variants that spontaneously arise in the human oral cavity may not persist.

**INTRODUCTION**

Controlling microbial biofilm infections in humans is challenging in part due to the presence of extracellular polymeric substances (EPS) in which the organisms are embedded, the presence of persister cells as well as subpopulations of cells with low metabolic activity, and exchange of antimicrobial resistance genes through horizontal gene transfer. In the human oral cavity, accumulation of biofilms on the teeth can lead to destruction of tooth mineral when there is frequent and significant localized acidification by organic acids produced as metabolic end products from fermentable carbohydrates (1). The initiation and progression of dental caries can be controlled to a degree through non-specific mechanical removal of the biofilms or by treatment with various chemical agents that include fluoride, xylitol or chlorhexidine (CHX) (2, 3). While the effectiveness of CHX in caries prevention is not definitively established and there can be undesirable side effects (e.g. staining, irritation of host tissues) (4), CHX remains a standard against which antimicrobial therapies are compared for their efficacy in the suppression of caries-causing bacteria, such as *Streptococcus mutans* (5–7). Currently in the United States, mouth rinses containing 0.12% CHX are the only CHX-containing products marketed for caries prevention (4).

CHX is a symmetric bis-biguanide agent comprising two chloroguanide chains that are connected by a central hexamethylene chain and has diverse medical applications as a disinfectant for surfaces and as an antiseptic for topical applications. CHX carries two positive charges at physiological pH and is attracted to the bacterial cell surface, where it may electrostatically interact with negatively charged phospholipids (8). Depending on the concentration of CHX, it can reduce bacterial membrane fluidity or disrupt the structural integrity of the membrane(s), causing increased permeability, leakage of cell contents, loss of proton motor force and cell death as its main mode of action (9). In one study (10), it was found that CHX induces the formation of dented spots on the surface of the cell wall of both *Escherichia coli* and *Bacillus subtilis*. The authors posited that the structural changes of the bacterial cell wall may be due to the action of CHX with membrane cardiolipin and phosphatidylethanolamine to disturb the normal arrangement and integrity of the phospholipid bilayer structure and its associated proteins (10). A similar finding was recently made in *S. mutans*, where vesicle-like structures were apparent on the cell surface after treatment with 0.2% CHX (11). In the same study, treatment of 72-hour *S. mutans* biofilms resulted in 80% of the population displaying loss of membrane integrity, assessed by staining with propidium iodide (PI) and analysis of the population using flow cytometry (11).

While CHX likely puts substantial selective pressure on *S. mutans* and the oral microbiota in general, particularly if used repeatedly for oral care, the organisms appear to remain susceptible to CHX, with few reports of resistance (12). However one recent study (6) reported that when CHX was continuously flowed on biofilms formed on polystyrene blocks *in vitro*, detached cells displayed a higher minimal inhibitory concentration (MIC) to CHX (1.25 $\mu$g mL$^{-1}$), whereas most host-derived isolates of *S. mutans* evaluated had MICs < 1$\mu$g mL$^1$ (6). Earlier studies with other oral streptococci, including *Streptococcus sanguinis* and *Streptococcus mitis,* showed that repeated exposure to CHX resulted in increased MICs to CHX, but these findings were not deemed to be of clinical significance (13–17). In enterococci, increased tolerance to antiseptics like CHX in hospital settings has inspired research into mechanisms for increased CHX tolerance, including studies of CHX effects on the transcriptome (18) and analysis of the impact of CHX on a transposon mutant library (19). More specifically, Bhardwaj and colleagues (18) used RNA-Seq to show that VanA-type vancomycin resistance genes are upregulated in *Enterococcus faecium* following exposure to CHX. In *E. faecium*, two genes encoding for a two-component regulatory system were found to contribute to CHX tolerance (19). Thus, it is clear that Firmicutes have the capacity to develop increased tolerance

to CHX, but the mechanisms by which tolerance may develop, especially in streptococci, is only
beginning to be explored.
There is a continued need to be alert to the emergence of new oral isolates that display
reduced susceptibility to CHX due to its increased usage within the clinic (20). Adaptations to
CHX may also lead to cross-resistance to other antimicrobials, as was noted recently with
clinical strains of *Klebsiella pneumoniae* that could acquire resistance to the last-resort antibiotic
colistin after exposure to CHX (21). To date, the emergence of tolerant or resitant oral bacteria
due to repeated use of CHX in the dental practice has not been fully studied, although
determination of whether exposure to CHX can result in acquisition of resistance to other
antimicrobials and the molecular mechanisms by which CHX may confer resistance in these
organisms is of growing interest (22). In this study, we focus on the caries-associated pathogen
*S. mutans*, show that passage on increasing concentrations of CHX gives rise to variants that
display reduced susceptibility to CHX, and begin to examine the genetic basis for reduced CHX
susceptibility. This research serves as a starting point to investigate the mechanisms of action
of CHX and how streptococci cope with this chemical stressor that will allow us to generate
information that supports the development of more effective treatments for removal of
cariogenic bacteria from oral biofilm communities.
110
111   **RESULTS AND DISCUSSION**
112
*Selection for Chlorhexidine Reduced Susceptibility Variants.* To begin our study, we initially
screened nine strains of *S. mutans* that were acquired from different sources and that displayed
phenotypic heterogeneity in established virulence-related traits (23, 24). The MICs for the nine
strains ranged from 0.25 – 0.625 $\mu$g mL$^{-1}$ CHX **(Supplemental Figure 1)**, and none greatly
exceeded the previously reported 0.5 $\mu$g mL$^{-1}$ MIC for *S. mutans* (12). For the continuation of
the research, we chose to work with strain UA159, which was orginally isolated in 1982 from a
child with active caries and serves as the wild-type strain for many laboratories that study *S.*
*mutans*. We next wanted to determine how *S. mutans* may adapt to repeated CHX exposures
over time, and whether isolates that display reduced susceptibility may arise. To accomplish
this, UA159 was passaged on BHI agar plates containing various concentrations of CHX
**(Figure 1A)**. Colonies able to grow on the highest concentration CHX agar plate were picked,
regrown in BHI broth under non-selective conditions, and then plated on a higher concentration
of CHX. The selections were then repeated with one additional round using higher
concentrations of CHX and the colonies were stored for further analysis.
To acquire the first generation of spontaneously-arising CHX $\underline{r}$educed $\underline{s}$usceptibility *S.*
*mutans* $\underline{v}$ariants (RSVs), 1x10$^6$ colonly forming units (CFUs) of *S. mutans* UA159 grown in BHI
broth to OD$_{600\ nm}$ = 0.5 (mid-exponential log phase) was spread on BHI agar plates containing
0.1, 0.25 or 0.5 $\mu$g mL$^{-1}$ CHX. While the plates containing 0.1 or 0.25 $\mu$g mL$^{-1}$ CHX yielded lawns
of bacteria, individual colonies were obtained on the 0.5 $\mu$g mL$^{-1}$ agar plates. No colonies could
be obtained in the first round of plating on BHI agar plates supplemented with 0.75 $\mu$g mL$^{-1}$
CHX. Ten individual bacterial colonies were selected from the 0.5 $\mu$g mL$^{-1}$ plate and designated
as RSV1s (reduced susceptibility variants from the 1$^{st}$ generation). These colonies were
regrown to OD$_{600\ nm}$ = 0.5 in broth, diluted to 1x10$^6$ CFUs and 0.1 mL of each of the 10 RSV1
cultures were spread on individual BHI agar plates containing either 0.75, 1.0 or 1.25 $\mu$g mL$^{-1}$
CHX, establishing 10 independent lineages moving forward to the second generation (RSV2s).
Individual colonies were obtained on plates with all three concentrations of CHX. One colony
was randomly selected from each of the 1.25 $\mu$g mL$^{-1}$ CHX plates to move forward to the third
generation (RSV3s). After growth to OD$_{600\ nm}$ = 0.5 and dilution to 1x10$^6$ CFUs, these 10 RSV2s
were then plated on 1.5, 1.625 or 1.75 $\mu$g mL$^{-1}$ CHX. Again, colonies were obtained on all
concentrations. At this point, one randomly-selected colony from each of the RSV3 plates (1.75

µg mL$^{-1}$ CHX) was stored for further analysis. Confirmation that the variants were indeed *S.*
*mutans* was obtained by amplifying and sequencing a 1.5 kbp region from the 16S rDNA
sequence. Seven selected RSV3s that returned as *S. mutans* were subjected to further
analysis.
To confirm that the selected RSV3s had reduced susceptibility to CHX, an MIC test was
conducted **(Figure 1B)**. While the parental strain could only grow in BHI broth containing 0.5 µg
mL$^{-1}$ CHX after 24 h, all RSV3s were able to grow in BHI containing >0.5 µg mL$^{-1}$ CHX.
Interestingly, all seven RSV3s tested displayed variable MICs, with the most tolerant isolate,
RSV3-6, able to grow in BHI broth containing 1.50 µg mL$^{-1}$ CHX, a 3-fold increase from the
parental strain. Other isolates, such as RSV3-1 and RSV3-8 were only able to grow in BHI broth
containing 0.75 µg mL$^{-1}$ CHX after 24 h, a half-fold increase from parental. These data indicated
to us that the seven selected RSV3s are diverse in that they were able to propagate in different
concentrations of CHX, possibly indicative of the acquisition of different mutations within the
genome.
157

158
**Figure 1. Acquisition of Chlorhexidine Reduced Susceptability Variants.** **(A)** Schematic of methods used to obtain the reduced
susceptibility variants to chlorhexidine (RSVs). ~1x10$^6$ CFU of *S. mutans* UA159 grown to mid-exponential phase was spread plated
on BHI agar plates containing either 0.1, 0.25 or 0.5 µg mL$^{-1}$ CHX. Ten different single colonies were selected from the 0.5 µg mL$^{-1}$
CHX plate to establish 10 independent lineages (RSV1). Cultures of these colonies were then plated on separate 0.75, 1.0 or 1.25
µg mL$^{-1}$ CHX BHI agar plates (RSV2). One randomly selected colony from each plate was then cultured in BHI and plated on
separate 1.5, 1.625 or 1.75 µg mL$^{-1}$ CHX. At this point, one randomly selected colony from each of the RSV3 plates was stored for
further analysis and characterization. **(B)** MIC determination of the parental along with the RSV3s grown for 24 h in BHI with the
designated concentrations of CHX. Red dots denote highest CHX concentration that the strains displayed growth (OD$_{600\,nm}$ > 0.1).
Four biological replicates of each isolate were tested during the experiment and was replicated four times.

168
*Attributes of the RSV3s.* We reasoned that potential mutations could have pleiotropic effects, so
several key properties of the variants were examined. We found that a majority of the RSV3s,
except RSV3-1 and RSV3-9, had a shorter lag phase compared to the parental UA159 **(Figure**
**2A)**. However, all RSV3s displayed longer doubling times than the parental strain, ranging from
56 ± 2 min for RSV3-3 to 145 ± 26 min for RSV3-1, in contrast to 51 ± 2 for the parental **(Table**
**1)**. When growth rates in the presence of 0.75 µg mL$^{-1}$ CHX were determined, the parental
strain displayed little growth over a 24 h period (doubling time 623 ± 69 min, final yield 0.29 ±
0.09 OD$_{600\,nm}$), whereas all isolates displayed average doubling times under 150 min and final
yields higher than 0.45 **(Figure 2B)**. Some variants, including RSV3-3, RSV3-8 and RSV 3-10,
showed little difference in doubling times in the presence or absence of 0.75 µg mL$^{-1}$ CHX.
These data confirm that the selected isolates have reduced susceptibility to CHX, and provide
support for the idea that the disparate growth characteristics may reflect variation in the
underlying genetic and physiologic bases for increased MICs to CHX.

One critical virulence attribute of *S. mutans* is acid tolerance (aciduricity), which allows
organisms to continue metabolic processes at lower pH values than acid-sensitive oral
organisms, leading to ecological shifts that promote caries development (25–27). Additionally,
when *S. mutans* is cultured at acidic pH values, the fatty acid profile of the membrane is
modified from mainly short-chained saturated fatty acids to a membrane with increased
proportions of mono-unsaturated fatty acids with longer chains (28). Interestingly, all isolates
had a shorter lag time compared to the parental when diluted from BHI at neutral pH into BHI
that had been acidified with HCl to a pH of 5.5 **(Figure 2C)**. However, the doubling time of the
RSVs was comparable to that of the parental, with the parental obtaining the highest final yield.
When 0.75 µg mL$^{-1}$ CHX was included in the acidified broth, the parental strain failed to grow
after 48 h, while all variants displayed growth **(Figure 2D)**. Some variants, including RSV3-3
and RSV3-7, actually displayed shorter doubling times and higher final yields when grown with
CHX, compared with no CHX present. One possible explanation is that the variants have
changes in the the phospholipid profile that allows them to be more toleratant of, and/or more
rapidly adapt to, the low pH challenge.
The ability of the RSVs to compete with the parental strain and to form biofilms was also
assessed. For competitive index measurements, a marked parental that included integration of
an antibiotic resistance marker from pMC340B, which allow for single copy integration of a
kanamycin resistance marker in the *mtlA – phnA* locus (29), was competed against unmarked
parental and the RSV3s **(Figure 2E)**. In planktonic cultures, RSV3-3, RSV3-6 and RSV3-10
were significantly more competitive against the parental strain after 24 h. These differences are
consistent with the varying growth rates displayed by the parental and variants. In terms of
biofilm formation in the presence of sucrose, all RSV3s other than RSV3-6 and RSV3-9 had
significantly less biomass accumulation compared to the parental after 24 h of establishment
and growth **(Figure 2F)**. After 48 h, variants RSV3-6 and RSV3-9 showed stable biomass
accumulation that now surpassed the parental strain. Interestingly, RSV3-7, RSV3-8 and RSV3-
10 displayed a 3-fold decrease in biomass compared to parental strain at the 48 h time point.
The mechanisms by which the acquired mutations and/or changes in gene expression impact
the biofilm phenotype of the RSV3s will need to be further explored, but the differences
observed in biofilm forming capacity futher highlights their diversity.
212
*RSV3s have reduced susceptibility to other antimicrobials.* Two recent studies with *E. faecium*
and Methicillin-Resistant *Staphylococcus aureus* (MRSA) demonstrated reduced antibiotic
susceptibility in strains that are less sensitive to CHX or the biguanide antiseptic polyhexanide,
respectively (30, 31). To determine if reduced susceptibility to CHX could lead to cross-
resistance against other antimicrobials in our variants, we determined MICs of the RSV3s for six
different antibiotics (amoxicillin, azithromycin, bacitracin, clindamycin, daptomycin, vancomycin)
that utilize different modes of action using the Etest Strip Assay **(Table 2)**. We saw no change in
susceptibility to antibiotics that inhibit peptidoglycan cross-linking (amoxicillin, vancomycin), but
changes against cell-wall targeting bacitracin and daptomycin were evident **(Supplemental**
**Figure 2)**. All variants, except for RSV3-9, displayed increased susceptibility to bacitracin, and
all variants except for RSV3-1 showed higher MICs to daptomycin (2-6 fold increase). The latter
result is consistent with the fact that CHX and daptomycin disrupt cell membrane potential and
fluidity (30, 31). Interestingly, the RSV3s also showed reduced susceptibility to clindamycin,
which inhibits protein synthesis through binding to the 50S ribosomal subunit (32), perhaps
associated with decreased permeability or increased export of the drug in the RSV3s.
228

**Figure 2. Growth attributes of the RSV3s.** Growth of the parental UA159 strain (black line, squares) and each of the RSV3s (colored lines, circles) in either **(A)** BHI, **(B)** BHI with 0.75 µg mL$^{-1}$ CHX, **(C)** BHI pH 5.5, and **(D)** BHI pH 5.5 with 0.75 µg mL$^{-1}$ CHX. **(E)** Competitive index assay between the parental strain with the RSV3s in BHI medium after 24 h, competed with a marked UA159 that included integration of plasmid pMC340B (Km$^R$) that was used for selection. Data plotted is in the competitive index (Log10) of the unmarked parental or the RSV3s. * = P value $\leq$ 0.1 by Student's T-Test compared to the parental. **(F)** Accumulated biofilm biomass for all individual strains grown for either 24 (black dots) or 48 hours (red dots) in CDM medium supplemented with 20 mM glucose and 5 mM sucrose. 12 individual biofilms from three biological replicates were measured. The experiment was replicated twice.

We next wanted to examine the susceptibility of the RSV3s to two compounds commonly used to prevent oral infectious diseases: triclosan and sodium fluoride. Triclosan is a broad-spectrum antibacterial, but unlike CHX, triclosan is more compatible with other ingredients commonly used in toothpastes. Triclosan also does not stain teeth or have an unpleasant taste, as does CHX, although CHX appears to have more potent antibacterial properties (33). Triclosan was also part of a recent ban by the FDA on 19 biocidal chemicals that were deemed nonbeneficial to household wash products (34). No observable growth differences were seen when the variants were inoculated with 8 µg mL$^{-1}$ triclosan **(Figure 3A)**, but when grown with 16 µg mL$^{-1}$ triclosan all RSV3s grew more poorly than strain UA159 **(Figure 3B)**. Only variants RSV3-3, RSV3-8 and RSV3-10 displayed final yields similar to the parental -- all others did not reach as high a final optical density. Fluoride is widely used in toothpastes and mouthwashes, and is typically added to municipal drinking water. Fluoride can

not only arrest the demineralization of the tooth enamel, but at high concentrations (13 – 647
mM) it inhibits growth and metabolism of *S. mutans* (35, 36). Growth of the variants compared
to the parental was moderately impaired with addition of 100 μg mL$^{-1}$ sodium fluoride **(Figure
3C)**, but substantial growth deficiencies were observed with 300 μg mL$^{-1}$ sodium fluoride.
Collectively, these observations show that the RSV3s may display higher MICs to certain
antibiotics, but they also have increased sensitivity to agents that are commonly used in oral
health formulations.
259

260
261 **Figure 3. Changes in susceptibility to clinically-relevant antimicrobials.** Growth of the parental UA159 strain (black line;
262 squares) and each of the RSV3s (colored lines; circles) in either **(A)** 8 μg mL$^{-1}$ triclosan, **(B)** 16 μg mL$^{-1}$ triclosan, **(C)** 100 μg mL$^{-1}$
263 sodium fluoride and **(D)** 300 μg mL$^{-1}$ sodium fluoride. Note differences in time scale for x-axis.
264

<u>*RSV3s are Less Competitive Against Commensal Oral Streptococci.*</u> As noted above (6), there
is potential for spontaneously-arising variants that show decreased susceptibility to CHX that
are selected within *S. mutans* populations *in vitro*, but such isolates have not been recovered
from dental plaque. One possibility is that there are selective pressures imposed by host factors
and the oral microbiome that keep spontaneous CHX reduced susceptibility variants of *S.*
*mutans* from taking hold in the human oral cavity.
To explore this idea, we investigated the antagonistic interactions of our RSV3s with
species of commensal streptococci that are generally abundant and associated with dental
health in humans. We measured bacteriocin production by our variants using a deferred
antagonism assay in which the bacteria are stabbed on agar plates, grown overnight, and then
the plates are overlaid with soft agar containing 1x10$^7$ cells of a sensitive indicator strain; in this
case either *S. sanguinis* SK150 or *Streptococcus gordonii* DL1. Bacteriocin production was
assessed by measuring the area of growth inhibition shown by the indicator strain. When
overlaid with *S. sanguinis* SK150, we observed 1.2- to 3-fold decreases in the area of growth
inhibition, indicating less bacteriocin production **(Figures 4A and 4C)**. A more significant
difference was observed when *S. gordonii* DL1 was used as the indicator strain, as all isolates
displayed ~ 3 fold decrease compared to parental **(Figure 4B and 4D)**. We also performed a
282

**Figure 4. Competition of RSV3s against oral streptococci.** Agar plate images of bacteriocin overlay assays displaying areas of growth inhibition against indicator strains **(A)** *S. sanguinis* SK150 and **(B)** *S. gordonii* DL1. Quantification of bacteriocin overlay assays against indicator strains **(C)** *S. sanguinis* SK150 and **(D)** *S. gordonii* DL1. Quantification was conducted using the ImageJ measuring tool and represents the average of three biological replicates from two independent experiments. All RSV3 areas of growth inhibition were statistically significant from parental (P value ≤ 0.003). Growth of the parental strain (black line; squares) and each of the RSV3s (colored lines; circles) in either **(E)** 0.001% $H_2O_2$ or **(F)** 0.003% $H_2O_2$. Outgrowth of parental and each variant in individual *ex vivo* biofilms co-inoculated with cell-containing saliva (CCS) after **(G)** 24 h and **(H)** 48 h. Outgrowth was calculated by subtracting the initial inoculation CFUs at the beginning of the experiment (t = 0 h) from the CFUs enumerated at the time point indicated. Biofilms from three biological replicates of each strain were grown. CFUs for all RSV3s were statistically significant from parental (P value ≤ 0.03 and 0.02). Each time point was replicated twice.

plate-based competition assay where *S. sanguinis* SK150 and our RSV3s were spotted next to each other at the same time and grown for 48 h **(Supplemental Figure 3)**. Whereas no zone of inhibition was seen in the parental when *S. sanguinis* SK150 was grown adjacent, a zone of inhibition was observed in all but two of the isolates, RSV3-1 and RSV3-8, indicating that the RSV3s did not compete as well as the parental under this condition. One source of antagonism by commensal streptococci against *S. mutans* that could account of the antagonism is hydrogen

peroxide production (37, 38). To determine if the variants had marked differences in their
sensitivity to $H_2O_2$ that would impact their fitness in such assays, the strains were cultured in
BHI supplemented with either 0.001% or 0.003% $H_2O_2$. While differences were not observed in
growth when 0.001% $H_2O_2$ was used **(Figure 4E)**, all variants displayed decreased growth
rates, compared to parental, when challenged with 0.003% $H_2O_2$ **(Figure 4F)**. These data
support that the spontaneous mutations that confer reduced susceptibility to CHX come at a
competitive fitness cost for the variants in the form of decreased bacteriocin production and
increased $H_2O_2$ susceptibility. These changes may account, at least in part, for the absence of
CHX resistance among *S. mutans* isolates from human dental plaque samples. Other factor(s)
not examined here may also contribute to decreased antagonism by the RS3s.
To further test the hypothesis that the RSV3s had diminished competitive fitness
compared to the parental strain, we next examined their ability to establish and persist in an *ex*
*vivo*, multi-species biofilm consortium. To complete this experiment, we chose to propagate the
parental and each variant seperatly in independent biofilms that were co-inoculated with cell-
containing saliva (CCS), collected and pooled from four healthy donors. The CCS model has
been previously shown to produce a highly diverse microbial community that is predominantly
composed of *Streptococcus* spp. (39–41). CFUs of each strain were determined at the time of
initial inoculation (t = 0 h) and after either 24 or 48 h of incubation (t = 24 or 48 h) by plating
serially-diluted cells on BHI (total cells) or mitis salivarius-bacitracin (MSB) agar supplemented
with 1% tellurite and 20% sucrose (42) **(Supplemental Figures 4 and 5)**. All variants, except
for RSV3-1, grew in this multi-species biofilm. We measured the outgrowth of all *S. mutans*
strains, calculated by subtracting the initial inoculated CFUs at t = 0 h from the final CFUs. At t =
24 h, the RSV3 variants displayed a 2- to 7-fold reduction in outgrowth compared to the parental
(mean = 4-fold) **(Figure 4G)**. The variants did not persist as well as the parental strain in the
CCS biofilms at 48 h, with 12- to 94-fold lower CFUs measured (average of 55-fold) **(Figure**
**4H).** Total cells were also enumerated from the biofilms by plating on non-selective BHI agar
and the cell count did not significantly differ between any of the samples (data not shown*).* The
CCS assay therefore provided additional evidence that the RSV3s are impaired for growth in
conditions that more closely mimic the complex communities found within the oral cavity, when
compared to the parental strain. The simplest interpretation of these results is that RSV3s
produce less bacteriocins, decreasing their antagonistic capability towards other streptococci
present within the CCS during times of establishment, and that their ability to tolerate hydrogen
peroxide over time and possibly other antagonistic factors produced by oral bacteria (37, 43) is
concurrently diminished, thereby lessening their ability to persist in this model.
335
*Whole Genome Sequencing Reveals Mutations in RSVs.* To determine what mutations are
carried in each of the characterized RSV3s, whole genome sequencing was performed on
extracted chromosomal DNA as described in the material and methods section. Our analysis for
single nucleotide polymorphisms (SNPs) revealed four genes of interest that carried mutations
in at least three or more of the sequenced RSV3s. The presence of the SNPs were confirmed
by PCR and sequencing in SMU.804, SMU.1542c, SMU.1576c and SMU.1588c.
The only gene within the genome to carry SNPs in all seven of the sequenced RSV3s
was SMU.1542c **(Table 3)**. SMU.1542c is homologous to *dgkB* (*yerQ*) of *Bacillus subtilis*, a
soluble diacylglycerol kinase (DGK) that phosphorylates diacylglycerol (DAG) generated by the
transfer of the *sn*-1-glycerol-P headgroup from phosphatidylglycerol to lipoteichoic acid (LTA)
(44, 45). Intriguingly, DgkB from *B. subtilis* is essential for growth, yet four different non-
synonymous SNPs were identified between our seven sequenced RSV3s that occurred over the
length of the gene and were not confined to one particular region. It is not known whether the
lethality of the loss of *dgkB* in *B. subtilis* is due to accumulation of toxic DAG in the membranes
or from the absence of LTA (45). We suspected that the resulting non-synonymous SNPs in the
apparent *S. mutans dgkB* homologue in RSV3s may result in changes to the efficiency of the

enzyme, rather than complete loss of function as a recent genome-wide screen in *S. mutans* by
Tn-Seq found SMU.1542c to be essential (46). To test this hypothesis, we performed a western
blot on the extracted LTA from the parental strain and the RSV3s **(Figure 5A)**. Differential
accumulation of LTA in the variant strains were noted. RSV3-1 and RSV3-10 appeared to
accumulate more LTA and RSV3-6, RSV3-7, RSV3-8 and RSV3-9 accumulated less LTA than
the parental strain. Thus, it does appear that our variants show different patterns in the amount
of LTA produced that may be related to the SNPs. Further analysis will be required to determine
if the LTA and lipid profiles are altered in the RSV3s, as has been reported for *E. faecium* (30).

We analyzed the SNPs occurring within *dgkB* of our variants to determine how they
might affect DgkB function within *S. mutans* by comparing to characterized Dgk enzymes,
including the lipid kinase YegS from *Escherichia coli*, DgkB of *S. aureus* to which the structure
has been solved (44), as well as the ten known isoforms of human DGKs **(Supplemental
Figure 6)**. We noted that the amino acid (aa) substitution G131S occurring within RSV3-6,
RSV3-7 and RSV3-9 arose within beta sheet 6 and was a naturally occurring substitution found
within human cerimide kinase CERK, a member of the DGK superfamily (Pfam00781). The
mutation causing G164D, found in RSV3-1, was a substitution occurring within DGKζ, a human
type-2 DGK. Substitution D249E found in RSV3-3 occurred at the end of alpha helix 7 and is
present in YegS of *E. coli*. Finally, aa substitution A304D of RSV3-8 and RSV3-10 occurred
within a conserved residue of the DGKs (44) after beta sheet 17, with no known symmetry
between the DGKs examined. None of the aa substitutions were within regions of the protein
that impacted nucleotide (ATP/ADP) -binding regions nor along the DgkB dimerization interface.
Therefore, we are unable to postulate how the identified SNPs of the RSV3s within SMU.1542c
may alter the function of DgkB. However, since a majority of the SNPs lead to amino acid
substitutions found in other DgkB homologs, combined with the fact that these SNPs do not lead
to abberant growth defects in their respective variant as demonstrated using multiple growth
tests, we assume that the DgkB enzyme in the RSV3s is still functional.

SMU.1588c is annotated to be a glycolipid synthesis enzyme, specifically a 1,2-
diacylglycerol 3-glucosyltransferase that synthesizes glucose($\alpha$1-3)-DAG from uridine
diphosphate glucose (UDP-glucose) and termed *lafA* in *Listeria monocytogenes* (47). It is
encoded within an operon with SMU.1589c that shares homology to *lafB* of *L. monocytogenes*,
responsible for the addition of a galactose moiety to form galctosyl-($\alpha$1-2)-glucosyl-($\alpha$1-3)-DAG
using UDP-galactose as a substrate. In *Streptococcus agalactiae*, the LTA anchor is $\alpha$1-2 linked
$Glu_2$-DAG and is produced by a two-enzyme system similar to L. *monocytogenes* (48–50). We
encountered two different non-synonymous SNPs within the *lafA* homolog SMU.1588c: 1) aa
substitution P433A in RSV3-6 and RSV3-8, and 2) W391STOP in RSV3-10, which leads to
truncation of the protein at aa 391. Interestingly, the protein sequence of the well characterized
*Acholeplasma laidlawii* 1,2-diacylglycerol 3-glucosyltransferase also ends at position 391 when
the protein sequences are aligned **(Figure 5B)**. It is unknown what function the additional C-
terminal amino acids of the *S. mutans* 1,2-diacylglycerol 3-glucosyltransferase might serve, but
the additional domain(s) is present in *Streptococcus*, *Staphylococcu*s, *Enterococcus* and
*Listeria,* but not in *Bacillus*.

Two other genes, SMU.804 and SMU.1576c, were found to have SNPs in three of the
RSV3s. RSV3-7, RSV3-8 and RSV3-10 harbored SNPs in SMU.804, while RSV3-7, RSV3-8
and RSV3-9 harbored SNPs in SMU.1576c. Both genes are listed as hypothetical proteins with
no known function. To determine if these two genes may be involved in a CHX reduced
susceptibility phenotype that is observed in our variants, we individually replaced each gene
with a non-polar kanamycin antibiotic marker to produce strains Δ804 and Δ1576c. The mutant
strains did not display any changes in growth compared to the parental strain when grown in
0.50 µg mL $^{-1}$ or 0.75 µg mL $^{-1}$ CHX, nor did the loss of either gene impact the strain's MIC to
daptomycin or change antagonistic interactions with commensal streptococci as assessed with

**Figure 5. RSV3s SNPs associate with changes in LTA.** **(A)** Western blot of parental and the RSV3s using a 1:5000 dilution of Gram-positive bacterial LTA monoclonal antibody. Cells were grown in BHI medium to late exponential log phase ($OD_{600\ nm}$ ~ 0.8) prior to harvest. LTA was extracted by adding 0.5 mL of 2x sample buffer to 0.1 g of cell pellet and boiling for 1 hr. 10 μL was then added to a 4-20% TGX pre-cast protein gel. Molecular mass standards (in kDa) are shown to the left. **(B)** Amino acid (aa) alignment (Clustal Omega) between the *lafA* gene (SMU.1588c) of *A. laidlawii, S. mutans*, and *S. pneumoniae* (*D39*, spd_0961). Red bolded aa within the *S. mutans* sequence signify those residues that were found to be mutated within the RSV3s by whole genome sequence analysis, with the corresponding amino acid substitution found above. STOP = stop codon. Sequences were aligned using the web tool Clustal Omega (https://www.ebi.ac.uk/Tools/msa/clustalo/).

the bacteriocin overlay and plate-based competition assays **(Supplemental Figure 7)**.
Together, these data suggest that while SNPs accumulated in both of these genes in three
different variants, they were either not solely responsible for the phenotypes of the variants
and/or the mutations did not cause complete loss of function.
Other intriguing SNPs **(Supplemental Table 1)**, such as those found in the *rcrP* gene of
RSV3-1 (51) or the *rcrRPQ* peptides of RSV3-7 (52) were documented. The *rcrRPQ* operon and
two peptides encoded in *rcrQ* play roles in stress tolerance and genetic competence, but SNPs
in *rcrRPQ* occurred in only one of the variants sequenced. We cannot rule out, however, that
these single SNPs or a combinatorial effect of the multiple SNPs in individual variants are
responsible for the observed phenotypes. However, we have been unable to assign direct
causation of the displayed phenotypes by any particular SNP. A future study designed to assess
global transcriptomic changes in the variants compared to the parental strain may help to
resolve the basis for the reduced susceptibility of growth inhibition by CHX and may provide
valuable insights into mechanisms of action of the compound and ways to enhance its efficacy
in oral biofilm control.
428
429 **SUMMARY**
430
The use of antimicrobials in either mouthwashes or dentifrices in conjunction with
standard mechanical removal of oral biofilms through brushing has been common practice in
the dental setting since the 1970s (53). Still today, use of antimicrobials such as CHX in
varnishes or even incorporation of CHX in nanoparticles is being proposed for treatments to
reduce the appearance of white spot lesions in patients with fixed orthodontic appliances. Even
though CHX has been familiar in the dental setting for decades, we still know very little about its
mode of action against cariogenic bacteria, such as *S. mutans,* and whether or not a tolerant
and/or resistant phenotypes against CHX can arise and the consequences that brings to the
homeostasis of the oral microbiome. The work reported here finally begins to explore these
issues and finds that after only a few rounds of passaging on CHX-containing agar can
spontaneously-arising variants develop with reduced susceptibility to CHX. However, further
investigation and characterization of these variants determines that while these variants do in
fact have the potential to tolerate and persist at low pH values, reduced susceptibility to CHX
comes at a fitness cost to the bacteria in terms of being able to compete against other
commensal streptococci within oral biofilms, as well as increased susceptibility to other common
oral antimicrobials such as triclosan and sodium fluoride.
While broad-spectrum antimicrobials that reduce the overall bacterial load will most likely
always be found within the healthcare setting, the increase in antimicrobial resistance to
common antibiotics and the realization that removal of the natural flora can have unintended
consequences such as allowing pathogens into unoccupied niche(s), necessitate continuing the
search for novel preventive and therapeutic interventions. The unintended finding that reduced
CHX susceptibility through accumulated SNPs can lead to increased vulnerability to triclosan
and sodium fluoride opens the door to future research in determining the mechanisms of action
for designation of a new drug target that complements existing treatments. Together, a focus on
these mechanistically-driven studies in microbial physiology and genetics will strengthen our
knowledge base, provide novel insights into how we can more effectively target cariogenic
bacteria, and translate to better health outcomes in the clinic to increase our quality of life.

464    **MATERIALS and METHODS**
465
*Bacterial Strains and Growth Conditions.*  *S. mutans* wild-type strain UA159 and its derivatives
**(Supplemental Table 2)** were grown in either brain heart infusion (BHI – Difco) or FMC (54)
medium.  The medium was supplemented with 1 mg mL$^{-1}$ of kanamycin when needed. Unless
otherwise noted, cultures were grown overnight in BHI medium with the appropriate antibiotics
at 37°C in a 5% $CO_2$ aerobic atmosphere.  The next day, cultures were harvested by
centrifugation, washed twice in 1 mL of phosphate-buffered saline (PBS), and resuspended in
PBS to remove all traces of BHI. Cells were then diluted in the desired medium before
beginning each experiment. For growth rate comparisons, overnight cultures were diluted 1:50
and grown to mid-exponential phase ($OD_{600\,nm}$ = 0.5–0.6). Then, cultures were re-diluted 1:100
into 0.3 mL of BHI and added to multiwell plates along with any condition to be tested (0.75 µg
ml$^{-1}$ CHX, 8 and 16 µg ml$^{-1}$ Triclosan, 100 and 300 µg ml$^{-1}$ NaF, and 0.001% and 0.003% $H_2O_2$).
Cultures were overlaid with 50 µl of sterile mineral oil to reduce the growth inhibitory effects of
air on growth of *S. mutans*. The optical density at 600 nm ($OD_{600\,nm}$) was monitored at 30 min
intervals for 24 h using a Bioscreen C lab system (Helsinki, Finland) at 37°C with shaking for 10
s before each reading. Three biological replicates measured in technical triplicates were
measured for each experiment and the averaged growth rates of the replicates was used to
calculate doubling times and final yields.
483
*Passaging of S. mutans on BHI Agar Plates Containing CHX.* *S. mutans* wild-type strain UA159
was first grown overnight in BHI medium. The next morning, the overnight cultures were diluted
1:50 and grown to mid-exponential growth phase ($OD_{600\,nm}$ = 0.5–0.6) before being back-diluted
to 1X10$^6$ CFU. The 1X10$^6$ CFU was plated on BHI agar plates containing either 0.1, 0.25 or 0.50
µg mL$^{-1}$ CHX ( > 99.5%, Sigma-Aldrich). The colonies returned on these BHI agar plates was
considered the first generation (RSV1s). 10 randomly selected colonies that appeared on the
0.50 µg mL$^{-1}$ CHX plate were then chosen for overnight inoculation to repeat the process. For
the second generation, 1X10$^6$ CFU of each outgrowth was plated on 10 separate BHI agar
plates containing either 0.75, 1.0 or 1.25 µg mL$^{-1}$ CHX. One single colony from each of the 10
agar plates containing 1.25 µg mL$^{-1}$ CHX was then chosen for overnight inoculation to repeat
the process for a third and final time. 1X10$^6$ CFU of each outgrowth grown to mid-exponential
growth phase was plated on 10 separate BHI agar plates containing either 1.5, 1.625 or 1.75 µg
mL$^{-1}$ CHX. One randomly selected colony from each of the 10 agar plates containing 1.75 µg
mL$^{-1}$ CHX was then selected to be further characterized.
498
*Microtiter Plate Growth Assays.* CHX concentrations to be tested were first added to BHI
medium with addition of DMSO only serving as a negative control. 198 µl of each concentration
was then pipetted across a row of a 96-well microtiter plate. Next, 2 µl of either *S. mutans* wild-
type strain UA159 (parental) or its derivatives to be tested grown to mid-exponential growth
phase ($OD_{600\,nm}$ = 0.5–0.6) in BHI medium was used to inoculate the wells of the plate. The
$OD_{600\,nm}$ of the cultures was measured after 24 h incubation at 37°C in a 5% $CO_2$ aerobic
atmosphere with a Synergy 2 multimode microplate reader (BioTek). The experiment was
performed at least four times.
507
*Competitive Fitness Index Assays.* To determine fitness of the RSV3s, cultures were grown
overnight alongside *S. mutans* wild-type strain UA159 (parental) and UA159 harboring the
chromosomally-integrated vector pMC304B that carries an antibiotic resistant kanamycin
cassette which can be used for selection purposes.  The next day, all strains were grown to mid-
exponential growth phase ($OD_{600\,nm}$ = 0.5–0.6) after a 1:50 dilution from the overnights. A
1:1000 dilution from the mid-exponential growth phase cultures was then used to inoculate 1 mL
of BHI in which either the RSV3s or parental were competed against UA159 harboring

pMC304B. At this time, a small aliquot was serially diluted and plated on either BHI (total cell count) or BHI containing 1 mg mL$^{-1}$ kanamycin (UA159 harboring pMC304B count) to determine initial cell count at the beginning of the experiment. After 24 h of growth at 37°C in a 5% $CO_2$ aerobic atmosphere, cultures were again serially diluted and plated to determine final cell count. All colony forming units (CFUs) were counted after 48 h of agar plate incubation at 37°C in a 5% $CO_2$ aerobic atmosphere. To determine the cell count of either the RSV3s or wild-type UA159, the UA159 harboring pMC304B count was subtracted from the total cell count. The competitive fitness index from the perspective of either the RSV3s or wild-type UA159 was calculated with the following equation: Log10( (RSV3 cell count $_{final}$ / UA159 cell count $_{final}$) / (RSV3 cell count $_{initial}$ / UA159 cell count $_{initial}$). This experiment was repeated twice with three biological replicates for each experiment.

*Measurements of Biofilm Biomass.* Selected strains were grown from overnight cultures to mid-exponential growth phase after a 1:25 dilution in CDM broth at 37ºC in a 5% $CO_2$ atmosphere. Cells were then diluted 1:100 into CDM medium containing 20 mM glucose and 5 mM sucrose as a carbohydrate source. 0.2 mL of this dilution was loaded into 96 well polystyrene microtiter plates and incubated in a 5% $CO_2$ atmosphere at 37ºC for either 24 or 48 h. After, the medium was decanted, and the plates were washed twice with 0.2 mL of sterile water to remove planktonic and loosely bound cells. The adherent bacteria were stained with 0.15 mL of 0.1% crystal violet for 15 min at room temperature. After rinsing twice with 0.2 mL of water, the bound dye was extracted from the stained biofilm using 0.2 mL of ethanol:acetone (8:2) solution. The extracted dye was then diluted 1:20 into water so that the measured values would be within a linear range of the plate reader. Biofilm formation was quantified by measuring the absorbance of the solution at OD $_{575\ nm}$ using a Synergy 2 multimode microplate reader (BioTek). This experiment was repeated twice with 3 independent biological replicates that included 4 technical replicates each.

*ETEST Antibiotic Susceptibility Testing.* Selected strains were grown from overnight cultures to mid-exponential growth phase after a 1:20 dilution in BHI broth at 37ºC in a 5% $CO_2$ atmosphere. 10 µL from the mid-exponential culture was then added to 10 mL soft BHI agar (0.75% agarose) and overlaid on to BHI agar plates. Once solidified, ETEST strips (bioMérieux, Inc., Durham, N.C.) containing either Amoxicillin, Azithromycin, Bacitracin, Clindamycin, Daptomycin or Vancomycin when then placed in the center of the agar with sterile forceps. After 48 h of growth at 37°C in a 5% $CO_2$ aerobic atmosphere, the agar plates were examined, imaged and lowest MIC recorded where no growth was visible. Two independent observers (J.K. and M.G.) interpreted the ETEST MICs. While a specific blinding protocol was not used, neither observer had knowledge of the other's measurements prior to making their own MIC determinations. Discrepancies were resolved by consensus. Each strain was plated three times from three independent biological replicates.

*Bacteriocin Overlay Assays.* Selected strains were grown from overnight cultures to mid-exponential growth phase after a 1:20 dilution in BHI broth at 37ºC in a 5% $CO_2$ atmosphere. Using a toothpick, the strains were stabbed into BHI agar and incubated for 24 h in a 5% CO2 atmosphere at 37°C. The following day, 10 mL soft BHI agar (0.75% agarose) was inoculated with either 10 µL *Streptococcus sanguinis* SK150 or *Streptococcus gordonii* DL1 overnight culture. The soft agar with bacterial indicator strain was overlaid onto the BHI agar plates with stabbed colonies. After 24 h, plates were imaged and the area of zones of clearance were calculated using ImageJ image analysis software. This experiment was repeated twice with three independent biological replicates in each experiment. Statistical analysis was performed using the Student's T-Test.

*Agar Spot Competition Assays.* Selected strains were grown from overnight cultures to mid-
exponential growth phase after a 1:20 dilution in BHI broth at 37ºC in a 5% $CO_2$ atmosphere.
Each strain (7 µL) was then inoculated adjacent to one another on agar plates as follows: *S.*
*sanguinis* SK150 first and to the left, followed by *S. mutans* UA159 (parental) or RSV3 strains
after the SK150 spot had dried. The plates were incubated for 48 h at 37°C, in 5% $CO_2$ prior to
imaging. This experiment was repeated twice with three independent biological replicates in
each experiment.
573
*Outgrowth of Bacterial Strains in ex vivo Cell-Containing Saliva Biofilm Model.* *S. mutans* UA159
(parental) and RSV3 variants from overnight cultures in BHI were diluted 1:20 into CDM medium
containing 20 mM glucose and grown until the $OD_{600\ nm}$ reached 0.5. At this point, all strains
were inoculated individually (n = 3) into CDM supplemented with 20 mM glucose and 5 mM
sucrose using 1:1000 dilutions, along with an aliquot of cell-containing saliva (CCS) that was
diluted 1:50 into CDM immediately after thawing from -80°C storage. Saliva samples for the
CCS were originally collected and pooled from four healthy adult volunteers, all non-smokers
that had not taken antibiotics for at least 3 months prior to collection, as described previously
(IRB201500497 at the University of Florida) (55). 0.35 mL of the co-inoculated medium was
loaded into an ibidi 15 µ-Slide 8-well glass bottom slide and incubated at 37°C in a 5% $CO_2$
aerobic atmosphere for either 24 h or 48 h, while another 0.1 mL was used for serially dilution
and plating on mitis salivarius-bacitracin (MSB) agar supplemented with 1% tellurite, 20%
sucrose and 0.2 units $mL^{-1}$ bacitracin to select for *S. mutans* (42) to determine initial CFU count.
After 24 or 48 h of incubation, biofilms were washed twice with 1x PBS to remove loosely bound
cells and then recovered by scraping off the attached cells with a pipette tip in 0.3 mL of 1x PBS
and moved to a 5 mL glass tube**.** 0.7 mL of 1x PBS (total 1 mL) was added prior to four 30 s
rounds of sonication with resting on ice in between cycles. Cell suspensions were then serially
diluted and plated on MSB agar. CFUs were enumerated after the plates were incubated for 72
h at 37°C in a 5% $CO_2$ aerobic atmosphere. A blank CCS control (no inoculation of *S. mutans*)
was run side-by-side and plated on MSB agar plates at the beginning and end of the
experiments. Between 100 and 1000 total cells were recovered, representing background levels
of endogenous organisms that could grow on the selective agar. These counts were subtracted
from the *S. mutans* cell counts. Outgrowth in CCS was calcuted by subtracting the initial
inoculation CFU count from the final cell count and significance calculated by student's *t*-test
comparing each individual RSV3 variant to the parental. Serial dilutions of all biofilms were also
plated on BHI agar to obtain a reference for total bacteria in the biofilms – the cell count of all
biofilms (~ $1 \times 10^9$ CFUs) did not differ significantly between time points or samples. Three
independent biological replicates of each strain were tested simultaneously and the experiment
was performed twice.
603
*Construction of Bacterial Strains.*  Mutant strains of *S. mutans* **(Supplemental Table 3)** were
created using a PCR ligation mutagenesis approach as previously described (Lau *et al*., 2002).
Briefly, 700 bp fragments both upstream and downstream of the gene of interest were amplified
with designed primers and containing a BamHI restriction site in the region closest to the gene.
After restriction digest, a non-polar kanamycin resistant cassette was ligated within the arms.
This ligated fragment was then transformed into *S. mutans* UA159 with selection on BHI agar
containing 1 mg $mL^{-1}$ kanamycin. Successful replacement was confirmed both by size selection
colony PCR and Sanger sequencing (Eurofins Genomics, Louisville, KY, USA). Restriction and
DNA-modifying enzymes were obtained from New England Biolabs. PCRs were carried out with
100 ng of chromosomal DNA by using *Taq* DNA polymerase, and PCR products were purified
with the QIAquick kit (QIAGEN).
615

*Whole Genome Sequencing and SNP Calling.* For whole genome shotgun sequencing, genomic DNA were isolated from both the parental UA159 and each of the seven RSV3s using Wizard® Genomic DNA Purification Kit (Promega, Madison, WI, USA) with some modifications. The total cell lysate from 10 mL of overnight cultures grown in BHI was then harvested and the supernatant containing the DNA sample was transferred to a fresh tube containing room temperature isopropanol. The supernatant was rotated at room temperature about an hour or until the thread-like strands of DNA formed a visible mass. DNA was purified in nuclease-free water after two washes in 70% ethanol. Total DNA concentration was measured using a NanoDrop™ One Microvolume UV-Vis Spectrophotometer (ThermoFisher Scientific, Waltham, MA, USA) and DNA integrity was determined by 260/280 ratio. 2-5 ng of DNA from each variant was prepared for next-generation whole genome shotgun sequencing using the Illumina Nextera-XT library preparation and indexing kit. Libraries were normalized by hand rather than with the Nextera-XT kit beads, and were pooled at a final concentration of 2 nM and sequenced on an Illumina MiSeq using the Illumina MiSeq v2 kit with paired-end sequencing and 250-bp reads. Reads were de-multiplexed by the Illumina 468 software. The pipeline a5 (58) was used to remove adapters from, quality-trim, and assemble the raw fastq files into contigs, using default settings. The quality of assembled genomes was checked using QUAST (59). The assembled contigs from each respective variant were pairwise aligned against the deposited GenBank *S. mutans* UA159 file (http://www.ncbi.nlm.nih.gov/nuccore/NC_004350.2) using the Mauve multiple genome alignment tool (version 2.4.0) (40) and SNPs unique to that variant from the parental were called through visual inspection.  SNPs in genes SMU.804, SMU.1542c, SMU.1576c and SMU.1588c were confirmed via Sanger sequencing (Eurofins Genomics, Louisville, KY, USA) (Supplemental Table 3). The raw reads from all genomes we sequenced for this study are available to download from the NCBI under BioProject project accession number PRJNA515165 **(Supplemental Table 4)**.

*Western blotting.* Bacterial strains were grown in BHI at 37°C until they reached an $OD_{600\ nm}$ ~ 0.8. Cells were harvested by centrifugation at 4000 g for 10 min, spent media was discarded, and cell pellets were stored overnight at -20°C. The following day, pellets were re-suspended in 0.5 mL 2X SDS sample buffer (100 mM Tris, pH 6.8, 4% SDS, 20% glycerol, 0.004% bromophenol blue, 10% 2-mercaptoethanol) corresponding to 0.1 g of cell pellet weight. The resuspension was then boiled for 1 h. Following, the remaining debris was sedimented by centrifugation and ten µL of each sample was loaded onto a 4-20% precast TGX protein gel (BioRad). Protein samples were separated by SDS-polyacrylamide gel electrophoresis (SDS-PAGE) and then transferred to a polyvinylidene difluoride (PVDF) membrane using a Trans-Blot® Turbo™ Transfer System (BioRad). LTA was detected using a Gram-positive Bacteria LTA monoclonal antibody (1:5000 primary dilution; Thermo Fisher Scientific) and a goat-anti-mouse immunoglobulin G (IgG) antibody (1:5000 dilution; SeraCare Life Sciences, USA). Western blot signals were detected using the SuperSignal™ West Pico Chemiluminescent Substrate kit (Thermo Fisher Scientific) and visualized with a FluorChem 8900 imaging system (Alpha Innotech, USA).

**ACKNOWLEDGEMENTS**

We thank Dr. Robert Shields for assistance with the SNP analysis and for thoughtful discussions related to the manuscript. Research reported in this publication was supported by the National Institute of Dental and Craniofacial Research of the National Institutes of Health under Award Numbers R01 DE023339, R01 DE13239, T90 DE21990 and F32 DE028479.

**REFERENCES**

1.  **Pitts NB**, **Zero DT**, **Marsh PD**, **Ekstrand K**, **Weintraub JA**, **Ramos-Gomez F**, **Tagami J**, **Twetman S**, **Tsakos G**, **Ismail A**. 2017. Dental caries. Nat Rev Dis Prim **3**:17030.

2.  **Wang Y**, **Li J**, **Sun W**, **Li H**, **Cannon RD**, **Mei L**. 2017. Effect of non-fluoride agents on the prevention of dental caries in primary dentition: A systematic review. PLoS One **12**:e0182221.

3.  **Duangthip D**, **Jiang M**, **Chu CH**, **Lo EC**. 2015. Non-surgical treatment of dentin caries in preschool children – systematic review. BMC Oral Health **15**:44.

4.  **Autio-Gold J**. 2008. The Role of Chlorhexidine in Caries Prevention. Oper Dent **33**:710–716.

5.  **Lee D-W**, **Jung J-E**, **Yang Y-M**, **Kim J-G**, **Yi H-K**, **Jeon J-G**. 2016. The antibacterial activity of chlorhexidine digluconate against *Streptococcus mutans* biofilms follows sigmoidal patterns. Eur J Oral Sci **124**:440–446.

6.  **Liu J**, **Ling J-Q**, **Zhang K**, **Huo L-J**, **Ning Y**. 2012. Effect of sodium fluoride, ampicillin, and chlorhexidine on *Streptococcus mutans* biofilm detachment. Antimicrob Agents Chemother **56**:4532–5.

7.  **Jeon J-G**, **Rosalen PL**, **Falsetta ML**, **Koo H**. 2011. Natural products in caries research: current (limited) knowledge, challenges and future perspective. Caries Res **45**:243–63.

8.  **Gilbert P**, **Moore LE**. 2005. Cationic antiseptics: diversity of action under a common epithet. J Appl Microbiol **99**:703–715.

9.  **Kampf G**. 2018. Chlorhexidine Digluconate, p. 429–534. *In* Antiseptic Stewardship. Springer International Publishing, Cham.

10. **Cheung H-Y**, **Wong MM-K**, **Cheung S-H**, **Liang LY**, **Lam Y-W**, **Chiu S-K**. 2012. Differential Actions of Chlorhexidine on the Cell Wall of *Bacillus subtilis* and *Escherichia coli*. PLoS One **7**:e36659.

11. **Cieplik F**, **Kara E**, **Muehler D**, **Enax J**, **Hiller K-A**, **Maisch T**, **Buchalla W**. 2018. Antimicrobial efficacy of alternative compounds for use in oral care toward biofilms from caries-associated bacteria *in vitro*. Microbiologyopen e00695.

12. **Järvinen H**, **Tenovuo J**, **Huovinen P**. 1993. In vitro susceptibility of *Streptococcus mutans* to chlorhexidine and six other antimicrobial agents. Antimicrob Agents Chemother **37**:1158–9.

13. **Emilson CG**, **Ericson T**, **Heyden G**, **Lilia J**. 1972. Effect of chlorhexidine on human oral streptococci. J Periodontal Res **7**:189–191.

14. **Emilson CG**, **Fornell J**. 1976. Effect of toothbrushing with chlorhexidine gel on salivary microflora, oral hygiene, and caries. Scand J Dent Res **84**:308–19.

15. **Schiott CR**, **Loe H**. 1972. The sensitivity of oral streptococci to chlorhexidine. J Periodontal Res **7**:192–194.

16. **Wade WG**, **Addy M**. 1989. *In vitro* Activity of a Chlorhexidine–Containing Mouthwash Against Subgingival Bacteria. J Periodontol **60**:521–525.

17. **Maynard JH**, **Jenkins SM**, **Moran J**, **Addy M**, **Newcombe RG**, **Wade WG**. 1993. A 6-month home usage trial of a 1 % chlorhexidine toothpaste. (II). Effects on the oral microflora. J Clin Periodontol **20**:207–211.

18. **Bhardwaj P**, **Ziegler E**, **Palmer KL**. 2016. Chlorhexidine Induces VanA-Type Vancomycin Resistance Genes in Enterococci. Antimicrob Agents Chemother **60**:2209–21.

19. **Guzmán Prieto AM**, **Wijngaarden J**, **Braat JC**, **Rogers MRC**, **Majoor E**, **Brouwer EC**, **Zhang X**, **Bayjanov JR**, **Bonten MJM**, **Willems RJL**, **van Schaik W**. 2017. The Two-Component System ChtRS Contributes to Chlorhexidine Tolerance in *Enterococcus faecium*. Antimicrob Agents Chemother **61**:AAC.02122-16.

20. **Horner C**, **Mawer D**, **Wilcox M**. 2012. Reduced susceptibility to chlorhexidine in

staphylococci: is it increasing and does it matter? J Antimicrob Chemother **67**:2547–
2559.

21.  **Wand ME**, **Bock LJ**, **Bonney LC**, **Sutton JM**. 2017. Mechanisms of Increased
Resistance to Chlorhexidine and Cross-Resistance to Colistin following Exposure of
*Klebsiella pneumoniae* Clinical Isolates to Chlorhexidine. Antimicrob Agents Chemother
**61**:e01162-16.

22.  **Cieplik F**, **Jakubovics NS**, **Buchalla W**, **Maisch T**, **Hellwig E**, **Al-Ahmad A**. 2019.
Resistance Toward Chlorhexidine in Oral Bacteria – Is There Cause for Concern? Front
Microbiol **10**:587.

23.  **Palmer SR**, **Miller JH**, **Abranches J**, **Zeng L**, **Lefebure T**, **Richards VP**, **Lemos JA**,
**Stanhope MJ**, **Burne RA**. 2013. Phenotypic heterogeneity of genomically-diverse
isolates of *Streptococcus mutans*. PLoS One **8**:e61358.

24.  **Cornejo OE**, **Lefébure T**, **Bitar PDP**, **Lang P**, **Richards VP**, **Eilertson K**, **Do T**,
**Beighton D**, **Zeng L**, **Ahn S-J**, **Burne RA**, **Siepel A**, **Bustamante CD**, **Stanhope MJ**.
2013. Evolutionary and population genomics of the cavity causing bacteria *Streptococcus*
*mutans*. Mol Biol Evol **30**:881–93.

25.  **Belli WA**, **Marquis RE**. 1991. Adaptation of *Streptococcus mutans* and *Enterococcus*
*hirae* to acid stress in continuous culture. Appl Environ Microbiol **57**:1134–8.

26.  **Hamilton IR**, **St Martin EJ**. 1982. Evidence for the involvement of proton motive force in
the transport of glucose by a mutant of *Streptococcus mutans* strain DR0001 defective in
glucose-phosphoenolpyruvate phosphotransferase activity. Infect Immun **36**:567–75.

27.  **Marsh PD**. 2003. Are dental diseases examples of ecological catastrophes? Microbiology
**149**:279–294.

28.  **Fozo EM**, **Quivey RG**. 2004. Shifts in the membrane fatty acid profile of *Streptococcus*
*mutans* enhance survival in acidic environments. Appl Environ Microbiol **70**:929–36.

29.  **Liu Y**, **Zeng L**, **Burne RA**. 2009. AguR is required for induction of the *Streptococcus*
*mutans* agmatine deiminase system by low pH and agmatine. Appl Environ Microbiol
**75**:2629–37.

30.  **Bhardwaj P**, **Hans A**, **Ruikar K**, **Guan Z**, **Palmer KL**. 2018. Reduced Chlorhexidine and
Daptomycin Susceptibility in Vancomycin-Resistant *Enterococcus faecium* after Serial
Chlorhexidine Exposure. Antimicrob Agents Chemother **62**:e01235-17.

31.  **Renzoni A**, **Von Dach E**, **Landelle C**, **Diene SM**, **Manzano C**, **Gonzales R**, **Abdelhady**
**W**, **Randall CP**, **Bonetti EJ**, **Baud D**, **O'Neill AJ**, **Bayer A**, **Cherkaoui A**, **Schrenzel J**,
**Harbarth S**, **François P**. 2017. Impact of Exposure of Methicillin-Resistant
*Staphylococcus aureus* to Polyhexanide *In Vitro* and *In Vivo*. Antimicrob Agents
Chemother **61**:e00272-17.

32.  **Wilson DN**. 2014. Ribosome-targeting antibiotics and mechanisms of bacterial
resistance. Nat Rev Microbiol **12**:35–48.

33.  **Blinkhorn A**, **Bartold PM**, **Cullinan MP**, **Madden TE**, **Marshall RI**, **Raphael SL**,
**Seymour GJ**. 2009. Is there a role for triclosan/copolymer toothpaste in the management
of periodontal disease? Br Dent J **207**:117–125.

34.  **McNamara PJ**, **Levy SB**. 2016. Triclosan: an Instructive Tale. Antimicrob Agents
Chemother **60**:7015–7016.

35.  **Bender GR**, **Thibodeau EA**, **Marquis RE**. 1985. Reduction of acidurance of
streptococcal growth and glycolysis by fluoride and gramicidin. J Dent Res **64**:90–5.

36.  **Liao Y**, **Brandt BW**, **Li J**, **Crielaard W**, **Van Loveren C**, **Deng DM**. 2017. Fluoride
resistance in *Streptococcus mutans*: a mini review. J Oral Microbiol **9**:1344509.

37.  **Kreth J**, **Zhang Y**, **Herzberg MC**. 2008. Streptococcal antagonism in oral biofilms:
*Streptococcus sanguinis* and *Streptococcus gordonii* interference with *Streptococcus*
*mutans*. J Bacteriol **190**:4632–40.

38.  **Kreth J**, **Merritt J**, **Shi W**, **Qi F**. 2005. Competition and coexistence between

*Streptococcus mutans* and *Streptococcus sanguinis* in the dental biofilm. J Bacteriol
**187**:7193–203.

39. **Edlund A**, **Yang Y**, **Hall AP**, **Guo L**, **Lux R**, **He X**, **Nelson KE**, **Nealson KH**, **Yooseph**
**S**, **Shi W**, **McLean JS**. 2013. An *in vitro* biofilm model system maintaining a highly
reproducible species and metabolic diversity approaching that of the human oral
microbiome. Microbiome **1**:25.

40. **Kolderman E**, **Bettampadi D**, **Samarian D**, **Dowd SE**, **Foxman B**, **Jakubovics NS**,
**Rickard AH**. 2015. L-Arginine Destabilizes Oral Multi-Species Biofilm Communities
Developed in Human Saliva. PLoS One **10**:e0121835.

41. **Fernández CE**, **Aspiras MB**, **Dodds MW**, **González-Cabezas C**, **Rickard AH**. 2017.
The effect of inoculum source and fluid shear force on the development of *in vitro* oral
multispecies biofilms. J Appl Microbiol **122**:796–808.

42. **Clancy KA**, **Pearson S**, **Bowen WH**, **Burne RA**. 2000. Characterization of Recombinant,
Ureolytic *Streptococcus mutans* Demonstrates an Inverse Relationship between Dental
Plaque Ureolytic Capacity and Cariogenicity. Infect Immun **68**:2621–2629.

43. **Bowen WH**, **Burne RA**, **Wu H**, **Koo H**. 2017. Oral Biofilms: Pathogens, Matrix, and
Polymicrobial Interactions in Microenvironments. Trends Microbiol.

44. **Miller DJ**, **Jerga A**, **Rock CO**, **White SW**. 2008. Analysis of the *Staphylococcus aureus*
DgkB Structure Reveals a Common Catalytic Mechanism for the Soluble Diacylglycerol
Kinases. Structure **16**:1036–1046.

45. **Jerga A**, **Lu Y-J**, **Schujman GE**, **de Mendoza D**, **Rock CO**. 2007. Identification of a
soluble diacylglycerol kinase required for lipoteichoic acid production in *Bacillus subtilis*. J
Biol Chem **282**:21738–45.

46. **Shields RC**, **Zeng L**, **Culp DJ**, **Burne RA**. 2018. Genomewide Identification of Essential
Genes and Fitness Determinants of *Streptococcus mutans* UA159. mSphere **3**:e00031-
18.

47. **Webb AJ**, **Karatsa-Dodgson M**, **Gründling A**. 2009. Two-enzyme systems for glycolipid
and polyglycerolphosphate lipoteichoic acid synthesis in *Listeria monocytogenes*. Mol
Microbiol **74**:299–314.

48. **Reichmann NT**, **Gründling A**. 2011. Location, synthesis and function of glycolipids and
polyglycerolphosphate lipoteichoic acid in Gram-positive bacteria of the phylum
Firmicutes. FEMS Microbiol Lett **319**:97–105.

49. **Theilacker C**, **Sanchez-Carballo P**, **Toma I**, **Fabretti F**, **Sava I**, **Kropec A**, **Holst O**,
**Huebner J**. 2009. Glycolipids are involved in biofilm accumulation and prolonged
bacteraemia in *Enterococcus faecalis*. Mol Microbiol **71**:1055–1069.

50. **Doran KS**, **Engelson EJ**, **Khosravi A**, **Maisey HC**, **Fedtke I**, **Equils O**, **Michelsen KS**,
**Arditi M**, **Peschel A**, **Nizet V**. 2005. Blood-brain barrier invasion by group B
*Streptococcus* depends upon proper cell-surface anchoring of lipoteichoic acid. J Clin
Invest **115**:2499–507.

51. **Seaton K**, **Ahn S-J**, **Sagstetter AM**, **Burne RA**. 2011. A transcriptional regulator and
ABC transporters link stress tolerance, (p)ppGpp, and genetic competence in
*Streptococcus mutans*. J Bacteriol **193**:862–74.

52. **Ahn S-J**, **Kaspar J**, **Kim JN**, **Seaton K**, **Burne R a**. 2014. Discovery of novel peptides
regulating competence development in *Streptococcus mutans*. J Bacteriol **196**:3735–
3745.

53. **Loe H**, **Rindom Schiott C**. 1970. The effect of mouthrinses and topical application of
chlorhexidine on the development of dental plaque and gingivitis in man. J Periodontal
Res **5**:79–83.

54. **Chang JC**, **LaSarre B**, **Jimenez JC**, **Aggarwal C**, **Federle MJ**. 2011. Two Group A
Streptococcal Peptide Pheromones Act through Opposing Rgg Regulators to Control
Biofilm Development. PLoS Pathog **7**:e1002190.

55.   **Chen L**, **Chakraborty B**, **Zou J**, **Burne RA**, **Zeng L**. 2019. Amino Sugars Modify
Antagonistic Interactions between Commensal Oral Streptococci and *Streptococcus*
*mutans*. Appl Environ Microbiol AEM.00370-19.

56.   **Lau PC.**, **Sung CK**, **Lee JH**, **Morrison DA**, **Cvitkovitch DG**. 2002. PCR ligation
mutagenesis in transformable streptococci: application and efficiency. J Microbiol
Methods **49**:193–205.

57.   **Kaspar J**, **Ahn S-J**, **Palmer SR**, **Choi SC**, **Stanhope MJ**, **Burne RA**. 2015. A Unique
ORF within the *comX* gene of *Streptococcus mutans* Regulates Genetic Competence and
Oxidative Stress Tolerance. Mol Microbiol **96**:463–482.

58.   **Tritt A**, **Eisen JA**, **Facciotti MT**, **Darling AE**. 2012. An Integrated Pipeline for de Novo
Assembly of Microbial Genomes. PLoS One **7**:e42304.

59.   **Gurevich A**, **Saveliev V**, **Vyahhi N**, **Tesler G**. 2013. QUAST: quality assessment tool for
genome assemblies. Bioinformatics **29**:1072–1075.

60.   **Darling AE**, **Tritt A**, **Eisen JA**, **Facciotti MT**. 2011. Mauve Assembly Metrics.
Bioinformatics **27**:2756–2757.

**TABLES**

836    TABLE 1. Growth characteristics of RSV3s under different conditions.

| Strain | Growth Characteristics | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| | BHI pH 7.0 | | BHI pH 7.0 + 0.75 µg mL-1 CHX | | BHI pH 5.5 | | BHI pH 5.5 + 0.75 µg mL-1 CHX | |
| | Doubling Time (min) | Final Yield (OD $_{600\,nm}$) | Doubling Time (min) | Final Yield (OD $_{600\,nm}$) | Doubling Time (min) | Final Yield (OD $_{600\,nm}$) | Doubling Time (min) | Final Yield (OD $_{600\,nm}$) |
| Parental | 51 ± 2 | 0.61 ± 0.08 | 623 ± 69 | 0.29 ± 0.09 | 178 ± 12 | 0.65 ± 0.01 | No growth | 0.04 ± 0.02 |
| RSV3-1 | 145 ± 26 | 0.45 ± 0.11 | 126 ± 19 | 0.41 ± 0.10 | 186 ± 11 | 0.45 ± 0.02 | 142 ± 14 | 0.23 ± 0.07 |
| RSV3-3 | 56 ± 2 | 0.73 ± 0.14 | 78 ± 7 | 0.50 ± 0.04 | 259 ± 25 | 0.41 ± 0.01 | 142 ± 12 | 0.48 ± 0.02 |
| RSV3-6 | 141 ± 16 | 0.55 ± 0.08 | 139 ± 17 | 0.44 ± 0.05 | 132 ± 14 | 0.32 ± 0.02 | 145 ± 8 | 0.31 ± 0.02 |
| RSV3-7 | 108 ± 11 | 0.67 ± 0.05 | 143 ± 32 | 0.57 ± 0.11 | 142 ± 44 | 0.47 ± 0.02 | 119 ± 11 | 0.59 ± 0.09 |
| RSV3-8 | 66 ± 7 | 0.72 ± 0.03 | 92 ± 13 | 0.59 ± 0.11 | 189 ± 34 | 0.54 ± 0.01 | 121 ± 12 | 0.59 ± 0.04 |
| RSV3-9 | 97 ± 8 | 0.63 ± 0.04 | 133 ± 12 | 0.60 ± 0.01 | 160 ± 10 | 0.34 ± 0.01 | 215 ± 23 | 0.39 ± 0.11 |
| RSV3-10 | 63 ± 6 | 0.56 ± 0.12 | 74 ± 5 | 0.52 ± 0.01 | 107 ± 7 | 0.55 ± 0.03 | 113 ± 22 | 0.59 ± 0.03 |

838    TABLE 2. Changes in antibiotic sensitivity for RSV3s.

| Antibiotic | Mechanism of Action | MIC Determined by ETest® Strips ($\mu$g mL$^{-1}$) | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| | | Parental | RSV3-1 | RSV3-3 | RSV3-6 | RSV3-7 | RSV3-8 | RSV3-9 | RSV3-10 |
| Amoxicillin | Inhibition of peptidoglycan cross-linkage | 0.094 | 0.047 | 0.094 | 0.094 | 0.064 | 0.094 | 0.094 | 0.094 |
| Azithromycin | Inhibition of protein synthesis | 3 | 3 | 4 | 1.5 | 3 | 4 | 4 | 4 |
| Bacitracin | Interference of peptidoglycan precursor transport to cell wall | 128 | 48 | 64 | 96 | 96 | 96 | 192 | 96 |
| Clindamycin | Inhibition of protein synthesis | 0.064 | 0.125 | 0.094 | 0.125 | 0.094 | 0.125 | 0.19 | 0.125 |
| Daptomycin | Insertion into cell membrane; loss of membrane potential | 2 | 1.5 | 3 | 4 | 4 | 3 | 12 | 3 |
| Vancomycin | Inhibition of peptidoglycan cross-linkage | 1.5 | 1 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 |

839
840

841     TABLE 3. SMU.1542c Mutations in RSV3s

| RSV Strain Containing Mutation | Genome SNP[*] | Genome Position[%] | 1542c Location[$] | 1542c SNP^ | 1542c aa Location[&] | 1542c aa Change[+] |
|---|---|---|---|---|---|---|
| 6, 7, 9 | c > t | 1471014 | 391 | g > a | 131 | G > S |
| 8, 10 | g > t | 1470494 | 911 | c > a | 304 | A > D |
| 1 | c > t | 1470914 | 491 | g > a | 164 | G > D |
| 3 | c > t | 1470659 | 746 | g > a | 249 | G > E |

842
[*]Refers to the base change on the forward chromosome strand; 1542c is encoded in the reverse orientation
[%]Location of the base pair that encodes the SNP in reference to the entire genome
[$]Location of the base pair that encodes the SNP in reference to the 1542c gene alone
^Refers to the base change on the reverse chromosome strand in which 1542c is encoded
[&]Location of the amino acid that encodes the SNP in reference to the 1542c gene alone
[+]Refers to the amino acid change within the 1542c coding sequence due to the SNP
849
850
851

852    TABLE 4. SMU.1588c Mutations in RSV3s

| RSV Strain Containing Mutation | Genome SNP[*] | Genome Location[%] | 1588c Location[$] | 1588c SNP[^] | 1588c aa Location[&] | 1588c aa Change[+] |
|---|---|---|---|---|---|---|
| 6, 8 | g > c | 1518523 | 1295 | c > g | 433 | P > A |
| 10 | c > t | 1518644 | 1174 | g > a | 391 | W > STOP |

853
[*]Refers to the base change on the forward chromosome strand; 1542c is encoded in the reverse orientation
[%]Location of the base pair that encodes the SNP in reference to the entire genome
[$]Location of the base pair that encodes the SNP in reference to the 1542c gene alone
[^]Refers to the base change on the reverse chromosome strand in which 1542c is encoded
[&]Location of the amino acid that encodes the SNP in reference to the 1542c gene alone
[+]Refers to the amino acid change within the 1542c coding sequence due to the SNP

**Table S1.** SNPs of the RSV3s.

| RSV3-1 | | | | | | |
|---|---|---|---|---|---|---|
| Number | SNP pattern | Gene | Name | Function | Location | |
| 1 | ct | SMU.922 | *rcrP* | ABC transporter | NC_004350 | 875439 |
| 2 | at | SMU.974 | *potB* | putative spermidine/putrescine ABC transporter | NC_004350 | 920843 |
| 3 | ct | SMU.1542c | | lipid kinase | NC_004350 | 1470914 |

| RSV3-3 | | | | | | |
|---|---|---|---|---|---|---|
| Number | SNP pattern | Gene | Name | Function | Location | |
| 1 | ca | SMU.577 | *lytS* | histidine kinase | NC_004350 | 540225 |
| 2 | ga | SMU.895 | | DNA damage inducible protein | NC_004350 | 849956 |
| 3 | ct | SMU.1344c | | malonyl-CoA acyl-carrier protein transacylase | NC_004350 | 1284607 |
| 4 | ct | SMU.1542c | | lipid kinase | NC_004350 | 1470659 |
| 5 | ga | SMU.1989c | *rpoC* | DNA-directed RNA polymerase | NC_004350 | 1857079 |

| RSV3-6 | | | | | | |
|---|---|---|---|---|---|---|
| Number | SNP pattern | Gene | Name | Function | Location | |
| 1 | ag | IGR SMU.685 - SMU.687c | | hypothetical protein | NC_004350 | 652281 |
| 2 | gc | SMU.1004 | *gtfB* | glucosyltransferase | NC_004350 | 953114 |
| 3 | tg | SMU.1004 | *gtfB* | glucosyltransferase | NC_004350 | 953123 |
| 4 | gt | SMU.1004 | *gtfB* | glucosyltransferase | NC_004350 | 953124 |
| 5 | at | SMU.1004 | *gtfB* | glucosyltransferase | NC_004350 | 953133 |
| 6 | ag | SMU.1004 | *gtfB* | glucosyltransferase | NC_004350 | 953144 |
| 7 | tc | SMU.1004 | *gtfB* | glucosyltransferase | NC_004350 | 953145 |
| 8 | at | SMU.1004 | *gtfB* | glucosyltransferase | NC_004350 | 953151 |
| 9 | ta | SMU.1004 | *gtfB* | glucosyltransferase | NC_004350 | 953153 |
| 10 | tg | SMU.1004 | *gtfB* | glucosyltransferase | NC_004350 | 953154 |
| 11 | tg | SMU.1004 | *gtfB* | glucosyltransferase | NC_004350 | 953157 |
| 12 | ag | SMU.1004 | *gtfB* | glucosyltransferase | NC_004350 | 953158 |
| 13 | cg | SMU.1004 | *gtfB* | glucosyltransferase | NC_004350 | 953166 |
| 14 | ct | SMU.1542c | | lipid kinase | NC_004350 | 1471014 |
| 15 | gc | SMU.1588c | | hexosyltransferase | NC_004350 | 1518523 |
| 16 | ac | SMU.1990c | *rpoB* | DNA-directed RNA polymerase subunit | NC_004350 | 1862650 |
| 17 | ac | IGR SMU.2121c - SMU.2123 | | hypothetical protein | NC_004350 | 1990740 |

| RSV3-7 | | | | | | |
|---|---|---|---|---|---|---|
| Number | SNP pattern | Gene | Name | Function | Location | |

| | | | | | | |
|---|---|---|---|---|---|---|
| 1 | t C | SMU.25 | recO | DNA repair protein | NC_00435 0 | 29442 |
| 2 | t A | SMU.25 IGR | | DNA repair protein - phosphate acyltransferase | NC_00435 0 | 29489 |
| 3 | aT | SMU.25 IGR | | DNA repair protein - phosphate acyltransferase | NC_00435 0 | 29540 |
| 4 | aC | SMU.25 IGR | | DNA repair protein - phosphate acyltransferase | NC_00435 0 | 29559 |
| 5 | cA | SMU.25 IGR | | DNA repair protein - phosphate acyltransferase | NC_00435 0 | 29598 |
| 6 | aG | SMU.44 | | DNA mismatch repair | NC_00435 0 | 47444 |
| 7 | aG | SMU.44 | | DNA mismatch repair | NC_00435 0 | 47530 |
| 8 | aG | SMU.44 IGR | | DNA mismatch repair - hypothetical protein | NC_00435 0 | 47892 |
| 9 | gT | SMU.44 IGR | | DNA mismatch repair - hypothetical protein | NC_00435 0 | 49100 |
| 10 | aT | pyrG - fbaA IGR | | fructose bisphosphate aldolase | NC_00435 0 | 100116 |
| 11 | aT | SMU.112c | | transcriptional regulator | NC_00435 0 | 112507 |
| 12 | t A | purA - sgaT IGR | | adenylosuccinate synthetase - PTS | NC_00435 0 | 261148 |
| 13 | t A | purA - sgaT IGR | | adenylosuccinate synthetase - PTS | NC_00435 0 | 261178 |
| 14 | gC | SMU.502 | | hypothetical protein | NC_00435 0 | 470965 |
| 15 | t C | SMU.504 | dam | site-specific DNA methylase | NC_00435 0 | 474498 |
| 16 | aG | SMU.515 - SMU.516 IGR | | hypothetical protein(s) | NC_00435 0 | 484132 |
| 17 | aG | SMU.562c - SMU.563 IGR | dpE | ATP-dependent protease | NC_00435 0 | 527485 |
| 18 | t G | SMU.562c - SMU.563 IGR | dpE | ATP-dependent protease | NC_00435 0 | 527506 |
| 19 | aC | SMU.562c - SMU.563 IGR | dpE | ATP-dependent protease | NC_00435 0 | 527563 |
| 20 | cT | SMU.597 | pbp2 b | pencillin binding protein 2b | NC_00435 0 | 554999 |
| 21 | t C | SMU.597 | pbp2 b | pencillin binding protein 2b | NC_00435 0 | 555170 |
| 22 | aC | SMU.616 | | hypothetical protein | NC_00435 0 | 578092 |
| 23 | aG | SMU.618 - SMU.620 IGR | | hypothetical protein(s) | NC_00435 0 | 578744 |
| 24 | cT | SMU.618 - SMU.620 IGR | | hypothetical protein(s) | NC_00435 0 | 578748 |
| 25 | t A | SMU.618 - SMU.620 IGR | | hypothetical protein(s) | NC_00435 0 | 578750 |
| 26 | aG | SMU.683 | | ATP-binding protein | NC_00435 0 | 648149 |
| 27 | aT | SMU.758c - SMU.759 IGR | | hypothetical protein - protease | NC_00435 0 | 710076 |
| 28 | aT | SMU.758c - SMU.759 IGR | | hypothetical protein - protease | NC_00435 0 | 710126 |
| 29 | t A | SMU.773c - SMU.774 IGR | | tRNA-synthetase - hypothetical protein | NC_00435 0 | 724043 |
| 30 | gT | SMU.773c - SMU.774 IGR | | tRNA-synthetase - hypothetical protein | NC_00435 0 | 724044 |
| 31 | t C | SMU.773c - SMU.774 IGR | | tRNA-synthetase - hypothetical protein | NC_00435 0 | 724046 |
| 32 | aC | SMU.773c - SMU.774 IGR | | tRNA-synthetase - hypothetical protein | NC_00435 0 | 724048 |
| 33 | aG | SMU.773c - SMU.774 IGR | | tRNA-synthetase - hypothetical protein | NC_00435 0 | 724049 |

| Number | SNP pattern | Gene | Name | Function | Location | |
|---|---|---|---|---|---|---|
| 34 | aC | SMU.773c - SMU.774 IGR | | tRNA-synthetase - hypothetical protein | NC_004350 | 724050 |
| 35 | cA | SMU.773c - SMU.774 IGR | | tRNA-synthetase - hypothetical protein | NC_004350 | 724053 |
| 36 | tA | SMU.773c - SMU.774 IGR | | tRNA-synthetase - hypothetical protein | NC_004350 | 724057 |
| 37 | aG | SMU.773c - SMU.774 IGR | | tRNA-synthetase - hypothetical protein | NC_004350 | 724059 |
| 38 | gA | SMU.773c - SMU.774 IGR | | tRNA-synthetase - hypothetical protein | NC_004350 | 724061 |
| 39 | aT | SMU.773c - SMU.774 IGR | | tRNA-synthetase - hypothetical protein | NC_004350 | 724094 |
| 40 | aC | SMU.895 | | DNA-damage inducible protein | NC_004350 | 850090 |
| 41 | tG | SMU.923 - SMU.924 | | Rcr peptides SNP? | NC_004350 | 878588 |
| 42 | tC | SMU.992 - SMU.993 IGR | | hypothetical protein - GTPase | NC_004350 | 939662 |
| 43 | tC | topA - gld IGR | | DNA topoisomerase - tRNA | NC_004350 | 949062 |
| 44 | aC | SMU.1010c | citC | citrate ligase | NC_004350 | 965767 |
| 45 | tA | SMU.1010c | citC | citrate ligase | NC_004350 | 965816 |
| 46 | aT | SMU.1033 - SMU.1034c IGR | | site-specific tyrosine recombinase | NC_004350 | 982851 |
| 47 | cG | SMU.1197 IGR | | hypothetical protein | NC_004350 | 1140398 |
| 48 | gT | SMU.1469 - SMU.1470c IGR | | hypothetical protein | NC_004350 | 1395952 |
| 49 | tG | SMU.1469 - SMU.1470c IGR | | hypothetical protein | NC_004350 | 1395963 |
| 50 | aC | covR - glnQ IGR | | response regulator - ABC transporter | NC_004350 | 1447027 |
| 51 | aG | covR - glnQ IGR | | response regulator - ABC transporter | NC_004350 | 1447078 |
| 52 | gT | SMU.1542c | | lipid kinase | NC_004350 | 1470494 |
| 53 | tG | SMU.1576c | | hypothetical protein | NC_004350 | 1503438 |
| 54 | gT | SMU.1708 | trkA | potassium transporter | NC_004350 | 1619470 |
| 55 | aC | SMU.1848 | | hypothetical protein | NC_004350 | 1745705 |
| 56 | aT | SMU.1895c - 1897 iGR | | hypothetical protein - ABC transporter | NC_004350 | 1783686 |

| RSV3-8 | | | | | | |
|---|---|---|---|---|---|---|
| Number | SNP pattern | Gene | Name | Function | Location | |
| 1 | gA | SMU.43 | | site-specific DNA methyltransferase modification protein | NC_004350 | 46486 |
| 2 | cT | SMU.369c - SMU.370 IGR | | hypothetical protein - ABC transporter | NC_004350 | 352139 |
| 3 | tA | SMU.470 | | hypothetical protein | NC_004350 | 439254 |
| 4 | aT | SMU.773c - SMU.774 IGR | | tRNA-synthetase - hypothetical protein | NC_004350 | 724145 |
| 5 | tG | SMU.804 | | hypothetical protein | NC_004350 | 750816 |
| 6 | tG | topA - gld IGR | | DNA topoisomerase - tRNA | NC_004350 | 949107 |
| 7 | cT | SMU.1542c | | lipid kinase | NC_004350 | 1471014 |
| 8 | gC | SMU.1588c | | hexosyltransferase | NC_004350 | 1518523 |
| 9 | aC | SMU.2121c - SMU.2123 | | hypothetical protein | NC_004350 | 1990749 |

| | | IGR | | | 0 | 0 |
|---|---|---|---|---|---|---|

| RSV3-9 | | | | | | |
|---|---|---|---|---|---|---|
| Number | SNP pattern | Gene | Name | Function | Location | |
| 1 | gT | SMU.750c | | hypothetical protein | NC_004350 | 703503 |
| 2 | cG | SMU.875c | | transposase | NC_004350 | 826253 |
| 3 | tC | SMU.1005 | *gtfC* | glucosyltransferase | NC_004350 | 957858 |
| 4 | cT | SMU.1005 | *gtfC* | glucosyltransferase | NC_004350 | 957915 |
| 5 | aG | SMU.1005 | *gtfC* | glucosyltransferase | NC_004350 | 957924 |
| 6 | gC | SMU.1134c | | phosphate transporter | NC_004350 | 1076258 |
| 7 | tA | SMU.1332c - SMU.1334c IGR | | transposase – phosphopantetheinyl transferase | NC_004350 | 1255056 |
| 8 | tC | SMU.1332c - SMU.1334c IGR | | transposase - phosphopantetheinyl transferase | NC_004350 | 1255058 |
| 9 | aT | SMU.1336c | *pksD* | hypothetical protein | NC_004350 | 1257526 |
| 10 | tC | SMU.1407c | | Transposase | NC_004350 | 1337500 |
| 11 | cT | SMU.1542c | | lipid kinase | NC_004350 | 1471014 |
| 12 | tA | SMU.1966c - SMU.1967c IGR | | periplasmic sugar binding protein - ssb2 | NC_004350 | 1843377 |
| 13 | gT | SMU.1966c - SMU.1967c IGR | | periplasmic sugar binding protein - ssb2 | NC_004350 | 1843380 |
| 14 | aC | SMU.2121c - SMU.2123 IGR | | hypothetical protein | NC_004350 | 1990740 |
| 15 | aC | SMU.2163 | *htrA* | serine protease | NC_004350 | 2030550 |

| RSV3-10 | | | | | | |
|---|---|---|---|---|---|---|
| Number | SNP pattern | Gene | Name | Function | Location | |
| 1 | cT | SMU.895 | | DNA-damage-inducible protein | NC_004350 | 850077 |
| 2 | gA | SMU.1188c | *lepB* | signal peptidase | NC_004350 | 1129851 |
| 3 | aT | SMU.1188c | *lepB* | signal peptidase | NC_004350 | 1129852 |
| 4 | gT | SMU.1188c | *lepB* | signal peptidase | NC_004350 | 1129856 |
| 5 | tC | SMU.1188c | *lepB* | signal peptidase | NC_004350 | 1129860 |
| 6 | aT | SMU.1188c | *lepB* | signal peptidase | NC_004350 | 1129868 |
| 7 | gA | SMU.1188c | *lepB* | signal peptidase | NC_004350 | 1129870 |
| 8 | tC | SMU.1337c | | alpha/beta hydrolase | NC_004350 | 1258225 |
| 9 | gT | SMU.1542c | | lipid kinase | NC_004350 | 1470494 |
| 10 | cT | SMU.1588c | | hexosyltransferase | NC_004350 | 1518644 |
| 11 | cT | SMU.1773c | | hypothetical protein | NC_004350 | 1677179 |

**Table S2.** Strains Used in This Study.

| Strain | Respective Characteristics* | Reference |
|---|---|---|
| *S. mutans* strains | | |
| UA159 | Wild-type | Lab Stock |
| RSV3-1 | Third-generation CHX passaged variant | This study |
| RSV3-3 | Third-generation CHX passaged variant | This study |
| RSV3-6 | Third-generation CHX passaged variant | This study |
| RSV3-7 | Third-generation CHX passaged variant | This study |
| RSV3-8 | Third-generation CHX passaged variant | This study |
| RSV3-9 | Third-generation CHX passaged variant | This study |
| RSV3-10 | Third-generation CHX passaged variant | This study |
| $\Delta$804 | Non-polar Km marker replacement of SMU.804; $Km^R$ | This study |
| $\Delta$1576c | Non-polar Km marker replacement of SMU.1576c; $Km^R$ | This study |
| *S. gordonii* strains | | |
| DL1 | Wild-type | Lab Stock |
| *S. sanguinis* strains | | |
| SK150 | Wild-type | Lab Stock |
| *S. australis*-like strains | | |
| A12 | Wild-type | Lab Stock |

* Km, Kanamycin

**Table S3.** Primers Used in This Study.

| Primer Name | Sequence (5' -> 3')* | Use |
| --- | --- | --- |
| 16S-F | AGA GTT TGA TCC TGG CTC AG | 16S rRNA Sequencing of RSV3s |
| 16S-R | TAC GGG TAC CTT GTT ACG ACT | 16S rRNA Sequencing of RSV3s |
| | | |
| 804-F | CCT GTG GTT TGT CGC CTA GTC TC | Forward Amplification of SMU.804 for Sequencing |
| 804-R | GCC ACT ATT GTG GAC TGA CTT GCA C | Reverse Amplification of SMU.804 for Sequencing |
| 804-Seq1 | GTT TTT CAA TGC ACC TTA CCT CC | Nested Sequencing Primer 1 of SMU.804 |
| 804-Seq2 | ACA AGG ATG TGA TTA TTC CGA AG | Nested Sequencing Primer 2 of SMU.804 |
| 804-Seq3 | GAG GAG AAT AAG ATA GAT ATT TCA GG | Nested Sequencing Primer 3 of SMU.804 |
| 804-Seq4 | GTT TAG CAA CAC TGT TAT TGC | Nested Sequencing Primer 4 of SMU.804 |
| | | |
| 804-A | CGT CGT CAA ATC GCT GAC TGT C | PCR for Arm 1 in Construction of ∆804 |
| 804-B | CAT G<u>GG ATC C</u>TG ACT TTT GAA GCG | PCR for Arm 1 in Construction of ∆804 |
| 804-C | GAG <u>GGA TCC</u> TTC ATT CTC AGT TCC GAC | PCR for Arm 2 in Construction of ∆804 |
| 804-D | GGA AGA AGC ACA AAA GCA TGG CAT GG | PCR for Arm 2 in Construction of ∆804 |
| | | |
| 1542c-F | GTC TTG GTG CTA ATG TAG CTG G | Forward Amplification of SMU.1542c for Sequencing |
| 1542c-R | CTG TCC GCT CTG CCA TTG C | Reverse Amplification of SMU.1542c for Sequencing |
| 1542c-Seq1 | AAC CAG TCT GGT CAT AGC GG | Nested Sequencing Primer 1 of SMU.1542c |
| 1542c-Seq2 | GAA GCT GCA CGT GCC AGT CAA GC | Nested Sequencing Primer 2 of SMU.1542c |
| 1542c-Seq3 | TCA TAT GGA TAT TGG TAG GGC | Nested Sequencing Primer 3 of SMU.1542c |
| 1542c-Seq4 | GGA GGC TTT GAA ACG ATT GC | Nested Sequencing Primer 4 of SMU.1542c |
| | | |
| 1576c-F | CAT TTG ATA TAG ATC CAA ATG ATG AGG | Forward Amplification of SMU.1576c for Sequencing |
| 1576c-R | CAA GTG AAG GAA TTT CAA TCT CAT C | Reverse Amplification of SMU.1576c for Sequencing |
| | | |
| 1576c-A | CGA AAT TTT AGA AGT TTT GGA TCA GG | PCR for Arm 1 in Construction of ∆1576c |
| 1576c-B | CTA TTT <u>GGA TCC</u> CCA TAT ATT AAA TCT CC | PCR for Arm 1 in Construction of ∆1576c |
| 1576c-C | GTA <u>GGA TCC</u> AGT CTT AAA AAT TGA GGG | PCR for Arm 2 in Construction of ∆1576c |
| 1576c-D | CAT TAG CTG TAG CAA GTC CAT GG | PCR for Arm 2 in Construction of ∆1576c |
| | | |
| 1588c-F | GGC TGG ATA ACT GAT GAA GCA G | Forward Amplification of SMU.1588c for Sequencing |
| 1588c-R | GGC CTG CAC TTT CTC AGC TGC C | Reverse Amplification of SMU.1588c for Sequencing |
| 1588c-Seq1 | AGA GGC AGG TTA TTT AGT CGC CC | Nested Sequencing Primer 1 of SMU.1588c |
| 1588c-Seq2 | CGC AGA CAG AAT TTA GTC TGG G | Nested Sequencing Primer 2 of SMU.1588c |
| 1588c-Seq3 | GAC ATT AGT CAA GAA GAT ATT GCC | Nested Sequencing Primer 3 of SMU.1588c |
| 1588c-Seq4 | GCC GAT TAT CGC TCA GTC CAA TCC | Nested Sequencing Primer 4 of SMU.1588c |

*Underlines represent BamHI Restriction Sites used in cloning

**Table S4.** NCBI BioProject PRJNA515165.

| BioSample | Experiment | Sample Name | Library Name | Organism | Strain | RSV3 isolate |
|---|---|---|---|---|---|---|
| SAMN10743465 | SRX5250866 | JK1 | JK1_S40 | Streptococcus mutans | UA159 | 1 |
| SAMN10743467 | SRX5250863 | JK3 | JK3_S42 | Streptococcus mutans | UA159 | 3 |
| SAMN10743470 | SRX5250857 | JK6 | JK6_S45 | Streptococcus mutans | UA159 | 6 |
| SAMN10743471 | SRX5250859 | JK7 | JK7_S46 | Streptococcus mutans | UA159 | 7 |
| SAMN10743472 | SRX5250858 | JK8 | JK8_S47 | Streptococcus mutans | UA159 | 8 |
| SAMN10743473 | SRX5250867 | JK9 | JK9_S48 | Streptococcus mutans | UA159 | 9 |
| SAMN10743474 | SRX5250866 | JK10 | JK10_S49 | Streptococcus mutans | UA159 | 10 |
| SAMN10743475 | SRX5250860 | JKW | JKW_S39 | Streptococcus mutans | UA159 | parental |

**Figure S1**

**Supplemental Figure 1. MIC Testing of Smu Isolates to CHX.** Selected

*Streptococcus mutans* (Smu) isolates along with the wild-type strain UA159 grown for

24 h in BHI with the designated concentrations of CHX in microtiter plates to determine

minimal inhibitory concentration (MIC). Red dots denote highest CHX concentration that

the strains displayed growth ($OD_{600\ nm}$ > 0.1). Four biological replicates of each isolate

were tested during the experiment.

**Figure S2**

**Supplemental Figure 2. Results of MIC Testing Using ETest Strips.** To determine the MIC of various antibiotics using different mechanisms of action on the parental UA159 and RSV3s, 10 µL from a mid-exponential culture of the respective strain was added to 10 mL soft BHI agar (0.75% agarose) and overlaid on to BHI agar plates. Once solidified, ETEST strips (bioMérieux, Inc., Durham, N.C.) containing either Amoxicillin, Azithromycin, Bacitracin, Clindamycin, Daptomycin or Vancomycin when then laid over the agar with sterile forceps. After 48 h of growth at 37°C in a 5% $CO_2$ aerobic atmosphere, the agar plates were examined, imaged and lowest MIC recorded where no growth was visible. Each image above is one representative image from three independent biological replicates tested.

**Figure S3**

**Supplemental Figure 3. Agar Plate-based Antagonism of RSV3 Strains by**

***Streptococcus sanguinis* SK150.** Agar plate competition assay between *S. sanguinis*

SK150 (labeled S. s, left colony) and either the parental strain UA159 or the RSV3s

(right colony). Colonies were spotted at the same time on BHI agar plates after growth

in CDM medium to an $OD_{600\ nm}$ = 0.5. The plates were incubated for 48 h at 37°C in a

5% $CO_2$ aerobic atmosphere prior to imaging.

**Figure S4**

**Supplemental Figure 4. Diversity of Cell-Containing Saliva Biofilm.** A cell-

containing saliva control biofilm (no inoculation of parental *S. mutans* UA159 or RSV3s)

was grown for 24 h in the chemically defined medium CDM supplemented with 20 mM

glucose and 5 mM sucrose within an ibidi 15 µ-Slide 8-well glass bottom slide. After 24

h, the biofilm was washed and removed by scraping off attached cells with a pipette tip

in 1x PBS. Cells were sonicated 4x 30 s prior to serial dilution and plating on BHI agar

plates. Left agar plate image is the outgrowth of cells after 24 hours in incubation at

37°C in a 5% $CO_2$ aerobic atmosphere, right agar plate image is 72 hours incubation.

**Figure S5**

**Supplemental Figure 5. Recorded initial and final CFUs of *S. mutans* parental UA159 and RSV3s within a CCS biofilm model.** *S. mutans* UA159 (parental) and RSV3 isolates were all grown in CDM medium from overnight cultures to an $OD_{600\ nm}$ = 0.5. At this point, all strains were inoculated individually into CDM supplemented with 20 mM glucose and 5 mM sucrose at a 1:1000 dilution along with an aliquot of cell-containing saliva (CCS) at a 1:50 dilution. 0.35 mL of this inoculated medium was loaded into an ibidi 15 μ-Slide 8-well glass bottom slide and incubated at 37°C in a 5% $CO_2$ aerobic atmosphere for either (A) t = 24 h or (B) t = 48 h. Biofilms were washed to remove loosely bound cells and then removed by scraping off attached cells with a pipette tip in 1x PBS**.** Cells were serially diluted after sonication and plated on mitis salivarius-bacitracin agar supplemented with potassium tellurite and 20% sucrose to select for *S. mutans* only. CFUs were counted after 72 h incubation at 37°C in a 5% $CO_2$.

**Figure S6**

```
S. aureus     ---MRKRARIIYNPTSGKEQFKRELPDALIKLEKAGYETSAYATEKIGD-ATLEAERAMH  56
S. mutans     MVRKQKRARLIYNPTSGQEIMKKNVAEVLDILESFGYETSAFQTTPRADSAKNEAARASQ  60
              :****:*******:* :*::: :.*  **. *****: *   .* *. ** ** :

S. aureus     ENYDVLIAAGGDGTLNEVVNGIAEKPNRPKLGVIPMGTVNDFGRALHIPNDIMGA-LDVI  115
S. mutans     AGFDLVIAAGGDGTINEVVSGIAPLKRRPKVAIIPTGTTNDFARALKIPRGNPVEATKLI  120
              .:*::*********:****.***   .***:.:** **.***.***:**..      .:*

                      G131S                                G164D
S. aureus     IEGHSTKVDIGK-MNNRYFINLAAGGQLTQVSYETPSKLKSIVGPFAYYIKGFEMLPQMK  174
S. mutans     GKNQTIHMDIGRAYNDKYFINIAAAGSLTELTYSVPSQLKTMFGYLAYLVKGAELLPRVR  180
              :.:: ::***:   *::****:**.*.**:::*..**:**::.* :** :** *:**:::

S. aureus     AVDLRIEYDGNVFQGEALLFFLGLTNSMAGFEKLVPDAKLDDGYFTLIIVEKSNLAELGH  234
S. mutans     NVPVKVFYDHGVFEGEASMIFVALTNSVGGFETIAPDAKLDDGKFTLILVKTANLFELTD  240
              * ::: ** .**:*** ::*:.****:.***.:.******** ****:*::*** ** .

                      G249E
S. aureus     IMTLAS-RGEHTKHPKVIYEKAKAINIS----SFTDLQLNVDGEYGGKLPANFLNLERHI  289
S. mutans     LIRLVLTGGRHINHKKIEYIKTSYLKIIPQGNPPKRMMINLDGEYGGDAPIELRNLKNHI  300
              :: *.   *.* :* *: * *:. ::*      . : :*:******. * :: **:.**

                      A304D
S. aureus     DVFAPNDIVNEELINNDHVDDNLIE-------E---------------  315
S. mutans     EFFANTDEISDDAITLDTEELVLEAIAQKFAREAESLDEMDKTKEQD  347
              :.** .* :.:: *. *  :  *           *
```

**Supplemental Figure 6. Alignment of DgkB.** Alignment of the DgkB protein coding sequence between *Staphylococcus aureus* and *S. mutans* (1542c). Bolded red amino acids in the *S. mutans* sequence signify those residues that are mutated in the RSV3 variants, with the corresponding amino acid substitution found above. Sequences were aligned using the web tool Clustal Omega (https://www.ebi.ac.uk/Tools/msa/clustalo/).

**Figure S7**

**Supplemental Figure 7. Assaying of Δ804 and Δ1576c Phenotypes.** Non-polar

kanamycin antibiotic marker replacement mutants of Δ804 and Δ1576c were compared

to the parental UA159 strain for the following phenotypes: (A) growth in BHI, (B) growth

in BHI with 0.50 µg mL $^{-1}$ CHX, (C) growth in BHI with 0.75 µg mL $^{-1}$ CHX, (D) growth in

BHI with 1.00 µg mL $^{-1}$ CHX, (E) ETEST Strip Assay with Daptomycin, (F) bacteriocin

overlay assay with *S. sanguinis* SK150 as the indicator strain, and (G) plate-based

antagonism assay.