Characterization of Human Dosage-Sensitive Transcription Factor Genes

Zhihua Ni[1,2], Xiao-Yu Zhou[1], Sidra Aslam[1], Deng-Ke Niu[1*]

[1]MOE Key Laboratory for Biodiversity Science and Ecological Engineering and Beijing Key
Laboratory of Gene Resource and Molecular Development, College of Life Sciences, Beijing Normal
University, Beijing, China

[2]College of Life Sciences, Hebei University, Baoding, China

* Correspondence:
Deng-Ke Niu
dkniu@bnu.edu.cn, dengkeniu@hotmail.com

Keywords: Transcription factor, dosage-sensitive, ohnolog, conserved copy number,
haploinsufficiency, disease, nuclear receptor, C2H2-ZF.

ABSTRACT

Copy number changes in protein-coding genes are detrimental if the consequent changes in protein
concentrations disrupt essential cellular functions. The dosage sensitivity of transcription factor (TF)
genes is particularly interesting because their products are essential in regulating the expression of
genetic information. From four recently curated datasets of dosage-sensitive genes (genes with
conserved copy numbers across mammals, ohnologs, and two datasets of haploinsufficient genes),
we compiled a dataset of the most reliable dosage-sensitive (MRDS) genes and a dataset of the most
reliable dosage-insensitive (MRDIS) genes. The MRDS genes were those present in all four datasets,
while the MRDIS genes were those absent from any one of the four datasets and with pLI values <
0.5 in both of the haploinsufficient gene datasets. Enrichment analysis of TF genes among the MRDS
and MRDIS gene datasets showed that TF genes are more likely to be dosage-sensitive than other
genes in the human genome. The nuclear receptor family was the most enriched TF family among the
dosage-sensitive genes. TF families with very few members were also deemed more likely to be
dosage-sensitive than TF families with more members. In addition, we found a certain number of
dosage-insensitive TFs. The most typical were the KRAB domain-containing zinc-finger proteins
(KZFPs). Gene Ontology (GO) enrichment analysis showed that the MRDS TFs were enriched for
many more terms than the MRDIS TFs; however, the proteins interacting with these two groups of
TFs did not show such sharp differences. Furthermore, we found that the MRDIS KZFPs were not
significantly enriched for any GO terms, whereas their interacting proteins were significantly
enriched for thousands of GO terms. Although KZFPs and some other MRDIS TFs likely have lower
functional complexity than MRDS TFs, they were under-annotated in the GO database. We also
examined whether the presence of a repressor domain and an activation domain underlies the
mechanism of dosage sensitivity of TFs. Further characterizations revealed significant differences
between MRDS TFs and MRDIS TFs in the lengths and nucleotide compositions of DNA-binding
sites as well as in expression level, protein size, and selective force.

1   INTRODUCTION

Gene duplication and loss in evolution and gene copy number polymorphisms at the population level have been widely observed in both animals and plants (Innan and Kondrashov, 2010; Schrider and Hahn, 2010; Panchy et al., 2016). The copy number of a particular gene present in a genome is termed the gene dosage. A dosage change in a gene can happen within one locus through inactivation of one or two alleles or among two or more loci through gene duplication and loss of duplicates. Gene dosage sensitivity is the model of the deleterious effects resulting from genomic alteration of gene dosage (Rice and McLysaght, 2017a).

If the protein products of both alleles are required for a normal phenotype, heterozygous loss of function results in an abnormal phenotype. Haploinsufficiency, a measure of intolerance to such heterozygous loss-of-function variations, is a widely studied model of gene dosage sensitivity. Here, we briefly describe two sources of data on human gene haploinsufficiency used in this study. According to the sensitivity to loss-of-function variation, each human gene can be assigned to one of three natural categories: null (in which loss-of-function variation, regardless of whether it is heterozygous or homozygous, is completely tolerated by natural selection), recessive (in which heterozygous loss-of-function variation is tolerated but homozygous loss-of-function (HLOF) variation is not), and haploinsufficient (in which heterozygous loss-of-function variation is not tolerated). Lek et al. (2016) assumed that tolerant genes have the expected amount of truncating variation and took the empirical mean observed/expected rates of truncating variation for recessive disease genes and severe haploinsufficient genes to represent the average outcome of the homozygous and heterozygous intolerant scenarios, respectively; they then built a three-state model and designed a metric, the probability of being loss of function-intolerant (pLI). This metric can be used to evaluate the intolerance to heterozygous protein-truncating variation of each gene. With a cutoff of a $pLI > 0.9$, the researchers identified 3230 haploinsufficient genes by analyzing 60706 human exomes. In addition to previously identified haploinsufficient genes associated with diseases, their dataset included some genes that had not yet been assigned to any human diseases. The low incidence of these genes in human populations indicates that heterozygous loss of function confers some survival or reproductive disadvantage. More recently, Shihab et al. (2017) integrated genomic and evolutionary information from several large databases and predicted the existence of 7841 haploinsufficient genes in the human genome using a machine learning approach called HIPred. This dataset was comparably larger than that of Lek et al. mostly because Shihab et al. used a relaxed cutoff of a $pLI > 0.5$.

As the dosage sensitivity of a gene results from effects related to the ratio of its product to other cellular components, small-scale duplication events involving dosage-sensitive genes are deleterious. Therefore, the duplicability of a gene in evolution relative to all other genes in the same genome can also be used as a measure of dosage sensitivity (Rice and McLysaght, 2017a). Pairs of genes in the same genome originating from small-scale duplication are termed paralogs, while those originating from whole-genome duplications are termed ohnologs (Glover et al., 2016). If a gene has only ohnologs but no paralogs, its copy number changes during evolution are very likely under dosage constraints. Makino and McLysaght (2010) compiled a list of 7294 ohnologs in the human genome and confirmed that these ohnologs are in chromosomal regions with low copy number variations. Later, Rice and McLysaght (2017b) found that the copy numbers of genes in pathogenic copy number-variation regions of the human genome are more conserved across 13 mammalian genomes than those of genes in benign regions. Therefore, the 7014 genes they identified to have the most conserved copy numbers across the 13 mammalian genomes could be regarded as dosage-sensitive genes.

Transcription factors, in a narrow sense, are proteins that control the rate of transcription by binding
to specific DNA sequences (Lambert et al., 2018). Mutations affecting the DNA-binding domains
(DBDs) of TFs disrupt the accurate control of gene expression and cause diseases. It should be noted
that some proteins that regulate gene expression but do not directly bind DNA are also called TFs.
Previous studies have shown that some TFs work in a dose-dependent manner; heterozygous loss-of-
function mutations of TFs are also associated with severe phenotypic disorders (Engelkamp and van
Heyningen, 1996; Seidman and Seidman, 2002). In addition, in some cases, individuals with
heterozygous deletions of particular TFs (e.g., THRB) do not exhibit abnormal phenotypes
(Engelkamp and van Heyningen, 1996; Seidman and Seidman, 2002). A strong majority of genes in
the human genome are dosage insensitive (Makino et al., 2013; Lek et al., 2016; Rice and
McLysaght, 2017a; Rice and McLysaght, 2017b; Shihab et al., 2017). However, considering that the
function of a TF is to ensure the expression of a target gene in the right cell at the right time and in
the right amount in response to intracellular, intercellular or environmental signals, it seems odd for a
TF to exist whose dose reduction or duplication does not cause any phenotypic changes. In 2002,
Seidman and Seidman (2002) surveyed the molecular causes of human syndromes. Among 491 TF
genes, only 27 were confirmed to exhibit phenotypic haploinsufficiency in the syndromic records
available at that time. Members of several TF gene families appear more likely to be sensitive to
copy number changes than members of other families. Are the previously identified dosage-
insensitive TF genes (Engelkamp and van Heyningen, 1996; Seidman and Seidman, 2002) just
exceptions, or is dosage sensitivity limited to certain unique families of TF genes? Recently, Lambert
et al. (2018) curated a comprehensive list of human TFs by manually examining lists of putative TF
from several sources, including previous manual curations, domain searches, the Gene Ontology
(GO) database , and crystal and NMR structure data on proteins in complex with DNA taken from
the Protein Data Bank. This most updated list includes 1639 TFs categorized into 65 families.
Another study in the same year curated 1665 TFs, a very similar number (Hu et al., 2018). Taking
advantage of an unprecedented wealth of data with regard to both the annotation of TF genes in the
human genome and the identification of dosage-sensitive genes, we re-evaluated the dosage
sensitivity of human TF genes. By comparing dosage-sensitive and dosage-insensitive TF genes, we
characterized the mechanistic properties of the sensitive genes, including their DNA quaternary
structures, DNA-binding sites, DBDs, behaviors in protein-protein interactions (PPIs), GO
enrichments, expression patterns, selective forces (nonsynonymous difference [d_N]/synonymous
difference [d_S] values) experienced in evolution, etc.

2 MATERIALS AND METHODS

A list of 1639 human TFs, their classifications into different families, their DNA-binding motifs, and
their quaternary structures were obtained from Lambert et al. (2018). Data on the DBDs of human
TFs also contributed by Lambert et al. (2018) were downloaded from
http://humantfs.ccbr.utoronto.ca/download.php. Data on transactivation domains and transcriptional
repressor domains were retrieved from the UniProt Knowledgebase (UniProtKB, UniProt release
2019_04) (The UniProt Consortium, 2019). Data on Krüppel-associated box (KRAB) domain-
containing zinc-finger proteins (KZFPs) were obtained from Imbeault et al. (2017). Data on PPIs
were obtained from the STRING database (Szklarczyk et al., 2018).

A list of 1317 HLOF-tolerant human genes was obtained from Saleheen et al. (2017). From Lek et al.
(2016), we retrieved the probability of a gene to be tolerant to both heterozygous loss-of-function and
HLOF mutations (pLI and pNull values) of all human genes. With a cutoff of > 0.9 for both metrics,
1226 HLOF-tolerant genes and 3230 haploinsufficient genes were obtained. A larger dataset of
human haploinsufficient genes (7841) was obtained from Shihab et al. (2017). A list of 7294

ohnologs in the human genome was obtained from Makino and McLysaght (2010) and Makino et al.
(2013), and a list of 7014 copy number-conserved genes was obtained from Rice and McLysaght
(2017b). The orthologous relationships between human and mouse genes and their d_N and d_S values
were retrieved from BioMart (Ensembl version GRCh37,
http://grch37.ensembl.org/biomart/martview). Gene expression data were downloaded from the
Human Protein Atlas (Version 18.1, https://www.proteinatlas.org/) (Uhlén et al., 2015), and the
length of each protein-coding sequence was retrieved from Ensembl (version GRCh37,
ftp://ftp.ensembl.org/pub/grch37/update/fasta/homo-sapiens/). Only transcripts with full open reading
frames and multiples of three nucleotides were retained. For the multiple alternative splicing
isoforms, we retained only the longest transcripts. In total, 22810 protein-coding genes have been
annotated in version GRCh37.p13 of the human genome.

In enrichment analyses, chi-square tests (expected value > 5) and Fisher's exact tests (expected value
≤ 5) were used to test whether a particular set of TF genes was significantly over- or
underrepresented in a dataset of dosage-sensitive genes or HLOF-tolerant genes. As tens of chi-
square or Fisher's exact tests were performed per dataset, some of the obtained p values were likely
to be less than 0.05 purely by chance. Therefore, the false discovery rate (FDR)-adjusted p values
were computed using the Benjamini-Hochberg (BH) procedure.

Enrichments and comparisons of GO terms were performed using the R package clusterProfiler with
its default settings (Yu et al., 2012). In this program, adjusted p values were also estimated to prevent
a high FDR during multiple testing.

149   3   RESULTS

According to Lambert et al. (2018), 1639 annotated TF genes are in the human genome. Among
them, 1570 TF genes were categorized into 64 families according to the DBDs they encoded. In
addition, 69 TFs lacking recognizable DBDs were collectively categorized into a family termed
"Unknown". The largest family was the C2H2-ZF family, with 747 members, followed by the
homeodomain (196 members), bHLH (108 members), bZIP (54 members), forkhead (49 members),
nuclear receptor (46 members), HMG/Sox (30 members), ETS (27 members), T-box (17 members),
AT-hook (16 members), Homeodomain+POU (16 members), Myb/SANT (14 members), THAP
finger (12 members), CENPB (11 members), E2F (11 members), BED ZF (10 members), GATA (10
members), and Rel (10 members) families, among others. In addition, we were also interested in
whether TF gene families with very few members have some uniqueness with regard to dosage
sensitivity. For this reason, we defined four additional categories of small TF gene families, each
with ≤ 5, ≤ 7, ≤ 9, or ≤ 11 members. All the TF members within each category were grouped together
in statistical analyses like a gene family.

163   3.1   Enrichment of TF Genes among the Dosage-Sensitive Genes

Using a machine learning approach called HIPred, Shihab et al. (2017) predicted 7841
haploinsufficient genes in the human genome. Among them, 7824 genes have been annotated in
version GRCh37.p13 of the human genome. By analyzing the variations across 60706 human
exomes, Lek et al. (2016) identified 3230 haploinsufficient genes.

Ohnologs are duplicates of genes whose duplication can be tolerated only in whole-genome
duplication events. Their duplicability is limited by their dosage relative to other genes of the same
genome. Using an all-against-all blastp search of human, zebrafish (*Danio rerio*), green spotted
puffer (*Tetraodon nigroviridis*), stickleback (*Gasterosteus aculeatus*), medaka (*Oryzias latipes*),

Japanese puffer (*Takifugu rubripes*) and sea vase (*Ciona intestinalis*, an ascidian) protein sequences,
Makino and McLysaght (2010) identified 7294 ohnologs in the human genome. Among them, 7074
genes have been annotated in the version GRCh37.p13 of the human genome.

In addition, Rice and McLysaght (2017b) identified 7014 human genes whose copy numbers are
conserved across 13 mammalian genomes (*Bos taurus*, *Callithrix jacchus*, *Canis lupus familiaris*,
Equus caballus, *Felis catus*, *Gorilla gorilla*, *Macaca mulatta*, *Mus musculus*, *Oryctolagus cuniculus*,
Ovis aries, *Pan troglodytes*, *Rattus norvegicus* and *Sus scrofa*) and showed evidence that these copy
number-conserved genes are dosage-sensitive.

We noticed that the datasets of dosage-sensitive genes obtained through different methods varied
substantially (Figure 1). The dosage sensitivity of 853 genes was consistently supported by the four
independent studies, so we regarded these genes as the most reliable dosage-sensitive (MRDS) genes
(Supplementary Table 1). The MRDS gene dataset contained 122 TF genes representing 32 TF gene
families. Statistical analysis showed that TF genes were significantly overrepresented in the MRDS
gene dataset (chi-square = 20.1, df = 1, BH-adjusted $p = 3 \times 10^{-4}$). This significant
overrepresentation was observed in nine TF gene families, including the nuclear receptor,
Grainyhead, bHLH, C2H2-ZF;Homeodomain, T-box, AP-2, RFX, Rel, and paired box families (BH-
adjusted $p < 0.05$ for all cases, Table 1). The small TF gene families were also overrepresented in the
MRDS gene dataset regardless of whether they were defined by ≤ 5, ≤ 7, ≤ 9, or ≤ 11 members (BH-
adjusted $p < 0.05$ for all cases, Table 1).

Furthermore, we observed 9459 genes that were not detected to be dosage-sensitive in any of the four
studies on gene dosage sensitivity (Makino and McLysaght, 2010; Lek et al., 2016; Rice and
McLysaght, 2017b; Shihab et al., 2017). To obtain a dataset of the most reliable dosage-insensitive
(MRDIS) genes, we discarded the genes with a pLI value > 0.5 in either the dataset of Shihab et al.
(2017) or the dataset of Lek et al. (2016) from these 9459 genes. In total, 5579 MRDIS genes were
obtained, including 368 TF genes (Supplementary Table 2). Statistical analysis showed that TF genes
were not significantly overrepresented or significantly underrepresented among the MRDIS genes
(chi-square = 1.4, df = 1, BH-adjusted $p = 0.569$). These 368 MRDIS TF genes were distributed in 25
TF families (Table 2). Among the MRDIS TFs, members of the gene families C2H2-ZF and CENPB
were significantly overrepresented, with a greater incidence than expected (BH-adjusted $p < 0.05$ for
both cases). In the Homeodomain gene family, significantly fewer MRDIS genes (15) were observed
than expected (47.9) (chi-square = 17.2, df = 1, BH-adjusted $p = 0.001$). No nuclear receptor genes
were observed in the MRDIS gene dataset, but the expected value by chance was 11.3. Thus, nuclear
receptor genes were significantly underrepresented in the MRDIS gene dataset (chi-square = 11.3,
df=1, BH-adjusted $p = 0.009$). In all other TF gene families, the numbers of TF members were not
significantly different from the expected values (BH-adjusted $p > 0.05$ for all cases). The small TF
gene families were underrepresented in the MRDIS gene dataset regardless of whether they were
defined by ≤ 5, ≤ 7, ≤ 9, or ≤ 11 members (BH-adjusted $p < 0.05$ for all cases, Table 2).

Furthermore, we surveyed the abundance of KZFPs, which form a major subfamily of C2H2-ZF, in
the above two datasets separately. Among the MRDS genes, there were no KZFP genes, although the
expected value by chance was 14 (chi-square = 14.4, df = 1, $p = 1.5 \times 10^{-4}$). Furthermore, the KZFP
genes were significantly overrepresented in the MRDIS gene dataset (observed/expected = 209/94,
chi-square = 43.7, df = 1, $p = 3.8 \times 10^{-11}$).

3.2   MRDS TF Genes Are Enriched for a Variety of GO Terms, but MRDIS TF Genes Are
Enriched for Only Three

GO enrichment analysis showed that the MRDS TFs were significantly enriched for 43 molecular
function terms, 10 cellular component terms and 581 biological process terms (Figure 2 and
Supplementary Table 3-5), consistent with the various roles of the TFs in regulating gene expression.
In sharp contrast, the MRDIS TFs were significantly enriched for only three molecular function
terms: 1) DNA-binding transcription repressor activity, RNA polymerase II-specific; 2) DNA-
binding transcription activator activity, RNA polymerase II-specific; and 3) TF activity, RNA
polymerase II proximal promoter sequence-specific DNA binding (Figure 2 and Supplementary
Table 3-5). For each term, the matched TFs accounted for fewer than 10% of the MRDIS TFs; the
majority of MRDIS genes did not significantly match any GO terms. These three GO terms for which
MRDIS TFs were enriched were also the significant GO terms for the MRDS TFs (Figure 2).

3.3   MRDS TFs Interact with More Proteins than MRDIS TFs

We downloaded human PPI data from the STRING database (Szklarczyk et al., 2018). The most
recent version of STRING, 11.0, contains PPI data for 19566 human proteins, including 121 MRDS
TFs and 356 MRDIS TFs. On average, each MRDS TF was found to interact with 851 proteins,
while each MRDIS TF was found to interact with 263 proteins. Mann-Whitney U tests showed that
the difference was statistically significant ($p < 10^{-6}$). In addition, we observed that 25% of the
proteins interacting with the MRDS TFs were TFs while 17% of the proteins interacting with the
MRDIS TFs were TFs (Mann-Whitney U test, $p < 10^{-6}$).

Next, we performed GO enrichment analysis of the proteins interacting with the MRDS TFs and
MRDIS TFs. Generally, the results were consistent with the commonly believed functions of TFs in
gene expression regulation (Supplementary Table 6-8). Both the proteins interacting with MRDS TFs
and those interacting with MRDIS TFs were enriched for numerous terms indicating gene expression
regulation, such as 0000982 (TF activity, RNA polymerase II proximal promoter sequence-specific
DNA binding), 0007389 (pattern specification process), 0005667 (TF complex). In addition, there
were some differences in the ranks of the enriched terms between the two groups of proteins. Among
the top 20 enriched terms, the two groups shared 15, 11, and 12 common terms in the molecular
function, biological process, and cellular component categories, respectively. However, it is difficult
to draw conclusions on the functional differences between MRDS TFs and MRDIS TFs based on
their interacting proteins.

The GO enrichment analyses of the MRDIS TFs and their interacting proteins indicated entirely
different functional roles. One possible explanation is that the MRDIS TFs are more poorly annotated
in the GO database than their interacting proteins. As the functions of most KZFPs are unknown
(Imbeault et al., 2017), KZFPs are likely under-annotated in the GO database. From the dataset of
Imbeault et al. (2017), we retrieved 399 KZFPs. None of these KZFPs was found in the MRDS TF
dataset, while 209 KZFPs were found in the MRDIS TF dataset. GO enrichment analysis showed that
the MRDIS KZFPs were not significantly enriched for any GO terms (BH-adjusted $p > 0.05$ for all
the terms in the molecular function, biological process, and cellular component categories). In the
STRING database (Szklarczyk et al., 2018), we found PPI records for all 208 MRDIS KZFPs. In
total, the proteins interacting with the 208 MRDIS KZFPs were significantly enriched for 130
molecular function terms, 1299 biological process terms, and 139 cellular component terms (BH-
adjusted $p < 0.05$ for cases, Supplementary Table 9-11). The potential functions of the MRDIS
KZFPs in regulating gene expression were evident from the most enriched GO terms of their
interacting proteins, such as 0005667 (TF complex), 0044798 (nuclear TF complex), 0000790
(nuclear chromatin), 0090575 (RNA polymerase II TF complex), 0000982 (TF activity, RNA
polymerase II proximal promoter sequence-specific DNA binding), 0001228 (DNA-binding

transcription activator activity, RNA polymerase II-specific), 0001077 (proximal promoter DNA-
binding transcription activator activity, RNA polymerase II-specific), 0001227 (DNA-binding
transcription repressor activity, RNA polymerase II-specific), 0001078 (proximal promoter DNA-
binding transcription repressor activity, RNA polymerase II-specific), 0003713 (transcription
coactivator activity), 0003714 (transcription corepressor activity), 0001158 (enhancer sequence-
specific DNA binding), 0035326 (enhancer binding), and 0070491 (repressing TF binding). These
results indicate that some MRDIS TFs are under-annotated in the GO database. Furthermore, the
Mann-Whitney U test showed that MRDIS KZFPs interact with significantly fewer proteins than
other MRDIS TFs (average values: 204 $vs.$ 346, $p < 10^{-6}$), indicating that the under-annotated
MRDIS TFs have lower functional complexity than the well-annotated ones.

3.4   Differences in the DBDs of MRDS TFs and MRDIS TFs

Protein domains are the significant tertiary structures of a protein that are generally believed to have
more functional implications than the rest of the protein chain. A TF gene with two or more functions
is more likely to be sensitive to dosage changes than one with fewer functions because a dosage
constraint on any function will cause the gene to be dosage-sensitive. For this reason, we compared
the number of DBDs between MRDS TFs and MRDIS TFs. The majority of human TFs had only one
DBD in each protein. There were only 43 two-DBD TFs, including seven MRDS TFs and two
MRDIS TFs. Although a significantly percentage of MRDS TFs than MRDIS TFs were two-domain
proteins (chi-square = 4.30, df = 1, $p = 0.038$), the difference in dosage sensitivity between MRDS
TFs and MRDIS TFs could not be attributed to the number of domains.

To test whether the dosage sensitivity of human TFs is achieved through dual functions of both
activator and repressor properties, we searched the keywords "transcription factor," "activator
domain," "repressor domain," "activation domain," and "repression domain" in the UniProtKB
database (The UniProt Consortium, 2019). Among the 122 MRDS TFs, 21 TFs were found to have
activation domains, while eight TFs were found to have repressor domains. Three MRDS TFs
(*HIF1A*, *SP1*, and *TP63*) have both activation domains and repressor domains. Among the 368
MRDIS TFs, three have activation domains, and five have repressor domains. Two MRDIS TFs
(*ZBTB32* and *YY2*) have both activation domains and repressor domains. According to Imbeault et al.
(2017), the KRAB domains of KZFPs can repress transposable elements by recruiting transcriptional
regulators such as TRIM28. Therefore, KRAB domains were also regarded as repressor domains in
the present study. From Imbeault et al. (2017), we retrieved 399 KRAB-containing proteins, 209 of
which were MRDIS TFs. None of these 209 TFs was found to have activation domains. However,
MRDS TFs were found to be more likely to work as activators than MRDIS TFs (17.2% $vs.$ 0.82%,
chi-square = 14.9, $p = 10^{-4}$), whereas MRDIS TFs are more likely to work as repressors (6.56% $vs.$
58.2%, chi-square = 41.1, $p = 1.4 \times 10^{-10}$ if regarding KRAB as a repressor domain; the difference
was not significant if we did not regard KRAB as a repressor domain).

We also compared the sizes of DBDs between MRDS TFs and MRDIS TFs. Among the 77 MRDS
TFs whose domains have been annotated, the average domain size was 118 amino acid residues. In
contrast, the average domain size of the 279 MRDIS TFs was only 52 amino acid residues. Mann-
Whitney U tests showed that the difference was highly significant ($p = 2.1 \times 10^{-6}$). Furthermore, we
compared the sizes of the activation domains and the repressor domains. Regardless of whether the
comparisons were performed within MRDS TFs, within MRDIS TFs, or within a combined dataset
of MRDS TFs and MRDIS TFs, we did not detect any significant difference in either activation
domain size or repressor domain size (Mann-Whitney U test, $p > 0.05$ for all cases).

3.5   Quaternary Structures of MRDS TFs and MRDIS TFs

We surveyed the quaternary structures of TFs using the data of Lambert et al. (2018) and found three
types of proteins: low-specificity DNA-binding proteins, monomers or homomultimers, and obligate
heteromers. The vast majority of the TFs were monomers or homomultimers. In total, 96.7% of
MRDS TFs were monomers or homomultimers, and 97.6% of MRDIS TFs were monomers or
homomultimers. Therefore, the dosage sensitivity of TFs is not related to the type of quaternary
structure.

3.6   Differences in the DNA-binding Sites of MRDS TFs and MRDIS TFs

From Lambert et al. (2018), we obtained the DNA-binding sites of 102 MRDS TFs and 235 MRDIS
TFs. On average, the DNA-binding sites of the MRDS TFs had 12 nucleotides, whereas those of the
MRDIS TFs had 13 nucleotides. Although the difference seems slight, it was statistically significant
(Mann-Whitney U test, $p = 7.4 \times 10^{-5}$, Table 3). Furthermore, we compared the nucleotide
compositions of the DNA-binding sites. The DNA-binding sites of both MRDS TFs and MRDIS TFs
had greater numbers of A than any of the other three nucleotides (Mann-Whitney U test, $p < 10^{-6}$ for
all comparisons). However, the DNA-binding sites of MRDS TFs had significantly fewer A residues
than those of MRDIS TFs (Mann-Whitney U test, $p < 10^{-6}$). Furthermore, the DNA-binding sites of
MRDS TFs had more G residues than those of MRDIS TFs (Mann-Whitney U test, $p = 0.048$). No
significant differences were observed in C and T.

3.7   Expression Patterns of MRDS TFs and MRDIS TFs

One previous study has shown that dosage-sensitive genes are generally expressed at high levels
(Gout et al., 2010; Rice and McLysaght, 2017a). From the Human Protein Atlas (Uhlén et al., 2015),
we retrieved the mRNA expression levels of 119 MRDS TFs and 348 MRDIS TFs in 37 cell
samples. After averaging the mRNA expression levels of each TF among these 37 samples, we
compared the expression levels between the 119 MRDS TFs and 348 MRDIS TFs. Consistent with
previous studies, we found that the mRNA expression levels of MRDS TFs (6.36 ± 0.679) were
significantly higher than those of MRDIS TFs (6.36 ± 0.679 *vs.* 6.26 ± 1.767; Mann-Whitney U test,
$p = 0.001$).

Although mRNA abundance is significantly correlated with protein abundance, it is far from a perfect
parameter to represent or predict protein expression levels (Greenbaum et al., 2003; Gry et al., 2009;
Maier et al., 2009). Therefore, we re-examined the relationship between dosage sensitivity and gene
expression level using protein abundance data. In the Human Protein Atlas database, the expression
level of a protein, if it has ever been studied, is defined as high, medium, low, or not detected in each
studied cell sample. To ensure that expression data existed for all the studied genes in the studied cell
samples, we selected 31 cell samples in which 684 MRDS genes (including 92 MRDS TF genes) and
2736 MRDIS genes (including 150 MRDIS TF genes) had ever been studied. For each gene, we
surveyed its expression level in the 31 cell samples. For example, the expression level of gene
ENSG00000095951, a member of the C2H2-ZF gene family, was high in nine samples, medium in
12 samples, low in eight samples, and not detected in two samples. We first compared the presence
of the MRDS genes and MRDIS genes in each of the four categories. On average, each MRDS gene
had high expression levels in 4.5 samples, while each MRDIS had high expression levels in 3.0
samples. The Mann-Whitney U test showed that the difference was statistically significant ($p < 10^{-6}$;
Table 4). Similarly, at the medium and low levels, MRDS genes were detected in significantly more
samples than MRDIS genes (Mann-Whitney U test, $p < 10^{-6}$ for both cases; Table 4). In the "not

detected" category, MRDS genes were assigned to significantly fewer samples than MRDIS genes
(Mann-Whitney U test, $p < 10^{-6}$; Table 4). However, no significant differences were detected
between MRDS TF genes and MRDIS TF genes in any of the four categories (Mann-Whitney U test,
$p > 0.05$ for all cases). Thus, we found that dosage-sensitive genes in general have higher expression
levels than dosage-insensitive genes, confirming the findings of a previous study. However, our
results suggest that dosage-sensitive TF genes do not have higher expression levels than dosage-
insensitive TF genes.

3.8   Protein Size Differences between MRDS TFs and MRDIS TFs

Rice and McLysaght (2017a) found that ohnologs have much longer coding sequences than genes
duplicated by small-scale duplication. We re-examined this phenomenon using our datasets. In
version GRCh37.p13 of the human genome, we obtained the coding sequences (CDSs) of 5569
(among the 5579) MRDIS genes. The CDSs of the 853 MRDS genes were significantly longer than
those of the MRDIS genes (Mann-Whitney U test, $p < 10^{-6}$), with a difference in average values of
more than 2-fold (2640 $vs.$ 1310, Table 5). The MRDS TF genes and the MRDIS TF genes also
differed significantly in CDS length (Mann-Whitney U test, $p = 10^{-6}$), but there was a smaller
difference in their average values (2189 $vs.$ 1577, Table 5). Furthermore, we noticed that TF genes
had significantly longer CDSs than other protein-coding genes (Mann-Whitney U test, $p < 10^{-6}$,
Table 5).

3.9   MRDS TF Genes Experience Stronger Selective Pressures than MRDIS Genes

Schuster-Böckler et al. (2010) showed that dosage-sensitive genes are under strong selective force. In
this study, we used the d_N/d_S value for each human gene relative to its mouse ortholog as a measure
of the selective pressure of each gene. Valid d_N/d_S values were obtained for 4188 MRDIS genes and
all 853 MRDS genes. Mann-Whitney U tests showed that the MRDS genes experienced significantly
stronger selective pressure than the MRDIS genes ($p < 10^{-6}$, Table 6). The MRDS TF genes and
MRDIS TF genes exhibited the same patterns (Mann-Whitney U test, $p < 10^{-6}$, Table 6). Compared
with other genes in the human genome, TF genes are under stronger selective forces (Mann-Whitney
U test, $p < 10^{-6}$, Table 6).

3.10   HLOF-Tolerant TF Genes Are Not Rarer than Other Human Genes

It has been determined experientially that complete homozygous loss of TF genes rarely occurs in
humans (Engelkamp and van Heyningen, 1996). Here, we used recent datasets of human HLOF-
tolerant genes to examine whether TF genes are sparser than other genes in the human genome. A
series of large-scale sequencing studies have been carried out on healthy human adults to identify
HLOF-tolerant genes (Lim et al., 2014; Sulem et al., 2015; Lek et al., 2016; Narasimhan et al., 2016;
Saleheen et al., 2017; Bartha et al., 2018). We retrieved the data from the two most extensive studies
(Lek et al., 2016; Saleheen et al., 2017). In addition to the abovementioned pLI metric, Lek et al.
(2016) designed another metric, pNull. With a cutoff of a pNull > 0.9, 1226 genes were identified as
extremely loss of function-tolerant from their dataset. In version GRCh37.p13 of the human genome,
22810 protein-coding genes have been annotated; thus, 5.4% of human genes are HLOF-tolerant.
Among the 1639 TF genes, we found 103 genes (6.3%) in this HLOF-tolerant gene dataset. The TF
genes were not significantly over- or underrepresented among the HLOF-tolerant genes (BH-adjusted
p value = 1). However, we found that one TF gene family, C2H2-ZF, was significantly enriched in
HLOF-tolerant genes (observed value = 91, expected value =40, BH-adjusted p value = 0.0006).
Furthermore, we found that KZFPs, a subfamily of C2H2-ZF, had an even higher observed/expected

ratio (81/21, chi-square = 34.8, df = 1, $p = 3.6 \times 10^{-9}$). In a more recent study, Saleheen et al. (2017)
sequenced the protein-coding regions of 10503 adult participants with a high rate of consanguinity
and identified 1317 distinct genes for which loss-of-function mutations in both copies were tolerated,
including 85 TF genes. Statistical analysis did not show significant over- or underrepresentation of
TF genes among the HLOF-tolerant genes (expected value = 104, chi-square = 1.9, BH-adjusted p
value = 1). The same result was observed when the enrichment of each TF gene family (except for
KZFPs, a subfamily of C2H2-ZF) was analyzed separately (BH-adjusted p value > 0.05 for all cases).
The observed number of KZFPs in this HLOF-tolerant gene dataset was more than twice the value
expected by chance (55 vs. 22, chi-square = 14.0, df = 1, $p = 1.9 \times 10^{-4}$).

4   DISCUSSION

TFs are a group of proteins with unique functional roles to ensure the temporally and spatially
accurate expression of the genetic information encoded in the genome. Intuitively, changes in their
gene copy numbers should be hazardous because changes in the concentrations of TFs in cells will
disturb the normal expression patterns of the TF target genes. Consistent with this idea, complete
homozygous loss of TF genes is rarely observed in humans (Engelkamp and van Heyningen, 1996).
Using the two most extensive datasets of human HLOF-tolerant genes (Lek et al., 2016; Saleheen et
al., 2017), we found that TF genes are not significantly underrepresented among HLOF-tolerant
genes. At least with regard to their percentages, TF genes are not more indispensable than other
genes in the human genome. However, a specific TF family, C2H2-ZF, seems to be more dispensable
than other genes in the human genome. In particular, the genes of the large KZFP subfamily, which
accounts for 51% of the C2H2-ZF gene family, are significantly more dispensable than other genes in
the human genome. Imbeault et al. (2017) showed that the majority of KZFPs bind transposable
elements and repress these transposable elements rather than regulating the expression of other
protein-coding genes. With the rapid accumulation and decay of transposable elements during
evolution (Blass et al., 2012; Wallau et al., 2014), the KZFP subfamily has also likely experienced
dynamic gain of new members and loss of obsolete members.

Compared with complete homozygous losses, gain of a new copy or heterozygous loss of one gene
copy in a diploid genome is expected to have a gentler effect. Seidman and Seidman (2002) surveyed
the haploinsufficiency of human TFs using the data available at that time. Deleterious phenotypic
effects resulting from heterozygous losses were confirmed in only 27 of the 491 TF genes. From four
recent datasets on human dosage-sensitive genes, including datasets obtained from both heterozygous
loss studies and gene duplication studies (Makino et al., 2013; Lek et al., 2016; Rice and McLysaght,
2017b; Shihab et al., 2017), we defined a dataset of the most reliable dosage-sensitive genes, the
MRDS genes, by selecting the dosage-sensitive genes shared by all four datasets. As a control, we
also defined a dataset of the most reliable dosage-insensitive genes, the MRDIS genes, by selecting
the genes that were not considered dosage-sensitive in any of the four datasets. Upon surveying the
abundances of the TF genes in these two datasets, we observed some commonalities (Table 1-2).

First, TF genes were more likely to be dosage-sensitive than other genes in the human genome. The
significant enrichment of TF genes in the dosage-insensitive gene datasets was expected given the
roles of TFs in ensuring accurate expression of their target genes.

Second, the nuclear receptor is a very unique TF gene family with regard to dosage sensitivity.
Members of this family were significantly overrepresented in the MRDS gene dataset and absent
from the MRDIS gene dataset. Nuclear receptors can activate target genes after binding nonpolar
ligands such as estrogen, progesterone, retinoic acid, oxysterols, and thyroid hormone. In contrast to

other TFs, nuclear receptors respond directly to extracellular changes by binding ligands that are
diffusible across the plasma membrane (Sladek, 2011). A distinguishing characteristic of steroid
hormones is the dose-response curve manifested in their regulation of gene expression (Szapary et
al., 1999; Simons, 2006). Fixed dosages of the nuclear receptors for steroid hormones might be the
premise of the dose-response curve.

Third, TF gene families with fewer numbers were more likely to be enriched among the dosage-
sensitive genes. This finding was supported by observations of overrepresentation of small-family TF
genes in the MRDS gene dataset and underrepresentation of them in the MRDIS gene dataset. We
did not observe any common characteristics among these small-family TFs. Instead, we suspect that
the dosage sensitivity of these TF genes might act as a selective force against the expansion of these
families and maintain the small number of members during evolution.

Fourth, many TF genes were dosage-insensitive, and the KZFP genes seemed to be the most dosage-
insensitive. Considering the roles of TF genes in the regulation of gene expression, the existence of a
large number of dosage-insensitive TF genes was unexpected. One possible explanation is that some
TFs do not directly regulate the expression levels of other genes but simply contribute, more or less,
to the maintenance of a nuclear environment compatible with the expression of required genes.
Nonfunctional interactions within the accessible portions of the genome have been observed for
many TFs (Fisher et al., 2012; Slattery et al., 2014). Some TFs, such as KZFPs, have been shown to
be involved in the repression of transposable elements (Thomas and Schneider, 2011; Imbeault et al.,
2017; Yang et al., 2017). As there is no purifying selection against the accumulation of degenerative
mutations in repressed transposable elements, most transposable elements quickly lose their
transposition potential during evolution. Dosage changes in TFs that target transposable elements
with low or no transposition potential should not have severe effects. In addition, if large proportions
of TF binding events are nonfunctional, the effects of dosage changes in TF genes on the expression
of target genes should be buffered. Nonfunctional binding (Todeschini et al., 2014) might therefore
underlie the mechanism of dosage insensitivity of some TF genes.

Then, we characterized the dosage-sensitive TFs by comparing the MRDS TFs and the MRDIS TFs.

GO enrichment analysis showed that MRDS TFs were significantly enriched for 634 GO terms,
whereas MRDIS TFs were significantly enriched for only three molecular function terms
(Supplementary Table 3-5), indicating that MRDS TFs have a much greater variety of functions than
MRDIS TFs. Consistent with this result, we found that MRDS TFs interact with many more proteins
than MRDIS TFs. GO enrichment analysis of the proteins interacting with MRDS TFs and MRDIS
TFs, however, showed that both groups of proteins were enriched for numerous terms indicating gene
expression regulation. The KZFP family is a unique TF family; no member of this family fell in the
MRDS category. The MRDIS KZFPs were not significantly enriched for any GO terms, whereas the
proteins found to interact with the MRDIS KZFPs were significantly enriched for thousands of GO
terms (Supplementary Table 9-11). The most enriched GO terms of the proteins interacting with the
MRDIS KZFPs provide evidence that the MRDIS KZFPs are actively involved in regulating gene
expression. In addition, we found that MRDIS KZFPs interact with fewer proteins than other MRDIS
TFs. It might be concluded that KZFPs and perhaps some other MRDIS TFs have lower functional
complexity than MRDS TFs; however, these TFs are under-annotated in the GO database compared
with MRDS TFs.

In plants, the TF WUS provides an insightful example of the mechanism of dosage sensitivity of TFs
(Hofhuis and Heidstra, 2018). WUS can bind different cofactors and activates its target gene CLV3

when its protein is expressed at low levels, while it represses the target gene when its protein is at
high levels. For the WUS gene, both an increase and a decrease in its dosage would disturb its
function. The dosage of the WUS gene has been fixed by its dual functions as an activator and a
repressor. We speculated that these dual functions could be achieved by either a protein complex
containing both activator subunits and repressor subunits or a protein containing both repressor
domains and activation domains. Our survey of the quaternary structures of TFs showed that the vast
majority of both MRDS TFs and MRDIS TFs are monomeric or homomultimeric rather than
heteromeric. Furthermore, we found that the majority of human TFs have a single DBD. For a few
MRDS TFs and a few MRDIS TFs, both activation domains and repressor domains have been
annotated in the UniProtKB database (The UniProt Consortium, 2019). However, the activation or
repression functions of most TFs have not yet been annotated in the database; therefore, more data is
needed before a solid conclusion can be made.

The DNA-binding sites of MRDS TFs were shorter and contained fewer A and slightly more G
residues than those of the MRDIS TFs.

Previous analyses have characterized dosage-sensitive genes as being highly expressed, encoding
large proteins, and being under strong selective force (Gout et al., 2010; Schuster-Böckler et al.,
2010; Rice and McLysaght, 2017a). In this study, we confirmed most of these conclusions by
comparing MRDS genes and MRDIS genes as well as MRDS TF genes and MRDIS TF genes;
however, we found that MRDS TFs do not have significantly higher protein expression levels than
MRDIS TFs. Dosage sensitivity could not be explained by expression burden, at least among the TF
genes. The small d_N/d_S values and the consequently high selective pressures experienced by the
dosage-sensitive genes indicate that functional constraints might be tightly associated with dosage
constraints.

Using the dosage-sensitive gene and dosage-insensitive gene datasets we compiled based on four
recent datasets of dosage-sensitive genes, we found that human TFs were significantly enriched
among the dosage-sensitive genes. In addition, many TFs were found to be dosage-insensitive. The
most dosage-sensitive TF gene family was the nuclear receptor family, while the most dosage-
insensitive TF gene family was the KZFP subfamily. Further characterization of these genes revealed
both intrinsic differences between the dosage-sensitive genes and the dosage-insensitive genes and
the relatively limited knowledge regarding some dosage-insensitive genes, such as KZFPs.

5   Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

6   Author Contributions

DKN conceived the study and wrote the manuscript. ZN retrieved the data from online databases and
performed all the statistical analyses. XYZ and SA each repeated some of the analyses. All authors
read and approved the final manuscript.

7   Funding

This work was supported by the National Natural Science Foundation of China (31671321).

8   Acknowledgments

REFERENCES

Bartha, I., di Iulio, J., Venter, J.C., and Telenti, A. (2018). Human gene essentiality. *Nature Reviews Genetics* 19**,** 51-62.
doi: 10.1038/nrg.2017.75.

Blass, E., Bell, M., and Boissinot, S. (2012). Accumulation and Rapid Decay of Non-LTR Retrotransposons in the
Genome of the Three-Spine Stickleback. *Genome Biology and Evolution* 4(5)**,** 687-702. doi:
10.1093/gbe/evs044.

Engelkamp, D., and van Heyningen, V. (1996). Transcription factors in disease. *Current Opinion in Genetics &*
Development 6(3)**,** 334-342. doi: 10.1016/s0959-437x(96)80011-6.

Fisher, W.W., Li, J.J., Hammonds, A.S., Brown, J.B., Pfeiffer, B.D., Weiszmann, R., et al. (2012). DNA regions bound at
low occupancy by transcription factors do not drive patterned reporter gene expression in *Drosophila*.
Proceedings of the National Academy of Sciences of the United States of America 109(52)**,** 21330-21335. doi:
10.1073/pnas.1209589110.

Glover, N.M., Redestig, H., and Dessimoz, C. (2016). Homoeologs: What are they and how do we infer them? *Trends in*
Plant Science 21(7)**,** 609-621. doi: 10.1016/j.tplants.2016.02.005.

Gout, J.-F., Kahn, D., Duret, L., and Paramecium Post-Genomics Consortium (2010). The relationship among gene
expression, the evolution of gene dosage, and the rate of protein evolution. *PLOS Genetics* 6(5)**,** e1000944. doi:
10.1371/journal.pgen.1000944.

Greenbaum, D., Colangelo, C., Williams, K., and Gerstein, M. (2003). Comparing protein abundance and mRNA
expression levels on a genomic scale. *Genome Biology* 4(9)**,** 117-226. doi: 10.1186/gb-2003-4-9-117.

Gry, M., Rimini, R., Strömberg, S., Asplund, A., Pontén, F., Uhlén, M., et al. (2009). Correlations between RNA and
protein expression profiles in 23 human cell lines. *BMC Genomics* 10(1)**,** 365. doi: 10.1186/1471-2164-10-365.

Hofhuis, H.F., and Heidstra, R. (2018). Transcription factor dosage: more or less sufficient for growth. *Current Opinion*
in Plant Biology 45**,** 50-58. doi: 10.1016/j.pbi.2018.05.008.

Hu, H., Miao, Y.-R., Jia, L.-H., Yu, Q.-Y., Zhang, Q., and Guo, A.-Y. (2018). AnimalTFDB 3.0: a comprehensive
resource for annotation and prediction of animal transcription factors. *Nucleic Acids Research***,** gky822-gky822.
doi: 10.1093/nar/gky822.

Imbeault, M., Helleboid, P.-Y., and Trono, D. (2017). KRAB zinc-finger proteins contribute to the evolution of gene
regulatory networks. *Nature* 543**,** 550-554. doi: 10.1038/nature21683.

Innan, H., and Kondrashov, F. (2010). The evolution of gene duplications: classifying and distinguishing between
models. *Nature Reviews Genetics* 11**,** 97. doi: 10.1038/nrg2689.

Ishikawa, K., Makanae, K., Iwasaki, S., Ingolia, N.T., and Moriya, H. (2017). Post-translational dosage compensation
buffers genetic perturbations to stoichiometry of protein complexes. *PLOS Genetics* 13(1)**,** e1006554. doi:
10.1371/journal.pgen.1006554.

Lambert, S.A., Jolma, A., Campitelli, L.F., Das, P.K., Yin, Y.M., Albu, M., et al. (2018). The human transcription
factors. *Cell* 172(4)**,** 650-665. doi: 10.1016/j.cell.2018.01.029.

Lek, M., Karczewski, K.J., Minikel, E.V., Samocha, K.E., Banks, E., Fennell, T., et al. (2016). Analysis of protein-coding
genetic variation in 60,706 humans. *Nature* 536**,** 285-291. doi: 10.1038/nature19057.

Lim, E.T., Würtz, P., Havulinna, A.S., Palta, P., Tukiainen, T., Rehnström, K., et al. (2014). Distribution and Medical
Impact of Loss-of-Function Variants in the Finnish Founder Population. *PLOS Genetics* 10(7)**,** e1004494. doi:
10.1371/journal.pgen.1004494.

Lundgren, M., Chow, C.-M., Sabbattini, P., Georgiou, A., Minaee, S., and Dillon, N. (2000). Transcription factor dosage
affects changes in higher order chromatin structure associated with activation of a heterochromatic gene. *Cell*
103(5)**,** 733-743. doi: 10.1016/S0092-8674(00)00177-X.

Maier, T., Guell, M., and Serrano, L. (2009). Correlation of mRNA and protein in complex biological samples. *FEBS*
Letters 583(24)**,** 3966-3973. doi: 10.1016/j.febslet.2009.10.036.

Makino, T., and McLysaght, A. (2010). Ohnologs in the human genome are dosage balanced and frequently associated
with disease. *Proc Natl Acad Sci U S A* 107(20)**,** 9270-9274. doi: 10.1073/pnas.0914697107.

Makino, T., McLysaght, A., and Kawata, M. (2013). Genome-wide deserts for copy number variation in vertebrates.
Nature Communications 4**,** 2283. doi: 10.1038/ncomms3283.

Narasimhan, V.M., Hunt, K.A., Mason, D., Baker, C.L., Karczewski, K.J., Barnes, M.R., et al. (2016). Health and
population effects of rare gene knockouts in adult humans with related parents. *Science* 352(6284)**,** 474-477. doi:
10.1126/science.aac8624.

Ni, Z., Zhou, X.-Y., Aslam, S., and Niu, D.-K. (2019). Dosage-sensitivity of human transcription factor genes. *bioRxiv***,**
528554. doi: 10.1101/528554.

Panchy, N., Lehti-Shiu, M., and Shiu, S.H. (2016). Evolution of Gene Duplication in Plants. *Plant Physiology* 171(4)**,**
2294-2316.

Pascual, G., and Glass, C.K. (2006). Nuclear receptors versus inflammation: mechanisms of transrepression. *Trends in*
Endocrinology & Metabolism 17(8)**,** 321-327. doi: 10.1016/j.tem.2006.08.005.

Rice, A.M., and McLysaght, A. (2017a). Dosage-sensitive genes in evolution and disease. *BMC Biology* 15**,** 10. doi:
10.1186/s12915-017-0418-y.

Rice, A.M., and McLysaght, A. (2017b). Dosage sensitivity is a major determinant of human copy number variant
pathogenicity. *Nature Communications* 8**,** 14366. doi: 10.1038/ncomms14366.

Saleheen, D., Natarajan, P., Armean, I.M., Zhao, W., Rasheed, A., Khetarpal, S.A., et al. (2017). Human knockouts and
phenotypic analysis in a cohort with a high rate of consanguinity. *Nature* 544(7649)**,** 235-239.

Schnable, J., Pedersen, B., Subramaniam, S., and Freeling, M. (2011). Dose–sensitivity, conserved non-coding sequences,
and duplicate gene retention through multiple tetraploidies in the grasses. *Frontiers in Plant Science* 2(2). doi:
10.3389/fpls.2011.00002.

Schrider, D.R., and Hahn, M.W. (2010). Gene copy-number polymorphism in nature. *Proceedings of the Royal Society B-*
Biological Sciences 277(1698)**,** 3213-3221. doi: 10.1098/rspb.2010.1180.

Schuster-Böckler, B., Conrad, D., and Bateman, A. (2010). Dosage sensitivity shapes the evolution of copy-number
varied regions. *PLOS ONE* 5(3)**,** e9474. doi: 10.1371/journal.pone.0009474.

Seidman, J.G., and Seidman, C. (2002). Transcription factor haploinsufficiency: when half a loaf is not enough. *Journal*
of Clinical Investigation 109(4)**,** 451-455. doi: 10.1172/JCI15043.

Shihab, H.A., Rogers, M.F., Campbell, C., and Gaunt, T.R. (2017). HIPred: an integrative approach to predicting
haploinsufficient genes. *Bioinformatics* 33(12)**,** 1751-1757. doi: 10.1093/bioinformatics/btx028.

Simons, S.S. (2006). How much is enough? Modulation of dose-response curve for steroid receptor-regrulated gene
expression by changing concentrations of transcription factor. *Current Topics in Medicinal Chemistry* 6(3)**,** 271-
285. doi: 10.2174/156802606776173465.

Sladek, F.M. (2011). What are nuclear receptor ligands? *Molecular and Cellular Endocrinology* 334(1)**,** 3-13. doi:
10.1016/j.mce.2010.06.018.

Slattery, M., Zhou, T.Y., Yang, L., Machado, A.C.D., Gordan, R., and Rohs, R. (2014). Absence of a simple code: how
transcription factors read the genome. *Trends in Biochemical Sciences* 39(9)**,** 381-399. doi:
10.1016/j.tibs.2014.07.002.

Springer, M., Weissman, J.S., and Kirschner, M.W. (2010). A general lack of compensation for gene dosage in yeast.
Molecular Systems Biology 6**,** 368. doi: 10.1038/msb.2010.19.

Sulem, P., Helgason, H., Oddson, A., Stefansson, H., Gudjonsson, S.A., Zink, F., et al. (2015). Identification of a large set
of rare complete human knockouts. *Nature Genetics* 47(5)**,** 448-U430. doi: 10.1038/ng.3243.

Szapary, D., Huang, Y., and Simons, J.S.S. (1999). Opposing effects of corepressor and coactivators in determining the
dose-response curve of agonists, and residual agonist activity of antagonists, for glucocorticoid receptor-
regulated gene expression. *Molecular Endocrinology* 13(12)**,** 2108-2121. doi: 10.1210/mend.13.12.0384.

Szklarczyk, D., Gable, A.L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., et al. (2018). STRING v11: protein–
protein association networks with increased coverage, supporting functional discovery in genome-wide
experimental datasets. *Nucleic Acids Research* 47(D1)**,** D607-D613. doi: 10.1093/nar/gky1131.

The UniProt Consortium (2019). UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Research* 47(D1)**,** D506-
D515. doi: 10.1093/nar/gky1049.

Thomas, J.H., and Schneider, S. (2011). Coevolution of retroelements and tandem zinc finger genes. *Genome Research*
21(11)**,** 1800-1812. doi: 10.1101/gr.121749.111.

Todeschini, A.-L., Georges, A., and Veitia, R.A. (2014). Transcription factors: specific DNA binding and specific gene
regulation. *Trends in Genetics* 30(6)**,** 211-219. doi: 10.1016/j.tig.2014.04.002.

Torres, E.M., Springer, M., and Amon, A. (2016). No current evidence for widespread dosage compensation in *S.*
cerevisiae. *Elife* 5**,** e10996. doi: 10.7554/eLife.10996.

Uhlén, M., Fagerberg, L., Hallström, B.M., Lindskog, C., Oksvold, P., Mardinoglu, A., et al. (2015). Tissue-based map of
the human proteome. *Science* 347(6220)**,** 1260419. doi: 10.1126/science.1260419.

Wallau, G.L., Capy, P., Loreto, E., and Hua-Van, A. (2014). Genomic landscape and evolutionary dynamics of mariner
transposable elements within the Drosophila genus. *BMC Genomics* 15(1)**,** 727. doi: 10.1186/1471-2164-15-727.

Yang, P., Wang, Y., and Macfarlan, T.S. (2017). The role of KRAB-ZFPs in transposable element repression and
mammalian evolution. *Trends in Genetics* 33(11)**,** 871-881. doi: 10.1016/j.tig.2017.08.006.

Yu, G., Wang, L.-G., Han, Y., and He, Q.-Y. (2012). clusterProfiler: an R Package for Comparing Biological Themes
Among Gene Clusters. *OMICS: A Journal of Integrative Biology* 16(5)**,** 284-287. doi: 10.1089/omi.2011.0118.

Table 1. TF genes in the most reliable dosage-sensitive (MRDS) gene dataset.

TF family	Total number in human genome	Observed number of MRDS TFs	Expected number in this dataset	BH-adjusted p
Nuclear receptor	46	20	1.72	1.5×10^{-14}
Grainyhead	6	4	0.22	6.0×10^{-4}
C2H2-ZF;Homeodomain	4	3	0.15	0.002
T-box	17	5	0.64	0.003
AP-2	5	3	0.19	0.004
bHLH	108	12	4.04	0.005
RFX	8	3	0.30	0.016
Rel	10	3	0.37	0.030
Paired box	4	2	0.15	0.040
All TFs	1639	122	61.29	2.6×10^{-4}
Small-family TFs				
$\leq$ 5 members	87	15	3.25	0.032
$\leq$ 7 members	126	23	4.71	0.004
$\leq$ 9 members	186	31	6.96	0.001
$\leq$ 11 members	238	37	8.90	5.9×10^{-4}

The chi-square test (expected value > 5) and Fisher's exact test (expected value ≤ 5) were used to test
the overrepresentation or underrepresentation of the TF genes in this dataset. The Benjamini-
Hochberg (BH) procedure was used to compute the false discovery rate-adjusted p values. The
MRDS genes were defined as the common dosage-sensitive genes among the four datasets obtained
by Shihab et al. (2017), Lek et al. (2016), Makino et al. (2013) and Rice and McLysaght (2017b).

Table 2. TF genes in the most reliable dosage-insensitive (MRDIS) gene dataset.

TF family	Total number in human genome	Observed number of MRDS TFs	Expected number in this dataset	BH-adjusted p
C2H2-ZF	747	281	183	4×10^{-4}
Homeodomain	196	15	48	0.001
Nuclear receptor	46	0	11	0.009
CENPB	11	8	2.7	0.010
All TFs	1639	368	401	0.569
Small-family TFs				
≤ 5 members	87	6	21	0.030
≤ 7 members	126	8	31	0.004
≤ 9 members	186	15	45	0.002
≤ 11 members	238	27	58	0.009

The chi-square test (expected value > 5) and Fisher's exact test (expected value ≤ 5) were used to test
the overrepresentation or underrepresentation of the TF genes in this dataset. The Benjamini-
Hochberg (BH) procedure was used to compute the false discovery rate-adjusted p values. The
MRDIS genes are the genes with pLI values < 0.5 in either the dataset of Shihab et al. (2017) or the
dataset of Lek et al. (2016) that were not considered as dosage-sensitive genes in any of the four
datasets obtained by Shihab et al. (2017), Lek et al. (2016), Makino et al. (2013) and Rice and
McLysaght (2017b).

Table 3. Differences in the DNA-binding sites of the MRDS TFs and MRDIS TFs.

	MRDS TFs (mean ± SD)	MRDIS TFs (mean ± SD)	p
Length (nucleotides)	12.0 ± 3.84	12.7 ± 4.60	7.4×10^{-5}
A (%)	28.2 ± 10.5	30.1 ± 10.7	$< 10^{-6}$
C (%)	24.1 ± 13.3	23.5 ± 12.0	0.899
G (%)	24.3 ± 10.9	23.3 ± 10.1	0.047
T (%)	23.4 ± 10.3	23.1 ± 9.58	0.671

652 The p values were calculated using Mann-Whitney U tests.

654 Table 4. Differences in the protein expression patterns of the MRDS TFs and MRDIS TFs.

	MRDS genes	MRDIS genes	p	MRDS TFs	MRDIS TFs	p
Number of genes	684	2736		92	150	
Number of cell samples in which the proteins are						
at high levels	4.5 ± 6.6	3.1 ± 4.8	$< 10^{-6}$	4.6 ± 6.6	3.8 ± 5.4	0.520
at medium levels	9.2 ± 6.6	7.6 ± 6.7	$< 10^{-6}$	9.3 ± 7.2	9.8 ± 6.2	0.416
at low levels	5.9 ± 4.3	4.8 ± 4.2	$< 10^{-6}$	5.4 ± 4.3	6.2 ± 4.4	0.116
not detected	11.3 ± 9.9	15.5 ± 11.1	$< 10^{-6}$	11.8 ± 10.8	11.1 ± 9.5	0.960

655   In total, 31 cell samples were studied for each gene. The MRDS genes and the MRDIS genes are
656   defined in Tables 1 and 2 as well as in the main text. The p values were calculated using Mann-
657   Whitney U tests.

659   Table 5. Comparison of coding sequence lengths.

	Number of genes	Mean $\pm$ SD (bp)	p
MRDS genes	853	2640 ± 1783	$< 10^{-6}$
MRDIS genes	5569	1310 ± 1956	
MRDS TFs	122	2189 ± 1444	10^{-6}
MRDIS TFs	368	1577 ± 698	
TFs	1608	1777 ± 1205	$< 10^{-6}$
Other genes	19634	1710 ± 1826	

660   The MRDS genes and the MRDIS genes are defined in Tables 1 and 2 as well as in the main text.
661   The p values were calculated using Mann-Whitney U tests.

663   Table 6. Comparison of selective pressures.

	Number of genes	$d_N/d_S \pm$ SD	p
MRDS genes	853	0.062 ± 0.057	$< 10^{-6}$
MRDIS genes	4188	0.257 ± 1.536	
MRDS TFs	122	0.066 ± 0.054	$< 10^{-6}$
MRDIS Fs	226	0.256 ± 0.217	
TFs	1374	0.134 ± 0.141	$< 10^{-6}$
Other genes	16804	0.161 ± 0.776	

The MRDS genes and the MRDIS genes are defined in Tables 1 and 2 as well as in the main text.
The p values were calculated using Mann-Whitney U tests.

Figure Legends

Figure 1. Venn diagram displaying the variations among dosage-sensitive gene datasets obtained
through different methods. The 853 dosage-sensitive genes shared by the four datasets obtained by
Shihab et al. (2017), Lek et al. (2016), Makino et al. (2013) and Rice and McLysaght (2017b) were

regarded as the most reliable dosage-sensitive (MRDS) genes. To obtain a dataset of the most reliable
dosage-insensitive (MRDIS) genes, the 9459 genes that were absent from any of the four datasets
were further filtered by discarding the genes with a pLI value > 0.5 in either the dataset of Shihab et
al. (2017) or the dataset of Lek et al. (2016). In total, 5579 MRDIS genes were obtained.

Figure 2. Clear differences in the Gene Ontology enrichment between MRDS TFs and MRDIS TFs.
Due to space limitations, only the most significant terms of the MRDS TFs are displayed. The color
of each circle represents the significance of the enrichment adjusted using the Benjamini-Hochberg
(BH) procedure, and the size of each circle represents the percentage of genes associated with that
term.

Figure 1

Figure 2