CO$_2$ conversion to methane and biomass in obligate methylotrophic methanogens in

marine sediments

Xiuran Yin[1,2,3]*, Weichao Wu[2,4]*‡, Mara Maeke[1,3], Tim Richter-Heitmann[1], Ajinkya C.

Kulkarni[1,2,3], Oluwatobi E. Oni[1,2], Jenny Wendt[2,4], Marcus Elvert[2,4] and Michael W. Friedrich[1,2]

[1]Microbial Ecophysiology Group, Faculty of Biology/Chemistry, University of Bremen, Bremen,

Germany

[2]MARUM - Center for Marine Environmental Sciences, Bremen, Germany

[3]International Max-Planck Research School for Marine Microbiology, Max Planck Institute for Marine

Microbiology, Bremen, Germany

[4]Department of Geosciences, University of Bremen, Bremen, Germany

Running title: Methane formation from CO$_2$ in *Methanococcoides*

Correspondence:

Michael W. Friedrich,

Microbial Ecophysiology Group, Faculty of Biology/Chemistry, University of Bremen, PO

Box 33 04 40, D-28334 Bremen, Germany

Email: michael.friedrich@uni-bremen.de

* These authors contributed equally to this work.

‡ Current address: Department of Biogeochemistry of Agroecosystems, University of Goettingen,

Goettingen, Germany

Abstract

Methyl substrates are important compounds for methanogenesis in marine sediments but diversity and carbon utilization by methylotrophic methanogenic archaea have not been clarified. Here, we demonstrate that RNA-stable isotope probing (SIP) requires [13]C-labeled bicarbonate as co-substrate for identification of methylotrophic methanogens in sediment samples of the Helgoland mud area, North Sea. Using lipid-SIP, we found that methylotrophic methanogens incorporate 60 to 86% of dissolved inorganic carbon (DIC) into lipids, and thus considerably more than what can be predicted from known metabolic pathways (~40% contribution). In slurry experiments amended with the marine methylotroph *Methanococcoides methylutens*, up to 12% of methane was produced from CO_2, indicating that CO_2-dependent methanogenesis is an alternative methanogenic pathway and suggesting that obligate methylotrophic methanogens grow in fact mixotrophically on methyl compounds and DIC. Thus, the observed high DIC incorporation into lipds is likely linked to CO_2-dependent methanogenesis, which was triggered when methane production rates were low. Since methylotrophic methanogenesis rates are much lower in marine sediments than under optimal conditions in pure culture, CO_2 conversion to methane is an important but previously overlooked methanogenic process in sediments for methylotrophic methanogens.

Introduction

Methanogenesis is the terminal step of organic matter mineralization in marine sediments [1]. There are three main pathways producing methane, i.e., hydrogenotrophic (H_2/CO_2), acetoclastic (acetate) and methylotrophic (e.g., methanol, methylamine) methanogenesis [2] with the former two pathways considered dominant [3]. However, the importance of methylated compounds for methanogenesis in marine sediments has been acknowledged in recent years. Geochemical profiles and molecular analysis have shown that methylotrophic methanogenesis is the most significant pathway for methane formation in hypersaline sediments [4,5] and in the sulfate reduction zone (SRZ) in marine environments [6,7] where methanol concentration of up to 69 µM had been measured [8,9]. Especially in the SRZ, methylated compounds are regarded as non-competitive substrates for methanogenesis, since sulfate reducing microorganisms apparently do not compete with methanogens for these compounds [10]. In sediments of the Helgoland mud area, specifically, high relative abundances of potential methylotrophic methanogens were observed [11] of which many are unknown. The potential for methylotrophic methanogenesis was recently even predicted from two metagenome-assembled genomes of uncultivated Bathyarchaeota, assembled from a shotgun metagenome [12].

The formation of methane via the three main pathways in methanogenic archaea has been studied intensively [13,14,15]. Much less is known regarding assimilation of carbon into biomass under *in situ* conditions and to which extend different carbon sources in the environment are utilized. Discrepancies between predicted pathways known and the actual carbon metabolism measured appear to be based on different cellular functions of carbon dissimilation and assimilation, originate from reaction equilibria operative, intermediate carbon cross utilization as well as interplay between different microbial communities [14, 16,17,18,19]. For example,

mixotrophically growing cultures of *Methanosarcina barkeri* form their biomass equally from

methanol and CO_2, however, almost all the methane is formed from methanol rather than from

CO_2 [20] since methanol is disproportionated to methane and CO_2 according to the following

reaction:

$4 CH_3OH \rightarrow 3 CH_4 + CO_2 + 2 H_2O$ (1)

But apart from such culture studies using the nutritionally versatile *Methanosarcina barkeri*,

which is capable of hydrogenotrophic, acetoclastic and methylotrophic methanogenesis, the

respective contribution of CO_2 and methylated carbon substrates to biomass formation during

methylotrophic methanogenesis, especially for "obligate" methylotrophic methanogens, in

natural sediments has not been studied to date.

Nucleic acids (RNA ~20%, DNA ~3%, of dry biomass, respectively), lipids (7–9%) and proteins

(50–55%) are crucial cell components in living microorganisms [21], and thus, are suitable

markers of carbon assimilation. In order to characterize carbon assimilation capabilities, stable

isotope probing (SIP) techniques exist, among which RNA-SIP is very powerful for identifying

active microorganisms based on separating [13]C-labeled from unlabeled RNA using isopycnic

centrifugation [22,23]. In combination with downstream sequencing analysis, RNA-SIP provides

high phylogenetic resolution in detecting transcriptionally active microbes [24,25] but is limited

in its sensitivity by requiring more than 10% of [13]C incorporation into RNA molecules for

sufficient density gradient separation [26]. To date, a number of SIP studies successfully

detected methylotrophic bacteria [27,28,29] but the detection of methylotrophic methanogens by

RNA-SIP with [13]C labeled methyl compounds might be hampered by mixotrophic growth [20],

i.e., simultaneous assimilation of carbon from methylated compounds and CO_2.

In contrast to RNA-SIP, lipid-SIP has a lower phylogenetic resolution, but can detect very

sensitively δ^{13}C-values in lipid derivatives by gas chromatography combustion isotope ratio mass

spectrometry (GC-IRMS), thereby facilitating quantitative determination of small amounts of

assimilated carbon [30,31].

In this study, we aimed to identify methylotrophic methanogens by RNA-SIP and elucidate

carbon assimilation patterns in marine sediments. We hypothesized that the large pool of ambient

dissolved inorganic carbon (DIC) in sediments [6] alters carbon utilization patterns in

methylotrophic methanogens compared to pure cultures. To address these questions, we tracked

carbon dissimilation into methane and quantified assimilation into lipids by lipid-SIP in slurry

incubations and pure cultures. We found an unexpectedly high degree of methanogenesis from

DIC by previously considered obligate methylotrophic methanogens, i.e., using only methyl

groups for methane formation. This mixotrophic methanogenesis might be the basis for our

observation that more inorganic carbon was assimilated into biomass than could be expected

from known pathways.

Materials and Methods

Sediment incubation setup for SIP

Sediment was collected from the Helgoland mud area (54°05.23'N, 007°58.04'E) by gravity

coring in 2015 during the RV HEINCKE cruise HE443. The geochemical profiles were

previously described [11]. Sediments of the SRZ (16–41 cm) and MZ (238–263 cm) from

gravity core HE443/077-1 were selected for incubations. 50 mL of 1:4 (w/v) slurries were

prepared anoxically by mixing sediments with sterilized artificial sea water without sulfate (26.4

g L^{-1} NaCl, 11.2 g L^{-1} MgCl$_2$·6H$_2$O, 1.5 g L^{-1} CaCl$_2$·2H$_2$O and 0.7 g L^{-1} KCl). Slurries were

dispensed into sterile 120-mL serum bottles and sealed with butyl rubber stoppers. Residual

oxygen was removed by exchanging bottle headspace 3 times with N_2 gas. A 10-day pre-

incubation was performed, followed by applying vacuum (3 min at 100 mbar) to remove most of

the headspace CO_2. Triplicate incubations were conducted by supplementing 1 mM [13]C-labeled

methanol and unlabeled 10 mM sodium bicarbonate, or 1 mM unlabeled methanol and 10 mM

[13]C-labeled sodium bicarbonate ([13]C-labeled substrates provided by Cambridge Isotope

Laboratories, Tewksbury, Massachusetts, USA) at 10 °C.

Pure culture setup

The carbon assimilation patterns were compared between SIP sediment incubations and the

obligate methylotrophic methanogen, *Methanococcoides methylutens*. *M. methylutens* strain

MM1 (DSM 16625) was obtained from the German Collection of Microorganisms and Cell

Cultures (DSMZ, Braunschweig, Germany). Initial cultivation was performed using Medium 280

according to DSMZ protocols. After several transfers of the culture in anoxic marine Widdel

medium [32], 5% of the culture were inoculated into fresh Widdel medium supplemented with

30 mM methanol, trace element solution SL 10 [33], and 50 mM sodium bicarbonate (i.e., DIC)

with carbon sources containing 5% of [13]C-label. Pure cultures were grown at 30 °C. Cells were

harvested after methane formation had ceased by centrifugation and used for lipid-SIP analysis.

Slurry incubations inoculated with *M. methylutens*

To test methanogenesis from CO_2, a series of incubations were performed with *M. methylutens* in

autoclaved (n=3) sediment slurry from the SRZ with different amendments of electron donor

(H_2), electron shuttles (humic acid; anthraquinone-2,6-disulfonic acid - AQDS), and electron

acceptors/electron conductors (hematite, α-Fe_2O_3; magnetite, Fe_3O_4; Lanxess, Germany).

Incubations were separately prepared with 50% H_2 in headspace, 100 μM AQDS, 30 mM

magnetite, 30 mM hematite and 500 mg L^{-1} humic acid (Sigma-Aldrich, Steinheim, Germany). 5%

of the culture was inoculated into these setups each, and amended with 20 mM unlabeled

methanol and 10% of ^{13}C-labeled DIC for measuring carbon partitioning into methane. The

control incubation comprised autoclaved slurry, 50% H_2 in headspace and 20 mM methanol

without addition of *M. methylutens*. All experiments were setup with a total volume of 50 mL in

120-mL serum bottles sealed with butyl rubber stoppers, and incubated at 30 °C.

Gas analysis

The concentration of methane in the headspace of bottles was measured by gas chromatography

as previously described [34]. Headspace H_2 in incubation bottles was determined with a

reduction gas detector (Trace Analytical, Menlo Park, California, USA). 1 mL was used for

injection. The parameters were as follows: carrier gas (nitrogen) 50 mL min^{-1}, injector

temperature 110 °C, detector 230 °C, column (Porapak Q 80/100) 40 °C.

The $\delta^{13}C$ values of methane and CO_2 in the headspace as well as of DIC in the slurries were

determined using a Thermo Finnigan Trace GC connected to a DELTA Plus XP IRMS (Thermo

Scientific, Bremen, Germany) as described previously [35]. Prior to analyses of $\delta^{13}C$-DIC, DIC

in 1 mL of slurry supernatant was converted to CO_2 by adding 100 μL phosphoric acid (85%,

H_3PO_4) overnight at room temperature.

Nucleic acids extraction, quantification and DNase treatment

The nucleic acids were extracted according to Lueders et al. [36]. Briefly, 2 mL of wet sediment

without supernatant was used for cell lysis by bead beating, nucleic acid purification by phenol-

chloroform-isoamyl alcohol extraction and precipitation with polyethylene glycol. For the RNA

extract, DNA was removed by using the RQ1 DNase kit (Promega, Madison, Wisconsin, USA).

DNA and RNA were quantified fluorimetrically using Quant-iT PicoGreen and Quant-iT

RiboGreen (both Invitrogen, Eugene, Oregon, USA), respectively.

Isopycnic centrifugation, gradient fractionation and reverse transcription

Isopycnic centrifugation and gradient fractionation were performed according to the previously

described method with modifications [36]. In brief, 600 to 800 ng RNA from each sample was

loaded with 240 μL formamide, 6 mL cesium trifluoroacetate solution (CsTFA, GE Healthcare,

Buckinghamshire, UK) and gradient buffer solution. The density of the centrifugation medium

was adjusted to ~1.80 g mL^{-1} based on refractive index. RNA was density separated by

centrifugation at 124 000 g at 20 °C for 65 h using an Optima L-90 XP ultracentrifuge (Beckman

Coulter, Brea, California, USA). 13 fractions were obtained from each sample, and RNA in each

fraction was precipitated with 1 volume of isopropanol. RNA was quantified and reverse

transcription was conducted using the high capacity cDNA reverse transcription kit (Applied

Biosystems, Foster City, California, USA).

Quantitative PCR (qPCR)

Archaeal 16S rRNA and *mcrA* genes were quantified using primer sets 806F/912R and ME2

mod/ME3´Fs 1011 (Table S1), respectively, on a Step One Plus qPCR thermocycler (Applied

Biosystems, Foster City, USA); *mcrA* encodes the alpha subunit of methyl coenzyme M

reductase, a key enzyme of methanogenic and methanotrophic archaea [37]. Standard curves

were based on the 16S rRNA gene of *M. barkeri* and the *mcrA* gene clone A4-67 for archaea and

methanogens, respectively. Each 20 μL reaction mixture consisted of 10 μL of MESA Blue

qPCR Master Mix (Eurogentec, Seraing, Belgium), 300 nM primers (600 nM for *mcrA* genes),

0.2 mg mL^{-1} bovine serum albumin (Roche, Mannheim, Germany), 1 ng DNA templates or 2 μL

of cDNA samples. The qPCR protocol comprised an initial denaturation for 5 min at 95 °C and

40 cycles amplification (95 °C for 30 sec, 58 °C for 30 sec and 72 °C for 40 sec). The detection

thresholds were 100–1 000 gene copies with an efficiency of 90–110%.

Sequencing and bioinformatics analysis

For community composition analysis of single labeling incubations (one substrate labeled, the

other unlabeled), "heavy" (1.803–1.823 g mL^{-1}, combination of fraction 3, 4, and 5) and "light"

(1.777–1.780 g mL^{-1}, fraction 11) fractions of RNA-SIP samples were selected according to

CsTFA buoyant densities. PCR targeting the V4 region of archaeal 16S rRNA genes was

conducted with KAPA HiFi HotStart PCR kit (KAPA Biosystems, Cape Town, South Africa)

and barcoded primer sets Arc519F/806R (Table S1). Thermocycling was performed as follows:

95 °C for 3 min; 35 cycles at 98 °C for 20 sec, 61 °C for 15 sec and 72 °C for 15 sec; 72 °C for 1

min. Library construction and sequence read processing were described as previously [34].

Lipid analysis

Total lipids were extracted from ~4 g of freeze-dried sediment samples from single labeling

incubations (one substrate labeled, the other unlabeled) using a modified Bligh-Dyer protocol

[38]. Intact polar archaeal ether lipids were purified by preparative high performance liquid

chromatography with fraction collection according to the method by Zhu et al. [39]. The intact

archaeal lipid fraction was converted to hydrocarbon derivatives including phytane, phytenes,

biphytane, and biphytanes containing cycloalkyl rings (Figure 3C, Figure S1) according to the

method by Liu et al. [40]. Archaeal lipid derivatives were quantified by GC-FID (Thermo

Scientific, Bremen Germany) and the carbon isotope composition was measured via GC-IRMS

on a Thermo Finnigan Trace GC connected with a DELTA V Plus IRMS system via GC Isolink

(Thermo Scientific, Bremen Germany). The detailed chromatographic and mass spectrometric

parameters were described by Kellermann et al. [41].

$\delta^{13}C$ calculation

The proportion of methane from DIC (f_{DIC/CH_4}) was calculated based on the fractional

abundance of ^{13}C (^{13}F) of methane, methanol (MeOH) and DIC in the incubation with ^{13}C-DIC

and MeOH. According to a two-end member model, DIC and MeOH are two main carbon

sources for methane production expressed as follows:

$f_{DIC/CH_4}\,^{13}F_{DIC} + \left(1 - f_{DIC/CH_4}\right)^{13}F_{MeOH} = {}^{13}F_{CH_4}$ (Eq.1)

$f_{DIC/CH_4} = \dfrac{^{13}F_{CH_4} - {}^{13}F_{MeOH}}{^{13}F_{DIC} - {}^{13}F_{MeOH}} \times 100\%$ (Eq.2)

where ^{13}F is obtained from the δ notation according to F = R/(1+R) and R = (δ/1000+1) *

0.011180 [42]. $^{13}F_{CH_4}$ and $^{13}F_{DIC}$ were the fractional ^{13}C abundance of methane and DIC at

harvest time, and $^{13}F_{MeOH}$ that of MeOH in the medium at the start.

^{13}C label incorporation ratios from MeOH or DIC in single labeling experiments were calculated

from the ^{13}C abundance increase relative to the ^{13}C label strength via Eq. 3 and Eq. 4. X_{MeOH} and

X_{DIC} signify the ^{13}C incorporation ratio from MeOH and DIC, respectively. $^{13}F_{t_{end}}$ and $^{13}F_{t_0}$ are

the ^{13}C fractional abundance of lipids harvested at t_{end} and t_0.

$X_{MeOH} = \dfrac{^{13}F_{t_{end}} - {}^{13}F_{t_0}}{^{13}F_{MeOH}}$ (Eq.3)

$X_{DIC} = \dfrac{^{13}F_{t_{end}} - {}^{13}F_{t_0}}{^{13}F_{DIC}}$ (Eq.4)

Given that the single labeling incubations were conducted with the same treatment, i.e., 1 mM

methanol and 10 mM DIC, the relative proportion of DIC for lipids biosynthesis ($f_{DIC/lipid}$) was

estimated from the ^{13}C incorporation ratios (X_{MeOH} and X_{DIC}) in these single labeling

incubations as follow:

$$f_{DIC/lipid} = \frac{X_{DIC}}{X_{DIC}+X_{Me}}$$ (Eq.5)

Results

Methanogenesis and gene copy numbers in ^{13}C-amended sediment slurry incubations

In order to examine carbon labeling into RNA and lipids of methylotrophic methanogens in

anoxic marine environments, sediment slurries from two depths amended with or without ^{13}C-

methanol (1 mM) and ^{13}C-DIC (10 mM) were incubated at 10 °C. Samples from SRZ and MZ

incubations showed methanogenesis started after 20 and 10 days, respectively (Figure 1A).

Amended ^{13}C-DIC was diluted into the sediment endogenous DIC pool to about 70–84%, which

was more obvious in samples from MZ than SRZ (Table 1). In SRZ sediment incubations with

^{13}C-DIC and unlabeled methanol, up to 10.3% of methane originated from ^{13}C-DIC (Table 1).

The dynamics of the archaeal communities in all incubations was tracked by qPCR of archaeal

16S rRNA genes and *mcrA* genes after methanogenesis ceased (Figure 1B). Archaeal and *mcrA*

gene copy numbers increased strongly by 10–14 and 19–30 times for all incubations,

respectively (Figure 1B).

Carbon assimilation into RNA and identification of metabolically active archaea

In preliminary sediment incubations, SIP experiments with ^{13}C-methanol had shown that RNA

could not be labeled to a sufficiently high extent to become detectable in heavy gradient fractions

(e.g., >1.803 g mL^{-1}) after isopycnic separation of RNA. Contrastingly, methane formation was

strong and archaeal and *mcrA* gene copies increased likewise indicating that methylotrophic

methanogens were active (Figures 1A and B). Because of mixotrophic assimilation capabilities

in methylotrophic methanogens, i.e., utilizing methylated compounds and DIC, a series of SIP

slurry experiments were conducted with combinations of methanol and DIC in order to improve

the sensitivity of RNA-SIP: double ^{13}C-label (methanol + DIC), single ^{13}C-label (one of the

substrates labeled), both substrates unlabeled, and a ^{13}C-DIC control (Figure 2, Figure S2). After

density separation of RNA, different degrees of RNA labeling were detected in isotopically

heavy gradient fractions, e.g., >1.803 g mL^{-1} (Figure 2). Strongest ^{13}C-labeling, as indicated by

largest amounts of RNA found in gradient fractions > 1.803 g mL^{-1}, was detected in RNA from

incubations with double ^{13}C-labeling (Figure 2A), followed by single-label incubations with ^{13}C-

DIC. For single-label ^{13}C-methanol incubations, however, RNA fraction shifts according to

density were minor compared to unlabeled incubations, indicating limited assimilation of ^{13}C-

methanol into RNA of methylotrophic methanogens.

In order to estimate ^{13}C-labeling levels of methanogens in single SIP experiments (^{13}C-methanol

or ^{13}C-DIC), a series of molecular techniques were applied including qPCR of cDNA in heavy

fractions of RNA-SIP samples, archaeal 16S rRNA sequencing from RNA-SIP fractions and

δ^{13}C value determination of methanogen lipids, e.g., phytanes derived from intact polar archaeol-

based molecules. In incubations amended with ^{13}C-DIC and unlabeled methanol, archaeal 16S

rRNA gene copies were one order of magnitude higher in the heavier gradient fractions (i.e.,

1.803–1.823 g mL^{-1}) than that in ^{13}C-methanol incubations (Figure 2B). Correspondingly,

Illumina sequencing of RNA revealed that sequences identified as related to the genera

Methanococcoides and *Methanolobus* were more dominant in the heavy than in the light

fractions from incubations amended with unlabeled methanol and ^{13}C-DIC (Figure 2C, S3).

Light fractions were overall mainly composed of anaerobic methanotrophic (ANME) archaea,

Bathyarchaeota and Lokiarchaeota except for methylotrophic methanogens (Figure 2C). For ^{13}C-

DIC control incubations, abundances of methanogens were low in SIP samples (Figure S3). For

unlabeled methanol incubations, given the low amount of labeled RNA in heavy fractions, no

amplicons were obtained, but light fractions showed a high abundance of methylotrophic

methanogens (Figure S3). Classifications were confirmed by phylogenetic clustering of cloned

16S rRNA gene fragments (about 800 base pairs) with OTU sequences representing

Methanococcoides and *Methanolobus* spp. (Figure S4). Sequences of these methanogens

accounted for more than 97% of total archaea in heavy gradient fractions. However, known

hydrogenotrophic methanogens were undetectable (Figure 2C) although 5 and 10% of methane

was formed from DIC in incubations with MZ and SRZ sediments, respectively (Table 1).

In parallel to RNA-SIP, lipid-SIP incubations with SRZ sediment slurries demonstrated δ^{13}C

values of phytane and phytenes being more positive in ^{13}C-DIC and unlabeled methanol

treatment than that in ^{13}C-methanol amendments, while the opposite was found in MZ sediment

incubations (Figure 3A). After elimination of ^{13}C-DIC dilution effects by ambient inorganic

carbon, DIC contributions to lipids ranged from 59.3% to 86.1% in SRZ sediment incubations,

which was constantly higher than that of MZ sediment incubations (52.7% to 56.4%).

To understand how carbon is assimilated into lipids by methylotrophic methanogens, pure

culture incubations of *M. methylutens* were performed with 5% of the ^{13}C-labeled substrates (i.e.,

DIC or MeOH) and the dominating archaeal lipids archaeol (AR) and hydroxyarchaeol (OH-AR)

were directly analyzed without cleavage. In contrast to sediment incubations, lipids showed

lower $f_{\text{DIC/lipid}}$ (~49%) based on the carbon incorporation in single labeling incubations (Figure

3B).

Methylotrophic methanogenesis from DIC in autoclaved slurry incubations

The unexpectedly high proportion of methane formed from DIC in methanol amended sediment

slurry incubations (Table 1) prompted us to investigate the underlying mechanism in more detail.

Thus, autoclaved sediment slurries were used as a surrogate of natural sediment, but with all

microorganisms killed, and inoculated with the obligate methylotroph *M. methylutens*. Hematite

and magnetite known to serve as electron acceptors or conductors [43,44] were added along with

humic acid, and AQDS as electron shuttles, as well as an additional electron donor (H_2), which

are all known to stimulate methanogenesis [43,44,45,46].

Methane concentrations in incubations with hematite and humic acid were higher than that of the

other incubations after 7 days (Figure 4A). Although methane production rates were low,

methane proportions from DIC in treatments with *M. methylutens* alone, H_2, AQDS and

magnetite were much higher ($f_{DIC/CH4}$, ~10%) than that in incubations with hematite and humic

acid (~2%) (Figure 4B). Linear regression showed a strong correlation between methane

production rate and CO_2-dependent methanogenesis by methylotrophic methanogens on day 3

and 5 of the incubations, indicating that lower methanogenesis rates triggered higher levels of

methane formation derived from [13]C-DIC (Figure 4C).

Discussion

In this study, we utilized RNA-SIP employing [13]C-DIC and methanol and successfully identified

methylotrophic methanogens in both, SRZ and MZ sediments of the Helgoland mud area in the

North Sea. We demonstrated that the addition of [13]C-DIC is necessary to detect label in RNA of

methylotrophic methanogens rather than using [13]C-methanol as energy substrate alone. We

further evaluated carbon utilization patterns of the methylotrophic methanogens by lipid-SIP and

identified an unexpectedly high DIC assimilation into characteristic lipids within SRZ sediment.

Isotope probing experiments unexpectedly revealed that up to 12% of methane was formed from

DIC by the presumably "obligate" methylotrophic methanogen, *M. methylutens*, thereby

suggesting an explanation for the elevated DIC incorporation into biomass.

Carbon assimilation by methylotrophic methanogens in sediment incubations

Nucleic acids-SIP techniques depend on [13]C-labeling levels of DNA or RNA molecules, from

which carbon assimilation can be reconstructed and compared to the known pathway of nucleic

acid biosynthesis from methyl-groups in methanogens [47,48,49,50,51,52]. The current

pathways show that only one carbon atom stems from methanol in ribose-5-phosphate while 25%

to 40% of carbon in nucleobases originates from the methyl carbon of the substrate (Figure 5).

This is corroborated by our RNA-SIP experiments using [13]C-labeled methanol alone, but RNA

was not found to be labeled effectively enough for density separation and further sequence

analysis. However, by additionally using [13]C-DIC, we found high 16S rRNA copy numbers

(Figure 2B) and a high representation of known methylotrophic methanogens (Figure 2C) in the

heavy RNA gradient fractions, successfully recovering [13]C-labeled RNA of methylotrophic

methanogens in the SRZ and MZ sediments of the Helgoland mud area. Consequently, RNA

labeling will be more effective in methylotrophic methanogenic archaea by DIC than methanol.

Combined with downstream analysis including qPCR, 16S rRNA sequencing and cloning, we

directly show that members of the genus *Methanococcoides* were the predominantly active

methylotrophic methanogens in SRZ incubations, while *Methanococcoides* together with

Methanolobus were dominating in MZ incubations. Hence, addition of [13]C-labeled DIC or a

combination of both substrates labeled enables tracking of methylotrophic methanogens via

RNA-SIP techniques. For carbon assimilation into nucleic acids of these methanogens, both

proposed biosynthesis pathway of nucleic acid and labeling strategy of RNA-SIP confirmed

inorganic carbon as the main carbon source for nucleic acids.

Since carbon assimilation into biomass cannot be quantified by RNA-SIP, lipid-SIP was used. In

lipid-SIP analysis, we evaluated [13]C-incorporation into intact polar archaeol- and

hydroxyarchaeol diether molecules, which are the dominant lipids produced by moderately

thermophilic methanogenic archaea [53,54,55], via phytane and phytene side-chain analysis

(Figure 3). These moieties were the only ones being [13]C-labeled while tetraether-derived

biphytane and cycloalkylated biphytanes as indicators of archaea such as Thaumarchaeota [56],

anaerobic methanotrophs [57,58] or Bathyarchaeota [59] did not show a [13]C incorporation

(Figure S5). This was corroborated by our sequencing results demonstrating that methylotrophic

methanogens were the dominant archaea in the heavy fractions and that the relative abundances

of other archaea were very low or even below detection (Figure 2C).

By evaluating [13]C incorporation into methanogen-derived phytane and phytenes, lipid-SIP

provides insight into methanogen activities and carbon utilization. As the main precursors of

ether lipids in archaea, biosynthesis of isopentenyl diphosphate (IPP) and dimethylallyl

diphosphate (DMAPP) proceeds via the modified mevalonate pathway [60,61,62]. In this

pathway, mevalonate-5-phosphate is decarboxylated to IPP, in which three out of five carbon

atoms are derived from methanol (Figure 6). DMAPP is further converted to geranylgeranyl

diphosphate (GGPP), which receives 60% of its carbon from methanol, suggestive of a lower

DIC contribution to isoprenoid chains than methanol. This was supported by the fact that

archaeol and hydroxyarchaeol contained more methanol-derived than DIC-derived carbon using

a pure culture of *M. methylutens* (Figure 3B). However, unlike the proposed lipid biosynthesis

pathway and the pure culture, clearly more DIC was assimilated into lipids than methanol in both

sediment incubations, which was most prominent in the sediment from the SRZ (Figure 3A). We,

moreover, detected that ~10% of methane produced was derived from DIC during the SIP

experiments and using *M. methylutens* in autoclaved sediment slurry incubations (Table 1, Figure

4). From the lipid biosynthesis pathway it is very likely that part of the DIC is converted to

methyl-tetrahydrosarcinapterin (CH_3-H_4SPT) and further reduced to methane, and thus, such

CH_3-H_4SPT generated from CO_2 will be simultaneously available for lipid biosynthesis (Figure 6)

leading to the ^{13}C-enrichment of the lipid pool.

CO_2 reduction to methane by obligate methylotrophic methanogens

There are two types of CO_2-dependent methanogenesis: i) Hydrogenotrophic methanogenesis by

the orders of *Methanopyrales*, *Methanococcales*, *Methanobacteriales*, *Methanomicrobiales*,

Methanocellales and *Methanosarcinales* [2,3]. These methanogens contain F_{420}-reducing [NiFe]-

hydrogenase to catalyze F_{420} reduction by H_2 [63]. ii) Mediation by interspecies electron transfer

between bacteria and some members of the *Methanosarcinales*. CO_2 reduction to methane was

observed in *Methanosaeta* and *Methanosarcina* during syntrophic growth with *Geobacter*

species on alcohols (ethanol, propanol, and butanol), as electrons generated from *Geobacter* are

directly transferred to methanogens to reduce CO_2 [64,65,66,67].

Based on our SIP incubations with SRZ sediment showing 10% of methane generation from DIC

at low H_2 partial pressure (< 0.3 Pa) (Table 1) and the overall lack of hydrogenotrophic

methanogens in RNA-SIP fractions (Figure 2C), we argue that H_2-dependent methanogenesis

does not play a role [2]. Similarly, in autoclaved sediment slurry (Figure 4B), the "obligate"

methylotroph *M. methylutens* generated methane from CO_2 without a hydrogen (or electron) supplying partner microorganism.

Members of the genus *Methanococcoides* are considered as obligate methylotrophic methanogens since no F_{420}-reducing [NiFe]-hydrogenase was detected in their genomes [60, 68], ruling out hydrogenotrophic methanogenesis in sediment incubations. Nevertheless, part of the methane formed during methylotrophic methanogenesis by *M. methylutens* was from CO_2, especially when methane production rates were low (Figure 4C). Apparently, at high rates of methanol dissimilation to CO_2, the reverse pathway of CO_2 reduction to methane was outcompeted. Higher rates of methylotrophic methanogenesis can be achieved potentially by amendments in autoclaved slurries using hydrogen as electron donor, electron conductors (hematite, magnetite) and electron shuttles (humic acid, AQDS); these compounds are known to affect electron utilizations in microorganisms and stimulate methanogenesis [43,44,45,46]. In our incubations, we found humic acid and hematite most strongly stimulating methylotrophic methanogenesis. Although the underlying mechanism is beyond the scope of the current paper, methylotrophic methanogens in our incubations could take advantage of hematite as potential electron conductor [43,44] or humic acid as electron shuttle [46] as indicated by a higher rate of methanogenesis compared to the other treatments (Figure 4A).

It has been shown that 3% of methane was produced from CO_2 during methylotrophic methanogenesis of *Methanosarcina barkeri* (i.e., a facultative methylotroph) without the addition of H_2 [20], which is similar to about 2.5% of methane generated from CO_2 by *M. methylutens* (i.e., a presumed "obligate" methylotroph) in our study (Table S2). However, methane generation from CO_2 increased to 10% in SRZ sediment incubations (Table 1), highlighting the importance of the activity of concomitant CO_2 reduction during methylotrophic methanogenesis

in marine sediments. It seems that the substantially higher methanogenesis rates in pure cultures

decrease the amount of methane produced from CO_2. In marine sediment methylotrophic

methanogenesis rates are likely lower than those in pure cultures because of the limitation in

methylated substrates [6, 14], strongly suggesting that methane generation from CO_2 by

methylotrophic methanogens is underestimated under *in situ* conditions. Thus, CO_2 conversion to

methane has to be considered when estimates of *in situ* methylotrophic methanogenesis in

marine sediments are performed.

Conclusions

In this study, we have shown that [13]C-DIC is required as co-substrate for successful

identification of methylotrophic methanogens by RNA-SIP in marine sediments. DIC is the main

carbon source for biosynthesis of nucleic acids in these methanogens and thus using [13]C-

methanol as energy and carbon substrate alone is insufficient in SIP experiments. Given the

intricacies of known assimilatory pathways in methanogenic archaea as a functional group, it

might be necessary to at least check for the possibility of DIC as a main assimilatory carbon

component in all methanogens for successful SIP experiments. In general, it seems that archaea

have a propensity for using DIC as a carbon source for assimilation [59, 69], possibly as an

evolutionary adaptation to environments with limited availability of organic carbon [70]; in our

laboratory, we are currently unveiling more DIC assimilating microorganisms in marine

sediments.

But beyond known pathways, we detected an unexpectedly high amount of methane (> 10%)

formed from DIC. This finding strongly suggests that the alleged obligate methylotroph studied

here was rather mixotrophically converting both available substrates (DIC, methanol) to methane.

This seems to be a conundrum, however, our detailed labeling studies showed that the kinetics of

substrate utilization apparently is a decisive factor in channeling more or less CO_2 into the

pathway of methanogenesis: more methane formed from CO_2 when the overall kinetics were

slow, and vice versa. It is not obvious currently, what the benefit for the cell is, since more

methane could be formed when methanogenesis proceeds by disproportionation of methanol.

From an ecological perspective, DIC is a much more pertinent substrate than methanol (or other

methyl compounds) in marine sediments [4, 6], and thus, we speculate that more DIC reduction

by obligate methylotrophic methanogens occurs *in situ* than is currently known. A larger

proportion of DIC-dependent methanogenesis from methyl compounds should also impact

carbon fractionation, and thus, delta $^{13}CH_4$ might be overprinted by such mixotrophic

methanogenesis. Thus, the CO_2 reduction to methane and assimilation into biomass by obligate

methylotrophic methanogens plays a much more important role in the environment than was

previously known.

Acknowledgments

This study was supported by the Research Center/Cluster of Excellence 'The Ocean in the Earth

System' (MARUM) funded by the Deutsche Forschungsgemeinschaft (DFG) and by the

University of Bremen. Xiuran Yin and Weichao Wu were funded by the scholarship from China

Scholarship Council (CSC). We thank the captain, crew and scientists of R/V HEINCKE

expeditions HE443.

Conflict of interest

The authors declare that they have no conflict of interest.

Data availability

Sequencing data have been submitted to GenBank Short Reads Archive with accession numbers from SRR8207425 to SRR8207442.

Reference

1. Ferry JG, Lessner DJ. Methanogenesis in marine sediments. Ann N Y Acad Sci. 2008;1125:147-57.

2. Thauer RK, Kaster AK, Seedorf H, Buckel W, Hedderich R. Methanogenic archaea: ecologically relevant differences in energy conservation. Nat Rev Microbiol. 2008;6:579-591.

3. Liu Y, Whitman WB. Metabolic, phylogenetic, and ecological diversity of the methanogenic archaea. Ann N Y Acad Sci. 2008;1125:171-89.

4. Zhuang G-C, Elling FJ, Nigro LM, Samarkin V, Joye SB, Teske A, et al. Multiple evidence for methylotrophic methanogenesis as the dominant methanogenic pathway in hypersaline sediments from the Orca Basin, Gulf of Mexico. Geochim Cosmochim Acta. 2016;187:1-20.

5. Lazar CS, Parkes RJ, Cragg BA, L'Haridon S, Toffin L. Methanogenic diversity and activity in hypersaline sediments of the centre of the Napoli mud volcano, Eastern Mediterranean Sea. Environ Microbiol. 2011;13:2078-91.

6. Zhuang G-C, Heuer VB, Lazar CS, Goldhammer T, Wendt J, Samarkin VA, et al. Relative importance of methylotrophic methanogenesis in sediments of the Western Mediterranean Sea. Geochimica et Cosmochimica Acta. 2018;224:171-86.

7. Maltby J, Steinle L, Löscher CR, Bange HW, Fischer MA, Schmidt M, et al. Microbial methanogenesis in the sulfate-reducing zone of sediments in the Eckernförde Bay, SW Baltic Sea. Biogeosciences. 2018;15:137-57.

8.   Yanagawa K, Tani A, Yamamoto N, Hachikubo A, Kano A, Matsumoto R, et al.

Biogeochemical cycle of methanol in anoxic deep-sea sediments. Microbes Environ.

2016;31:190-3.

9.   Zhuang G-C, Lin Y-S, Elvert M, Heuer VB, Hinrichs K-U. Gas chromatographic

analysis of methanol and ethanol in marine sediment pore waters: Validation and

implementation of three pretreatment techniques. Mar Chem. 2014;160:82-90.

10.   Oremland RS, Polcin S. Methanogenesis and sulfate reduction: competitive and

noncompetitive substrates in estuarine sediments. Appl Environ Microbiol.

1982;44:1270-6.

11.   Oni O, Miyatake T, Kasten S, Richter-Heitmann T, Fischer D, Wagenknecht L, et al.

Distinct microbial populations are tightly linked to the profile of dissolved iron in the

methanic sediments of the Helgoland mud area, North Sea. Front Microbiol. 2015;6:365.

12.   Evans PN, Parks DH, Chadwick GL, Robbins SJ, Orphan VJ, Golding SD, et al.

Methane metabolism in the archaeal phylum Bathyarchaeota revealed by genome-centric

metagenomics. Science. 2015;350:432-8.

13.   Weimer PJ, Zeikus JG. Acetate metabolism in *Methanosarcina barkeri*. Arch Microbiol.

1978;119:175-82.

14.   Summons RE, Franzmann PD, Nichols PD. Carbon isotopic fractionation associated

with methylotrophic methanogenesis. Org Geochem. 1998;28:465-75.

15.   Hippe H, Caspari D, Fiebig K, Gottschalk G. Utilization of trimethylamine and other N-

methyl compounds for growth and methane formation by *Methanosarcina barkeri*. Proc

Natl Acad Sci USA. 1979;76:494-8.

16. Teeling H, Glockner FO. Current opportunities and challenges in microbial metagenome

analysis–a bioinformatic perspective. Brief Bioinform. 2012;13:728-42.

17. Singer E, Wagner M, Woyke T. Capturing the genetic makeup of the active microbiome

in situ. ISME J. 2017;11:1949-63.

18. Neelakanta G, Sultana H. The use of metagenomic approaches to analyze changes in

microbial communities. Microbiol Insights. 2013;6:37-48.

19. Thauer RK. Anaerobic oxidation of methane with sulfate: on the reversibility of the

reactions that are catalyzed by enzymes also involved in methanogenesis from CO_2. Curr

Opin Microbiol. 2011;14:292-9.

20. Weimer PJ, Zeikus JG. One carbon metabolism in methanogenic bacteria. Arch

Microbiol. 1978;119:47-57.

21. Feijo Delgado F, Cermak N, Hecht VC, Son S, Li Y, Knudsen SM, et al. Intracellular

water exchange for measuring the dry mass, water mass and changes in chemical

composition of living cells. PLoS One. 2013;8:e67590.

22. Friedrich MW. Stable-isotope probing of DNA: insights into the function of uncultivated

microorganisms from isotopically labeled metagenomes. Curr Opin Biotechnol.

2006;17:59-66.

23. Lueders T. Stable isotope probing of hydrocarbon-degraders. Handbook of Hydrocarbon

and Lipid Microbiology 2010. p. 4011-26.

24. Grob C, Taubert M, Howat AM, Burns OJ, Dixon JL, Richnow HH, et al. Combining

metagenomics with metaproteomics and stable isotope probing reveals metabolic

pathways used by a naturally occurring marine methylotroph. Environ Microbiol.

2015;17:4007-18.

25. Fortunato CS, Huber JA. Coupled RNA-SIP and metatranscriptomics of active

chemolithoautotrophic communities at a deep-sea hydrothermal vent. ISME J.

2016;10:1925-38.

26. Manefield M, Whiteley AS, Ostle N, Ineson P, Bailey MJ. Technical considerations for

RNA-based stable isotope probing an approach to associating microbial diversity with

microbial community function. Rapid Commun Mass Spectrom. 2002;16:2179-83.

27. Vandieken V, Thamdrup B. Identification of acetate-oxidizing bacteria in a coastal

marine surface sediment by RNA-stable isotope probing in anoxic slurries and intact

cores. FEMS Microbiol Ecol. 2013;84:373-86.

28. Lueders T, Wagner B, Claus P, Friedrich MW. Stable isotope probing of rRNA and

DNA reveals a dynamic methylotroph community and trophic interactions with fungi

and protozoa in oxic rice field soil. Environ Microbiol. 2003;6:60-72.

29. Neufeld JD, Schafer H, Cox MJ, Boden R, McDonald IR, Murrell JC. Stable-isotope

probing implicates *Methylophaga* spp and novel *Gammaproteobacteria* in marine

methanol and methylamine metabolism. ISME J. 2007;1:480-91.

30. Wegener G, Kellermann MY, Elvert M. Tracking activity and function of

microorganisms by stable isotope probing of membrane lipids. Curr Opin Biotechnol.

2016;41:43-52.

31. Boschker HTS, Nold SC, Wellsbury P, Bos D, de Graaf W, Pel R, et al. Direct linking of

microbial populations to specific biogeochemical processes by ^{13}C-labelling of

biomarkers. Nature. 1998;392:801-5.

32. Widdel F, Pfennig N. Studies on dissimilatory sulfate-reducing bacteria that decompose

fatty acids I. Isolation of new sulfate-reducing bacteria enriched with acetate from saline

environments. Description of *Desulfobacter postgatei* gen. nov., sp. nov. Arch Microbiol.

1981;134:282-5.

33. Widdel F, Kohring GW, Mayer F. Studies on dissimilatory sulfate-reducing bacteria that

decompose fatty acids III. Characterization of the filamentous gliding *Desulfonema*

limicola gen. nov. sp. nov., and *Desulfonema magnum* sp. nov. Arch Microbiol.

1983;134:286-94.

34. Aromokeye DA, Richter-Heitmann T, Oni OE, Kulkarni A, Yin X, Kasten S, et al.

Temperature controls crystalline iron oxide utilization by microbial communities in

methanic ferruginous marine sediment incubations. Front Microbiol. 2018;9:2574.

35. Ertefai TF, Heuer VB, Prieto-Mollar X, Vogt C, Sylva SP, Seewald J, et al. The

biogeochemistry of sorbed methane in marine sediments. Geochim Cosmochim Acta.

2010;74:6033-48.

36. Lueders T, Manefield M, Friedrich MW. Enhanced sensitivity of DNA- and rRNA-based

stable isotope probing by fractionation and quantitative analysis of isopycnic

centrifugation gradients. Environ Microbiol. 2003;6:73-8.

37. Friedrich MW. Methyl-coenzyme M reductase genes: unique functional markers for

methanogenic and anaerobic methane-oxidizing Archaea. Methods Enzymol.

2005;397:428-42.

38. Sturt HF, Summons RE, Smith K, Elvert M, Hinrichs KU. Intact polar membrane lipids

in prokaryotes and sediments deciphered by high-performance liquid

chromatography/electrospray ionization multistage mass spectrometry–new biomarkers

for biogeochemistry and microbial ecology. Rapid Commun Mass Spectrom.

2004;18:617-28.

39. Zhu C, Lipp JS, Wörmer L, Becker KW, Schröder J, Hinrichs K-U. Comprehensive
glycerol ether lipid fingerprints through a novel reversed phase liquid chromatography–
mass spectrometry protocol. Org Geochem. 2013;65:53-62.

40. Liu X-L, Lipp JS, Simpson JH, Lin Y-S, Summons RE, Hinrichs K-U. Mono- and
dihydroxyl glycerol dibiphytanyl glycerol tetraethers in marine sediments: Identification
of both core and intact polar lipid forms. Geochim Cosmochim Acta. 2012;89:102-15.

41. Kellermann MY, Yoshinaga MY, Wegener G, Krukenberg V, Hinrichs K-U. Tracing the
production and fate of individual archaeal intact polar lipids using stable isotope probing.
Org Geochem. 2016;95:13-20.

42. Boschker HTS, Middelburg JJ. Stable isotopes and biomarkers in microbial ecology.
FEMS Microbiol Ecol. 2002;40:85–95.

43. Kato S, Hashimoto K, Watanabe K. Methanogenesis facilitated by electric syntrophy via
(semi)conductive iron-oxide minerals. Environ Microbiol. 2012;14:1646-54.

44. Kato S, Nakamura R, Kai F, Watanabe K, Hashimoto K. Respiratory interactions of soil
bacteria with (semi)conductive iron-oxide minerals. Environ Microbiol. 2010;12:3114-
23.

45. Bond DR, Lovley DR. Reduction of Fe(III) oxide by methanogens in the presence and
absence of extracellular quinones. Environ Microbiol. 2002;4:115–24.

46. Lovley DR, Fraga JL, Coates JD, L. B-HE. Humics as an electron donor for anaerobic
respiration. Environ Microbiol. 1999;1:89-99.

47. Choquet CG, Richards JC, Patel GB, Sprott GD. Ribose biosynthesis in methanogenic
bacteria. Arch Microbiol. 1994a;161:481-8.

48. Choquet CG, Richards JC, Patel GB, Sprott GD. Purine and pyrimidine biosynthesis in

methanogenic bacteria. Arch Microbiol. 1994b;161:471-80.

49. Ekiel I, Smith ICP, Sportt GD. Biosynthetic pathways in *Methanospirillum hungatei* as

determined by ^{13}C nuclear magnetic resonance. J Bacteriol. 1983;156:316-26.

50. Nyce GW, White RH. dTMP biosynthesis in archaea. J Bacteriol. 1996 178:914–6.

51. Soderberg T. Biosynthesis of ribose-5-phosphate and erythrose-4-phosphate in archaea:

a phylogenetic analysis of archaeal genomes. Archaea. 2005;1:347-52.

52. Sorokin DY, Makarova KS, Abbas B, Ferrer M, Golyshin PN, Galinski EA, et al.

Discovery of extremely halophilic, methyl-reducing euryarchaea provides insights into

the evolutionary origin of methanogenesis. Nat Microbiol. 2017;2:17081.

53. Nishihara M, Koga Y. Hydroxyarchaetidylserine and hydroxyarchaetidyl-myo-inositol

in *Methanosarcina barkeri*: polar lipids with a new ether core portion. Biochim Biophys

Acta. 1991;1082:211-7.

54. Nishihara M, Utagawa M, Akutsu H, Koga Y. Archaea contain a novel diether

phosphoglycolipid with a polar head group identical to the conserved core of eucaryal

glycosyl phosphatidylinosito. J Biol Chem. 1992;267:12432-5.

55. Pancost RD, McClymont EL, Bingham EM, Roberts Z, Charman DJ, Hornibrook ERC,

et al. Archaeol as a methanogen biomarker in ombrotrophic bogs. Org Geochem.

2011;42:1279-87.

56. Elling FJ, Könneke M, Lipp JS, Becker KW, Gagen EJ, Hinrichs K-U. Effects of growth

phase on the membrane lipid composition of the thaumarchaeon *Nitrosopumilus*

maritimus and their implications for archaeal lipid distributions in the marine

environment. Geochim Cosmochim Acta. 2014;141:579-97.

57.   Blumenberg M, Seifert R, Reitner J, Pape T, Michaelis W. Membrane lipid patterns

typify distinct anaerobic methanotrophic consortia. Proc Natl Acad Sci USA.

2004;101:11111–6.

58.   Rossel PE, Lipp JS, Fredricks HF, Arnds J, Boetius A, Elvert M, et al. Intact polar lipids

of anaerobic methanotrophic archaea and associated bacteria. Org Geochem.

2008;39:992-9.

59.   Yu T, Wu W, Liang W, Lever MA, Hinrichs K-U, Wang F. Growth of sedimentary

Bathyarchaeota on lignin as an energy source. Proc Natl Acad Sci USA. 2018;115:6022-

7.

60.   Allen MA, Lauro FM, Williams TJ, Burg D, Siddiqui KS, De Francisci D, et al. The

genome sequence of the psychrophilic archaeon, *Methanococcoides burtonii*: the role of

genome evolution in cold adaptation. ISME J. 2009;3:1012-35.

61.   Grochowski LL, Xu H, White RH. *Methanocaldococcus jannaschii* uses a modified

mevalonate pathway for biosynthesis of isopentenyl diphosphate. J Bacteriol.

2006;188:3192-8.

62.   Nichols DS, Miller MR, Davies NW, Goodchild A, Raftery M, Cavicchioli R. Cold

adaptation in the Antarctic archaeon *Methanococcoides burtonii* involves membrane

lipid unsaturation. J Bacteriol. 2004;186:8508-15.

63.   Thauer RK, Kaster AK, Goenrich M, Schick M, Hiromoto T, Shima S. Hydrogenases

from methanogenic archaea, nickel, a novel cofactor, and H_2 storage. Annual review of

biochemistry. 2010;79:507-36.

64.   Lovley DR. Happy together: microbial communities that hook up to swap electrons.

ISME J. 2017;11:327-36.

65.   Rotaru AE, Shrestha PM, Liu F, Markovaite B, Chen S, Nevin KP, et al. Direct

617   interspecies electron transfer between *Geobacter metallireducens* and *Methanosarcina*

618   *barkeri*. Appl Microbiol Biotechnol. 2014;80:4599-605.

619   66.   Rotaru A-E, Shrestha PM, Liu F, Shrestha M, Shrestha D, Embree M, et al. A new

620   model for electron flow during anaerobic digestion: direct interspecies electron transfer

621   to *Methanosaeta* for the reduction of carbon dioxide to methane. Energy Environ Sci.

622   2014;7:408-15.

623   67.   Wang LY, Nevin KP, Woodard TL, Mu BZ, Lovley DR. Expanding the diet for DIET:

624   electron donors supporting direct interspecies electron transfer (DIET) in defined co-

625   cultures. Front Microbiol. 2016;7:236.

626   68.   Guan Y, Ngugi DK, Blom J, Ali S, Ferry JG, U. S. Draft genome sequence of an

627   obligately methylotrophic methanogen, *Methanococcoides methylutens*, isolated from

628   marine sediment. Genome Announc. 2014;2:e01184-14.

629   69.   Kellermann MY, Wegener G, Elvert M, Yoshinaga MY, Lin YS, Holler T, et al.

630   Autotrophy as a predominant mode of carbon fixation in anaerobic methane-oxidizing

631   microbial communities. Proc Natl Acad Sci USA. 2012;109:19321-6.

632   70.   Weiss MC, Sousa FL, Mrnjavac N, Neukirchen S, Roettger M, Nelson-Sathi S, et al. The

633   physiology and habitat of the last universal common ancestor. Nat Microbiol.

634   2016;1:16116.

635   71.   Qin S, Sun Y, Tang Y. Early hydrocarbon generation of algae and influences of

636   inorganic environments during low temperature simulation. Energ Explor Exploit.

637   2008;26:377-396.

638   72.   Koga Y, Morii H. Biosynthesis of ether-type polar lipids in archaea and evolutionary

639   considerations. Microbiol Mol Biol Rev. 2007;71:97-120.

640   73.   Ekiel I, Sportt GD, Patel GB. Acetate and CO_2 assimilation by *Methanothrix concilii*. J

641   Bacteriol. 1985;162:905-8.

 Figure legends

Figure 1. Dynamics of methane formation and archaeal populations in stable isotope probing

incubations with SRZ and MZ sediment samples. (A) Methane concentrations in SIP incubations.

Methane data is presented as average values (n = 3, error bar = SD). (B) Gene copy numbers of

archaea (16S rRNA genes) and methanogens (*mcrA* gene). Gene copies were quantified based on

DNA extracts at harvest. Fold increase of gene copies were indicated above each histogram by

comparing gene copies on day 0 after preincubation (n = 3, error bar = SD). DIC: dissolved

inorganic carbon, i.e., bicarbonate; MeOH: methanol.

Figure 2. Density distribution of RNA, gene copy numbers, and community composition from

SIP incubations with SRZ and MZ sediment after isopycnic separation. (A) RNA profiles from

different RNA-SIP experiments. (B) Gene copy numbers of archaeal cDNA in heavy fractions

(1.803 g mL^{-1} to 1.823 g mL^{-1}) from RNA-SIP experiments. (C) Relative abundances of density

separated archaeal 16S rRNA from single-labeling incubations in light (1.771 g mL^{-1} to 1.800 g

mL^{-1}) and heavy (1.803 g mL^{-1} to 1.835 g mL^{-1}) gradient fractions.

Figure 3. Lipid-SIP experiments from sediment incubations (natural community) and pure

cultures in Widdel medium. Lipid δ^{13}C values were measured in homogenized samples after

methanogenesis had ceased. (A) δ^{13}C values of phytanes in sediment incubations with 70% ^{13}C-

DIC. Phytane originates from intact polar archaeol lipids and phytenes (phytene I and phytene II)

derive from intact polar hydroxyarchaeol lipids. $f_{DIC/lipid}$ is indicated on the top of bars from

single labeling incubations based on Eq.5. (B) δ^{13}C values of archaeol (AR) and hydroxyarchaeol

(AR-OH) in pure culture of *M. methylutens* treated with 5% ^{13}C-labeled substrates (methanol or

DIC). (C) Structures of archaeal lipids. Enclosed structures of phytenes in Figure C were

tentatively assigned according to GC-MS mass spectra (Figure S6) [71]. Data is expressed as

average values (n = 3, error bar=SD).

Figure 4. Methane production from DIC during methylotrophic methanogenesis in autoclaved

slurry supplemented with pure culture of *M. methylutens*. (A) Total methane concentrations in

headspace. (B) Proportion of methane derived from DIC. Methane proportion from DIC

(f_{DIC/CH_4}) was calculated according to Eq.2. Data is expressed as average values (n = 3, error bar

= SD). (C) Linear correlation between methanogenesis rate and (f_{DIC/CH_4}) after 3 and 5 days.

Day 3: Pearson's r = -0.92, $P < 0.001$, C1(0.95) = -0.79 > r > -0.97; Day 5: Pearson's r = -0.85,

$P < 0.001$, C1(0.95) = -0.62 > r > -0.94. M. met: *M. methylutens*.

Figure 5. Carbon assimilation into nucleic acids in methylotrophic methanogens. Biosynthesis of

nucleotide moieties, the pyrimidine and purine bases, as well as the C5-carbon from [13]C-labeled

methanol in methylotrophic methanogens based on previous studies [47,48,49,50,51,52] with

final carbon contribution from methanol added besides the compounds. Black arrows indicate

ribose synthesis and the blue arrows represent synthesis of base moieties in nucleosides. The

reverse gluconeogenesis pathway is displayed in green and the reverse ribulose monophosphate

pathway in pink.

Figure 6. Carbon assimilation into lipids in methylotrophic methanogens. Methylotrophic

methanogenesis pathway from methanol (yellow) and carbon assimilation pattern into isoprenoid

chains of archaeal lipids (blue) with carbon contribution from [13]C-methanol added besides the

compounds. The pathway of archaeal lipid biosynthesis is based on previous studies [60, 72,73].

Table 1. [13]C fractional abundance and H_2 partial pressures in SIP incubations. Data is presented

as average values (n = 3).

Sediment	Substrates	$^{13}F_{DIC}$ (%)	f_{DIC/CH_4} (%)[a]	H_2 (Pa)[b]	Incubation time (d)
SRZ	DIC + [13]C-MeOH	3.7 ± 0.4	89.3 ± 0.3	NA	43
MZ	DIC + [13]C-MeOH	3.0 ± 0.0	96.4 ± 0.1	NA	19
SRZ	MeOH + [13]C-DIC	83.6 ± 0.6	10.3 ± 0.2	0.1 ± 0.1	43
MZ	MeOH + [13]C-DIC	69.8 ± 0.7	3.4± 0.1	0.3 ± 0.0	19

688 [a]Methane proportion from DIC (f_{DIC/CH_4}) in "methanol + [13]C-DIC" incubations was based on

689 Eq.2. [b]H_2 partial pressure was measured on day 23 and 16 for incubation SRZ and MZ sediments,

690 respectively. NA, not analyzed.

Figures

Figure 1. Dynamics of methane formation and archaeal populations in stable isotope probing incubations with SRZ and MZ sediment samples. (A) Methane concentrations in SIP incubations. Methane data is presented as average values (n = 3, error bar = SD). (B) Gene copy numbers of archaea (16S rRNA genes) and methanogens (*mcrA* gene). Gene copies were quantified based on DNA extracts at harvest. Fold increase of gene copies were indicated above each histogram by comparing gene copies on day 0 after preincubation (n = 3, error bar = SD). DIC: dissolved inorganic carbon, i.e., bicarbonate; MeOH: methanol.

Figure 2. Density distribution of RNA, gene copy numbers, and community composition from SIP incubations with SRZ and MZ sediment after isopycnic separation. (A) RNA profiles from different RNA-SIP experiments. (B) Gene copy numbers of archaeal cDNA in heavy fractions (1.803 g mL^{-1} to 1.823 g mL^{-1}) from RNA-SIP experiments. (C) Relative abundances of density separated archaeal 16S rRNA from single-labeling incubations in light (1.771 g mL^{-1} to 1.800 g mL^{-1}) and heavy (1.803 g mL^{-1} to 1.835 g mL^{-1}) gradient fractions.

Figure 3. Lipid-SIP experiments from sediment incubations (natural community) and pure cultures in Widdel medium. Lipid δ^{13}C values were measured in homogenized samples after methanogenesis had ceased. (A) δ^{13}C values of phytanes in sediment incubations with 70% ^{13}C-DIC. Phytane originates from intact polar archaeol lipids and phytenes (phytene I and phytene II) derive from intact polar hydroxyarchaeol lipids. $f_{DIC/lipid}$ is indicated on the top of bars from single labeling incubations based on Eq.5. (B) δ^{13}C values of archaeol (AR) and hydroxyarchaeol (AR-OH) in pure culture of *M. methylutens* treated with 5% ^{13}C-labeled substrates (methanol or DIC). (C) Structures of archaeal lipids. Enclosed structures of phytenes in Figure C were tentatively assigned according to GC-MS mass spectra (Figure S6) [71]. Data is expressed as average values (n = 3, error bar=SD).

Figure 4. Methane production from DIC during methylotrophic methanogenesis in autoclaved

slurry supplemented with pure culture of *M. methylutens*. (A) Total methane concentrations in

headspace. (B) Proportion of methane derived from DIC. Methane proportion from DIC

(f_{DIC/CH_4}) was calculated according to Eq.2. Data is expressed as average values (n = 3, error bar

= SD). (C) Linear correlation between methanogenesis rate and (f_{DIC/CH_4}) after 3 and 5 days.

Day 3: Pearson's r = -0.92, $P < 0.001$, C1(0.95) = -0.79 > r > -0.97; Day 5: Pearson's r = -0.85,

$P < 0.001$, C1(0.95) = -0.62 > r > -0.94. M. met: *M. methylutens*.

Figure 5. Carbon assimilation into nucleic acids in methylotrophic methanogens. Biosynthesis of nucleotide moieties, the pyrimidine and purine bases, as well as the C5-carbon from [13]C-labeled methanol in methylotrophic methanogens based on previous studies [47,48,49,50,51,52] with final carbon contribution from methanol added besides the compounds. Black arrows indicate ribose synthesis and the blue arrows represent synthesis of base moieties in nucleosides. The reverse gluconeogenesis pathway is displayed in green and the reverse ribulose monophosphate pathway in pink.

Figure 6. Carbon assimilation into lipids in methylotrophic methanogens. Methylotrophic methanogenesis pathway from methanol (yellow) and carbon assimilation pattern into isoprenoid chains of archaeal lipids (blue) with carbon contribution from [13]C-methanol added besides the compounds. The pathway of archaeal lipid biosynthesis is based on previous studies [60, 72,73].