Title

Tempo of gene regulation in wild and cultivated *Vitis* species shows coordination

between cold deacclimation and budbreak

Running Title

Tempo of gene regulation in coordination with phenology

Authors:

Alisson P. Kovaleski[1,2*] and Jason P. Londo[1,2*]

[1] School of Integrative Plant Science – Horticulture Section, Cornell University – New

York State Agricultural Experiment Station, 630 W. North Street, Geneva, New York,

USA.

[2] United States Department of Agriculture, Agricultural Research Service, Grape

Genetics Research Unit, 630 W. North Street, Geneva, New York, USA.

[*] Corresponding authors

APK – e-mail: ap874@cornell.edu; phone: (315) 787-2484

JPL – e-mail: jason.londo@ars.usda.gov; phone: (315) 787-2463

Date of submission: 23 Jan 2019

Number of tables: 1

Number of figures: 9

Number of supplementary figures: 10

Total word count: 6499

Tempo of gene regulation in wild and cultivated *Vitis* species shows

coordination between cold deacclimation and budbreak

Highlight

Faster deacclimation and budbreak phenology is related to a faster regulon

rather than higher expression of specific genes. ABA is a master regulator of

deacclimation.

Abstract

Dormancy release, loss of cold hardiness and budbreak are critical aspects of

the annual cycle of deciduous perennial plants. Molecular control of these processes is

not fully understood, and genotypic variation may be important for climate adaptation.

Single-node cuttings from wild (*Vitis amurensis*, *V. riparia*) and cultivated *Vitis*

genotypes (*V. vinifera* 'Cabernet Sauvignon', 'Riesling') were collected from the field

during winter and placed under forcing conditions. Cold hardiness was measured daily,

and buds were collected for RNA-Seq until budbreak. Field-collected single-node

cuttings of 'Riesling' were treated with abscisic acid (ABA), and cold hardiness and

budbreak at 7 °C were tracked. Wild *Vitis* genotypes had faster deacclimation and

budbreak than *V. vinifera*. Temperature-sensing related genes were quickly and

synchronously differentially expressed in all genotypes. ABA synthesis was down-

regulated in all genotypes, and exogenous ABA prevented deacclimation. Ethylene- and

oxidative stress-related genes were transiently up-regulated. Growth-related genes

were up-regulated and showed staggering similar to deacclimation and budbreak of the

four genotypes. The gene expression cascade that occurs during deacclimation and

budburst phenology of fast (wild) and slow (cultivated) grapevines appears coordinated

and temporally conserved. This may extend to other temperate woody species and

suggest constraints on identification of process-specific keystone genes.

Key words

Budbreak, cold hardiness, deacclimation, dormancy, phenology, RNA-Seq, time-

series, *Vitis* spp. (grapevine).

Introduction

Low temperatures during winter are the most important factor limiting plant distribution (Burke *et al.*, 1976). Perennial plants withstand low temperatures by forming wood and overwintering structures (i.e., dormant buds), utilizing dormancy to survive during non-growth-conducive environmental conditions. Dormancy onset is induced by changing day length, or day length and low temperatures (Cooke *et al.*, 2012), and the transition from endogenous growth suppression (endodormancy) to environmental growth suppression (ecodormancy) occurs through accumulation of time in low, non-freezing temperatures – termed chilling accumulation (Lang *et al.*, 1987; Cooke *et al.*, 2012). Chilling greatly impacts timing of budbreak, and therefore budbreak phenology depends on temperatures during both winter (chilling) and spring (growing degree-days). This temperature-controlled interplay has been extensively studied in plants (Prentice *et al.*, 1992). While dormant buds have been the subject of many studies regarding dormancy control (Olukolu *et al.*, 2009; Ophir *et al.*, 2009; van Dyk *et al.*, 2010; Hedley *et al.*, 2010; Díaz-Riquelme *et al.*, 2012; Busov *et al.*, 2016; Sudawan *et al.*, 2016; Wu *et al.*, 2017; Meitha *et al.*, 2018), these have not considered the potential role of deacclimation, the loss of winter cold hardiness, as it impacts budbreak. Despite the foundation of dormancy research and importance of this trait for changing climate, the majority of cold hardiness studies focus on chill or freeze stress in annuals or actively growing tissues of perennials (Ruelland *et al.*, 2002; Li *et al.*, 2004; Nguyen *et al.*, 2017; Liu *et al.*, 2017a,b; Ban *et al.*, 2017; Londo *et al.*, 2018).

Dormancy is required for development of deep cold hardiness, and many processes associated with abiotic stress may be critical during dormancy. Transcriptomic tools (e.g., RNA-Seq) have revealed some of the gene expression processes that occur in dormant tissues. Dehydration-responsive element-binding 1 (DREB1)/C-repeat-binding factors (CBF) are known elicitors of cold-responsive genes following low temperature perception (Liu *et al.*, 1998). Other stress related transcription factors [other DREBs, heat shock factors (HSFs), NAC and WRKY families] are also involved in the signaling, much of these shared between heat, drought, and biotic stresses (Liu *et al.*, 1998, 2017a; Birkenbihl *et al.*, 2012). The circadian clock is also likely involved in regulation of bud dormancy (Penfield, 2008; Cooke *et al.*, 2012), and

dormancy release and budbreak have shared processes with germination, such as

hypoxia signaling via ETHYLENE-RESPONSIVE TRANSCRIPTION FACTORS (ERFs)

regulation (Penfield, 2008; Meitha *et al.*, 2018). Although known as central in

senescence and dormancy, the role of abscisic acid (ABA) in bud dormancy is not fully

understood (Cooke *et al.*, 2012). ABA promotes the induction of dormancy by blocking

cell-cell communication (Tylewicz *et al.*, 2018). Exogenous ABA applications during the

growing season have been shown to enhance hardiness earlier in the fall (Li and Dami,

2016) through increased expression of DREB1/CBF TFs (Rubio *et al.*, 2018), and

budbreak can be delayed by ABA when applied in dormant buds (Zheng *et al.*, 2015).

Considering recent evidence that budbreak phenology follows similar kinetics as

deacclimation (Kovaleski *et al.*, 2018), elucidating key gene regulation cascades during

these processes could help identify candidate genes for climate adaptation.

Cold hardiness kinetics and budbreak phenology are tightly linked (Kovaleski *et*

al., 2018), and understanding these two related traits may contribute to increased

sustainability of crop species as climate variation increases, especially when acute cold

weather episodes are expected to increase (Kolstad *et al.*, 2010). As buds transition

between dormancy stages, they alter the metabolic availability of sugars, carbon, and

reactive oxygen species (ROS) without any visual cues (Lang *et al.*, 1987). Therefore,

understanding the regulation and timing of these changes requires an understanding of

gene expression cascades. One major limitation to understanding important gene

expression changes during dormancy arises from the time scale of dormancy, spanning

throughout the entire winter. Traditional pairwise contrasts may be problematic over

large time scales as subtle changes in expression will be undetectable between sample

points, but significant over time. However, pairwise contrasts continue to be the routine

method for determining differentially expressed genes in time-series studies (Fu *et al.*,

2018; Meitha *et al.*, 2018). New methods of examining gene expression data are

needed to detect those changes that are significant over long periods of time.

Grapevine is the highest production value fruit crop worldwide with an estimated

67B (USD) impact (FAO, 2018). Sustainability is expected to be severely challenged by

climate change even in traditional production regions (Cook and Wolkovich, 2016;

Leolini *et al.*, 2018). Phenology modelling of cultivated grapevine, *Vitis vinifera*,

demonstrates wide variation in budbreak timing (Leolini *et al.*, 2018) but genetic control

of phenology is unknown in grapevine. Wild grapevine species that are frequently used

in breeding programs to increase cold hardiness have higher rates of deacclimation

during the spring when compared to *V. vinifera* (Londo and Kovaleski, 2017; Kovaleski

et al., 2018). Comparisons between *Vitis* species could help identify molecular drivers of

cold hardiness and deacclimation. In addition, the quantitative nature of deacclimation

also responds to chill accumulation (Kovaleski *et al.*, 2018). Therefore, understanding

the genetic regulation of cold hardiness loss could elucidate some of the mechanisms

that control dormancy in buds.

This study leverages phenological differences between species (*V. amurensis*, *V.*

riparia, *V. vinifera*) and between cultivars within species (*V. vinifera* 'Cabernet

Sauvignon' and 'Riesling') to uncover key aspects of gene regulation that occurs during

deacclimation. The objective was to determine the transcriptional regulation and

relationship between loss of cold hardiness and growth. We hypothesized that

genotypes with early spring budbreak phenology may have faster transcriptomic

responses tied to growth and development when placed under forcing conditions when

contrasted with slower phenology genotypes. Alternatively, differences between

genotypes with rapid or slow phenology could be due to acceleration of gene regulation

cascades, or acceleration of "bottleneck" genes which trigger specific regulatory

cascades. Additionally, we hypothesize that genes related to maintaining the cold hardy

and dormant phenotype will be quickly down-regulated in all genotypes; genes related

to deacclimation will either be transiently expressed or have a slow increase or

decrease, but mostly synchronized between genotypes; and genes related to growth

and budbreak will be staggered. Finally, we explore the use of ABA as a deacclimation

inhibitor.

Materials and Methods

Plant material

Canes of four genotypes of grapevine were used in this study: two wild species,

V. amurensis (PI588635) and *V. riparia* (PI588275) were collected from the USDA Plant

Genetic Resources Unit in Geneva, New York; and two cultivars of cultivated

grapevines (*V. vinifera*), 'Cabernet Sauvignon' and 'Riesling', from a local vineyard

(42.845N, 77.004W). The two locations are approximately 6 km apart. The three

species represent major clades in *Vitis*: American, Eurasian, and Asian (Fig. 1A). The *V.*

riparia accession was originally collected in North Dakota, USA (https://www.ars-

grin.gov/), while *V. vinifera* 'Cabernet Sauvignon' and 'Riesling' are cultivars

domesticated in France and Germany, respectively. *V. amurensis* is native to temperate

climates in East Asia, although no collection location information was available.

Dormant canes were collected in mid-winter [February 2015; 1030 chill units (North

Carolina model; Shaltout and Unrath, 1983)], and chopped into single node cuttings.

The cuttings were placed with cut ends in cups of water and placed within a growth

room with forcing conditions: 22 °C and 16/8h light/dark.

Loss of cold hardiness and budbreak

Measurement of bud cold hardiness was conducted following standard

differential thermal analysis (DTA) methods to detect lethal freezing temperature of

grapevine buds (Mills *et al.*, 2006). Buds were excised from canes, placed on

thermoelectric modules and subjected to decreasing temperatures at -4 °C hour^{-1}

(n=8). The release of heat that results from freezing of water inside the bud, the low

temperature exotherm (LTE), was recorded using a Keithley data logger (Tektronix,

Beaverton, OR). Cold hardiness was assessed over several days and rates were

determined using linear regression of LTE data. A subsample of 10 replicate single

node cuttings were left to fully deacclimate and reach budbreak. Phenology was visually

assessed, and budbreak was noted when buds reached stage 3 of the modified E-L

scale (Coombe and Iland, 2005).

RNAseq Libraries and data analysis

Three replicates of three buds each were collected directly into LN, daily for each

genotype, until buds reached E-L stage 3; corresponding to day 10, 12, 15, and 17 for

V. amurensis, *V. riparia*, 'Riesling', and 'Cabernet Sauvignon', respectively. Day 0

samples were field-collected. Total RNA was extracted using Sigma Spectrum kits

(Sigma-Aldrich, St. Louis, MO, USA), and strand-specific libraries were prepared

(Borodina *et al.*, 2011). Libraries were multiplexed (25-plex) and sequenced at 100 bp

single-end reads on HiSeq 2500 (Illumina, Inc., San Diego, CA, USA) at Cornell

University's Institute of Biotechnology (Ithaca, NY, USA).

Raw reads were de-multiplexed using FastX_BarcodeSplitter.pl (Gordon and

Hannon, 2010), quality trimmed using Cutadapt (Martin, 2011), aligned to the *V. vinifera*

12xV2 genome (https://urgi.versailles.inra.fr/Species/Vitis) and the V3 annotation

(Canaguier *et al.*, 2017) using STAR (Dobin *et al.*, 2013), and uniquely aligned reads

were quantified using HTSeq-count (Anders *et al.*, 2015). Sequence data are available

at NCBI-GEO (https://www.ncbi.nlm.nih.gov/geo/), series entry GSE124820.

Within each genotype, counts were normalized and grouped by days using the

edgeR library in R (ver. 3.3.0, R Foundation for Statistical Computing). Multidimensional

scaling and correlation plots were produced to visually verify outliers in libraries. After

outlier removal, remaining data were analyzed using the *DESeq2* library (Love *et al.*,

2014). In *DESeq2*, a full model containing a fourth order polynomial for time (t) as a

continuous variable and intercept, was compared to a null model of intercept only, using

likelihood ratio test. Genes with an adjusted-$P \leq 0.10$ [FDR correction (Benjamini and

Hochberg, 1995)] were used for subsequent filtering. Using parameter estimates

extracted from *DESeq2*, predicted normalized counts [counts per million (CPM)] were

calculated for the time span of each genotype's experiment (e.g., 0-10 days for *V.*

amurensis, 0-17 days for *V. vinifera* 'Cabernet Sauvignon', as integers). Considering

that counts for any given gene are a function of time (e.g.,

$CPM(t) = \beta_0 + \beta_1 t + \beta_2 t^2 + \beta_3 t^3 + \beta_4 t^4$), the filters used were: $max\{CPM(t) \mid 0 \leq t \leq t_f\} \geq 10$

(i.e., maximum predicted CPM at any time point ≥ 10), where t_f is the final time point for

any given genotype (e.g., for 'Cabernet Sauvignon', $t_f = 10$); and

$log_2(max\{CPM(t) \mid 0 \leq t \leq t_f\} / min\{CPM(t) \mid 0 \leq t \leq t_f\}) \geq 2$ (i.e., logFC≥ 2). Remaining

genes were considered relevant differentially expressed genes (DEGs).

Predicted logFC for each day calculated using *DESeq2* model estimates were

used for cluster analysis of DEGs using Cluster 3.0 (ver. 3.0, Human Genome Center,

University of Tokyo). Data were centered based on gene means, then on array means,

followed by normalization by genes and arrays. Twenty *k*-Means clusters were used for

genes, using Euclidian distances as the similarity metric, and 100 runs. The average

behavior of the genes placed in each cluster was visually inspected, and placed into 4

general categories: *up-regulated*, relative expression increasing over time; *down-*

regulated, relative expression decreasing over time; *transient* expression, peak

expression occurred within the duration of experiment; and *random*, where most of the

clusters had genes with a transient low expression within the duration of the experiment.

The resultant DEGs for each genotype were tested for pathway enrichment using

VitisPathways (Osier, 2016), which leverages biological pathways and molecular

functions in grapevine from the VitisNet database (Grimplet *et al.*, 2009). Enriched

pathways were determined using 1000 permutations, and a permuted-P <0.10. Pathway

enrichment analysis was conducted for each genotype using all DEGs, as well as DEGs

in the up-regulated, down-regulated, and transient expression groups. Enriched

pathways that were shared between genotypes were observed using Cytoscape (ver.

3.6.1; Shannon *et al.*, 2003). To plot trends in gene expression for pathways of interest,

genes predicted to encode or encoding the same protein were aggregated, and mean

expression was used for each day. Therefore, protein names are used as the notation

to refer to expression levels, with few exceptions (e.g., transcription factors). Qualitative

comparisons of expression behavior were done on a relative basis, using the maximum

mean counts within the duration of the experiment as 100% for each genotype.

ABA treatment of single node cuttings

Single node cuttings of *V. vinifera* 'Riesling' were prepared from material

collected from the same vineyard in 4 April 2017 [1630 chill units (Shaltout and Unrath,

1983)]. The cuttings were placed in a growth chamber with average temperature of 7 ºC

(2–7–12–7 ºC for 6h each) and a 0/24h light/dark photoperiod. ABA [ProTone® SG

(20% S-ABA), Valent BioSciences Corp., Libertyville, USA] treatments were applied as

the solution inside cups where cuttings were placed. Two concentrations were used,

1mM and 5mM, and a distilled water control. After 42 days, remaining cuttings were

washed in running water, the bottom ~1 cm of stems was cut, and cuttings were placed

in cups with water and moved to forcing conditions (22 °C, 16h/8h light/dark). Cold

hardiness was measured twice a week until 38 days, and three more sample dates of

unequal distance until day 56 (or two for the control). Phenology was evaluated in 5

buds for each treatment using the modified E-L scale (Coombe and Iland, 2005) until

buds reached stage 8 (2 separated leaves). Pairwise treatment comparisons were

evaluated for both cold hardiness (day 0 through 38) and phenology (day 27 through

56) within days and means were separated using Tukey's HSD test (α=0.01). A logistic

regression was used to compare when each treatment reached budbreak (E-L stage 3),

using the *drc* library (Ritz *et al.*, 2015).

Results

Deacclimation and budbreak (RNA-Seq)

Wild species had a higher initial cold hardiness upon collection (−29.8 and −28.0

°C for *V. amurensis* and *V. riparia*, respectively) compared to *V. vinifera* genotypes

(−23.4 and −23.9 °C for 'Cabernet Sauvignon' and 'Riesling', respectively; Fig. 1B).

Both wild species also had higher deacclimation rates than the two *V. vinifera*

genotypes (2.87, 2.81, 1.76, and 1.58 °C day^{-1} for *V. amurensis*, *V. riparia*, and *V.*

vinifera 'Cab. Sauvignon' and 'Riesling', respectively). Budbreak occurred latest and

sparsely on *V. vinifera* genotypes, starting at 13 days and spanning through day 17 for

'Riesling' and 'Cabernet Sauvignon' (Fig. 1B). Budbreak under forcing conditions

occurred after 8 days in *V. amurensis* and 10 days for *V. riparia*.

Analyses of Differential Gene Expression

The number of DEGs ranged from 8326 (*V. riparia*) to 9123 (*V. vinifera* 'Cabernet

Sauvignon'; Supplementary Fig. S1A). DEGs uniquely expressed in each species were

1348 (16.0%) for *V. amurensis*, 1006 (12%) for *V. riparia*, and 2232 (21.7%) for *V.*

vinifera ('Cabernet Sauvignon' + 'Riesling'). The majority of DEGs, 4551, were shared

among all four genotypes. Cluster analysis of shared DEGS grouped 1192 DEGs as

down-regulated, and 1065 DEGs as up-regulated (Supplementary Fig. S1B,C). Principal

components (PC) 1 and 2 generated through prcomp() [within pcaPlot() using

ntop=1000] separate global gene expression data points explaining 61% and 11% of

the variance, respectively, when using all genotypes. Plotting PC2 clearly separates *V.*

vinifera genotypes from *V. amurensis* and *V. riparia* (Fig. 2A,B). A third PC (*not shown*;

6% variance) separates *V. amurensis* from *V. riparia*. There appears to be a

discontinuity at the higher end of PC1 when points are compared by time, where the

variance is more closely related between the final timepoints in *V. vinifera* (15 d and 17

d for 'Riesling' and 'Cabernet Sauvignon', respectively) and the final timepoints in *V.*

amurensis (10 d) and *V. riparia* (12 d) (Fig. 2A). Time was transformed to a thermal time

measurement using deacclimation rate to normalize the response of each genotype to

temperature, therefore creating a thermal time unit of *Deacclimation Degree-Days* [*DDD*

$= k_{deacc} \times time\ (d)$], where k_{deacc} is the rate of deacclimation. Timepoint transformation to

DDD better scales the differences between genotypes in the distribution of PC1 (Fig.

2B). Global trends in gene expression are also similar across genotypes in respect to

phenology as observed in heatmap of gene expression (Supplementary Fig. **S2**). A

simple linear regression of PC1 by DDD results in an r^2=78.4% (Fig. 2C), further

demonstrating that the majority of the variation in gene expression comes from a time

component, and that differences between the genotypes are derived from their inherent

faster (wild species) or slower (cultivated species) response to temperature as

measured by the deacclimation rate.

Pathway enrichment analysis was performed with gene lists composed of all

DEGs to examine pathways where both up and down regulation may be occurring, but

also for those in the down-regulated, up-regulated, and transiently expressed DEGs

groups. There were 21 enriched pathways when all genes were queried, 22 for down-

regulated patterns, 15 for up-regulated patterns, and 5 for transient patterns that were

shared among all genotypes (Table 1). Overall, there were 43 enriched pathways that

were shared among all genotypes: 16 were defined as associated with Metabolism, four

with Environmental Information Processing, two with Cellular Processes, four with

Transport, and 17 with Transcription Factors. Important pathways identified included

many related to plant hormone production and signaling, sugar metabolism, and growth-

related processes like cell cycling and photosynthesis. Critical genes in these pathways

are reported below.

Within the ABA biosynthesis pathway, zeaxanthin epoxidase (ZEP), 9-cis-

epoxycarotenoid dioxygenase (NCED), and (ABA8'OH) were down-regulated while ABA

glucosidase (ABA BG), was strongly up-regulated (Fig. 3). ABA signaling genes, such

as *ABA-RESPONSIVE ELEMENT BINDING PROTEIN 2* (*AREB2*) and *KEEP ON*

GOING (*KEG*), were down-regulated. Timing of *AREB2* downregulation was

synchronized between genotypes, while the *KEG* was reduced faster in wild species

than in *V. vinifera*. Both *ABA INSENSITIVE 1* (*ABI1*) and *ABA REPRESSOR 1* (*ABR1*)

appear to have transient expressions. *MEDIATOR OF ABA-REGULATED DORMANCY*

1 (*MARD1*) and Phospholipase D (PLD) genes are both generally up-regulated.

Ethylene synthesis was transiently up-regulated within a few days of

deacclimation. 1-aminocyclopropane-1-carboxylate oxidase (ACC oxidase) and ACC

synthase had peak expression after placement in warm conditions, or a temporary up-

regulation (Fig. 4). Ethylene synthesis repressor gene *E8*, however, was up-regulated in

all genotypes. Ethylene signaling, which includes ERFs, WRKY, and HSF pathways

were primarily down-regulated (Supplementary Fig. S3, S4).

Key components of jasmonate and gibberellin (GA) synthesis pathways were

observed (Supplementary Fig. S5). Jasmonate synthesis was down-regulated, with

precursors being led to medium chain fatty acids via up-regulation of lipoxygenase.

MEJAE, which converts methyl-jasmonate to jasmonic acid, was down-regulated,

whereas Jasmonate-O-methyltransferase, which catalyzes the opposite reaction, was

up-regulated. For GAs, GA-3 β-dioxygenase is up-regulated in all genotypes, while GA-

2 β-dioxygenase is down-regulated, indicating a trend towards synthesis of bioactive

gibberellins. For Auxins, indole-3-acetic acid-amino acid hydrolases (IAA-AA-

hydrolases), which reactivates IAA, was up-regulated in all genotypes, reaching

maximum expression earlier in *V. amurensis* than other genotypes (Supplementary Fig.

S6). *IAA6*, *IAA19*, *AUXIN RESPONSE FACTOR 5* (*ARF5*) and *AUXIN-RESISTANT 1*

(*AUX1*) were up-regulated, with relative expression increasing faster in *V. amurensis*

than other genotypes. Auxin responsive *PIN-FORMED* (*PIN*) *1* was down-regulated

over time for all genotypes, while *PIN3* was up-regulated.

The degradation of cytokinin through cytokinin dehydrogenase (CKX) was up-

regulated for all genotypes over time (Fig. 5). In addition, *PURINE PERMEASE 1*, a

cytokinin transporter, was down-regulated. *AUTHENTIC RESPONSE REGULATOR*

(*ARR*) *11* was down-regulated, and *ARR17* was up-regulated, further indicating a

decrease in cytokinin content. Cyclin D, which is repressed by cytokinin responses and

is the first cyclin involved in the cell cycle, was up-regulated earlier compared to cyclins

A and B for all genotypes (Fig. 5). In fact, cyclins A and B have near 0 expression levels

before cyclin D is up-regulated. Genes related to cell expansion and cell division

(cyclins and cytoskeleton proteins) were up-regulated and staggered in accordance to

species: *V. amurensis*, followed by *V. riparia*, and *V. vinifera* (Fig. 5).

Circadian rhythm pathway genes were generally down-regulated in all genotypes

(Supplementary Fig. S7). *CIRCADIAN 1* (*CIR1*), *EARLY FLOWERING 3* (*ELF3*), and

ARABIDOPSIS PSEUDO-RESPONSE REGULATOR 5 (*APRR5*) had a strong down-

regulation in the first few days, whereas *CIRCADIAN CLOCK ASSOCIATED 1* (*CCA1*)

and Catalase *CAT3* reduced at a slower rate. In the phenylpropanoid biosynthesis

pathway, phenylalanine ammonia lyase was initially down-regulated, followed by up-

regulation that occurs earlier in *V. amurensis*, followed by *V. riparia*, and the *V. vinifera*

genotypes (Supplementary Fig. S8). Trihydroxystilbene synthase (stilbene synthase)

was quickly down-regulated in all genotypes, but faster in the wild species than *V.*

vinifera. Of the 44 predicted genes for stilbene synthase, between 29 and 35 were

DEGs for down-regulation in all genotypes.

Sugar metabolism genes encoding Starch and Glycogen [starch] synthases, and

beta amylase were down-regulated (Fig. 6). However, Glycogen [starch] synthase and

beta amylase are synchronized in all genotypes, whereas starch synthase reduces its

expression earlier in wild genotypes. Sucrose synthase is initially down-regulated in all

genotypes, while a return to high levels of expression is seen in *V. riparia* and *V.*

vinifera, but not in *V. amurensis*. Fructokinase and hexokinase are generally up-

regulated in all genotypes, leading sugars into glycolysis. Similarly, genes in fatty acid

biosynthesis and metabolism pathways were generally up-regulated over time

(Supplementary Fig. S9), including those involved in β-oxidation. *FATTY ACID*

DESATURASE 5 (*FAD5*), however, was strongly down-regulated for all genotypes.

In photosynthesis related pathways, genes encoding light harvesting complex I

(LHCI) and II (LHCII), and photosystem I (PSI) subunits also follow the trend of growth-

related genes, with earlier up-regulation in wild species (Supplementary Fig. S10).

RESPIRATORY BURST OXIDASE HOMOLOG PROTEIN B (*RBOHB*; all genotypes)

and *RBOHE* (all but *V. riparia*) had transient up-regulation in the earlier days. *RBOHF*

was also up-regulated in the first few days, returning to ~50% relative expression. Peak

expression of *RBOHF* occurs a few days prior to budbreak in *V. amurensis*, *V. riparia*,

and *V. vinifera* 'Cabernet Sauvignon'. *RBOHD* appeared to have an increasing

expression trend.

Vacuole localized aquaporins *TONOPLAST INTRINSIC PROTEIN* (*TIP*) *3;1* and

TIP3;2 were down-regulated, while *TIP1;3* and *TIP4;1* were up-regulated (Fig. 7).

CYCLIC NUCLEOTIDE-GATED ION CHANNEL PROTEIN 15 (*CNGC15*) was down-

regulated in all genotypes.

ABA effects on cold hardiness

ABA treatments had no effect on budbreak or deacclimation until 13 days post

application (Fig. 8A,B). Starting on day 16, ABA treatments prevented further loss of

cold hardiness, while deacclimation continued in the water control (Fig. 8B). Both ABA

concentrations maintained significantly more cold hardiness than the control, with no

difference amongst themselves. After ABA treatment was removed, previously treated

buds resumed deacclimation. Regarding phenology, control buds were significantly

more developed starting on day 44 in pairwise comparisons with ABA treated buds,

although buds in the control treatment average E-L stage > 1 starting on day 30 (Fig.

8A). Control buds reached budbreak (E-L stage 3) at 49.0±1.5 days after the beginning

of the experiment as estimated by a logistic regression. For 1mM ABA, budbreak

occurred at 73.2±3.3 days, while buds in 5mM did not resume growth before the

experiment ended at 84 days.

Discussion

Deacclimation and growth are processes that occur in tandem in grapevine,

although most or all of the cold hardiness is lost before growth is visible. As these

processes occur incrementally over many days, a novel approach for time-series data

analysis to find DEGs was used. To our knowledge, this is the first study exploring long-

term RNA-Seq data for the exploration of genes and pathways important for

deacclimation and budbreak. We used species with contrasting phenotypes to generate

hypotheses and explore the genes that are likely related to the cold hardy/dormant

phenotype, those related to growth and budbreak, and those that may be shared between both.

The approach developed for data analysis in this study was important to detect many trends in the long-term RNA-Seq data for differential expression. Here we treated time as a continuous variable, modeling counts with a fourth degree polynomial to allow transient expression of genes to be properly predicted, and using the predicted counts for filtering. Typically, studies will either use pairwise comparisons between adjacent timepoints (e.g., Meitha *et al.*, 2018) or between each timepoint and a 0h control (e.g., Fu *et al.*, 2018). The first approach may result in a low number of DEGs (Días-Riquelme *et al.*, 2012), while the second only accounts for changes in gene expression that are different from a "basal" condition. Differences in time between sample points [e.g., 3h-72h (Meitha *et al.*, 2018); 3h-120h (Fu *et al.*, 2018)] also argues that time may be more appropriate as a continuous variable.

While the approach in Meitha *et al.* (2018) would more likely detect genes with transient expression or genes with extreme changes in expression, and the approach of Fu *et al.* (2018) might miss transiently expressed genes with maximum and minimum expression occurring after the first time point, our approach makes no assumption as to the timing of significant expression change.

Sampling within a 24h period can result in the detection of "false" DEGs due to circadian oscillations in genes (Grundy *et al.*, 2015). In our case, samples were always taken within a 3h window. Irregular sampling could benefit from using time as a continuous variable and sine/cosine functions that may elucidate some of the variation in gene expression due to the circadian rhythm (Cronn *et al.*, 2017). Using continuous variables for time for RNA-Seq studies with more than 3 data points might be the most appropriate method, and even time-series data with lower replicate numbers [e.g., 2 biological replicates (Sozzani *et al.*, 2010)] could benefit from this type of analysis for more rigor than pairwise comparisons considering how degrees of freedom are distributed for continuous variables.

Our objective was to observe asynchronous behavior in gene expression relative to asynchronicity of the deacclimation pattern, therefore we analyzed the data in response to time within each genotype. It is clear that correcting gene expression data

by inherent phenological development rates may be necessary for other experiments

(Fig 2A,B,C). The results presented here demonstrate that not only does phenology in

grapevine correlate with deacclimation kinetics (Kovaleski *et al.*, 2018), so do the

patterns of gene expression that occur during this physiological transition. As wild

grapevines are used in cold hardy hybrid cultivar breeding, it is important to recognize

that these cold hardy genotypes also contribute faster deacclimation and budbreak

phenology (Fig 1A; Londo and Kovaleski, 2017; Kovaleski *et al.*, 2018) controlled by the

accelerated expression of the deacclimation cascade (Supplementary Fig. S2).

Temperature sensing must be the first step directing the physiology of the bud

towards growth. Changing the environment from field to growth chamber (–7 °C to 22

°C) between day 0 and day 1 resulted in rapid temperature sensing-related

transcriptional changes. Three lines of evidence for this rapid temperature sensing were

observed. Membrane rigidity has long been suggested as the most upstream

temperature sensor for thermal signaling (Horváth *et al.*, 1998, 2012; Saidi *et al.*, 2010;

Bahuguna and Jagadish, 2015). Our results agree with this hypothesis as fast down-

regulation of *FAD5* and general up-regulation of fatty acid biosynthesis genes

(Supplementary Fig. S9) likely lead to increased content of saturated fatty acids and

decreased membrane fluidity (Wallis and Browse, 2002; Falcone *et al.*, 2004; Filek *et*

al., 2017). CNGCs are also important in thermosensing as channel proteins for Ca^{2+}

signaling, and have been implicated in dormancy release of buds (Pang *et al.*, 2007;

Bahuguna and Jagadish, 2015). Although plasma membrane localized CNGCs have

clear roles in cold sensing (Bahuguna and Jagadish, 2015), here we observed down-

regulation of *CNGC15* [nucleus localized (DeFalco *et al.*, 2016)] was detected in all

genotypes. This suggests that in grapevine buds, Ca^{2+} stored in the nuclear envelope

lumen or endoplasmic reticulum may contribute to signaling during the dormant period.

Finally, PLD expression increased in response to dormancy release. PLD has been

observed to be up-regulated during cold and freezing stress (Ruelland *et al.*, 2002) as

well as many other stress conditions (Meijer and Munnik, 2003), and is indicated as a

signaling protein in thermal sensing (Bahuguna and Jagadish, 2015). PLD activity is

also positively influenced by Ca^{2+} influx during cold stress (Meijer and Munnik, 2003;

Wang and Nick, 2017). Overexpression of PLD results in higher freezing tolerance (Li *et*

al., 2004), and more rapid and sensitive response to ABA and stress (Sang *et al.*, 2001;

Meijer and Munnik, 2003). Considering our treatment can be seen as a release from

cold stress, but PLD expression increased, we speculate that effects of PLD in cold

stress-relief signaling are likely related to changes in plasma membrane composition

through selective hydrolysis of membrane phospholipids (Wang, 1999).

Following sensing of warm temperatures, we expected a decrease in cold-related

transcription factors would occur. Interestingly, none of the *CBF/DREB1* genes were

differentially expressed, and within *DREB1*s only *DREB1F/DDF2* was differentially

expressed. It is possible that CBFs are involved in cold hardiness gain and in response

to cold stress (Liu *et al.*, 2017a; Ban *et al.*, 2017; Rubio *et al.*, 2018), but are not

maintained in a high level of transcription throughout the winter. Alternatively, DREB2

homologs were differentially expressed and down-regulated. This result disagrees with

Liu *et al.* (1998) and Nguyen *et al.* (2017), who suggested *DREB1* and *DREB2* are two

independent *DREB* families that lead to signal transduction pathways for low

temperature and dehydration conditions, respectively**.** Consequently, when buds were

moved into forcing conditions, down-regulation of *CBF/DREB1*s transcripts was not

detected. Instead, lack of vascular continuity in early stages of budbreak (Xie *et al.*,

2018) may result in some level of bud drought stress, contributing to the DREB family

patterns observed.

Pathways and genes associated with stress were downregulated, such as ABA,

ethylene, ROS burst and detoxification, and circadian clock genes. ABA biosynthesis is

a requirement for cold acclimation and thermotolerance (Gilmour and Thomashow,

1991; Larkindale *et al.*, 2005; Penfield, 2008), and ABA is a clear antagonist of

deacclimation and growth. Both the ABA biosynthesis and the signaling pathways were

down-regulated in this study. The pivotal role of the ABA synthesis reduction for the loss

of hardiness is clear (Fig. 3). Both ZEP and especially NCED, the rate-limiting enzyme

in ABA biosynthesis (Grundy et al., 2015), were quickly down-regulated in all

genotypes, indicating carotenoids are likely no longer contributing to synthesis of new

ABA during deacclimation. ABR1, an AP2 TF repressor of ABA response (Pandey *et al.*,

2005), was quickly up-regulated from day 0 to the highest relative level of expression in

all genotypes at day 1, indicating the move to warm temperatures quickly signals for

reduction of ABA synthesis as well as repression of ABA responses from endogenous

pools. Supporting the role of ABA in reinforcing dormancy, high exogenous ABA

concentrations halted deacclimation (Fig. 8), although there was no net gain in

hardiness as in Rubio *et al.* (2018). ABA BG expression was upregulated while the rest

of the synthesis and signaling pathway was downregulated, suggesting that reactivation

of inactive vacuolar ABA may play a role in preventing dormancy release during

endodormancy. Given there was no lag in deacclimation response it is unlikely that

much vacuolar storage of inactive ABA remained when our study was conducted.

Future studies should separately examine the fates and concentrations of newly

synthesized ABA and reactivated ABA during deacclimation, and determine if these

separate pools could be leveraged as a biomarker for dormancy status. Another

important role of ABA during dormancy appears to be in controlling water status in the

bud, which is critical for low temperature survival (George and Burke, 1977). TIP3;1 and

TIP3;2 – ABA regulated aquaporins (Mao and Sun, 2015) – were down regulated as

opposed to the general up-regulation of TIP aquaporins (Fig. 7). While other aquaporins

have been implicated in chill stress responses (Bilska-Kos *et al.*, 2016), regulation of

TIP3s in our study demonstrates that ABA may be affecting water movement in the bud

during the dormant season, maintaining cold hardiness.

Steps leading to ethylene biosynthesis appear transiently up-regulated in the

initial days of warm period in this study. This may be due to quick loss of ABA's

antagonistic effect on the synthesis of ethylene. Ethylene, along with H_2O_2, lead to

activation of antioxidative stress genes in grapevine (Vergara *et al.*, 2012). H_2O_2 is

involved in the regulation of cold-responsive genes expression during acclimation

(Fedurayev *et al.*, 2018), during forced budbreak with hydrogen cyanamide (Or *et al.*

1999; Sudawan *et al.* 2016), and temporary increases in ROSs have been described to

precede budbreak (Ophir *et al.*, 2009; Pérez and Lira, 2005; Khalil-Ur-Rehman *et al.*,

2017; Meitha *et al.*, 2018), demonstrating contrasting roles. RBOHF is involved in

ethylene and ABA signaling through H_2O_2 production (Kwat *et al.*, 2003; Desikan *et al.*,

2006), and the expression pattern in our study appears to be a marker for budbreak.

While gene expression in relation to hypoxia and ethylene have been previously

described during budbreak (Ophir *et al.*, 2009; Meitha *et al.*, 2018), this study identified

a potentially novel role of stilbene synthase. Stilbene synthase is responsible for

resveratrol production in the phenylpropanoid pathway, which is an ROS scavenger.

The down-regulation of the majority of the 44 genes encoding stilbene synthase in all 4

genotypes after collection from the field suggests resveratrol production is an important

part of cold hardiness or dormancy maintenance. Stilbene synthase has previously

been described as up-regulated in grapevine leaves during cold or freeze stress (Londo

et al., 2018), but is also involved in other types of stress during the growing season

(Carvalho *et al.*, 2015). Similarly, studies examining cold tolerant *Passiflora edulis* (Liu

et al., 2017a) and ecodormant *Pyrus pyrifolia* (Bai *et al.*, 2013) support the hypothesis of

phenylpropanoids as cold hardiness and dormancy metabolites.

Regulation of circadian clock genes is important for abiotic stress tolerance

(Grundy *et al.*, 2015), and alternative splicing has been indicated as a mechanism of

signaling in cold response (Calixto *et al.*, 2016). Most genes in the circadian rhythm

pathway were down-regulated over time. Both *CCA1* and its promoter *CIR1* are down-

regulated in all genotypes. *CCA1* is a transcription factor that regulates low temperature

responses through control of *CBF* gene expression (Grundy *et al.*, 2015), and also

represses *TOC1* expression. Higher expression of *CCA1* in the field may be disrupting

of the circadian clock (Ramos *et al.*, 2005), but may also be required during dormancy

in order to maintain the circadian rhythm without strong effects of day length or light.

Growth and cell expansion pathways were upregulated rapidly and sequentially

after day 1. The trend towards synthesis of bioactive GA suggested by down-regulation

of GA-2 β-dioxygenase and up-regulation of GA-3 β-dioxygenase (Supplementary Fig.

S5) was also reported by Bai *et al.* (2013) comparing eco- and endodormant buds.

Those authors also observed an up-regulation of Jasmonate-O-methyltransferase as

seen here (Supplementary Fig. S5). This interplay between MEJAE and Jasmonate-O-

methyltransferase suggest that methyl-jasmonate is the preferred form of jasmonate

during budbreak.

Genes in carbohydrate metabolism likely have significant roles during the

dormancy period in grapevine buds (Khalil-Ur-Rehman *et al.*, 2017). While Starch and

Sucrose Metabolism pathway is enriched with down-regulated genes (Table **1**), and

enzymes in the degradation of starch are generally down-regulated, fructokinase and

hexokinase are both up-regulated (Fig. 6). This indicates that pools of sugar are being

used in glycolysis to generate energy, but also that sucrose used in sugar metabolism is

being sourced outside the bud. The up-regulation of Enoyl CoA hydratase and Acetyl

CoA c-Acyltransferase in the fatty acid metabolism (Supplementary Fig. S9) indicates

that energy is also being produced from free fatty acids such as those resulting from the

degradation of phospholipids by PLD (Wang, 1999).

The same temporal staggering observed for the budbreak phenotype was

expected to be reflected in gene expression patterns for genes related to growth. This

correlation is clearly observed in the up-regulation of cyclins and cytoskeleton proteins

in all genotypes (Fig. 5), following observed timing of budbreak (Fig. 1B). Within

genotypes, earlier expression of cyclin D indicates that the G_1 and S phases of the cell

cycle (Mironov *et al.*, 1999) occur concomitantly to deacclimation. However, the initially

very low expression of both cyclin A and B indicates that G_2 and M phases only occur

after cold hardiness is lost, suggesting deacclimation and budbreak are cell expansion-

but not cell division-dependent, and expression of these genes may be used as markers

for imminent budbreak.

What determines the phenotype difference between fast and slow

deacclimators? It appears that both sensing of the stimulus for dormancy release

(change in environment) and down-regulation of cold-hardiness- and dormancy-

maintaining genes occurred simultaneously between all genotypes. This is observed in

the synchronous down-regulation of *CNGC15*, *FAD5*, ABA synthesis, and stilbene

synthase, among others. Differences between the four genotypes in both phenotypes

observed (deacclimation and budbreak) must therefore arise from how quickly

genotypes are able to re-establish growth. This is exemplified in the staggering of

expression seen in cyclins, cytoskeleton- and auxin-related genes. Many players have

been implicated in the control of growth, but here we indicate those that may also play a

role in deacclimation, or promote growth through promoting deacclimation. We thus

propose a hypothetical model for events occurring during release of dormancy in

grapevine buds (Fig. 9). Upon sensing of growth permissive temperatures, likely by

membrane fluidity, ABA synthesis is down-regulated. With the release of inhibition by

ABA, a temporary increase in ethylene synthesis and signaling leads to a transient burst

of ROS. An influx of water and cell expansion slowly increase water potentials,

decreasing cold hardiness. At the same time, energy is generated from fatty acids and

cane-stored carbohydrates, and growth hormone pathways are up-regulated. Ultimately,

with cell expansion the appearance of budbreak occurs, along with the start of cell

division. As tissues emerges from buds, photosynthesis-related proteins are ready to

start generate energy – although the sink to source switch in breaking buds occurs

beyond the time analyzed here.

Understanding physiological transitions such as deacclimation and budbreak are

essential for improving crop sustainability in the face of changing climate. We identify

important genes and pathways related to maintaining cold hardiness, the loss of cold

hardiness (deacclimation), the release of dormancy and budbreak by exploring

phenotypic or temporal differences between *Vitis* spp. genotypes. It is likely that

processes described here are conserved in other perennial woody species which show

similar budbreak behavior in response to chilling [e.g., *Prunus persica* (Fan *et al.*,

2010)]. In this study, we describe a new way to deal with variable length, time-series

data such that a single list of DEGs is produced from all time-points. In our discussion

we demonstrate that molecular processes involved in gain of cold hardiness are not just

in reverse mode during deacclimation (e.g., PLD regulation, CBFs). ABA and its

signaling were confirmed to be closely downstream from temperature sensing, and

upstream from what defines cold hardiness with exogenous ABA. Therefore, ABA has

the potential for mitigation of unseasonal or early deacclimation and budbreak in

warmer winters. Finally, we discovered differences in deacclimation rate and initial cold

hardiness contribute to variation in budbreak phenology between genotypes through

differential gene regulation. Implications of this association may directly link to spring

phenology markers and improved budbreak prediction in agricultural conditions as they

experience climate variation.

Supplementary Data

Fig. S1. Venn diagrams of shared genes.

Fig. S2. Heatmap of top 1000 genes.

Fig. S3. WRKY transcription factors.

Fig. S4. HSF transcription factors.

Fig. S5. Jasmonate and gibberellin biosynthesis pathways.

Fig. S6. Auxin biosynthesis pathway and auxin-related genes.

Fig. S7. Circadian rhythm pathway.

Fig. S8. Phenylpropanoid biosynthesis pathway.

Fig. S9. Fatty acids synthesis and metabolism.

Fig. S10. Photosynthesis-related genes.

Acknowledgements

This work was partially supported by CAPES, Coordenação de Aperfeiçoamento

de Pessoal de Nível Superior, Brazil, award number 12945/13-7, and by U.S.

Department of Agriculture appropriated project 1910-21220-006-00D.

Table 1 Enriched VitisNet pathways shared among *V. amurensis*, *V. riparia*, and

V. vinifera 'Cabernet Sauvignon' and 'Riesling' during deacclimation and budbreak.

Pathway enrichment analysis was conducted in VitisPathways using all differentially

expressed genes (DEGs), as well as DEGs in the up-regulated, down-regulated, and

transient expression groups.

All DEGs	Down-regulated	Up-regulated	Transient
Fatty acid biosynthesis	Galactose metabolism	Fatty acid biosynthesis	Fatty acid metabolism
Photosynthesis antenna proteins	Tyrosine metabolism	Photosynthesis	Anthocyanin biosynthesis
Tyrosine metabolism	Glutathione metabolism	Photosynthesis antenna proteins	Auxin signaling
Zeatin biosynthesis	Starch and sucrose metabolism	Flavonoid biosynthesis	Ethylene signaling
ABA signaling	Inositol phosphate metabolism	Cell cycle	Auxin transport
Auxin Transport	Carotenoid biosynthesis	Regulation of actin cytoskeleton	
Transport electron carriers	Nitrogen metabolism	Auxin transport	
Thylakoid targeting pathway	Phenylpropanoid biosynthesis	Transport electron carriers	
AP2-EREBP	ABA biosynthesis	Thylakoid targeting pathway	
AUXIAA	ABA signaling	Porters cat7 to 17	
BZIP	Ethylene signaling	BHLH	
C2C2-DOF	Circadian rhythm	C2C2-GATA	
C2C2-GATA	AP2-EREBP	C2C2-YABBY	
C2C2-YABBY	C2C2-DOF	MYB	
GRAS	DBP	OFP	
HSF	GRAS		
NAC	HSF		
TCP	MYB		
Orphans zf-b box	NAC		
OFP	WRKY		
	Orphans zf-b box		
	Other zf-C3HC4		

References

References

Alleweldt G, Spiegel-Roy P, Reisch B. 1990. Grapes (*Vitis*). In: Moore JN,
Ballington JR, eds. Genetic resources of temperate fruit and nut crops. Wageningen,
NL. Acta Hort **290**, 291–337.

Anders S, Pyl PT, Huber W. 2015. HTSeq--a Python framework to work with
high-throughput sequencing data. Bioinformatics **31**, 166–169.

Bahuguna RN, Jagadish KSV. 2015. Temperature regulation of plant
phenological development. Environmental and Experimental Botany **111**, 83–90.

Bai S, Saito T, Sakamoto D, Ito A, Fujii H, Moriguchi T. 2013. Transcriptome
analysis of Japanese pear (*Pyrus pyrifolia* Nakai) flower buds transitioning through
endodormancy. Plant and Cell Physiology **54**, 1132–1151.

**Ban Q, Wang X, Pan C, Wang Y, Kong L, Jiang H, Xu Y, Wang W, Pan Y, Li
Y, *et al.*** 2017. Comparative analysis of the response and gene regulation in cold
resistant and susceptible tea plants. PLOS ONE **12**, e0188514.

Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical
and powerful approach to multiple testing. Journal of the Royal Statistical Society.
Series B **57**, 289–300.

Bilska-Kos A, Szczepanik J, Sowiński P. 2016. Cold induced changes in the
water balance affect immunocytolocalization pattern of one of the aquaporins in the
vascular system in the leaves of maize (*Zea mays* L.). Journal of Plant Physiology **205**,
75–79.

Birkenbihl RP, Diezel C, Somssich IE. 2012. Arabidopsis WRKY33 is a key
transcriptional regulator of hormonal and metabolic responses toward *Botrytis cinerea*
infection. Plant Physiology **159**, 266–285.

Borodina T, Adjaye J, Sultan M. 2011. Chapter five–a strand-specific library
preparation protocol for RNA sequencing. Methods in enzymology. Academic Press.
v.**500**, 79–98.

Burke MJ, Gusta LV, Quamme HA, Weiser CJ, Li PH. 1976. Freezing and
Injury in Plants. Annual Review of Plant Physiology **27**, 507–528.

Busov V, Carneros E, Yakovlev I. 2016. EARLY BUD-BREAK1 (EBB1) defines

a conserved mechanism for control of bud-break in woody perennials. Plant Signaling

and Behavior **11**, e1073873.

Calixto CPG, Simpson CG, Waugh R, Brown JWS. 2016. Alternative splicing

of barley clock genes in response to low temperature. PLOS ONE **11**, e0168028.

Canaguier A, Grimplet J, Di Gaspero G, Scalabrin S, Duchêne E, Choisne N,

Mohellibi N, Guichard C, Rombauts S, Le Clainche I. 2017. A new version of the

grapevine reference genome assembly (12X.v2) and of its annotation (VCost.v3).

Genomics Data **14**, 56–62.

Carvalho LC, Vidigal P, Amâncio S. 2015. Oxidative stress homeostasis in

grapevine (*Vitis vinifera* L.). Frontiers of Environmental Science and Engineering in

China **3**, 20.

Cook B, Wolkovich EM. 2016. Climate change decouples drought from early

wine grape harvests in France. Nature Climate Change **6**, 715–720.

DOI:10.1038/NCLIMATE2960.

Cooke JEK, Eriksson ME, Junttila O. 2012. The dynamic nature of bud

dormancy in trees: environmental control and molecular mechanisms: Bud dormancy in

trees. Plant, Cell and Environment **35**, 1707–1728.

Coombe BG, Iland P. 2005. Grapevine phenology. In Dry PR, Coombe BG, eds.

Viticulture, vol 1: Resources. Winetitles, Ashford, Australia, 210–248.

Cronn R, Dolan PC, Jogdeo S, Wegrzyn JL, Neale DB, St. Clair JB, Denver

DR. 2017. Transcription through the eye of a needle: daily and annual cyclic gene

expression variation in Douglas-fir needles. BMC Genomics **18**, 558. DOI

10.1186/s12864-017-3916-y

DeFalco TA, Moeder W, Yoshioka K. 2016. Opening the gates: Insights into

cyclic nucleotide-gated channel-mediated signaling. Trends in Plant Science **21**, 903–

906.

Desikan R, Last K, Harrett-Williams R, Tagliavia C, Harter K, Hooley R,

Hancock JT, Neill SJ. 2006. Ethylene-induced stomatal closure in Arabidopsis occurs

via AtrbohF-mediated hydrogen peroxide synthesis. The Plant Journal **47**, 907–916.

DOI: 10.1111/j.1365-313X.2006.02842.x

Díaz-Riquelme J, Grimplet J, Martínez-Zapater JM, Carmona MJ. 2012.
Transcriptome variation along bud development in grapevine (*Vitis vinifera* L.). BMC
Plant Biology **12**, 181.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P,
Chaisson M, Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner.
Bioinformatics **29**, 15–21.

van Dyk MM, Soeker MK, Labuschagne IF, Rees DJG. 2010. Identification of a
major QTL for time of initial vegetative budbreak in apple (*Malus x domestica* Borkh.).
Tree genetics and genomes **6**, 489–502.

Falcone DL, Ogas JP, Somerville CR. 2004. Regulation of membrane fatty acid
composition by temperature in mutants of Arabidopsis with alterations in membrane lipid
composition. BMC Plant Biology **4**, 17.

Fan S, Bielenberg DG, Zhebentyayeva TN, Reighard GL, Okie WR, Holland
D, Abbott AG. 2010. Mapping quantitative trait loci associated with chilling requirement,
heat requirement and bloom date in peach (*Prunus persica*). New Phytologist **185**, 917–
930. DOI: 10.1111/j.1469-8137.2009.03119.x

FAO. 2018. FAOSTAT. Food and Agriculture Organization of the United Nations,
Statistics Division. <http://www.fao.org/faostat/> (Accessed 18 Dec 2018)

Fedurayev PV, Mironov KS, Gabrielyan DA, Bedbenov VS, Zorina AA,
Shumskaya M, Los DA. 2018. Hydrogen peroxide participates in perception and
transduction of cold stress signal in *Synechocystis*. Plant and Cell Physiology **59**, 1255–
1264.

Filek M, Rudolphi-Skórska E, Sieprawska A, Kvasnica M, Janeczko A. 2017.
Regulation of the membrane structure by brassinosteroids and progesterone in winter
wheat seedlings exposed to low temperature. Steroids **128**, 37–45.

Fu JJ, Geng JC, Miao YJ, Xu YM, Hu TM, Yang PZ. 2018. Transcriptomic
analyses reveal genotype- and organ-specific molecular responses to cold stress in
Elymus nutans. Biologia plantarum **62**, 671–683.

George MF, Burke MJ. 1977. Cold hardiness and deep supercooling in xylem of
shagbark hickory. Plant Physiology **59**, 319–325.

Gilmour SJ, Thomashow MF. 1991. Cold acclimation and cold-regulated gene

expression in ABA mutants of *Arabidopsis thaliana*. Plant Molecular Biology **17**, 1233–

1240.

Gordon A, Hannon GJ. 2010. Fastx-toolkit. *FASTQ/A short-reads preprocessing*

tools (unpublished). <http://hannonlab. cshl. edu/fastx_toolkit> (Accessed 26 Dec 2018)

Grimplet J, Cramer GR, Dickerson JA, Mathiason K, Van Hemert J, Fennell

AY. 2009. VitisNet: "omics" integration through grapevine molecular networks. PLOS

ONE **4**, e8365.

Grundy J, Stoker C, Carré IA. 2015. Circadian regulation of abiotic stress

tolerance in plants. Frontiers in Plant Science **6**, 648.

Hedley PE, Russell JR, Jorgensen L, Gordon S, Morris JA, Hackett CA,

Cardle L, Brennan R. 2010. Candidate genes associated with bud dormancy release in

blackcurrant (Ribes nigrum L.). BMC Plant Biology **10**, 202.

Horváth I, Glatz A, Nakamoto H, Mishkind ML, Munnik T, Saidi Y,

Goloubinoff P, Harwood JL, Vigh L. 2012. Heat shock response in photosynthetic

organisms: membrane and lipid connections. Progress in Lipid Research **51**, 208–220.

Horváth I, Glatz A, Varvasovszki V, Török Z, Páli T, Balogh G, Kovács E,

Nádasdi L, Benkö S, Joó F, *et al.* 1998. Membrane physical state controls the

signaling mechanism of the heat shock response in *Synechocystis* PCC 6803:

Identification of hsp17 as a "fluidity gene". Proceedings of the National Academy of

Sciences **95**, 3513–3518.

Khalil-Ur-Rehman M, Wang W, Xu Y-S, Haider MS, Li C-X, Tao J-M. 2017.

Comparative Study on Reagents Involved in Grape Bud Break and Their Effects on

Different Metabolites and Related Gene Expression during Winter. Frontiers in Plant

Science **8**, 1340.

Kolstad EW, Breiteig T, Scaife AA. 2010. The association between

stratospheric weak polar vortex events and cold air outbreaks in the Northern

Hemisphere. Quarterly Journal of the Royal Meteorological Society **136**, 886–893.

Kovaleski AP, Reisch BI, Londo JP. 2018. Deacclimation kinetics as a

quantitative phenotype for delineating the dormancy transition and thermal efficiency for

budbreak in *Vitis* species. AoB PLANTS **10**, ply066. DOI: 10.1093/aobpla/ply066

Kwak JM, Mori IC, Pei ZM, Leonhardt N, Torres MA, Dangl JL, Bloom RE,

Bodde S, Jones JDG, Schroeder JI. 2003. NADPH oxidase *AtrbohD* and *AtrbohF*

genes function in ROS-dependent ABA signaling in *Arabidopsis*. The EMBO Journal **22**,

2623–2633.

Lang GA, Early JD, Martin GC, Darnell RL. 1987. Endo-, Para-, and

Ecodormancy: physiological terminology and classification for dormancy research.

HortScience **22**, 371–377.

Larkindale J, Hall JD, Knight MR, Vierling E. 2005. Heat stress phenotypes of

Arabidopsis mutants implicate multiple signaling pathways in the acquisition of

thermotolerance. Plant Physiology **138**, 882–897.

Leolini L, Moriondo M, Fila G, Costafreda-Aumedes S, Ferrise R, Bindi M.

2018. Late spring frost impacts on future grapevine distribution in Europe. Field Crops

Research **222**,197-208. DOI: 10.1016/j.fcr.2017.11.018

Li S, Dami IE. 2016. Responses of *Vitis vinifera* 'Pinot gris' Grapevines to

Exogenous Abscisic Acid (ABA): I. Yield, Fruit Quality, Dormancy, and Freezing

Tolerance. Journal of Plant Growth Regulation **35**, 245–255.

Li W, Li M, Zhang W, Welti R, Wang X. 2004. The plasma membrane-bound

phospholipase Ddelta enhances freezing tolerance in *Arabidopsis thaliana*. Nature

biotechnology **22**, 427–433.

Liu Q, Kasuga M, Sakuma Y, Abe H, Miura S, Yamaguchi-Shinozaki K,

Shinozaki K. 1998. Two transcription factors, DREB1 and DREB2, with an EREBP/AP2

DNA binding domain separate two cellular signal transduction pathways in drought- and

low-temperature-responsive gene expression, respectively, in Arabidopsis. The Plant

Cell **10**, 1391–1406.

Liu S, Li A, Chen C, Cai G, Zhang L, Guo C, Xu M. 2017a. De novo

transcriptome sequencing in *Passiflora edulis* Sims to identify genes and signaling

pathways involved in cold tolerance. Forests, Trees and Livelihoods **8**, 435.

Liu J, Xu Y, Zhang L, Li W, Cai Z, Li F, Peng M, Li F, Hu B. 2017b. De novo

assembly and analysis of the transcriptome of *Rumex patientia* L. during cold stress.

PLOS ONE **12**, e0186470.

Londo JP, Kovaleski AP. 2017. Characterization of wild North American

grapevine cold hardiness using differential thermal analysis. American Journal of

Enology and Viticulture **68**, 203–212.

Londo JP, Kovaleski AP, Lillis JA. 2018. Divergence in the transcriptional

landscape between low temperature and freeze shock in cultivated grapevine (*Vitis*

vinifera). Horticulture Research **5**, 10.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome Biology **15**, 550.

Mao Z, Sun W. 2015. Arabidopsis seed-specific vacuolar aquaporins are

involved in maintaining seed longevity under the control of ABSCISIC ACID

INSENSITIVE 3. Journal of Experimental Botany **66**, 4781–4794.

Martin, M. 2011. Cutadapt removes adapter sequences from high-throughput

sequencing reads. EMBnet. journal **17**, 10–12. DOI: 10.14806/ej.17.1.200

Meijer HJG, Munnik T. 2003. Phospholipid-based signaling in plants. Annual

Review of Plant Biology **54**, 265–306.

**Meitha K, Agudelo-Romero P, Signorelli S, Gibbs DJ, Considine JA, Foyer

CH, Considine MJ**. 2018. Developmental control of hypoxia during bud burst in

grapevine. Plant, Cell and Environment **41**, 1154–1170.

Mills LJ, Ferguson JC, Keller M. 2006. Cold-hardiness evaluation of grapevine

buds and cane tissues. American Journal of Enology and Viticulture **57**, 194–200.

Mironov V, De Veylder L, Van Montagu M, Inzé D. 1999. Cyclin-dependent

kinases and cell division in plants— the nexus. The Plant Cell **11**, 509–521.

**Nguyen HC, Cao PB, San Clemente H, Ployet R, Mounet F, Ladouce N,

Harvengt L, Marque C, Teulieres C**. 2017. Special trends in CBF and DREB2 groups

in Eucalyptus gunnii vs Eucalyptus grandis suggest that CBF are master players in the

trade-off between growth and stress resistance. Physiologia Plantarum **159**, 445–467.

**Olukolu BA, Trainin T, Fan S, Kole C, Bielenberg DG, Reighard GL, Abbott

AG, Holland D**. 2009. Genetic linkage mapping for molecular dissection of chilling

requirement and budbreak in apricot (*Prunus armeniaca* L.). Genome **52**, 819–828.

Ophir R, Pang X, Halaly T, Venkateswari J, Lavee S, Galbraith D, Or E. 2009.

Gene-expression profiling of grape bud response to two alternative dormancy-release

stimuli expose possible links between impaired mitochondrial activity, hypoxia, ethylene-ABA interplay and cell enlargement. Plant Molecular Biology **71**, 403–423.

Or E, Nir G, Vilozny I. 1999. Timing of hydrogen cyanamide application to grapevine buds. Vitis **38**, 1–6.

Osier MV. 2016. VitisPathways: gene pathway analysis for V. vinifera. Vitis: Journal of Grapevine Research **55**, 129–133.

Pandey GK, Grant JJ, Cheong YH, Kim BG, Li L, Luan S. 2005. ABR1, an APETALA2-domain transcription factor that functions as a repressor of ABA response in Arabidopsis. Plant Physiology **139**, 1185–1193.

Pang X, Halaly T, Crane O, Keilin T, Keren-Keiserman A, Ogrodovitch A, Galbraith D, Or E. 2007. Involvement of calcium signalling in dormancy release of grape buds. Journal of Experimental Botany **58**, 3249–3262.

Penfield S. 2008. Temperature perception and signal transduction in plants. New Phytologist **179**, 615–628.

Pérez FJ, Lira W. 2005. Possible role of catalase in post-dormancy bud break in grapevines. Journal of Plant Physiology **162**, 301–308.

Prentice IC, Cramer W, Harrison SP, Leemans R, Monserud RA, Solomon AM. 1992. A global biome model based on plant physiology and dominance, soil properties and climate. Journal of Biogeography **19**, 117–134.

Ramos A, Pérez-Solís E, Ibáñez C, Casado R, Collada C, Gómez L, Aragoncillo C, Allona I. 2005. Winter disruption of the circadian clock in chestnut. Proceedings of the National Academy of Sciences USA **102**, 7037–7042.

Ritz C, Baty F, Streibig JC, Gerhard D. 2015. Dose-response analysis using R. PLOS ONE **10**, e0146021. DOI: 10.1371/journal.pone.0146021

Rubio S, Noriega X, Pérez FJ. 2018. Abscisic acid (ABA) and low temperatures synergistically increase the expression of CBF/DREB1 transcription factors and cold-hardiness in grapevine dormant buds. Annals of Botany **XX**, 1–9. DOI: 10.1093/aob/mcy201

Ruelland E, Cantrel C, Gawer M, Kader J-C, Zachowski A. 2002. Activation of phospholipases C and D is an early response to a cold exposure in Arabidopsis suspension cells. Plant Physiology **130**, 999–1007.

Saidi Y, Peter M, Finka A, Cicekli C, Vigh L, Goloubinoff P. 2010. Membrane

lipid composition affects plant heat sensing and modulates Ca^{2+}-dependent heat shock

response. Plant Signaling and Behavior **5**, 1530–1533.

Sang Y, Zheng S, Li W, Huang B, Wang X. 2001. Regulation of plant water loss

by manipulating the expression of phospholipase Dα: Phospholipase D in plant water

loss. The Plant Journal **28**, 135–144.

Shaltout AD, Unrath CR. 1983. Rest completion prediction model for

'Starkrimson Delicious' apples. Journal of the American Society for Horticultural Science

108, 957–961.

Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N,

Schwikowski B, Ideker T. 2003. Cytoscape: a software environment for integrated

models of biomolecular interaction networks. Genome Research **13**, 2498–2504.

Sozzani R, Cui H, Moreno-Risueno MA, Busch W, Van Norman JM, Vernoux

T, Brady SM, Dewitte W, Murray JAH, Benfey PN. 2010. Spatiotemporal regulation of

cell-cyle genes by SHORTROOT links patterning and growth. Nature **466**, 128–132.

DOI:10.1038/nature09143

Sudawan B, Chang C-S, Chao H-F, Ku MSB, Yen Y-F. 2016. Hydrogen

cyanamide breaks grapevine bud dormancy in the summer through transient activation

of gene expression and accumulation of reactive oxygen and nitrogen species. BMC

Plant Biology **16**, 202.

Tylewicz S, Petterle A, Marttila S, Miskolczi P, Azeez A, Singh RK, Immanen

J, Mähler N, Hvidsten TR, Eklund DM, *et al.* 2018. Photoperiodic control of seasonal

growth is mediated by ABA acting on cell-cell communication. Science **360**, 212–215.

Vergara R, Parada F, Rubio S, Pérez FJ. 2012. Hypoxia induces H_2O_2

production and activates antioxidant defence system in grapevine buds through

mediation of H_2O_2 and ethylene. *Journal of Experimental Botany* **63**, 4123–4131.

Wallis JG, Browse J. 2002. Mutants of Arabidopsis reveal many roles for

membrane lipids. Progress in Lipid Research **41**, 254–278.

Wan Y, Schwaninger HR, Baldo AM, Labate JA, Zhong GY, Simon CJ. 2013.

A phylogenetic analysis of the grape genus (*Vitis* L.) reveals broad reticulation and

concurrent diversification during Neogene and quaternary climate change. BMC

Evolutionary Biology **13**, 141. DOI: 10.1186/1471-2148-13-141

Wang X. 1999. The role of phospholipase D in signaling cascades. Plant

Physiology **120**, 645–651.

Wang L, Nick P. 2017. Cold sensing in grapevine-Which signals are upstream of

the microtubular 'thermometer'. Plant, Cell and Environment **40**, 2844–2857.

Wu R, Wang T, Warren BAW, Allan AC, Macknight RC, Varkonyi-Gasic E.

2017. Kiwifruit SVP2 gene prevents premature budbreak during dormancy. Journal of

Experimental Botany **68**, 1071–1082.

Xie Z, Forney CF, Bondada B. 2018. Renewal of vascular connections between

grapevine buds and canes during bud break. Scientia Horticulturae **233**, 331–338.

Zheng C, Halaly T, Acheampong AK, Takebayashi Y, Jikumaru Y, Kamiya Y,

Or E. 2015. Abscisic acid (ABA) regulates grape bud dormancy, and dormancy release

stimuli may act through modification of ABA metabolism. Journal of Experimental

Botany **66**, 1527–1542.

Figure Legends

Fig. 1. Distribution, cold hardiness and deacclimation of four *Vitis* spp. genotypes. (a)

Distribution map of the centers of diversity for *Vitis* spp. in gray – American, Eurasian,

and Asian (Redrawn from Alleweldt *et al.*, 1990; Wan *et al.*, 2013). Colored area

denotes origin for each genotype used: *V. vinifera* 'Cabernet Sauvignon' and 'Riesling'

were domesticated in France and Germany, respectively; *V. amurensis* (PI588635) is

native to temperate climates in East Asia; and *V. riparia* (PI588275) was originally

collected from the wild in North Dakota, USA. Orange dot shows where genotypes are

cultivated and were collected from (43N, 77W). (b) Cold hardiness, indicated by

symbols and predicted linear deacclimation for each genotype. Arrow heads (*V.*

amurensis and *V. riparia*) or trapezoid (*V. vinifera*) at 0 °C line indicate timing of

budbreak (E-L stage 3; Coombe and Iland, 2005). Deacclimation rates were 2.87, 2.81,

1.76, and 1.58 °C day^{-1} for *V. amurensis*, *V. riparia*, and *V. vinifera* 'Cabernet

Sauvignon' and 'Riesling', respectively.

Fig. 2. Principal components (PCs) for differential expression of four *Vitis* spp.

genotypes. Principal components were generated using prcomp() and variance

stabilized normalized counts. Distribution of genotypes and data points in PC1 and PC2,

with points colored by (a) time in days and (b) thermal time in deacclimation degree-

days [$DDD = k_{deacc} \times time$, where k_{deacc} is the rate of deacclimation]. (c) Linear

association of PC1 and DDD ($r^2 = 78.4\%$).

Fig. 3. Reduced VitisNet ABA biosynthesis and signaling pathway. Mini-graphs indicate

relative level of expression of any given protein for *V. amurensis* (red), *V. riparia*

(yellow), and *V. vinifera* 'Cabernet Sauvignon' (purple) and 'Riesling' (green) during

deacclimation and budbreak. Multiple DEGs encoding for the same protein had their

expression aggregated. Multiple arrows indicate transition or signaling steps omitted.

Detail shows expanded graph for axis information.

Fig. 4. Reduced VitisNet ethylene biosynthesis and signaling pathway, and ethylene

response factors (ERFs). Mini-graphs indicate relative level of expression of any given

protein for *V. amurensis* (red), *V. riparia* (yellow), and *V. vinifera* 'Cabernet Sauvignon'

(purple) and 'Riesling' (green) during deacclimation and budbreak. Multiple DEGs

encoding for the same protein had their expression aggregated. Arrows indicate

transition; circular arrows indicate positive regulation; and "T" indicates negative

regulation. For details on graph axes and symbols see Fig. **3**.

Fig. 5. Reduced VitisNet cytokinin biosynthesis and signaling pathway, and growth

associated genes. Mini-graphs indicate relative level of expression of any given protein

for *V. amurensis* (red), *V. riparia* (yellow), and *V. vinifera* 'Cabernet Sauvignon' (purple)

and 'Riesling' (green) during deacclimation and budbreak. Multiple DEGs encoding for

the same protein had their expression aggregated. Arrows indicate transition; circular

arrows indicate positive regulation; and "T" indicates negative regulation. Multiple

arrows indicate transition or signaling steps omitted. For details on graph axes and

symbols see Fig. **3**.

Fig. 6. Reduced VitisNet starch, sucrose, and galactose metabolism pathways. Mini-

graphs indicate relative level of expression of any given protein for *V. amurensis* (red),

V. riparia (yellow), and *V. vinifera* 'Cabernet Sauvignon' (purple) and 'Riesling' (green)

during deacclimation and budbreak. Multiple DEGs encoding for the same protein had

their expression aggregated. Arrows indicate transition. Multiple arrows indicate

transition steps omitted. For details on graph axes and symbols see Fig. **3**.

Fig. 7. Relative expression of channel protein genes during deacclimation and budbreak

in *V. amurensis* (red), *V. riparia* (yellow), and *V. vinifera* 'Cabernet Sauvignon' (purple)

and 'Riesling' (green).

Fig. 8. Cold hardiness and budbreak of *V. vinifera* 'Riesling' buds treated with abscisic

acid (ABA). (a) Stage of development (E-L number; Coombe and Iland, 2005) and (b)

cold hardiness of *V. vinifera* 'Riesling' buds in 0, 1, and 5 mM abscisic acid solution at

an average temperature of 7 °C 0/24h light/dark photoperiod. After 42 days (vertical

dotted line) samples were moved to 22 °C 16/8h light/dark photoperiod. Error bars

represent standard deviations of the mean.

Fig. 9. Schematic representation of proposed molecular control of deacclimation and

growth based on gene expression data from this study. ABA appears as a master

regulator of cold hardiness and dormancy maintenance, inhibiting commonly known

routes that promote budbreak. Arrows indicate transition; circular arrows indicate

positive regulation; and "T" indicates repression. Solid lines indicate validated

interactions. Role of ABA in repression of deacclimation observed here highlighted in

blue. Dashed lines are new interactions supported by gene expression data.

Figure 1

Figure 2

977
978

Figure 3

Figure 4

Figure 5

983
984

Figure 6

Glycogen starch synthase

UDP-Glucose

ADP-Glucose

Sucrose synthase

Starch synthase

Amylose

UDP-Galactose

α-amylase

β-amylase

Inositol-3-α galactosyltransferase

Dextrin

Maltose

Galactinol sucrose galactosyltransferase

Starch

Sucrose

Rafinose

β-Fructose

α-D-Glucose

Fructokinase

Hexokinase

β-Glucose-6-P

α-Glucose-6-P

Glycolysis

985
986

Figure 7

Figure 8

Figure 9