Social hierarchy position in female mice is associated with

plasma corticosterone levels and hypothalamic gene

expression

Cait M. Williamson[1], Won Lee[1], Alexandra R. Decasien[2,3], Alesi Lanham[1], Russell

D. Romeo[4] & James P. Curley[1, 5*]

[1] Department of Psychology, Columbia University, New York, NY, 10027, USA

[2] Department of Anthropology, New York University, New York, NY 10003, USA

[3] New York Consortium in Evolutionary Primatology, New York, NY, 10024, USA

[4] Department of Psychology, Barnard College, New York, NY, 10027, USA

[5] Department of Psychology, UT Austin, Austin, Texas, 78712, USA

* Corresponding author: curley@utexas.edu

Abstract

Social hierarchies emerge when animals compete for access to resources such as food, mates or physical space. Wild and laboratory male mice have been shown to develop linear hierarchies, however, less is known regarding whether female mice have sufficient intrasexual competition to establish significant social dominance relationships. In this study, we examined whether groups of outbred CD-1 virgin female mice housed in a large vivaria formed social hierarchies. We show that females use fighting, chasing and mounting behaviors to rapidly establish highly directionally consistent social relationships. Notably, these female hierarchies are less linear, steep and despotic compared to male hierarchies. Female estrus state was not found to have a significant effect on aggressive behavior, though dominant females had elongated estrus cycles (due to increased time in estrus) compared to subordinate females. Plasma estradiol levels were equivalent between dominant and subordinate females. Subordinate females had significantly lower levels of basal corticosterone compared to dominant females. Analyses of gene expression in the ventromedial hypothalamus indicated that subordinate females have elevated ERα, ERβ and OTR mRNA compared to dominant females. This study provides a methodological framework for the study of the neuroendocrine basis of female social aggression and dominance in laboratory mice.

Introduction

The contextual and neurobiological factors that influence male intrasexual aggression and social dominance have been well-studied across species[1–6]. Conversely, female aggression and social dominance have been relatively understudied, with most work focused on maternal aggression expressed by females when they are pregnant or during the early postpartum period where the behavioral focus is on maternal defense of offspring[7,8]. Few studies have investigated the contextual and neurobiological factors that influence female-female aggression outside of reproduction in rodents[9,10].

Social hierarchies are likely to emerge whenever there is competition between individuals for resources such as food, water, territory or access to mates[11]. The more intense this competition is, the more likely it is that a highly linear social hierarchy will develop. In mammals, male social hierarchies are common as inter-sexual competition is typically dramatically higher in males compared to females, though there are notable exceptions such as hyenas where females have high levels of intra-sexual conflict and form strong female hierarchies[11]. Female hierarchies have also been observed in other species that have female intrasexual competition for access to resources including degus[12], bison[13], caribou[14], red deer[15], vervet monkeys[16], and chimpanzees[17]. Less is known about the formation of social hierarchies in female wild mice, though some population studies suggest that females do generate some form of social hierarchy with dominant aggressive females establishing territories and subordinate females being unable to do so when population sizes increase[18,19]. Conversely, when population density is very low it appears that female wild mice have relatively little intra-sexual competition and do not form hierarchies[20]. Female-female aggression also appears to be low if females have social experience with each other prior to the intra-sexual competition[21,22]. In laboratory mice, it has been documented that females in small groups of up to five can establish social ranks based on social interactions[23] or performance in the tube-test[24,25].

Previously, we have explored the complex group dynamics and neurobiology of male social hierarchies, demonstrating that male outbred CD-1 mice living in groups of up to 30 individuals will form highly linear social hierarchies when living in a large laboratory-based vivarium[5,26,27]. As relatively little is known about whether large groups of non-reproductively active female mice will form social hierarchies in the laboratory, we aimed to explore this question by housing groups of 12 virgin outbred CD-1 female mice in large vivaria. One historical reason why female behavior is so vastly understudied in comparison to male behavior in laboratory rodents is due to concerns that female behavior is more variable than males due to fluctuations in steroid hormone levels across the female estrus cycle. Indeed, estrus state has been shown to influence many behavioral states including anxiety-like behavior and exploration[28], motivation, addiction[29] and fear[30]. In rodents, some species also show variation in aggressive behavior across the estrous cycle[31]. Female California deer mice[32], rats[33,34] and hamsters[35–38] are less likely to show aggressive behavior during estrus than diestrus, although other studies have found no effect of estrus state on aggressive behavior[39–42]. There is also mixed evidence for estrous effects on aggression in female house mice.[43] [44]. Given the potential significance of estrous state on female dominance and subordinate behaviors, we examined whether the estrous state of females is associated with the frequency of aggressive behavior within social hierarchies.

The neurobiological basis of female intrasexual aggression among non-reproductive females is receiving
increased attention though much less is still known compared to male intrasexual aggression[9,10]. As in
males, brain regions in the social behavior network (medial amygdala (meA), bed nucleus of the stria
terminalis (BNST), lateral septum (LS), medial preoptic area (mPOA), anterior hypothalamus (AH),
ventromedial hypothalamus (VMH) and periaqueductal grey (PAG)) as well as the mesocorticolimbic
dopamine pathway have been found to form the basis of the neural circuit regulating aggression, though
there are some important sex differences[10]. In particular, it is well-established that the VMH is a key
modulator of aggression in non-reproductive female rodents, as the VMH is activated following aggressive
encounters between female California mice[32]. Electrical stimulation[45] or lesions[46,47] of the VMH have been
shown to increase aggression, whereas estradiol or progesterone implants into the VMH decrease
aggression[48–50]. Recently, distinct populations of estrogen-receptor alpha (ERα) expressing neurons in the
ventrolateral VMH have been shown to be critical for the expression of aggression[51]. In females, the mPOA
is recognized as being critical for the expression of parental behaviors and recently has been shown to be
involved in the expression of maternal aggression[52]. This region also plays a role in regulating non-
reproductive female aggression. Lesions to the mPOA decrease aggression and increase submissive
behavior in hamsters[53] and direct injection of progesterone or oxytocin into the mPOA inhibits female
aggression[54,55].

Estrogen-sensitive neuropeptide targets are key modulators of social behavior. In the brain, estrogen acts
at both ERα and estrogen-receptor beta (ERβ), which are expressed throughout the social behavior
network, including the VMH and mPOA[56]. Activation of these receptors results in the triggering of second
messenger systems that can modulate cell function[10,57,58]. Estradiol, the major estrogen steroid hormone,
has been primarily associated with promoting aggressive behaviors[59–61] but is also involved in many facets
of social behavior, such as social learning and social recognition[62]. Estradiol acts to alter the expression of
gene products in the hypothalamus, including progesterone receptors (PR), oxytocin receptors (OTR),
opioid receptors, and gonadotropin-releasing hormone (GnRH). These genes have been demonstrated to
regulate social behavior[63]. Progesterone acting at progesterone receptors has been shown to modulate
social recognition, with progesterone treatment impairing social recognition memory[64]. In females,
studies have found that oxytocin inhibits aggression, and genetic deletion of oxytocin results in increased
aggressive behavior toward other females[65]. More generally, oxytocin has been implicated in the
motivation to affiliate, perhaps acting to ameliorate aggressive tendencies[66]. Opioid receptors, specifically
mu-opioid receptor (OPRM1), have been shown to be involved in motivation to initiate social contact,
with OPRM1 knockout pups showing decreased motivation to seek out their mothers[67] and OPRM1
knockout males failing to exhibit typical social exploration behaviors[68]. Further, OPRM1 allele variations
are associated with differences in sociability and social dominance[69].

The current study used an established paradigm developed in our lab applied to the study of male social
hierarchy dynamics to study female social hierarchy behavior and begin to disentangle underlying
neurobiological and neuroendocrine mechanisms. We investigated the hierarchical structure of eight
groups of twelve females as well as plasma corticosterone and plasma estradiol concentrations for
dominant and subordinate mice within these hierarchies. In male mice, social stressors such as social
defeat and social stability are known to lead to increases in basal levels of corticosterone but findings from
female mice are more variable[70,71]. We have previously found that subordinate male mice have elevated

corticosterone levels compared to dominant male mice but only if the dominant males are highly despotic.
We predicted that female subordinate mice may also show higher levels of basal corticosterone compared
to dominant females, although previous studies have not found a consistent relationship between social
dominance and plasma corticosterone in female mice[23,72]. We further examined gene expression
differences between dominant and subordinate individuals in the VMH and mPOA of the hypothalamus
across six genes known to modulate various aspects of social behavior and moderated by the action of
estrogen: ERα, ERβ, PR, OTR, OPRM1, and GnRH. The aim of this work was to establish a feasible
methodology for the study of female aggression and dominance, as well as their neurobiological and
neuroendocrine mechanisms, outside of the reproductive period in laboratory mice.

Results

Hierarchy measures and organization

Sociomatrices of win-loss data for all female groups as well as the emergence of individual dominance
ranks over time are presented in **Figure 1**. Summary statistics of several aspects of the hierarchical
structure of each group are provided in **Table 1**. We found that seven out of the eight social groups formed
a significantly linear and steep social hierarchy as measured by modified Landau's *h',* triangle transitivity
and steepness. All eight social groups had significantly high directional consistency of agonistic behavior,
indicating that the majority of wins were directed from more dominant to more subordinate individuals.
Female groups had relatively low despotism values, indicating that alpha females were not exerting
complete dominance over all other females. This interpretation was confirmed by moderate Gini
Coefficient values for wins, demonstrating that the number of wins made by dominant females was fairly
evenly distributed between the top ranked females. Initial body weight measured at the beginning of
group housing was not related to final dominance rank in any group (Spearman Rank correlation tests: all
p>0.12).

Emergence of hierarchies over time

Despite having no previous social experience with each other, 4 of 8 social groups had a significantly linear
hierarchy by the end of Day 1 that continued throughout the 14-day observation period (**Figure 2**). Two
further social groups were significantly linear by the end of Day 2 and thereafter. One cohort did not have
a stable linear hierarchy until Day 9, although this hierarchy did show significant triangle transitivity on
Days 1 and 4 suggesting some linearity in the first week of co-housing. The social group that did not have
a significant hierarchy by the end of the observation period did have some linear organization, having a
significantly linear hierarchy on Day 3.

Sex differences in mouse social hierarchies

Female social hierarchies were significantly different from male social hierarchies in several aspects of
their dominance organization (see Methods about male comparison group). Female social hierarchies
were significantly less linear by triangle transitivity (W=14, p<0.05) and had lower directional consistency
(W=0, p<0.001) than male social hierarchies (**Figure 3**). The distribution of power was more even in
females than in males. Alpha females were significantly less despotic than alpha males (W = 1, p<0.001),

and the Gini coefficient of wins (W=0, p<0.001) was significantly lower in female hierarchies than male
hierarchies. There was no difference in the steepness of hierarchies (W=31, p=0.460), the Gini coefficient
of losses (W=32, p=0.515.) or Landau's modified h' value (W=23, p=0.146).

Frequency of each agonistic behavior over the group housing period

149   The hourly rate of each agonistic behavior is shown in **Figure S2**. Fighting and chasing were observed at
150   significantly higher rates than mounting. The rate of fighting behavior showed a significant decrease by
151   day over the group housing period (b_{day}= -0.62 [-0.89, -0.36]) while those of chasing and mounting
152   behaviors did not show significant changes by day (chasing: b_{day}= 0.08 [-0.29, 0.44]; mounting: b_{day}= 0.10
153   [-0.03, 0.23]).

Relationships between mounting, chasing and fighting

To determine if the directionality of fighting, chasing and mounting was consistent between individuals
we performed QAP correlation tests on the fighting, chasing and mounting sociomatrices (**Figure S2**).
Fighting and chasing sociomatrices were highly correlated across all cohorts (data presented as median
[IQR] across all eight cohorts: r=0.79 [0.75, 0.80], all p<0.001). Chasing and mounting sociomatrices were
also significantly correlated across all cohorts (r=0.44 [0.38, 0.54], all p<0.025). Fighting and mounting
sociomatrices were correlated for 7/8 cohorts (r=0.37[.35,.39], all p<.025 for significant correlations). See
supplemental **Figure S2** for individual sociomatrices.

Directional consistency of each agonistic behavior

163   All groups showed highly significant directionally consistent behavior for all behaviors (all p<0.001).
164   Notably, the most directionally consistent behavior was mounting behavior (median [IQR] = 0.87 [0.83,
165   0.93]), which was more consistent than chasing (0.75 [0.70, 0.77]) and significantly more consistent than
166   fighting (0.72 [0.69, 0.74]) (**Figure S3A**; Friedman's test X^2=7,df=2, p=0.03; post-hoc test mounting vs
167   fighting p= 0.033, mounting vs chasing p= 0.112).

Frequency of each agonistic behavior across individuals and ranks

As expected, the Gini coefficient for the total number of attacks and chases made by the females in each
group was moderately high, indicating that a few females are responsible for a disproportionate number
of these aggressive acts (**Figure S3B and S3C**). Unexpectedly, the Gini coefficient for mounting other
females was even higher than for fighting or chasing across cohorts (Friedman Test: X^2 = 12, df=2, p=0.002;
post hoc tests p<0.01). This analysis demonstrates that in each social group a small number of females are
responsible for a very large proportion of all mounting acts. The Gini coefficient for the total number of
attacks and chases received by each female in each social group was moderately low. This finding indicates
that most females are the recipients of fights and chases and these events are relatively evenly distributed
across females in each social group. Again, it was unexpected that the Gini coefficient of mounts received
was significantly higher than that for receiving the other two behaviors (Friedman Test: X^2 = 9.2, df=2,
p=0.01; post hoc tests p<0.05). This finding indicates that being mounted is far more unequally distributed
across females in each group than being chased or being attacked: certain females in each social group
are the targets for a disproportionately high number of mounts from other females.

We further examined if there is an effect of social rank on the hourly occurrence rate of each agonistic
behavior either given or received. Fighting and chasing, but not mounting, were exhibited at significantly
higher hourly rates by more dominant females (**Figure 4**, fighting given: b_{rank}= -1.11 [-1.44, -0.81]; chasing
given: b_{rank}= -1.13 [-1.46, -0.85]; mounting given: b_{rank}= -0.25 [-0.61, 0.12]). Similarly, there were significant
effects of social rank on the hourly rate of receiving fighting and chasing but not mounting, showing more
subordinate females received more fighting and chasing behaviors (fighting given: b_{rank}= -0.38 [0.19, 0.58];
chasing given: b_{rank}= 0.47 [0.25, 0.71]; mounting given: b_{rank}= 0.10 [-0.20, 0.42]). Although animals of each
rank differed in the absolute frequencies of each agonistic behavior used, there was no effect of rank on
the proportion of each behavior used (**Figure S4**). That is, all animals use each behavior proportionally
equivalently.

Estrus cycle and female behavior

The median (+/- IQR) proportion of time that individuals were in each estrous state were - estrus: 33.3%
[25.8%, 46.7%]; proestrus: 21.4% [13.3%, 36.4%], diestrus: 14.3% [0.08%, 23.1%], metestrus: 21.4%
[14.3%, 28.6%] (**Figures. 5A and 5B**). Female mice with higher social rank were more likely to be in estrus
compared to metestrus or proestrus (log-odds of being in estrus vs. metestrus: 0.48 [0.02, 0.95]; estrus
vs. proestrus: 0.36 [018, 1.57]). Mice did not significantly differ by social rank in the likelihood to be in
estrus compared to diestrus (estrus vs. diestrus: 0.50 [-0.03, 1.06]), or between other states (metestrus
vs. proestrus: 0.32 [-0.36, 1.05]; metestrus vs. diestrus: 0.02 [-0.57, 0.58]; proestrus vs. diestrus: -0.22 [-
0.78, 0.35]).

Overall there was no large effect of daily estrus state on the hourly rate of giving or receiving aggression.
There was small effect that mice showed a higher hourly rate of giving aggression when in metestrus
compared to proestrus ($b_{metestrus-proestrus}$: 0.19 [0.005, 0.35]). We further examined the effect of daily estrus
state on the hourly rate of giving or receiving each agonistic behavior individually (fighting, chasing,
mounting). For the rate of giving chases, mice had a higher rate when they are in metestrus compared to
diestrus and proestrus ($b_{metestrus-diestrus}$: 0.22 [0.03, 0.40]; $b_{metestrus-proestrus}$: 0.19 [0.01, 0.36]). Compared to
during diestrus, mice received higher rates of mounting when they were in estrus or metestrus ($b_{estrus-}$
$_{diestrus}$: 0.31 [0.09, 0.53]; $b_{metestrus-diestrus}$: 0.33 [0.08, 0.58]). The hourly rates of other behaviors given or
received did not differ across different estrus states.

Plasma corticosterone and estradiol levels

Plasma corticosterone levels were found to be significantly higher for subordinate individuals as compared
to dominant individuals (**Figure 6A**, $b_{subordinate-dominant}$: 153.9 ng/ul [107.6, 200.0]). We did not find an effect
of estrus cycle state on corticosterone level. There was no effect of dominant-subordinate status on
plasma estradiol levels (**Figure 6B**, $b_{subordinate-dominant}$: -2.97 pg/ul [-8.03, 1.99]). Mice that were in diestrus
state on the day of blood collection had significantly higher estradiol levels compared to those that were
in metestrus or proestrus ($b_{diestrus-metestrus}$: 6.79 pg/ul [0.29, 13.2]; $b_{diestrus-proestrus}$: 10.1 pg/ul [3.07, 17.5]).
There was no effect of estrus state on the estradiol levels among other states.

Gene expression in the VMH and the mPOA

219   In the VMH, there was small but significant effect of dominant-subordinate status on the levels of
220   expression in the VMH of ERα, ERβ, and OTR genes (**Figure 7**), with subordinate mice showing higher
221   expression than dominants. There were no significant differences in expression levels of OPRM1 and PR
222   genes in the VMH. There were no significant differences between dominant and subordinate individuals
223   in the expression any of the genes examined in the mPOA.

225   Discussion

Here we show that female mice living in social groups of up to 12 females are capable of forming
significantly linear dominance hierarchies that are stable for up to 14 days. Seven out of eight female
social groups had significantly linear hierarchies, as measured by Landau's h-value and triangle transitivity,
and significantly steep hierarchies as measured by the relative differences in David's scores. All eight
female social groups had significantly high directional consistency of agonistic behavior demonstrating
that dominant individuals more often won competitive interactions against subordinate individuals.
Similar to males, female mammals can form stable dominant-subordinate relationships whenever there
is intrasexual competition for resources[23,24,73–75]. In our study of virgin female mice, there is no intrasexual
competition for access to mates or food and water, however, female mice may compete for access to
preferred areas of the vivaria (such as nestboxes) or bedding. Our results also extend the findings from
studies of round-robin tube-tests conducted in groups of 5 or 8 female mice that report that females were
capable of forming linear social hierarchies based on wins and losses in that competitive exclusion test[24,25].
These findings are also consistent with studies of wild mice that have shown that females will form
territorially based social hierarchies when living in large environments with relatively high population
densities[18,19]. It appears that there may be a critical population density limit required for there to be
sufficient intrasexual competition for females to engage in sufficient aggression to establish hierarchies,
as one recent study found that groups of six females living in enclosures of $7m^2$ did not form social
hierarchies[20].

Our results demonstrate that females use fighting, chasing and mounting behaviors to establish and
maintain dominance relationships. The highest rates of fighting were observed during the establishment
of hierarchies (which typically occurred rapidly within the first two days), and the rate of fighting
significantly decreased thereafter. The rates of chasing and mounting did not decrease across the housing
period. Unsurprisingly, given the existence of a linear hierarchy, higher ranked individuals exhibited higher
rates of fighting and chasing than lower ranks. Similarly, lower-ranked individuals received higher amounts
of each of these behaviors. With respect to mounting behavior, we did not observe a significant overall
effect of rank on the rate of giving or receiving mounting. Indeed, there has been some controversy as to
the proximal function of female-female mounting in rodents and whether it should be considered to be a
dominance behavior, a sexual behavior, a masculinized behavior or something else[76–78]. Our data suggest
that mounting is being used by some females as a dominance behavior but not consistently by all
individuals. That is, females are more likely to mount females subordinate to themselves but not all
females engage in this behavior. We observed that the group sociomatrices for fighting, chasing and
mounting behaviors given and received all highly correlated with each other, indicating that the direction

and magnitude of these behaviors within each social relationship was consistent. Secondly, we observed
that the directional consistency of mounting behavior (0.87) was even stronger than the directional
consistency for fighting (0.72) and chasing (0.75). This finding indicates that mounting behavior
consistently occurred 87% of the time in the direction of more dominant to more subordinate females.
We also noted that females of all ranks, despite having different absolute rates of each behavior, used
each of the three behaviors in roughly equal proportions -suggesting that higher- or lower-ranked females
do not preferentially utilize mounting behavior over the other two as a dominance behavior. Additionally,
the Gini coefficient of behavior given and received was significantly higher for mounting as compared to
the other two agonistic behaviors. Thus, in each social group the majority of mounts given and received
were by specific individuals, and these interactions occurred in a directionally consistent manner.

Female mouse hierarchies exhibit several differences when compared to male social hierarchies. Although
both sexes produce linear hierarchies, those of females are less linear and have lower directional
consistency than male hierarchies. Male hierarchies tend to be despotic, with one alpha male exhibiting
the vast majority of all aggressive acts[5,26,27] . We did not observe such despotism in female hierarchies as
evidenced by the lower despotism scores and lower Gini coefficient of wins. More dominant females in
social groups tend to more equally distribute aggression towards other lower-ranked females. These sex
differences may be rooted in the ancestral biology of the house mice. In the wild, there are fundamental
differences in the reproductive strategies of males and females. Males have high levels of intersexual
competition and have high reproductive skew[79]. Females generally exhibit higher parental investment in
offspring and have reduced conflict over reproductive opportunities[75,80,81]. Indeed, even when given the
opportunity to compete for males, female mice do not necessarily increase their rates of aggression[20].
Across mammals, female adaptations for intrasexual competition can involve more subtle behaviors such
as low-level persistent aggression instead of overt displays of physical aggression[75]. Our data suggest that
in laboratory mice there persists low-level aggression that results in females forming stable social
hierarchies even with unlimited access to food, water, space and nesting material and no access to males.

We found that the estrous state of each female did not have a large effect on the likelihood of females
giving or receiving aggression. Females in all states were equally likely to bite or be bitten. However, we
did observe a very small effect for females in metestrus to have a higher rate of chasing than those females
in proestrus or diestrus. Previous work in rodents has suggested that estrus state can influence the
propensity of females to engage in intrasexual aggression although these effects are inconsistent and
appear to be highly influenced by many other contextual factors. For instance, female mice have been
reported to show higher aggression in the resident-intruder test during metestrus and proestrus
compared to estrus and distrus[43]. Similarly, female California mice show their highest aggression during
diestrus compared to proestrus and estrus[32]. Such a decline in aggression during estrus has been
suggested to facilitate mating, yet female mice in estrus have been shown to be capable of exhibiting
aggression and forming dominant-subordinate relationships[82]. Conversely, no changes in female-female
aggression across the estrous cycle have also been observed in rats and hamsters[41,42]. Although there was
no effect of estrous state on the likelihood to mount other females, we did find that female mice in estrus
or metestrus received significantly more mounts than those females in diestrus. This finding is somewhat
consistent with evidence from several species including baboons[83], squirrel monkeys[84], hanuman
langurs[85], rabbits[86] and rats[77] that mounted females tend to be subordinates in estrus.

We observed that group-housed females had extended estrous cycles with prolonged periods of diestrus
and estrus. The mouse estrus cycle is regulated by luteinizing hormone and follicle-stimulating hormone
released from the pituitary in response to gonadotrophin releasing hormone released from the
hypothalamus, which is in turn under control from estrogen and progesterone. The duration of this cycle
is usually 4-6 days for females housed in isolation or in pairs, but can be much longer for group-housed
females – an effect known as the Lee-Boot effect[87–90]. For instance, female mice living in groups of 8 have
been shown to have estrous cycles lasting up to 14 days[87] and females in groups of 30 have estrus cycles
up to 40 days in duration[90]. These cycles are typically extended due to longer periods of time spent in
diestrus, but they may also become disrupted. Following exposure to males, females quickly enter estrus
and become sexually receptive[90]. Our results are largely congruent with these earlier studies in that we
do observe extended estrous cycles in group-housed animals. Notably, many females in our study were in
estrous for prolonged periods of time, and this effect was larger in dominant females compared to
subordinate females. It is possible that this is an adaptive mechanism by which dominant females ensure
more or earlier mating opportunities than subordinate females, however it is unclear as to what
physiological processes underpin the extended length of estrous cycles we observed.

We found no significant differences between dominant and subordinate mice in plasma estradiol.
Peripheral estrogens are known to promote aggression in both males and females[10,91,92], and therefore
we predicted there would be higher estradiol levels in dominant females. However, our hormone samples
were taken at the end of group housing on Day 14 at a time when the social hierarchies had stabilized and
aggression was at a low level. Further, estrogens can have many, sometimes opposite, effects, depending
on where in the brain and on what receptors they are acting[62,93]. It is therefore perhaps not surprising that
estradiol levels found in plasma at one time point do not differ between dominant and subordinate
individuals.

Conversely, we did find that subordinate females had significantly higher levels of plasma corticosterone
than dominant females. This effect was moderately large and much larger than the effect that we had
previously observed in male hierarchies, where significant differences in plasma corticosterone levels
between dominants and subordinates only occur in highly despotic hierarchies[27]. In females, all
hierarchies were considered to be very low on the despotism scale. These results are interesting in the
context of the established literature on sex differences in stress responses. Several studies have shown
that male mice show a much more robust physiological response to stressors such as social defeat, social
instability and chronic stress compared to female mice[70,71]. However, these results are not always
consistent, and sometimes females do show increases in corticosterone depending on the social context
of the stressor[94–96]. It has also been reported that highly aggressive territorial females living in large groups
have elevated corticosterone compared to non-aggressive females[72], though one other study found
results consistent to ours that subordinate female mice had higher corticosterone than dominant females
in small groups of up to five albino mice[23]. The functional significance of the elevated corticosterone in
subordinate females living in our relatively stable social housing remains to be addressed, but these data
suggest the potential for studying female social hierarchies as a model of social stress.

The neurobiology of female intrasexual aggression has been relatively under-studied compared to that of
male intrasexual aggression, although this area has started to receive increased attention[9,10]. Here we

found that the expression of ERα, ERβ and OTR was moderately higher in subordinate female mice
compared to dominant female mice in the VMH. The VMH is known to be a critical regulator of female
aggression. Early studies in rats and hamsters demonstrated that lesions of the VMH lead to increased
aggression by females[47,59], whereas implants into the VMH with either estradiol or progesterone reduce
aggression[42,49,97]. More recently in mice, estrogen-receptor expressing neuronal populations in the
ventrolateral VMH have been found to become active when females bite versus mount other females[51].
This work builds on previous studies that demonstrated an important role for central estrogen receptors
in regulating male and female aggression. ERα knockout male mice show reduced aggression towards
other males[98], whereas females lacking ERα expression are more aggressive to other females[99,100]. Loss of
ERβ has also been associated with increased aggression in males[101,102]. Central administration of selective
ERα agonists to female mice increases aggressive attacks in a resident-intruder paradigm[103], while
selective ERβ agonists treatment leads to an increase in non-aggressive social behaviors[104]. The role of
each of these receptors in coordinating female aggressive behavior is clearly complex and context-
dependent, but it is possible that our observed increased VMH expression in both ERα and ERβ in
subordinate mice may underlie their inhibition of aggression. However, given the multitude of functions
of these receptors in this region it is possible that these differences may be unrelated to aggression and
associated with other social behaviors[105].

Likewise, oxytocin acting on oxytocin receptors has a range of effects on social behavior in females, and
its roles in promoting or inhibiting aggression are highly contextually dependent. In non-lactating females
oxytocin generally appears to inhibit aggression[10]. OT knockout mice show reduced aggression to each
other[65] and central or injections of OT into the MPOA or AH can reduce aggression[54,106]. These findings
may be congruent with our finding that subordinate females have higher VMH OTR expression than
dominant females. Notably, we did not observe any association between social status and mRNA levels of
OTR or any other gene in the MPOA. Though some lesion studies in other rodents have reported a role
for the MPOA in female aggressive behavior[53], our results would suggest that plasticity in gene expression
in response to an individual's social status occurs primarily in the VMH.

Conclusion

In the present study, we establish that outbred CD-1 female mice living in groups of 12 individuals are
capable of forming significant linear hierarchies. These hierarchies are linear and directionally consistent,
but less despotic than male social hierarchies. These hierarchies emerge quickly and are stable over 14
days and are relatively unaffected by the estrous cycle. All group-housed females also show an extended
estrous cycle, and dominants spend longer in estrus than subordinate females. Subordinate females have
significantly higher levels of plasma corticosterone than dominant females, suggesting that subordinate
females are more susceptible to the social stress of group living than male mice. We also find that
subordinate females have higher levels of ERα, ERβ, and OTR mRNA than dominant females in the VMH,
suggesting that these genes in this region may facilitate in part the reduced aggression displayed by these
females. This work furthers our understanding of group female social behavior, begins to explore sex
differences between male and female social hierarchy formation and maintenance and provides evidence
that the actions of estrogen may play a role in modulating female social hierarchy behavior.

Methods

Subjects and housing

A total of 96 female outbred CD1 mice were obtained from Charles River Laboratories at 7 weeks of age. Mice were housed in the animal facility in the Department of Psychology at Columbia University, with constant temperature (21-24°C) and humidity (30-50%). The room was kept on a 12/12 light/dark cycle, with white light (light cycle) on at 2400 hours and red lights (dark cycle) on at 1200 hours. All mice were uniquely marked by dying their fur with a blue, nontoxic animal marker (Stoelting Co.), enabling individuals to be identified throughout the study. These marks remain for up to 12 weeks and only require one application. All procedures were conducted with approval from the Columbia University Institutional Animal Care and Use Committee (IACUC Protocol No. AC-AAAP5405).

Social behavior observations

Following arrival at the animal facility, mice were housed in groups of 3 for 2 weeks in standard sized cages. At 9 weeks of age, groups of 12 mice were weighed and placed into large, structurally complex, custom built vivaria (length 150cm, height 80cm, width 80cm; Mid-Atlantic; **Figure S5**). The vivaria were constructed as described in Williamson et al.[26]. Each vivarium consists of an upper level constructed of multiple shelves connected by plastic tubes and covered in pine bedding and a lower level comprised of 5 interconnected standard sized cages filled with pine bedding and connected by a system of plastic tubes. Mice can access all levels of the vivarium at any time through this interconnecting system of ramps and tunnels. Standard chow and water were provided ad libitum on the top level of the vivarium. Social groups were created such that in each group of 12 females, each individual had previous social experience with maximum only one other individual and at least 6 females per group had absolutely no experience with any of the other individuals. Mice were placed in the vivarium at the onset of the dark cycle on Day 1 of the experiment and were observed by trained observers for 2 hours directly following introduction to the group and for 2 hours each day for the next two weeks (Day 1 – Day 14). Observations always occurred during the dark cycle at some point during the first 6 hours of lights off (red light). During these live observations, observers used all occurrence sampling to record the winner and loser in all instances of fighting, chasing, mounting, subordinate posture, and induced-flee behaviors (see **Table S1** for an ethogram of these behaviors). Winners of each encounter were considered to be those that chased, bit, mounted, or forced another individual to exhibit subordinate behavior. If behaviors between two females co-occurred within 2 seconds of each other they were recorded with the priority fighting, chasing, mounting, subordinate posture, flee. This method has been used previously in our lab to understand the social organization of groups of male mice[5,6,26,27,107,108]. Vaginal smears were collected from every mouse each evening at the same time of day (six to eight hours post lights-off). To collect the samples, trained lab members removed mice from the vivaria individually and placed them back as soon as the sample was collected. Collecting samples from each social group interrupted the group for less than 5 minutes. Smear samples were analyzed under a microscope by a single trained lab member and double checked by a second lab member to verify accuracy. Mice were weighed, final estrus smears taken, and euthanized via decapitation 2 hours post lights off on Day 15. Trunk blood was collected into heparinized tubes, immediately placed on ice, centrifuged at 4°C in a refrigerated centrifuge, and plasma separated and

frozen at -80°C until analyzed for corticosterone and estradiol levels via radioimmunoassay. Brains were
collected and flash frozen in hexane and stored at -80°C until dissection. At the end of group housing, the
2 most dominant and 2 most subordinate individuals were determined using the Glicko Rating System[26,109]
as well as David's Scores[26,110]. Plasma hormone and brain mRNA levels were measured for these two most
dominant and two most subordinate individuals in each group, except for in two cohorts where it was
difficult to distinguish the beta and gamma female so three dominant individuals and two subordinate
individuals were used.

Hormone assays

425   Plasma corticosterone and plasma estradiol concentrations were measured using commercially available
426   kits (MP Biomedicals) and conducted using the manufacturer's specifications. For the corticosterone
427   assay, the average inter-assay coefficient of variation was 9.3%, the lowest detectable was 24.78 ng/ul,
428   and the highest detectable was 938.34 ng/ul. For the estradiol assay, the coefficient of variation was 7.2%,
429   the lowest detectable was 8.53 pg/ul, and the highest detectable was 2455.79 pg/ul.

Gene expression

Brains were stored at -80° C until dissection. Samples of the medial preoptic area (mPOA) and
ventromedial hypothalamus (VMH) were collected using a Harris Micro-Punch with reference to the
coronal plane from the Mouse Brain Atlas[111] and the Allen Brain Atlas[112]. The mPOA was collected as one
1mm diameter area along the midline from Bregma +0.14 mm to -0.7 mm. The VMH was collected as one
1mm diameter area from each hemisphere from Bregma -1.34 mm to -1.82mm. RNA was isolated from
both brain regions using the AllPrep RNA Micro Kit (Qiagen) and reverse transcribed to cDNA using the
SuperScript III First-Strand Synthesis System for RT-PCR applications. Quantitative RT-PCR was performed
with 1ul of cDNA using an ABI 7500 Fast Thermal Cycler and the Fast SYBR Green Master Mix reagent
(Applied Biosystems). All primer probes (Sigma-Aldrich) were designed to span exon boundaries ensuring
amplification of only mRNA. The following validated quantitative PCR primers were used for mRNA
analysis: estrogen receptor alpha (ERα – Forward: CGTGTGCAATGACTATGCCTCT; Reverse:
TGGTGCATTGGTTTGTAGCTGG), estrogen receptor beta (ERβ – Forward: GTCAGGCACATCAGTAACAAGGG;
Reverse: ATTCAGCATCTCCAGCAGCAGGTC), progesterone receptor (PR – Forward:
GCGAGAGACAACTGCTTTCAGT; Reverse: CAAACACCATCAGGCTCATCCA), gonadotropin releasing
hormone (GnRH – Forward: AGCACTGGTCCTATGGGTTG; Reverse: GGTTCTGCCATTTGATCCAC), oxytocin
receptor (OTR – Forward: TTCTTCGTGCAGATGTGGAG; Reverse: CCAAGAGCATGGCAATGATG), opioid
receptor μ 1 (OPRM1 – Forward: AATGTTCATGGCAACCACAA; Reverse: TTTGAGCAGGTTCTCCCAGT).

Statistical analysis

449   All statistical analyses were undertaken in R v.3.5.0[113].

Group dominance structure and social organization

For each cohort, six measures of dominance structure and organization were calculated: Landau's
modified *h'*, directional consistency, steepness, triangle transitivity, despotism and Gini's coefficient of
wins and losses. The methods of calculation for these measures are detailed in Williamson et al.[26], but
briefly: Landau's modified *h'*, directional consistency, and steepness are calculated using frequency
win/loss sociomatrices, which are created using the total frequency of wins and losses recorded for each
individual over the observation period. Landau's modified h' evaluates the extent to which individuals in
a hierarchy can be linearly ordered[110] and ranges from 0-1, with a value of 1 indicating a completely linear
hierarchy. Triangle transitivity measures the proportion of relations between all triads that are transitive
(i.e. if individual A is dominant to individual B and individual B is dominant to individual C, then individual
A is dominant to individual C; a perfect hierarchy would have all transitive triads)[114]. It is calculated using
a binary win/loss sociomatrix, where 1s are assigned to individuals in rows that won more often against
individuals in columns and 0s are assigned to individuals in rows that lost more often to individuals in
columns. Triangle transitivity ranges from 0-1, with 1 indicating that all triads are transitive (i.e. a perfectly
linear hierarchy). Steepness measures the unevenness of relative individual dominance within the
hierarchy. This is calculated from the relative distribution of David's Scores, a win proportion measure
adjusted for strength of opponents[115]. It ranges from 0-1 with a score closer to 1 indicating that power is
not equitably distributed across the hierarchy, but rather lies in the hands of a few powerful individuals
at the top. Directional consistency measures the degree to which all agonistic interactions occur in the
direction from the more dominant to more subordinate individual in the pair. It also ranges from 0-1, with
1 indicating that all agonistic interactions occur in the direction of dominant to subordinate. Significance
testing for Landau's modified *h'*, triangle transitivity, steepness and directional consistency were carried
out using appropriate randomization methods[26]. P-values represent the proportion of times that values
were observed in randomized data that were greater than or equal to the observed values from empirical
data. Despotism is the proportion of all wins by the dominant male over the total number of aggressive
interactions over the observation period. It is a value between 0-1, with 1 indicating that the alpha male
performed 100% of all aggression within the group. Gini-coefficient is a measure of equality versus
inequality in a distribution. We calculated the Gini-coefficients for the frequency of wins and losses by
each animal across cohorts. It ranges from 0-1. Values closer to 1 indicate more inequality meaning that
a higher number of wins/losses are associated with relatively few individuals. Values closer to 0 indicate
that the frequency of wins/losses are equally distributed across all individuals. We examined the
association between initial body weight on Day 1 of group housing and final social rank for each social
group using Spearman Rank correlation tests.

Landau's modified *h'*, triangle transitivity, directional consistency and despotism were calculated using
the R package 'compete' v.0.1[116]. Steepness was calculated using the R package 'steepness' v.0.2.2[117]. Gini
coefficients were calculated using the 'ineq'[118] package.

Comparison between fighting, chasing and mounting behaviors

Raw frequency sociomatrices were constructed for each cohort based separately on fighting, chasing and
mounting behaviors. To determine how correlated each matrix (fighting, chasing or mounting) was to
each other within each cohort we performed a Quadratic Assignment Procedure (QAP) test with 1000
Monte Carlo randomizations of the data using the 'sna' package v.2.4[119]. From these matrices the
directional consistency and Gini-coefficient of wins and losses was also calculated for each behavior for
each cohort. To test if differences in these values existed between behaviors across cohorts we used
Friedman Tests. Significant differences between behaviors were determined using Nemenyi Post-Hoc
Tests using the 'PMCMR' R package v.4.3[120].

Emergence of hierarchies and individual ranks

To determine how quickly each cohort formed a linear hierarchy we calculated Landau's modified *h'* and
triangle transitivity as well as running significance tests for the aggregated win-loss data for each cohort
up to each day. The emergence of individual ranks across time was identified using Glicko ratings. All
individuals begin with a Glicko rating of 2200, and points are added or subtracted based on winning or
losing against other individuals. The degree of points won or lost is dependent upon the difference in
ratings between the two individuals. If an individual with a high Glicko rating defeats an individual with a
low Glicko rating, relatively few points would be added to their total and relatively few points would be
subtracted from the defeated individual. If an individual with a low Glicko rating defeats an individual with
a high Glicko rating, a larger number of points would be added to their total and subtracted from the loser.
Glicko ratings of all individuals in each group are recalculated after every behavioral interaction. We used
a constant value of 3 in our calculations of Glicko ratings. See Williamson et al.[26] for more information.
Glicko ratings were calculated using the 'PlayerRatings' package v.1.0 in R[121].

Comparison of female social hierarchy behavior to male social hierarchy behavior

To measure differences in social hierarchy structure between male and female social groups, we
compared female data from this study with previously published data on male social hierarchies from
Williamson et al.[26]. We recalculated hierarchy measures using the first fourteen days of observation data
from the 10 groups of 12 male CD-1 mice who were housed and observed in exactly the same manner as
the female groups in the current study. We used Wilcoxon rank sum tests to compare the values of
Landau's modified h', triangle transitivity, steepness, directional consistency, despotism and Gini-
coefficient of wins and losses between male and female groups.

Analysis of hourly occurrence rate of each agonistic behavior

All Bayesian linear and generalized linear regressions were fitted using R package 'brms'[122]. For each
fighting, chasing, and mounting behavior observed throughout the group housing period for each cohort,
we tested whether the day of group housing affects hourly occurrence rate of each behavior by fitting the
data with a gaussian distribution. Then we analyzed if the hourly occurrence rate of each behavior differs
by individual final social rank using a two-process hurdle-gamma family. In this model, the probability of

having zeros as the occurrence rates are modelled with a binomial error distribution with logit link
function and non-zero non-integer continuous values are fitted with a gamma error distribution with log
link[123]. We chose to use this model as the hourly occurrence rate data contains a significant number of
zeros as subordinate individuals often barely initiate agonistic behaviors and alpha individuals barely
receive aggression especially once the hierarchies are established. Further in this model we assume the
difference of any effect of individual ranks is not necessarily equidistant between ranks. We therefore
treated individual social rank as a monotonic predictor rather than a continuous variable throughout the
entire statistical analysis[124]. The beta coefficients estimated from monotonic models presented in this
study indicate the direction and the size (range between the lowest and highest categories of the ordinal
fixed factor e.g. rank) of the effects.

Analysis of estrus cycle state

We tested whether a mouse with higher social rank in the hierarchy is more likely to be in one estrus state
compared to another by fitting a multinomial logistic mixed effect model with estrus status as the
outcome variable, individual social rank as a monotonic predictor and cohort ID and subject ID as random
effects. We tested whether an individual is more likely to give (hourly given aggression rate) or receive
(hourly received aggression rate) each agonistic behavior when the mouse is in a certain estrus status
using a hurdle-gamma model. First, the probability of the hourly occurrence rate being zero was predicted
for each individual with daily Glicko rank of each mouse as a monotonic predictor to test whether the
observed number of zeros in the data could be explained by rank. After this hurdle, the non-zero values
of the hourly occurrence rate were fitted with a predictor of estrus state for each individual across each
group housing day. Cohort ID, subject ID and day of group housing were set as random effects for both
processes.

Analysis of plasma corticosterone and estradiol levels

Using leave-one-out cross-validation information criteria (LOOIC)[125], we compared four models with
different combination of fixed effects: i) dominant-subordinate status, ii) last measured estrus cycle
before blood collection, iii) dominant-subordinate status and last measured estrus state, iv) the
interaction of social status and estrus state. All models were fitted with cohort ID as a random effect. For
corticosterone data, a model with dominant-subordinate status only resulted in the best fit. A model with
the status and the estrus state resulted in the best fit for estradiol data.

Analysis of gene expression levels

Using the R package 'MCMC.qpcr'[126], we analyzed the differences in gene expression between dominant
and subordinate mice by fitting a generalized mixed effect model with Poisson-lognormal distribution and
a Bayesian Markov Chain Monte Carlo sampling approach. This approach provides advantages compared
to standard delta-CT analysis as it accounts for random variation between duplicates, increases power by
analyzing data for all target genes in one model, and does not require control genes[127]. Briefly, the raw
threshold cycle (CT) values were converted into molecule count data with consideration of the
amplification efficiency of each gene. We fitted the model with dominant-subordinate status category as

a fixed factor and cohort ID and subject ID as random factors. We confirmed linearity of the model by
inspecting diagnostic plots.

Data availability statement

All raw data and code used in this paper are publicly available at GitHub
https://github.com/jalapic/females

References

1.   Lenkov, K., Lee, M. H., Lenkov, O. D., Swafford, A. & Fernald, R. D. Epigenetic DNA Methylation

Linked to Social Dominance. *PLOS ONE* **10**, e0144750 (2015).

2.   Maruska, K. P. & Fernald, R. D. Social Regulation of Gene Expression in the Hypothalamic-Pituitary-

Gonadal Axis. *Physiology* **26**, 412–423 (2011).

3.   Rosvold, H. Enger, H., Mirsky, A. F. & Pribram, K. H. Influence of amygdalectomy on social behavior

in monkeys. *J. Comp. Physiol. Psychol.* **47**, 173–178 (1954).

4.   Wang, F. *et al.* Bidirectional Control of Social Hierarchy by Synaptic Efficacy in Medial Prefrontal

Cortex. *Science* **334**, 693–697 (2011).

5.   Williamson, C. M., Franks, B. & Curley, J. P. Mouse Social Network Dynamics and Community

Structure are Associated with Plasticity-Related Brain Gene Expression. *Front. Behav. Neurosci.* **10**,

(2016).

6.   Williamson, C. M., Romeo, R. D. & Curley, J. P. Dynamic changes in social dominance and mPOA

GnRH expression in male mice following social opportunity. *Horm. Behav.* **87**, 80–88 (2017).

7.   Erskine, M. S., Barfield, R. J. & Goldman, B. D. Intraspecific fighting during late pregnancy and

lactation in rats and effects of litter removal. *Behav. Biol.* **23**, 206–218 (1978).

8.   Haney, M., Debold, J. F. & Miczek, K. A. Maternal aggression in mice and rats towards male and

female conspecifics. *Aggress. Behav.* **15**, 443–453 (1989).

9.   Been, L. E., Gibbons, A. B. & Meisel, R. L. Towards a neurobiology of female aggression.

Neuropharmacology (2018). doi:10.1016/j.neuropharm.2018.11.039

10.   Duque-Wilckens, N. & Trainor, B. C. Behavioral Neuroendocrinology of Female Aggression. *Oxf.*

Res. Encycl. Neurosci. (2017). doi:10.1093/acrefore/9780190264086.013.11

11.   Holekamp, K. E. & Strauss, E. D. Aggression and dominance: an interdisciplinary overview. *Curr.*

Opin. Behav. Sci. **12**, 44–51 (2016).

12.   Correa, L. A., Frugone, M. J. & Soto-Gamboa, M. Social dominance and behavioral consequences of

intrauterine position in female groups of the social rodent Octodon degus. *Physiol. Behav.* **119**,

161–167 (2013).

13.   Rutberg, A. T. Dominance and Its Fitness Consequences in American Bison Cows. *Behaviour* **96**, 62–

91 (1986).

14.   Barrette, C. & Vandal, D. Social Rank, Dominance, Antler Size, and Access To Food in Snow-Bound

Wild Woodland Caribou. *Behaviour* **97**, 118–145 (1986).

15.   Ceacero, F. *et al.* Benefits for Dominant Red Deer Hinds under a Competitive Feeding System: Food

Access Behavior, Diet and Nutrient Selection. *PLOS ONE* **7**, e32780 (2012).

16.   Whitten, P. L. Diet and dominance among female vervet monkeys (Cercopithecus aethiops). *Am. J.*

Primatol. **5**, 139–159 (1983).

17.   Murray, C. M., Eberly, L. E. & Pusey, A. E. Foraging strategies as a function of season and rank

among wild female chimpanzees (Pan troglodytes). *Behav. Ecol.* **17**, 1020–1028 (2006).

18.   Chovnick, A., Yasukawa, N. J., Monder, H. & Christian, J. J. Female behavior in populations of mice

in the presence and absence of male hierarchy. *Aggress. Behav.* **13**, 367–375 (1987).

19.   Yasukawa, N. J., Monder, H., Leff, F. R. & Christian, J. J. Role of female behavior in controlling

population growth in mice. *Aggress. Behav.* **11**, 49–64 (1985).

20.   Weidt, A., Gygax, L., Palme, R., Touma, C. & König, B. Impact of male presence on female sociality

and stress endocrinology in wild house mice (Mus musculus domesticus). *Physiol. Behav.* **189**, 1–9

(2018).

21.   Palanza, P., Della Seta, D., Ferrari, P. F. & Parmigiani, S. Female competition in wild house mice

depends upon timing of female/male settlement and kinship between females. *Anim. Behav.* **69**,

1259–1271 (2005).

22.   Rusu, A. S. & Krackow, S. Kin-preferential cooperation, dominance-dependent reproductive skew,

and competition for mates in communally nesting female house mice. *Behav. Ecol. Sociobiol.* **56**,

298–305 (2004).

23.   Schuhr, B. Social structure and plasma corticosterone level in female albino mice. *Physiol. Behav.*

40, 689–693 (1987).

24.   van den Berg, W. E., Lamballais, S. & Kushner, S. A. Sex-Specific Mechanism of Social Hierarchy in

Mice. *Neuropsychopharmacology* **40**, 1364–1372 (2015).

25.   Varholick, J. A., Bailoo, J. D., Palme, R. & Würbel, H. Phenotypic variability between Social

Dominance Ranks in laboratory mice. *Sci. Rep.* **8**, 6593 (2018).

26.   Williamson, C. M., Lee, W. & Curley, J. P. Temporal dynamics of social hierarchy formation and

maintenance in male mice. *Anim. Behav.* **115**, 259–272 (2016).

27.   Williamson, C. M., Lee, W., Romeo, R. D. & Curley, J. P. Social context-dependent relationships

between mouse dominance rank and plasma hormone levels. *Physiol. Behav.* **171**, 110–119 (2017).

28.   Palanza, P., Gioiosa, L. & Parmigiani, S. Social stress in mice: Gender differences and effects of

estrous cycle and social dominance. *Physiol. Behav.* **73**, 411–420 (2001).

29.   Roberts, D. C. S., Bennett, S. a. L. & Vickers, G. J. The estrous cycle affects cocaine self-

administration on a progressive ratio schedule in rats. *Psychopharmacology (Berl.)* **98**, 408–411

(1989).

30.   Gray, P. Effect of the estrous cycle on conditioned avoidance in mice. *Horm. Behav.* **8**, 235–241

(1977).

31.   Floody, O. R. Hormones and Aggression in Female Mammals. in *Hormones and Aggressive Behavior*

39–89 (Springer, Boston, MA, 1983). doi:10.1007/978-1-4613-3521-4_3

32.   Davis, E. S. & Marler, C. A. C-FOS changes following an aggressive encounter in female California

mice: A synthesis of behavior, hormone changes and neural activity. *Neuroscience* **127**, 611–624

(2004).

33.   Ho, H.-P. *et al.* The Serotonin Reuptake Inhibitor Fluoxetine Reduces Sex Steroid-Related

Aggression in Female Rats: An Animal Model of Premenstrual Irritability?

Neuropsychopharmacology **24**, 502–510 (2001).

34.   Seward, J. P. Aggressive behavior in the rat. I. General characteristics; age and sex differences. *J.*

Comp. Psychol. **38**, 175–197 (1945).

35.   Ciaccio, L. A., Lisk, R. D. & Reuter, L. A. Prelordotic behavior in the hamster: A hormonally

modulated transition from aggression to sexual receptivity. *J. Comp. Physiol. Psychol.* **93**, 771–780

(1979).

36.   Floody, O. R. & Pfaff, D. W. Aggressive behavior in female hamsters: The hormonal basis for

fluctuations in female aggressiveness correlated with estrous state. *J. Comp. Physiol. Psychol.* **91**,

443–464 (1977).

37.   Ward Kislak, J. & Beach, F. A. INHIBITION OF AGGRESSIVENESS BY OVARIAN HORMONES.

Endocrinology **56**, 684–692 (1955).

38.   Wise, D. A. Aggression in the female golden hamster: Effects of reproductive state and social

isolation. *Horm. Behav.* **5**, 235–250 (1974).

39.   Barr, G. A., Gibbons, J. L. & Moyer, K. E. Male-female differences and the influence of neonatal and

adult testosterone on intraspecies aggression in rats. *J. Comp. Physiol. Psychol.* **90**, 1169–1183

(1976).

40.   Cordero, M. I., Ansermet, F. & Sandi, C. Long-term programming of enhanced aggression by

657   peripuberty stress in female rats. *Psychoneuroendocrinology* **38**, 2758–2769 (2013).

658   41.   Jong, T. R. de, Beiderbeck, D. I. & Neumann, I. D. Measuring Virgin Female Aggression in the

659   Female Intruder Test (FIT): Effects of Oxytocin, Estrous Cycle, and Anxiety. *PLOS ONE* **9**, e91701

660   (2014).

661   42.   Takahashi, L. K. & Lisk, R. D. Intrasexual interactions among female golden hamsters (Mesocricetus

662   auratus) over the estrous cycle. *J. Comp. Psychol. Wash. DC 1983* **98**, 267–275 (1984).

663   43.   Hyde, J. S. & Sawyer, T. F. Estrous cycle fluctuations in aggressiveness of house mice. *Horm. Behav.*

664   **9**, 290–295 (1977).

665   44.   Koonce, C. J. & Frye, C. A. Female mice with deletion of Type One 5α-reductase have reduced

666   reproductive responding during proestrus and after hormone-priming. *Pharmacol. Biochem.*

667   *Behav.* **122**, 20–29 (2014).

668   45.   Kruk, M. R. Ethology and pharmacology of hypothalamic aggression in the rat. *Neurosci. Biobehav.*

669   *Rev.* **15**, 527–538 (1991).

670   46.   Albert, D. J., Jonik, R. H. & Walsh, M. L. Hormone-dependent aggression in male and female rats:

671   Experiential, hormonal, and neural foundations. *Neurosci. Biobehav. Rev.* **16**, 177–192 (1992).

672   47.   Malsbury, C. W., Strull, D. & Daood, J. Half-cylinder cuts antero-lateral to the ventromedial nucleus

673   reduce sexual receptivity in female golden hamsters. *Physiol. Behav.* **21**, 79–87 (1978).

674   48.   Meisel, R. L. & Sterner, M. R. Progesterone inhibition of sexual behavior is accompanied by an

675   activation of aggression in female Syrian hamsters. *Physiol. Behav.* **47**, 415–417 (1990).

676   49.   Meisel, R. L., Sterner, M. R. & Diekman, M. A. Differential hormonal control of aggression and

677   sexual behavior in female Syrian hamsters. *Horm. Behav.* **22**, 453–466 (1988).

50.   Takahashi, L. K. & Lisk, R. D. Estrogen action in anterior and ventromedial hypothalamus and the modulation of heterosexual behavior in female golden hamsters. *Physiol. Behav.* **34**, 233–239 (1985).

51.   Hashikawa, K. *et al.* Esr1+ cells in the ventromedial hypothalamus control female aggression. *Nat. Neurosci.* **advance online publication**, (2017).

52.   Bosch, O. J. & Neumann, I. D. Brain vasopressin is an important regulator of maternal behavior independent of dams' trait anxiety. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 17139–17144 (2008).

53.   Hammond, M. A. & Rowe, F. A. Medial preoptic and anterior hypothalamic lesions: Influences on aggressive behavior in female hamsters. *Physiol. Behav.* **17**, 507–513 (1976).

54.   Harmon, A. C., Huhman, K. L., Moore, T. O. & Albers, H. E. Oxytocin Inhibits Aggression in Female Syrian Hamsters. *J. Neuroendocrinol.* **14**, 963–969 (2002).

55.   Takahashi, L. K. & Lisk, R. D. Diencephalic Sites of Progesterone Action for Inhibiting Aggression and Facilitating Sexual Receptivity in Estrogen-Primed Golden Hamsters. *Endocrinology* **116**, 2393–2399 (1985).

56.   Cushing, B. S. Estrogen Receptor Alpha Distribution and Expression in the Social Neural Network of Monogamous and Polygynous Peromyscus. *PLOS ONE* **11**, e0150373 (2016).

57.   Murakami, G. Distinct Effects of Estrogen on Mouse Maternal Behavior: The Contribution of Estrogen Synthesis in the Brain. *PLOS ONE* **11**, e0150728 (2016).

58.   Soma, K. K., Rendon, N. M., Boonstra, R., Albers, H. E. & Demas, G. E. DHEA effects on brain and behavior: Insights from comparative studies of aggression. *J. Steroid Biochem. Mol. Biol.* **145**, 261–272 (2015).

59.   Albert, D. J., Petrovic, D. M. & Walsh, M. L. Ovariectomy attenuates aggression by female rats cohabitating with sexually active sterile males. *Physiol. Behav.* **45**, 225–228 (1989).

60.   Rosvall, K. A. *et al.* Neural sensitivity to sex steroids predicts individual differences in aggression:

702   implications for behavioural evolution. *Proc. R. Soc. Lond. B Biol. Sci.* rspb20120442 (2012).

703   doi:10.1098/rspb.2012.0442

704   61.   Rubenstein, D. R. & Wikelski, M. Steroid hormones and aggression in female Galápagos marine

705   iguanas. *Horm. Behav.* **48**, 329–341 (2005).

706   62.   Ervin, K. S. J. *et al.* Estrogen involvement in social behavior in rodents: Rapid and long-term actions.

707   *Horm. Behav.* **74**, 53–76 (2015).

708   63.   Pfaff, D. W. *et al.* Estrogens, brain and behavior: studies in fundamental neurobiology and

709   observations related to women's health. *J. Steroid Biochem. Mol. Biol.* **74**, 365–373 (2000).

710   64.   Bychowski, M. E. & Auger, C. J. Progesterone impairs social recognition in male rats. *Horm. Behav.*

711   **61**, 598–604 (2012).

712   65.   Ragnauth, A. K. *et al.* Female oxytocin gene-knockout mice, in a semi-natural environment, display

713   exaggerated aggressive behavior. *Genes Brain Behav.* **4**, 229–239 (2005).

714   66.   Campbell, A. Attachment, aggression and affiliation: The role of oxytocin in female social behavior.

715   *Biol. Psychol.* **77**, 1–10 (2008).

716   67.   Moles, A., Kieffer, B. L. & D'Amato, F. R. Deficit in Attachment Behavior in Mice Lacking the μ-

717   Opioid Receptor Gene. *Science* **304**, 1983–1986 (2004).

718   68.   Wöhr, M., Moles, A., Schwarting, R. K. W. & D'Amato, F. R. Lack of social exploratory activation in

719   male μ-opioid receptor KO mice in response to playback of female ultrasonic vocalizations. *Soc.*

720   *Neurosci.* **6**, 76–87 (2011).

721   69.   Briand, L. A. *et al.* Mouse Model of OPRM1 (A118G) Polymorphism Increases Sociability and

722   Dominance and Confers Resilience to Social Defeat. *J. Neurosci.* **35**, 3582–3590 (2015).

723   70.   Dadomo, H. *et al.* What is stressful for females? Differential effects of unpredictable

724   environmental or social stress in CD1 female mice. *Horm. Behav.* **98**, 22–32 (2018).

71.   Jacobson-Pick, S., Audet, M.-C., McQuaid, R. J., Kalvapalle, R. & Anisman, H. Social agonistic

distress in male and female mice: changes of behavior and brain monoamine functioning in

relation to acute and chronic challenges. *PloS One* **8**, e60133 (2013).

72.   Chapman, J. C., Christian, J. J., Pawlikowski, M. A. & Michael, S. D. Analysis of steroid hormone

levels in female mice at high population density. *Physiol. Behav.* **64**, 529–533 (1998).

73.   Kaufmann, J. H. On the Definitions and Functions of Dominance and Territoriality. *Biol. Rev.* **58**, 1–

20 (1983).

74.   Rowell, T. E. The concept of social dominance. *Behav. Biol.* **11**, 131–154 (1974).

75.   Stockley, P. & Bro-Jørgensen, J. Female competition and its evolutionary consequences in

mammals. *Biol. Rev.* **86**, 341–366 (2011).

76.   Dagg, A. I. Homosexual behaviour and female-male mounting in mammals—a first survey.

Mammal Rev. **14**, 155–185 (1984).

77.   Fang, J. & Clemens, L. G. Contextual determinants of female–female mounting in laboratory rats.

Anim. Behav. **57**, 545–555 (1999).

78.   Kimchi, T., Xu, J. & Dulac, C. A functional circuit underlying male sexual behaviour in the female

mouse brain. *Nature* **448**, 1009–1014 (2007).

79.   Dean, M. D., Ardlie, K. G. & Nachman, M. W. The frequency of multiple paternity suggests that

sperm competition is common in house mice (Mus domesticus). *Mol. Ecol.* **15**, 4141–4151 (2006).

80.   Bronson, F. H. The Reproductive Ecology of the House Mouse. *Q. Rev. Biol.* **54**, 265–299 (1979).

81.   Bronson, F. H. Mammalian Reproduction: An Ecological Perspective. *Biol. Reprod.* **32**, 1–26 (1985).

82.   Brain, P. F., Haug, M. & Parmigiani, S. The aggressive female rodent: Redressing a "scientific" bias.

in *Of Mice and Women: Aspects of Female Aggression* 27–35 (Academic Press, San Diego, 1992).

83.   Anthoney, T. R. The Ontogeny of Greeting, Grooming, and Sexual Motor Patterns in Captive

Baboons (Superspecies Papio Cynocephalus). *Behaviour* **31**, 358–372 (1968).

84.   Talmage-Riggs, G. & Anschel, S. Homosexual Behavior and Dominance Hierarchy in a Group of

750   Captive Female Squirrel Monkeys (Saimiri sciureus). *Folia Primatol. (Basel)* **19**, 61–72 (1973).

751   85.   Borries, C., Sommer, V. & Srivastava, A. Dominance, age, and reproductive success in free-ranging

752   female hanuman langurs (Presbytis entellus). *Int. J. Primatol.* **12**, 231–257 (1991).

753   86.   Albonetti, M. E. & Dessi-Fulgheri, F. Female-female mounting in the European rabbit. *Z*

754   *Saugetierkunde* **55**, 128–138 (1990).

755   87.   Champlin, A. K. SUPPRESSION OF OESTRUS IN GROUPED MICE: THE EFFECTS OF VARIOUS

756   DENSITIES AND THE POSSIBLE NATURE OF THE STIMULUS. *Reproduction* **27**, 233–241 (1971).

757   88.   Ma, W., Miao, Z. & Novotny, M. V. Role of the Adrenal Gland and Adrenal-Mediated Chemosignals

758   in Suppression of Estrus in the House Mouse: The Lee-Boot Effect Revisited. *Biol. Reprod.* **59**,

759   1317–1320 (1998).

760   89.   Van Der Lee, S. & Boot, L. M. Spontaneous pseudopregnancy in mice. *Acta Physiol. Pharmacol.*

761   *Neerl.* **4**, 442–444 (1955).

762   90.   Whitten, W. K. OCCURRENCE OF ANOESTRUS IN MICE CAGED IN GROUPS. *J. Endocrinol.* **18**, 102–

763   107 (1959).

764   91.   Lenz, K. M., Nugent, B. M. & McCarthy, M. M. Sexual Differentiation of the Rodent Brain: Dogma

765   and Beyond. *Front. Neurosci.* **6**, (2012).

766   92.   Wu, M. V. *et al.* Estrogen Masculinizes Neural Pathways and Sex-Specific Behaviors. *Cell* **139**, 61–

767   72 (2009).

768   93.   Schulz, K. M. & Sisk, C. L. The organizing actions of adolescent gonadal steroid hormones on brain

769   and behavioral development. *Neurosci. Biobehav. Rev.* **70**, 148–158 (2016).

770   94.   Harris, A. Z. *et al.* A Novel Method for Chronic Social Defeat Stress in Female Mice.

771   *Neuropsychopharmacology* **43**, 1276–1283 (2018).

95.   Jarcho, M. R., Massner, K. J., Eggert, A. R. & Wichelt, E. L. Behavioral and physiological response to

onset and termination of social instability in female mice. *Horm. Behav.* **78**, 135–140 (2016).

96.   Schmidt, M. V. *et al.* A novel chronic social stress paradigm in female mice. *Horm. Behav.* **57**, 415–

420 (2010).

97.   Takahashi, L. K., Lisk, R. D. & Burnett II, A. L. Dual estradiol action in diencephalon and the

regulation of sociosexual behavior in female golden hamsters. *Brain Res.* **359**, 194–207 (1985).

98.   Ogawa, S., Lubahn, D. B., Korach, K. S. & Pfaff, D. W. Behavioral effects of estrogen receptor gene

disruption in male mice. *Proc. Natl. Acad. Sci.* **94**, 1476–1481 (1997).

99.   Ogawa, S., Taylor, J. A., Lubahn, D. B., Korach, K. S. & Pfaff, D. W. Reversal of Sex Roles in Genetic

Female Mice by Disruption of Estrogen Receptor Gene. *Neuroendocrinology* **64**, 467–470 (1996).

100. Ogawa, S. *et al.* Roles of Estrogen Receptor-α Gene Expression in Reproduction-Related Behaviors

in Female Mice*This work was supported by the Harry Frank Guggenheim Foundation (to S.O.), the

University of Missouri-Columbia molecular biology program (to D.B.L.), and NIH Grant HD-05751

(to D.W.P.). *Endocrinology* **139**, 5070–5081 (1998).

101. Nomura, M. *et al.* Genotype/Age Interactions on Aggressive Behavior in Gonadally Intact Estrogen

Receptor β Knockout (βERKO) Male Mice. *Horm. Behav.* **41**, 288–296 (2002).

102. Ogawa, S. *et al.* Survival of reproductive behaviors in estrogen receptor β gene-deficient (βERKO)

male and female mice. *Proc. Natl. Acad. Sci.* **96**, 12887–12892 (1999).

103. Clipperton-Allen, A. E., Almey, A., Melichercik, A., Allen, C. P. & Choleris, E. Effects of an estrogen

receptor alpha agonist on agonistic behaviour in intact and gonadectomized male and female

mice. *Psychoneuroendocrinology* **36**, 981–995 (2011).

104. Clipperton-Allen, A. E., Cragg, C. L., Wood, A. J., Pfaff, D. W. & Choleris, E. Agonistic behavior in

males and females: Effects of an estrogen receptor beta agonist in gonadectomized and gonadally

intact mice. *Psychoneuroendocrinology* **35**, 1008–1022 (2010).

105. Ogawa, S., Tsukahara, S., Choleris, E. & Vasudevan, N. Estrogenic regulation of social behavior and

sexually dimorphic brain formation. *Neurosci. Biobehav. Rev.* (2018).

doi:10.1016/j.neubiorev.2018.10.012

106. McCarthy, M. M. Oxytocin inhibits infanticide in female house mice (Mus domesticus). *Horm.*

Behav. **24**, 365–375 (1990).

107. Curley, J. P. Temporal pairwise-correlation analysis provides empirical support for attention

hierarchies in mice. *Biol. Lett.* **12**, 20160192 (2016).

108. Lee, W., Khan, A. & Curley, J. P. Major urinary protein levels are associated with social status and

context in mouse social hierarchies. *Proc R Soc B* **284**, 20171570 (2017).

109. Glickman, M. E. Parameter estimation in large dynamic paired comparison experiments. *J. R. Stat.*

Soc. Ser. C Appl. Stat. **48**, 377–394 (1999).

110. De Vries, H. An improved test of linearity in dominance hierarchies containing unknown or tied

relationships. *Anim. Behav.* **50**, 1375–1389 (1995).

111. Paxinos, G. & Franklin, K. B. J. *The Mouse Brain in Stereotaxic Coordinates*. (Gulf Professional

Publishing, 2004).

112. Lein, E. S. *et al.* Genome-wide atlas of gene expression in the adult mouse brain. *Nature* **445**, 168–

176 (2007).

113. R Core Team. *R: A language and environment for statistical computing.* (2017).

114. Shizuka, D. & McDonald, D. B. A social network perspective on measurements of dominance

hierarchies. *Anim. Behav.* **83**, 925–934 (2012).

115. Gammell, M. P., Vries, H. de, Jennings, D. J., Carlin, C. M. & Hayden, T. J. David's score: a more

appropriate dominance ranking method than Clutton-Brock et al.'s index. *Anim. Behav.* **66**, 601–

605 (2003).

116. Curley, J. P. *compete: Organizing and Analyzing Social Dominance Hierarchy Data.* (2016).

117. Leiva, D. & de Vries, H. *steepness: Testing steepness of dominance hierarchies*. (2014).

118. Zeileis, A. *ineq: Measuring inequality, concentration, and poverty*. (2014).

119. Butts, C. T. *sna: Tools for Social Network Analysis*. (2016).

120. Pohlert, T. *PMCMR: Calculate Pairwise Multiple Comparisons of Mean Rank Sums*. (2018).

121. Stephenson, A. & Sonas, J. *PlayerRatings: Dynamic updating methods for player ratings estimation*.

(2012).

122. Bürkner, P.-C. brms: An R Package for Bayesian Multilevel Models Using Stan. *J. Stat. Softw.* **1**, 1

(2017).

123. Pohlmeier, W. & Ulrich, V. An Econometric Model of the Two-Part Decisionmaking Process in the

Demand for Health Care. *J. Hum. Resour.* **30**, 339–361 (1995).

124. Bürkner, P.-C. & Charpentier, E. Monotonic Effects: A Principled Approach for Including Ordinal

Predictors in Regression Models. (2018). doi:10.31234/osf.io/9qkhj

125. Gelman, A., Hwang, J. & Vehtari, A. Understanding predictive information criteria for Bayesian

models. *Stat. Comput.* **24**, 997–1016 (2014).

126. Matz, M. V. *MCMC.qpcr: Bayesian Analysis of qRT-PCR Data*. (2016).

127. Matz, M. V., Wright, R. M. & Scott, J. G. No Control Genes Required: Bayesian Analysis of qRT-PCR

Data. *PLOS ONE* **8**, e71448 (2013).

Acknowledgements

839   We thank Dr. Frances Champagne for advice and suggestions in writing the manuscript and Curley Lab
840   students for help with behavioral observations.

Author contributions statement

842   CMW, WL, ARD and JPC conceived and planned the experiments. CMW, WL, ARD, RDR and AL carried
843   out the behavioral work and husbandry. RDR planned and carried out the hormone analysis. CMW and
844   ARD carried out the gene expression experiment. CMW, WL and JPC analyzed the data. CMW, WL, & JPC

845 wrote the paper. All authors provided critical feedback and helped shape the research, analysis and
846 manuscript.

Additional information

Competing interests

849 The authors declare there are no competing interests.

Table 1. Social hierarchy measures for each cohort (A-H). Females have significantly linear hierarchies (as measured by Landau's Modified h' value and triangle transitivity (ttri)) as well as significantly steep hierarchies (steepness). Aggressive behavior is also significantly directionally consistent (DC). Significant values are bolded. Despotism and the Gini-Coefficient of Wins and Losses measure how evenly distributed aggressive behavior is across ranks. Female hierarchies are not highly despotic.

Cohort ID	Landau's h'	*ttri*	Steepness	DC	Despotism	GC Wins	GC Losses
A	0.19	0.08	0.30	**0.54**	0.15	0.24	0.22
B	**0.85**	**0.88**	**0.61**	**0.76**	0.26	0.43	0.29
C	**0.71**	**0.82**	**0.55**	**0.70**	0.23	0.45	0.27
D	**0.67**	**0.68**	**0.48**	**0.69**	0.22	0.43	0.23
E	**0.49**	**0.48**	**0.47**	**0.68**	0.22	0.40	0.19
F	**0.46**	**0.36**	**0.46**	**0.72**	0.25	0.46	0.21
G	**0.82**	**0.95**	**0.55**	**0.79**	0.34	0.49	0.21
H	**0.78**	**0.90**	**0.51**	**0.57**	0.16	0.34	0.20

Figure legends

Figure 1. Row 1) Raw sociomatrices of wins and losses for each cohort (A-H). Each value represents the total number of wins by the individual in each row against the individual in each column. The degree of redness represents the frequency of wins. Individuals are ordered by I&SI rank. **Row 2)** Binarized sociomatrices of wins and losses for each cohort. A 1 in a cell represents that the individual in the row won more fights than they lost against the individual in the column. The degree of redness represents the directional consistency of wins and losses for each dyad. Individuals are ordered by I&SI rank. **Row 3)** The temporal patterning of dominance. The Glicko Rating of dominance (y-axis) against each successive behavioral event (x-axis). Each line represents the dominance rating of one individual in each cohort. The lines are colored from black (eventual most dominant individual) to red (eventual least dominant individual).

Figure 2. Female dominance hierarchies emerge rapidly. The triangle transitivity (A) and Landau's Modified h' value across the fourteen days of the housing period. Each line represents a different female cohort. Red lines indicate significant values.

Figure 3. Comparison between male and female social hierarchies. Boxplots compare modified Landau's h', triangle transitivity, steepness, directional consistency, despotism, and Gini coefficient (GC) of wins and losses from 8 female social groups (orange) and 10 male social groups (purple).

Figure 4. The effect of final social rank on the hourly rate of each agonistic behavior given (left) and received (right). Lines represent the median posterior means of fighting, chasing and mounting events per hour of observation by rank across the fourteen days of observation. The shaded areas show the 95% CI.

Figure 5 Row 1) The proportion of the fourteen days of observations in each estrus state by rank across each cohort (diestrus = black; metestrus = purple; proestrus = orange; estrus = yellow). **Row 2)** The estrus state assigned by vaginal smear for each rank across cohorts (diestrus = black; metestrus = purple; proestrus = orange; estrus = yellow; unable to determine = white).

Figure 6. Plasma Corticosterone and Estradiol levels for dominant and subordinate females. **(A)** Subordinate females have significantly higher levels of plasma corticosterone than dominant females. **(B)** There is no difference between dominant and subordinate individuals in plasma estradiol levels. Text in each panel indicates posterior means and 95% CI between dominant and subordinate females ($b_{subordinate-dominant}$). Red text indicates significant differences.

Figure 7. ERα, ERβ, and OTR mRNA levels in the VMH are higher for subordinate compared to dominant females. Solid dots and whiskers indicate posterior means and 95% credible intervals respectively for gene expression differences between dominant and subordinate females in each brain region and gene ($b_{subordinate-dominant}$). Text on the left notes posterior means and 95% CI for each comparison. Red text indicates significant differences.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7