Microbial phenotypic heterogeneity in response to a metabolic toxin: continuous, dynamically shifting distribution of formaldehyde tolerance in *Methylobacterium extorquens* populations

Jessica A. Lee[1,5,6,7], Siavash Riazi[2,5,6], Shahla H. Nemati[3,5], Andreas E. Vasdekis[3,5,6], Benjamin J. Ridenhour[4,5], Christopher H. Remien[4,5], and Christopher J. Marx[1,5,6]

Short title

Dynamics of microbial phenotypic heterogeneity in toxin tolerance

Affiliations

[1]Department of Biological Sciences, University of Idaho, Moscow, ID

[2]Bioinformatics and Computational Biology Graduate Program, University of Idaho, Moscow, ID

[3]Department of Physics, University of Idaho, Moscow, ID

[4]Department of Mathematics, University of Idaho, Moscow, ID

[5]Center for Modeling Complex Interactions, University of Idaho, Moscow, ID

[6]Institute for Bioinformatics and Evolutionary Studies, University of Idaho, Moscow, ID

[7]current address: Global Viral, San Francisco, CA

Contributions

Jessica A. Lee*: conceptualization, formal analysis, investigation, methodology, visualization, writing—original draft preparation

Siavash Riazi*: conceptualization, formal analysis, methodology, software, visualization, writing—review and editing

Shahla H. Nemati: investigation, visualization, writing—review and editing

Andreas E. Vasdekis: funding acquisition, investigation, visualization, writing—review and editing

Benjamin J. Ridenhour: formal analysis, software, supervision, writing—review and editing

Christopher H. Remien: conceptualization, funding acquisition, formal analysis, supervision, writing—review and editing

Christopher J. Marx: conceptualization, funding acquisition, project administration,

supervision, writing—review and editing

* These authors contributed equally to this work.

Corresponding authors

Jessica A. Lee: jessica.audrey.lee@gmail.com

Christopher J. Marx: cmarx@uidaho.edu

Acknowledgments

We are grateful to Jannell Bazurto, Eric Bruger, Sergey Stolyar, Nicholas Shevalier, and

Joshua Wirtz for assistance in conducting *M. extorquens* experiments; to Dipti Nayak for

advice and resources facilitating those experiments; to Craig Miller for advice on

analyzing colony counts; to William Harcombe and Jeremy Chacón for guidance on

time-lapse imaging of colony growth using flatbed scanners; to Mark Lamourine for help

with software and Dan Schneider for contributions to hardware for running the scanners;

and to Pål Johnsen, Alex Bradley and Jannell Bazurto for helpful comments on the

manuscript. Flow cytometry was conducted at the IBEST Optical Imaging Core at the

University of Idaho under the guidance of Ann Norton. Genomic DNA sample

preparation and sequencing were managed by the IBEST Genomics Resources Core at

the University of Idaho by Dan New, Samuel Hunter, and Matt Fagnan. IBEST is

supported in part by NIH COBRE grant P30GM103324. Funding for this work came

from an Army Research Office MURI sub-award to CJM (W911NF-12-1-0390), and a

CMCI Pilot Grant to CJM, CHR, and AEV (parent NIH award P20GM104420).

Abstract

While microbiologists often make the simplifying assumption that genotype determines phenotype in a given environment, it is becoming increasingly apparent that phenotypic heterogeneity (in which one genotype generates multiple phenotypes simultaneously even in a uniform environment) is common in many microbial populations. The importance of phenotypic heterogeneity has been demonstrated in a number of model systems involving binary phenotypic states (e.g., growth/non-growth); however, less is known about systems involving phenotype distributions that are continuous across an environmental gradient, and how those distributions change when the environment changes. Here, we describe a novel instance of phenotypic diversity in tolerance to a metabolic toxin within wild-type populations of *Methylobacterium extorquens,* a ubiquitous phyllosphere methylotroph capable of growing on the methanol periodically released from plant leaves. The first intermediate in methanol metabolism is formaldehyde, a potent cellular toxin that is lethal in high concentrations. We have found that at moderate concentrations, formaldehyde tolerance in *M. extorquens* is heterogeneous, with individual cells' minimum tolerance levels ranging between 0 mM and 8 mM. This form of heterogeneity is continuous in terms of threshold (in the range of maximum tolerances possible), yet binary in outcome (at a given formaldehyde concentration, cells either grow normally or die, with no intermediate phenotype), and it is not associated with any detectable genetic mutations. Moreover, tolerance distributions within the population are dynamic, changing over time in response to growth conditions. We characterized this phenomenon using bulk liquid culture experiments, colony growth tracking, flow cytometry, time-lapse microscopy, and genome resequencing. Finally, we used mathematical modeling to better understand the processes by which cells change phenotype, and found evidence for both stochastic, bidirectional phenotype diversification and responsive, directed phenotype shifts, depending on the growth substrate and the presence of toxin.

Introduction

Microbes are individuals. Even in seemingly simple unicellular organisms, phenotype is not always the straightforward product of genotype and environment; cells with identical genotypes in identical environments may nonetheless demonstrate cell-to-cell diversity in the expression of any of a number of traits. Frequently overlooked in everyday microbiology experiments, the phenomenon of cell-to-cell phenotypic heterogeneity has drawn increasing attention in recent decades both from a systems biology perspective and from an evolutionary perspective, as well as for its consequences to applied fields such as medicine (e.g., antibiotic persistence [1]; cancer cell drug tolerance [2,3]) and biological engineering [4].

Some forms of population heterogeneity might be considered trivial: molecular interactions within cells are inherently noisy. All genes might be expected to be expressed at slightly different levels among different cells [5–7], and historical contingency (e.g., pole age, asymmetrical division of macromolecules) can also create inherent diversity within microbial populations, independent of signals from the environment [8–10]. Naturally, evolution imposes some pressure on organisms to limit the noise in pathways that are essential for life; what is more remarkable is that some pathways seem to be selected for increased noise, and in many cases that noise is further amplified by feedback circuits, enabling a population to split into discretely different phenotypes. Specifically, genes involved in stress response and in metabolism have been found to show higher heterogeneity in expression than those in other pathways [11], and many of the well-understood examples of binary phenotypes involve stress response (antibiotic persistence [12]; sporulation [13]), or carbon transport and use (lactose utilization in *E. coli* [14]; diauxic switch in *S. cerevisiae* [15,16]). While some forms of phenotypic noise may have little fitness impact, phenotypic heterogeneity involving binary phenotypes is argued in many cases to offer an evolutionary advantage, as a strategy to facilitate diversifying bet-hedging or division of labor [11,17]. In many cases, the genetic basis of phenotypic differentiation is known, and laboratory evolution studies have demonstrated how populations can evolve the timing or

frequency of that differentiation in response to environmental selection [18,19].

Moreover, it is argued that phenotypic heterogeneity influences the rate of genotypic

evolution either by creating an "epigenetic load" that contributes to extinction [20], or by

allowing populations to adapt faster to changing environments and by increasing the

opportunities for mutations to arise and reach fixation [21,22].

Besides cases of binary phenotypes or modest phenotypic noise around a population

mean, a third possibility involves phenotypes that fall along a continuous gradient.

Fewer such phenomena have been described [11], but some examples include

populations in which cells have a range of levels of stress tolerance: for instance, a

genetically chloramphenicol (Cm)-resistant *E. coli* strain exhibits a wide, continuously-

varying distribution of maximum concentrations of Cm that individual cells can tolerate.

The population splits into growing and non-growing subpopulations in the presence of

Cm, and the proportion of growing cells varies according to the environmental Cm

concentration [23]. This effect is mediated by a positive-feedback interaction between

intracellular chloramphenicol acetyltransferase (CAT) activity and innate cell growth;

thus the ultimate fate of an individual cell depends on its initial internal CAT activity,

which can vary continuously among cells. If such a population were to experience shifts

in environmental Cm concentrations, the population distribution of per-cell CAT activity

levels could presumably shift as a consequence of either cellular responses (e.g., CAT

upregulation), or simply by selection against sensitive cells; however, this has not been

described. Analogous population-level phenotype distribution shifts have been explored

only through mathematical modeling, for instance in the case of human cancer cells

exhibiting a gradient of tolerances to cytotoxic drugs [24]. In this case, it is assumed that

random epimutations result in small phenotypic variations in drug tolerance, and that

drug exposure leads to selection upon that diversity. However, experimental work would

be necessary to verify whether modeling accurately predicts cell population dynamics,

or whether other processes—such as those in which cells sense drug concentrations

and respond with phenotype shifts—might also play a role. Examples of phenotypic

heterogeneity such as these two cases, in which a population with a continuous

phenotype distribution interacts dynamically with the environment to undergo dramatic

shifts in population distributions, pose complex—yet unanswered—questions about the

importance of phenotypic heterogeneity to population-level fitness in diverse

environments.

Here we present a novel example of a continuously-distributed phenotype in an

environmentally relevant microorganism, *Methylobacterium extorquens*, and describe

the dynamics of that phenotype distribution in response to shifts in its growth

environment. *M. extorquens*, a species of *Alphaproteobacteria* found ubiquitously on

plant leaves, is a methylotroph: it can grow on reduced single-carbon compounds such

as methanol, which is emitted from leaves through the activity of plant pectin

methylesterases [25]. The first intermediate in methanol metabolism is the potent toxin

formaldehyde [26,27], an electrophile that can cause cellular damage through its

reactions with macromolecules, and is lethal to microorganisms [28]. In *M. extorquens*,

formaldehyde is produced in the periplasm through the oxidation of methanol by

methanol dehydrogenase (MDH); in the cytoplasm it is then oxidized via a

tetrahydromethanopterin (H_4MPT)-dependent pathway. *M. extorquens* therefore exists

in a tension between the two goals of rapid substrate utilization and prevention of the

accumulation of a toxic intermediate. The importance of formaldehyde oxidation for

single-carbon metabolism has been demonstrated by the inability of H_4MPT-pathway

mutants to grow in the presence of methanol when they possess a functional MDH [29].

However, although MDH activity is constitutive, it has also been observed that

downstream single-carbon assimilation pathways are up-regulated only in the presence

of methanol; the consequence is that when cells previously grown on a multicarbon

substrate are first exposed to methanol, formaldehyde production initially outpaces

consumption so drastically that it is excreted into the medium [26]. This is just one

example of many potential situations in which cellular formaldehyde accumulation poses

a threat to *M. extorquens*, and yet we know very little about how the species copes with

formaldehyde toxicity.

As an initial step toward understanding the effect of formaldehyde toxicity on *M.*

extorquens, we conducted time-kill experiments in which we exposed cells to a range of

formaldehyde concentrations in batch liquid culture and monitored their viability over
time. To our knowledge, previous research on formaldehyde toxicity in *M. extorquens*
has consisted only of single-timepoint shock experiments, or a pulse of methanol added
to succinate-grown cultures to cause cellular formaldehyde production [29]. We hoped
that making time-resolved measurements in extreme formaldehyde conditions would
shed light on both large-scale patterns of toxicity and on cell behavior near the minimum
bactericidal concentration (MBC). Indeed, quite unexpectedly, we found cell-to-cell
variation in the MBC within isogenic populations of *M. extorquens*. We confirmed this to
be a phenomenon of phenotypic diversity at the single-cell level, and investigated its
dynamic population-level consequences, using a combination of liquid culture
experiments, colony growth tracking, flow cytometry, time-lapse microscopy, genome
resequencing, and mathematical modeling.
191
192
193 **<u>Results</u>**
194
Formaldehyde-induced death occurs at an exponential, concentration-dependent
rate at 5 mM and above
In order to better understand the physiological effects of the toxic metabolite
formaldehyde on *M. extorquens*, we initially conducted a series of experiments in which
we added formaldehyde to methanol growth medium and assessed the relationship
between toxin concentration and mortality in well-mixed liquid batch culture. We found
that concentrations ≥5 mM elicited loss of viability (as measured in colony-forming units,
CFUs) at an exponential rate, and the rate of death increased with increasing
formaldehyde concentration (Fig. 1). A concentration of 15 mM was sufficient to
eliminate all detectable viable cells within 1.5 hours (approximately half of one
generation time). These time-kill curves indicate that formaldehyde-induced death can
be modeled as a single-hit process [30], as is often observed for other multi-target lethal
agents such as radiation and heat [31], and some bactericidal antibiotics [32,33]. Given
that the precise mechanism of formaldehyde-induced mortality remains unknown but
likely involves multiple cellular targets, it is noteworthy that formaldehyde-induced death

over time did not seem to involve substantial "shoulders" or "tails" as is sometimes
observed with other agents [33], and that we observed no saturation of death rate at the
concentrations tested. In analogy to bactericidal antibiotics, our results suggested that
the minimum bactericidal concentration (MBC) of formaldehyde for *M. extorquens* is 5
mM.
215
During exposure to moderate formaldehyde concentrations, population trajectory
abruptly shifts from decreasing to increasing over time
Follow-up experiments on longer timescales (3–4 days) revealed that assessing MBC
was in fact not straightforward: lower concentrations had an effect on *M. extorquens*
growth as well. Exposure to formaldehyde concentrations between 3 and 5 mM allowed
normal growth of *M. extorquens* as measured by optical density (OD_{600}), but only after
an apparent lag time of several hours to days. Higher formaldehyde concentrations
induced longer lags, but growth subsequently resumed, and formaldehyde
concentration had no effect on growth rate (for 0, 3, 4, and 5 mM respectively, specific
growth rate (*r*) was 0.212±0.038, 0.208±0.045, 0.238±0.002, and 0.230±0.018 h^{-1},
where ± indicates 95% confidence interval; *p*=0.237 for the effect of treatment group on
growth rate by ANOVA) (Fig. S1). To better understand the apparent lag in these
conditions, we measured cell viability over time during formaldehyde exposure
experiments. CFU measurements revealed that the apparent "lag time" was in fact not a
lag, but rather a period of exponentially decreasing cell counts followed by an abrupt
transition to increasing counts (Fig. 2, round gray symbols and line). These dynamics
repeated themselves across multiple biological replicates at multiple formaldehyde
concentrations, with remarkable consistency in the timing of the transition between
population decrease and increase. At a concentration of 4 mM formaldehyde, this
transition occurred at approximately 20 hours. Upon recovery, the increase in CFUs
was exponential and indistinguishable from growth rates on methanol in the absence of
external formaldehyde (*r*=0.215±0.001 for 0 mM; *r*=0.202±0.030 for 4 mM; *p*=0.063 by
Welch's t-test) (Fig. 2).
239

We explored several hypotheses to explain the abrupt decrease-then-increase transition

we observed during formaldehyde exposure. These included: 1) consumption of

formaldehyde by the cells; 2) the existence of formaldehyde-resistant genetic mutants;

3) phenotypic changes in the plating efficiency of cells due to formaldehyde-induced

damage and its repair; 4) the existence of a subpopulation of phenotypically (but not

genetically) formaldehyde-tolerant cells.

Population recovery occurs while formaldehyde concentrations in the medium

remain high

One explanation for the abrupt change from population decline to population increase

might be a change in environmental conditions: for instance, cooperative behavior to

metabolize a toxin to sublethal levels can result in population rescue even if not all cells

are tolerant, as has been observed with antibiotics [34]. We therefore investigated the

possibility that formaldehyde consumption had reduced the toxin concentration in the

medium: in formaldehyde exposure experiments with 4 mM formaldehyde, we

monitored formaldehyde concentrations in the medium for 80 hours, a period that

encompassed the death and regrowth phases. Although *M. extorquens* is capable of

metabolizing formaldehyde, we found that a measurable decrease in the concentration

of the toxin only occurred at the very end of the growth period in batch culture, more

than 40 hours after the decrease-then-increase transition (Fig. 2, black symbols and

line). A change in the toxicity of the environment was therefore not responsible for the

change in population trajectory.

Genetic mutations are not responsible for population recovery

The next most parsimonious explanation was that the observed growth was due to a

small pre-existing subpopulation of formaldehyde-tolerant mutants whose existence

became apparent only after the death of the sensitive majority. To assess this

possibility, we grew cells in the presence of 4 mM formaldehyde for 80 hours as

described above, then subcultured them into fresh medium without formaldehyde for 6

generations (using succinate as the growth substrate), followed by another subculture

into formaldehyde-containing medium. The population decrease-then-increase

dynamics were recapitulated, indicating that formaldehyde tolerance was not

transmitted in a manner consistent with genetic heritability and that the descendants of

cells tolerant to 4 mM formaldehyde were re-sensitized in its absence (see "Transitions

between tolerance phenotypes," below, for more detail).

In addition, we prepared genomic DNA from cells that had grown in the presence of

formaldehyde (the 80-hour timepoint of a 4 mM formaldehyde exposure experiment),

and used this for whole-genome resequencing. The resequenced genome was

compared to the published genome sequence of wild-type *M. extorquens* PA1 and to

that of an *M. extorquens* population grown without formaldehyde; no evidence was

found for SNPs, deletions, insertions, or gene duplications in the formaldehyde-selected

population. These sequencing data, along with the instability of the formaldehyde

tolerance phenotype, indicate that the heterogeneous formaldehyde tolerance we have

observed in *M. extorquens* is not due to genetic mutations.

Population recovery is not due to changes in plating efficiency

Having ruled out both environmental change and genetic mutations as explanations for

the sharp transition between population decrease and increase during formaldehyde

exposure, we pursued the possibility that the observed dynamics might be due to a

change in phenotype. As our evidence for population dynamics thus far was based on

counts of colony-forming units on agar medium, one possibility was that the phenotypic

change might be related to the ability of cells to form colonies. In this scenario,

formaldehyde exposure might cause cellular damage resulting in a decrease in cells'

ability to form colonies and their entry into a viable-but-not-culturable (VBNC) state [35],

and the inflection point at 20 hours represented the beginning of recovery of these same

cells, rather than turnover of the population.

To investigate whether there had been population turnover, we carried out a cell

proliferation assay, which enabled us to interrogate single-cell growth dynamics in liquid

culture without plating (as in [36]). We used a nontoxic fluorescent membrane linker dye

to stain the population prior to conducting a 4 mM formaldehyde exposure experiment

as described above, and then assessed the trajectory of per-cell fluorescence over time
by flow cytometry. In exponentially growing populations, the membrane dye of each
parent cell was divided between its two daughter cells, and per-cell fluorescence
decreased uniformly across the population as the number of cells increased (Fig. 3,
upper left). In non-growing populations, as when *M. extorquens* was exposed to 20 mM
formaldehyde, membrane dye remained in the cells and did not fade, and per-cell
fluorescence remained the same (Fig. 3, lower left). However, in an *M. extorquens*
population treated with 4 mM formaldehyde, two populations were evident: one group of
cells that retained the same membrane fluorescence throughout the experiment, and
another group of cells that increased in number and decreased in per-cell membrane
fluorescence, indicating normal growth in the presence of formaldehyde (Fig. 3, right).
At 4 mM, the growing population was observable only after ~37 hours of incubation,
consistent with a population that began in extremely low abundance relative to the non-
growing population. These data indicate that there was population turnover, and that the
increase in viable cell counts was in fact due to the growth of a tolerant, rare
subpopulation of cells that remained active in the presence of formaldehyde.
318
Phenotypically tolerant subpopulation is present prior to formaldehyde exposure
Given our evidence against genetic mutants, a fourth hypothesis was that the observed
growth was due to a small subpopulation of cells that were phenotypically (but not
genetically) highly tolerant to formaldehyde. Stress-tolerant individuals may arise in
microbial populations stochastically, or they may do so in response to an environmental
signal [37]. To investigate whether the hypothesized formaldehyde-tolerant cells were
present in the original population or were induced during the course of formaldehyde
exposure, we monitored the abundance of formaldehyde-tolerant CFU over time.
Specifically, we repeated the 4 mM formaldehyde exposure experiment and plated the
cells harvested at each timepoint on selective agar culture medium (4 mM
formaldehyde, allowing the growth of only tolerant cells) and permissive medium
(without formaldehyde, to enumerate all cells). We found that at the beginning of the
experiment, the *M. extorquens* population already contained a detectable subpopulation
of cells that were able to form colonies in the presence of 4 mM formaldehyde. This

subpopulation comprised only a small portion of the total number of cells (a frequency of

~10^{-4} in the total population of ~2×10^6 cells) (Fig. 2, orange diamond symbols and lines).

And while the total abundance of cells decreased at an exponential rate between 0 and

20 hours, the formaldehyde-tolerant subpopulation increased at a constant rate for

nearly the entire course of the experiment, such that after 20 hours the population was

dominated by cells tolerant to 4 mM formaldehyde.

Formaldehyde-tolerant subpopulation shows no evidence of formaldehyde-

induced cell damage

The fact that the tolerant subpopulation increased at a rate commensurate with

standard growth in formaldehyde-free media suggests that these tolerant cells may

have escaped formaldehyde-induced mortality and any damage that might impede or

delay growth. However, some stress-tolerant phenotypes, such as persister cells, are

characterized by slow growth or even absence of growth [12]. We therefore looked

more closely at the growth phenotypes of the tolerant subpopulation: specifically,

whether they differed in their rate of colony growth, or in the time required to initiate a

colony, relative to normal unstressed cells. We repeated the 4 mM formaldehyde

exposure experiment described above, this time using time-lapse imaging to monitor the

growth of colonies resulting from timepoint samples plated on both selective and

permissive plates. Plates were incubated on a flatbed photo scanner and images

captured hourly and processed to extract per-colony statistics (Fig. S2), in a manner

similar to previous studies investigating cell damage and dormancy at the single-cell

level [38–40].

We found that within the total population, which consisted of sensitive cells at a

frequency of 10^{-5} at the beginning of the experiment, formaldehyde exposure prior to

plating did indeed induce a lag in colony formation, such that each hour of formaldehyde

exposure led to an increase in colony arisal time (the time necessary to form a

detectable colony) of approximately 4.80 hours (r^2=0.766 by linear regression, *p*<0.005)

(Fig. 4). This pattern held true for the first 16 hours of exposure, until most of the

sensitive cells lost viability entirely. At 16 hours, we observed a clear bimodality on the

permissive plates, with approximately half of the cells arising late and the other half

arising early, and the abundance of the early arisers on the permissive plates was

consistent with the abundance of the tolerant subpopulation (all colonies on the

selective plates). However, on the selective plates, we observed no relationship

between arisal time and the length of formaldehyde exposure prior to plating

(slope=-0.091 hours lag per hour exposure, r^2=0.0639, p<0.005), supporting the

hypothesis that the tolerant subpopulation is in a different physiological state that does

not experience formaldehyde damage in the same way as the sensitive subpopulation.

Furthermore, we observed no trend in average growth rate due to formaldehyde

exposure time (slope=-2.83x10^{-4} hours lag per hour exposure, r^2=6.62x10^{-4}, p=0.123),

although colonies from samples that had been exposed for longer had more varied

growth rates (there was a positive linear relationship between exposure time and the log

of the median average deviation of growth rate, p<0.05 for both permissive and

selective plates).

Tolerant subpopulation behavior is observable at the single-cell level

As a final step in confirming single-cell phenotypic heterogeneity in formaldehyde

tolerance, we observed growth of single cells directly using high-resolution time-lapse

phase-contrast microscopy of cultures embedded in agar pads. This method allowed us

to observe the division times and potential morphological aberrations of individual cells

and their progeny in microcolonies over the first 12 hours of growth. Because the cells

tolerant to 4 mM formaldehyde are only present at a frequency of approximately 10^{-4} in

the initial population, we conducted formaldehyde exposure at a lower concentration,

2.5 mM formaldehyde, at which plating experiments (see below) suggested that ~1-4%

of cells would be able to grow. Indeed, we found that 11 out of 546 cells (1.97%) were

able to grow at that formaldehyde concentration, compared to 100% of the 256 cells

observed in the no-formaldehyde condition (Fig. 5). In addition, all the cells that grew in

the presence of formaldehyde did so normally: we found no significant effect of

formaldehyde on cell division time (median doubling time at 0 mM: 2.58 hours; at 2.5

mM: 2.58 hours; p=0.262 by Mann-Whitney U-test). We did observe that cells in

formaldehyde took slightly longer to complete the first division (median lag time was

1.25 hours later in formaldehyde; $p<0.001$, Mann-Whitney U); this may indicate minor

cell damage or a very modest inhibitory effect of formaldehyde in the medium, or it may

have resulted from slight differences in the preparation of the cells (as it was technically

infeasible to conduct the two experiments simultaneously). Of the variance in doubling

times among individual cells, most was explained by the microcolony to which the cell

belonged ($p=0.001$, PERMANOVA) but not by formaldehyde treatment ($p=0.323$),

indicating that there is some heritability in growth rates (Fig. 5). Finally, we did not

witness any partial-growth or impaired-growth phenotypes: the 535 cells that were

unable to grow in the presence of formaldehyde showed no detectable elongation or

other change in morphology.

Formaldehyde-tolerant cells grow at normal rates on methanol but marginally

slower on succinate

Our single-cell observations of growth suggested that formaldehyde-tolerant cells grow

normally; the formaldehyde tolerance phenotype thus differs from other microbial stress-

tolerance phenotypes previously described (e.g., sporulation, persistence) in that it does

not require cells to sacrifice proliferative ability by entering a non-growing state in order

to survive stressful conditions. Moreover, our observations of colony arisal times

suggested that tolerant cells avoid the damage incurred by formaldehyde-sensitive

cells. We therefore investigated the possibility that other fitness tradeoffs might exist, by

comparing non-selected *M. extorquens* cells with those of a tolerant population (i.e.,

selected by growth in liquid culture at 4 mM formaldehyde).

We observed no difference in the resistance of the tolerant population to several other

chemical stressors (hydrogen peroxide, and the antimicrobial compounds rifampicin,

vancomycin, cefoxitin, novobiocin, nalidixic acid, ciprofloxacin, erythromycin,

kanamycin, gentamicin, chloramphenicol, colistin) during growth on agar medium (data

not shown). However, we did observe differences in growth rate when comparing the

tolerant and naive (non-selected) populations side-by-side in liquid batch culture,

depending on whether the growth substrate was methanol (15 mM) or the non-

methylotrophic substrate succinate (3.5 mM). In methanol medium, the two populations

grew at similar rates (naive: $r=0.222\pm0.023$; tolerant: $r=0.202\pm0.051$ h^{-1}); however, on

succinate medium, the naive population grew faster (naive: $r=0.270\pm0.002$ h^{-1}; tolerant:

$r=0.215\pm0.030$ h^{-1}) (Fig. S3). The differences in growth rate among the four treatments

were in this case not found to exhibit statistically significant differences by ANOVA

($F=2.617$, $p=0.123$ for the model; $p=0.940$ for the planned contrast between the two

populations on methanol and $p=0.136$ on succinate). However, the apparent slight

advantage of the naive population over the tolerant on succinate was attributable to the

fact that the naive population grew faster on methanol than on succinate, consistent

with observations commonly made in our lab (Fig. S7) and reported in the literature [27].

If the formaldehyde-tolerant population does not show the typical increased growth rate

on succinate, it would indicate that tolerance is associated with methylotrophic

metabolism, and may provide a clue as to the conditions in which tolerant cells could be

out-competed by sensitive cells in the environment.

Phenotypic heterogeneity in formaldehyde tolerance is a continuous phenotype

As described previously, many cases of phenotypic heterogeneity display binary

phenotypes (e.g., persistent or not [12]). On the other hand, a few display a continuous

distribution (e.g., resistance to chloramphenicol along a gradient of possible MBC levels

[23]). While formaldehyde exposure results in a binary outcome for each cell—either

growth, or cessation and death—we sought to quantify the ratio of these outcomes

along a gradient of formaldehyde concentrations, by plating *M. extorquens* onto agar

medium containing formaldehyde at concentrations between 1 and 10 mM in

increments of 1 mM. The results showed no evidence of a bimodal distribution along the

concentration axis; rather, tolerance is continuous, peaks at 0 mM, and declines

exponentially and predictably with increasing formaldehyde concentration (Fig. 6). We

were able to detect cells with tolerance levels as high as 6 mM. While we found growth

stage to have an effect on the abundance of tolerant cells (populations in exponential

growth were shifted toward slightly higher tolerance), the qualitative shape of the

distribution remained the same. Furthermore, populations that were cultured in the

same way and sampled at the same growth stage reproduced similar formaldehyde

tolerance distributions across multiple days (Fig. S4). Notably, all colonies that grew on

formaldehyde agar medium were of uniform shape and size, suggesting that tolerant

cells shared similar arisal times and growth rates, as was previously observed at 4 mM

(Fig. 4). In addition, cell proliferation assays with the membrane-intercalating dye

performed at concentrations of 2, 3, and 5 mM formaldehyde qualitatively recapitulated

the previously described population turnover seen with the assay at 4 mM, and

supported the relationship between concentration and abundance of tolerant cells (Fig.

S5). Thus, although the consequence of formaldehyde upon an individual cell is binary

(growth or not, rather than a range of growth rates), the distribution of tolerance (i.e.,

MBC values) within a population is continuous.

Transitions in tolerance phenotypes over time depend on growth conditions

Given our previous observations that growth conditions can change the shape of the

formaldehyde tolerance distribution in a population, and that formaldehyde tolerance,

once selected for, can be lost, we sought to characterize more precisely the processes

by which tolerance distributions might shift in a population. Specifically, we hoped to

better understand the relative importance of formaldehyde-mediated selection (whereby

formaldehyde exposure kills sensitive cells) and active changes made by cells to alter

phenotype. We began by monitoring the abundance of each subpopulation, at tolerance

levels between 0 and 10 mM in 1 mM intervals, over time during a 4 mM formaldehyde

exposure experiment. We monitored for 20 hours, the time it takes for most sensitive

subpopulations to lose viability entirely (Fig. 2). As expected, we found that in these

conditions, all subpopulations tolerant to <4 mM decrease in abundance and the

subpopulations tolerant to ≥4 mM increase, resulting after 20 hours in a tolerance

distribution with a maximum at 4 mM (Fig. 7, S6).

In order to measure the rate at which phenotypic tolerance in the population returns to

its original distribution, we conducted a formaldehyde-free regrowth experiment.

Specifically, we transferred the selected, high-tolerance population to liquid medium

without formaldehyde and monitored the changes in tolerance distributions for the next

24 hours in two different conditions: one with methanol as the sole carbon substrate,

and the other with succinate. We observed a marked difference between the two growth

conditions in their effect on population tolerance distributions over time. In the succinate
medium, only the populations with low tolerance increased in abundance, whereas
those with high tolerance *decreased* in abundance, so the shape of the distribution
shifted back toward that of naive *M. extorquens* cells (Fig. 7, S6). The observation that
tolerant cells decreased in abundance even during an increase in the overall population
suggests that cells were shifting in phenotype from high tolerance to low. In contrast, in
the methanol medium, all tolerant subpopulations increased in abundance at the same
rate: the overall shape of the distribution, with its high proportion of tolerant cells, stayed
the same. Growth in methanol medium thus maintains phenotypic formaldehyde
tolerance in a population that is *already* tolerant, even though it does not induce or
select for tolerance in sensitive populations. This unexpected substrate-based
hysteresis (historical dependence) may be due to the small amount of formaldehyde
produced inside the cell during methylotrophic metabolism, which might trigger cells,
either through a stress-response mechanism or through regulation of methylotrophic
metabolism, to remain in a tolerant phenotype even if external formaldehyde is not
present in the growth medium.
504
Mathematical modeling elucidates cellular phenotype transition processes
underlying population-level shifts
To better understand what biological processes might be responsible for the
observations of the three experiments described above, we developed a mathematical
model to test several hypotheses. We examined the dependence of death rate upon
formaldehyde concentration in the medium and tolerance phenotype of the cell. We also
asked whether shifts in the tolerance distribution could be explained by growth and
selective death alone, or involved other processes. To this end, we tested the effect of
introducing two processes by which cells might actively transition along the 1-
dimensional axis of "phenotype space": one involving random phenotype transitions in
any direction (a Brownian motion, or diffusion, process), and one involving directed
transitions toward either higher or lower tolerance (an advection process) (Fig. 8). We
evaluated models based on their ability to reproduce the dynamics from the three
experimental conditions shown in Figure 7: selection during a 4 mM formaldehyde

exposure experiment, and formaldehyde-free regrowth of the selected high-tolerance

population on either succinate or methanol.

We modeled the dynamics of *M. extorquens* populations during exposure to

formaldehyde with a partial differential equation:

$$\partial_t N(x,t) = r_c N(x,t) - H(x,F)N(x,t) + D\,\partial_{xx}N(x,t) + v\,\partial_x N(x,t) \qquad (1)$$

The population is structured by phenotype, with $N(x,t)$ denoting the concentration of

cells (CFU*mL^{-1}) with formaldehyde tolerance x (mM) at time t (hours). The

formaldehyde tolerance of a cell is defined as the maximum concentration of

formaldehyde in which the cell can grow. The model tracks cells in a well-mixed, closed

population as they grow on substrate c at per capita rate r_c (h^{-1}), die at per capita rate

$H(x,F)$, and change phenotype (Fig. 8). We assume that phenotypic transitions occur

due to two processes, diffusion with coefficient D (mM^{2}*h^{-1}) and advection with rate v

(mM*h^{-1}) (where mM refers to tolerance level). The death function $H(x,F)$ describes per

capita death rate of cells as a function of formaldehyde concentration and is given by:

$$H(x,F) = \alpha(F - bx) \text{ if } x < F \qquad\qquad\qquad\qquad (2)$$

where F (mM) is the formaldehyde concentration, α (mM^{-1}*h^{-1}, where mM refers to

formaldehyde in the medium) is the death rate, and b (mM tolerance / mM

formaldehyde) specifies the sensitivity of the death rate to a cell's formaldehyde

tolerance level.

For each of the three experimental conditions separately, we used maximum likelihood

to fit the parameters α, b, D, and v to the data, and used a likelihood ratio test on the

nested models to determine the best model structure. For the selection scenario, we

began with a 1-parameter model with $\alpha > 0$ and the other parameters equal to 0, and

tested 2-, 3-, and 4-parameter models sequentially, at each step choosing the model

with the highest likelihood as long as it was significantly better than the simpler model

(results in Table 1). For regrowth (where there is no death), we fit only v and D. In each

of the three experimental conditions, our model was able to reproduce the experimental

observations extremely well (pseudo-R^2 = 0.973, 0.993, 0.991 for the formaldehyde

selection, methanol regrowth, and succinate regrowth conditions respectively) (Table 1).

Phenotype transition processes change according to growth conditions

We examined not only what the best-fit value was for each of the four parameters of

interest, but also whether there was support (by likelihood ratio test) for including each

of the parameters. The three experimental conditions differed from one another in both

considerations, suggesting that the rate and nature of phenotype transition processes

change depending on environment.

In the formaldehyde selection regime, the best model included both death parameters

(a=0.202±0.022 h^{-1}, b=0.770±0.147), but for the phenotype transition parameters, only

diffusion (D=0.019±0.006 mM^{2}*h^{-1}) and not advection. This indicates that the changes

we observed in the formaldehyde tolerance distribution during formaldehyde exposure

are due not only to death, but also involve phenotype shifts consistent with diffusion.

Diffusion leads to the spread of phenotypes consistent with what was observed in the

model after 20 hours, including the presence of population density at x=2 and x=3 mM

(which otherwise would have decreased to below detection in the absence of transitions

from higher-tolerance phenotypes) and some cells at x>6 (which were not observed in

the initial population, but may have resulted from transitions from lower-tolerance

phenotypes) (Fig. 9).

For the scenarios involving regrowth of a selected population on formaldehyde-free

medium, the parameter estimates were markedly different from those in the selection

condition, and from each other. For methanol growth, we found support only for very

mild advection toward lower tolerance (v=0.018±0.006 mM*h^{-1}). For succinate growth,

we found support for both advection (v=0.285±0.026 mM*h^{-1}) and diffusion

(D=0.033±0.010 mM^{2}* h^{-1}); the advection term for succinate was an order of magnitude

greater than that for methanol, indicating strong shifts in the direction of lower tolerance

when high-tolerance cells are grown on succinate. This agrees with our earlier

qualitative observations that growth on succinate, but not on methanol, leads *M.*

extorquens populations to lose formaldehyde tolerance rapidly and in general to

undergo diversifying phenotype transitions. Furthermore, it supports our hypothesis that

formaldehyde tolerance is associated with methylotrophic growth, and implies that

further work is needed not only to understand the mechanism of phenotypic

formaldehyde tolerance shifts in *M. extorquens* but also the regulation of those

processes.

The inclusion of b in the model specifically allowed us to test the possibility that a cell's

formaldehyde tolerance might determine not only the threshold concentration above

which it dies, but also the rate at which it dies (Fig S8). That is, if $b=0$, the death rate is

equivalent for all cells regardless of tolerance level as long as tolerance is below the

threshold; but for $0<b\leq1$, cells with lower tolerance levels die more quickly than those

with higher tolerance, and the strength of this dependence upon x scales with the value

of b. The value of b that best fit our data was 0.770 (Table 1), indicating that

formaldehyde-tolerant cells may receive some protection from the tolerance phenotype

even at concentrations above their MBC. However, both the experimental results and

the model simulation showed a bimodal phenotype distribution after formaldehyde

exposure, consistent with a weak dependence of death rate from tolerance level (Fig.

S8). Future experiments with a greater number of observations at low-tolerance

phenotypes would better elucidate the relationship between tolerance and death

dynamics.

<u>Discussion</u>

We have described here a novel example of phenotypic tolerance to a metabolic toxin

that is continuously distributed across individual cells in a clonal microbial population,

where the phenotypic heterogeneity is present even when all cells inhabit the same

environment, but its distribution undergoes dynamic shifts when the population

experiences different environmental conditions. Wild-type populations of genetically

identical *Methylobacterium extorquens* cells grown in well-mixed liquid medium contain

individuals with maximum formaldehyde tolerance levels ranging from 0 mM to 6 mM

(and tolerance at up to 8 mM has been observed after selection); although the

distribution is continuous and exponentially decreasing, individuals show binary

growth/non-growth phenotypes at any given formaldehyde concentration, and may

transition among phenotypic states through both bidirectional phenotype diversification

and responsive, directed phenotype shifts.

When might *M. extorquens* experience formaldehyde in its natural environment, and at

what concentrations? *M. extorquens* excretes formaldehyde during the first stages of

the switch between multi-carbon and single-carbon metabolism [26]; in addition,

formaldehyde is a metabolic intermediate in the consumption of many lignin-derived

aromatic compounds [41,42], and we have observed lignin degraders to excrete

formaldehyde into the growth medium at millimolar levels during growth in batch liquid

culture on methoxylated aromatic compounds (Lee & Marx, unpublished). Thus, it is

possible that, in the environment, formaldehyde concentrations in the millimolar range

might accumulate transiently on the microscale, especially within cell aggregates such

as those observed on plant leaves [43]. However, it is also possible that formaldehyde

tolerance is an outward manifestation of a cellular state that has quite a different

relevance in the native environment of *M. extorquens*: high tolerance to local (external)

formaldehyde may indicate a high capacity for tolerating (internal) formaldehyde

generated through methylotrophic metabolism. As such, this phenomenon may provide

insight into processes that are general to many organisms whose central metabolic

pathways involve toxic intermediates [44,45].

The mechanisms by which high-tolerance cells can maintain normal growth in the

presence of seemingly lethal concentrations of formaldehyde, and by which that ability

is transmitted to progeny, remain yet to be discovered. Phenotypic formaldehyde

tolerance could involve a specific stress-response system. Alternatively, it may involve

components of methylotrophic metabolism, as suggested by the association between

maintenance of formaldehyde tolerance and methanol growth, and the potentially lower

fitness of tolerant cells during succinate growth. Previous reports have found phenotypic

heterogeneity in the related strain *M. extorquens* AM1 in growth rate, cell size, gene

expression levels, and ability to switch between carbon substrates [8,46,47]; those

forms of diversity may have a similar basis to the phenomenon discussed here.

Because of the crucial role that the H_4MPT pathway of formaldehyde oxidation plays in

removing cellular formaldehyde and thus allowing methylotrophic growth [29], very small

variations in the activity of H_4MPT pathway enzymes might have a significant impact on

a cell's capacity to tolerate externally-provided formaldehyde. Assuming that

formaldehyde transport into the cytoplasm is diffusion-driven (as no means of active

transport has yet been discovered), phenotypic diversity in formaldehyde oxidation

capacity would result in a range of internal concentrations across cells. Furthermore, if

cells that begin to experience formaldehyde-mediated damage to proteins and other

macromolecules begin to lose their capacity to oxidize formaldehyde, this would

generate a positive feedback circuit that could ultimately determine a binary outcome to

whether a cell lives or dies. The fact that we observed no elongation at all from the cells

that could not grow at 2.5 mM suggests that this postulated positive feedback

mechanism acts on a very fast timescale (i.e., minutes, not hours). This mechanism

could explain both the continuous relationship between concentration and abundance of

tolerant cells, and the binary nature of the "live normally or die immediately" distribution

observed at each specific formaldehyde level, similar to the topology of interactions

observed in the case of chloramphenicol resistance described earlier [23].

Viewed this way, formaldehyde tolerance exhibits potential parallels with other

examples in which phenotype switching is mediated by a bistable switch: for instance,

pyrimidine-mediated colony type switching in *Bacillus subtilis* [48], chloramphenicol

resistance [23], and lactose utilization *E. coli* [49]. In all these cases, the balance

between the concentrations of multiple intracellular components determines the

phenotype of a cell, with cells pushed away from the threshold into one of two states by

a positive feedback loop, and often maintained there by hysteresis (as we observed in

the maintenance of formaldehyde tolerance among cells growing on methanol, Fig. 7).

Making changes that alter the ratios of the components, through control of either gene

expression or of the environment, alters the frequency of cells near the threshold and

thereby the ratio of cells in each of the two bistable states (as we observed when

altering formaldehyde concentrations in the medium). Furthermore, these systems are

analogous to threshold traits observed in animals, in which observed discontinuous

phenotypes (e.g., two phenotypic morphs) arise from a continuously distributed

underlying trait [50]. The threshold model of quantitative genetics describes a constant

threshold that determines which values of the underlying trait generate each of the

phenotypic morphs; evolutionary and environmental changes can lead to shifts in the

underlying trait distributions that result in different ratios of the two morphs being

observed in different environments. As we have shown here, the continuously

distributed trait of *M. extorquens* formaldehyde tolerance similarly changes with

environment; however, in this case the threshold (environmental formaldehyde

concentration, which separates tolerant cells from sensitive) is external and may also be

manipulated. *M. extorquens* therefore provides a convenient model system in which to

probe the dynamics of threshold traits.

Both stress response proteins and metabolic enzymes are inherited during cell division,

so either pathway provides a hypothesis for the heritability of formaldehyde tolerance

that we observed after formaldehyde was removed. Lineage dependence has been

observed in numerous cases of phenotypic heterogeneity, and is often cited as

evidence that the phenotype of interest is dictated by a heritable component of the cell

with a moderately long lifetime (e.g., pole age [8], stress-protective protein aggregates

[51]). In these cases, the random transitions in phenotype that we modeled as diffusion

can be explained by asymmetric partitioning of cell contents [52], whereas the directed

transitions that we modeled as advection could be due to responsive up- or down-

regulation of the production/degradation or (in)activation of the molecules involved.

The balance between stochastic and responsive phenotype differentiation processes,

as mechanisms for increasing population fitness in unpredictable environments, has

often been discussed as a question of the relative costs and benefits of each. Both

mathematical modeling [53,54] and laboratory evolution [18,55] have demonstrated that

either mechanism—or both simultaneously—can be selected for, with evolutionary

outcomes dictated by the cost of sensing, the timing of environmental change, the

reliability of environmental cues, and the spatial structure of the community. The fact

that we observed non-directed general diversification in our populations in some growth

conditions, and environment-responsive, directed phenotype shifts in others, strongly

suggests that that differentiation in formaldehyde tolerance in *M. extorquens* is not due

simply to unavoidable molecular noise, but rather it is a regulated process conferring a

fitness advantage.

If it exists, the nature of such a fitness advantage remains unclear; it is tempting to ask

why *M. extorquens* does not maintain a population composed solely of high-tolerance

cells. The evolutionary basis for phenotypic heterogeneity in genetically identical

populations of microorganisms is frequently ascribed to diversifying bet-hedging, in

which a species in an unpredictably changing environment constitutively generates

progeny with multiple phenotypes to ensure that at least a few will thrive in any

circumstances [11,56]. The surface of the plant leaf, the native environment of *M.*

extorquens, is indeed unpredictable: cells depend for growth upon a combination of

gaseous methanol excreted from plant stomata [57] and other metabolites such as

simple organic acids produced by the plant or by other microorganisms [58]. The

emission of methanol, dependent on the metabolic state of the plant host and the

conductivity of the stomata, undergoes large temporal variation [59]. It is therefore

plausible that this environment could select for heterogeneity in formaldehyde toxicity

response and in growth on non-methylotrophic substrates. An alternative evolutionary

explanation is division of labor, which is an advantageous strategy when a particular

activity is beneficial to the population but incurs some cost to the individual carrying it

out [54]. In our experiments, the tolerant subpopulation is capable of detoxifying

formaldehyde from the growth medium; although in batch culture this occurs too late to

prevent the sensitive cells from dying (Fig. 2), this might occur differently in a spatially-

structured community such as cell aggregates on the surface of a plant leaf.

Importantly, both bet-hedging and division of labor assume fitness tradeoffs among the

phenotypes being diversified. Although tolerant cells may be slower than sensitive cells

at growth on a multicarbon substrate (Fig. S8), it is unclear whether this disadvantage is

substantial enough to explain the low frequency of high-tolerance cells in our *M.*

extorquens populations. Further mathematical modeling could be used to quantify the

effects of these growth tradeoffs and to explore conditions that could lead to the

evolution of the steady-state distribution that we have observed in the lab. Furthermore,

while phenotypic heterogeneity has been studied in great depth both in laboratory

populations and in populations simulated by mathematical modeling, there remains a

dearth of research on organisms in the natural environments in which they evolved

[56,60]. Future experiments examining the dynamics of phenotypic formaldehyde

tolerance among *M. extorquens* cells growing on plant leaves, and among related

methylotrophs in other environmental niches, will be essential to establishing the

environmental relevance of this phenomenon.

Materials and Methods

Bacterial strains and culture conditions

All experiments were conducted with *M. extorquens* PA1 CM2730, an otherwise wild-

type strain that contains a deletion of the cellulose synthesis operon to prevent cell

clumping [61]. All cultures were grown at 30 °C in MPIPES mineral medium (30 mM

PIPES, 1.45 mM K_2HPO_4, 1.88 mM NaH_2PO_4, 0.5 mM $MgCl_2$, 5.0 mM $(NH_4)_2SO_4$, 0.02

mM $CaCl_2$, 45.3 µM $Na_3C_6H_5O_7$, 1.2 µM $ZnSO_4$, 1.02 µM $MnCl_2$, 17.8 µM $FeSO_4$, 2 µM

$(NH_4)_6Mo_7O_{24}$, 1 µM $CuSO_4$, 2 µM $CoCl_2$, 0.338 µM Na_2WO_4, pH 6.7) [61], either in

liquid form or with the addition of 15 g/L Bacto-Agar (BD Diagnostics) for solid medium.

Growth substrates consisted either of methanol provided at 15 mM (in liquid medium) or

125 mM (in solid medium); or succinate provided at lower concentrations to ensure the

same molarity of carbon as in methanol conditions: 3.5 mM succinate (in liquid) or 15

mM (in solid). Unless otherwise noted, all liquid cultures were grown as a 20 mL volume

in 50 mL-capacity conical flasks, capped with Suba-Seal rubber septa (Sigma-Aldrich)

to prevent the escape of volatile compounds, with shaking at 250 rpm. Extensive testing

in our lab has shown that cultures grown thus are not limited by oxygen availability and

grow at the same rate as in flasks with loose lids. Prior to the beginning of each

experiment, cultures were streaked from freezer stock onto solid medium and allowed to

grow for 4 days to form colonies, and for each biological replicate a single colony was

used to inoculate 5 mL of liquid medium in a culture tube and allowed to grow 24 hours.

This overnight culture, containing ~2×10^8 CFU/mL at stationary phase, was used as the

inoculum for the growth experiment.

Formaldehyde exposure experiments

Growth and tolerance distribution dynamics in the presence of low concentrations of

formaldehyde were assessed by inoculating stationary-phase culture at a 1:64 dilution

(for an initial population of ~3×10^6 CFU/mL) into fresh MPIPES + methanol medium

containing formaldehyde at the specified concentration (4 mM unless otherwise stated).

Cultures were grown under the conditions described above (20 mL MPIPES, Suba-Seal

septa, shaking at 30 °C) and sampled periodically to measure cell viability and

formaldehyde concentration. Sampling was conducted as follows: 100 µL of culture was

removed through the septum using a sterile syringe and transferred to a trUVue low-

volume cuvette (Bio-Rad) to read optical density at 600 nm using a SmartSpec Plus

spectrophotometer (Bio-Rad). An additional 200 µL of culture was removed and

transferred to a microcentrifuge tube. The culture was centrifuged for 1 minute at 14,000

x g; the supernatant was removed and saved for formaldehyde measurement (see

below) and the cell pellet was resuspended in 200 µL fresh MPIPES medium without

carbon substrate or formaldehyde. Cells were then subjected to serial 1:10 dilutions in

MPIPES to a final dilution of 10^{-6}. From each of the seven dilutions, three replicates of

10 µL were pipetted onto culture plate containing MPIPES-methanol solid medium to

form spots (total: 21 spots per sample per plate type). In experiments measuring

formaldehyde tolerance distribution, multiple plate types were used, each type

containing a different concentration of formaldehyde (see "Formaldehyde tolerance

distributions," below). The spots were allowed to dry briefly in a laminar flow hood, then

lids were replaced and plates were stored in plastic bags and incubated at 30 °C for 4

days before colonies were counted. For each replicate set of seven spots, the two
highest-dilution spots with countable colonies were enumerated and summed, then
multiplied by 1.1 times the lower of the two dilution factors to calculate the original
number of colony-forming units (CFU) in the sample. For each sample, the mean and
standard deviation of the three replicate spot series was calculated.
802
For the measurement of death rates in the presence of high concentrations of
formaldehyde, cells were grown in MPIPES medium with methanol and no
formaldehyde until they reached an OD_{600} of 0.1 (~ 1×10^8 CFU/mL, mid-exponential
phase), then formaldehyde was added to the desired final concentration and
immediately mixed well. Growth conditions were as described above and samples were
taken every 20 minutes to measure formaldehyde concentrations and cell viability.
Growth rates were measured by fitting a linear relationship between time and the binary
logarithm of CFU/mL using the lm function; differences among growth rates were
assessed using either the anova function or the t.test function, in the stats package.
812
Fresh 1 M formaldehyde stock was made weekly by combining 0.3 g paraformaldehyde
powder (Sigma Aldrich), 9.95 mL ultrapure water, and 50 µL 10 N NaOH solution in a
sealed tube, and immersing in a boiling water bath for 20 minutes to depolymerize.
Formaldehyde was measured using the method of Nash [62]. In brief, equal volumes of
sample (or standard) and Nash Reagent B (2 M ammonium acetate, 50 mM glacial
acetic acid, 20 mM acetylacetone) were combined in a microcentrifuge tube and
incubated for 6 minutes at 60 ºC. Absorbance was read on a spectrophotometer at 412
nm. For experiments involving large numbers of samples, the same assay was
conducted in a 96-well polystyrene flat-bottom culture plate (Olympus Plastics): 100 µL
each of sample and Reagent B were combined in each well of a culture plate and the
plate was incubated at 60 ºC for 10 minutes before reading absorbance at 432 nm on a
Wallac 1420 Victor2 multilabel plate reader (Perkin Elmer). A clean plate was used for
each assay; each plate contained each sample in triplicate as well as a standard curve
run in triplicate.
827

Genome resequencing of formaldehyde-tolerant subpopulation

Whole-genome sequencing was conducted on an *M. extorquens* population selected for

high formaldehyde tolerance, to determine whether there were any genetic changes

associated with the formaldehyde tolerance phenotype. In addition, the genomes of two

non-selected *M. extorquens* populations were prepared, sequenced, and analyzed at

the same time using the same methods, as controls to enable us to distinguish

mutations specific to formaldehyde tolerance from any that may have accrued in our

laboratory strain since the sequencing of the published genome.

Tolerant cells were selected via a 4 mM formaldehyde exposure experiment as

described above, and allowed to grow until stationary phase; the tolerance distribution

of each population was assayed to confirm that cells were indeed 100% tolerant to 4

mM formaldehyde (see "Formaldehyde tolerance distributions," below). From each of 3

replicate populations, 2 mL of culture (~2.2×10^8 CFU/sample) were harvested. Genomic

DNA was extracted using the Epicentre Masterpure Complete DNA and RNA

Purification Kit (Epicentre/Illumina), following the manufacturer protocol for DNA

purification, and the three populations were pooled. Library preparation was carried out

by the IBEST Genomic Resources Center at the University of Idaho (Moscow, ID):

genomic DNA was used to make shotgun libraries using the TruSeq PCR-free Library

Prep Kit (Illumina) with the HiSeq-length insert option (short), amplified using KAPA

(Illumina), cleaned using magnetic beads (MagBio), and quantified using fluorometry

prior to pooling. The pooled library was quality-checked by Fragment Analyzer

(Advanced Analytical) and quantified with the KAPA Library Quantification Kit for ABI

Prism (Kapa Biosystems). Sequencing was conducted in a 1x100 run on a HiSeq 4000

at the University of Oregon Genomics and Cell Characterization Core Facility (Eugene,

OR); reads were demultiplexed by the facility using bcl2fastq Conversion Software

(Illumina).

Genomic data was analyzed for evidence of mutations using breseq 0.32.1 [63] with

default settings, using the published genome of *M. extorquens* PA1 (NC_010172 [64])

as a reference, and comparing with a similar dataset from an *M. extorquens* CM2730

population that had been grown without formaldehyde. No predicted mutations were

found in the formaldehyde-tolerant population that were not also found in the non-

formaldehyde population. Three genomic loci (MEXT_RS13110, MEXT_RS12285/

MEXT_RS12290 intergenic region, MEXT_RS02695) carried SNPs identified as

"marginal mutation predictions" at a frequency of <33%, but further analysis revealed

these to be due to assembly errors in extremely repeat-rich regions that appear in both

the tolerant and control samples. Areas containing marginal mutation predictions were

further checked by PCR amplification and Sanger sequencing of the original DNA.

Primers used were 5'- CTCTCCGCCGAAGTGGT-3' and 5'-

GCCTTCCTCGGGTTCAAGGG-3' (for MEXT_RS02695); 5'-

CAGGGAACGCTCGTAGAGG-3' and 5'- CCACCGTGAAACGCACCGTA-3' (for

MEXT_RS12285-RS12290); and 5'-GTAGACCGCCTCCGAGACTT-3' and 5'-

GTAGACCGCCTCCGAGACTT-3' (for MEXT_RS13110). Genome resequencing data

have been deposited in the NCBI SRA database under BioProject PRJNA504295.

Cell proliferation assay

Growth versus non-growth phenotypes in bulk liquid culture were assessed using a cell

proliferation assay with the non-toxic green fluorescent membrane linker dye PKH67

(Sigma-Aldrich). To ensure that *Methylobacterium* cells could be easily distinguished

from background events in flow cytometry, experiments were conducted using *M.*

extorquens CM3839, a strain identical to CM2730 but constitutively expressing the red

fluorescent protein mCherry at the *hpt* locus [65]. Stationary-phase cultures were

stained and washed following the manufacturer protocol for PKH67, modified only in

that all centrifugation steps were carried out for 1 minute at 14,000 x g. OD_{600} was

measured after staining to account for any loss of cells, and inoculation density was

adjusted to ensure an initial population of ~$3x10^6$ CFU/mL. Stained cells were used in a

formaldehyde exposure experiment as described above, with treatments at 0, 1, 2, 3, 4,

5, and 20 mM formaldehyde. Unstained cells were grown alongside the stained cells as

a control to assess whether staining affected growth rate or viability; no measurable

difference in growth was detected. Samples were taken periodically by syringe and

washed of formaldehyde as described above; instead of plating for viability, they were

resuspended in 1 M (3%) formaldehyde as a fixative and stored at 4 °C until analysis by

flow cytometry. Immediately prior to analysis, cells were centrifuged once more and

resuspended in fresh medium to remove excess formaldehyde.

Flow cytometry was conducted in the IBEST Imaging Core at the University of Idaho

using a CytoFLEX S benchtop flow cytometer (Beckman Coulter); each sample was

analyzed at a flow rate of 10 µL/min for 3 minutes to ensure that an equal volume was

examined from each. Output was gated to allow only events with a mCherry signal

(ECD-area channel, excitation: 488 nm, emission: 610/620 bp) $>10^3$. Per-cell membrane

fluorescence was measured in the FITC-area channel (excitation: 488 nm; emission:

525/40 bp). Data analysis was conducted in R v.3.4.3 [66] using the flowCore package

[67] to interpret fcs files and the ggplot2 package [68] to generate plots.

Formaldehyde tolerance distributions

The distribution of formaldehyde tolerance phenotypes within a population was

assessed by plating cell cultures onto agar plates containing formaldehyde. MPIPES

medium was prepared with agar, autoclaved, and cooled to 50 ºC; then methanol and

formaldehyde to the desired final concentration were rapidly mixed in, and the agar was

poured into 100 mm petri dishes. The dish lids were immediately replaced and plates

were cooled on the benchtop. Plates were stored at 4 ºC and used within 1 week of

pouring. CFU were plated and enumerated as described above. This method has a limit

of detection of 1.65×10^{-7} (an abundance of 34 CFU/mL is necessary to observe 1 cell

per 30 µL plated, and the total cell population tested was 2×10^8 CFU/mL; therefore the

least-abundant subpopulation that could be detected, disregarding the effects of

Poisson distributions at lower λ, is one with an average frequency of 1.65×10^{-7} within

the total population).

To assess whether formaldehyde concentrations in agar culture plates changed over

time due to volatilization, plates containing 0, 2, 4, 6, 8, and 10 mM of formaldehyde

were assayed for formaldehyde content before and after a 3-day incubation, stored

together in the same bag, at 30 ºC. A small amount of agar (~0.1 g) was excised from

the plate, melted, diluted 1:10 in MPIPES medium, and assayed using the Nash

protocol (described above). Each plate was assayed in triplicate. No change in

concentration was detected in any of the plates (Fig. S9).

Colony arisal time and growth rates

To measure the effect of formaldehyde damage on colony time and growth rates, colony

growth was monitored using Epson Perfection V600 flatbed photo scanners.

Formaldehyde exposure experiments were carried out on *M. extorquens* as described

above, but diluted samples were spread-plated rather than spotted onto culture plates:

100 µL of one dilution was spread on each plate. Lids were lined with sterile black felt to

reduce condensation and increase contrast, and plates were placed agar-side-down on

scanner beds. Each culture contained precisely 30 mL of agar medium, to ensure

uniformity of nutrient supply and hydration status across all plates; temperature probes

were included between plates on scanner beds to monitor temperatures for consistency.

Scanners were placed in a 30 °C incubator and image acquisition was controlled by a

computer running Linux Mint, using a cron job for scheduling and a custom bash script

employing the utility scanimage to take images once per hour.

Images were processed using a custom Python 3.5.6 script employing scikit-image

v.0.12.1 [69] to identify colonies and measure their areas in pixels. Subsequent data

analysis was conducted in R: double colonies and non-colony objects were removed

from the dataset, colony arisal time was measured as the first timepoint at which colony

area measured greater than 100 pixels, and colony growth rate was measured by fitting

a linear relationship between time and the binary logarithm of colony area using the lm

function. Variability among growth rates was assessed as the median average deviation

(MAD) within the colonies from a single timepoint, using the mad function in the stats

package. The effect of formaldehyde exposure time on colony arisal time, colony growth

rate, and MAD was calculated using simple linear models using the lm function. Further

details of both image analysis and colony growth statistics are included in Supplemental

Materials (Fig. S2).

Time-lapse microscopy

To analyze the formaldehyde-dependent heterogeneous response in lag-phase and elongation rates of *M. extorquens*, we employed single-cell time-lapse microscopy using a phase-contrast inverted microscope (Leica, DMi8) equipped with an automated stage. For image acquisition, we employed a 60x (PH2, NA: 0.7) magnification objective and a sCMOS camera (Hamamatsu, ORCA-Flash 4). Time-lapse imaging was performed under ambient conditions (regulated at 26 ºC ± 1 ºC) with a 5-minute period using automated routines (Molecular Devices, Methamorph). All reported experiments were performed once.

For imaging with single-cell resolution, 2 µL of 10x diluted stationary-phase cells (approximately 6×10^5 CFU) grown in liquid MPIPES-methanol without formaldehyde were introduced on 1 mm thick agar pads pre-deposited on a glass coverslip. The culture solution was allowed to dry for approximately 10 minutes within a biosafety cabinet and was subsequently covered with a second coverslip. Individual bacteria were therefore immobilized at the agar-coverslip interface, allowing them to grow in two dimensions. Specific locations (200x150 µm^2) were monitored in parallel, selected by evaluating the distance between individual cells at the beginning of each experiment so that expanding micro-colonies would not overlap spatially at later time points. On average, 40 microcolonies per location were imaged, with approximately 8-10 individual cells per microcolony in the final timepoint. ImageJ [70] and manual curation were employed for cell segmentation and tracking (Fig. S10).

To prepare the pads, MPIPES-agar medium was prepared as described above ("Bacterial strains and culture conditions"), and melted at 70 ºC for approximately 2 hours using a convection oven. Subsequently, methanol was introduced to the agar medium to a final concentration of 125 mM. Specific to the formaldehyde tolerance experiments, formaldehyde was also added to a final concentration of 2.5 mM. In all experiments, a polydimethylsiloxane (PDMS, Sylgard 184, Dow Corning) frame was employed to cover all free edges of the agar pads. This step was employed to minimize formaldehyde evaporation during time-lapse imaging. To fabricate these membranes,

PDMS monomer was mixed with its catalyst at a 10:1 ratio, degassed, and cured at 70

ºC for approximately 2 hours. Subsequently, the cured PDMS was cut to an area of

25×50 mm^2, including a 21×47 mm^2 internal aperture, where the melted agar was

subsequently introduced.

Analysis of cell doubling times and microcolony lag times was conducted in R. The

wilcox.test function was used to compare the formaldehyde-treated population with the

control population by Mann-Whitney Wilcoxon test [71]. The relative effects of

formaldehyde treatment and colony lineage were assessed by permutational

multivariate ANOVA using distance matrices (PERMANOVA) [72], with the adonis

function in the vegan package [73].

For all microscopy experiments, an aliquot of the same culture used for the microscopy

experiment was simultaneously grown in batch liquid culture with same concentration of

formaldehyde, and CFUs in both the total and formaldehyde-tolerant populations were

tracked over time, as a control. In all cases, the growth of the liquid culture proceeded

as expected and the frequency of formaldehyde-tolerant cells matched that observed in

the microscopy experiment.

1001

Assays on selected formaldehyde-tolerant subpopulations: fitness trade-offs

formaldehyde-free regrowth experiments

For characterization of the formaldehyde-tolerant subpopulation, a 4 mM formaldehyde

exposure experiment was conducted as described above to select for cells with a

minimum tolerance level of 4 mM. Once cultures had reached stationary phase

(approximately 80 hours), they were used within 4 hours for further experiments.

1008

Assessment of tolerance to antimicrobial drugs and to hydrogen peroxide were carried

out using a disk susceptibility test: liquid exponential-phase cultures grown on methanol

in standard conditions were diluted to an OD_{600} of 0.3, then spread using a sterile swab

onto the surface of MPIPES-methanol-agar plates and allowed to dry. A BBL Sensi-Disc

(Becton-Dickinson), impregnated with one antibiotic compound at a set concentration,

was then placed on top of the plate. Plates were incubated at 30 °C for 48 hours to

allow a lawn to grow; the diameter of the clearing zone around the antibiotic disk was

then measured and compared between the naive and selected high-tolerance

populations.

1018

To compare growth rates on different carbon substrates, three selected subpopulations,

and three naive populations not previously exposed to formaldehyde, were diluted 1:64

into fresh MPIPES medium containing either methanol or succinate, and each of those

was transferred into the wells of a 48-well tissue culture plate (Corning Costar) with 640

µL culture in each of three replicate wells. Plates were incubated on a Liconic LPX44

incubator shaker at 650 rpm, and OD_{600} was read using a Wallac 1420 Victor2 plate

reader. Growth rates were calculated in R using timepoints during the period of

exponential growth (between 5-18 hours for methanol growth and 5-22 hours for

succinate), by fitting a linear relationship between time and the binary logarithm of the

OD using the lm function in R. Rates were calculated for each well individually; for each

biological replicate, the mean and standard deviation of the three replicate wells was

calculated. Statistical tests of differences among growth rates consisted of analysis of

variance using the anova function from the stats package, and planned contrasts

calculated using the lsmeans and contrast functions from the lsmeans package.

1033

To assess tolerance distribution dynamics during formaldehyde-free regrowth, three

selected subpopulations were each diluted 1:64 into two batches of fresh MPIPES liquid

medium, one batch containing methanol and the other succinate. Dilutions were grown

in standard flask conditions as described above, but with loose caps instead of Suba

Seals. Cultures were sampled every 4 hours for serial dilution and plating as described

above, with each sample plated onto each of 7 plates containing different levels of

formaldehyde (0, 2, 4, 6, 8, 10, or 12 mM). Plates were incubated and colonies were

counted as described above.

1042

Mathematical model

Growth, death, and phenotype transitions of *M. extorquens* populations were modeled

using a partial differential equation (PDE; Equation 1). The full code for parameter fitting

and model selection is included as an R notebook in the Supporting Information, File

S1. Over the time periods for which we ran the model simulation, there was no effective

change in the concentration of either of the growth substrates (methanol, succinate;

each of which have extremely low half-saturation constants [74,75]) or formaldehyde

(Fig. 2); these compounds were therefore not explicitly included as time-dependent

variables in the model. Furthermore, given that tolerance either has no effect upon

growth on methanol, or a mild effect upon succinate growth, we use a single value of r_c

for all values of *x*.

1054

Modeling was conducted in R. The PDE was solved numerically by vectorized ODEs,

where each ODE corresponded to a discrete bin within tolerance space of 0.01 mM

formaldehyde. A finite difference grid was created using the setup.grid.1D function, and

advection and diffusion were calculated using the advection.1D or tran.1D function as

appropriate for the different models, in the ReacTran package v. 1.4.3.1 [76]. The

vectorized ODEs were solved using the ode.1D function from the package deSolve

v1.21 [77], with the lsoda method [78]. Zero-flux boundary conditions for Equation 1 are

given by: $\partial_x N(0,t){=}0$ and $\partial_x N(L,t){=}0$. The lower boundary was set as *x*=0 because

tolerance cannot fall below 0. The upper boundary (*x*=*L*) for each growth condition was

set higher than the highest experimentally observed value, sufficiently large so as not to

constrain the upward transition of cells in phenotypic space.

1066

Initial conditions

The experimentally measured distributions of formaldehyde tolerances in the original

populations were used as initial conditions in all model runs. Because we found that

populations exhibit a slight shift toward higher average tolerance within 2 hours of

transitioning from the stationary (non-growing) to the exponential (growing) phase (Fig.

6), and because our model specifically focuses on growth phenomena and does not

capture behavior in stationary phase, we chose the 2-hour tolerance distribution as our

initial condition (Fig. S7) in order to avoid artefacts that would result from using the 0-

hour stationary-phase distribution. Three biological replicates were generated and the

average of the three used to generate the distribution.

1077

Because the model is continuous, whereas the experimental data were obtained at a

resolution of 1.0 or 2.0 mM formaldehyde in tolerance space, the initial conditions for

the model were generated by fitting a monotone cubic spline using the function

splinefun in R with method "hyman," to interpolate cell abundances for values of x at

intervals of 0.01 mM (Fig. S7). Subsequently, for comparison of model results with

experimental results, the model results were summed in 1 mM or 2 mM intervals to re-

obtain coarse bins matching the resolution of the experimental data (as in Fig. 9).

1085

In the model, x denotes tolerance as the *maximum* tolerance level of a cell or

subpopulation. However, our experimental CFU counts are cumulative in that regard

(*i.e.*, the cells that form colonies on culture plates containing 3 mM formaldehyde

include those with a maximum tolerance of not just 3 mM but also 4 mM, 5 mM, and 6

mM). For the purposes of carrying out the model, we therefore transformed the

empirical tolerance distributions in the experimental datasets to calculate the actual

(non-cumulative) number of CFU at each phenotype level using the following formula:

1093

1094 $$N(x) = \widehat{N}(x) - \widehat{N}(x + h) \qquad\qquad (3)$$

1095

where $N(x)$ is the number of cells that uniquely have a given tolerance level x, $\widehat{N}(x)$ is

the number of cells measured experimentally as CFU on culture plates with

formaldehyde concentration x, and h is the step size between categories in data (1 mM

for Fig. 7a; 2 mM for Figs. 7b-c). This transformation to the non-cumulative distribution

was used after spline fitting to calculate the initial distributions for the model, and model

results are shown in this format in Fig. 9a-b and Fig. S8b. However, for all other

purposes, including parameter fitting as well as display in Figs. 9c-d and S11, model

results were transformed back to the cumulative distribution for comparison with the

experimental results. Both forms of the data are displayed in the R notebook (File S1).

1105

A further adjustment to data from all timepoints was made to account for the

experimental limit of detection (34 CFU/mL) in measuring formaldehyde tolerance

distributions: growth of originally undetected low-frequency cells in the high-tolerance

phenotypes could potentially be mistaken as transition into those phenotypes from

lower-tolerance phenotypes. To assess whether such undetected cells would make a

difference to the model results, we generated a set of "extended" CFU counts in which

we made the extreme correction of adding 1 CFU to the experimental observations (the

equivalent of 90.9 CFU/mL for that replicate, or 30.3 CFU/mL if averaged across the

three replicates) at high formaldehyde concentrations where the observation had been 0

CFU. This correction was made to the data from all timepoints, according to the

following rules:

1. If 0 colonies were observed in all technical replicates, add 1 colony to one of the

replicates.

2. Only do (1) if either:

a) there is an observation of ≥1 colony at a higher concentration than the one being

considered, OR

b) it is the first concentration beyond the last observation of ≥1 colony.

1123

We carried out parameter fitting and model selection for the 4 mM formaldehyde

selection experiment using both the original dataset and the extended one (as

described in the previous paragraph). Although the use of the extended dataset resulted

in minor differences in the estimated parameters (Tables 1 and S1, Fig. S11), the same

form of the model was favored (the 3-parameter model with α, d, and b), and the

pseudo-R^2 (0.973) was slightly better than for the version using the original dataset

(0.970). We interpreted this to mean that the inclusion or omission of rare undetected

cells at high tolerance levels makes little difference to any biologically relevant

conclusions, but that our correction may allow for a better model fit. We therefore

proceeded with only the extended dataset for the models describing formaldehyde-free

regrowth, and all parameter and statistical values given in the text are those for the

extended data.

1136

Parameter estimation

1137

Growth rates (r_c for substrate c, either methanol or succinate, h^{-1}) and their standard

errors were estimated from fitting experimental data of growth of a wild-type, naive

population on either succinate or methanol as a primary carbon source, in the absence

of formaldehyde (Fig. S7). Linear regression was used to fit the relationship between

cell count and time for the exponential portion of three replicate growth curves, using

the lm function in R. The growth rate on methanol was 0.195±0.001 h^{-1} and the growth

rate on succinate was 0.267±0.005 h^{-1}, where ± denotes 95% confidence interval.

1145

The death rate (α), dependence of death on tolerance (b), diffusion (D), and advection

(v) parameters were estimated using maximum likelihood. Due to the exponential nature

of bacterial growth, and because our data sets contain a number of zeroes, we

transformed our data using the hyperbolic arcsine function ($asinh(x) = ln(x + \sqrt{1 + x^2})$)

[60], which is approximately logarithmic but defined at x=0. The functions lm and logLik

in R were used to calculate log-likelihood under a linear model; optimization was carried

out in R using optim with the Nelder-Mead method. Standard errors of the parameters

were calculated from the Hessian matrix. Fitted values for D and v are given in Table 1

and Table S1. In addition, we generated a phenotype-independent estimate of α using

death rate data from a separate set of time-kill experiments (such as those in Fig. 1) at

formaldehyde concentrations between 3 and 20 mM. Time-kill curves were conducted

as described above, the death rate at each concentration (h^{-1}) was calculated using

linear regression, and the relationship between concentration and death rate was then

also calculated using linear regression. By this method, we estimated α as 0.189±0.010

h^{-1}*mM^{-1}, which fell within the range of values predicted by fitting α for the models we

tested, and within the 95% confidence interval of the estimate of α in the best model.

This phenotype-independent estimate was not used in modeling, but helped to verify

that the values we obtained by parameter fitting were reasonable.

1164

Model evaluation

For each experimental condition, we used a likelihood ratio test on the nested models

using a forward, stepwise procedure to choose the model that best fit the experimental

data. For the formaldehyde selection scenario, the "absolute death" model (with α as

the only parameter) was used as the null model, and each 2-parameter model (with

alpha and either b, D, or v) was compared against it. Of the three 2-parameter models,

we chose the one with the highest likelihood as long as it was significantly better than

the null model. If a 2-parameter model was chosen, it became the null model and the

procedure was repeated to determine whether a 3-parameter model was supported,

and subsequently the 4-parameter model. For the regrowth scenarios, where no death

due to formaldehyde is possible, we omitted α and b. We first compared two 1-

parameter models (v only or D only) against a null model containing neither, and then

evaluated whether adding the second parameter was significantly better. LR was

calculated as

$LR = -2(LL_0 - LL_1)$

where LL_0 is the log-likelihood of a reduced model and LL_1 is the log-likelihood of the

model being tested. Statistical significance was assessed using the chi-squared test,

with degrees of freedom given by the difference in the number of parameters between

the model being tested and the null model.

References

1.   Harms A, Fino C, Sørensen MA, Semsey S, Gerdes K. Prophages and growth
       dynamics confound experimental results with antibiotic-tolerant persister cells.
       mBio. 2017;8: e01964-17. doi:10.1128/mBio.01964-17

2.   Shaffer SM, Dunagin MC, Torborg SR, Torre EA, Emert B, Krepler C, et al. Rare
       cell variability and drug-induced reprogramming as a mode of cancer drug
       resistance. Nature. 2017;546: 431–435. doi:10.1038/nature22794

3.   Lorenzi T, Chisholm RH, Desvillettes L, Hughes BD. Dissecting the dynamics of
       epigenetic changes in phenotype-structured populations exposed to fluctuating
       environments. J Theor Biol. 2015;386: 166–176. doi:10.1016/j.jtbi.2015.08.031

4.   Vasdekis AE, Silverman AM, Stephanopoulos G. Origins of cell-to-cell
       bioprocessing diversity and implications of the extracellular environment revealed
       at the single-cell level. Sci Rep. 2015;5: 17689. doi:10.1038/srep17689

5.   Elowitz MB, Levine AJ, Siggia ED, Swain PS. Stochastic gene expression in a
       single cell. Science. 2002;297: 1183–1186. doi:10.1126/science.1070919

6.   Kiviet DJ, Nghe P, Walker N, Boulineau S, Sunderlikova V, Tans SJ. Stochasticity
       of metabolism and growth at the single-cell level. Nature. 2014;514: 376–379.
       doi:10.1038/nature13582

7.   Newman JRS, Ghaemmaghami S, Ihmels J, Breslow DK, Noble M, DeRisi JL, et
       al. Single-cell proteomic analysis of *S. cerevisiae* reveals the architecture of
       biological noise. Nature. 2006;441: 840. doi:10.1038/nature04785

8.   Bergmiller T, Ackermann M. Pole age affects cell size and the timing of cell division
       in *Methylobacterium extorquens* AM1. J Bacteriol. 2011;193: 5216–5221.
       doi:10.1128/JB.00329-11

9.   Levy SF, Ziv N, Siegal ML. Bet hedging in yeast by heterogeneous, age-correlated
expression of a stress protectant. PLOS Biol. 2012;10: e1001325.
doi:10.1371/journal.pbio.1001325

10. Huh D, Paulsson J. Random partitioning of molecules at cell division. Proc Natl
Acad Sci U S A. 2011;108: 15004–15009. doi:10.1073/pnas.1013171108

11. Ackermann M. A functional perspective on phenotypic heterogeneity in
microorganisms. Nat Rev Microbiol. 2015;13: 497–508. doi:10.1038/nrmicro3491

12. Balaban NQ, Merrin J, Chait R, Kowalik L, Leibler S. Bacterial persistence as a
phenotypic switch. Science. 2004;305: 1622–1625. doi:10.1126/science.1099390

13. Veening J-W, Stewart EJ, Berngruber TW, Taddei F, Kuipers OP, Hamoen LW.
Bet-hedging and epigenetic inheritance in bacterial cell development. Proc Natl
Acad Sci. 2008;105: 4393–4398. doi:10.1073/pnas.0700463105

14. Choi PJ, Cai L, Frieda K, Xie XS. A stochastic single-molecule event triggers
phenotype switching of a bacterial cell. Science. 2008;322: 442.
doi:10.1126/science.1161427

15. Venturelli OS, Zuleta I, Murray RM, El-Samad H. Population diversification in a
yeast metabolic program promotes anticipation of environmental shifts. PLOS Biol.
2015;13: e1002042. doi:10.1371/journal.pbio.1002042

16. Wang J, Atolia E, Hua B, Savir Y, Escalante-Chong R, Springer M. Natural
variation in preparation for nutrient depletion reveals a cost–benefit tradeoff. PLOS
Biol. 2015;13: e1002041. doi:10.1371/journal.pbio.1002041

17. Eldar A, Elowitz MB. Functional roles for noise in genetic circuits. Nature.
2010;467: 167–173. doi:10.1038/nature09326

18. Fridman O, Goldberg A, Ronin I, Shoresh N, Balaban NQ. Optimization of lag time
underlies antibiotic tolerance in evolved bacterial populations. Nature. 2014;513:
418–421. doi:10.1038/nature13469

19. Rainey PB, Beaumont HJ, Ferguson GC, Gallie J, Kost C, Libby E, et al. The
evolutionary emergence of stochastic phenotype switching in bacteria. Microb Cell
Factories. 2011;10: S14. doi:10.1186/1475-2859-10-S1-S14

20. Chisholm RH, Lorenzi T, Desvillettes L, Hughes BD. Evolutionary dynamics of
phenotype-structured populations: from individual-level mechanisms to population-
level consequences. Z Für Angew Math Phys. 2016;67: 100. doi:10.1007/s00033-
016-0690-7

21. Klironomos FD, Berg J, Collins S. How epigenetic mutations can affect genetic
evolution: Model and mechanism. BioEssays. 2013;35: 571–578.
doi:10.1002/bies.201200169

22. Levin-Reisman I, Ronin I, Gefen O, Braniss I, Shoresh N, Balaban NQ. Antibiotic
tolerance facilitates the evolution of resistance. Science. 2017;355: 826–830.
doi:10.1126/science.aaj2191

23. Deris JB, Kim M, Zhang Z, Okano H, Hermsen R, Groisman A, et al. The innate
growth bistability and fitness landscapes of antibiotic-resistant bacteria. Science.
2013;342: 1237435. doi:10.1126/science.1237435

24. Lorenzi T, Chisholm RH, Clairambault J. Tracking the evolution of cancer cell
populations through the mathematical lens of phenotype-structured equations. Biol
Direct. 2016;11: 43. doi:10.1186/s13062-016-0143-4

25. Fall R, Benson AA. Leaf methanol — the simplest natural product from plants.
Trends Plant Sci. 1996;1: 296–301. doi:10.1016/S1360-1385(96)88175-0

26. Skovran E, Crowther GJ, Guo X, Yang S, Lidstrom ME. A systems biology
approach uncovers cellular strategies used by *Methylobacterium extorquens* AM1
during the switch from multi- to single-carbon growth. PLoS ONE. 2010;5: e14091.
doi:10.1371/journal.pone.0014091

27. Nayak DD, Marx CJ. Genetic and phenotypic comparison of facultative
methylotrophy between *Methylobacterium extorquens* strains PA1 and AM1. PLoS
ONE. 2014;9: e107887. doi:10.1371/journal.pone.0107887

28. Chen NH, Djoko KY, Veyrier FJ, McEwan AG. Formaldehyde stress responses in
bacterial pathogens. Front Microbiol. 2016;7. doi:10.3389/fmicb.2016.00257

29. Marx CJ, Chistoserdova L, Lidstrom ME. Formaldehyde-detoxifying role of the
tetrahydromethanopterin-linked pathway in *Methylobacterium extorquens* AM1. J
Bacteriol. 2003;185: 7160–7168. doi:10.1128/JB.185.23.7160-7168.2003

30. Atwood KC, Norman A. On the interpretation of multi-hit survival curves. Proc Natl
Acad Sci U S A. 1949;35: 696–709.

31. Peleg M. Microbial survival curves — the reality of flat "shoulders" and absolute
thermal death times. Food Res Int. 2000;33: 531–538. doi:10.1016/S0963-
9969(00)00088-0

32. Regoes RR, Wiuff C, Zappala RM, Garner KN, Baquero F, Levin BR.
Pharmacodynamic functions: a multiparameter approach to the design of antibiotic
treatment regimens. Antimicrob Agents Chemother. 2004;48: 3670–3676.
doi:10.1128/AAC.48.10.3670-3676.2004

33. Levin BR, Udekwu KI. Population dynamics of antibiotic treatment: a mathematical
model and hypotheses for time-kill and continuous-culture experiments. Antimicrob
Agents Chemother. 2010;54: 3414–3426. doi:10.1128/AAC.00381-10

34. Yurtsev EA, Chao HX, Datta MS, Artemova T, Gore J. Bacterial cheating drives the
population dynamics of cooperative antibiotic resistance plasmids. Mol Syst Biol.
2013;9: 683. doi:10.1038/msb.2013.39

35. Lennon JT, Jones SE. Microbial seed banks: the ecological and evolutionary
implications of dormancy. Nat Rev Microbiol. 2011;9: 119–130.
doi:10.1038/nrmicro2504

36. Kotte O, Volkmer B, Radzikowski JL, Heinemann M. Phenotypic bistability in
Escherichia coli's central carbon metabolism. Mol Syst Biol. 2014;10: 736–736.
doi:10.15252/msb.20135022

37. Boxtel C van, Heerden JH van, Nordholt N, Schmidt P, Bruggeman FJ. Taking
chances and making mistakes: non-genetic phenotypic heterogeneity and its
consequences for surviving in dynamic environments. J R Soc Interface. 2017;14:
20170141. doi:10.1098/rsif.2017.0141

38. Levin-Reisman I, Fridman O, Balaban NQ. ScanLag: High-throughput
quantification of colony growth and lag time. J Vis Exp JoVE. 2014;
doi:10.3791/51456

39. Guillier L, Pardon P, Augustin J-C. Automated image analysis of bacterial colony
growth as a tool to study individual lag time distributions of immobilized cells. J
Microbiol Methods. 2006;65: 324–334. doi:10.1016/j.mimet.2005.08.007

40. Ernebjerg M, Kishony R. Distinct growth strategies of soil bacteria as revealed by
large-scale colony tracking. Appl Environ Microbiol. 2012;78: 1345–1352.
doi:10.1128/AEM.06585-11

41. Mitsui R, Kusano Y, Yurimoto H, Sakai Y, Kato N, Tanaka M. Formaldehyde
fixation contributes to detoxification for growth of a nonmethylotroph, *Burkholderia*
cepacia TM1, on vanillic acid. Appl Environ Microbiol. 2003;69: 6128–6132.
doi:10.1128/AEM.69.10.6128-6132.2003

42. Sudtachat N, Ito N, Itakura M, Masuda S, Eda S, Mitsui H, et al. Aerobic vanillate
degradation and C1 compound metabolism in *Bradyrhizobium japonicum*. Appl
Environ Microbiol. 2009;75: 5012–5017. doi:10.1128/AEM.00755-09

43. Peredo EL, Simmons SL. Leaf-FISH: microscale imaging of bacterial taxa on
phyllosphere. Front Microbiol. 2018;8. doi:10.3389/fmicb.2017.02669

44. Voordeckers K, Kominek J, Das A, Espinosa-Cantú A, Maeyer DD, Arslan A, et al.

Adaptation to high ethanol reveals complex evolutionary pathways. PLOS Genet.

2015;11: e1005635. doi:10.1371/journal.pgen.1005635

45. Costa E, Pérez J, Kreft J-U. Why is metabolic labour divided in nitrification? Trends

Microbiol. 2006;14: 213–219. doi:10.1016/j.tim.2006.03.006

46. Strovas TJ, Sauter LM, Guo X, Lidstrom ME. Cell-to-Cell heterogeneity in growth

rate and gene expression in *Methylobacterium extorquens* AM1. J Bacteriol.

2007;189: 7127–7133. doi:10.1128/JB.00746-07

47. Strovas TJ, Lidstrom ME. Population *in Methylobacterium extorquens* AM1.

Microbiology. 2009;155: 2040–2048. doi:10.1099/mic.0.025890-0

48. Gallie J, Libby E, Bertels F, Remigi P, Jendresen CB, Ferguson GC, et al.

Bistability in a metabolic network underpins the de novo evolution of colony

switching in *Pseudomonas fluorescens*. PLOS Biol. 2015;13: e1002109.

doi:10.1371/journal.pbio.1002109

49. Ozbudak EM, Thattai M, Lim HN, Shraiman BI, van Oudenaarden A. Multistability

in the lactose utilization network of *Escherichia coli*. Nature. 2004;427: 737–740.

doi:10.1038/nature02298

50. Roff DA. The evolution of threshold traits in animals. Q Rev Biol. 1996;71: 3–35.

doi:10.1086/419266

51. Govers SK, Mortier J, Adam A, Aertsen A. Protein aggregates encode epigenetic

memory of stressful encounters in individual *Escherichia coli* cells. PLOS Biol.

2018;16: e2003853. doi:10.1371/journal.pbio.2003853

52. Bandyopadhyay A, Wang H, Ray JCJ. Lineage space and the propensity of

bacterial cells to undergo growth transitions. PLOS Comput Biol. 2018;14:

e1006380. doi:10.1371/journal.pcbi.1006380

53. Kussell E, Leibler S. Phenotypic diversity, population growth, and information in
fluctuating environments. Science. 2005;309: 2075–2078.
doi:10.1126/science.1114383

54. Arnoldini M, Mostowy R, Bonhoeffer S, Ackermann M. Evolution of stress response
in the face of unreliable environmental signals. PLOS Comput Biol. 2012;8:
e1002627. doi:10.1371/journal.pcbi.1002627

55. Beaumont HJE, Gallie J, Kost C, Ferguson GC, Rainey PB. Experimental evolution
of bet hedging. Nature. 2009;462: 90. doi:10.1038/nature08504

56. Draghi J. Links between evolutionary processes and phenotypic robustness in
microbes. Semin Cell Dev Biol. 2018; doi:10.1016/j.semcdb.2018.05.017

57. Abanda-Nkpwatt D, Müsch M, Tschiersch J, Boettner M, Schwab W. Molecular
interaction between *Methylobacterium extorquens* and seedlings: growth
promotion, methanol consumption, and localization of the methanol emission site. J
Exp Bot. 2006;57: 4025–4032. doi:10.1093/jxb/erl173

58. Ryffel F, Helfrich EJN, Kiefer P, Peyriga L, Portais J-C, Piel J, et al. Metabolic
footprint of epiphytic bacteria on *Arabidopsis thaliana* leaves. ISME J. 2016;10:
632–643. doi:10.1038/ismej.2015.141

59. Nemecek-Marshall M, MacDonald RC, Franzen JJ, Wojciechowski CL, Fall R.
Methanol emission from leaves (Enzymatic detection of gas-phase methanol and
relation of methanol fluxes to stomatal conductance and leaf development). Plant
Physiol. 1995;108: 1359–1368. doi:10.1104/pp.108.4.1359

60. Ackermann M. Microbial individuality in the natural environment. ISME J. 2013;7:
465–467. doi:10.1038/ismej.2012.131

61. Delaney NF, Kaczmarek ME, Ward LM, Swanson PK, Lee M-C, Marx CJ.
Development of an optimized medium, strain and high-throughput culturing
methods for *Methylobacterium extorquens*. PLOS ONE. 2013;8: e62957.
doi:10.1371/journal.pone.0062957

62. Nash T. The colorimetric estimation of formaldehyde by means of the Hantzsch
reaction. Biochem J. 1953;55: 416–421.

63. Deatherage DE, Barrick JE. Identification of mutations in laboratory-evolved
microbes from next-generation sequencing data using *breseq*. Engineering and
Analyzing Multicellular Systems. Humana Press, New York, NY; 2014. pp. 165–
188. doi:10.1007/978-1-4939-0554-6_12

64. Marx CJ, Bringel F, Chistoserdova L, Moulin L, Farhan Ul Haque M, Fleischman
DE, et al. Complete genome sequences of six strains of the genus
Methylobacterium. J Bacteriol. 2012;194: 4746–4748. doi:10.1128/JB.01009-12

65. Michener JK, Vuilleumier S, Bringel F, Marx CJ. Transfer of a catabolic pathway for
chloromethane in *Methylobacterium* strains highlights different limitations for
growth with chloromethane or with dichloromethane. Front Microbiol. 2016;7.
doi:10.3389/fmicb.2016.01116

66. R Core Team. R: A language and environment for statistical computing. [Internet].
Vienna, Austria: R Foundation for Statistical Computing; 2018. Available:
http://www.R-project.org

67. Hahne F, LeMeur N, Brinkman RR, Ellis B, Haaland P, Sarkar D, et al. flowCore: a
Bioconductor package for high throughput flow cytometry. BMC Bioinformatics.
2009;10: 106. doi:10.1186/1471-2105-10-106

68. Wickham, H. ggplot2: Elegant graphics for data analysis [Internet]. New York:
Springer-Verlag; 2016. Available: http://ggplot2.org

69. van der Walt S, Schönberger JL, Nunez-Iglesias J, Boulogne F, Warner JD, Yager
N, et al. scikit-image: image processing in Python. PeerJ. 2014;2: e453.
doi:10.7717/peerj.453

70. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image
analysis. Nat Methods. 2012;9: 671–675.

71.   Wilcoxon F. Individual comparisons by ranking methods. Biom Bull. 1945;1: 80–83.
doi:10.2307/3001968

72.   Anderson MJ. A new method for non-parametric multivariate analysis of variance.
Austral Ecol. 2001;26: 32–46. doi:10.1111/j.1442-9993.2001.01070.pp.x

73.   Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, et al. The
vegan package. Community Ecol Package. 2007;10: 631–637.

74.   McAllister CF, Lepo JE. Succinate transport by free-living forms of *Rhizobium*
japonicum. J Bacteriol. 1983;153: 1155–1162.

75.   Anthony C, Zatman LJ. The microbial oxidation of methanol. 2. The methanol-
oxidizing enzyme of *Pseudomonas* sp. M27. Biochem J. 1964;92: 614–621.

76.   Soetaert K, Meysman F. Reactive transport in aquatic ecosystems: Rapid model
prototyping in the open source software R. Environ Model Softw. 2012;32: 49–60.
doi:10.1016/j.envsoft.2011.08.011

77.   Stoetaert K, Petzoldt T, Setzer W. Solving differential equations in R: package
deSolve. J Stat Softw. 2010;33: 1–25. doi:10.18637/jss.v033.i09

78.   Hindmarsh AC. ODEPACK, A systematized collection of ODE solvers. In:
Stepleman RS, editor. Scientific Computing. Amsterdam: North-Holland; 1983. pp.
55–64.

1407

1408

1409 <u>**Figures with legends**</u>

1410
Figure 1. Formaldehyde kills *M. extorquens* at an exponential, concentration-
dependent rate.
Formaldehyde was added at the indicated concentrations to liquid cultures of *M.*
extorquens cells growing in minimal medium with methanol, and abundance of viable
cells was measured as colony-forming units (CFU) over time. Note that negligible
growth is expected to have occurred during the course of this experiment, as the 180-
minute duration was less than one generation (~3.5 hrs) for *M. extorquens* in these
conditions.

1419

Figure 2. Re-growth of _M. extorquens_ after population decline in the presence of formaldehyde is due to a pre-existing sub-population of formaldehyde-tolerant cells.

Stationary-phase cells were inoculated into fresh medium containing methanol and 4 mM formaldehyde. The abundance of viable cells in the two different populations was assessed over time by removing and washing cells, then plating onto both permissive medium (without formaldehyde: "all cells") and selective medium (with 4 mM formaldehyde: "tolerant to 4 mM"). CFU = colony-forming units. Each line represents one biological replicate; error bars show the standard deviation of three replicate platings. Formaldehyde in the liquid medium during the incubation period was measured by a colorimetric assay on subsamples after removing cells by centrifugation.

1431
Figure 3. **Cell proliferation assay shows dynamics consistent with the**
coexistence of both growing and non-growing subpopulations, with no turnover
between the two.
Cells were stained with PKH67 fluorescent membrane dye, then allowed to grow in
minimal medium with methanol and either 0, 4, or 20 mM formaldehyde. Histograms
show per-cell fluorescence of the cells (events measured by flow cytometry) present in
30 µL of culture at each timepoint; colors denote the time of sampling in hours (note that
different color scales are used in different panels). Top left: without formaldehyde, all
cells underwent doubling, diluting their membrane fluorescence so that the median
fluorescence decreased as population increased. Bottom left: at high concentrations of
formaldehyde, no cells grew, leaving per-cell fluorescence unchanged. Right: in the
presence of 4 mM formaldehyde, most cells did not grow, but a few did; consequently, a
small growing population with lower per-cell fluorescence became detectable at 37
hours and continued to increase in abundance thereafter. Results of experiments
conducted at other formaldehyde concentrations are shown in Fig. S5.

1447
Figure 4. **Cell damage by formaldehyde results in delayed colony arisal for the**
majority of cells, but not for the tolerant subpopulation.
Cells from a formaldehyde exposure experiment (liquid MPIPES medium with 4 mM
formaldehyde) were sampled at 2- to 4-hour intervals, washed, and plated onto both
permissive medium (no formaldehyde, allowing the growth of all cells) and selective
medium (4 mM formaldehyde, allowing the growth of only the tolerant subpopulation).
A) Images of colonies on plates. Colony size heterogeneity was evident only on
permissive medium with cultures exposed to formaldehyde for 16 hours, consistent with
a population containing both sensitive cells that formed colonies late due to
formaldehyde-induced damage (small colonies) and tolerant cells that formed colonies
early (large colonies). All images are shown at the same magnification level; *dil*=dilution
factor prior to plating. B) Relationship between formaldehyde exposure and colony
growth characteristics. Shading indicates abundance of colony-forming units in each
population (see Fig. 2); samples were diluted prior to plating for an average of 500
colonies per plate. Left panel: gray line shows linear regression of arisal time on
exposure time for the first 12 hours. Every hour of exposure to formaldehyde led to a
~4.8-hour delay in colony arisal time among sensitive cells. At 16 hours, the population
consisted of both damaged and tolerant cells; after 20 hours, all cells were tolerant due
to the death of the damaged cells. Right panel: among tolerant cells, formaldehyde
exposure had no effect on arisal time. Insets: exposure time affected only the variability
among colony growth rates, but not their median.

Figure 5. Time-lapse microscopy reveals binary (i.e., growth or non-growth) phenotypes in response to formaldehyde.
A) Example images: cells were embedded in agar medium with methanol and either 0 mM (top) or 2.5 mM (bottom) formaldehyde and monitored for 9 hours (~3 generations). At 0 mM, 256 cells were observed and all underwent at least one doubling; at 2.5 mM, 546 cells were observed and 11 (1.97%) underwent at least one doubling, in accordance with our predictions for this formaldehyde concentration (see Fig. 6). B) Histograms of cell division time (across all generations) and lag time (time between deposition and first cell division, for each microcolony) for cells that grew. No difference was observed in cell division time between the two treatments (p=0.262, Mann-Whitney Wilcoxon test). However, cells in formaldehyde took approximately 1.25 hours longer to reach the first cell division (p<0.001, Mann-Whitney). C) Scatterplot of individual cell doubling times; each position along the x-axis represents a single microcolony, ordered by mean doubling time (shown in black symbols). Individual doubling time of each cell was strongly predicted by the colony it came from (p=0.001) but not by formaldehyde treatment (p=0.323, PERMANOVA).

1487
Figure 6. **Subpopulations of formaldehyde-tolerant cells are distributed within a wild-type population with continuous, exponentially-decreasing frequency.**
M. extorquens cells not previously exposed to formaldehyde were plated onto methanol agar medium containing a range of formaldehyde concentrations at 1-mM intervals. The frequency of tolerant cells is expressed as the ratio of the colony-forming units (CFU) on formaldehyde medium at the specified concentration to the CFU on formaldehyde-free (0 mM) medium. Error bars denote the standard deviation of replicate experiments from 5 different dates (shown individually in Fig. S4). Detection limit is indicated by the dashed horizontal line.

1497
Figure 7. **The distribution of formaldehyde tolerance within an *M. extorquens***
population changes over time depending on growth conditions.
Plots show total abundance (not frequency) of cells tolerant to each level of
formaldehyde, as assessed by plating onto selective medium; each colored line
represents one timepoint and error bars represent the standard deviation of three
plating replicates. For clarity, only one biological replicate is shown; results from other
replicates are shown in Fig. S6. Populations were tested for tolerance at up to 10 mM
(panel A) or 12 mM (panel B), but 0 CFU were detected above the tolerance levels
shown here. A) Exposure of a naive population to 4 mM formaldehyde results in rapid
decline of subpopulations with tolerance levels <4 mM and selective growth of
subpopulations with tolerance levels ≥4 mM. B) When the population from A), enriched
in tolerant cells, is transferred to medium without formaldehyde, tolerance distribution
dynamics depend on the growth substrate provided. If growth occurs on methanol, all
subpopulations grow equally well: the enrichment of formaldehyde-tolerant populations
is retained for the full 24 hours (~7 generations) of observation. If growth occurs on
succinate, subpopulations with high tolerance decline in abundance and those with low
tolerance increase: the population reverts to its original naive distribution.

1515
Figure 8. Schematic of processes described by mathematical model of tolerance
distribution dynamics.
Cells exist in 1-dimensional phenotype space along a continuum from sensitive to
tolerant, with x denoting the maximum concentration of formaldehyde (F) at which a cell
can grow. Under normal growth (at rate r), progeny cells carry the same tolerance
phenotype as their parents. Exposure to formaldehyde results in the death of low-
tolerance ($x<F$) phenotypes at a rate described by $H(x, F)$. In the process of diffusion,
cells and their progeny shift to adjacent tolerant states according to the diffusion
constant D, resulting in the broadening of the population's tolerance distribution. In
advection, cells and their progeny move in a single direction in tolerance space at rate
v, resulting in an overall shift in the population's distribution toward either lower or higher
average tolerance.

1528
Figure 9. Mathematical modeling reproduces growth, death, and phenotype
transition dynamics of *M. extorquens* population under multiple conditions.
A) and B) Heat maps showing model simulations of population dynamics. Model
parameters as given in Table 1. Note that model results are continuous in phenotype
space, and non-cumulative (the abundance at concentration *x* shows only the number
of cells for which that is the maximum concentration tolerable, not the number of all cells
that can grow at that concentration; see Methods for details). C) and D) Comparison of
model results (lines) and experimental data (points). Experimental data are averages of
3 biological replicates; model results have been binned at 1- or 2-mM intervals, and
summed to form cumulative distributions, to facilitate comparison. A and C) 4 mM
formaldehyde exposure experiment, resulting in selection of cells with >4 mM tolerance.
B and D) Formaldehyde-free regrowth experiment, in which the selected high-tolerance
population is transferred to medium without formaldehyde and either methanol or
succinate as the carbon substrate, resulting in different shifts in phenotype distribution.

Table 1. Comparison of possible models describing formaldehyde-tolerance phenotype transition processes of *M. extorquens* populations.

For each combination of culture condition and growth substrate, we used a stepwise procedure to evaluate nested models using a likelihood ratio test. Shown below are the best-fit values for the four fitted parameters in each of those models, and test results for each model. α: dependence of death rate on formaldehyde ($h^{-1}*mM^{-1}$). v: advection rate ($mM*h^{-1}$). D: diffusion constant (mM^2*h^{-1}). (for v and D, mM denotes tolerance). b: dependence of death rate on individual tolerance level (mM tolerance / mM formaldehyde in medium). For the likelihood ratio test, name of the model used as the null model, as well as the χ^2 value and p-value, are given. Gray shading: the best-supported model for that experimental scenario. Pseudo-R^2 values for those models were: for formaldehyde selection, 0.973; for methanol regrowth, 0.993; for succinate regrowth, 0.991.

Model	Experimental scenario		Parameters				Likelihood Ratio Test		
	Condition	Substrate	α	b	v	D	null model	χ^2	p
F1	selection	Methanol + Formaldehyde	0.152	n/a	n/a	n/a	n/a	n/a	n/a
F2a	selection	Methanol + Formaldehyde	0.177	0.800	n/a	n/a	F1	0.914	0.339
F2b	selection	Methanol + Formaldehyde	0.156	n/a	-0.020	n/a	F1	17.455	3E-05
F2c	selection	Methanol + Formaldehyde	0.160	n/a	n/a	0.026	F1	18.480	2E-05
F3a	selection	Methanol + Formaldehyde	0.202±0.022	0.770±0.147	n/a	0.019±0.006	F2b	9.379	0.002
F3b	selection	Methanol + Formaldehyde	0.165	n/a	-0.034	0.007	F2b	1.647	0.199
F4	selection	Methanol + Formaldehyde	0.204	0.791	0.007	0.022	F3a	0.029	0.865
M0	regrowth	Methanol	n/a	n/a	n/a	n/a	n/a	n/a	n/a
M1a	regrowth	Methanol	n/a	n/a	0.018±0.006	n/a	M0	7.843	0.005
M1b	regrowth	Methanol	n/a	n/a	n/a	4.94×10^9	M0	0.000	1.000
M2	regrowth	Methanol	n/a	n/a	1.85×10^{-2}	4.60×10^{-8}	M1a	-0.008	1.000
S0	regrowth	Succinate	n/a	n/a	n/a	n/a	n/a	n/a	n/a
S1a	regrowth	Succinate	n/a	n/a	0.189	n/a	S0	97.197	<0.001
S1b	regrowth	Succinate	n/a	n/a	n/a	6.43×10^9	S0	0.000	1.000
S2	regrowth	Succinate	n/a	n/a	0.285±0.026	0.033±0.010	S1a	15.918	<0.001

Supporting Information

Figure S1. Formaldehyde concentrations of ≤5 mM allow growth of *M. extorquens* at a normal rate, but only after a period of lag; higher concentrations lead to longer lag times.

Figure S2. Image processing pipeline to generate colony growth data from formaldehyde-exposed cultures.

Figure S3. Formaldehyde tolerance may be associated with lower fitness on a multicarbon substrate.

Figure S4. Formaldehyde tolerance distributions in *Methylobacterium* populations are robust across experimental replicates, but vary depending on growth conditions.

Figure S5. Cell proliferation assays support the hypothesis that growth of *M. extorquens* in the presence of formaldehyde is due to a small subpopulation of tolerant cells, and that the abundance of tolerant cells decreases with increasing formaldehyde.

Figure S6. The distribution of formaldehyde tolerance within an *M. extorquens* population changes over time depending on growth conditions.

Figure S7. Estimation of growth rates and initial conditions for use in the mathematical model.

Figure S8. The parameter *b* (dependence of death rate on formaldehyde tolerance) determines the shape of the population's phenotypic tolerance distribution after exposure to formaldehyde.

Figure S9. Formaldehyde concentrations in agar growth medium are stable over time and reflective of similar concentrations in liquid medium.

Figure S10. Time-lapse microscopy: cell segmentation and tracking.

Figure S11. Models using extended and original tolerance distributions perform similarly.

Table S1. Results of model selection using original data set for fitting (distribution not extended to account for experimental limit of detection).

File S1. Modeling phenotypic switching in *Methylobacterium extorquens*: R notebook