Fundamental contribution and host range determination of ANP32 protein

family in influenza A virus polymerase activity

Haili Zhang[#], Zhenyu Zhang[#], Yujie Wang, Meiyue Wang, Xuefeng Wang, Xiang

Zhang, Shuang Ji, Cheng Du, Hualan Chen, Xiaojun Wang*

State Key Laboratory of Veterinary Biotechnology, Harbin Veterinary

Research Institute, the Chinese Academy of Agricultural Sciences, Harbin

150069, China

* Corresponding author

Email: wangxiaojun@caas.cn

[#] These authors contributed equally to this work

Running Title: ANP32 proteins are indispensable for influenza virus

ABSTRACT

The polymerase of the influenza virus is part of the key machinery necessary for viral replication. However, the avian influenza virus polymerase is restricted in mammalian cells. The cellular protein ANP32A has been recently found to interact with viral polymerase, and to both influence polymerase activity and interspecies restriction. Here we report that either ANP32A or ANP32B is indispensable for influenza A virus RNA replication. The contribution of ANP32B is equal to that of ANP32A, and together they play a fundamental role in the activity of mammalian influenza A virus polymerase, while neither human ANP32A nor ANP32B support the activity of avian viral polymerase. Interestingly, we found that avian ANP32B was naturally inactive, leaving ANP32A alone to support viral replication. Two amino acid mutations at sites 129-130 in chicken ANP32B lead to the loss of support of viral replication and weak interaction with the viral polymerase complex, and these amino acids are also crucial in the maintenance of viral polymerase activity in other ANP32 proteins. Our findings strongly support ANP32A&B as key factors for both virus replication and adaption.

KEYWORDS: ANP32A; ANP32B; influenza A virus; polymerase activity; RNA replication; interspecies transmission

IMPORTANCE

The key host factors involved in the influenza A viral the polymerase activity and RNA replication remain largely unknown. Here we provide evidence that ANP32A and ANP32B from different species are powerful factors in the maintenance of viral polymerase activity. Human ANP32A and ANP32B contribute equally to support human influenza virus RNA replication. However, unlike avian ANP32A, the avian ANP32B is evolutionarily non-functional in supporting viral replication because of a 129-130 site mutation. The 129-130 site plays an important role in ANP32A/B and viral polymerase interaction, therefore determine viral replication, suggesting a novel interface as a potential target for the development of anti-influenza strategies.

INTRODUCTION

Virus transmission from its natural host species to a different species reflects mechanisms of molecular restriction, evolution, and adaptation. Birds harbor most of the influenza A viruses, which are typically replication-restricted in human hosts because of receptor incompatibility and limited viral polymerase activity in human cells. Although many host factors have been reported to be involved in viral replication (1-8), the key mechanisms that determine viral polymerase activity and host range are poorly understood.

Influenza A viral ribonucleoprotein (vRNP), the viral minimum replicon, comprises the viral genome, the heterotrimeric RNA polymerase PB1, PB2, PA, and the nucleoprotein (NP), and carries out viral RNA transcription and replication in infected cells. Avian influenza A polymerases have very limited activity in mammalian cells, indicating an unknown host-specific restriction mechanism that directly affects viral RNA replication. The adaptation of avian viruses to mammals, such as occurred with the H5N1 and H7N9 avian viruses, occurs along with substitutions on the viral polymerase, mainly on the PB2 subunit (E627K and other signatures), which enhance viral replication (9-22). Many host factors have been reported to interact with the vRNP complex to help viral replication (7, 23-33). Of these proteins, chicken acidic nuclear phosphoprotein 32 family member A (chANP32A) has been reported to specifically promote avian influenza replication due to the 33 amino acids insert

(8). This 33 amino acid insert includes a hydrophobic SUMO interaction motif,

which connects to host SUMOylation to partially contribute to the promotion of

avian viral polymerase activity (34). Furthermore, ANP32A from birds has

splicing variants without SUMO interaction motif also support viral replication

and polymerase adaption (35). Interestingly, a previous study showed that

ANP32A&B (acidic nuclear phosphoprotein 32 family, members A&B) have

been found to regulate viral RNA synthesis (7). These studies suggest that

ANP32A plays an important role in viral replication.

The acidic (leucine-rich) nuclear phosphoprotein 32kDa (ANP32) family

comprises several members, including ANP32A, ANP32B, and ANP32E, which

have various functions in the regulation of gene expression, intracellular

transportation, and cell death (36). Although ANP32A is considered to be an

important co-factor of the influenza virus polymerase and to influence the viral

host range, the roles of different ANP32 members in viral replication, and the

extent to which the proteins are involved in the activity of the polymerase

remain unclear. In this study, by using CRISPR/Cas9 knockout screening, we

found that ANP32A and ANP32B play fundamental roles in the facilitation of

influenza A viral RNA synthesis, and that both ANP32A and ANP32B contribute

equally. The mammalian ANP32 proteins give no or only limited support to the

avian virus polymerase. Human ANP32A&B do not support the replication of

the avian influenza virus, but this restriction can be overcome by E627K

substitution in the viral PB2 protein. Furthermore, we found that chicken

ANP32B has no effect on polymerase activity, which can be ascribed to

mutations in two key amino acid residues of ANP32A&B that determine their

activity in supporting viral RNA replication. Together, these data reveal

fundamental roles for ANP32A and ANP32B in supporting influenza virus A

polymerase activity as well as a site key for their function, and show that both

ANP32A and B determine viral polymerase adaptation and host range.

RESULTS

HuANP32A&B Are Critical Host Factors that Determine Viral Polymerase

Activity and Virus Replication

Previous studies have reported that several host factors, including BUB3,

CLTC, CYC1, NIBP, ZC3H15, C14orf173, CTNNB1, ANP32A, ANP32B,

SUPT5H, HTATSF1, and DDX17, interact with influenza viral polymerase and

that some of these factors have an effect on viral polymerase activity (4, 23,

37). All these observations were based on the gene transient knockdown

technique, meaning that the results may vary because of the different

knockdown efficiency of target genes. Using a CRISPR/Cas9 system, however,

allows rapid knockout of certain genes and accurate evaluation of target

proteins. To identify the critical roles of the above-mentioned host factors in

influenza viral replication, we used a CRISPR/Cas9 system to establish a series

of knockout 293T cell lines, and a model viral-like luciferase RNA was

expressed together with viral polymerase PB1, PB2, PA, and NP, to determine

the polymerase activity in these cell lines. Firstly, we tested the polymerase

activity of 2009 pandemic H1N1 virus A/Sichuan/01/2009(H1N1$_{SC09}$) (38) on

these knockout 293T cells, and found that individual knockout of NIBP, ZC3H15,

or DDX17 results in a 2-4 fold decrease in viral polymerase activity, in which is

consistent with previous results (23, 37). However, in none of the other single

gene knockout cells was viral polymerase activity blocked (Fig. 1A). We did not

observe any reduction of viral polymerase activity in ANP32A or ANP32B

knockout cells, which was a surprising result as ANP32A and ANP32B were

reported to be important host factors supporting viral polymerase activity (7, 8,

34, 35). Since ANP32A and ANP32B have high similarity in both structure and

known functions, we predicted that in single knockout cell lines of ANP32A or

ANP32B, the presence of either one of these two proteins could support viral

replication in the absence of the other. We then developed an ANP32A and

ANP32B double knockout cell line to confirm this hypothesis. In human ANP32A

(huANP32A) knockout 293T cells (AKO cells), and human ANP32B

(huANP32B) knockout 293T cells (BKO cells), the viral polymerases have

similar activities to wild-type 293T cells, but when both ANP32A and ANP32B

were knocked out (DKO) (Fig. S1 in the supplemental material), the polymerase

activity was abolished (about 10000-fold reduction) (Fig 1B), although the

polymerase was expressed equally in the WT, AKO, BKO, and DKO cells (Fig.

1B). We then tested the polymerase activities of 2013 China H7N9 human

isolate A/Anhui/01/2013 (H7N9$_{AH13}$) (39), and H1N1 virus A/WSN/1933 (WSN)

on AKO, BKO and DKO cell lines, and found that the viral polymerase complex

lost all activity in the DKO cells, but not in the AKO or BKO cells (Fig. 1C and

1D). We further confirmed the effect of huANP32A&B on viral infectivity by

infecting 293T cells and DKO cells with WSN virus. In DKO cells, but not AKO

or BKO cells, the infectivity of WSN decreased about 100000-fold (Fig. 1E).

These results indicate a crucial role for both ANP32A&B in viral RNA replication.

The knockout of both ANP32A and ANP32B led to dramatic loss of viral

polymerase activity (about 10000 fold), which is distinct from the previous report,

which used a gene knockdown method and resulted in about 3-5 fold reduction

of viral polymerase activity (8), indicating that ANP32A and ANP32B are of

fundamental importance in viral replication. We found that reconstitution of

either ANP32A or ANP32B, or both of them, in the DKO cells restored the viral

polymerase activities of H1N1$_{SC09}$, H7N9$_{AH13}$, and WSN viruses (Fig. 2A to 2C).

Expression of huANP32A or huANP32B in DKO cells supported the H1N1$_{SC09}$

viral polymerase activity in a dose-dependent manner (Fig. 2D). Reconstitution

of huANP32A or huANP32B or both of them in DKO cells restored full viral

infectivity (Fig. 2E). We confirmed that the required level of expression of

huANP32A or B is very low, and overdose expression has a negative effect (Fig.

S2), suggesting an explanation of the confusing phenotype previously observed,

that in normal 293T cells overexpression of ANP32 protein decreased viral

polymerase activity (8). These data indicate that ANP32A and ANP32B are key

factors required for polymerase activity, that they have similar functions in the

support of viral replication, and that they can function independently and

contribute equally to influenza virus polymerase activity.

Influenza A virus replication starts from the transcription and replication of

the negative single-stranded viral RNA (vRNA) by the vRNP complex. vRNA is

transcribed into positive complementary RNA (cRNA) and mRNA, then the

cRNA is used as a template to amplify into new vRNA (40, 41). We found that

vRNA, cRNA, and mRNA synthesis were dramatically reduced in the DKO cells.

However, when huANP32A was reconstituted in the DKO cells, vRNA, cRNA,

and mRNA synthesis fully recovered (Fig. 3A to 3C). We observed similar

results from reconstitution of ANP32B or both (Fig. 3A to 3C), indicating that

huANP32A&B are key factors in triggering the replication of the viral genome.

We next used an eight-plasmid reverse genetics system in H1N1$_{SC09}$ and

tested the virus production in the supernatant of the transfected cells using an

antigen capture ELISA (42). The result showed that the DKO cells had very low

NP production in the cell supernatant, compared with the 293T, AKO, or BKO

cells (Fig. 3D). The NP production was recovered when the huANP32A and/or

huANP32B was expressed in the DKO cells (Fig. 3E). Taken together, these

results suggest that huANP32A and huANP32B determine viral RNA replication

efficiency and subsequent viral production in 293T cells.

Support of Influenza A Viral Replication by ANP32A or B from Different

Species

ANP32A&B are members of the evolutionarily conserved ANP32 family,

which has various functions in the regulation of gene expression, intracellular

transportat, and cell death (36). We tried to knock out ANP32A&B in both

chicken DF1 cells and human A549 cells, but the knockout caused cell death,

indicating both ANP32A&B have crucial roles in cell proliferation in these cells.

ANP32A&B exist in almost all eukaryotic cells, with the exception of early

eukaryotic life (yeast and other fungi) (43). Molecular phylogenetic analysis of

ANP32 proteins revealed an evolutionary relationship among different species

(Fig. S3). We then investigated the support of ANP32A or ANP32B from

different species for viral polymerase activity in DKO cells. ANP32A from human,

chicken, duck, turkey, zebra finch, mouse, pig, and horse, and ANP32B from

human and chicken, were expressed individually with minigenomes of either

$H1N1_{SC09}$, human isolate $H7N9_{AH13}$, H3N2 Canine influenza virus

A/canine/Guangdong/1/2011($H3N2_{GD11}$), H3N8 equine influenza virus

A/equine/Xinjiang/1/2007(H3N8$_{XJ07}$), A/equine/Jilin/1/1989(H3N8$_{JL89}$), or H9N2

chicken virus A/chicken/Zhejiang/B2013/2012(H9N2$_{ZJ12}$), in DKO cells (Fig 4

and Fig. S4A). We found interestingly that chicken and other avian ANP32A

proteins, which contain an additional 33 amino acids compared to the ANP32

proteins of mammals (8), supported viral polymerase activities in all cases,

whereas the ANP32A proteins from human, pig, and horse, as well as human

ANP32B, supported mammalian influenza virus polymerase activities, but not

that of chicken H9N2$_{ZJ12}$, or the H3N8$_{JL89}$ (an avian-like virus). This result is

consistent with previous reports that avian ANP32A can promote avian viral

polymerase activity in human cells (8, 35). PB2 is the most important

polymerase subunit, and affects host range (25, 44). Almost all the avian

viruses had a glutamic acid (E) at PB2 residue 627, while it could be rapidly

selected as a lysine (K) when the virus adapted to mammals, accompanied with

increased pathogenicity and transmission abilities. The PB2 E627K mutation

has long been regarded as a key signature of the avian influenza virus in

overcoming the block to replicate in mammalian cells (9, 20, 21, 45-49). We

observed that the K627E mutation in H7N9$_{AH13}$ lost support from mammalian

ANP32 proteins (Fig. S4A and S4B), while E627K mutations in isolates of avian

origin (horse H3N8$_{JL89}$, or chicken H9N2$_{ZJ12}$) dramatically changed viral fitness

to mammalian ANP32 proteins (Fig. S4B and S4C). These results revealed that

ANP32A&B play important roles in influenza A viral replication across different

species, and that mammalian ANP32 proteins provide poor support for avian

influenza viral RNA replication.

A Novel 129-130 Site Vital for Influenza Polymerase Activity in ANP32A&B

Surprisingly, chicken ANP32B did not support any viral RNA replication, and

mouse ANP32A showed limited support to H1N1$_{SC09}$ and H3N8$_{XJ07}$, indicating

a functional loss in these molecules (Fig. 4). The ANP32 protein family has a

conserved structure containing an N-terminal leucine-rich repeat (LRR) and a

C-terminal low-complexity acidic region (LCAR) domain (36). ANP32A&B have

been reported to interact directly with the polymerase complex (7, 35). The key

functional domains of ANP32 proteins remain largely unknown. We noticed that

the chicken ANP32B (chANP32B) was functionally inactive and hypothesized

that certain amino acids may be responsible for this phenotype. Alignment of

huANP32B, pgANP32B, and chANP32B revealed scattered substitutions on

the proteins (Fig. S5A). Chimeric clones between chicken and human ANP32B

were constructed and evaluated (Fig. S5B). We found that the replacement of

amino acids 111-160 of huANP32B with those of chANP32B aborted the activity

of this protein, while chANP32B with human fragment 111-161 gained the ability

to boost viral polymerase activity (Fig. S5C and S5D). Further comparing of

ANP32B fragments 111-161 of chicken, human, and pig showed eight amino

acid substitutions (Fig. 5A). Of these, a combined mutation N129I/D130N, but

not others, on huANP32B impaired H1N1 polymerase activity (Fig. 5B), and the

N129I showed stronger impairment than did the D130N mutation (Fig. 5C). We

confirmed this phenotype on H7N9 polymerase activity (Fig. 5D). The 129-130

mutations also aborted the support of huANP32A of H1N1 and H7N9

polymerases (Fig. 5E and 5F). Furthermore, substitutions at 129-130 in

chANP32A and chANP32B reversed the support of these proteins of the H7N9

virus human isolate (Fig. 5G). Interestingly, in the content of a chicken H7N9

viral minigenome, although mutation of 129-130 of chANP32A impaired the

polymerase activity, the chANP32B reverse mutation did not restore its support

to chicken-like H7N9 viral RNA replication (Fig. 5H), indicating that chicken

virus replication may require both the 129-130 site and an extra 33-aminal acid

peptide as in chANP32A. The expression levels of mutant ANP32 proteins were

assessed using western blotting (Fig. S6). Most terrestrial mammalian

ANP32A&B proteins have a 129-ND-130 signature, but most of the avian

ANP32Bs have 129-IN-130 residues (Table S1). Murine ANP32A (muANP32A)

has 129-NA-130 and this may explain the impaired support of H1N1

polymerase by muANP32A (Fig. 4 and Fig. S7). To further confirm the function

of the 129-130 site, we made inactivating mutations of huANP32A and

huANP32B at the sites 129-130 (N129I, D130N) on the 293T genome using

CRISPR-Cas9 mediated recombination (Fig. 6A and Fig. S8); the double

mutated proteins expressed well and fully impaired the replication of both

H1N1$_{SC09}$ and H7N9$_{AH13}$ (Fig. 6B and 6C). Interestingly, the single ANP32A or

ANP32B mutations lead to slightly decreased polymerase activities compared

with wildtype 293T cells, suggesting competition between the mutated proteins

and the native proteins for viral polymerase binding.

The 129-130 Site in ANP32B Affects Its Interaction with Viral Polymerase

It has been reported that ANP32A variants interact with the polymerase

subunit PB2 and that the avian ANP32A can enhance avian viral RNP assembly

in human cells (35). ANP32A and ANP32B can interact with polymerase only

when 3 subunits of the polymerase are present (7). ANP32B proteins from

different species have a conserved structure that comprises a N-terminal

leucine-rich repeat (LRR) region and a C terminal low complexity acidic region

(LCAR) (Fig. 7A). To investigate the function of the 129-130 site in ANP32B in

the interaction of this protein with viral polymerase, we used variants of

huANP32B, chANP32B to co-transfect with WSN minireplicon into DKO cells.

Co-immunoprecipitations detected strong interactions between huANP32A and

the polymerase subunits PB1 and PA (Fig. 7B, Lane 1), but weak interactions

when chANP32B was present (Fig. 7B, Lane 2). No interactions were detected

when PB1 was absent (Fig. 7B, Lane 3), indicating this interaction occurred

between ANP32B and the viral trimeric polymerase complex, which is

consistent with previous findings (7, 34). As controls, we also observed that the

LCAR truncations of huANP32B were unable to interact with the viral

polymerase (Fig. 7B, Lane 4-6), in agreement with previous observations (34).

When we mutated the functional human ANP32B at the site 129-130 to the

chicken ANP32B signature, the huANP32B lost the ability to interact with viral

polymerase. Conversely, chANP32B gained the ability to co-

immumoprecipitate with viral polymerase when its 129-130 sites were mutated

to the human ANP32B signature (Fig. 7C). Together these results revealed a

fundamental function of ANP32A and ANP32B in supporting influenza A viral

polymerase activity and a novel 129-130 site of ANP32A&B in different species

that may influence influenza virus replication.

DISCUSSION

Although the host factors that are involved in influenza A virus replication

have been long investigated, and genome-wide screening has shown that many

host proteins interact with viral polymerase, the key mechanisms that determine

viral polymerase activity in different cells remain largely unclear. ANP32A and

ANP32B have been previously identified binding with viral polymerase and

promoting human influenza viral vRNA synthesis (7). Furthermore, a recent

study showed for the first time that chicken ANP32A can rescue avian

polymerase activity in human cells because the chicken ANP32A harbors an

extra stretch of 33 amino acids which is absent in mammalian ANP32A (8).

However, the potencies of ANP32A and ANP32B (as well as other host factors)

in viral replication are not well investigated in different hosts. In our study, we

used CRISPR/Cas9 knockout cells to screen the candidate host proteins

involved in viral RNA replication. We identified that ANP32A and ANP32B are

key host co-factors that determine influenza A virus polymerase activity. Without

ANP32A&B, the viral polymerase activity decreased by 10000 fold and the viral

infectivity decreased by more than 100000 fold. In contrast, the DDX17

knockout cells showed about 3-fold decreased viral polymerase activity (Fig.

1A), which was in agreement with prior reports (23). We found that ANP32A or

ANP32B can independently restore viral polymerase activity, indicating that

both have a similar function in viral replication (7).

Influenza viruses and their hosts have a long co-evolution history. ANP32

family members are expressed in animal and plant cells, with both conserved

LRR and LCAR domains (43). Three conserved ANP32 members (ANP32A,

ANP32B, and ANP32E) in vertebrates were reported to have several functions

in cells. We found that knockout of both ANP32A and ANP32B abolished the

influenza viral replication, while the ANP32E may not be involved in viral

replication. Interestingly, we found that the chicken ANP32A keeps the

conserved function to support both avian and mammalian virus polymerase

activity, while human ANP32A or B did not support avian viral replication. Thus,

we demonstrate here that ANP32A and ANP32B are key host factors that play

a fundamental role in influenza A viral RNA replication.

A recent study revealed that avian ANP32A has a hydrophobic SUMO

interaction motif (SIM) in an extra 33 amino acid insert region which connects

to host SUMOylation to specifically promote avian viral polymerase activity (34).

However, this SIM-like domain is not present in all of the ANP32B proteins from

different hosts, and neither is it found in mammalian ANP32A. We showed that

most ANP32Bs from different species are functional as ANP32As in support of

viral replication, but chANP32B is naturally unable to support polymerase

activity (Fig. 4). We finally found that the amino acids 129I and 130N of

chANP32B are responsible for the loss of this function. Sequence analysis and

mutagenesis studies suggest that the 129-130 sites are important for

maintaining the function of avian ANP32A and human ANP32A/B in viral

replication (Fig. 5). The co-immunoprecipitation assay showed that mutations

on these 129-130 sites changed the interaction efficiency between the ANP32

proteins and viral polymerase complex (Fig. 7). Mutating the 129-130 sites in

chANP32B to the functional signature 129N and 130D enables chANP32B to

support polymerase activity in human viruses, but not chicken viruses with PB2

627E, indicating that the ANP32B may undergo selection in two areas during

co-evolution with the avian influenza virus: chANP32B does not harbor an extra

insert as chANP32A and the 129-130 mutations. This result also suggests that

avian ANP32A is the only protein from the avian ANP32 family to support avian

viral replication.

Together, out data give new insights into the functions of ANP32A and

ANP32B. The 129-130 substitution could be used as a novel target to modify

the genome of animals to develop influenza-A-resistant transgenic chickens, or

other animals, which will benefit the husbandry industry, as well as animal and

human health. Further investigation into the molecular mechanisms that

determine how ANP32 proteins work with the viral polymerase complex, for

example, the structure of the ANP32 and vRNP complex, and the fitness of

different virus subtypes would contribute to the understanding viral

pathogenesis and host defense.

MATERIALS AND METHODS

Cells, Viruses, and Plasmids

Human embryonic kidney (293T, ATCC CRL- 3216) and Madin-Darby

canine kidney (MDCK, CCL-34™) cells were maintained in Dulbecco's modified

Eagle's medium (DMEM, Hyclone) with 10% fetal bovine serum (FBS; Sigma),

1% penicillin and streptomycin (Gibco), and kept at 37°C with 5% CO_2. Certain

reagents were kindly provided by the following persons: polymerase plasmids

of H1N1 human influenza A virus A/Sichuan/01/2009 (H1N1$_{SC09}$), and H7N9

A/Anhui/01/2013 (H7N9$_{AH13}$) by Dr. Hualan Chen; H1N1 human influenza virus

A/WSN/1933 (WSN) by Dr Kawoaka; H3N2 canine influenza virus

A/canine/Guangdong/1/2011 (H3N2$_{GD12}$) by Dr Shoujun Li from China Southern

Agriculture University; and H9N2 avian influenza virus

A/chicken/Zhejiang/B2013/2012 (H9N2$_{ZJ12}$) by Dr Zejun Li from Shanghai

Veterinary Research Institute of CAAS. H3N8 equine influenza viruses

A/equine/Jilin/1/1989 (H3N8$_{JL89}$) and A/equine/Xinjiang/1/2007 (H3N8$_{XJ07}$)

were preserved in our lab. The reverse genetics system based on the pBD

vector for the H1N1$_{SC09}$ virus was established in Dr Hualan Chen's lab. The

pCAGGS plasmids of containing full length ANP32A isoforms of several species

were generated by gene synthesis (Synbio technologies, China) according to

the sequences deposited in GenBank, including chicken ANP32A (chANP32A,

XM_413932.5, XP_413932.3), human ANP32A(huANP32A, NM_006305.3,

NP_006296.1), zebra finch ANP32A (zfANP32A, XM_012568610.1,

XP_012424064.1), duck ANP32A (dkANP32A, XM_005022967.1,

XP_005023024.1), turkey ANP32A (tyANP32A, XM_010717616.1,

XP_010715918.1), pig ANP32A (pgANP32A, XM_003121759.6,

XP_003121807.3), mouse ANP32A (muANP32A, NM_009672.3,

NP_033802.2), equine ANP32A (eqANP32A, XM_001495810.5,

XP_001495860.2), chicken ANP32B (chANP32B, NM_001030934.1,

NP_001026105.1) and human ANP32B (huANP32B, NM_006401.2,

NP_006392.1). Site-directed mutants of these sequences were generated

using overlapping PCR and identified by DNA sequencing. Mutants of pcAGGS-

huANP32B-Δ216/190/165 and pcDNA3.1-PA-V5 were constructed according

to the online In-Fusion ®HD Cloning Kit User Manual

(http://www.clontech.com/CN/Products/Cloning_and_Competent_Cells/Clonin

g_Kits/xxclt_searchResults.jsp). Briefly, the fragments of the

pcAGGS/pcDNA3.1 vector and each target gene were amplified with a 15 bp

homologous arm and were then fused using the In-Fusion HD Enzyme

(Clontech, Felicia, CA, USA). To create the pcAGGS-huANP32B-Δ216/190/165

plasmids, pcAGGS-huANP32B was used as the template to amplify the

pcAGGS vector. This sequence was then fused with different truncated

huANP32B fragments (huANP32B-Δ216/190/165). To obtain pcDNA3.1-PA-V5

plasmid, pBD- H1N1$_{SC09}$-PA was used as the template to amplify the PA-V5

sequence, and then fused with pcDNA3.1 vector.

Knockout Cell Lines

To generate knockout cell lines for host proteins BUB3(AF081496.1),

CLTC(NM_004859.3), CYC1 (CR541674.1), NIBP (BC006206.2), ZC3H15

(NM_018471.2), C14orf173 (DQ395340.1), CTNNB1 (NM_001904.3), ANP32A

(NM_006305.3), ANP32B (NM_006401.2), SUPT5H (U56402.1), HTATSF1

(NM_014500), and DDX17 (NM_006386, NM_001098504) (4, 23, 37), the

gRNAs design tool (http://crispr.mit.edu/) was used for gRNAs design and off-

target prediction (50). DNA fragments that contained the U6 promoter, gRNAs

specific for host factors, a guide RNA scaffold, and U6 termination signal

sequence were synthesized and subcloned into the pMD18-T backbone vector.

The Cas9-eGFP expression plasmid (pMJ920) was a gift from Jennifer Doudna

(Addgene plasmid # 42234) (51). Briefly, 293T cells in 6-well plates were

transfected with 1.0 µg pMJ920 plasmids and 1.0 µg gRNA expression

plasmids by Lipofectamine® 2000 Transfection Reagent (Invitrogen,

Cat.11668-027) using the recommended protocols. GFP-positive cells were

sorting by fluorescence-activated cell sorting (FACS) at 48 h post-transfection,

then monoclonal knockout cell lines were screened by western blotting and/or

DNA sequencing.

Generation of Site-directed Amino-acid-substituted 293T Cell Line

High efficiency guide sequences for ANP32A and ANP32B which bind

upstream and downstream with close proximity to the target (129/130 ND) were

chosen. The gRNA expression plasmids were constructed as described above.

An 80nt oligo with the desired mutations at the target site was used as the donor

DNA. 293T cells were transfected with 1 μg pMJ920 plasmids, 1 μg gRNA

expression plasmids, and 50 pmol donor DNA. 48 h later, GFP positive cells

were isolated by FACS, and site-directed mutagenesis clones were identified

after screening by sequencing and western blotting using Anti-PHAP1 antibody

(ab51013) and Anti-PHAPI2 / APRIL antibody EPR14588 (ab200836). Cell lines

with double gene mutations were generated by second round transfection and

selection.

Polymerase Assay

A minigenome reporter, which contains the firefly luciferase gene flanked

by the non-coding regions of the influenza HA gene segment with a human polI

promoter and a mouse terminator sequence (52), was transfected with viral

polymerase and NP expression plasmids to analyze the polymerase activity.

Mutants of PB2 genes were generated using overlapping PCR and identified

by DNA sequencing. To determine the effect of ANP32 proteins on viral

polymerase activity, 293T or KO cells in 12-well plates were transfected with

plasmids of the PB1 (80 ng), PB2 (80 ng), PA (40 ng) and NP (160 ng), together

with 80 ng minigenome reporter and 10 ng *Renilla* luciferase expression

plasmids (pRL-TK, kindly provided by Dr Luban*)*, using Lipofectamine 2000

transfection reagent (Invitrogen) according to the manufacturers' instructions.

Cells were incubated at 37°C. Cells were lysed with 100ul of passive lysis buffer

(Promega) at 24 h after transfection, and firefly and *Renilla* luciferase activities

was measured using a Dual-luciferase kit (Promega) with Centro XS LB 960

luminometer (Berthold technologies). The function of ANP32 was examined

using a polymerase assay by co-transfection of DKO cells with different ANP32

proteins for 24 h. All the experiments were performed independently at least

three times. Results represent the mean ± SEM of the replicates within one

representative experiment. The expression levels of polymerase proteins on

different cell lines were detected by western blotting, using specific mouse

monoclonal antibodies for NP and PB1 proteins, anti-V5 tag antibody (ab27671)

for PA-V5 protein.

RNA isolation, Reverse Transcription, and Quantification by RT- PCR

Total RNA from 293T cells was extracted using an RNeasy mini kit (Qiagen),

following the manufacturer's instructions. For the synthesis of first strand cDNA

derived from firefly luciferase RNAs driven by influenza polymerase, equal

concentrations of RNA were subject to cDNA synthesis using a Reverse

Transcription Kit (PrimeScript™ RT reagent Kit with a gDNA Eraser (Perfect

Real Time), Cat.RR047A). Primers used in the RT reaction were as follows: 5'-

CATTTCGCAGCCTACCGTGGTGTT-3' for the firefly luciferase vRNA; 5'-

AGTAGAAACAAGGGTG-3' for the firefly luciferase cRNA; oligo-dT20 for the

firefly luciferase mRNA (53). The cDNA samples were subjected to

quantification by real-time PCR with specific primers F (5'-

GATTACCAGGGATTTCAGTCG-3') and R (5'-GACACCTTTAGGCAGACCAG-

3') using SYBR ® Premix Ex TaqTM II (Tli RNaseH Plus) (TaKaRa Cat:

RR820A). Fold change in RNA was calculated by double-standard curve

methods and β-actin as an internal control.

Quantitative ELISA for Determination of Virus Production

The ELISA assay has been previously described(42). Briefly, a 96-well

microtiter plate (Costar, Bodenheim, Germany) was coated with 1μg/well of the

mouse monoclonal anti-IAV NP protein antibody in phosphate-buffered saline

(PBS), incubated overnight at 4 °C or 37 °C for 2 h. The plate was washed three

times with washing buffer (PBS containing 0.1% Tween-20, PBST) and blocked

with 200 μl 5% calf serum at 37 °C for 2 h. After washing three times with PBST

buffer, 100 μl of virus/VLPS-containing samples were added in dilution buffer

(PBS containing 10% calf serum and 0.1% Triton X-100) and incubated at 37 °C

for 1 h. The plate was then washed, and 100 μl of a 1:2000 dilution of HRP-

conjugated anti-IAV NP protein mAb was added. After incubation at 37 °C for

0.5 h, the plate was washed again and incubated with freshly prepared TMB

peroxidase substrate (Galaxy Bio, Beijing, China) for 10 min at room

temperature. The reaction was stopped by adding 2 M H_2SO_4, and the optical

density at 450 nm wavelength (OD450) was measured using the VersaMax

Microplate Reader (BioTek, Winooski, USA). Dilution buffer was taken as a

blank control and the purified IAV-NP protein was double diluted eight times in

dilution buffer to obtain the standard curve. The virus production was calculated

using the standard curve.

Influenza Virus Infection and Infectivity

Human influenza virus WSN was rescued from a 12-plasmid rescue system

(54). Briefly, 293T culture in a 6-well plate was transfected with 0.1μgof each

pPoll plasmids and 1ug each of pCAGGS-NP, pCAGGS-PA, pCAGGS-PB1,

and pCAGGS-PB2 using Lipofectamine 2000 (Invitrogen) in Opti-MEM. Eight

hours post-transfection, medium was changed to DMEM plus 10% FBS. The

supernatant was harvested after 48 h and injected into 9-day-old specific

pathogen-free embryonated eggs for virus propagation. Eggs were incubated

at 35 °C for 72 h, the allantoic fluid from eggs was detected by

haemagglutination assay and the titers of virus on 293T cells were determined

using the method of Reed and Muench (55). Then 293T cells, huANP32A

knockout 293T cells (AKO cells), huANP32B knockout 293T cells (BKO cells),

and huANP32A &B double knocked out 293T cells (DKO) were infected at a

multiplicity of infection (MOI) of 0.01 for 1 h, washed twice with PBS, and then

cultured at 37°C in Opti-MEM containing 0.5% FBS and tosylsulfonyl

phenylalanyl chloromethyl ketone (TPCK)-trypsin (Sigma) at 1mg/ml. At the

indicated time points, the culture supernatant was harvested. Viral titers in

MDCK cells were determined as described above.

Immunoprecipitation and Western Blotting

For immunoprecipitation and western blotting, transfected cells were lysed

using an ice-cold lysis buffer (50 mM Hepes-NaOH [pH 7.9], 100 mM NaCl, 50

mM KCl, 0.25% NP-40, and 1 mM DTT), and centrifuged at 13,000× g and 4 °C

for 10 min. After centrifugation, the crude lysates were incubated with Anti-

FLAG M2 Magnetic Beads (SIGMA-ALDRICH, M8823) or Anti-NP Magnetic

Beads (MCE Protein A/G Magnetic Beads and NP antibody from our lab) at 4°C

for 2 hr. After incubation, the resins were collected using a magnetic separator

and washed three times with PBS. The resin-bound materials were eluted using

a 3X Flag peptide or by boiling in the SDS-PAGE loading buffer, subjected to

SDS-PAGE and then transferred onto nitrocellulose membranes. Membranes

were blocked with 5% milk powder in Tris-buffered saline (TBS) for 2 h.

Incubation with the first anti-mouse antibody (Anti-Flag antibody from SIGMA

(F1804), Anti-V5 antibody from Abcam (ab27671), Anti-NP and PB1 antibody

from our lab) was performed for 2 h at room temperature (RT), followed by

washing three times with TBST. The secondary antibody (Sigma, 1:10,000) was

then applied and samples were incubated at RT for 1 h. Subsequently,

membranes were washed three times for 10 min with TBST. Signals were

detected using a LI-COR Odyssey Imaging System (LI-COR, Lincoln, NE, USA).

Statistics. Statistical analysis was performed in GraphPad Prism, version 5

(Graph Pad Software, USA). Statistical differences between groups were

assessed by One-way ANOVA followed by a Dunnett's post-test. All the

experiments were performed independently at least three times. Error bars

represent the standard deviation (SD) or the standard error of the mean (SEM)

in each group, as indicated in figure legends. NS, not significant ($p > 0.05$),

*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

ACKNOWLEDGEMENTS

539 We thank W. Barclay, SJ. Li, ZJ. Li, J. Luban, and J. Doudna for providing the

540 plasmids and advice. We thank CJ. Li and DM. Zhao for discussions.

FUNDING INFORMATION

This work was supported by the grant from the Natural Science Foundation of

China to HL Chen (No. 31521005).

REFERENCES

1.   **Brass AL, Huang IC, Benita Y, John SP, Krishnan MN, Feeley EM, Ryan BJ, Weyer**

JL, van der Weyden L, Fikrig E, Adams DJ, Xavier RJ, Farzan M, Elledge SJ. 2009.

The IFITM proteins mediate cellular resistance to influenza A H1N1 virus, West Nile

virus, and dengue virus. Cell **139:**1243-1254.

2.   **Konig R, Stertz S, Zhou Y, Inoue A, Hoffmann HH, Bhattacharyya S, Alamares JG,**

Tscherne DM, Ortigoza MB, Liang Y, Gao Q, Andrews SE, Bandyopadhyay S, De

Jesus P, Tu BP, Pache L, Shih C, Orth A, Bonamy G, Miraglia L, Ideker T, Garcia-

Sastre A, Young JA, Palese P, Shaw ML, Chanda SK. 2010. Human host factors

required for influenza virus replication. Nature **463:**813-817.

3.   **Watanabe T, Kawakami E, Shoemaker JE, Lopes TJS, Matsuoka Y, Tomita Y,**

Kozuka-Hata H, Gorai T, Kuwahara T, Takeda E, Nagata A, Takano R, Kiso M,

Yamashita M, Sakai-Tagawa Y, Katsura H, Nonaka N, Fujii H, Fujii K, Sugita Y,

Noda T, Goto H, Fukuyama S, Watanabe S, Neumann G, Oyama M, Kitano H,

Kawaoka Y. 2014. Influenza Virus-Host Interactome Screen as a Platform for Antiviral

Drug Development. Cell Host & Microbe **16:**795-805.

4.   **Bradel-Tretheway BG, Mattiacio JL, Krasnoselsky A, Stevenson C, Purdy D,**

Dewhurst S, Katze MG. 2011. Comprehensive proteomic analysis of influenza virus

polymerase complex reveals a novel association with mitochondrial proteins and RNA

polymerase accessory factors. J Virol **85:**8569-8581.

5.   **Shapira SD, Gat-Viks I, Shum BO, Dricot A, de Grace MM, Wu L, Gupta PB, Hao**

T, Silver SJ, Root DE, Hill DE, Regev A, Hacohen N. 2009. A physical and regulatory

map of host-influenza interactions reveals pathways in H1N1 infection. Cell **139:**1255-

1267.

6.   **Karlas A, Machuy N, Shin Y, Pleissner K-P, Artarini A, Heuer D, Becker D, Khalil**

H, Ogilvie LA, Hess S, Maurer AP, Muller E, Wolff T, Rudel T, Meyer TF. 2010.

Genome-wide RNAi screen identifies human host factors crucial for influenza virus

replication. Nature **463:**818-822.

7.   **Sugiyama K, Kawaguchi A, Okuwaki M, Nagata K.** 2015. pp32 and APRIL are host

cell-derived regulators of influenza virus RNA synthesis from cRNA. Elife **4**.

8.   **Long JS, Giotis ES, Moncorge O, Frise R, Mistry B, James J, Morisson M, Iqbal**

M, Vignal A, Skinner MA, Barclay WS. 2016. Species difference in ANP32A underlies

influenza A virus polymerase host restriction. Nature **529:**101-104.

9.   **Subbarao EK, London W, Murphy BR.** 1993. A single amino acid in the PB2 gene of

influenza A virus is a determinant of host range. J Virol **67:**1761-1764.

10.   **Xiao C, Ma W, Sun N, Huang L, Li Y, Zeng Z, Wen Y, Zhang Z, Li H, Li Q, Yu Y,**

Zheng Y, Liu S, Hu P, Zhang X, Ning Z, Qi W, Liao M. 2016. PB2-588 V promotes the

mammalian adaptation of H10N8, H7N9 and H9N2 avian influenza viruses. Sci Rep

6:19474.

11.   **Song W, Wang P, Mok BW, Lau SY, Huang X, Wu WL, Zheng M, Wen X, Yang S,**

Chen Y, Li L, Yuen KY, Chen H. 2014. The K526R substitution in viral protein PB2

enhances the effects of E627K on influenza virus replication. Nat Commun **5:**5509.

12.   **Sediri H, Schwalm F, Gabriel G, Klenk HD.** 2015. Adaptive mutation PB2 D701N

promotes nuclear import of influenza vRNPs in mammalian cells. Eur J Cell Biol

94:368-374.

13.   **Manz B, de Graaf M, Mogling R, Richard M, Bestebroer TM, Rimmelzwaan GF,**

Fouchier RA. 2016. Multiple Natural Substitutions in Avian Influenza A Virus PB2

Facilitate Efficient Replication in Human Cells. J Virol **90:**5928-5938.

14.   **Liu Q, Qiao C, Marjuki H, Bawa B, Ma J, Guillossou S, Webby RJ, Richt JA, Ma W.**

2012. Combination of PB2 271A and SR polymorphism at positions 590/591 is critical

for viral replication and virulence of swine influenza virus in cultured cells and in vivo.

J Virol **86:**1233-1237.

15.   **Zhang H, Li X, Guo J, Li L, Chang C, Li Y, Bian C, Xu K, Chen H, Sun B.** 2014. The

PB2 E627K mutation contributes to the high polymerase activity and enhanced

replication of H7N9 influenza virus. J Gen Virol **95:**779-786.

16.   **Mok CK, Lee HH, Lestra M, Nicholls JM, Chan MC, Sia SF, Zhu H, Poon LL, Guan**

Y, Peiris JS. 2014. Amino acid substitutions in polymerase basic protein 2 gene

contribute to the pathogenicity of the novel A/H7N9 influenza virus in mammalian hosts.

J Virol **88:**3568-3576.

17.   **Hu M, Yuan S, Zhang K, Singh K, Ma Q, Zhou J, Chu H, Zheng BJ.** 2017. PB2

substitutions V598T/I increase the virulence of H7N9 influenza A virus in mammals.

Virology **501:**92-101.

18.   **Li Z, Chen H, Jiao P, Deng G, Tian G, Li Y, Hoffmann E, Webster RG, Matsuoka Y,**

Yu K. 2005. Molecular basis of replication of duck H5N1 influenza viruses in a

mammalian mouse model. J Virol **79:**12058-12064.

19.   **Gao Y, Zhang Y, Shinya K, Deng G, Jiang Y, Li Z, Guan Y, Tian G, Li Y, Shi J, Liu**

L, Zeng X, Bu Z, Xia X, Kawaoka Y, Chen H. 2009. Identification of amino acids in HA

and PB2 critical for the transmission of H5N1 avian influenza viruses in a mammalian

host. Plos Pathog **5:**e1000709.

20.   **Steel J, Lowen AC, Mubareka S, Palese P.** 2009. Transmission of Influenza Virus in

a Mammalian Host Is Increased by PB2 Amino Acids 627K or 627E/701N. Plos Pathog

5(1):e1000252.

21.   **Hatta M, Gao P, Halfmann P, Kawaoka Y.** 2001. Molecular basis for high virulence of

Hong Kong H5N1 influenza A viruses. Science **293:**1840-1842.

22.   **Bussey KA, Bousse TL, Desmet EA, Kim B, Takimoto T.** 2010. PB2 residue 271

plays a key role in enhanced polymerase activity of influenza A viruses in mammalian

host cells. J Virol **84:**4395-4406.

23.   **Bortz E, Westera L, Maamary J, Steel J, Albrecht RA, Manicassamy B, Chase G,**

Martinez-Sobrido L, Schwemmle M, Garcia-Sastre A. 2011. Host- and strain-specific

regulation of influenza virus polymerase activity by interacting cellular proteins. MBio

2:e00151-00111.

24.   **Diot C, Fournier G, Santos MD, Magnus J, Komarova A, Werf SVD, Munier S,**

Naffakh N. 2016. Influenza A Virus Polymerase Recruits the RNA Helicase DDX19 to

Promote the Nuclear Export of Viral mRNAs. Scientific Reports **6:**33763.

25. **Eisfeld AJ, Neumann G, Kawaoka Y.** 2015. At the centre: influenza A virus

ribonucleoproteins. Nat Rev Microbiol **13:**28-41.

26. **Gabriel G, Klingel K, Otte A, Thiele S, Hudjetz B, Arman-Kalcek G, Sauter M,**

Shmidt T, Rother F, Baumgarte S, Keiner B, Hartmann E, Bader M, Brownlee GG,

Fodor E, Klenk HD. 2011. Differential use of importin-alpha isoforms governs cell

tropism and host adaptation of influenza virus. Nature Communications **2**(2):156.

27. **Landeras-Bueno S, Jorba N, Pérez-Cidoncha M, Ortín J.** 2011. The Splicing Factor

Proline-Glutamine Rich (SFPQ/PSF) Is Involved in Influenza Virus Transcription. Plos

Pathog **7:**e1002397.

28. **Marcosvillar L, Pazo A, Nieto A.** 2016. Influenza Virus and Chromatin: Role of the

CHD1 Chromatin Remodeler in the Virus Life Cycle. Journal of Virology **90:**3694.

29. **Momose F, Naito T, Yano K, Sugimoto S, Morikawa Y, Nagata K.** 2002. Identification

of Hsp90 as a stimulatory host factor involved in influenza virus RNA synthesis. J Biol

Chem **277:**45306-45314.

30. **Shaw ML, Stertz S.** 2017. Role of Host Genes in Influenza Virus Replication. In: Tripp

R., Tompkins S. (eds) Roles of Host Gene and Non-coding RNA Expression in Virus

Infection. Current Topics in Microbiology and Immunology, vol 419. Springer, Cham.

31. **Su WC, Hsu SF, Lee YY, Jeng KS, Lai MM.** 2015. A Nucleolar Protein, Ribosomal

RNA Processing 1 Homolog B (RRP1B), Enhances the Recruitment of Cellular mRNA

648 in Influenza Virus Transcription. Journal of Virology **89:**11245.

649 32. **Ver LS, Marcos-Villar L, Landeras-Bueno S, Nieto A, Ortín J.** 2015. The Cellular

650 Factor NXP2/MORC3 Is a Positive Regulator of Influenza Virus Multiplication. Journal

651 of Virology **89:**10023-10030.

652 33. **Zhang J, Li G, Ye X.** 2010. Cyclin T1/CDK9 Interacts with Influenza A Virus Polymerase

653 and Facilitates Its Association with Cellular RNA Polymerase II. Journal of Virology

654 **84:**12619-12627.

655 34. **Domingues P, Hale BG.** 2017. Functional Insights into ANP32A-Dependent Influenza

656 A Virus Polymerase Host Restriction. Cell reports **20:**2538-2546.

657 35. **Baker SF, Ledwith MP, Mehle A.** 2018. Differential Splicing of ANP32A in Birds Alters

658 Its Ability to Stimulate RNA Synthesis by Restricted Influenza Polymerase. Cell reports

659 **24:**2581-2588 e2584.

660 36. **Reilly PT, Yu Y, Hamiche A, Wang L.** 2014. Cracking the ANP32 whips: important

661 functions, unequal requirement, and hints at disease implications. Bioessays **36:**1062-

662 1071.

663 37. **Watanabe T, Kawakami E, Shoemaker JE, Lopes TJ, Matsuoka Y, Tomita Y,**

664 **Kozuka-Hata H, Gorai T, Kuwahara T, Takeda E, Nagata A, Takano R, Kiso M,**

665 **Yamashita M, Sakai-Tagawa Y, Katsura H, Nonaka N, Fujii H, Fujii K, Sugita Y,**

666 **Noda T, Goto H, Fukuyama S, Watanabe S, Neumann G, Oyama M, Kitano H,**

667 **Kawaoka Y.** 2014. Influenza virus-host interactome screen as a platform for antiviral

668 drug development. Cell Host Microbe **16:**795-805.

38.   **Zhang Y, Zhang Q, Kong H, Jiang Y, Gao Y, Deng G, Shi J, Tian G, Liu L, Liu J,**

Guan Y, Bu Z, Chen H. 2013. H5N1 hybrid viruses bearing 2009/H1N1 virus genes

transmit in guinea pigs by respiratory droplet. Science **340:**1459-1463.

39.   **Zhang Q, Shi J, Deng G, Guo J, Zeng X, He X, Kong H, Gu C, Li X, Liu J, Wang G,**

Chen Y, Liu L, Liang L, Li Y, Fan J, Wang J, Li W, Guan L, Li Q, Yang H, Chen P,

Jiang L, Guan Y, Xin X, Jiang Y, Tian G, Wang X, Qiao C, Li C, Bu Z, Chen H. 2013.

H7N9 influenza viruses are transmissible in ferrets by respiratory droplet. Science

341:410-414.

40.   **Hay AJ, Skehel JJ, McCauley J.** 1982. Characterization of influenza virus RNA

complete transcripts. Virology **116:**517-522.

41.   **Young RJ, Content J.** 1971. 5'-terminus of influenza virus RNA. Nat New Biol **230:**140-

142.

42.   **Wang M, Zhang Z, Wang X.** 2018. Strain-Specific Antagonism of the Human H1N1

Influenza A Virus against Equine Tetherin. Viruses **10**(5):264.

43.   **Matilla A, Radrizzani M.** 2005. The Anp32 family of proteins containing leucine-rich

repeats. Cerebellum **4:**7-18.

44.   **Almond JW.** 1977. A single gene determines the host range of influenza virus. Nature

270:617-618.

45.   **Gao R, Cao B, Hu Y, Feng Z, Wang D, Hu W, Chen J, Jie Z, Qiu H, Xu K, Xu X, Lu**

H, Zhu W, Gao Z, Xiang N, Shen Y, He Z, Gu Y, Zhang Z, Yang Y, Zhao X, Zhou L,

Li X, Zou S, Zhang Y, Li X, Yang L, Guo J, Dong J, Li Q, Dong L, Zhu Y, Bai T, Wang

S, Hao P, Yang W, Zhang Y, Han J, Yu H, Li D, Gao GF, Wu G, Wang Y, Yuan Z, Shu

Y. 2013. Human infection with a novel avian-origin influenza A (H7N9) virus. N Engl J

Med **368:**1888-1897.

46.   **Chen H, Bright RA, Subbarao K, Smith C, Cox NJ, Katz JM, Matsuoka Y.** 2007.

Polygenic virulence factors involved in pathogenesis of 1997 Hong Kong H5N1

influenza viruses in mice. Virus Res **128:**159-163.

47.   **Shinya K, Hamm S, Hatta M, Ito H, Ito T, Kawaoka Y.** 2004. PB2 amino acid at

position 627 affects replicative efficiency, but not cell tropism, of Hong Kong H5N1

influenza A viruses in mice. Virology **320:**258-266.

48.   **Mehle A, Doudna JA.** 2008. An inhibitory activity in human cells restricts the function

of an avian-like influenza virus polymerase. Cell Host Microbe **4:**111-122.

49.   **Shi J, Deng G, Kong H, Gu C, Ma S, Yin X, Zeng X, Cui P, Chen Y, Yang H, Wan X,**

Wang X, Liu L, Chen P, Jiang Y, Liu J, Guan Y, Suzuki Y, Li M, Qu Z, Guan L, Zang

J, Gu W, Han S, Song Y, Hu Y, Wang Z, Gu L, Yang W, Liang L, Bao H, Tian G, Li

Y, Qiao C, Jiang L, Li C, Bu Z, Chen H. 2017. H7N9 virulent mutants detected in

chickens in China pose an increased threat to humans. Cell Res **27:**1409-1421.

50.   **Hsu PD, Scott DA, Weinstein JA, Ran FA, Konermann S, Agarwala V, Li YQ, Fine**

EJ, Wu XB, Shalem O, Cradick TJ, Marraffini LA, Bao G, Zhang F. 2013. DNA

targeting specificity of RNA-guided Cas9 nucleases. Nat Biotechnol **31:**827-+.

51.   **Jinek M, East A, Cheng A, Lin S, Ma EB, Doudna J.** 2013. RNA-programmed

genome editing in human cells. Elife **2**: e00471.

52.   **Pleschka S, Jaskunas R, Engelhardt OG, Zurcher T, Palese P, Garcia-Sastre A.**

1996. A plasmid-based reverse genetics system for influenza A virus. J Virol **70:**4188-

4192.

53.   **Kawakami E, Watanabe T, Fujii K, Goto H, Watanabe S, Noda T, Kawaoka Y.** 2011.

Strand-specific real-time RT-PCR for distinguishing influenza vRNA, cRNA, and mRNA.

J Virol Methods **173:**1-6.

54.   **Neumann G, Watanabe T, Ito H, Watanabe S, Goto H, Gao P, Hughes M, Perez DR,**

Donis R, Hoffmann E, Hobom G, Kawaoka Y. 1999. Generation of influenza A viruses

entirely from cloned cDNAs. Proc Natl Acad Sci U S A **96:**9345-9350.

55.   **Muench H, Reed L.** 1938. A simple method of estimating fifty percent endpoints. 27(3):

493-497.

56.   **Zhu W, Zhu Y, Qin K, Yu Z, Gao R, Yu H, Zhou J, Shu Y.** 2012. Mutations in

polymerase genes enhanced the virulence of 2009 pandemic H1N1 influenza virus in

mice. PLoS One **7:**e33383.

FIG 1. HuANP32A&B are indispensable for influenza A virus polymerase activity and viral replication. (A) Wild type 293T cells and single gene knockout 293T cell lines, including BUB3, CLTC, CYC1, NIBP, ZC3H15, C14orf173, CTNNB1, ANP32A, ANP32B, SUPT5H, HTATSF1, and DDX17, were transfected with firefly minigenome reporter, *Renilla* expression control, and the polymerase of H1N1$_{SC09}$. Luciferase activity was measured 24 h after transfection and data are firefly Luciferase gene activity normalized to *Renilla* Luciferase gene activity. (Statistical differences between samples are indicated, according to a one-way ANOVA followed by a Dunnett's test; NS= not significant, *P < 0.05, **P < 0.01, ****P < 0.0001. Error bars represent the SEM within one representative experiment.). **(B to D)** 293T, huANP32A knockout cells (AKO), huANP32B knockout cells (BKO), and huANP32A&B double knockout cells (DKO), were transfected with firefly minigenome reporter, *Renilla* expression control, and either H1N1$_{SC09}$ polymerase (**B**), H7N9$_{AH13}$ polymerase (**C**), or WSN polymerase (**D**). Luciferase activity was measured at 24 h post transfection. (**B** to **D,** data are firefly Luciferase gene activity normalized to that of *Renilla*, Statistical differences between cells are indicated, following a one-way ANOVA and subsequent Dunnett's test; NS = not significant, **P < 0.01, ***P < 0.001, ****P < 0.0001. Error bars represent the SEM of the replicates within one representative experiment.). The expression levels of H1N1$_{SC09}$ polymerase proteins were assessed by western blotting **(B)**. **(E)** 293T, AKO, BKO and DKO cells were infected with WSN virus at an MOI of 0.01. The supernatants were sampled at 0, 12, 24, 36, and 48 h post infection and the virus titers were

determined by means of endpoint titration in MDCK cells. Error bars indicate SD from

three independent experiments. *P < 0.05, **P < 0.01, ***P < 0.001.

FIG 2. Reconstitutions of huANP32A&B rescue polymerase activity and

replication of influenza A virus. 20 ng huANP32A, 20 ng huANP32B,10 ng

huANP32A and 10 ng huANP32B, or 20 ng empty vector were co-transfected with

either H1N1$_{SC09}$ polymerase (**A**), H7N9$_{AH13}$ polymerase (**B**) or WSN polymerase (**C**)

into DKO cells, luciferase activity was assayed at 24 h after transfection. (**D**) Increasing

doses of huANP32A or huANP32B were co-transfected with H1N1$_{SC09}$ polymerase into

DKO cells. Luciferase activity was measured at 24 h post transfection. The expression

of ANP32 proteins was assessed by western blotting. (**A** to **D,** data are firefly Luciferase

gene activity normalized to that of *Renilla*, Statistical differences between cells are

indicated, following a one-way ANOVA and subsequent Dunnett's test; NS = not

significant, **P < 0.01, ***P < 0.001, ****P < 0.0001. Error bars represent the SEM of

the replicates within one representative experiment). (**E**) DKO cells were transfected

with 1ug huANP32A or/and B, or empty vector. After 24 h, the cells were infected with

WSN virus at an MOI of 0.01. The supernatants were sampled at 0, 12, 24, 36, and 48

h post infection and the virus titers in these supernatants were determined as above.

Error bars indicate SD from three independent experiments. NS = not significant, *P <

0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

FIG 3. HuANP32A&B determine viral RNA replication efficiency and viral

production. (A-C) 293T, or DKO cells, were transfected with the H1N1$_{SC09}$

minigenome reporting system together with either 20 ng empty vector or huANP32A.

The cells were incubated at 37 °C for 24 h before reverse transcription followed by

quantitative PCR (qRT–PCR) for vRNA, mRNA and cRNA of the Luciferase gene. Error

bars represent the SD from three independent experiments. NS = not significant,

****$P < 0.0001$. **(D** and **E)** Replication kinetics of H1N1$_{SC09}$ in ANP32 knockout cells.

293T, AKO, BKO, and DKO cells in 6-well plates were transfected with 0.5 μg each of

the eight plasmids of H1N1$_{SC09}$ (**D**), or, in DKO cells, 40 ng huANP32A, 40ng

huANP32B, 20 ng each of huANP32A and huANP32B, or 40 ng empty vector were co-

transfected with the eight-plasmids of H1N1$_{SC09}$ (**E**). The cells were cultured at 37 °C

and supernatants collected at 0, 12, 24, 48 and 60 h post-transfection and subjected

to virus production assay by ELISA, as described in 'Methods'. Error bars represent

the SD from three independent experiments. Error bars indicate SD from three

independent experiments. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$.

FIG 4. Support of influenza A viral replication by ANP32A or B from different

species. Twenty nanograms of either empty vector or ANP32A or B from one of several

different species was co-transfected with a minigenome reporter, *Renilla* expression

control, and human influenza virus polymerase from H1N1$_{SC09}$ (**A**), H7N9$_{AH13}$ (**B**),

canine influenza virus H3N2$_{GD11}$ (**C**), equine influenza virus H3N8$_{XJ07}$ (**D**) and H3N8$_{JL89}$

(**E**), or avian influenza virus H9N2$_{ZJ12}$ (**F**) into DKO cells. Luciferase activity was

measured 24 h later. (Data are firefly Luciferase gene activity normalized to that of

Renilla, Statistical differences between cells are indicated, following a one-way ANOVA

and subsequent Dunnett's test; NS = not significant, *P < 0.05, **P < 0.01, ***P < 0.001,

****P < 0.0001. Error bars represent the SEM of the replicates within one

representative experiment.) dk, duck; ty, turkey; zf, zebra finch; mu, mouse; pg, pig;

eq, equine.

FIG 5. Key amino acids in ANP32A&B determine the activity of influenza viral

polymerases. (A) Amino acid sequence comparison for chicken, human and pig

ANP32B proteins (amino acids 110–161). Amino acid sequences are available in

GenBank. The black dots indicate the different amino acid within these proteins. **(B** to

D) huANP32B mutants were co-transfected with polymerase plasmids from H1N1$_{SC09}$

(**B** and **C**), or H7N9$_{AH13}$ (**D**), plus minigenome reporter and *Renilla* luciferase control

into DKO cells. Luciferase activity was then assayed 24 h after transfection. **(E** and **F)**

huANP32A mutants were co-transfected with polymerase plasmids from H1N1$_{SC09}$ (**E**)

or H7N9$_{AH13}$ (**F**), plus a minigenome reporter and *Renilla* luciferase control, into DKO

cells. **(G** and **H)** chANP32A or B mutants were co-transfected with polymerase

plasmids from H7N9$_{AH13}$ with either PB2 627K (**G**), or 627E (**H**) respectively. Luciferase

activity was analyzed as mentioned above. (Data are firefly Luciferase gene activity

normalized to that of *Renilla*, Statistical differences between cells are given, following

one-way ANOVA and subsequent Dunnett's test; NS = not significant, *P < 0.05,

**P < 0.01, ***P < 0.001, ****P < 0.0001. Error bars represent the SEM of the replicates

within one representative experiment.)

FIG 6. Amino-acids N129I/D130N substitute into ANP32A&B impaired influenza

polymerase activity in 293T cells. (A) The endogenous protein expressions of

different cell lines were identified by Western blotting using antibodies against β-actin,

huANP32A and huANP32B. (B and C) Selected 293T cell lines were transfected with

firefly minigenome reporter, *Renilla* expression control, and polymerase from $H1N1_{SC09}$

(B) or $H7N9_{AH13}$ (C). Cells were assayed for Luciferase activity. WT, wild-type 293T

cells; A5#, ANP32A mutant cells; B32#, ANP32B mutant cells; AB93#,

ANP32A/ANP32B double mutant cells. (Data are firefly Luciferase gene activity

normalized to that of *Renilla*, Statistical differences between cells are given, following

a one-way ANOVA and subsequent Dunnett's test; NS = not significant, *P < 0.05,

**P < 0.01, ***P < 0.001, ****P < 0.0001. Error bars represent the SEM of the replicates

within one representative experiment.)

FIG 7. The 129-130 site in ANP32B contributes to the interaction with influenza

polymerase. (A) A schematic describing the structure of ANP32B. LRR, leucine-rich repeat,

including 4 LRR repeats and one C-terminal LRR domain (LRRCT); LCAR, low-complexity

acidic region; NLS, nuclear localization signal. (B) and (C) 293T cells were transfected with

different ANP32-Flag constructs, together with viral polymerase subunits PA, PB1, and

PB2. Coimmunoprecipitation of the anti-Flag antibodies and the proteins was assessed

using western blotting.

SUPPORTING INFORMATION

FIG S1. The identification of huANP32A and huANP32B knockout cells. 293T cells

were transfected with pMJ920 vector (plasmid expressing eGFP and Cas9) and

gRNAs targeting huANP32A or huANP32B to generate huANP32A knockout cells

(AKO), huANP32B knockout cells (BKO), and huANP32A&B double knockout cells

(DKO). The mRNA levels of huANP32A and B in different cell lines were determined

using qRT-PCR using primers specific for huANP32A (**A**) and huANP32B (**B**). The

primer sequences are given in 'Materials and Methods'. Fold change in RNA was

calculated by double-standard curve methods. β-actin was used as an internal control.

Statistical differences between samples are given, following one-way ANOVA and

subsequent Dunnett's test; NS = not significant, ****$P < 0.0001$. Error bars indicate SD

from three independent experiments. (**C**) The endogenous proteins were identified by

western blotting using antibodies against β-actin, huANP32A and huANP32B,

described as "Materials and Methods".

FIG S2. Overdose expression of huANP32A&B impaired viral polymerase activity.

Different doses of huANP32A or huANP32B were co-transfected with H1N1$_{SC09}$

polymerase (**A**) and H7N9$_{AH13}$ polymerase (**B**) into DKO cells. Luciferase activity was

measured 24 h after transfection and data are firefly Luciferase gene activity

normalized to that of *Renilla*. (Statistical differences between samples are given,

following one-way ANOVA and subsequent Dunnett's test; NS = not significant, *P <

0.05, ***P < 0.001, ****P < 0.0001. Error bars represent the SEM of the replicates within

one representative experiment.). The expression levels of the huANP32A and

huANP32B proteins were confirmed by western blotting.

FIG S3. Phylogenetic tree of amino acid sequences of ANP32A (A) and ANP32B

(B). Trees were generated by the neighbor-joining method and evaluated by 1000

bootstrap pseudo replicates. The bar scale represents the genetic distance. The

credibility value for each node is shown.

FIG S4. Expression of flag-tagged ANP32A or B proteins and PB2 E627K

mutation overcomes human ANP32A&B restriction. (A) 1μg of each ANP32

plasmid was transfected into 293T cells using Lipofectamine® 2000. 48 h post

transfection, the cell lysates were analyzed using SDS-PAGE and Western Blotting

using antibodies against Flag peptide and β-actin. (B-D) DKO cells were transfected

with 20ng either empty vector or ANP32A or B from one of several different species,

together with minigenome reporter, *Renilla* expression control, and different viral

polymerase including: human H7N9$_{AH13}$ with PB2 K627E (**B**), an avian origin equine

influenza virus H3N8$_{XJ07}$ with PB2 E627K (**C**), and avian influenza virus H9N2$_{ZJ12}$ with

PB2 E627K (**D**). Luciferase activity was measured 24 h post transfection. (Data are

polymerase activity normalized to *Renilla*; Statistical differences between cells are

given, following one-way ANOVA and subsequent Dunnett's test; NS = not significant,

*P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001. Error bars represent the SEM of the

replicates within one representative experiment.) dk, duck; ty, turkey; zf, zebra finch;

mu, mouse; pg, pig; eq, equine.

FIG S5. Mapping of crucial sites in ANP32 proteins that influence viral

polymerase activity. (A) ANP32B sequences from chicken, human, and pig were

aligned using the Geneious R6 software. The notches are marked with dashes. The

similarity of amino acid identity is highlighted in different colors. (B) Schematic diagram

of chimeric clones between chicken and human ANP32B, constructed according to the

known domains (Separated by dotted lines.1-41aa, LRR1 region; 42-110aa, LRR2,3

&4 region; 111-161aa, LRRCT region; 162-251aa, LCAR region, as showed in a). The

bars show the origin of the genes as follows: white, huANP32B; blue, chANP32B. (56)

Chimeric clones were co-transfected with minigenome reporter, *Renilla* expression

control, and H1N1$_{SC09}$ polymerase into DKO cells. Luciferase activity measured 24 h

later. (Data are polymerase activity normalized to *Renilla*; Statistical difference

between samples are given labeled, following one-way ANOVA and subsequent

Dunnett's test; NS = not significant, ****$P < 0.0001$. Error bars represent the SEM of

the replicates within one representative experiment.)

FIG S6. Expression of flag-tagged ANP32A or B mutants.

1μg of each mutant ANP32 and wildtype ANP32 plasmids were transfected in 293T

cells by Lipofectamine® 2000. 48 h post transfection, the cells lysates were analyzed

by SDS-PAGE and Western Blotting using antibodies against Flag peptide and β-actin.

FIG S7. Mouse ANP32A 130A impaired its support of viral replication. Mutated

muANP32A or human ANP32 proteins were co-transfected with minigenome reporter,

Renilla expression control, and H1N1$_{SC09}$ polymerase into DKO cells. Luciferase

activity was assayed. (Data are polymerase activity normalized to *Renilla*; Statistical

differences between cells are given, following one-way ANOVA and subsequent

Dunnett's test; NS = not significant, *$P < 0.05$, ****$P < 0.0001$. Error bars represent the

SEM of the replicates within one representative experiment.)

FIG S8. Identification of ANP32A&B site-mutation 293T cell lines. N129I/D130N

substitutions in huANP32A and ANP32B on the 293T chromosome were generated

using a CRISPR/Cas9 system. Positive 293T colonies of N129I/D130N substitutions

on huANP32A and ANP32B were identified by sequencing. (A) Cloned cell A5# was

identified carrying the N129I/D130N on ANP32A but not ANP32B. (B) Cell B32# was

identified harboring N129I/D130N on ANP32B but not ANP32A. (C) Cell AB93# has

N129I/D130N on both ANP32A&B.

A

B

C

Table S1. Summary of amino acids at 129-130 of ANP32 proteins

| | Residue at amino acid position* | | | |
| | ANP32A | | ANP32B | |
Host	129	130	129	130
Gallus	N	D	I	N
Anas platyrhynchos	N	D	I	N
Homo sapiens	N	D	N	D
Gorilla gorilla	N	D	N	D
Felis catus	N	D	N	D
Canis lupus familiaris	N	D	N	D
Bos taurus	N	D	N	D
Sus scrofa	N	D	N	D
Mus musculus	N	A	S	D
Equus caballus	N	D	N	D

931 *The sequences of different species ANP32 proteins were retrieved from NCBI