Redefining the medicago sativa alphapartitivirus genome sequences

Short communication

Nicolás Bejerman[1,2*], Debat Humberto[1,2*], Verónica Trucco[1,2], Soledad de Breuil[1,2], Sergio Lenardon[1,2,3], Fabián Giolitti[1,2]

[*]both authors contributed equally to this work

[1] UFYMA-INTA-CONICET, Av. 11 de Septiembre 4755, Córdoba, 5020, Argentina

[2] IPAVE-CIAP-INTA, Av. 11 de Septiembre 4755, Córdoba, 5020, Argentina

[3] Facultad de Agronomía y Veterinaria (FAV), Universidad Nacional de Río Cuarto (UNRC), Enlaces Rutas 8 y 36, 5800, Río Cuarto, Córdoba, Argentina

Corresponding author: Nicolás Bejerman, nicobejerman@gmail.com, Tel: +54 9 351 4973636, Fax: +54 9 351 4974330

Abstract

In alfalfa samples analyzed by high-throughput sequencing, four *de novo* assembled contigs encoding

gene products showing identities to alphapartitiviruses proteins were found based on BLASTX analysis.

The predicted amino acid (aa) sequences of two contigs presented 99-100% identity to the RNA-

dependent RNA polymerase (RdRP) and the capsid protein (CP) of the recently reported medicago sativa

alphapartitivirus 1 (MsAPV1). In addition, the remaining contigs shared only 56% (CP) and 70% (RdRP)

pairwise aa identity with the proteins of MsAPV1, suggesting that the samples presented also a novel

Alphapartitivirus species. Further analyses based on complete genome segments termini and the

presence/absence of alphapartitivirus RNA in several samples and public alfalfa RNA datasets,

corroborated the identification of two different alphapartitviruses members. Results also likely indicate

that the reported MsAPV1 genome was previously reconstructed with genome segments of two different

alphapartitiviruses. Overall, we not only revisited the MsAPV1 genome sequence but also report a new

tentative alphapartitivirus species, which we propose the name medicago sativa alphapartitivirus 2. In

addition, the RT-PCR detection of both MsAPV1 and MsAPV2 in several alfalfa cultivars suggests a

broad distribution of both viruses.

Keywords: alphapartitivirus, medicago sativa, complete genomes, virus discovery

Short communication

In Argentina, alfalfa (*Medicago sativa* L.) is a primary forage crop and a major feed component in dairy and beef cattle production systems. In 2011, we observed alfalfa plants showing symptoms of shortened internodes (bushy appearance), leaf puckering and varying-sized vein enations on abaxial leaf surfaces (Bejerman et al., 2011). Deep sequencing of alfalfa plants collected in central region of Argentina showing the dwarfism symptoms, revealed the presence of four RNA viruses: alfalfa mosaic virus (AMV), alfalfa dwarf virus (ADV), alfalfa enamovirus-1 (AEV-1) and bean leaf roll virus (BLRV) (Bejerman et al., 2015; 2016; Trucco et al., 2014, 2016) and one DNA virus: alfalfa leaf curl virus (ALCV) (Bejerman et al., 2018). Furthermore, four assembled sequences (contigs) analyzed by BlastX searches shared significant identity (E-value = 0), to the capsid and replication proteins encoded by alphapartitiviruses.

Partitiviruses particles are isometric and have a double stranded RNA (dsRNA) bi-segmented genome; each dsRNA contains a single open reading frame (ORF), which encodes a RNA dependent RNA polymerase (RdRP) or a coat protein (CP) (Nibert et al., 2014). The *Partitiviridae* family has been taxonomically reorganized recently; it comprises five genera, *Alphapartitivirus*, *Betapartitivirus*, *Gammapartitivirus*, *Deltapartitivirus,* and *Cryspovirus* (Nibert et al., 2014). Partitiviruses corresponding to the same genus are usually reported as single infections; however partitiviruses co-infecting a single host have also been described (Lesker et al., 2013; Lyu et al., 2018; Ong et al., 2017; Sabanadzovic and Valverde, 2011). In the latter case, it is a challenge to pair the two genomic segments to identify and annotate the genomes without mixing up genomic segments from different partitiviruses (Ong et al., 2017). Alphapartitiviruses infect either plant or fungi. Plant-infecting alpahapartitiviruses are associated with latent infections of their hosts and are transmitted with a high frequency via the ovule and by pollen to the seed embryo, but cannot be transmitted by grafting or mechanical inoculation (Nibert et al., 2014). A new alphapartitivirus member infecting alfalfa, named as medicago sativa alphapartitvirus 1 (MsAPV1), was reported by analyzing public transcriptome dataset (Kim et al., 2018).

In this work we report a new alphapartitivirus member named as medicago sativa alphartitivirus 2 (MsAPV2) and redefined the reported MsAPV1 genome sequence. Furthermore, we detected both viruses in mixed infections in several alfalfa cultivars samples and public alfalfa RNA datasets suggesting a high prevalence of both viruses.

In 2011, symptomatic alfalfa plants showing dwarfism were collected from a field located in Manfredi

(central region of Argentina), and total RNA was extracted using TRIzol Reagent (Life Technologies)

according to manufacturer's instructions. This total RNA was sent to Fasteris Life Sciences SA (Plan-les-

Ouates, Switzerland), where the bands located between 21 and 30 bp were excised, purified, processed

and sequenced on an Illumina HiSeq 2000. The obtained raw reads were filtered by quality and trimmed

using the FASTX-Toolkit as implemented in http://hannonlab.cshl.edu/fastx_toolkit/index.html. The

remaining short reads were de novo assembled using Velvet 1.2.10 with the velvet optimiser parameter

available at https://www.ebi.ac.uk/~zerbino/velvet/. The obtained contigs were subjected to bulk BlastX

searches (E-value < 1e-5) against a local database of the non-redundant virus proteins. Four contigs

ranging from 1,650-1,840 nt and sharing high similarity (56-100%) with alphapartitivirus proteins were

retrieved, further inspected and annotated as described in Debat and Bejerman (2018). The contigs were

curated by mapping the filtered sequencing reads using the Geneious 8.1.9 (Biomatters, inc) mapper with

low sensitivity parameters and assessing the consensus by hand. The curated contigs were initially

designated as alphaA1 (1,872 nt, mean coverage 31.6X, supported by 2,782 reads, encoding an RdRP);

alphaA2 (1,687 nt, mean cov. 11.5X, supported by 876 reads, encoding a CP); alphaB1 (1,845 nt, mean

coverage 38.9X, supported by 3,357 reads, encoding an RdRP); and alphaB2 (1,714 nt, mean coverage

28.5X, supported by 2,314 reads, encoding a CP). AlphaA1 and alphaA2 shared 98.7-99.8 nt identity with

the reported MsAPV1 dsRNA1 (MF443256) and dsRNA2 (MF443257), respectively. Thus, we

tentatively grouped the identified contigs into two putative viruses corresponding to MsAPV1

(alphaA1/alphaA2) and MsAPV2 (alphaB1/alphaB2). However, evidence described below based on the

complete genome sequences and RNA levels indicated that the genome segments could be paired

differently. So, alphaA2 (almost identical to the reported MsAPV1 dsRNA2 (MF443257)) is in fact the

RNA2 of MsAPV2, and thus alphaB2 should be paired with alphaA1.

To obtain the complete sequences of these putative alphapartitivirus genome segments, their terminal

sequences were amplified by 5´and 3´rapid amplification of cDNA ends (RACE) that was carried out as

described previously by Phelan and James (2016) which resulted in assembly of full genome sequences

that were deposited in the GenBank database under accession numbers MK292286 (MsAPV1 dsRNA1),

MK292287 (MsAPV1 dsRNA2), MK292288 (MsAPV2 dsRNA1) and MK292286 (MsAPV2 dsRNA2).

Primers used to amplify the terminal regions are listed is Supplementary Table 1.

The complete sequences of the revisited MsAPV1 dsRNA1 (extended from the formerly designated

alphaA1 contig) and dsRNA2 (formerly designated alphaB2 contig) were determined to be 1942- and

1806- nt long, respectively (Fig. 1A) with GC contents of 38.62% and 43.79%, respectively; whereas the

complete sequences of the novel MsAPV2 dsRNA1 (alphaB1) and dsRNA2 (alphaA2) were determined

to be 1939- and 1764- nt long, respectively (Fig. 1A) with GC contents of 43.73% and 44.72%,

respectively. GC content is likely similar to those values reported for other partitiviruses (Phelan and

James, 2016). Each dsRNA contains a single open reading frame (ORF) with 5´ and 3´untranslated

regions (UTRs). Interestingly, multiple alignments of 5´UTRs revealed that these sequences are highly

conserved, being the first 9 nt at the 5´terminus of the dsRNA1 and dsRNA2 fragments of MsAPV1

identical, whereas the first 10nt at the 5´terminus of the dsRNA1 and dsRNA2 fragments of MsAPV2

also are identical, but different from the 5' termini of the MsAPV1 segments (Fig. 1B). Furthermore, the

5´UTR of MsAPV1 and MsAPV2 dsRNA1 and 2 were predicted to fold into stem-loop structures, when

examined using Mfold software (http://mfold.rna.albany.edu) (Fig. 1C). Similar stem-loop structures,

which are likely to play an important role in dsRNA replication and virion assembly, have been described

in other partitiviruses (Guo et al., 2017; Lesker et al., 2013; Osaki and Sasaki, 2018).

Sequence analysis of MsAPV1 and MsAPV2 complete dsRNA1 revealed that an expected single open

reading frame (ORF) encoding a putative 586-amino acid (aa) RdRP protein, where an RdRP conserved

domain was found (cd01699, e-values =3.73e-05 and 3.40e-07). Moreover, the multiple-protein

alignment of the deduced aa sequences of the RdRPs of MsAPV1, MsAPV2 and selected

alphapartitiviruses confirmed that this protein includes the six conserved motifs (III to VIII) described for

partitiviruses (Liu et al., 2015). Sequence analysis of MsAPV1 and 2 dsRNA2 revealed that it contained

a single ORF encoding a putative 499- and 491-aa CP protein, respectively.

Pairwise comparison using Sequence Demarcation Tool (SDT v1.2) (Muhire et al., 2014) showed that at

amino acid level the RdRP of alphapartitivirus 1 is 100% identical to that one encoded by the former

MsAPV1 (Fig. 2A), whereas its CP is just 56% identical to that one of the former MsAPV1 (Fig. 2B). On

the other hand, the RdRP of the alphapartitivirus 2 is just 70% identical to that one encoded by

alphapartitivirus 1 and the former MsAPV1 (Fig. 2A), whereas its CP is 99% identical to that one of

alphapartitivirus 1 and the former MsAPV1 (Fig. 2B), which likely suggest that MsAPV1 genome

segments could be paired differently that previously reported. Both alphapartitivirus 1 and 2 have a

degree of identity below 90% and 80% for RdRp and CP when compared with other partitiviruses,

respectively, which is the species demarcation threshold recommended recently for alphapartitiviruses

(Vainio et al., 2018).

To further assess our speculation that the reported MsAPV1 segments derived from two different viruses

we returned to the raw data which was the source of Kim et al (2018) study, which pooled eight libraries

before de novo assembly. In this direction, raw reads corresponding to RNAseq NGS libraries (SRA:

SRP017117), associated to NCBI Bioproject PRJNA179114 and additionally from an independent study

oriented to elucidate the molecular genetic basis of herbivory between butterflies and their host plants,

including alfalfa (Nallu et al 2018, PRJNA437291, SRA: SRP134094), were analyzed according to Debat

and Bejerman (2018). Briefly, the respective public datasets were downloaded and reads of each library

including our NGS data were mapped against the assembled virus genomes sequences using Bowtie2

v2.3.4.3 available at http://bowtie-bio.sourceforge.net/bowtie2/ with default settings and the fast end-to-

end preset. The obtained values were normalized as mapped reads per million (RPM) for each library.

Interestingly, we observed a differential pattern of presence and absence of the diverse alphapartitivirus

segments in each library (Table 1). When we analyzed in detail the transcriptome data used by Kim et al.,

(2018) we observed that two of the eight pooled libraries presented virus RNA derived from both

segments of the revisited MsAPV1, and only one library harbored virus reads of segment 2 of MsAPV2

(which was previously annotated as segment 2 of MsAPV1). Thus, we believe that the use of a pooled

transcriptome dataset for virus discovery could misrepresent the virus landscape associated to each

specific sample that led to a miss assignment of alphapartitiviruses genome segments. Moreover, when

we explored 18 alfalfa RNAseq datasets from bioproject PRJNA437291 we observed that in three

libraries only virus RNA from both genome segments of the redefined MsAPV1 were found, while in five

additional runs we detected virus RNA from both segments of MsAPV1 and MsAPV2, suggesting mixed

infections of alphapartitiviruses in those samples (Table 1). The detection of only both segments of the

redefined MsAPV1 in several samples likely indicates that the former MsAPV1 genome could be

redefined.

Following sequence annotation and the assessment of RNA levels, the redefined MsAPV1 and 2

predicted RdRp and CP aa sequences were aligned with the corresponding sequences of selected

partitiviruses, described in Supplementary Table 2, using MUSCLE (Edgar, 2004) as implemented in

Mega X (Kumar et al., 2018). ML phylogenetic trees were inferred using Mega X (Kumar et al., 2018)

with LG + G and LG + G + F amino acid substitution models, respectively, and 1000 bootstrap replicates.

Phylogenetic analyses supported the pairwise comparison results showing that MsAPV1 RdRP grouped

within alphapartitiviruses with the former MsAPV1 whereas MsAPV2 clustered with pyrus pyrifolia

partitivirus 2 (PpPV2) and rose partitivirus (RoPV) (Fig. 2C). On the other hand, MsAPV1 CP formed a

monophyletic cluster within alphapartitiviruses whereas MsAPV2 grouped with the former MsAPV1,

PpPV2 and RoPV (Fig. 2D). This suggest that MsAPV1 and MsAPV2 do not share a common immediate

ancestor, so it is tempting to speculate that they originated from two distinct ancestral partitiviruses

adapted to the same host.

Recently, it was reported the presence of MsAPV1 in alfalfas from USA (Nemchinov et al., 2018). Based

on our results it is tempting to suggest that these alfalfas could be infected with both MsAPV1 and

MsAPV2 given that the presence of alphapartitivirus RNA sequences in that study were assessed by

RNAseq reads mapping using as reference the published sequences by Kim et al., (2018).

In sum, it was reported that there is higher sequence identity between the corresponding 5' UTRs of CP

and RdRP segments belonging to the same alphapartitivirus than those belonging to different species

(Ong et al., 2017). Therefore we verified if genomes were consistently annotated in this study by

comparing the 5' UTRs of segments of MsAPV1 and MsAPV2, which was only possible by generation of

complete genome segments using RACE. The comparison revealed a 70.5% and 70.1% identity between

the 5' UTRs of CP and replicase segments of MsAPV1 and MsAPV2, respectively. Whereas identities

values when 5' UTRs of CP and replicase segments of MsAPV1 were compared against those of

MsAPV2 were lower than 60%. This supports our tentative pairing of the revisited MsAPV1 and

MsAPV2 genomes in this study. In addition, the bioproject previously analyzed by Kim et al., (2018) as

well as other alfalfa datasets were analyzed showing the presence of both MsAPV1 and MsAPV2 which

further supports our hypothesis. Plant-infecting partitiviruses are efficiently pollen/seed transmitted, but

are cryptic, so infected- plants are symptomless (Nibert et al., 2014), therefore they are not considered as

quarantine pest and they are not tested when alfalfa seeds are traded. Consequently, both MsAPV1 and

MsAPV2 could have been transmitted to many alfalfa cultivars inadvertently, resulting in their

dissemination all around the world through the exchange of alfalfa seeds, which led to its detection in

diverse contexts. In this regard, in order to assess a tentative prevalence landscape of alphapartitivirus in

alfalfa, we studied the occurrence of MsAPV1 and MsAPV2 by RT-PCR assays of several alfalfa

cultivars, using two set of virus-specific primers designed from dsRNA1 (Supplementary Table 1) in

order to detect MsAPV1 and MsAPV2 in alfalfa leafs. Interestingly, we detected both dsRNA1 sequences

in all samples tested, suggesting that the presence of both viruses as mixed infections is common in

alfalfa (Supp. Fig. 1).

In conclusion, we describe the complete genome sequences of two alphapartitivirus infecting alfalfa,

which we dubbed as medicago sativa alphapartitivirus 1 and medicago sativa alphapartitivirus 2. These

viruses were found in mixed and single infections in alfalfa samples from diverse regions of the world

and should be considered as new members of the genus *Alphapartitivirus*. We also implemented here a

simple tool based on RT-PCR detection to screen the presence of both viruses in alfalfa cultivars. Our

study provides new avenues to explore and confirm virus genome segments when partitiviruses (or any

multi-segmented viruses) are co-infecting a same sample, which appears to be genuinely common.

Compliance with ethical standards

Conflict of interest: All authors declare that they have no conflict of interest

Ethical Approval: This article does not contain any studies with human participants or animals performed

by any of the authors.

Funding: This work was partially supported by the PNPV 1135022 project of INTA.

References

Bejerman, N., Nome, C., Giolitti, F., Kitajima, E., de Breuil, S., et al. 2011. First report of a Rhabdovirus

infecting alfalfa in Argentina. Plant Dis 95, 771.

Bejerman, N., Giolitti, F., deBreuil, S., Trucco, V., Nome, C., Lenardon, S., Dietzgen, R.G. 2015.

Complete genome sequence and integrated protein localization and interaction map for alfalfa dwarf

virus, which combines properties of both cytoplasmic and nuclear plant rhabdoviruses. Virol 483,

275–283.

Bejerman, N., Giolitti, F., Trucco, V., de Breuil, S., Dietzgen, R., Lenardon,, S. 2016. Complete genome

sequence of a new enamovirus from Argentina infecting alfalfa plants showing dwarfism symptoms.

Arch Virol 161, 2029-2032.

Bejerman, N., Trucco, V., de Breuil, S., Rodriguez-Pardina, P., Lenardon, S., Giolitti, F. 2018. Genome

characterization of an Argentinean isolate of alfalfa leaf curl virus. Arch Virol 163, 799-803.

Debat, H., Bejerman., N. 2018. Novel bird's-foot trefoil RNA viruses provide insights into a clade of

legume associated enamoviruses. bioRxiv. doi: https://doi.org/10.1101/368944.

Edgar, R.C. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput.

Nucleic Acids Res 32, 1792–1797.

Guo, M., Bian, Y., Wang, J., Wang, G., Ma, X., Xu, Z. 2017. Biological and molecular characteristics of

a novel partitivirus infecting the edible fungus Lentinula edodes. Plant Dis. 101, 726–733.

Kim, H., Park, D., Hahn, Y. 2018. Identification of novel RNA viruses in alfalfa (Medicago sativa): an

Alphapartitivirus, a Deltapartitivirus, and a Marafivirus. Gene 638, 7-12.

Kumar, S., Stecher, G., Li, M., Knyaz, C., Tamura, K. 2018. MEGA X: Molecular evolutionary genetics

analysis across computing platforms. Mol. Biol. Evol. 35, 1547–1549.

Lesker, T., Rabenstein, F., Maiss, E. 2013. Molecular characterization of five betacryptoviruses infecting

four clover species and dill. Arch. Virol. 158, 1943–1952.

Liu, L., Wang, Q., Cheng, J., Fu, Y., Jiang, D., Xie, J. 2015. Molecular characterization of a bipartite

double-stranded RNA virus and its satellite-like RNA co-infecting the phytopathogenic fungus

Sclerotinia sclerotiorum. Front. Microbiol. 6, 406.

Lyu, R., Yi, Z., Tang, Q., Li, Y., Cheng, J., Fu, Y., et al. 2018. Two alphapartitiviruses co-infecting a

single isolate of the plant pathogenic fungus Rhizoctonia solani. Arch. Virol. 163, 515–520.

Muhire, B.M., Varsani, A., Martin, D.P. 2014. SDT: a virus classification tool based on pairwise

sequence alignment and identity calculation. PLoS One 9, e108277.

Nallu, S., Hill, J. A., Don, K., Sahagun, C., Zhang, W., Meslin, C., et al. 2018. The molecular genetic

basis of herbivory between butterflies and their host plants. Nat Ecol Evol. 2, 1418-1427.

Nemchinov, L.G., Lee, M.G., Shao, J. 2018. First Report of Alphapartitiviruses Infecting Alfalfa

(Medicago sativa L.) in the United States. Microbiol Resour Announc. 7, e01266-18.

Nibert, M. L., Said, A., Ghabrial, S. A., Maiss, E., Lesker, T., et al. 2014. Taxonomic reorganization of

family *Partitiviridae* and other recent progress in partitivirus research. Virus Res. 188, 128–141.

Ong, J.W.L., Li, H., Sivasithamparam, K., Dixon, K.W., Jones, M.G.K., Wylie, S.J. 2017 The challenges

of using high-throughput sequencing to track multiple bipartite mycoviruses of wild orchid-fungus

partnerships over consecutive years. Virol. 510, 297–304.

Osaki, H., Sasaki, A. 2018. A novel alphapartitivirus detected in Japanese pear. Virus Genes 54:149-154.

Phelan, J., James, D. 2016. Complete genome sequences of a putative new alphapartitivirus detected in

Rosa spp. Arch Virol. 161, 2623-2626.

Sabanadzovic, S., Valverde, A.R. 2011. Properties and detection of two cryptoviruses from pepper

(*Capsicum annuum*). Virus Genes. 43, 307–312.

Trucco, V., de Breuil, S., Bejerman, N., Lenardon, S., Giolitti, F. 2014. Complete nucleotide sequence of

Alfalfa mosaic virus isolated from alfalfa (*Medicago sativa* L.) in Argentina. Virus Genes 48, 562-

565.

Trucco, V., de Breuil, S., Bejerman, N., Lenardon, S., Giolitti, F. 2016. Bean leafroll virus (BLRV) in

Argentina: molecular characterization and detection in alfalfa fields. Eur J Plant Pathol 146, 207-

212.

Vainio, E.J., Chiba, S., Ghabrial, S.A., Maiss, E., Roossinck, M., Sabanadzovic, S., Suzuki, N., Xie, J.,

Nibert, M. 2018. ICTV Report Consortium. ICTV virus taxonomy profile: Partitiviridae. J. Gen.

Virol. 99, 17–18.

Fig. 1. **A**. Genomic organization of medicago sativa alphapartitivirus 1 and 2. The numbers above the

solid lines indicate the nucleotide positions of the beginning and end of the 5'- and 3'-UTRs. **B**. Multiple

alignments between 5´UTR of dsRNA1 and 2 of MsAPV1 and MsAPV2. **C**. Potential secondary

structures formed by 5'-terminal sequences of positive strands of two segments of MsAPV1 and

MsAPV2.

Fig.2. **A and B**. Amino acid percentage pairwise identities between MsAPV1 and MsAPV2 RdRp (**A**)

and CP (**B**) with those from selected partitiviruses members calculated using SDT v1.2 (Muhire et al.,

2014). **C.** Maximum likelihood phylogenetic tree of MsAPV1 and 2 and selected partitiviruses members

based on the RdRp amino acid sequences. **D**. Maximum likelihood phylogenetic tree of MsAPV1 and 2

and selected partitiviruses members based on the CP amino acid sequences. All trees were constructed

using MegaX software. LG + G model was used to construct the phylogenetic tree based on RdRp amino

acid sequences; whereas model LG + G + F was used to construct the phylogenetic tree based on CP

amino acid sequences. Bootstrap values out of 1000 replicates are given at the nodes. The bar below each

tree represents substitutions per site. Accession numbers of every virus used to perform SDT analysis and

construct phylogenetic trees are listed in Supplementary Table 2.

Table 1. Virus RNA levels corresponding to each genome segment of MsAPV1 & 2 in multiple public
datasets. Values express virus derived reads per million of total reads (RPM) based on mapping to the
assembled genome sequences. An arbitrary (and conservative) cut-off of 5 RPM was considered to indicate

BioProject	Run	BioSample	Experiment	Virus RPM			
				MsAPV1 RNA1	MsAPV1 RNA2	MsAPV2 RNA1	MsAPV2 RNA2
PRJNA179114	SRR612164	SAMN01804647	SRX203401	3	1	0	0
PRJNA179114	SRR612163	SAMN01804646	SRX203400	0	0	0	0
PRJNA179114	SRR612116	SAMN01804612	SRX203369	498	99	0	0
PRJNA179114	SRR612115	SAMN01804610	SRX203368	0	0	0	0
PRJNA179114	SRR612114	SAMN01804609	SRX203367	0	0	0	0
PRJNA179114	SRR612113	SAMN01804577	SRX203355	0	0	0	0
PRJNA179114	SRR611885	SAMN01804552	SRX203330	0	0	0	16
PRJNA179114	SRR611809	SAMN01804510	SRX203284	0	0	0	0
PRJNA437291	SRR6814494	SAMN08647220	SRX3771495	0	0	0	0
PRJNA437291	SRR6814495	SAMN08647220	SRX3771494	982	122	187	56
PRJNA437291	SRR6814496	SAMN08647220	SRX3771493	5202	625	837	284
PRJNA437291	SRR6814497	SAMN08647220	SRX3771492	2	0	0	0
PRJNA437291	SRR6814498	SAMN08647220	SRX3771491	2	0	0	0
PRJNA437291	SRR6814499	SAMN08647220	SRX3771490	1	0	0	0
PRJNA437291	SRR6814500	SAMN08647220	SRX3771489	1	0	0	0
PRJNA437291	SRR6814510	SAMN08647220	SRX3771479	0	0	0	0
PRJNA437291	SRR6814527	SAMN08647220	SRX3771462	2	0	0	0
PRJNA437291	SRR6814528	SAMN08647220	SRX3771461	0	0	0	0
PRJNA437291	SRR6814531	SAMN08647220	SRX3771458	7741	952	0	0
PRJNA437291	SRR6814532	SAMN08647220	SRX3771457	4123	865	1175	792
PRJNA437291	SRR6814533	SAMN08647220	SRX3771456	5719	758	0	0
PRJNA437291	SRR6814534	SAMN08647220	SRX3771455	4054	579	0	0
PRJNA437291	SRR6814535	SAMN08647220	SRX3771454	1	0	0	0
PRJNA437291	SRR6814536	SAMN08647220	SRX3771453	1	0	0	0
PRJNA437291	SRR6814537	SAMN08647220	SRX3771452	3423	705	930	663
PRJNA437291	SRR6814538	SAMN08647220	SRX3771451	5287	1108	1652	1182

286 virus RNA presence, which is indicated in green cell coloring.