Oscillations, criticality and responsiveness in complex brain networks

MinKyung Kim[1,2] and UnCheol Lee[1,2]

Corresponding author: UnCheol Lee (uclee@umich.edu)

[1]Department of Anesthesiology, University of Michigan Medical School, Ann Arbor, MI 48109, USA

[2]Center for Consciousness Science, University of Michigan Medical School, Domino's Farms, P.O. Box 385, Ann Arbor, MI 48105, USA

Abstract

Brain networks during unconscious states resulting from sleep, anesthesia, or traumatic injuries are associated with a limited capacity for complex responses to stimulation. Even during the conscious resting state, responsiveness to stimulus is highly dependent on spontaneous brain activities. Many empirical findings have been suggested that the brain responsiveness is determined mainly by the ongoing brain activity when a stimulus is given. However, there has been no systematic study exploring how such various brain activities with high or low synchronization, amplitude, and phase response to stimuli. In this model study, we simulated large-scale brain network dynamics in three brain states (below, near, and above the critical state) and investigated a relationship between ongoing oscillation properties and a stimulus decomposing the brain activity into fundamental oscillation properties (instantaneous global synchronization, amplitude, and phase). We identified specific stimulation conditions that produce varying levels of brain responsiveness. When a single pulsatile stimulus was applied to globally desynchronized low amplitude of oscillation, the network generated a large response. By contrast, when a stimulus was applied to specific phases of oscillation that were globally synchronized with high amplitude activity, the response was inhibited. This study proposes the oscillatory conditions to induce specific stimulation outcomes in the brain that can be systematically derived from networked oscillator properties, and reveals the presence of state-dependent temporal windows for optimal brain stimulation. The identified relationship will help advance understanding of the small/large responsiveness of the brain in different states of consciousness and suggest state-dependent methods to modulate responsiveness.

Author Summary

A responsiveness of the brain network to external stimulus is different across brain states such

as wakefulness, sleep, anesthesia, and traumatic injuries. It has been shown that responsiveness

of the brain during conscious state also varies due to the diverse transient states of the brain

characterized by different global and local oscillation properties. In this computational model study

using large-scale brain network, we hypothesized that the brain responsiveness is determined by

the interactions of networked oscillators when a stimulus is applied to the brain. We examined

relationships between responsiveness of the brain network, global synchronization levels, and

instantaneous oscillation properties such as amplitude and phase in different brain states. We

found specific stimulation conditions of the brain that produce large or small levels of

responsiveness. The identified relationship suggests the existence of temporal windows that

periodically inhibit sensory information processing during conscious state and develops state-

dependent methods to modulate brain responsiveness considering dynamically changed

functional brain network.

Introduction

The brain's responsiveness to external stimuli significantly depends on the overall brain state

and is dramatically restricted in unconscious states resulting from sleep, general anesthesia, or

traumatic injuries. Previous studies with transcranial magnetic stimulation (TMS) have

demonstrated that the effects of stimulation during sleep (NREM), anesthesia, and vegetative

states are simple and constrained to local areas, whereas more complex, diffuse responses are

typically found during the conscious state [1–4]. Furthermore, in the state of wakefulness, the

brain's response to stimuli has also been associated with oscillation properties, such as amplitude

and phase, at the moment when the stimulus is applied. In visual perception tasks, for example,

target detectability is dependent on the power and phase of ongoing alpha waves (~10Hz) [5–14]. Recent repetitive TMS (rTMS) studies have suggested that the instantaneous phase of sensorimotor μ-rhythm in the α band (8-12Hz) is an important determinant of stimulation efficacy [15,16].

Although a link between these oscillation properties and responsiveness has been suggested, there has been no systematic exploration of how a stimulus interacts with spontaneous brain activity and results in state-specific outcomes. In this paper, we hypothesized that the brain responsiveness is determined largely by the networked oscillation properties when a stimulus is applied to the brain. To test this hypothesis, we constructed a large-scale brain network dynamics using coupled oscillator model (Stuart-Landau) based on anatomically informed human brain network structure. In the model, three brain states were defined with modulating the criticality (below, near, and above critical state), which produces various networked oscillations in the brain network (large or small synchronization, amplitude, and phase). Here, we simulated alpha oscillations (~10Hz) that have been successfully simulated by simple coupled oscillators, and examined how well the characteristic responsiveness of various oscillations explains the observed empirical results without implementing details of state-specific neurophysiology. The alpha oscillation models have been successfully explained empirically observed brain network behaviors such as functional connectivity, traveling waves, and network state transitions in the electroencephalography (EEG), magnetoencephalography (MEG), and functional MRI (fMRI), which indicates that alpha oscillations may be fit well to study the responsiveness of the brain network [17–24]. To the best of our knowledge, the response behaviors of various alpha oscillations in the brain network have not been studied.

In the model, the critical state was defined as the state with the largest metastability, a measure of the variance of synchronization fluctuation across time [25]. When a system is near its critical

point, i.e., the balanced state between order and disorder, the largest fluctuation in dynamics is

observed [26,27]. During the conscious resting state in the brain, the dynamics are near the critical

point and marked by large fluctuations in synchrony [28,29]. Notably, disruptions to the resting

state (e.g., sleep, anesthesia, traumatic injuries) appear to mitigate the fluctuation variance by

causing a transition away from criticality [30].

Here we decomposed various brain activities into fundamental oscillation properties: (large/small)

instantaneous global synchronization, amplitude, and phase, modulating the criticality (below,

near, and above critical states) and investigated a relationship between oscillation properties and

a stimulus. We found specific oscillation properties that induce large/small responsiveness, which

is also consistent with the empirical observations for the presence of rhythmic temporal windows

of large/small responsiveness to a stimulus [5,6,31–34]. Our model study may provide a

theoretical framework to interpret complex responses of the brain in different states of

consciousness in terms of fundamental networked oscillation properties. In the following result

sections, we first show that the characteristics of the three brain states and the oscillation

properties at the stimulus onset. Second, we demonstrate distinct response behaviors of the

amplitude and phase synchronization to the pulsatile stimulus across states. Thus, we quantify

the changes of both amplitude and phase synchronization as responsiveness indexes. Next, we

show how the instantaneous global synchronization level correlates with the responsiveness of

amplitude and phase synchronization with different responsiveness measures. Specific phases

of alpha oscillations that induce large or small responsiveness are also demonstrated. Finally, we

compare the responsiveness of the regional stimulation with effective, random, and non-effective

stimulation conditions we identified in the model study. A schematic diagram of the study design

is presented in Fig 1.

Fig 1. Schematic diagram of the study design. In this study, we simulated alpha oscillations using coupled Stuart-Landau model on the human brain network structure that consists of 82 brain regions. The brain states were classified into below, near, and above critical states based on their global synchronization level and the metastability (the variance of synchronization fluctuation). The critical state C_p was defined as the state with the largest metastability. A global pulsatile stimulus was applied to various ongoing alpha oscillations in the three brain states. We then quantified a perturbation response of amplitude (PR_j^{amp}) and a phase synchronization (PR_j^{sync}) for node j after the stimulus onset (j=1, 2, …,82), in a binary fashion. To measure responsiveness to the stimulus, we first defined responsivity R_j, as the temporal average of PR_j^{amp} (PR_j^{sync}) during 600 msec (200 msec). Second, Lempel-Ziv complexity (LZc) of PR_j^{amp} (PR_j^{sync}) was calculated to measure the perturbational complexity of the brain's response to the applied stimulus. In doing so, we identified a relationship between the ongoing brain state and the degree of responsiveness and identified the effective stimulus conditions for inducing a large response. Finally, we applied the stimulus to the occipital region and compared the performance between the effective and random stimulation conditions.

Results

The characteristics of alpha oscillations for the three brain states

To study a relationship between oscillation properties at the stimulus onset and the responsiveness, we simulated networked alpha oscillations of the 82 brain regions using coupled Stuart-Landau model in three different brain states, C_b, C_p, and C_a (see Materials and methods for details). The alpha oscillations demonstrated various oscillation properties across states in terms of variance of instantaneous synchronization, instantaneous synchronization and amplitude. Figure 2A demonstrates the metastability of alpha oscillations in the three brain states, which defined as the variance of instantaneous global synchronization level $r(t)$ over time. The alpha oscillations at C_p show significantly large variances of $r(t)$ compared to those of C_b and C_a. Fig 2B and C demonstrate distinct oscillation properties of the three brain states at the stimulus onsets. The alpha oscillations at C_p have a balanced distribution of $r(t)$ in the middle of the two extreme cases. The distributions of $r(t)$ at C_b and C_a are characterized by small and large $r(t)$, respectively

(Fig 2B). The $r(t)$ values at the stimulus onsets are 0.26 ± 0.04 (Mean $\pm$ SD) for C_b, 0.45 ± 0.11

for C_p and 0.75 ± 0.08 for C_a. The amplitudes of the alpha oscillations of C_p have a broad and

balanced distribution, however, the alpha oscillations of C_b and C_a show narrow/relatively small

amplitudes and broad/biased distribution toward large amplitude, respectively (Fig 2C). The

amplitudes at the stimulus onsets are 1 ± 0.01 for C_b, 1.2 ± 0.07 for C_p and 1.83 ± 0.08 for C_a

(Mean $\pm$ SD). To simplify our model, we applied a simple form of stimulus (pulsatile) to all nodes

for 50 msec, which enables us to exclude the effects of stimulus type and network structure (i.e.,

hub and periphery) in our consideration. Total 500 stimulation trials were performed for each state.

Fig 2. The characteristics of alpha oscillations in the three brain states, C_b C_p, and C_a and
instantaneous oscillation properties at the stimulus onset for each state. (A) Mean and variance
(metastability) of instantaneous global synchronization $r(t)$ for C_b (blue), C_p (red), and C_a (yellow). The
critical state C_p is characterized by the largest metastability (Mean $\pm$ SD = 0.01 ± 0.005), compared to
smaller metastability values of C_b and C_a. (B) Distributions of instantaneous global synchronization $r(t)$ at
the stimulus onset for C_b (blue), C_p (red), and C_a (yellow). Total 500 stimulation trials were performed for
each state. The $r(t)$ values at C_b and C_a are characterized by small and large averages of $r(t)$, respectively.
The $r(t)$ at C_p is balanced in the middle of the two extreme cases. (C) Node amplitude distributions at
stimulus onset for C_b, C_P, and C_a. The amplitudes of alpha oscillations at C_b and C_a are biased with smaller
and larger amplitudes, while the amplitudes at C_p are relatively unbiased. These characteristic and state-
specific alpha oscillations may result in distinct responses to the stimulus.

Distinct responses of amplitude and phase synchronization

The effects of a stimulus at differing levels of global synchronization, amplitude, and phase of the

alpha oscillations across three distinct brain states, C_b, C_p, and C_a, were investigated. Perturbation

responses (PRs) of amplitude ($PR_j^{amp}(t)$) and phase synchronization ($PR_j^{sync}(t)$) of a node j (j = 1,

2, …, 82) at time t were defined as 1 if the change was significant, otherwise, as 0 (see Materials

and methods for details). Fig 3B and C present examples of PR_j^{amp} and PR_j^{sync}. The black dots

represent PR_j^{amp} (PR_j^{sync}) equal to 1 after the stimulation. We investigated the temporal properties

of average PR_j^{amp} and PR_j^{sync} over nodes for the three brain states. The results from 500 stimulus

applications across each brain state was divided into five groups, classified by the level of $r(t)$.

We found distinct response durations of amplitude and phase synchronization following the

stimulation. Notably, PR^{amp} persisted significantly longer than PR^{sync} (p<0.001, sign test, $PR^{amp} > P$

R^{sync} after 100ms for all C_b, C_p, and C_a). For instance, at C_b the PR^{amp} was maintained for

approximately 800 msec, whereas the PR^{sync} diminished before 200 msec. The longer persistence

of PR^{amp} was observed in all three states. This implies that amplitude and phase synchronization

in the brain network are governed by distinct response mechanisms, which might be associated

with distinct temporal dynamics of amplitude and phase synchronization in the resting state [35].

We tested differing stimulus strengths (p=10, 20, 30, 40, and 50) as well as stimulus durations

(50,100,150, 200, 250 msec) at C_p (S1 Fig and S2 Fig). From the following analyses, we quantify

the changes of both amplitude and phase synchronization as responsiveness indexes separately

setting different durations (600 and 200 msec).

Fig 3. Distinct perturbation responses (PRs) of amplitude and phase synchronization. (A) Examples
of global synchronization fluctuations for the three brain states, C_b, C_p, and C_a. Each black triangle indicates
the stimulus timing of one trial. PRs of (B) amplitude (PR^{amp}) and (C) phase synchronization (PR^{sync}) are
presented as examples of one stimulation trial. The stimulus was applied at time zero. A red shaded area
indicates the stimulus duration (= 50 msec). The black dots of each node indicate the time points at which
the stimulus induces significantly increased amplitude (phase synchronization); PR_j^{amp} (PR_j^{sync}) = 1. Average
(D) PR^{amp} and (E) PR^{sync} over all nodes for the trials with different instantaneous global synchronization
levels r(t) are compared, respectively. The instantaneous global synchronization levels were divided into
five groups: q(0)-q(20), q(20)-q(40), q(40)-q(60), q(60)-q(80), and q(80)-q(100) regarding the % quantiles.
The lines with shaded regions indicate the mean $\pm$ standard error. PR^{amp} and PR^{sync} are differentiable, that
is, the PR^{amp} maintains longer than the PR^{sync}.

Correlation between responsivity and instantaneous global synchronization level

We investigated the relationship between instantaneous global synchronization level $r(t)$ and the responsivity of amplitude and phase synchronization, respectively. The responsivity R_j^{amp} (R_j^{sync}) measures the total amount of PR_j^{amp} (PR_j^{sync}), defined as a significant change of amplitude (phase synchronization) of node j within the epoch (600 msec for R^{amp} and 200 msec for R^{sync}) following the stimulus. The epoch was determined to cover the maximum persistence of each responsivity in our simulation. The average responsivity for $< R^{amp} >$ and $< R^{sync} >$ was calculated by taking the average of R_j^{amp} and R_j^{sync} over all nodes in order to investigate the relationship with the global feature $r(t)$. The $< R^{amp} >$ and $< R^{sync} >$ for C_b, C_p, and C_a are presented with respect to $r(t)$ in Fig 4A and 4B. Both $< R^{amp} >$ and $< R^{sync} >$ at C_b (incoherent state) are larger than those of the other states with small variances (Mean $\pm$ SD: 0.7 ± 0.05 for $< R^{amp} >$ and 0.62 ± 0.03 for $< R^{sync} >$). By contrast, the responsivity at C_a (highly synchronized state) is diverse, with a relatively larger $r(t)$ overall. Spearman correlations between $r(t)$ and the $< R^{amp} >$ and $< R^{sync} >$ are presented in Fig 4C & 4D. At C_p, both $< R^{amp} >$ and $< R^{sync} >$ are significantly correlated with the $r(t)$ (Spearman correlations: $\rho^{Ramp} = -0.33, p < 0.001$ for $< R^{amp} >$; $\rho^{Rsync} = -0.53, p < 0.001$ for $<R^{sync} >$). At C_p, a globally desynchronized state showed larger responsivity for both amplitude and phase synchronization to the stimulus, whereas at the brain states far from the critical state, C_b and C_a, the responsivities were not highly correlated with the $r(t)$.

Fig 4. Correlations between instantaneous global synchronization level $r(t)$ and the responsivity of amplitude (R^{amp}) and phase synchronization (R^{sync}). The average responsivity of (A) amplitude and (B) phase synchronization for 500 stimulation trials is presented with respect to r(t) for the three brain states, C_b (blue), C_p (red), and C_a (yellow). Spearman correlations between r(t) and the global (C) amplitude

225 responsivity, ρ^{Ramp}, and (D) synchronization responsivity, ρ^{Rsync}, for C_b (blue), C_p (red), and C_a (yellow) are

226 presented. ***p<0.001 and **p<0.01. At C_p, applying the stimulus at smaller r(t) induces larger R^{amp} and

227 R^{sync}.

Correlation between perturbational Lempel-Ziv complexity and instantaneous global synchronization level

We investigated the relationship between instantaneous global synchronization level $\mathrm{r(t)}$ and the

perturbational complexities of amplitude and phase synchronization. We calculated the Lempel-

Ziv Complexity (LZc) [36] of the PR^{amp} (PR^{sync}) in response to the stimulus, which served as the

measure of perturbational complexity. LZc of the PR is similar to the Perturbational Complexity

Index (PCI), a measure that has been previously applied to experimental data with demonstrably

good performance for distinguishing consciousness from sleep, anesthesia, and pathologic

unconsciousness [2,4]. In the same way with $<\mathrm{R}^{amp}>$ and $<\mathrm{R}^{sync}>$, we first calculated the LZc

of each node with PR^{amp}_j and PR^{sync}_j (binary patterns) for their associated epochs (600 msec and

200 msec, respectively), and calculated the average over 82 nodes defining $<\mathrm{LZc}^{amp}>$ and

$<\mathrm{LZc}^{sync}>$. Fig 5A and 5B present $<\mathrm{LZc}^{amp}>$ and $<\mathrm{LZc}^{sync}>$ with respect to $\mathrm{r(t)}$ for the three

brain states (C_b, C_p, and C_a), and show the results of 500 stimulus trials for each state. Among the

three states, C_p has large $<\mathrm{LZc}^{amp}>$ and $<\mathrm{LZc}^{sync}>$ (See S3 Fig.), indicating that the largest

flexibility and responsiveness of the brain network occurred at the critical state for both amplitude

and phase synchronization. Spearman correlations between $\mathrm{r(t)}$ and $<\mathrm{LZc}^{amp}>$ and $<\mathrm{LZc}^{sync}$

$>$ were presented in Fig 5C & 5D. At C_p, a globally desynchronized state showed larger

perturbational complexities for both amplitude and phase synchronization to the stimulus. The $<$

$\mathrm{LZc}^{amp}>$ at C_p showed a small correlation with $\mathrm{r(t)}$ ($\rho^{LZcamp} = -0.14$, $**p < 0.01$), by contrast, $<$

$\mathrm{LZc}^{sync}>$ at C_p is significantly correlated with $\mathrm{r(t)}$ ($\rho^{LZcsync} = -0.60$, $*** p < 0.001$). The result

implies that, within the critical state, stimulation during desynchronized alpha oscillations induces

more complex perturbation patterns in phase synchronization.

Fig 5. Correlations between instantaneous global synchronization level r(t) and the perturbational

Lempel-Ziv Complexity (LZc) for the amplitude (LZc^{amp}) and the phase synchronization (LZc^{sync}).

The average (A) LZc^{amp} and (B) LZc^{sync} for 500 stimulus trials is presented with respect to the r(t) for C_b

(blue), C_p (red), and C_a (yellow). Spearman correlations between r(t) and (C) the average LZc^{amp}, $\rho^{LZc^{amp}}$

and (D) the average LZc^{sync}, $\rho^{LZc^{sync}}$ are presented for C_b (blue), C_p (red), and C_a (yellow), respectively.

**p<0.01 and ***p<0.001 for the Spearman correlation value. At C_p, applying stimulus at lower level of

instantaneous global synchronization induces larger perturbational complexities for both amplitude and

phase synchronization.

Specific phases of alpha oscillations with inhibited response to external stimuli

We expanded the investigation to explore the amplitude and phase dependencies across different

levels of responsiveness (R^{amp}, R^{sync}, LZc^{amp}, and LZc^{sync}) at C_p. We first classified the alpha

oscillation properties at the stimulus onset in C_p into four groups based on the instantaneous

global synchronization level (high/low) and amplitude (high/low). Nodes were first classified by

determining whether the observed instantaneous global synchronization was higher (HS) or lower

(LS) relative to the average instantaneous global synchronization across 500 stimulus trials. This

same approach was used to further classify the nodes via their measured amplitude (HA/LA)

relative to the average amplitude. We found that the responsiveness of LS & LA (red line) is larger

than that of HS & HA (blue line) in most phases (See black lines in Fig 6, p<0.005, Wilcoxon rank-

sum test). Interestingly, stimulating at specific phases from 60° to 180° of HS & HA shows

significantly smaller responsiveness, which is consistent in four different measures (R^{amp}, R^{sync},

LZc^{amp}, and LZc^{sync}) (Fig 6A-D). By contrast, the synchronization responsiveness (R^{sync} and

LZc^{sync}) of LS & LA have no phase dependence (Fig 6B and 6D). The results suggest that the

existence of periodic temporal windows (phases from 60° to 180 in each cycle of alpha wave) that

create network conditions of small responsiveness to external stimuli. The results for LS & HA

and HS & LA are presented in S4 Fig. Different stimulus strengths at C_p were also tested (S5 Fig

– S8 Fig). The results for the two other states, C_b and C_a are shown in S9 Fig and S10 Fig. As

expected, the strength of stimulus significantly affects the phase dependence at C_p. A strong

stimulus (larger than p=40 in S8 Fig.) begins to induce large changes in both amplitude and phase

synchronization diminishing the phase dependence (p=50 in S9 Fig). On the contrary, a weak

stimulus (p=10, S5 Fig.) produces small changes in amplitude and phase synchronization, thus,

the phase dependences are indistinctive across the conditions (LS&LA, LS&HA, HS&LA, and

HS&HA).

Fig 6. Specific alpha phases with inhibited response to the stimulus at C_p. For each panel, a red (blue)
thick line indicates the average (A) R^{amp}, (B) R^{sync}, (C) LZc^{amp}, (D) LZc^{sync} of nodes with lower (higher)
levels of instantaneous global synchronization and lower (higher) amplitude at the stimulus onset. Lower
and higher levels of instantaneous global synchronization (amplitude) were distinguished by the average
of instantaneous global synchronization levels of all trials (amplitudes of all nodes). Phases of nodes at the
stimulus onset were sorted in 30 phase bins. Thin colored lines indicate the standard errors of
responsiveness. Black lines present phase ranges with significantly different responsiveness between LS
& LA and HS & HA (p<0.005, Wilcoxon rank-sum test). Lower levels of global synchronization and amplitude
(LS & LA) show larger responsiveness to the stimulus than higher levels of global synchronization and
amplitude (HS & HA) in most phases. Notably, for HS & HA, stimulation applied at specific phases of alpha
waves from 60° to 180° showed lower responsiveness, implying the existence of temporal windows during
which there is little response to external stimuli.

Stimulation of occipital region with effective, random, and non-effective conditions

We tested if the effective stimulation conditions identified in the global stimulation also hold for a local stimulation. Here we applied a stimulus to nodes within the radius of 50 mm centered a left cuneus in the occipital region at C_p (13 regions; (left) cuneus, inferior parietal, isthmus cingulate, lateral occipital, lingual, pericalcarine, precuneus, superior parietal, (right) cuneus, isthmus cingulate, lingual, pericalcarine, precuneus) to roughly simulate a stimulation to visual area, and compared four responsiveness measures, R^{amp}, R^{sync}, LZc^{amp}, and LZc^{sync}, of effective, non-effective, and random stimulation conditions. For the effective stimulation, we applied a stimulus during phases from 240° to 360° of alpha oscillations with a low level of instantaneous global synchronization and low amplitude (Fig 7). For the random stimulation, we applied a stimulus to random instantaneous states that may correspond to conventional stimulation methods. For the non-effective stimulation, we stimulated at phases from 60° to 180° of alpha oscillation with a high level of instantaneous global synchronization and high amplitude. We found that the effective (non-effective) stimulation conditions induce larger (smaller) responsiveness for R^{amap}, R^{sync}, LZc^{sync} (Fig 7A, 7B, and 7D). The random stimulation conditions were in the middle of two conditions. The responsiveness of the three conditions was significantly different in R^{amap}, R^{sync}, and LZc^{sync} (Kruskal-Wallis test, p<0.001) and a multi-comparison test was performed using the Tukey-Kramer method (*p<0.05 and ***p<0.001). However, since LZc^{amp} for the global stimulus was barely correlated with the instantaneous global synchronization ($\rho^{LZcamp} = -0.14$ in Fig 5C), and not significantly distinguishable with the amplitude and phase (Fig 6C), LZc^{amp} for the local stimulus also showed relatively small values and can't be differentiated the three stimulation conditions. The results demonstrated that the effective and non-effective stimulation conditions we found in the global stimulation can be also applied to the local stimulation.

Fig 7. Comparison of the responsiveness due to a local stimulation with effective, random, and non-effective stimulation conditions. Average (A) R^{amp}, (B) R^{sync}, (C) LZc^{amp}, and (D) LZc^{sync} were calculated with three different conditions found in the global stimulation; (effective stimulation condition): alpha phases from 240° to 360° with low level of instantaneous global synchronization and low amplitude (red). (random stimulation condition): random selection from all stimulation trials (gray). (non-effective stimulation condition): alpha phases from 60° to 180° with high level of instantaneous global synchronization and high amplitude (blue). Effective (non-effective) stimulation conditions show a larger (smaller) responsiveness for R^{amp}, R^{sync}, and LZc^{sync}. The results of three conditions are significantly different for R^{amp}, R^{sync}, LZc^{sync} (Kruskal-Wallis test, p<0.001). Multiple comparison test was performed with the Tukey-Kramer method (*p<0.05 and ***p<0.001).

Discussion

Many empirical studies have reported that the brain responsiveness to a stimulus is dependent upon brain state and ongoing spontaneous brain activities. Here, we hypothesized that the empirically observed response behaviors of the brain is determined by the interaction way of networked oscillators to a stimulus. To test the hypothesis, we used a human brain network model of coupled oscillators and decomposed the networked oscillations into fundamental oscillation properties (instantaneous global synchronization, amplitude, and phase) within the three brain states (C_b, C_p, and C_a). Each state presents characteristic oscillation features: (below the critical state, C_b) a low amplitude and incoherent state, (near the critical point, C_p) balanced high and low amplitudes with a large synchronization fluctuation, and (above the critical point, C_a) a high amplitude and highly synchronized state. How such characteristic oscillations respond to a stimulus provides a theoretical framework to understand the periodic temporal windows of

large/small responsiveness in conscious resting state as well as the distinct responsiveness of

the brain in different states of consciousness.

Dependence on the criticality of the network: Empirical and computational model studies have

proposed that the brain during conscious resting state resides near the critical state, whereas the

brain states under significant alterations such as anesthesia and traumatic injuries are distant

from the critical state [28–30,37]. In this model study, we observed that the amplitude responsivity

(; the amount of significantly increased amplitudes) at C_b and C_a was larger than that at C_p(Fig 4A).

However, the perturbational complexities of amplitude (; the temporal complexity of significantly

increased amplitudes) at C_b and C_a were smaller than C_p. The brain network at C_p exhibits the

largest perturbational complexity of the amplitude (Fig 5A and S3 Fig A). The amplitude

responsiveness is similar with the TMS outcomes in empirical studies. The TMS-evoked activities

during unconsciousness showed high-amplitude and short-lasting simple response patterns [1,3],

whereas the EEG responses induced by TMS in the resting state showed more complex

perturbational patterns compared to anesthesia and NREM sleep [1–4]. The similarity between

the model and the empirical results proposes that the state-dependent responses of the human

brain to TMS may be due to the distinct response behaviors of the characteristic states

(synchronous/incoherent or high/low amplitudes) of networked oscillators at below, near, and

above critical state.

Dependence on the instantaneous global synchronization: In the model, we found—even

assuming criticality—that the brain network responsiveness correlates with the level of

instantaneous global synchronization at the critical state. Lower levels of instantaneous global

synchronization produced larger responses for both amplitude and phase synchronization,

whereas higher levels of instantaneous global synchronization induced smaller responses.

Considering the large synchronization fluctuations over time at the critical state, it is important for

a stimulus to be applied at the appropriate time in order to induce a larger responsiveness.

Applying a stimulus without consideration of this relationship may in turn produce greater

variability in the stimulation outcomes. In addition, finding the dependence on instantaneous

global synchronization also provides a novel insight on the role of synchronization fluctuation for

the information processing in the brain. In the epochs with lower levels of instantaneous

synchronization, the brain might not be able to integrate distributed information within the brain

network, however, the brain can be highly susceptible to external stimuli. On the contrary, in the

epochs of higher levels of instantaneous synchronization, the brain might easily integrate

distributed information within the brain network, but the brain might not be able to respond to

external stimuli. Regarding the large synchronization fluctuation at critical state, the response

behavior of a network at high/low level of instantaneous synchronization seems to create temporal

windows that can integrate internal or external information separately, which may prevent mixing

of internal and external information in the brain.

Dependence on the amplitude and phase of oscillation: In our model study, we found that the

responsiveness correlates to the amplitude and phase of ongoing alpha oscillations. We

decomposed the alpha waves into high/low instantaneous synchronization levels, high/low

amplitudes and phases ($0°$ to $360°$). The responsiveness of lower instantaneous synchronization

level & low amplitude (LS & LA) was larger than that of higher instantaneous synchronization level

& high amplitude (HS & HA), irrespective to the phases (Fig 6). Importantly, considering the alpha

oscillations of high synchronization and high amplitude, we found that there are specific phases

($60°$ to $180°$) exihibiting significantly lower responsiveness regardless of the responsiveness

measures. The results suggest another periodic temporal windows (phases from $60°$ to 180 in

each cycle of alpha wave) with inhibited response to external stimuli. The time scale of this

temporal window is much smaller than the time scales of synchronization fluctuations.

The amplitude and phase dependences have been observed in empirical studies. The amplitude

and phase of the alpha waves (8-13Hz) are associated with the processing of sensory stimuli,

especially in regards to visual perception [5–14]. Stimulation at lower amplitudes, and at phase

near the trough of alpha waves, showed increased target detectability [6,8,10,11,13,38].

Phosphenes artificially induced by TMS were more detectable in lower pre-stimulus alpha power

and specific alpha phase ranges [7,12,39]. Recent TMS studies also showed that a motor-evoked

potential (MEP) amplitude is dependent on the pre-stimulus cortical μ-rhythm phase [15,16]. The

established dependences on the amplitude and specific phases of alpha waves were consistent

regardless of the stimulus types. Our human brain network model showed that stimuli applied at

specific phases of alpha waves resulted in larger responsivity and perturbational complexity for

both amplitude and phase synchronization at the critical state. Interestingly, the phase

dependence was more pronounced when the stimulus was given at higher amplitudes of alpha

oscillations, which is supported by the visual target detectability experiment [10]. This result

emphasizes the importance of considering the effective stimulation conditions, especially when

the stimulus is applied to the alpha waves with high amplitude and high levels of synchronization.

We also tested if the conditions identified from the global stimulation still hold for a local stimulation.

We applied the same stimulus to the occipital region across three different conditions: effective

(low instantaneous synchronization level, low amplitude, and phases of 240°-360°), non-effective

(high instantaneous synchronization level, high amplitude, and phases of 60°-180°), and random

(the conventional stimulation ways without considering ongoing brain activity). Relative to the

random condition, the effective and non-effective conditions produced significantly larger and

smaller brain responsiveness, respectively. These results may explain why the brain stimulation

outcomes in conventional studies show such large inter-or intra-subject variability [40], which has

affected the reliability and consistency of previous brain stimulation studies. In addition, finding

the specific phases of the alpha oscillations for large or small responsiveness re-illuminates the

historical concept of the "neuronic shutter" described in 1960s [34], which implied the existence

of temporal windows that periodically prohibit sensory information processing during conscious

state [5,31–34,41,42]. Further study is required to identify a relevance of the temporal windows

of sensory information processing to the response behavior of networked alpha oscillations to

external stimuli.

Multi-scale mechanisms of large and small responsiveness: A desynchronized EEG, as

characterized by low amplitude, noisy, and fast-frequency activity, is associated with arousal and

increased awareness [43]. This desynchronized state emerges as the product of highly recurrent

and stochastic interactions within, and between, the excitatory-inhibitory neuronal populations at

both the cellular and network level [43]. At the cellular level, the stochastic interplay of excitatory

and inhibitory conductance sets neurons into a high-conductance state with enhanced

responsiveness. At the population level, neuronal ensembles also become highly responsive to

afferent inputs [43]. At the large-scale brain network level, the responsiveness is determined with

the characteristics of ongoing networked oscillators. However, it still remains to be answered how

the specific oscillation properties induce such large or small responsiveness. One of the

candidates to answer the question is the phase response curve (PRC) which has been studying

in physics. The PRC characterizes the way that a system with a collective periodic behavior

responds to external stimuli. The response of a periodically driven oscillating system to the stimuli

is measured by the phase shift from the original phase. The ongoing phase value dependence of

the phase shift (advancing or delaying the original phase) at which a stimulus occurs is an inherent

characteristic of any oscillatory systems. This method has been applying to many rhythmic

biological systems such as circadian rhythms, cardiac rhythms, and spiking neuron to study how

external stimuli perturb the original rhythms [44–46]. The previous analytic studies discovered the

influential factors causing phase shift and phase synchronization perturbation: a low (high) phase

coherence induces a large (small) phase shift [45]. Stimulation to the phases around stable fixed

point of the PRC increases phase coherence, whereas stimulation to the phases around unstable

fixed point decreases phase coherence. These properties also hold for networks with different

coupling functions, network structure, connectivity, and even for amplitude dynamics of Stuart-

Landau oscillators [47]. Our network model of alpha oscillations can be a complex real world

example of the simple PRC models that constrained by the hierarchical human brain network

connections. If the alpha oscillations possess globally coupled in the human brain network, the

response behavior of alpha oscillations should be governed by the same mechanism. A stability

analysis will be required to identify specific conditions of the human brain network that induce

large or small responsiveness. Furthermore, the future study will be needed to establish how

responsiveness across multiple scales, cellular, neural population, and large-scale brain network,

are related to each other.

In summary, this model study demonstrated for the first time the relationship between network

responsiveness to a stimulus and the decomposed oscillation properties at different network

states (below, near, and above critical state). We found that brain states with lower levels of

instantaneous global synchronization, lower amplitude, and specific phase ranges as well as a

criticality, are important factors to consider for inducing greater network responsiveness. This

study proposes systematically derived oscillatory conditions to induce specific stimulation

outcomes in the brain network. The identified relationship will help advance understanding of the

brain in states of small/large responsiveness and suggest methods by which to modulate

responsiveness.

Limitations: This model study has some limitations. First, our model simulated only alpha

oscillations. We assumed that the alpha oscillations are globally networked in the human brain

because many global features of alpha oscillations such as functional connectivity, traveling

waves, and network state transitions in empirical findings were successfully reproduced with

coupled oscillators of alpha frequency [17–24]. It also well fits to study the response behaviors of

the human brain network due to previous empirical findings [5–14]. Second, we only tested a

pulsatile stimulus. Despite the fact that we examined various strengths and durations of the

pulsatile stimulus, we cannot assure whether other types of stimulus, for instance, continuous

sinusoidal stimulus, gives the same result. Third, we applied a stimulus to all the nodes in the

brain network and later tested only the occipital stimulation in order to simplify the model study.

However, regarding the significant influence of network topology to local brain dynamics, it

requires to identify the effect of stimulation sites on the responsiveness.

Material and Methods

Simulation of networked alpha oscillations in human brain network

We constructed a large-scale functional brain network using coupled Stuart-Landau models on

the human anatomical brain structure to generate spontaneous oscillations. The Stuart-Landau

model with brain topology has been used to replicate the oscillatory dynamics of different types

of electromagnetic brain signals such as electroencephalography (EEG),

magnetoencephalography (MEG) and functional magnetic resonance imaging (fMRI)

[17,18,20,24,48]. The coupled Stuart-Landau model is defined as following:

$$\dot{z}_j(t) = \{\lambda_j + i\omega_j - |z_j(t)|^2\}z_j(t) + A_{ij}\sum_{k=1}^{N}K_{jk}z_k(t) - \tau_{jk} + \beta\xi_j(t) \qquad (1)$$

Here, the state of j_{th} oscillator, $j = 1,2,...,N$, is determined by a complex variable $z_j(t)$ at time t. N

is the total number of brain regions acquired from group-averaged diffusion tensor imaging (DTI)

with 82 nodes [49]. The $A_{jk} = 1$ if there is a connection between j_{th} and k_{th} oscillators, and $A_{jk} = 0$

if there is not, based on the structural brain network. Each node undergoes supercritical Hopf

bifurcation and the dynamics of the oscillator settles on a limit cycle if $\lambda_j > 0$, and on a stable focus

if $\lambda_j < 0$. Here, we fixed all λ_j as 1. The $\omega_j = 2\pi f_j$, is an initial angular natural frequency of each j_{th}

oscillator. We used a Gaussian distribution for the f_j with the mean 10 Hz and standard deviation

0.5 Hz to simulate the alpha bandwidth of human EEG activity [20–22,24]. We also used

homogeneous coupling term $K_{jk} = K$ between j_{th} and k_{th} oscillator from 0 to 0.4 with $\delta K = 0.002$

which determines the global connection strength among brain regions. The time delay between

oscillators $\tau_{jk} = D_{jk}/s$ was introduced with the average speed of axons in brain areas, $s = 7\ ms$

[50], and the distance D_{jk} between brain regions. The node j received the input from connected

nodes k after the time τ_{jk}. The time delays are various, but as long as the time delay is smaller

than a quarter of the period of oscillators, here $\tau_{jk} < {\sim}25\text{ms}$, results are not qualitatively different

[20]. A Gaussian white noise $\xi_j(t)$ for each node was added with the standard deviation $\beta = 1$.

We numerically solved the differential equations of the Stuart-Landau model using the 4th order

Runge-Kutta method with 1,000 discretization steps. The first 5 seconds of time series were

discarded and last 15 seconds were used for the analysis at each simulation. Therefore, each

brain region generates its own spontaneous oscillatory dynamics near alpha bandwidth at each

coupling strength K for one simulation.

Three brain states: below, near, and above critical State

We selected three representative brain states at different coupling strengths based on the level

of global synchronization and the metastability to understand the responsiveness of different

systems. We first calculated an order parameter $r(t)$ to obtain the level of global synchronization

at time t using phase difference $\Delta\theta_{jk}(t) = \theta_j(t) - \theta_k(t)$ at each coupling strength K.

$$r(t) = \left\langle \left| \frac{1}{N}\Sigma_{k=1}^{N} A_{jk} e^{i\Delta\theta_{jk}(t)} \right| \right\rangle \tag{2}$$

Here $r(t) = 1$ if all phases are equal, but $r(t)$ is nearly zero if all phases are randomly distributed

each other. Then we calculated the variance of $r(t)$ for each coupling strength as a metastability

to define the critical state C_p [25]. We considered that generated signals at certain coupling

strength with the largest metastability can be equivalent to the spontaneous alpha oscillations in

normal brain. There have been emerging evidences in computational model studies that the brain

dynamics at critical point show the largest spatiotemporal diversity [29,51]. Notably, a large

repertoire of metastable states exists in brain dynamics during conscious state [48]. With the

critical state, we selected two other representative states which are below (C_b) and above (C_a)

critical states. We defined the C_b and C_a as the states with 10% and 90% percentile values of

averaged order parameters over all coupling strengths, respectively. These states are the

representative states far from the critical states which show small metastability. Generated signals

in these three states were used for the analysis. We used 20 different initial frequency distributions

and selected three representative states for each frequency distribution.

Brain network stimulation procedure

The global pulsatile stimuli were induced as a stimulation with $u(t)$ as following:

$$\dot{z}_j(t) = \{\lambda_j + i\omega_j - |z_j(t)|^2\}z_j(t) + A_{ij}\sum_{k=1}^{N}K_{jk}z_k(t) - \tau_{jk} + \beta\xi_j(t) + u(t) \qquad (3)$$

$$u(t) = \begin{cases} p, & t_1 < t < t_2 \\ 0, & otherwise \end{cases} \qquad (4)$$

Here p is a strength of the stimulus during period $T = t_2 - t_1$. We fixed the $p = 30$ for the analysis

after testing the effect of various stimulus strengths (S1 Fig.). We also fixed the duration of

stimulus, $T = 50 \text{ msec}$ which is about a half of the period of cycle within alpha frequency range.

We tested the stimulus duration from 50 msec to 250 msec (S2 Fig.). We gave a same stimulus

to whole brain network to understand the dependence of response only derived from the network

dynamics of the system. The stimuli were induced at randomly selected timing t_1 for one trial.

Total 25 different timings were selected for each different frequency distribution with 10 iterations.

Therefore, C_b, C_p, and C_a have 500 stimulation trials with 500 different instantaneous brain states,

respectively. We decomposed the instantaneous brain states into the level of instantaneous

global synchronization, amplitude and phase of alpha waves to identify the relationship with the

responsiveness.

Determining significant changes of amplitude and phase synchronization

We assumed that the responses of amplitude and phase dynamics to the stimulus are different

in the system. Therefore, we investigated the perturbation response after the stimulus dividing

into two parts, amplitude and phase synchronization. First, we calculated the instantaneous

amplitude of j_{th} node, $|z_j(t)|$ for each trial. The instantaneous amplitudes for j_{th} node of one trial

were normalized by the mean and standard deviation of the baseline amplitudes of j_{th} node (1

second before the stimulus onset for each iteration). We considered the one tail $(1 - \alpha) * 100^{th}$

quantile with $\alpha = 0.05$ as a significantly increased amplitude after the stimulus onset. An amplitude

response of j_{th} node at time t was defined in a binary fashion: $PR_j^{amp}(t) = 1$ for the significantly

increased amplitude of node j; $PR_j^{amp}(t) = 0$, otherwise. Same procedures were performed for the

phase synchronization. The instantaneous phase synchronization for j_{th} node at time t was

obtained by taking average of the pairwise synchronization between node j and node k at time t,

$$D_j(t) = \left| \frac{1}{N_j} \sum_{k=1}^{N} A_{jk} e^{i\Delta\theta_{jk}(t)} \right| \qquad (5)$$

553   Here N_j is the number of connections of j_{th} node. A synchronization response after the stimulus

554   onset was also defined in a binary fashion: $PR_j^{sync}(t) = 1$ for the significantly increased phase

555   synchronization value; $PR_j^{sync}(t) = 0$, otherwise.

Responsiveness measures: Responsivity & Perturbational Complexity

We defined two different measures to quantify the responsiveness of different instantaneous

brain states to the stimulus. First, the amplitude responsivity R^{amp}_j and the synchronization

responsivity R^{sync}_j were defined by taking average of the perturbation response PR^{amp}_j and PR^{sync}_j

of node j during specific time duration t_{amp} (t_{sync}). It quantifies a total amount of significant

perturbation responses after the stimulus onset. Here we used t_{amp}= 600 msec and t_{sync}= 200

msec, which covers the maximum response changes after the stimulus onset.

Second, we also calculated the Lempel-Ziv complexity (LZc) of the perturbation response for

both amplitude and phase synchronization to understand how much information is contained in

the response. LZc of the perturbation response for node j was defined as following:

$$\mathrm{LZc}^{amp}_j = \frac{c^{amp}_j}{t_{amp}/\log_2 t_{amp}} \qquad (6)$$

$$\mathrm{LZc}^{sync}_j = \frac{c^{sync}_j}{t_{sync}/\log_2 t_{sync}} \qquad (7)$$

Here c^{amp}_j (c^{sync}_j) is the non-normalized Lempel-Ziv complexity of PR^{amp}_j (PR^{sync}_j) calculated by a

LZ76-algorithm [36], i.e., the number of unique patterns in the PR^{amp}_j (PR^{sync}_j) during time t_{amp} (

t_{sync}). We defined a perturbational complexity, LZc^{amp}_j (LZc^{sync}_j) as the normalized c^{amp}_j (c^{sync}_j).

We first investigated the relationships between the level of instantaneous global synchronization

$r(t)$ and the average R and LZc over all nodes at C_b, C_p, and C_a. A Spearman correlation with p-

value was calculated between the $r(t)$ and the average R and LZc at C_b, C_p, and C_a.

We also examined the instantaneous alpha amplitude/phase dependences of responsiveness (

R^{amp}, R^{sync}, LZc^{amp}, and LZc^{sync}) to the stimulus. Here we found the certain ongoing brain states

of alpha oscillations that can induce large or small responsiveness at C_p. We considered the

ongoing brain states with certain levels of instantaneous global synchronization, amplitudes, and

phases inducing large (small) responsiveness as effective (non-effective) stimulation conditions

and applied these conditions to stimulate the local areas described below.

Local stimulation effect

Next, we induced a stimulus to the left cuneus with regions within 50 mm radius in occipital

region at C_p (13 regions; (left) cuneus, inferior parietal, isthmus cingulate, lateral occipital, lingual,

pericalcarine, precuneus, superior parietal, (right) cuneus, isthmus cingulate, lingual,

pericalcarine, precuneus) to confirm that the specific stimulation conditions we found from the

global stimulation can be applied to the local stimulation. We selected the occipital region because

abundant previous sensory stimulation studies mostly targeted the occipital region for visual tasks

within alpha band. We compared the responsiveness measures, R^{amp}, R^{sync}, LZc^{amp} and LZc^{sync}

with effective, non-effective and random stimulation conditions. For the effective (non-effective)

stimulation, we applied the stimulus to the alpha phases from 240° to 360° (60° to 180°) with low

(high) level of instantaneous global synchronization and low (high) amplitude. The stimulus was

applied at random instantaneous states for the random stimulation condition. The stimulus was

applied to random instantaneous brain states of occipital region for the random stimulation. The

number of nodes for random stimulation was determined by a mean value of the number of nodes

for effective and non-effective conditions. We performed 500 different trials at C_p, in which each

trial consists of the stimuli with $p = 300$ for 50 msec at random timing t_1, and classified the nodes

with specific conditions. We used Kruskal-Wallis test to compare significant difference among

responsiveness of three conditions. Multiple comparison test was also performed (*p<0.05 and

***p<0.001).

Acknowledgments

600   This work is supported by grant No. R01 GM098578 (PIs: GM and UL) from the National Institutes

601   of Health. We are grateful to Justin Campbell for his proofreading.

Reference

1.   Ferrarelli F, Massimini M, Sarasso S, Casali A, Riedner BA, Angelini G, et al. Breakdown

in cortical effective connectivity during midazolam-induced loss of consciousness. Proc

Natl Acad Sci. 2010;107: 2681–2686. doi:10.1073/pnas.0913008107

2.   Casali AG, Gosseries O, Rosanova M, Boly M, Sarasso S, Casali KR, et al. A theoretically

based index of consciousness independent of sensory processing and behavior. Sci Transl

Med. 2013;5. doi:10.1126/scitranslmed.3006294

3.   Sarasso S, Rosanova M, Casali AG, Casarotto S, Fecchio M, Boly M, et al. Quantifying

cortical EEG responses to TMS in (Un)consciousness. Clin EEG Neurosci. 2014;45: 40–

49. doi:10.1177/1550059413513723

4.   Sarasso S, Boly M, Napolitani M, Gosseries O, Charland-Verville V, Casarotto S, et al.

Consciousness and complexity during unresponsiveness induced by propofol, xenon, and

ketamine. Curr Biol. Elsevier; 2015;25: 3099–3105. doi:10.1016/j.cub.2015.10.014

5.   Nunn CM, Osselton JW. The influence of the EEG alpha rhythm on the perception of visual

stimuli. Psychophysiology. 1974. pp. 294–303.

6.   Ergenoglu T, Demiralp T, Bayraktaroglu Z, Ergen M, Beydagi H, Uresin Y. Alpha rhythm of

the EEG modulates visual detection performance in humans. Cogn Brain Res. 2004;20:

376–383. doi:10.1016/j.cogbrainres.2004.03.009

7.   Romei V, Brodbeck V, Michel C, Amedi A, Pascual-Leone A, Thut G. Spontaneous

fluctuations in posterior α-band EEG activity reflect variability in excitability of human visual

areas. Cereb Cortex. 2008;18: 2010–2018. doi:10.1093/cercor/bhm229

8.   van Dijk H, Schoffelen J-M, Oostenveld R, Jensen O. Prestimulus Oscillatory Activity in the

Alpha Band Predicts Visual Discrimination Ability. J Neurosci. 2008;28: 1816–1823.

doi:10.1523/JNEUROSCI.1853-07.2008

9.   Busch NA, Dubois J, VanRullen R. The Phase of Ongoing EEG Oscillations Predicts Visual

Perception. J Neurosci. 2009;29: 7869–7876. doi:10.1523/JNEUROSCI.0113-09.2009

10.   Mathewson KE, Gratton G, Fabiani M, Beck DM, Ro T. To See or Not to See:

Prestimulus   Phase Predicts Visual Awareness. J Neurosci. 2009;29: 2725–2732.

doi:10.1523/JNEUROSCI.3963-08.2009

11.   Mathewson KE, Lleras A, Beck DM, Fabiani M, Ro T, Gratton G. Pulsed out of awareness:

EEG alpha oscillations represent a pulsed-inhibition of ongoing cortical processing. Front

Psychol. 2011;2: 1–15. doi:10.3389/fpsyg.2011.00099

12.   Dugue L, Marque P, VanRullen R. The Phase of Ongoing Oscillations Mediates the Causal

Relation between Brain Excitation and Visual Perception. J Neurosci. 2011;31: 11889–

11893. doi:10.1523/JNEUROSCI.1161-11.2011

13.   Milton A, Pleydell-Pearce CW. The phase of pre-stimulus alpha oscillations influences the

visual perception of stimulus timing. Neuroimage. The Authors; 2016;133: 53–61.

doi:10.1016/j.neuroimage.2016.02.065

14.   Benedetto A, Diego L-S, VanRullen R. 2017, Benedetto_et_al, Different responses of

spontaneous and stimulus-related alpha activity to ambient luminance changes.pdf.

15.   Zrenner C, Desideri D, Belardinelli P, Ziemann U. Real-time EEG-defined excitability states

determine efficacy of TMS-induced plasticity in human motor cortex. Brain Stimul. Elsevier

Ltd; 2017;11: 374–389. doi:10.1016/j.brs.2017.11.016

16. Schaworonkow N, Triesch J, Ziemann U, Zrenner C. EEG-triggered TMS reveals stronger

brain state-dependent modulation of motor evoked potentials at weaker stimulation

intensities. bioRxiv. 2018; doi:10.1101/251363

17. Deco G, Cabral J, Saenger VM, Boly M, Tagliazucchi E, Laufs H, et al. Perturbation of

whole-brain dynamics in silico reveals mechanistic differences between brain states.

Neuroimage. Elsevier Ltd; 2018;169: 46–56. doi:10.1016/j.neuroimage.2017.12.009

18. Deco G, Cabral J, Woolrich MW, Stevner ABA, van Hartevelt TJ, Kringelbach ML. Single

or multiple frequency generators in on-going brain activity: A mechanistic whole-brain

model of empirical MEG data. Neuroimage. Elsevier; 2017;152: 538–550.

doi:10.1016/j.neuroimage.2017.03.023

19. Jobst BM, Hindriks R, Laufs H, Tagliazucchi E, Hahn G, Ponce-Alvarez A, et al. Increased

Stability and Breakdown of Brain Effective Connectivity during Slow-Wave Sleep:

Mechanistic Insights from Whole-Brain Computational Modelling. Sci Rep. 2017;7: 1–16.

doi:10.1038/s41598-017-04522-x

20. Moon JY, Lee UC, Blain-Moraes S, Mashour GA. General relationship of global topology,

local dynamics, and directionality in large-scale brain networks. PLoS Comput Biol.

2015;11: 1–21. doi:10.1371/journal.pcbi.1004225

21. Moon JY, Kim J, Ko TW, Kim M, Iturria-Medina Y, Choi JH, et al. Structure shapes

dynamics and directionality in diverse brain networks: Mathematical principles and

empirical confirmation in three species. Sci Rep. 2017; doi:10.1038/srep46606

22. Kim M, Kim S, Mashour GA, Lee U. Relationship of Topology, Multiscale Phase

Synchronization, and State Transitions in Human Brain Networks. Front Comput Neurosci.

2017;11: 1–12. doi:10.3389/fncom.2017.00055

23.   Zhang H, Watrous AJ, Patel A, Jacobs J. Theta and Alpha Oscillations Are Traveling

Waves in the Human Neocortex. Neuron. Elsevier; 2018;98: 1269–1281.e4.

doi:10.1016/j.neuron.2018.05.019

24.   Kim H, Moon J, Mashour GA, Lee U. Mechanisms of hysteresis in human brain networks

during transitions of consciousness and unconsciousness : Theoretical principles and

empirical evidence. 2018; 1–22.

25.   Shanahan M. Metastable chimera states in community-structured oscillator networks.

Chaos An Interdiscip J Nonlinear Sci. 2010;20: 013108. doi:10.1063/1.3305451

26.   Chialvo DR. Emergent complex neural dynamics. Nat Phys. Nature Publishing Group;

2010;6: 744–750. doi:10.1038/nphys1803

27.   Cocchi L, Gollo LL, Zalesky A, Breakspear M. Criticality in the brain: A synthesis of

neurobiology, models and cognition. Prog Neurobiol. Elsevier Ltd; 2017;158: 132–152.

doi:10.1016/j.pneurobio.2017.07.002

28.   Kitzbichler MG, Smith ML, Christensen SR, Bullmore E. Broadband criticality of human

brain network synchronization. PLoS Comput Biol. 2009;5.

doi:10.1371/journal.pcbi.1000314

29.   Haimovici A, Tagliazucchi E, Balenzuela P, Chialvo DR. Brain organization into resting

state networks emerges at criticality on a model of the human connectome. Phys Rev Lett.

2013;110: 1–4. doi:10.1103/PhysRevLett.110.178101

30.   Hutt A, Lefebvre J, Hight D, Sleigh J. Suppression of underlying neuronal fluctuations

mediates EEG slowing during general anaesthesia. Neuroimage. 2018;179: 414–428.

doi:10.1016/j.neuroimage.2018.06.043

31.   VanRullen R, Koch C. Is perception discrete or continuous? Trends Cogn Sci. 2003;7: 207–

213. doi:10.1016/S1364-6613(03)00095-0

32.   VanRullen R. Perceptual Cycles. Trends Cogn Sci. Elsevier Ltd; 2016;20: 723–735.

doi:10.1016/j.tics.2016.07.006

33.   VanRullen   R.   Attention   Cycles.   Neuron.   Elsevier   Inc.;   2018;99:   632–634.

doi:10.1016/j.neuron.2018.08.006

34.   Callaway E, Alexander JD. The temporal coding of sensory data: An investigation of two

theories. J Gen Psychol. 1960;62: 293–309. doi:10.1080/00221309.1960.9920419

35.   Daffertshofer A, Ton R, Kringelbach ML, Woolrich M, Deco G. Distinct criticality of phase

and   amplitude   dynamics   in   the   resting   brain.   Neuroimage.   Elsevier   Ltd;   2018;   1–6.

doi:10.1016/j.neuroimage.2018.03.002

36.   Lempel A, Ziv J. On the complexity of finite sequences IEEE Transactions on Information

Theory Vol IT22. IEEE Trans Inf Theory1. 1976;22: 75–81. doi:10.1109/TIT.1976.1055501

37.   Fekete T, Omer DB, O'Hashi K, Grinvald A, van Leeuwen C, Shriki O. Critical dynamics,

anesthesia   and   information   integration: Lessons   from   multi-scale   criticality   analysis   of

voltage   imaging   data.   Neuroimage.   Elsevier   Inc.;   2018;183:   919–933.

doi:10.1016/j.neuroimage.2018.08.026

38.   Brüers S, VanRullen R. Alpha power modulates perception independently of endogenous

factors. Front Neurosci. 2018;12: 1–8. doi:10.3389/fnins.2018.00279

39.   Samaha J, Gosseries O, Postle BR. Distinct Oscillatory Frequencies Underlie Excitability

of   Human   Occipital   and   Parietal   Cortex.   J   Neurosci.   2017;37:   2824–2833.

doi:10.1523/JNEUROSCI.3413-16.2017

40.   Huang Z, Zhang J, Longtin A, Dumont G, Duncan NW, Pokorny J, et al. Is There a

714   Nonadditive Interaction Between Spontaneous and Evoked Activity? Phase-Dependence

715   and Its Relation to the Temporal Structure of Scale-Free Brain Activity. Cereb Cortex.

716   2017;27: 1037–1059. doi:10.1093/cercor/bhv288

717   41.   Harris KD, Thiele A. Cortical state and attention. Nat Rev Neurosci. Nature Publishing

718   Group; 2011;12: 509–523. doi:10.1038/nrn3084

719   42.   Cecere R, Rees G, Romei V. Individual differences in alpha frequency drive crossmodal

720   illusory perception. Curr Biol. The Authors; 2015;25: 231–235.

721   doi:10.1016/j.cub.2014.11.034

722   43.   Zerlaut Y, Destexhe A. Enhanced Responsiveness and Low-Level Awareness in

723   Stochastic Network States. Neuron. Elsevier Inc.; 2017;94: 1002–1009.

724   doi:10.1016/j.neuron.2017.04.001

725   44.   Granada A, Hennig RM, Ronacher B, Kramer A, Herzel H. Phase response curves

726   elucidating the dynamics of coupled oscillators. Methods Enzymol. 2009;454: 1–27.

727   doi:10.1016/S0076-6879(08)03801-9

728   45.   Hannay KM, Booth V, Forger DB. Collective phase response curves for heterogeneous

729   coupled oscillators. Phys Rev E - Stat Nonlinear, Soft Matter Phys. 2015;92: 1–9.

730   doi:10.1103/PhysRevE.92.022923

731   46.   Ko T-W, Ermentrout B. Phase Response Curves of Coupled Oscillators. 2008; 1–6.

732   doi:10.1103/PhysRevE.79.016211

733   47.   Levnajić Z, Pikovsky A. Phase resetting of collective rhythm in ensembles of oscillators.

734   Phys Rev E - Stat Nonlinear, Soft Matter Phys. 2010;82: 1–10.

doi:10.1103/PhysRevE.82.056202

48. Deco G, Kringelbach ML, Jirsa VK, Ritter P. The dynamics of resting fluctuations in the

brain: Metastability and its dynamical cortical core. Sci Rep. Springer US; 2017;7: 1–14.

doi:10.1038/s41598-017-03073-5

49. van den Heuvel MP, Sporns O. Rich-Club Organization of the Human Connectome. J

Neurosci. 2011;31: 15775–15786. doi:10.1523/JNEUROSCI.3539-11.2011

50. Caminiti R, Carducci F, Piervincenzi C, Battaglia-Mayer A, Confalone G, Visco-Comandini

F, et al. Diameter, Length, Speed, and Conduction Delay of Callosal Axons in Macaque

Monkeys and Humans: Comparing Data from Histology and Magnetic Resonance Imaging

Diffusion   Tractography.   J   Neurosci.   2013;33:   14501–14511.

doi:10.1523/JNEUROSCI.0761-13.2013

51. Hudetz AG, Humphries CJ, Binder JR. Spin-glass model predicts metastable brain states

that   diminish   in   anesthesia.   Front   Syst   Neurosci.   2014;8:   1–9.

doi:10.3389/fnsys.2014.00234

752   **Supporting information**

754 **S1 Fig. Average (A) PR^{amp} and (B) PR^{sync} over all trials with respect to the strength of the**

755 **pulsatile stimuli p.** The error bar indicates a standard deviation. Both PR^{amp} and PR^{sync} are

756 increased when p is increased.

S2 Fig. (Left panel) PR^{amp} and (Right panel) PR^{sync} of one trial with different durations of stimuli with the strength $p = 30$. The stimulus duration was varied from 50 to 250 msec with 50 msec intervals. Red rectangle indicates the duration of stimulus.

S3 Fig. Average perturbational complexity of (A) amplitude (LZc^{amp}) and (B) phase synchronization (LZc^{sync}) at C_b, C_p, and C_a. Kruskal-Wallis test was performed (***p<0.001 for both amplitude and phase synchronization LZc). Multi-comparison test was also performed using Tukey-Kramer method (***p<0.001).

S4 Fig. Alpha phase dependence of responsiveness at C_p (See Fig 6 in the main text). For each panel, a red (blue) thick line indicates the average (A) R^{amp}, (B) R^{sync}, (C) LZc^{amp}, (D) LZc^{sync} of nodes with lower (higher) level of instantaneous global synchronization and lower (higher) amplitude at the stimulus onset. Lower and higher levels of instantaneous global synchronization (amplitude) were distinguished by the average of instantaneous global synchronization levels of all trials (amplitudes of all nodes). Phases of nodes at the stimulus onset were sorted in 30 phase bins. Thin lines indicate the standard errors of responsiveness.4 different conditions were classified; LS & LA (red), LS & HA (orange), HS & LA (light blue), and HS & HA (blue).

S5 Fig. Alpha phase dependence of responsiveness at C_p for the stimuli strength $p = 10$. For each panel, a red (blue) thick line indicates the average (A) R^{amp}, (B) R^{sync}, (C) LZc^{amp}, (D) LZc^{sync} of nodes with lower (higher) level of instantaneous global synchronization and lower (higher) amplitude at the stimulus onset. Lower and higher levels of instantaneous global synchronization (amplitude) were distinguished by the average of instantaneous global synchronization levels of all trials (amplitudes of all nodes). Phases of nodes at the stimulus onset were sorted in 30 phase bins. Thin lines indicate the standard errors of responsiveness.4 different conditions were classified; LS & LA (red), LS & HA (orange), HS & LA (light blue), and HS & HA (blue). Responsiveness is small for all measures.

S6 Fig. Alpha phase dependence of responsiveness at C_p for the stimuli strength $p = 20$.
For each panel, a red (blue) thick line indicates the average (A) R^{amp}, (B) R^{sync}, (C) LZc^{amp}, (D)
LZc^{sync} of nodes with lower (higher) level of instantaneous global synchronization and lower
(higher) amplitude at the stimulus onset. Lower and higher levels of instantaneous global
synchronization (amplitude) were distinguished by the average of instantaneous global
synchronization levels of all trials (amplitudes of all nodes). Phases of nodes at the stimulus onset
were sorted in 30 phase bins. Thin lines indicate the standard errors of responsiveness.4 different
conditions were classified; LS & LA (red), LS & HA (orange), HS & LA (light blue), and HS & HA
(blue). The responsiveness curves are similar with the responsiveness curve with the strength
$p = 30$ except the LZc^{amp}.

S7 Fig. Alpha phase dependence of responsiveness at C_p for the stimuli strength $p = 40$.
For each panel, a red (blue) thick line indicates the average (A) R^{amp}, (B) R^{sync}, (C) LZc^{amp}, (D)
LZc^{sync} of nodes with lower (higher) level of instantaneous global synchronization and lower
(higher) amplitude at the stimulus onset. Lower and higher levels of instantaneous global
synchronization (amplitude) were distinguished by the average of instantaneous global
synchronization levels of all trials (amplitudes of all nodes). Phases of nodes at the stimulus onset
were sorted in 30 phase bins. Thin lines indicate the standard errors of responsiveness. 4 different
conditions were classified; LS & LA (red), LS & HA (orange), HS & LA (light blue), and HS & HA
(blue). The responsiveness curves are similar with the responsiveness curve with the strength
$p = 30$, but synchronization responsiveness measures, R^{sync} and LZc^{sync} are similar for all phase
ranges.

S8 Fig. Alpha phase dependence of responsiveness at C_p for the stimuli strength $p = 50$.
For each panel, a red (blue) thick line indicates the average (A) R^{amp}, (B) R^{sync}, (C) LZc^{amp}, (D)
LZc^{sync} of nodes with lower (higher) level of instantaneous global synchronization and lower
(higher) amplitude at the stimulus onset. Lower and higher levels of instantaneous global
synchronization (amplitude) were distinguished by the average of instantaneous global
synchronization levels of all trials (amplitudes of all nodes). Phases of nodes at the stimulus onset

815 were sorted in 30 phase bins. Thin lines indicate the standard errors of responsiveness. 4 different

816 conditions were classified; LS & LA (red), LS & HA (orange), HS & LA (light blue), and HS & HA

817 (blue). Lower amplitude shows larger amplitude responsiveness in (A) and (C), and lower

818 synchronization shows larger synchronization responsiveness in (B) and (D).

820 **S9 Fig. The instantaneous global synchronization level, node amplitude and phase**

821 **dependences of responsivity R at C_b, C_p, and C_a.** (A) First, 100 trials stimulating at lower (left)

822 / higher (right) levels of instantaneous global synchronization (Low synch and High synch,

823 respectively) in C_b, C_p and C_a were selected. Dots indicate 5% of nodes with the largest (smallest)

824 R^{amp} in one trial. Color of dots indicates the value of R^{amp} of the node. A distance from the origin

825 is the instantaneous node amplitude, and a phase angle is the node phase at the stimulus onset.

826 The phase is important to determine the R^{amp} at C_b. The phase ranges from 60° to 180° of the

827 signal have smaller R^{amp} for all three states. (B) Same as (A), but a color of dots indicates the

828 value of R^{sync} of the node. Nodes with lower amplitude tend to have larger R^{sync} in High synch at

829 C_p. Nodes with lower amplitude tend to have larger R^{sync} in Low synch at C_a.

831 **S10 Fig. The instantaneous global synchronization level, node amplitude and phase**

832 **dependences of perturbational complexity LZc at C_b, C_p, and C_a.** (A) First, 100 trials

833 stimulating at lower (left) / higher (right) levels of instantaneous global synchronization (Low synch

834 and High synch, respectively) in C_b, C_p and C_a were selected. Dots indicate 5% of nodes with the

835 largest (smallest) LZc^{amp} in one trial. Color of dots indicates the value of LZc^{amp} of the node. A

836 distance from the origin is the instantaneous node amplitude, and a phase angle is the node

837 phase at the stimulus onset. The ongoing phase dependence is shown in the LZc^{amp} at C_b. Nodes

838 with lower amplitude tend to show larger LZc^{amp} at C_p in High synch. (B) Same as (A), but a color

839 of dots indicates the value of LZc^{sync} of the node. Nodes with lower amplitude tend to have larger

840 LZc^{sync} in High synch at C_p. Nodes with lower amplitude appeared to have larger R^{sync} in Low

841 synch at C_a.

Model construction

Brain network structure

N=82

Anterior

Posterior

Coupled Stuart-Landau model

$$\dot{z}_j(t) = \left\{\lambda_j + i\omega_j - |z_j(t)|^2\right\} z_j(t) + A_{ij} \sum_{k=1}^{N} K_{jk} z_k(t - \tau_{jk}) + \beta \xi_j(t)$$

Alpha oscillations

State classification

States at different coupling strength

Synchronization fluctuations

Metastability $= Var(r(t))$

Global stimulation

Application of global pulsatile stimuli $u(t)$

$$\dot{z}_j(t) = \left\{\lambda_j + i\omega_j - |z_j(t)|^2\right\} z_j(t) + A_{ij} \sum_{k=1}^{N} K_{jk} z_k(t - \tau_{jk}) + \beta \xi_j(t) + \textcolor{red}{u(t)}$$

Level of instantaneous global synchronization $r(t)$

$$u(t) = \begin{cases} p, & t_1 < t < t_2 \\ 0, & otherwise \end{cases}$$

Signals

Instantaneous oscillation properties

- Amplitude $|z_j(t)|$
- Phase $\theta_j(t)$

Perturbation Response (PR)

Amplitude PR PR_j^{amp}

Phase synchronization PR PR_j^{sync}

• : $PR_j(t) = 1$
▌ : Stimulus duration
▌ : Response period after stimulus onset

Responsiveness; Responsivity & Perturbational complexity

PR^{amp}

PR^{sync}

1. **Responsivity:** $< PR >_t$
   - Amplitude responsivity; R_j^{amp}
   - Phase synchronization responsivity; R_j^{sync}

2. **Perturbational complexity:** Lempel-Ziv complexity (LZc) of PR
   - Amplitude complexity; LZc_j^{amp}
   - Phase synchronization complexity; LZc_j^{sync}

=> Relationship between ongoing oscillation properties & responsiveness

		Responsivity R	Perturbational Complexity LZc
Global oscillation property	Level of instantaneous global synchronization $r(t)$	$< R^{amp} >$ vs. $r(t)$	$< LZc^{amp} >$ vs. $r(t)$
		$< R^{sync} >$ vs. $r(t)$	$< LZc^{sync} >$ vs. $r(t)$
Local oscillation properties	Node amplitude	R_j^{amp} vs. $|z_j(t)|$	LZc_j^{amp} vs. $|z_j(t)|$
		R_j^{sync} vs. $|z_j(t)|$	LZc_j^{sync} vs. $|z_j(t)|$
	Node phase	R_j^{amp} vs. $\theta_j(t)$	LZc_j^{amp} vs. $\theta_j(t)$
		R_j^{amp} vs. $\theta_j(t)$	LZc_j^{sync} vs. $\theta_j(t)$

=> Find effective conditions to induce large R & LZc

Local stimulation

Local stimulation to occipital region with the effective conditions

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7