Generation of conditional auxin-inducible degron (AID) cells and tight control

of degron-fused proteins using the degradation inhibitor auxinole

Aisha Yesbolatova[1], Toyoaki Natsume[1], Ken-ichiro Hayashi[2]

and Masato T. Kanemaki[1]

1. National Institute of Genetics, Research Organization of Information and Systems

(ROIS), and Department of Genetics, The Graduate University for Advanced Studies

(SOKENDAI), Yata 1111, Mishima, Shizuoka 411-8540, Japan

2. Department of Biochemistry, Okayama University of Science, Okayama 700-0005,

Japan

All correspondence should be addressed to:

Masato T. Kanemaki, PhD

Molecular Cell Engineering Laboratory,

National Institute of Genetics, ROIS,

Yata 1111, Mishima, Shizuoka 411-8540, Japan

E-mail: mkanemak@nig.ac.jp

Phone: (+81) 55 981 5830

ORCID: https://orcid.org/0000-0002-7657-1649

Keywords: degron, protein degradation, auxin, genome editing, conditional mutant,

degradation inhibitor

Abstract

Controlling protein expression using a degron is drawing more attention because the protein of interest can be rapidly depleted in a reversible manner. We pioneered the development of the auxin-inducible degron (AID) technology by transplanting a plant-specific degradation pathway to non-plant cells. In human cells expressing an E3 ligase component, OsTIR1, it is possible to degrade a degron-fused protein with a half-life of 15–45 min in the presence of the phytohormone auxin. We reported previously the generation of human HCT116 mutants in which the C terminus of endogenous proteins was fused with the degron by CRISPR–Cas9-based knock-in. Here, we show new plasmids for N-terminal tagging and describe a detailed protocol for the generation of AID mutants of human HCT116 and DLD1 cells. Moreover, we report the use of an OsTIR1 inhibitor, auxinole, to suppress leaky degradation of degron-fused proteins. The addition of auxinole is also useful for rapid re-expression after depletion of degron-fused proteins. These improvements enhance the utility of AID technology for studying protein function in living human cells.

1. Introduction

Conditional depletion of a protein of interest (POI) is a powerful approach to analyse its function in vivo, especially for POIs that are essential for cell viability. Recently, conditional approaches using a degron have been drawing increased attention [1]. A degron-fused protein can be rapidly and efficiently degraded when needed, so that the primary defect arising from the depletion can be observed before the phenotype is complicated or compromised by the secondary defects. For this purpose, we pioneered the establishment of the auxin-inducible degron (AID) technology to control degron-fused proteins in yeast and mammalian cells (**Figure 1A**) [2]. When expressed in non-plant cells, TIR1 of rice (OsTIR1) forms a chimeric SCF (Skp1–Cul1–F box) complex with the other endogenous components. The SCF–OsTIR1 E3 ubiquitin ligase is only activated when IAA or NAA (a natural or synthetic auxin, respectively) is bound (**Figure 1A**). We identified a 7 kD degron termed mini-AID (mAID) and others identified similar AID degrons [3-5]. A POI fused with a mAID is recognized by SCF–OsTIR1 for ubiquitylation and subsequent proteasomal degradation (**Figure 1A**).

Recently, we showed that conditional human cells can be generated by tagging endogenous genes with a mAID cassette using CRISPR–Cas9-based gene tagging [6]. This technology has recently been applied in many studies. One particular example is the assessment of chromosome architectures [7-10]. Moreover, AID technology has been applied to other model organisms, such as fission yeast, fruit fly, nematode, zebrafish and the parasitic *Toxoplasma gondii* [11-15]. These studies support the idea that the AID technology can be used as a standard method to achieve conditional protein depletion. However, a drawback of this technology is that the expression level of mAID-fused proteins can be low in OsTIR1-expressing

cells [6]. This "basal degradation" might be caused by the presence of contaminating

auxin-like chemicals in bovine serum or culture media [1]. To achieve a tight control

of the expression of mAID-fused proteins, an improvement in this technology is

awaited.

Here, we describe a method to generate conditional human HCT116 and

DLD1 mutants by homology-directed repair (HDR)-mediated gene tagging using

CRISPR–Cas9 (**Figure 1B**). We offer a new series of plasmids for N- or C-terminal

tagging with mAID and other tags. To overcome the problems associated with basal

degradation, we used a TIR1 inhibitor called auxinole (**Figure 5A**) [16]. It was

possible to suppress basal degradation and rapidly recover expression after

depletion by supplementing culture media with auxinole.

2. Construction of CRISPR and donor plasmids for tagging

Figure 1B shows the procedures that were used to generate conditional AID

cells, which typically required one month of work. We previously reported mAID

tagging at the C terminus of a POI (**Figure 2B**) [6]. We now developed a procedure

to tag a POI with mAID at the N terminus (**Figure 2A**) [17]. We offer N- or C-terminal

tagging   plasmids   with   mAID   or   other   tags   at   Addgene

(https://www.addgene.org/Masato_Kanemaki/) and the National Bio-resource Center

(NBRP) (http://dna.brc.riken.jp/en/gsb0000en/rdb08468) (**Figure 2C**).

2.1. Construction of a CRISPR–Cas9 plasmid

To identify a CRISPR–Cas9 targeting site, we usually choose an appropriate

sequence within 50 bp upstream or downstream from the ATG or stop codon. We

use   the   following   target   finder   sites.   IDT   custom   Alt-R   guide   design:

https://sg.idtdna.com/site/order/designtool/index/CRISPR_CUSTOM. WEG CRISPR

finder: https://www.sanger.ac.uk/htgt/wge/. We mainly use pX330-U6-Chimeric_BB-

CBh-hSpCas9 (Addgene #42230) to express the SpCas9 nuclease and guide RNA

according to the protocol of Ran et al. [18]. However, it should be possible to use a

plasmid encoding other Cas9 variants. As discussed in the next section, it is

important to destroy or lose the target site within the genome by the HDR-mediated

insertion of a tagging cassette.

2.2. Construction of a donor plasmid

We described previously a method to construct donor plasmids by generating

homology arms (HAs) using long primers and gene synthesis [6]. A downside of this

strategy is that the HAs are relatively short (up to 200 bp each) and that it can be

costly. As an alternative (and economic) approach, we describe a method to clone

HAs from the genomic DNA (**Figure 3**). The N- or C-terminal coding region (about

1000 bp) is amplified by PCR, followed by cloning into a conventional cloning

plasmid (such as pBluescript II). After confirming the sequence, a cloning site for

inserting a tagging cassette with a selection marker is created by inverse PCR. The

cassette is cloned at the cloning site to complete the construction of the donor

plasmid, which contains the HA at both ends (about 500 bp each) (**Figure 3A and B**).

Importantly, the donor plasmid has to be designed to destroy the CRISPR target site

when inserted into the genome. In the case of targeting a noncoding locus, it is

possible to delete part of the target sequence, unless a functional noncoding element

is absent. In the case of targeting a coding locus within a gene, it is important to

introduce silent mutations, to avoid re-cutting after HDR-mediated insertion. Another

important point is that a cassette encoding a tag has to be cloned in frame with the

gene of interest, to be able to express a fusion protein. All cloning procedures can be

carried out using a standard cloning method. In cases in which standard cloning

using restriction enzymes and ligase is difficult, it is possible to employ a

recombinase-mediated cloning method, such as In-Fusion cloning or Gibson

assembly.

To select clones with a biallelic insertion efficiently, we transfected two donor

plasmids containing a Neo or Hygro marker, respectively, for dual-antibiotic selection

[6]. As an alternative approach, it is possible to generate clones with a biallelic

insertion using a single donor plasmid with a single selection marker. In this case,

both monoallelic and biallelic clones can be obtained.

3. Transfection and colony formation

We established near-diploid colon cancer cell lines, HCT116 and DLD1,

constitutively or conditionally expressing OsTIR1 (CMV- or Tet-OsTIR1, respectively)

by inserting the transgene at the safe AAVS1 locus [6]. These parental cell lines are

available from NBRP or our laboratory upon request. Alternatively, it is possible to

generate a new parental cell line. In this case, plasmids that can be used for

introducing OsTIR1 at AAVS1 are available at Addgene and NBRP (**Figure 2D**) [6].

A CRISPR–Cas9 plasmid for targeting the AAVS1 locus is also available from

Addgene (AAVS1 T2 CRISPR in pX330: #72833).

3.1. Transfection

1. Prepare a sub-confluent culture of low-passage-number HCT116 or DLD1

CMV/Tet-OsTIR1 cells.

2. Collect cells by trypsinization to prepare a cell suspension. Count and dilute the

cells to 0.5×10^5 cells/mL. Seed 1 mL of the diluted cells per well in a 12-well

plate and culture the cells at 37 ⁰C for two days. HCT116 cells are slow to attach

to the bottom. We allow two days of culture before transfection.

3. Prepare the transfection mixture by mixing 2 µL of 200 ng/µL CRISPR–Cas9

plasmid or TE (control), 2.5 µL of 200 ng/µL donor plasmid, 45.5 µL of Opti-MEM

I Reduced Serum Medium (ThermoFisher Scientific, 31985062) and 4 µL of

FuGENE HD Transfection Reagent (Promega, E2311), and incubate the mixture

at RT for 15 min before applying to the cells.

4. One day after transfection, collect the cells in 1 mL of medium after trypsinization.

Depending on the efficiency of CRISPR–Cas9 and donor insertion, the number of

colonies obtained will vary significantly. To pick up single colonies successfully,

prepare several dilutions of transfected cells in a 10 cm dish (dilutions range from

100 to 1000 times; 100 to 10 µl of cell suspension into one 10 cm dish containing

10 mL of culture medium).

3.2. *Antibiotic selection and colony formation*

The final concentration of antibiotics is shown below.

Neo: 700 µg/mL of G418 Sulfate (potency-based)

Hygro: 100 µg/mL of Hygromycin B Gold (InvivoGen, #ant-hg)

BSD: 10 µg/mL of Blasticidin S Hydrochloride

If using a donor harbouring Neo, add G418 soon after plating the transfected

cells. Hygromycin B Gold or Blasticidin S should be added one day after plating for

cell recovery. In case of dual selection, add G418 and Hygromycin B Gold at the

same time one day after plating. Replace culture media with a fresh one containing

an appropriate antibiotic every 3–4 days until colonies become visible (usually 11–13

days after plating).

4. Colony isolation, expansion in 96-well plates and genotyping

Visible colonies should form on the selection dishes after 11–13 days of

culture. If the tagging worked well, you should find more colonies on the plates

containing cells transfected with the CRISPR–Cas9 plasmid than in those containing

cells without the CRISPR–Cas9 plasmid. Although this is a good indicator of

successful tagging in many cases, gene tagging sometimes works well even if no

differences in colony number are observed. We usually pick at least 32 and 16

clones for single- and dual-antibiotic selection, respectively, to obtain clones

harbouring the tag on both alleles.

4.1. Colony isolation

1. Prepare a 96-well plate containing 10 µL of trypsin/EDTA solution in each well.

2. Replace the culture medium in the dish with PBS.

3. Under a stereo microscope, pick a single colony using a micropipette with the

volume set at 25 µL (Use a micropipette such as a Gilson pipetman P200.).

4. Transfer the colony to the 96-well plate containing trypsin/EDTA for trypsinization.

After transferring each set of 16 colonies, add 100 µL of media to each well, to

quench trypsin. Repeat this step to obtain a sufficient number of clones.

5. After isolating clones, resuspend cells in culture medium by pipetting. Culture the

isolated clones for 2–3 days until most clones become confluent.

6. Duplicate the 96-well plate after trypsinization, and culture the cells on the plates

for an additional 2–3 days.

One 96-well plate will be used to prepare a frozen stock and the other will be used to

prepare genomic DNA for PCR genotyping.

4.2. Frozen mini-stock

1. Remove the medium from one 96-well plate and wash cells once with 100 µL of

PBS. Add 10 µL of trypsin/EDTA and incubate at 37 ºC for 3 min.

2. Resuspend cells in 50 µL of medium and transfer them to 0.75 µL Matrix Storage

Tubes (ThermoFisher Scientific) containing 50 µL of Bambanker DIRECT

medium (Nippon Genetics, CS-06-001). Mix the cells by pipetting.

3. Close the tubes with a cap and store them at –80 ºC until PCR genotyping is

finished.

4.3. Genomic DNA preparation

1. Remove the medium from the 96-well plate and wash cells once with 100 µL of

PBS. Add 75 µL of the DirectPCR working solution (0.5 × DirectPCR Lysis

Reagent-Cell [Viagen Biotech, 302-C] containing 0.5 mg/ml of Proteinase K).

2. Seal the plate with an aluminium seal and incubate it at 55 ºC for >6 h in a

rocking incubator.

3. Place the plate in a Tupperware containing wet tissue papers and float the wet

chamber in a water bath at 85 ºC for 1.5 h, to inactivate proteinase K.

4. Spin the plate to collect the samples and use 1 µL for PCR genotyping.

4.4. Genotyping by PCR

1. Design appropriate primers to check the insertion by PCR (**Figure 4A** and **B**).

We designed a primer set to detect both the wild-type (WT) and inserted alleles

(primer set a), and another primer set to detect only the inserted allele (primer

set b).

2. Set up a PCR reaction using Tks Gflex DNA Polymerase (Takara Bio, R060A) (1

× Gflex PCR Buffer, 0.3 µM primers and 0.4 U of Tks Gflex DNA Polymerase in a

15 µL reaction). Add 1 µL of the genomic DNA to the 15 µL reaction mixture. As

controls, add genomic DNA extracted from the parental cells to the reaction

mixture. Perform PCR (30 cycles of 98 ºC for 10 s, 55 ºC for 15 s and 68 ºC for

0.5 min/kb).

3. Examine PCR products using agarose gel electrophoresis or the MultiNA

microchip electrophoresis system (Shimadzu MCE-202) (**Figure 4C**).

4. Identify clones showing the expected band patterns of biallelic insertion. An

example of N-terminal tagging of CENPC is shown in **Figure 4C**. Clones with

biallelic insertion are highlighted by a red circle (clones 2 and 8). These clones

can be checked further by genomic sequencing. Expand these clones of the

mini-stock kept at –80 ºC.

5. Testing degradation of mAID-fused proteins

If mAID-mClover or mAID-mCherry2 was fused to a POI, depletion of the

mAID-fused POI upon the addition of auxin can be checked easily by flow cytometry

or microscopy. If an antibody that detects the POI is available, it is possible to

assess the loss of the WT protein and confirm the expression of the mAID-fused

protein. Next, we describe our methods to check clones by Western blotting and flow

cytometry.

5.1. Confirmation by Western blotting

1. Grow clones that have been confirmed by PCR in a 6-well plate. To prepare a

control with mock treatment (DMSO), one clone should be grown in two wells.

Grow cells to 50% confluency.

2. Prepare a 500 mM stock solution of IAA in DMSO and keep aliquots at −20 ºC.

3. If testing cells constitutively expressing OsTIR1 (CMV-OsTIR1), add IAA to one

well at the final concentration of 200–500 µM. Cells growing in the other well

should be mock treated with DMSO. To test cells conditionally expressing

OsTIR1 (Tet-OsTIR1), add IAA and 0.2 µg/mL of doxycycline. Grow the treated

cells for 12–24 h.

4. Collect cells by trypsinization and wash them in culture media.

5. Collect cells in a microtube and wash once with PBS.

6. Mix the cell pellet in 50 µL of RIPA buffer (25 mM Tris-HCl [pH 7.6], 150 mM

NaCl, 1% NP40, 1% sodium deoxycholate and 0.1 % SDS) and incubate the tube

for 30 min on ice.

7. Remove debris by centrifugation at 10000 rpm for 5 min at 4 ºC. Collect the

supernatant and mix with the same volume of 2 × Laemmli SDS sample buffer

(Tris-HCl [pH 6.8], 4% SDS, 20% glycerol, 10% 2-mercaptoethanol and 0.004%

bromophenol blue) before incubating at 95 ºC for 5 min.

8. Typically, load 5 µL of the protein sample for Western blotting. Commercial

antibodies are available to detect mAID and OsTIR1 (MBL M214-3 and PD048,

respectively). The example of mAID-fused CENPC is shown in **Figure 4D**.

5.2. Confirmation by flow cytometry

1. Grow and treat cells in a 6-well plate as described in section 5.1.

2. Remove the media and wash cells once with PBS. After trypsinization, resuspend

cells in media and transfer them to a microtube. Remove most of the media after

spinning down cells and resuspend them in the residual medium by pipetting.

3. Resuspend the cells in 1 mL of 4% paraformaldehyde/PBS by pipetting. Incubate

in the dark at 4 ºC. Note that a methanol-free paraformaldehyde solution should

be used to preserve the fluorescence of mClover and mCherry2.

4. Before flow cytometric analysis, wash the fixed cells once with PBS containing

1% of BSA. Resuspend the washed cells in an appropriate volume of PBS

containing 1% of BSA and analyse them after filtration with a strainer to remove

aggregation.

6. Suppression of degradation using auxinole

We demonstrated previously that HCT116 cells in which mAID-mClover was

fused to the cytoplasmic dynein heavy chain 1 (DHC1) protein showed a growth

defect when OsTIR1 was expressed without IAA [6]. This was caused by partial

degradation of DHC1-mAID-mClover (DHC1-mACl). We subsequently noted that, in

some cases, the expression level of mAID-fused proteins was significantly reduced

in cells constitutively expressing OsTIR1 (data not shown). Moreover, in some cases,

we could only introduce a biallelic mAID into a POI in the conditional Tet-OsTIR1

background (and not in the constitutive CMV-OsTIR1 background) [6]. These results

indicated that mAID-tagged proteins could be ubiquitylated by OsTIR1 for partial

degradation even without the addition of exogenous IAA.

We aimed to suppress this basal degradation by using a chemical antagonist

of TIR1. Previously, we developed a potential TIR1 inhibitor named auxinole to

control auxin signalling in plants [16]. Auxinole contains a moiety that is analogous to

IAA and binds to the IAA-binding pocket of TIR1 (**Figure 5A**, shown in blue).

Because of steric inhibition by the additional dimethylphenylethyl moiety (**Figure 5A**,

shown in red), auxinole blocks the association of the degron domain of the AUX/IAA

proteins. For the experiments in this work, auxinole was dissolved in DMSO to make

a 200 mM stock solution, which was stored at -20 C°.

We initially tested whether auxinole affected the growth of HCT116 cells and

found that it did not alter the cell cycle or colony formation when added up to 200 μM

(data not shown). Subsequently, we tested the suppression of basal degradation

using DHC1-mACl cells in the HCT116 Tet-OsTIR1 background [6]. Addition of

doxycycline induced OsTIR1 expression, which caused partial depletion of DHC1-

mACl and mitotic arrest in many cells (**Figure 5B**). This showed that OsTIR1

expression, even in the absence of auxin, induced a mitotic phenotype that was

analogous to knockdown or inhibition of dynein [19, 20]. The addition of auxinole

together with doxycycline clearly suppressed the downregulation of DHC1-mACl and

the mitotic arrest (**Figure 5B**). To test whether DHC1-mACl could be rapidly depleted,

we added doxycycline with or without auxinole for 24 h. We monitored the

expression levels of DHC1-mACl by flow cytometry, and found that basal

degradation was mostly suppressed in the cells treated with doxycycline and

auxinole (**Figure 5C**, compare the boxes shown in red). Subsequently, the culture

media was replaced with fresh one containing doxycycline and IAA, but not auxinole.

Figure 5C shows that DHC1-mACI was rapidly degraded after medium replacement

and was mostly depleted within 4 h.

An advantage of AID technology is that the expression level of mAID-fused

proteins can be reversibly controlled [2]. We expected that auxinole would be useful

for re-expression after depletion, because IAA-bound OsTIR1 can remain active for a

while, even after the removal of IAA from the culture medium. To test this idea, we

used HCT116 CMV-OsTIR1 cells in which the cohesin subunit RAD21 was fused to

mAID-mClover (RAD21-mACI) [6]. Initially, we depleted RAD21-mACI by adding IAA

for 24 h (**Figure 6A**). Subsequently, we replaced the medium with fresh media with

or without auxinole, and collected time-course samples to monitor the expression

levels of RAD21-mACI by flow cytometry (**Figure 6B**). We found that recovery of

RAD21-mADI was significantly rapid and sharp when auxinole was added, compared

with cells without auxinole. These results suggest that the OsTIR1 inhibitor auxinole

is useful for the tight control of the expression of mAID-fused proteins in human cells.

7. Conclusion

We described a CRISPR–Cas9-based method that can be used to fuse

endogenous POIs to mAID and other tags. We developed new plasmids for N-

terminal tagging, so that it is now possible to tag the N and C termini of POIs (**Figure**

2). To suppress basal degradation in cells expressing OsTIR1, we used the OsTIR1

antagonist auxinole (**Figure 5A**). Even in the Tet-OsTIR1 background cells, it is now

possible to induce rapid degradation of mAID-fused proteins by inducing OsTIR1 in

the presence of auxinole (**Figure 5C**). Moreover, auxinole is useful for the re-

expression of mAID-fused POIs after depletion (**Figure 6B**). The use of auxinole

allows the rapid, tight and efficient control of the expression of mAID-fused POIs.

Other genetic systems also enable the control of degron- or tag-fused POIs using a

chemical ligand [1, 21-23]. However, to the best of our knowledge, there are no

inhibitors that allow the tight control of these systems. AID technology—now

combined with the degradation inducer, auxin, and the inhibitor, auxinole—will be

particularly useful to dissect biological networks, such as transcriptional cascades,

signal transduction and cell-cycle control systems, in which a primary defect caused

by the loss of a POI leads to secondary defects. We hope that the method described

in this manuscript will enhance the utility of AID technology for functional studies of

endogenous proteins in living cells.

Acknowledgements

We thank Venny Santosa, Akemi Mizuguchi and Tomoko Ashikawa for their

support. This study was supported by JSPS KAKENHI grants 17K15068 to TN, and

18H02170 and 18H04719 to MTK; and by research grants from JST A-STEP (grant

number AS2915150U), the Canon Foundation, the Asahi Glass Foundation and the

Takeda Science Foundation.

Figure legends

Figure 1. Schematic illustration of the AID system. (**A**) Indole-3-acetic acid (IAA)- or

1-naphtaleneacetic acid (NAA)-bound OsTIR1 associates with mini-AID (mAID) and

promote ubiquitylation of the mAID-fused protein for proteasomal degradation. (**B**)

Strategy used for generating conditional AID cells by tagging an endogenous POI.

Figure 2. Tagging of an endogenous POI using CRISPR–Cas9-based genome

editing. (**A**) Fusing a tag at the N terminus of a POI. CRISPR–Cas9 generates a

double-strand break (DSB) near the first ATG site for insertion of a donor harbouring

a marker-P2A-tag cassette. The fusion proteins will be processed at P2A to express

the marker and tag-fused POI. (**B**) Fusing a tag at the C terminus of a POI.

CRISPR–Cas9 generates a DSB near the stop site for insertion of a donor

harbouring a tag and a marker. The fusion protein and the marker will be expressed

independently. (**C**) List of N- and C-terminal tagging plasmids. (**D**) List of donor

plasmids used for introducing OsTIR1 at the AAVS1 locus.

Figure 3. Schematic illustration of donor plasmid construction. (**A**) A DNA fragment

containing the first ATG site (about 1 kb) is cloned into a cloning plasmid (such as

pBluescript II). After creating restriction enzyme sites (such as SalI and BamHI) by

inverse PCR, a DNA fragment containing marker-P2A-tag is cloned into the plasmid,

to generate a donor vector. (**B**) A DNA fragment containing the stop site (about 1 kb)

is cloned into a cloning plasmid. After creating a restriction enzyme site (such as

BamHI) by inverse PCR, a DNA fragment containing a tag and marker is cloned into

the plasmid, to generate a donor vector.

Figure 4. Confirmation of isolated clones. (**A**) Genotyping of N-terminally tagged

alleles by PCR. The primer set (a) amplifies a smaller PCR product (1–1.5 kb) from

the WT allele, while it amplifies a larger product (1–1.5 kb plus the size of the

insertion) from the tagged allele. The primer set (b) generates a PCR product from

the tagged allele exclusively (and not from the WT allele). Note that the primer set

(a) must be designed outside of the homology arms. (**B**) Genotyping of N-terminally

tagged alleles by PCR. The PCR strategy is analogous to that used to analyse N-

terminal tagging. (**C**) PCR genotyping of the CENPC allele, in which a Hygro-P2A-

mAID-mClover cassette was inserted at the N-terminal coding region. HCT116 CMV-

OsTIR1 parental cells were used. (**D**) Confirmation of the fusion protein by WB. WT

or mAID-CENPC clones in the HCT116 CMV-OsTIR1 background were treated with

DMSO or 500 µM IAA for 24 h. Anti-CENPC, anti-mAID and anti-tubulin antibodies

(MBL, PD030, M214-3 and M175-3, respectively) were used for detection.

Figure 5. Use of auxinole to control DHC1-mAID-mClover (DHC1-mACl) Tet-OsTIR1

cells. (**A**) The structure of IAA and auxinole. (**B**) Microscopic analysis of DHC1-mAC

Tet-OsTIR1 cells. The cells were treated with 0.2 µg/mL of doxycycline (Dox) or Dox

with 200 µM auxinole for 48 h before microscopy. (**C**) Flow cytometric analysis of

DHC1-mAC Tet-OsTIR1 cells. The cells were treated with 0.2 µg/mL of Dox or Dox

with 200 µM auxinole for 24 h before replacing the culture medium with Dox and 500

µM IAA.

Figure 6. Re-expression of RAD21-mAID-mClover (RAD21-mACl). (**A**) Scheme of

the experiment. The RAD21-mACl CMV-OsTIR1 cells were treated with 500 µM IAA

for 24 h before replacing the medium with fresh medium with or without 200 µM

auxinole. (**B**) Flow cytometric analysis of RAD21-mACl cells.

References

1. Natsume, T., and Kanemaki, M.T. (2017). Conditional Degrons for Controlling
Protein Expression at the Protein Level. Annu Rev Genet *51*, 83-102.
2. Nishimura, K., Fukagawa, T., Takisawa, H., Kakimoto, T., and Kanemaki, M.
(2009). An auxin-based degron system for the rapid depletion of proteins in
nonplant cells. Nat Methods *6*, 917-922.
3. Kubota, T., Nishimura, K., Kanemaki, M.T., and Donaldson, A.D. (2013). The
Elg1 replication factor C-like complex functions in PCNA unloading during
DNA replication. Mol Cell *50*, 273-280.
4. Morawska, M., and Ulrich, H.D. (2013). An expanded tool kit for the auxin-
inducible degron system in budding yeast. Yeast *30*, 341-351.

5.   Brosh, R., Hrynyk, I., Shen, J., Waghray, A., Zheng, N., and Lemischka, I.R.
(2016). A dual molecular analogue tuner for dissecting protein function in
mammalian cells. Nat Commun *7*, 11742.
6.   Natsume, T., Kiyomitsu, T., Saga, Y., and Kanemaki, M.T. (2016). Rapid
Protein Depletion in Human Cells by Auxin-Inducible Degron Tagging with
Short Homology Donors. Cell Rep *15*, 210-218.
7.   Rao, S.S.P., Huang, S.C., Glenn St Hilaire, B., Engreitz, J.M., Perez, E.M.,
Kieffer-Kwon, K.R., Sanborn, A.L., Johnstone, S.E., Bascom, G.D., Bochkov,
I.D., et al. (2017). Cohesin Loss Eliminates All Loop Domains. Cell *171*, 305-
320 e324.
8.   Nora, E.P., Goloborodko, A., Valton, A.L., Gibcus, J.H., Uebersohn, A.,
Abdennur, N., Dekker, J., Mirny, L.A., and Bruneau, B.G. (2017). Targeted
Degradation of CTCF Decouples Local Insulation of Chromosome Domains
from Genomic Compartmentalization. Cell *169*, 930-944 e922.
9.   Gibcus, J.H., Samejima, K., Goloborodko, A., Samejima, I., Naumova, N.,
Nuebler, J., Kanemaki, M.T., Xie, L., Paulson, J.R., Earnshaw, W.C., et al.
(2018). A pathway for mitotic chromosome formation. Science *359*.
10.   Bintu, B., Mateo, L.J., Su, J.H., Sinnott-Armstrong, N.A., Parker, M., Kinrot, S.,
Yamaya, K., Boettiger, A.N., and Zhuang, X. (2018). Super-resolution
chromatin tracing reveals domains and cooperative interactions in single cells.
Science *362*.
11.   Kanke, M., Nishimura, K., Kanemaki, M., Kakimoto, T., Takahashi, T.S.,
Nakagawa, T., and Masukata, H. (2011). Auxin-inducible protein depletion
system in fission yeast. BMC Cell Biol *12*, 8.
12.   Brown, K.M., Long, S., and Sibley, L.D. (2018). Conditional Knockdown of
Proteins Using Auxin-inducible Degron (AID) Fusions in Toxoplasma gondii.
Bio Protoc *8*.
13.   Trost, M., Blattner, A.C., and Lehner, C.F. (2016). Regulated protein depletion
by the auxin-inducible degradation system in Drosophila melanogaster. Fly
(Austin) *10*, 35-46.
14.   Zhang, L., Ward, J.D., Cheng, Z., and Dernburg, A.F. (2015). The auxin-
inducible degradation (AID) system enables versatile conditional protein
depletion in C. elegans. Development *142*, 4374-4384.
15.   Daniel, K., Icha, J., Horenburg, C., Muller, D., Norden, C., and Mansfeld, J.
(2018). Conditional control of fluorescent protein degradation by an auxin-
dependent nanobody. Nat Commun *9*, 3297.
16.   Hayashi, K., Neve, J., Hirose, M., Kuboki, A., Shimada, Y., Kepinski, S., and
Nozaki, H. (2012). Rational design of an auxin antagonist of the SCF(TIR1)
auxin receptor complex. ACS Chem Biol *7*, 590-598.
17.   Park, A., Won, S.T., Pentecost, M., Bartkowski, W., and Lee, B. (2014).
CRISPR/Cas9 allows efficient and complete knock-in of a destabilization
domain-tagged essential protein in a human cell line, allowing rapid
knockdown of protein function. PLoS One *9*, e95101.
18.   Ran, F.A., Hsu, P.D., Wright, J., Agarwala, V., Scott, D.A., and Zhang, F.
(2013). Genome engineering using the CRISPR-Cas9 system. Nat Protoc *8*,
2281-2308.
19.   Firestone, A.J., Weinger, J.S., Maldonado, M., Barlan, K., Langston, L.D.,
O'Donnell, M., Gelfand, V.I., Kapoor, T.M., and Chen, J.K. (2012). Small-
molecule inhibitors of the AAA+ ATPase motor cytoplasmic dynein. Nature
484, 125-129.

20.   Raaijmakers, J.A., Tanenbaum, M.E., and Medema, R.H. (2013). Systematic
dissection of dynein regulators in mitosis. J Cell Biol *201*, 201-215.
21.   Nabet, B., Roberts, J.M., Buckley, D.L., Paulk, J., Dastjerdi, S., Yang, A.,
Leggett, A.L., Erb, M.A., Lawlor, M.A., Souza, A., et al. (2018). The dTAG
system for immediate and target-specific protein degradation. Nat Chem Biol
14, 431-441.
22.   Banaszynski, L.A., Chen, L.C., Maynard-Smith, L.A., Ooi, A.G., and Wandless,
T.J. (2006). A rapid, reversible, and tunable method to regulate protein
function in living cells using synthetic small molecules. Cell *126*, 995-1004.
23.   Buckley, D.L., Raina, K., Darricarrere, N., Hines, J., Gustafson, J.L., Smith,
I.E., Miah, A.H., Harling, J.D., and Crews, C.M. (2015). HaloPROTACS: Use
of Small Molecule PROTACs to Induce Degradation of HaloTag Fusion
Proteins. ACS Chem Biol *10*, 1831-1837.
467

Figure 1

A

N-terminal tagging donor

B

C-terminal tagging donor

C

N-terminal tagging plasmid	Tag	Marker
pMK344	mAID	Hygro
pMK345	mAID-mClover	Hygro
pMK346	mAID-mCherry2	Hygro
pMK347	mAID	BSD
pMK348	mAID-mClover	BSD
pMK349	mAID-mCherry2	BSD
pMK375	Stag-3FLAG	Hygro
pMK376	mClover	Hygro
pMK377	mCherry2	Hygro
pMK401	mAID-NanoLuc	Hygro
pMK407	mAID-NanoLuc	BSD

C-terminal tagging plasmid	Tag	Marker
pMK277	mClover	Neo
pMK278	mClover	Hygro
pMK280	mCherry2	Neo
pMK281	mCherry2	Hygro
pMK283	Stag-3FLAG	Neo
pMK284	Stag-3FLAG	Hygro
pMK286	mAID	Neo
pMK287	mAID	Hygro
pMK289	mAID-mClover	Neo
pMK290	mAID-mClover	Hygro
pMK292	mAID-mCherry2	Neo
pMK293	mAID-mCherry2	Hygro
pMK388	mClover	BSD
pMK389	mCherry2	BSD
pMK390	Stag-3FLAG	BSD
pMK391	mAID	BSD
pMK392	mAID-mClover	BSD
pMK393	mAID-mCherry2	BSD
pMK394	mAID-Nluc	Neo
pMK408	mAID-Nluc	Hygro
pMK409	mAID-Nluc	BSD

D

AAVS1 introducing donor	Transgene	Marker
pMK232	CMV-OsTIR1	Puro
pMK243	Tet-OsTIR1	Puro
pMK364	CMV-OsTIR1	loxP-Puro-loxP
pMK365	Tet-OsTIR1	loxP-Puro-loxP

Figure 2

Figure 3

Figure 4

Figure 5

A

+ 500 µM IAA

IAA removal
+/- 200 µM auxinole

24 h

time-course sampling
(0, 1, 2, 3, 4, 5, and 6 h)

B

Figure 6