A nucleotide resolution map of Top2-linked DNA breaks in the yeast and human genome

William Gittens*, Dominic J. Johnson, Rachal M. Allison, Tim J. Cooper, Holly Thomas, and Matthew J Neale*

Genome Damage and Stability Centre, University of Sussex, UK, BN1 9RQ

*Correspondence to: m.neale@sussex.ac.uk, w.gittens@sussex.ac.uk

Abstract

DNA topoisomerases are required to resolve DNA topological stress. Despite this essential role, abortive topoisomerase activity generates aberrant protein-linked DNA breaks, jeopardising genome stability. Here, to understand the genomic distribution and mechanisms underpinning topoisomerase-induced DNA breaks, we map Top2 DNA cleavage with strand-specific nucleotide resolution across the *S. cerevisiae* and human genomes—and use the meiotic Spo11 protein to validate the broad applicability of this method to explore the role of diverse topoisomerase family members. Our data characterises Mre11-dependent repair in yeast, and defines two strikingly different fractions of Top2 activity in humans: tightly localised CTCF-proximal, and broadly distributed transcription-proximal, the latter correlated with gene length and expression. Moreover, single nucleotide accuracy enables us to reveal the influence primary DNA sequence has upon Top2 cleavage—distinguishing canonical DNA double-strand breaks (DSBs) from a major population of DNA single-strand breaks (SSBs) induced by etoposide (VP16) *in vivo*.

Introduction

DNA topoisomerases are a broad and ubiquitous family of enzymes that tackle topological constraints to replication, transcription, the maintenance of genome structure and chromosome segregation in mitosis and meiosis (Pommier et al., 2016). Although the specific mechanisms by which this is accomplished vary considerably across the family, key aspects are shared: including single or double-strand DNA cleavage to form a transient covalent complex (CC), which allows alteration of the topology of the nucleic acid substrate prior to religation (Wang, 2009). These processes are essential but carry with them a significant risk to genome stability because the CC may be stabilised as a permanent protein-linked DNA break by several physiological factors; such as the proximity of other

DNA lesions, the collision of transcription and replication complexes, denaturation of the topoisomerase, or by the binding of small molecules that inhibit religation (Reviewed in (Pommier et al., 2016; Nitiss, 2009)). Topoisomerase-induced DNA strand breaks have been proposed to constitute a significant fraction of the total damage to genomic DNA per day, and have been linked to the genesis and development of various cancers (Lin et al., 2009; Haffner et al., 2010), including a subset of therapy-related acute myeloid leukaemias (t-AML) caused by the use of Topoisomerase 2 (Top2) poisons in the chemotherapeutic treatment of primary cancers (Rowley and Olney, 2002; Cowell et al., 2012).

In *S. cerevisiae*, either Top1 or Top2 activity is sufficient to support transcription (Brill et al., 1987). However, whilst *top1Δ* cells are viable, Top2 is essential for sister chromatid segregation (Baxter and Diffley, 2008; DiNardo et al., 1984; Holm et al., 1985). In contrast to *S. cerevisiae*, all known vertebrate species encode two Top2 proteins (TOP2α and TOP2β). Interestingly, whilst TOP2α is essential for cellular proliferation—cells arrest in mitosis in its absence (Akimitsu et al., 2003)—TOP2β is not. Rather, TOP2β apparently plays an important role in promoting transcriptional programmes associated with neuronal development (Tiwari et al., 2012; Gómez-Herreros et al., 2014), a function that cannot be supported by TOP2α.

Accurate maps of the positions of topoisomerase-DNA covalent complexes (Top2 CCs) genome-wide and throughout the cell cycle provide insights into topological genome structural organisation, as well as providing the tools to study their repair should they become permanent DNA lesions. Whilst the dual catalytic sites present within the homodimeric Top2 enzyme suggest a primary role in DNA double-strand break (DSB) formation driven by a biological need for decatenation (Pommier et al., 2016), previous research has suggested that etoposide and other Top2 poisons may induce a population of DNA single-strand breaks (SSBs), due to independent inhibition of each active site (Wozniak and Ross, 1983; Long et al., 1986; Bromberg et al., 2003). Yet, despite the use of etoposide in chemotherapy, and the different toxicity of these two classes of DNA lesion, the prevalence of Top2-SSBs remains unclear.

More recently, general DSB mapping techniques have been applied to map etoposide-induced lesions (Canela et al., 2017; Yan et al., 2017). However, such methods are only able to map DSB ends, which therefore excludes and obscures etoposide-induced SSBs. Moreover, these methodologies lack specificity for protein-linked DNA ends, and require

nucleolytic processing to blunt 5' DNA termini as part of sample preparation (Canela et al.,
2017). When combined, these factors lead to loss of nucleotide resolution at the site of
Top2 cleavage.

To generate a more complete, and direct, picture of topoisomerase action across the
genome, we present here a technique for nucleotide resolution mapping of protein-linked
SSBs and DSBs (referred to collectively here as "Covalent Complexes"; CC), which we
demonstrate to be generally applicable in both yeast and human cell systems. We establish
with high confidence that the technique is able to map both the positions of Top2 CCs and
also CCs of DNA linked to the meiotic recombination protein Spo11, which is related to the
archaeal Topoisomerase VI (Bergerat et al., 1997). Furthermore, we provide insights into the
spatial distribution of human TOP2 CCs, including comparative analysis of their local
enrichment around transcription start sites (TSSs) and CTCF-binding motifs. We find that
TSS-proximal TOP2 CC levels are strongly correlated with transcription, a question that has
come under scrutiny recently (Canela et al., 2017). Finally, we present compelling evidence
that etoposide induces a majority of Top2-linked SSBs *in vivo*, corroborating previous
research and revealing, for the first time, that this is governed by primary DNA sequence at
the cleavage site.

Results

CC-seq enriches Spo11-linked DNA fragments

To elucidate the *in vivo* functions of topoisomerase-like enzymes, we set out to establish a
direct method (termed 'CC-seq') to enrich and map protein-DNA covalent complexes (CC)
genome-wide with nucleotide resolution. Silica fibre-based enrichment of CCs has been
used in the original identification of the meiotic DSB-inducer Spo11 (Keeney and Kleckner,
1995; Keeney et al., 1997). We first verified this enrichment principle using meiotic *sae2Δ*
cells, in which Spo11-linked DSBs accumulate at defined loci due to abrogation of the
nucleolytic pathway that releases Spo11 (Keeney and Kleckner, 1995). To demonstrate
specific enrichment of protein-linked molecules, genomic DNA from meiotic *S. cerevisiae*
cells was isolated in the absence of proteolysis, digested with *Pst*I restriction enzyme, and
isolated on glass-fibre spin columns (**Figure 1A, Methods**). We used eluted material to
assay a known Spo11-DSB hotspot by Southern blotting (**Figure 1B**). While DSB fragments
are a minor fraction of input material (~10% of total), and were absent in wash fractions,
DSBs accounted for >99% of eluted material, indicating ~1000-fold enrichment relative to
non-protein-linked DNA (**Figure 1B**).

CC-seq maps known Spo11-DSB hotspots genome-wide with high reproducibility

To generate a genome-wide map of Spo11-DSBs, genomic DNA from meiotic *sae2Δ* cells was sonicated to <400 bp in length, enriched upon the silica column, eluted, and ligated to DNA adapters in a two-step procedure that utilised the known phosphotyrosyl-unlinking activity of mammalian TDP2 to uncap the Spo11-bound end (**Figure 1A / Methods**) (Cortes Ledesma et al., 2009; Johnson et al., 2019). Libraries were paired-end sequenced and mapped to the *S. cerevisiae* reference genome (**Table S1**) alongside reads from a previous mapping technique ('Spo11-oligo-seq') that relies on the isolation of Spo11-linked oligonucleotides generated in wild-type cells during DSB repair (Pan et al., 2011). CC-seq revealed sharp, localised peaks ('hotspots') in *SPO11*+ cells that visually (**Figure 1C**) and quantitatively (**Figure S1A**, r=0.82) correlate with Spo11-oligo seq, and that are absent in the *spo11-Y135F* control strain in which Spo11-DSBs do not form (Bergerat et al., 1997), demonstrating that signal detected by CC-seq accurately reflects the distribution of Spo11 cleavages. Moreover, CC-seq biological replicates were highly correlated (**Figure S1B**; r = 0.98), demonstrating high reproducibility. Finally, when analysed at nucleotide resolution, CC-seq revealed high correlation (r = 0.85) between the frequency of Watson-mapping and Crick-mapping cleavage sites, when offset by a single bp (**Figure 1D and S1C**). This correlation is expected because of the 2 bp 5' overhang generated at Spo11-DSBs *in vivo* (**Fig1D**; e.g. (Liu et al., 1995))—demonstrating the nucleotide resolution accuracy of CC-seq, which is further supported by our observation of nucleotide composition preference at the site of cleavage (**Figure 1E**).

CC-seq enriches and maps Top2-linked DNA fragments from *S. cerevisiae*

Confident in the specificity of CC-seq to map meiotic protein-linked DSBs with high resolution and dynamic range, we employed CC-seq to characterise within *S. cerevisiae* the covalent complexes generated naturally by Topoisomerase 2 (Top2) that become stabilised upon exposure to etoposide, and are thus a proxy for Top2 catalytic activity (Minocha and Long, 1984; Chen et al., 1984). Because *S. cerevisiae* is relatively insensitive to etoposide, we utilised strains ('*pdr1Δ*') in which the action of major drug export pathways are downregulated due to modulation of *PDR1* activity (Stepanov et al., 2008). As expected, *pdr1Δ* dramatically increased cellular sensitivity to etoposide, which was further enhanced by mutation of *SAE2* and *MRE11* (**Figure 2A**), factors involved in the repair of covalent protein-linked DNA breaks (Neale et al., 2005; Cannavo et al., 2018; Hoa et al., 2016; Hartsuiker et al., 2009).

Next, we prepared sequencing libraries from mitotically growing etoposide-treated control,

sae2Δ, and *mre11Δ S. cerevisiae pdr1Δ* cells (**Table S2**). A human DNA spike-in was

included to enable calibration of relative signal between strains (**Methods**). Replicate

libraries displayed high reproducibility (r > 0.89; **Figures S2A-C**), so were pooled. In

untreated control cells, weak Top2 CC signal was distributed homogeneously across the

genome, with few strong peaks (**Figure 2B**). Upon etoposide treatment, sharp single-

nucleotide peaks arose at similar locations in all strains (**Figure 2B and S2D-E**), but with

increased amplitude in the repair mutants—thereby directly linking the MRX pathway to

repair of Top2 CCs in *S. cerevisiae*.

Global analyses revealed Top2 activity in *S. cerevisiae* to be relatively enriched in divergent

and tandem intergenic regions (IGRs) and depleted in intragenic and convergent IGRs

(**Figure 2B and C**), similar to, but less pronounced than, the patterns of Spo11 (**Figure 2B**

and C). Such connections to global gene organisation are likely driven by the need for

Spo11 and Top2 to interact with the DNA helix—an interpretation underpinned by an

anticorrelation with nucleosome occupancy (**Figure S1D and S3A**). Nevertheless, the fact

that *S. cerevisiae* Top2 CC signal is not correlated with proximal gene expression—neither

in untreated cells, nor following etoposide exposure (**Figure S3B**)—suggests that in the

gene-dense *S. cerevisiae* genome topological stress may be dealt with by Top2 at sites

dislocated from where it is generated. By contrast, Spo11 activity is positively correlated

with gene expression (**Figure S1E**), perhaps due to the influence transcription has upon the

generation of higher order chromatin loop structures (Schalbetter et al., 2018), which in turn

are known to pattern Spo11-DSBs (Sun et al., 2015).

Interestingly, Top2 CC signal remained weakly correlated over greater distances than

Spo11 (**Figure 2D**), suggesting localised 5-10 kb wide regions of enriched activity—much

larger than Spo11-DSB hotspots (generally <500 bp in size; **Figure 1C**; (Pan et al., 2011))—

suggesting factors other than just nucleosome occupancy influence local Top2 catalysis.

Top2 CCs are strongly correlated between Watson and Crick strands when offset by 3 bp

(**Figure 2E and S2F-K**)—as expected for the 4 bp overhang generated by Top2-DSBs—

further demonstrating the nucleotide resolution accuracy of CC-seq. Together these data

describe how CC-seq reveals a strand-specific nucleotide resolution map of etoposide-

induced Top2 CCs.

CC-seq maps TOP2 CCs in the Human genome

Having demonstrated that CC-seq is applicable for mapping two divergent types of topoisomerase-like covalent complexes in *S. cerevisiae*, we applied CC-seq to map TOP2 activity in a human cell system. Asynchronous, sub-confluent RPE-1 cells were treated with etoposide (VP16) in the presence of MG132 proteasome inhibitor, in order to limit potential proteolytic degradation of TOP2 CCs that might otherwise hamper enrichment (**Figure 1A**). Slot-blotting and immunodetection with anti-TOP2β antibody demonstrated complete recovery of input TOP2β-linked CCs within the column elution, with no TOP2β remaining in the flow through or being removed by the washes (**Figure 3A**). Eluted material was used to generate sequencing libraries from three replicate control (–VP16) and four replicate etoposide-treated samples (+VP16), and high-depth paired-end sequencing reads (**Table S2**) were aligned to the human genome (hg19; **Methods**).

Replicates displayed high correlation in the distribution of TOP2 CC signal at broad scale (r values ≥ 0.79; **Figure S4A-B**), and so were pooled. Visual inspection revealed that -VP16 and +VP16 signals are spatially-correlated (**Figure 3B**), but that +VP16 signal intensity is less uniform, with more dynamic range (**Figure S4C-D**). These differences are likely due to higher signal-to-noise enabled by enrichment of TOP2 CC following VP16 treatment. Like yeast Top2 libraries (**Figure 2E**), nucleotide resolution analysis of human CC-seq libraries displayed a skew towards Watson-Crick read-pairs that are offset by 3 bp, as expected for the 4 bp 5' overhang generated by TOP2 (**Figure S4E**). This skew is much greater in VP16-treated samples (**Figure S4E**), supporting the view that +VP16 libraries have greater Top2 CC enrichment.

The human genome encodes two type-IIA topoisomerases, TOP2α and TOP2β (**Introduction**). To demonstrate the specificity of CC-seq to map human TOP2 activity we used CRISPR-Cas9 to generate a human RPE-1-derived cell line with homozygous knockout mutations in the non-essential *TOP2B* gene, and arrested cells in G1 (**Figure S5A**) when TOP2α is not expressed (Heck et al., 1988; Woessner et al., 1991). TOP2α and TOP2β expression was undetectable in such G1-arrested cells (**Figure S5B**), and importantly, VP16-induced ɣ-H2AX foci were reduced ~7-fold relative to wild type control lines, indicating that most of the signal is TOP2β-dependent (**Figure S5C-D**).

Next, we applied CC-seq to asynchronous and G1-arrested wild type and *TOP2B*⁻ᐟ⁻ cells. In G1, VP16 exposure induced localised regions of enriched TOP2 CC formation in wild type

cells that were largely diminished or absent in *TOP2B*$^{-/-}$ cells (**Figure S5E**). By comparison,

TOP2β deletion did not prevent VP16-induced signal in asynchronous cells (**Figure S5E**), in

which TOP2α is still present. Together, these results indicate that CC-seq detects a mixture

of both TOP2α and TOP2β covalent complexes in wild type human cells depending on the

cell cycle phase.

Spatial distribution of Top2 activity in the human genome

The human genome is divided into a relatively gene-rich 'A' compartment and a relatively

gene-poor 'B' compartment, the 3D spatial segregation of which can be determined by Hi-

C (Lieberman-Aiden et al., 2009). In both untreated and VP16-treated asynchronous wild

type cells, TOP2 CC are enriched in the active A compartment (**Figure 3B and C**),

consistent with the role of TOP2 in facilitating transcription. TOP2 activity also correlated

with regions of negative DNA supercoiling (Naughton et al., 2013) (**Figure 3B**). These

observations provide a functional link between TOP2 activity, DNA topological stress, and

large-scale chromatin compartments.

We next focused our attention in more detail on the genome-wide pattern of TOP2 activity

enriched by VP16 treatment. Simple visual inspection (**Figure 3D**) suggested that TOP2 CC

signals are enriched around genome features previously identified as regions of high TOP2-

binding and etoposide-induced DNA strand breakage, including sites of CTCF-binding

(Uuskülä-Reimand et al., 2016; Canela et al., 2017) and active transcription start sites

(TSSs) marked by H3K4Me3 and H3K27Ac (Baranello et al., 2014; Yang et al., 2015).

Genomic maps of etoposide-induced DNA breaks were recently generated using a related

methodology, 'END-seq' (Canela et al., 2017), that is not specific for protein-linked TOP2

CC. Aggregating END-seq data around strong CC-seq positions revealed concordant peak

signals at broad scale (**Figure 3E**), demonstrating a general agreement in the positions

reported by the two techniques. However, at finer scale, END-seq positions are offset by 1-

15 bp in the 3' direction relative to CC-seq positions on each strand (**Figure 3F**)—

something not observed when aggregating CC-seq data around the same loci (**Figure 3F**).

This important difference is likely to be explained by the different populations of DNA

breaks mapped by each technique: specific TOP2 cleavage positions (CC-seq) versus DNA

breaks that have undergone limited nucleolytic resection (END-seq).

To investigate the relationship between CTCF and TOP2 activity, we filtered CTCF motifs to
include only those that overlapped an RPE-1 CTCF ChIP-seq peak (**Methods;** (Akimitsu et
al., 2003)). After aligning Watson and Crick motifs in the same orientation, we frequently
observed a strong TOP2 CC peak upstream of the motif centre and accompanied by more
distal, weaker peaks on both sides of the motif (**Figure 4A and S6A**). Loci containing
multiple CTCF-binding motifs display more complex TOP2 CC patterns (**Figure 4A and**
S6B) with enrichment on both sides of the double motif, as would be expected from the
aggregation of a heterogeneous population of CTCF and TOP2 activity present across
different cells.

We next stratified CTCF motifs into quantiles based on proximal CTCF binding intensity,
and aggregated CC-seq signals centred on these loci (**Figure 4A**). Most Top2 CC signal is
concentrated in two peaks located ±~54 bp relative to the centre of the CTCF motif, with
the upstream peak ~2-fold more intense (**Fig 4A, middle and bottom panels**). Remaining
Top2 CC signal is focused in a flanking array of weaker peaks. This signal pattern has been
reported recently for VP16-induced DSBs (Canela et al., 2017), and has been suggested to
result from a role for TOP2 in facilitating genome organisation by cohesin-dependent loop
extrusion operating in the context of CTCF and other boundary elements. In support of this
interpretation, total aggregated TOP2 CC signal across these regions is positively
correlated with CTCF occupancy as measured by ChIP-seq (**Figure 4A**); interestingly
however, TOP2 CCs were not correlated with interaction strength of associated loops
(**Figure S6C**), annotated previously by Hi-C (Darrow et al., 2016). The positions of TOP2 CC
surrounding CTCF motifs anticorrelate with the well-positioned nucleosomes at these sites
(**Figure 4B**). However, such anticorrelation with nucleosome positioning is not a unique
feature of CTCF loci, but is instead a general property of all TOP2 CC sites (**Figure 4C**),
consistent with maps of both Top2 CC (**Figure S3B**) and Spo11 (**Figure S1E;** (Pan et al.,
2011)) in *S. cerevisiae*.

In human cells, etoposide-induced TOP2 CCs are also enriched around gene TSSs (**Figure**
5), typically concentrated in broad peaks ~1-2 Kbp downstream and ~0.5-1 Kbp upstream
of the TSS (**Figure 5A**; (Yang et al., 2015)). Similar to elsewhere in the genome, in the region
immediately downstream of the TSS (0-400 bp), TOP2 CC signal anticorrelated with
nucleosome occupancy (**Figure S6D**). Stratifying TSSs based on gene expression
microarray data (Ganem et al., 2014) revealed a strong positive correlation (**Figure 5A**)—
something not identified when prior etoposide-induced DSB maps were compared to

nascent transcription as measured by GRO-Seq (Canela et al., 2017). Thus, while human

TOP2 activity is correlated with total transcription of the gene, it is not proportional to local

RNA polymerase activity when assayed at finer scale—suggesting a model where, as

postulated for *S. cerevisiae* (**Figure S3A**), superhelical stress is resolved at sites dislocated

from the process that generates them. Indeed, when analysed with an identical pipeline,

both TSS-proximal TOP2 CCs (**Figure 5A**) and etoposide-induced DSBs (Canela et al.,

2017) positively correlate with gene expression (**Figure S6E**), indicating that this correlation

is not due to technical differences in mapping methodologies. Based on the twin-domain

model of DNA supercoiling, topological stress is expected to increase not just with gene

expression level, but also with increasing gene length (Liu and Wang, 1987). In support of

this idea, we observe a strong positive correlation between human TOP2 CC enrichment

and gene length independent from gene expression (**Figure S7A**). Interestingly, no such

correlation with gene length is present in *S. cerevisiae* (**Figure S7B**), further supporting a

model whereby, in this compact gene-dense genome, Top2 activity in the IGR is neither

directly linked to expression nor length of the proximal gene.

Notably, the increased density of etoposide-induced human TOP2 CCs at strongly

expressed genes is 2.7-fold greater than at strongly bound CTCF motifs (**Figure 5B**

and C). Thus, whilst strongly-bound CTCF motifs (14,924) greatly outnumber active genes

(4,194)—leading, globally, to more TOP2 activity in total at CTCF sites (**Figure 5D**)—our

data reinforce the important role of topoisomerase activity during transcription.

Local DNA sequence and methylation status direct the formation of Top2 CCs

Previous reports suggest that etoposide induces a majority of SSBs, rather than DSBs, due

to independent poisoning of each active site in the Top2 homodimer (Bromberg et al.,

2003). In our yeast and human datasets, we similarly observe more single-strand Top2 CCs

(90.1% and 94.2%, respectively) than double-strand Top2 CCs (9.9% and 5.8%,

respectively) following etoposide exposure (**Figure 6A and S3C**).

Prior, fine-scale mapping of Top2 cleavage at specific substrates indicated that cleavage

patterns are influenced by local DNA sequence composition (Pommier et al., 1991;

Strumberg et al., 1999). To investigate the generality of such findings, and to precisely

determine how DNA sequence influences Top2 site preference across the yeast and human

genome, we computed the nucleotide resolution average base composition around Top2

CC SSB and DSB sites (**Fig 6B**)—something made possible by the positional accuracy of

315 our mapping technique. For DSBs, average DNA sequence patterns are rotationally

316 symmetrical around the dyad axis of cleavage (**Figure 6B, left)**, consistent with the known

317 homodimeric nature of eukaryotic Top2 (Pommier et al., 2016). Base positions with notable

318 skews are restricted to the region ±11 bp from the dyad axis ($p<10^{-4}$, Chi-squared test),

319 consistent with the DNA contacts made by Top2 (Lee et al., 1989; Thomsen et al., 1990;

320 Wu et al., 2011), yet with minor, species-specific differences at a few positions (±2, ±7, ±9;

321 **Figure 6C, top**), suggesting subtle differences in the ScTop2 and HsTOP2 DNA binding

322 surface. Positions -3 and +3 are the bases immediately 5' to the scissile phosphodiester

323 bonds cleaved by Top2 (**Figure 6B**). In the averaged DSB sequence motif, strong

324 preference for cytosine at these positions (69.0% of DSBs) on the scissile strand (**Figure**

325 **6B, left)** agrees with *in vitro* studies of Top2 mechanism—attributed to favourable base-

326 stacking interactions of etoposide with guanosine at the -1 position on the non-scissile

327 strand (Pommier et al., 1991; Strumberg et al., 1999; Wu et al., 2011).

329 Importantly, whilst SSBs display similar overall symmetrical sequence features as DSBs,

330 SSBs differed substantially at position +3, where the strong preference for guanine on the

331 scissile strand (and paucity of cytosine on the non-scissile strand) is absent (**Figure 6B,**

332 **right, and Figure 6C, bottom**). Furthermore, the preference for cytosine at the -3 position

333 on the scissile strand increases to 88.1% for SSBs. This clear difference in sequence bias

334 provides strong evidence that such positions truly reflect sites of preferred VP16-induced

335 SSBs vs DSBs *in vivo*.

337 The human genome is subject to methylation of cytosine (meC) at CpG sites. Dramatically,

338 meC completely abolishes the formation of VP16-induced TOP2 CCs (**Figure 6D**),

339 suggesting this bulky group may interfere sterically with the VP16-Top2-DNA ternary

340 complex, modifying TOP2 catalysis. Collectively, these observations demonstrate that local

341 DNA sequence composition—including epigenetic marks—are major determinants of both

342 TOP2 CC abundance and the differential generation of SSBs and DSBs by VP16 *in vivo*.

Discussion

346 Topoisomerases are ubiquitous gatekeepers to the genetic material—facilitating changes to

347 DNA topology, nuclear organisation, and chromosome segregation that are essential for the

348 generation and propagation of all life on Earth. Yet, topoisomerase dysfunction can also

349 prove harmful, generating difficult to repair protein-linked DNA breaks that create risks to

genome stability. Thus, understanding how, where, and when topoisomerase activity takes place is of great importance. Here, we demonstrate a robust, technically simple and sensitive method to map sites of preferred type IIA topoisomerase activity across the eukaryotic genomes of yeast and human cells at strand-specific nucleotide resolution. Our methodology joins an extensive toolkit of complementary NGS solutions (Canela et al., 2017; Crosetto et al., 2013; Yan et al., 2017), each with its own set of unique advantages. However, many of these techniques do not achieve nucleotide resolution, either due to the use of nucleolytic processing to blunt 5' DNA termini (Canela et al., 2017), and/or inability to directly map DNA breaks covalently linked to protein (Canela et al., 2017; Crosetto et al., 2013; Yan et al., 2017). Here, by utilising the tyrosyl phosphodiesterase activity of recombinant Tdp2 to unblock 5'-DNA termini non-nucleolytically, we are able to accurately map the position of not just Top2 CCs, but also related classes of topoisomerase-like enzymes such as Spo11, which acts uniquely in meiotic cells. Importantly, such specificity in our cleavage maps enable us to corroborate and extend—with completely independent methodology—prior research findings from meiosis (Pan et al., 2011), the impact that VP16 has on SSB versus DSB formation—and for the first time demonstrate the influence that primary DNA sequence composition and methylation status has on the process of Top2 catalysis *in vivo* in both yeast and human cells.

Top2 activity has long been associated with transcriptional activity (Yang et al., 2015; Brill et al., 1987; Baranello et al., 2014; Yu et al., 2017). In addition, recent research has identified etoposide-induced DSBs that colocalise with a subpopulation of chromatin-bound TOP2 on the loop-external side of CTCF- and Rad21-bound CTCF motifs (Canela et al., 2017; Uusküla-Reimand et al., 2016). TOP2 activity at these sites has been proposed to facilitate genome organisation driven by chromatin loop extrusion, and thereby to constitute a major fraction of the total nuclear TOP2 activity (Canela et al., 2017). Moreover, contrary to expectations, the frequency of etoposide-induced DSBs was found to be locally uncorrelated with nascent RNA levels (Canela et al., 2017)—generating some confusion in the understanding of where TOP2 activity is most prevalent and most important. Here, our independent CC-seq method and analyses permit us to revisit this question in explicit detail, and clarify the role of TOP2 in these processes. Importantly, whilst Top2 activity is certainly highly enriched at CTCF-bound genomic loci, the total amount of activity at any given CTCF locus is many times lower than observed at transcriptionally active genes. Nevertheless, it is critical to emphasise that because of the great abundance of CTCF-

bound sites compared to active genes, total aggregated TOP2 activity is globally similar for
CTCF-associated versus transcriptionally active regions of the genome.

Our data also permit us to investigate how other genomic features influence Top2 activity.
For example, at both TSS and CTCF-proximal regions—and indeed in the wider genome of
both yeast and human—Top2 cleavages are highly enriched in nucleosome-free DNA, in
support of prior findings made at a subset of genomic loci (Capranico et al., 1990; Udvardy
and Schedl, 1991; Käs and Laemmli, 1992). Notably, whilst yeast Top2 activity is also
enriched near TSSs, it is concentrated immediately (120-140 bp) upstream, unlike human
TOP2 activity, which is primarily concentrated in a broader peak 1-2 kb downstream, within
the gene body. Furthermore, unlike human TOP2, yeast Top2 activity is uncorrelated with
downstream gene expression. These differences may stem from the very different scale
and transcriptional organisation of the yeast and human genomes, and the strong
patterning of Top2 activity driven by accessibility to nucleosome-free intergenic promoter
regions in *S. cerevisiae*. In support of this interpretation, similar nucleosomal constraints
appear to substantially govern the activity of the related topoisomerase-like enzyme,
Spo11, during meiosis (Pan et al., 2011) (**Figure S1** herein).

Despite these differences in distribution of Top2 activity at this larger scale, the DNA
sequence preferences for yeast and human Top2 are similar, implying substantial
conservation of the DNA binding interface, which is supported by analysis of the published
crystal structure of VP16-TOP2-DNA (Wu et al., 2011). A nucleotide resolution genome-
wide map of *E. coli* gyrase activity was recently reported (Sutormin et al., 2018)), revealing a
sequence motif composed of a central 35 bp region containing the cleavage site, plus two
flanking 47 bp regions with ~10.5 bp periodicity centred ~±40 bp from the site of cleavage,
which are attributed to the known wrapping of proximal DNA around the gyrase C-terminal
domains (Reece and Maxwell, 1991). Strikingly, whilst we observe a core central motif for
Top2, there are no such flanking motifs over these ranges, in agreement with its lack of
DNA wrapping domains (**Figure S3D**). We do, however, observe two more-proximal
flanking regions centred ~±20 bp from the site of cleavage, each with ~10.5 bp periodicity
(**Figure S3E**), suggesting that Top2 also bends its DNA substrate—in agreement with a
recent biophysical study (Huang et al., 2017).

We have used CC-seq to probe MRX-dependent repair of Top2 CCs in *S. cerevisiae*,
corroborating and extending prior genetic experiments in this organism (Hamilton and

Maizels, 2010; Stepanov et al., 2008), and others (Hoa et al., 2016; Hartsuiker et al., 2009).

Importantly, calibration of our signal using an internal human standard enables the relative

quantification of Top2 CCs, and thus directly demonstrates for the first time that loss of

Mre11 activity increases cellular levels of Top2 CCs in *S. cerevisiae*. The antigen-

independent enrichment step allows calibration to be performed without genetic epitope-

tagging or reconstitution of recombinant protein-DNA complexes, making CC-seq ideally

suited to the future study of repair dynamics.

We have demonstrated that CC-seq maps CCs of both human TOP2α and TOP2β—

proteins that are conserved amongst all vertebrates and whose functional differences

remain a subject of interest within the field (Austin et al., 2018). Given the essential

requirement for TOP2α during mitosis (Akimitsu et al., 2003), and the apparent inability of

TOP2β to compensate for its loss (Grue et al., 1998), it will be particularly interesting to

determine the genomic distribution of TOP2α/TOP2β activities throughout the cell cycle,

and to relate this to 3D chromatin conformation measured by Hi-C. Whilst not essential for

cell proliferation, TOP2β plays an important role in promoting transcriptional programmes

associated with neuronal development (Tiwari et al., 2012), and this function cannot be

supported by TOP2α. Furthermore, loss of TDP2 function inhibits TOP2-dependent gene

transcription and leads to neurological symptoms including intellectual disability, seizures

and ataxia (Gómez-Herreros et al., 2014). Thus, accurate genomic maps of TOP2β CC

formation in these tissues will help to identify those genes most pertinent to the

development of neurodegeneration. Moreover, as demonstrated by our Spo11 maps, the

antigen-independent, high-efficiency protein-DNA enrichment process makes CC-seq

generally applicable for mapping not just Top2 activity in diverse organisms, but also for

mapping similar types of topoisomerase-DNA covalent complexes, or even those less

distinct DNA-protein crosslinks that arise upon ablation of the DNA-dependent protease

SPRTN/Wss1 (Stingele et al., 2015).

Acknowledgements

448 We thank Keith Caldecott, Antony Oliver and Peter Hornyak for sharing recombinant TDP2,

449 and Jon Nitiss for sharing the *pdr1Δ* strains.

Author Contributions

452 W.G., D.J. and M.J.N. conceived the project and developed the CC-seq methodology.

453 W.G., D.J., H.T. and R.A. generated whole genome CC-seq libraries. W.G. performed all

454 data processing and analysis, and all human cell work. D.J., H.T. and R.A. performed yeast
455 sensitivity and meiotic DSB assays. T.J.C. developed the mapping pipeline and associated
456 tools. W.G. and M.J.N. interpreted the observations and wrote the manuscript.
457

Competing interests

459 The authors declare no competing financial interests.
460

Funding statement

462 W.G., D.J., R.A., T.J.C., and M.J.N. were supported by an ERC Consolidator Grant
463 (#311336), the BBSRC (#BB/M010279/1) and the Wellcome Trust (#200843/Z/16/Z).
464

Data availability

466 All strains and cell lines listed in Table S3 and S4 are available on request. Sequencing
467 reads are available via the NCBI Sequence Read Archive (accession numbers pending).
468

Figure Legends

Figure 1. CC-seq maps covalent Spo11-linked DNA breaks in *S. cerevisiae* meiosis with nucleotide accuracy

A) Schematic of the CC-seq method.

B) Column-based enrichment of Spo11-linked DNA fragments detected by Southern blotting at the *his4*::*LEU2* recombination hotspot (**Methods**). Arrows indicate expected sizes of Spo11-DSBs.

C) Nucleotide resolution mapping of *S. cerevisiae* Spo11 hotspots by CC-seq or oligo-seq (Pan et al., 2011). Red and blue traces indicate Spo11-linked 5' DNA termini on the Watson and Crick strands, respectively. Grey arrows indicate positions of gene open reading frames.

D) Pearson correlation (r) of Spo11 CC-seq signal between Watson and Crick strands, offset by the indicated distances.

E) Average nucleotide composition over a 200 (top) and 30 bp (bottom) window centred on Spo11 breaks. Bases reported are for the top strand only. HpM = Hits per million mapped reads per base pair.

Figure 2. CC-seq maps covalent Top2-linked DNA breaks in *S. cerevisiae* cycling cells with nucleotide accuracy

A) Serial dilution spot tests of VP16 tolerance for the indicated strains.

B) Nucleotide resolution mapping of *S. cerevisiae* Top2 CCs by CC-seq of the indicated strains after treatment for 4 hours with 1 mM VP16. Spo11-CC data is plotted for comparison. Top2 CC data were calibrated using a human DNA spike-in (**Methods**). Red and blue traces indicate CC-linked 5' DNA termini on the Watson and Crick strands, respectively. Grey arrows indicate positions of gene open reading frames. Lower panels show an expanded view of the region from 31.5 to 40 Kbp.

C) Quantification of Top2 and Spo11 CC signal stratified by genomic region. The genome was divided into intra and intergenic regions; the intergenic region was further divided into divergent, tandem and convergent based on orientation of flanking genes. Spo11 and Top2 activity mapped by CC-seq is expressed as box-and-whisker plots of density (upper and lower box limit: 3rd and 1st quartile; bar: median; upper and lower whisker: highest and lowest values within 1.5-fold of the interquartile range), or as the percentage of total mapped reads.

D) Local correlation of Top2 or Spo11 CC-seq signals. Top2 or Spo11 CC-seq data were binned at 50 bp resolution and the Pearson correlation calculated between bins of increasing separation.

E) Pearson correlation (r) of Top2 CC-seq signal on Watson and Crick strands, offset by the indicated distance. HpM = Hits per million mapped reads per base pair.

Figure 3. CC-seq maps TOP2-linked DNA breaks in Human cells with nucleotide accuracy

A) Anti-TOP2β western slot blot of input, flow through, wash, and elution fractions. RPE-1 cells were treated or not with VP16, prior to processing according to **Figure 1A**.

B) Broad-scale maps of *H. sapiens* TOP2 CCs produced by CC-seq in RPE-1 cells ±VP16. Raw data were scaled, binned, smoothed and median subtracted prior to plotting (**Methods**). Chromatin compartments revealed by Hi-C eigenvector analysis (Darrow et al., 2016), and supercoiling revealed by bTMP ChIP-seq (Naughton et al., 2013) are shown for comparison.

C) Quantification of TOP2 CC in chromatin compartments A and B. Data are expressed as box-and-whisker plots of density as for **Figure 2C**.

D) Fine-scale mapping of *H. sapiens* TOP2 CCs by CC-seq in RPE-1 cells ±VP16. Red and blue traces indicate TOP2-linked 5' DNA termini on the Watson and Crick strands, respectively. Pale shaded areas are the same data smoothed according to local density (see **Methods**). RPE-1 CTCF and H3K4Me3 ChIP-seq data plus H3K27Ac ChIP-seq data overlaid from seven cell lines (ENCODE, 2012) is shown for comparison.

E) Medium-scale aggregate of END-seq mapped DSBs (Canela et al., 2017) surrounding nucleotide resolution CC-seq mapped TOP2 CCs.

F) Fine-scale strand-specific aggregate of END-seq mapped DSB ends (red and blue bars) and nucleotide resolution CC-seq mapped TOP2 CCs (grey bars) surrounding strong TOP2 CC sites.

Figure 4. CTCF-proximal TOP2 activity is tightly confined within nucleosome-depleted regions

A) Aggregation of TOP2 CCs in a 1 Kbp window centred on orientated CTCF motifs in human RPE-1 cells. Four example CTCF loci are shown orientated in the 5'-3' direction (top). A heatmap of all CTCF-motifs in the human genome, with 25 rows stratified by the strength of colocalising CTCF ChIP-seq peaks in RPE-1 cells (middle). The colour scale indicates average TOP2 CC density. Motifs are also stratified into 4 quartiles of CTCF-binding, and the average TOP2 CC distribution in each quartile plotted (bottom).

B) Aggregated TOP2 CC distribution (red line) in the highest quartile of CTCF-binding compared with the average MNase-seq signal (blue line).

C) Aggregated TOP2 CC distribution (black and red lines: single-nucleotide resolution and smoothed, respectively) and the average MNase-seq signal (blue line) surrounding strong TOP2 CC sites.

In (B) and (C) peaks in MNase-seq signal indicate inferred nucleosome positions (blue circles).

Figure 5. TSS-proximal TOP2 activity is strongly correlated with gene transcription

A) Aggregation of TOP2 CCs in a 10 Kbp window centred on orientated TSSs in human RPE-1 cells. Four example TSSs are shown orientated in the 5'-3' direction (top). A heatmap of all TSSs in the human genome, with 25 rows stratified by gene expression level in RPE-1 cells (middle). The colour scale indicates average TOP2 CC density. Motifs are also stratified into 4 quartiles of gene expression, and the average TOP2 CC distribution in each quartile plotted (bottom).

B) Comparison of CTCF-proximal and TSS-proximal TOP2 CC distributions in the highest quartile of CTCF-binding (pink) and gene-expression (blue), respectively.

C) Average TOP2 CC density in 10 Kbp regions centred on the highest quartile of TSS, CTCF, or

regions where both features are present. Data are expressed as box-and-whisker plots of density as

in **Figure 2C**. Statistical significance was determined using the KS test.

D) As in (C) but sum total TOP2 CCs found in these regions.

Figure 6. Local DNA sequence and methylation status direct the formation of Top2 CCs

A) The percentage of strong sites that are SSBs and DSBs in etoposide-treated *S. cerevisiae* and

human cells.

B) Average nucleotide composition over a 30 bp window centred on the DSB or (inferred) SSB dyad

axis, in etoposide-treated *S. cerevisiae* and human cells. Values reported are for the top strand only.

C) Heatmaps describing pairwise similarity of nucleotide composition patterns shown in (B). Rows 1-

4 are the absolute differences between: yeast and human SSB patterns, yeast and human DSB

patterns, yeast SSB and DSB patterns, and human SSB and DSB patterns, respectively.

D) The average number of TOP2 CCs at the +1 position relative to methylated and unmethylated

cytosines in human RPE-1 cells, ±VP16. Statistical significance was determined using KS test.

References

562
Akimitsu, N., Adachi, N., Hirai, H., Hossain, M. S., Hamamoto, H., Kobayashi, M., Aratani, Y.,
Koyama, H., and Sekimizu, K. (2003). Enforced cytokinesis without complete nuclear division in
embryonic cells depleting the activity of DNA topoisomerase IIalpha. Genes Cells *8*, 393-402.
Austin, C. A., Lee, K. C., Swan, R. L., Khazeem, M. M., Manville, C. M., Cridland, P., Treumann, A.,
Porter, A., Morris, N. J., and Cowell, I. G. (2018). TOP2B: The First Thirty Years. Int J Mol Sci *19*,
Baranello, L., Kouzine, F., Wojtowicz, D., Cui, K., Przytycka, T. M., Zhao, K., and Levens, D. (2014).
DNA break mapping reveals topoisomerase II activity genome-wide. Int J Mol Sci *15*, 13111-13122.
Baxter, J., and Diffley, J. F. (2008). Topoisomerase II inactivation prevents the completion of DNA
replication in budding yeast. Mol Cell *30*, 790-802.
Bergerat, A., de Massy, B., Gadelle, D., Varoutas, P. C., Nicolas, A., and Forterre, P. (1997). An
atypical topoisomerase II from Archaea with implications for meiotic recombination. Nature *386*, 414-
417.
Brill, S. J., DiNardo, S., Voelkel-Meiman, K., and Sternglanz, R. (1987). Need for DNA topoisomerase
activity as a swivel for DNA replication for transcription of ribosomal RNA. Nature *326*, 414-416.
Bromberg, K. D., Burgin, A. B., and Osheroff, N. (2003). A two-drug model for etoposide action
against human topoisomerase IIalpha. J Biol Chem *278*, 7406-7412.
Canela, A., Maman, Y., Jung, S., Wong, N., Callen, E., Day, A., Kieffer-Kwon, K.-R., Pekowska, A.,
Zhang, H., Rao, S. S. P., Huang, S.-C., Mckinnon, P. J., Aplan, P. D., Pommier, Y., Aiden, E. L.,
Casellas, R., and Nussenzweig, A. (2017). Genome Organization Drives Chromosome Fragility. Cell
170, 507-521.e18.
Cannavo, E., Johnson, D., Andres, S. N., Kissling, V. M., Reinert, J. K., Garcia, V., Erie, D. A., Hess,
D., Thomä, N. H., Enchev, R. I., Peter, M., Williams, R. S., Neale, M. J., and Cejka, P. (2018).
Regulatory control of DNA end resection by Sae2 phosphorylation. Nat Commun *9*, 4016.
Capranico, G., Jaxel, C., Roberge, M., Kohn, K. W., and Pommier, Y. (1990). Nucleosome positioning
as a critical determinant for the DNA cleavage sites of mammalian DNA topoisomerase II in
reconstituted simian virus 40 chromatin. Nucleic Acids Res *18*, 4553-4559.
Chen, G. L., Yang, L., Rowe, T. C., Halligan, B. D., Tewey, K. M., and Liu, L. F. (1984).
Nonintercalative antitumor drugs interfere with the breakage-reunion reaction of mammalian DNA
topoisomerase II. J Biol Chem *259*, 13560-13566.
Cortes Ledesma, F., El Khamisy, S. F., Zuma, M. C., Osborn, K., and Caldecott, K. W. (2009). A
human 5'-tyrosyl DNA phosphodiesterase that repairs topoisomerase-mediated DNA damage. Nature
461, 674-678.

Cowell, I. G., Sondka, Z., Smith, K., Lee, K. C., Manville, C. M., Sidorczuk-Lesthuruge, M., Rance, H. A., Padget, K., Jackson, G. H., Adachi, N., and Austin, C. A. (2012). Model for MLL translocations in therapy-related leukemia involving topoisomerase IIβ-mediated DNA strand breaks and gene proximity. Proc Natl Acad Sci U S A *109*, 8989-8994.

Crosetto, N., Mitra, A., Silva, M. J., Bienko, M., Dojer, N., Wang, Q., Karaca, E., Chiarle, R., Skrzypczak, M., Ginalski, K., Pasero, P., Rowicka, M., and Dikic, I. (2013). Nucleotide-resolution DNA double-strand break mapping by next-generation sequencing. Nat Methods *10*, 361-365.

Darrow, E. M., Huntley, M. H., Dudchenko, O., Stamenova, E. K., Durand, N. C., Sun, Z., Huang, S. C., Sanborn, A. L., Machol, I., Shamim, M., Seberg, A. P., Lander, E. S., Chadwick, B. P., and Aiden, E. L. (2016). Deletion of DXZ4 on the human inactive X chromosome alters higher-order genome architecture. Proc Natl Acad Sci U S A *113*, E4504-12.

DiNardo, S., Voelkel, K., and Sternglanz, R. (1984). DNA topoisomerase II mutant of Saccharomyces cerevisiae: topoisomerase II is required for segregation of daughter molecules at the termination of DNA replication. Proc Natl Acad Sci U S A *81*, 2616-2620.

ENCODE, P. C. (2012). An integrated encyclopedia of DNA elements in the human genome. Nature *489*, 57-74.

Ganem, N. J., Cornils, H., Chiu, S. Y., O'Rourke, K. P., Arnaud, J., Yimlamai, D., Théry, M., Camargo, F. D., and Pellman, D. (2014). Cytokinesis failure triggers hippo tumor suppressor pathway activation. Cell *158*, 833-848.

Gómez-Herreros, F., Schuurs-Hoeijmakers, J. H., McCormack, M., Greally, M. T., Rulten, S., Romero-Granados, R., Counihan, T. J., Chaila, E., Conroy, J., Ennis, S., Delanty, N., Cortés-Ledesma, F., de Brouwer, A. P., Cavalleri, G. L., El-Khamisy, S. F., de Vries, B. B., and Caldecott, K. W. (2014). TDP2 protects transcription from abortive topoisomerase activity and is required for normal neural function. Nat Genet *46*, 516-521.

Grue, P., Grässer, A., Sehested, M., Jensen, P. B., Uhse, A., Straub, T., Ness, W., and Boege, F. (1998). Essential mitotic functions of DNA topoisomerase IIalpha are not adopted by topoisomerase IIbeta in human H69 cells. J Biol Chem *273*, 33660-33666.

Haffner, M. C., Aryee, M. J., Toubaji, A., Esopi, D. M., Albadine, R., Gurel, B., Isaacs, W. B., Bova, G. S., Liu, W., Xu, J., Meeker, A. K., Netto, G., De Marzo, A. M., Nelson, W. G., and Yegnasubramanian, S. (2010). Androgen-induced TOP2B-mediated double-strand breaks and prostate cancer gene rearrangements. Nat Genet *42*, 668-675.

Hamilton, N. K., and Maizels, N. (2010). MRE11 function in response to topoisomerase poisons is independent of its function in double-strand break repair in Saccharomyces cerevisiae. PLoS One *5*, e15387.

Hartsuiker, E., Neale, M. J., and Carr, A. M. (2009). Distinct requirements for the Rad32(Mre11) nuclease and Ctp1(CtIP) in the removal of covalently bound topoisomerase I and II from DNA. Mol Cell *33*, 117-123.

Heck, M. M., Hittelman, W. N., and Earnshaw, W. C. (1988). Differential expression of DNA topoisomerases I and II during the eukaryotic cell cycle. Proc Natl Acad Sci U S A *85*, 1086-1090.

Hoa, N. N., Shimizu, T., Zhou, Z. W., Wang, Z. Q., Deshpande, R. A., Paull, T. T., Akter, S., Tsuda, M., Furuta, R., Tsusui, K., Takeda, S., and Sasanuma, H. (2016). Mre11 Is Essential for the Removal of Lethal Topoisomerase 2 Covalent Cleavage Complexes. Mol Cell *64*, 580-592.

Holm, C., Goto, T., Wang, J. C., and Botstein, D. (1985). DNA topoisomerase II is required at the time of mitosis in yeast. Cell *41*, 553-563.

Huang, W. C., Lee, C. Y., and Hsieh, T. S. (2017). Single-molecule Förster resonance energy transfer (FRET) analysis discloses the dynamics of the DNA-topoisomerase II (Top2) interaction in the presence of TOP2-targeting agents. J Biol Chem *292*, 12589-12598.

Johnson, D., Allison, R. M., Cannavo, E., Cejka, P., and Neale, M. (2019). Removal of Spo11 from meiotic DNA breaks in vitro but not in vivo by Tyrosyl DNA Phosphodiesterase 2. bioRxiv *http://dx.doi.org/10.1101/527333*,

Käs, E., and Laemmli, U. K. (1992). In vivo topoisomerase II cleavage of the Drosophila histone and satellite III repeats: DNA sequence and structural characteristics. EMBO J *11*, 705-716.

Keeney, S., Giroux, C. N., and Kleckner, N. (1997). Meiosis-specific DNA double-strand breaks are catalyzed by Spo11, a member of a widely conserved protein family. Cell *88*, 375-384.

Keeney, S., and Kleckner, N. (1995). Covalent protein-DNA complexes at the 5' strand termini of meiosis-specific double-strand breaks in yeast. Proc Natl Acad Sci U S A *92*, 11274-11278.

Lee, M. P., Sander, M., and Hsieh, T. (1989). Nuclease protection by Drosophila DNA topoisomerase II. Enzyme/DNA contacts at the strong topoisomerase II cleavage sites. J Biol Chem *264*, 21779-21787.

Lieberman-Aiden, E., van Berkum, N. L., Williams, L., Imakaev, M., Ragoczy, T., Telling, A., Amit, I., Lajoie, B. R., Sabo, P. J., Dorschner, M. O., Sandstrom, R., Bernstein, B., Bender, M. A., Groudine, M., Gnirke, A., Stamatoyannopoulos, J., Mirny, L. A., Lander, E. S., and Dekker, J. (2009). Comprehensive mapping of long-range interactions reveals folding principles of the human genome. Science *326*, 289-293.

Lin, C., Yang, L., Tanasa, B., Hutt, K., Ju, B. G., Ohgi, K., Zhang, J., Rose, D. W., Fu, X. D., Glass, C. K., and Rosenfeld, M. G. (2009). Nuclear receptor-induced chromosomal proximity and DNA breaks underlie specific translocations in cancer. Cell *139*, 1069-1083.

Liu, J., Wu, T. C., and Lichten, M. (1995). The location and structure of double-strand DNA breaks induced during yeast meiosis: evidence for a covalently linked DNA-protein intermediate. EMBO J *14*, 4599-4608.

Liu, L. F., and Wang, J. C. (1987). Supercoiling of the DNA template during transcription. Proc Natl Acad Sci U S A *84*, 7024-7027.

Long, B. H., Musial, S. T., and Brattain, M. G. (1986). DNA breakage in human lung carcinoma cells and nuclei that are naturally sensitive or resistant to etoposide and teniposide. Cancer Res *46*, 3809-3816.

Minocha, A., and Long, B. H. (1984). Inhibition of the DNA catenation activity of type II topoisomerase by VP16-213 and VM26. Biochem Biophys Res Commun *122*, 165-170.

Naughton, C., Avlonitis, N., Corless, S., Prendergast, J. G., Mati, I. K., Eijk, P. P., Cockroft, S. L., Bradley, M., Ylstra, B., and Gilbert, N. (2013). Transcription forms and remodels supercoiling domains unfolding large-scale chromatin structures. Nat Struct Mol Biol *20*, 387-395.

Neale, M. J., Pan, J., and Keeney, S. (2005). Endonucleolytic processing of covalent protein-linked DNA double-strand breaks. Nature *436*, 1053-1057.

Nitiss, J. L. (2009). Targeting DNA topoisomerase II in cancer chemotherapy. Nat Rev Cancer *9*, 338-350.

Pan, J., Sasaki, M., Kniewel, R., Murakami, H., Blitzblau, H. G., Tischfield, S. E., Zhu, X., Neale, M. J., Jasin, M., Socci, N. D., Hochwagen, A., and Keeney, S. (2011). A Hierarchical Combination of Factors Shapes the Genome-wide Topography of Yeast Meiotic Recombination Initiation. Cell *144*, 719-731.

Pommier, Y., Capranico, G., Orr, A., and Kohn, K. W. (1991). Local base sequence preferences for DNA cleavage by mammalian topoisomerase II in the presence of amsacrine or teniposide. Nucleic Acids Res *19*, 5973-5980.

Pommier, Y., Sun, Y., Huang, S. N., and Nitiss, J. L. (2016). Roles of eukaryotic topoisomerases in transcription, replication and genomic stability. Nat Rev Mol Cell Biol *17*, 703-721.

Reece, R. J., and Maxwell, A. (1991). DNA gyrase: structure and function. Crit Rev Biochem Mol Biol *26*, 335-375.

Rowley, J. D., and Olney, H. J. (2002). International workshop on the relationship of prior therapy to balanced chromosome aberrations in therapy-related myelodysplastic syndromes and acute leukemia: overview report. Genes Chromosomes Cancer *33*, 331-345.

Schalbetter, S. A., Fudenberg, G., Baxter, J., Pollard, K. S., and Neale, M. J. (2018). Principles of Meiotic Chromosome Assembly. bioRXiv *http://dx.doi.org/10.1101/442038*,

Stepanov, A., Nitiss, K. C., Neale, G., and Nitiss, J. L. (2008). Enhancing drug accumulation in Saccharomyces cerevisiae by repression of pleiotropic drug resistance genes with chimeric transcription repressors. Mol Pharmacol *74*, 423-431.

Stingele, J., Habermann, B., and Jentsch, S. (2015). DNA-protein crosslink repair: proteases as DNA repair enzymes. Trends Biochem Sci *40*, 67-71.

Strumberg, D., Nitiss, J. L., Dong, J., Kohn, K. W., and Pommier, Y. (1999). Molecular analysis of yeast and human type II topoisomerases. Enzyme-DNA and drug interactions. J Biol Chem *274*, 28246-28255.

Sun, X., Huang, L., Markowitz, T. E., Blitzblau, H. G., Chen, D., Klein, F., and Hochwagen, A. (2015). Transcription dynamically patterns the meiotic chromosome-axis interface. Elife *4*,

Sutormin, D., Rubanova, N., Logacheva, M., Ghilarov, D., and Severinov, K. (2018). Single-nucleotide-resolution mapping of DNA gyrase cleavage sites across the Escherichia coli genome. Nucleic Acids Res

Thomsen, B., Bendixen, C., Lund, K., Andersen, A. H., Sørensen, B. S., and Westergaard, O. (1990). Characterization of the interaction between topoisomerase II and DNA by transcriptional footprinting. J Mol Biol *215*, 237-244.

Tiwari, V. K., Burger, L., Nikoletopoulou, V., Deogracias, R., Thakurela, S., Wirbelauer, C., Kaut, J., Terranova, R., Hoerner, L., Mielke, C., Boege, F., Murr, R., Peters, A. H., Barde, Y. A., and

Schübeler, D. (2012). Target genes of Topoisomerase IIβ regulate neuronal survival and are defined by their chromatin state. Proc Natl Acad Sci U S A *109*, E934-43.

Udvardy, A., and Schedl, P. (1991). Chromatin structure, not DNA sequence specificity, is the primary determinant of topoisomerase II sites of action in vivo. Mol Cell Biol *11*, 4973-4984.

Uusküla-Reimand, L., Hou, H., Samavarchi-Tehrani, P., Rudan, M. V., Liang, M., Medina-Rivera, A., Mohammed, H., Schmidt, D., Schwalie, P., Young, E. J., Reimand, J., Hadjur, S., Gingras, A. C., and Wilson, M. D. (2016). Topoisomerase II beta interacts with cohesin and CTCF at topological domain borders. Genome Biol *17*, 182.

Wang, J. C. (2009). Untangling the double helix Cold Spring Harbor Laboratory Pr).

Woessner, R. D., Mattern, M. R., Mirabelli, C. K., Johnson, R. K., and Drake, F. H. (1991). Proliferation- and cell cycle-dependent differences in expression of the 170 kilodalton and 180 kilodalton forms of topoisomerase II in NIH-3T3 cells. Cell Growth Differ *2*, 209-214.

Wozniak, A. J., and Ross, W. E. (1983). DNA damage as a basis for 4'-demethylepipodophyllotoxin-9-(4,6-O-ethylidene-beta-D-glucopyranoside) (etoposide) cytotoxicity. Cancer Res *43*, 120-124.

Wu, C. C., Li, T. K., Farh, L., Lin, L. Y., Lin, T. S., Yu, Y. J., Yen, T. J., Chiang, C. W., and Chan, N. L. (2011). Structural basis of type II topoisomerase inhibition by the anticancer drug etoposide. Science *333*, 459-462.

Yan, W. X., Mirzazadeh, R., Garnerone, S., Scott, D., Schneider, M. W., Kallas, T., Custodio, J., Wernersson, E., Li, Y., Gao, L., Federova, Y., Zetsche, B., Zhang, F., Bienko, M., and Crosetto, N. (2017). BLISS is a versatile and quantitative method for genome-wide profiling of DNA double-strand breaks. Nat Commun *8*, 15058.

Yang, F., Kemp, C. J., and Henikoff, S. (2015). Anthracyclines induce double-strand DNA breaks at active gene promoters. Mutat Res *773*, 9-15.

Yu, X., Davenport, J. W., Urtishak, K. A., Carillo, M. L., Gosai, S. J., Kolaris, C. P., Byl, J. A. W., Rappaport, E. F., Osheroff, N., Gregory, B. D., and Felix, C. A. (2017). Genome-wide TOP2A DNA cleavage is biased toward translocated and highly transcribed loci. Genome Res *27*, 1238-1249.

Methods

Yeast strains, culture methods and treatment

The *Saccharomyces cerevisiae* yeast strains used in this study are described in **Table 2** and
were derived using standard genetic techniques. Strains used for Spo11 mapping are
isogenic to the SK1 subtype and carry the *sae2Δ::kanMX* gene disruption allele. Strains
used for Top2 mapping are isogenic to the BY4741 subtype and carry the *pdr1DBD-CYC8*
drug sensitivity cassette (Stepanov et al., 2008). For Spo11 DSB mapping, cells were
induced to undergo synchronous meiosis as follows: Cells were grown overnight to
saturation in 4 ml YPD medium (1% yeast extract, 2% peptone, 2% glucose supplemented
with 0.5 mM adenine and 0.4 mM uracil) at 30 ℃, then diluted to OD_{600} 0.2 in 200 ml YPA
medium (1% yeast extract, 2% peptone, 1% potassium acetate) and grown vigorously for
15 h at 30 ℃. Cells were then washed with water, resuspended in 200 ml sporulation
medium (2% potassium acetate supplemented with diluted amino acids), incubated
vigorously for 6 h at 30 ℃ and harvested by centrifugation. For etoposide treatment, cells
were grown overnight to saturation in 4 ml YPD medium at 30 ℃, then diluted to OD_{600} 0.5
in 100 ml YPD and grown until OD_{600} 2. 50 ml cultures were then incubated for a further 4 h
in the presence of either 1 mM etoposide or 2% DMSO, before harvesting by
centrifugation.

Human cell lines, culture methods and treatment

The human cell lines used in this study are described (**Supplementary Table 3**). Human
hTERT RPE-1 cells were obtained from ATCC and cultured at 37 ℃, 5% CO_2 and 3% O_2 in
Dulbecco's Modified Eagle's Medium DMEM/F-12 (Sigma), supplemented with 10% Fetal
Calf Serum (FCS) and 90 Units/ml Penicillin-Streptomycin. For all experiments (CC-seq,
Slot Blot, WB, IF, FACs) with asynchronous cell populations, RPE-1 cells were seeded at a
density of (3.5×10^3 cells/cm²l) and incubated for 72 h at 37 ℃, to ensure subconfluent log-
phase growth (~70% confluency) at the time of the experiment. For G1 cell populations,
WT and *TOP2B*[-/-] RPE-1 cells were seeded at a density of 5×10^3 cells/cm² and incubated for
48 h at 37 ℃ in DMEM/F12 containing 10% FCS, prior to a further 24 h incubation in
DMEM/F12 medium containing no FCS to ensure complete G1-phase arrest (verified by
FACs, see below). For experiments with proteasome inhibitor and etoposide (CC-seq, Slot
Blot), cells were preincubated with 5 μM MG132 (Sigma) for 90 min, trypsinised and
incubated in suspension with 5 μM MG132 and 100 μM etoposide (Sigma) for 20 min at 37

℃. For experiments with etoposide alone (IF), adherent cells were treated with 100 µM
etoposide for 20 min at 37 ℃.

Generation of *TOP2B*^{-/-} RPE-1 cells

The oligonucleotides 5'-CACCGCCGCAGCCACCCGACT and 5'-
AAACAGTCGGGTGGCTGCGGC (identified using Benchling; https://benchling.com) were
annealed and cloned into pX330 following BbsI restriction, as described previously (Hsu et
al., 2013). This SpCas9/ trugRNA co-expression plasmid was transiently expressed in RPE-
1 cells to target the 17 bp target sequence GCCGCAGCCACCCGACT (TGG) within exon 1
of *TOP2B*. Single clones were trypsinised and passaged to isolated culture vessels prior to
screening for absent protein by Western Blot (WB). All experiments involving *TOP2B*^{-/-} were
conducted using the RPE-1 clone T2B/6, in which no TOP2β is detectable by WB or IF.

Spot tests of chronic etoposide sensitivity

Single colonies of each *S. cerevisiae* strain were incubated overnight in 4 ml YPD at 30 ℃
with shaking. 0.1 ml of this starter was used to inoculate 4 ml YPD, prior to incubation for 5
hours at 30 ℃. Cultures were diluted to make a stock with an OD_{600} of 2.0, then this was
10-fold serially diluted five times. Each dilution was spotted onto plates containing 0, 0.1,
0.3 or 1.0 mM VP16, prior to incubation for three days at 30 ℃. Plates were imaged on a
2400 Photo scanner (Epson).

Southern blotting of meiotic Spo11 DSBs

Approximately 2 µg of genomic DNA (isolated by non-proteolysing Phenol-Chloroform
extraction, as described below) was digested at 37 °C overnight using *Pst*I restriction
enzyme (NEB) in NEBuffer 3.1 (100 mM NaCl, 50 mM Tris Base·HCl pH 7.9, 10 mM MgCl₂,
100 µg ml-1 BSA). Additional *Pst*I was added for 4 hours before the addition of NEB purple
loading dye to 1×. Digested samples were proteolysed using 1 mg ml-1 Proteinase K
(Sigma) at 60 ℃ for 30 minutes, left to reach room temperature before 10 µg was loaded on
a 0.7% 1× TAE agarose gel (40 mM Tris Base·HCl, 20 mM glacial acetic acid, 1 mM EDTA
pH 8.0) containing 50 µg ml-1 ethidium bromide. DNA was separated in 1× TAE at 60 V for
18 hours. The gel was imaged using InGenius (Syngene) bioimaging system to check
migration and then exposed to 180 mJ/m² UV in the Stratalinker (Stratagene). The gel was
then soaked in three times its volume of denaturation solution (0.5 M NaOH, 1.5 M NaCl) for
30 minutes and then transferred to Zetaprobe (Bio-Rad) membrane by means of a vacuum
at 55 mBar for 2 hours. After transfer the membrane was washed in water ten times and

then cross-linked by exposing the membrane to 120 mJ/m^2 UV in the Stratalinker. The

membrane was incubated in 30 ml of hybridisation buffer (0.5 M NaHPO$_4$ buffer pH 7.5, 7%

SDS, 1 mM EDTA, 1% BSA) at 65 ℃ for 1 hour. The *MXR2* probe for looking at the

HIS4::LEU2 locus was created from 50 ng of template DNA, 0.1 ng of Lambda DNA (NEB)

digested with *BstE*II (NEB), and water. The mix was denatured at 100 °C for 5 minutes then

put on ice. High Prime (Roche) was added in addition to 0.5-3 mBq of α-^{32}P dCTP and

incubated at 37 °C for 15 minutes. 30 µl 1× TE was added and the probe spun through a G-

50 spin column (GE Healthcare) at 400 × *g* for 2 minutes. The probe was then denatured by

incubating at 100 °C for 5 minutes and then put on ice before being added to 20 ml

hybridisation mixture. The original 30 ml hybridisation buffer was discarded and the 20 ml

containing the probe was added to the membrane and incubated overnight at 65 ℃. After

incubation, the membrane was washed five times with 100 ml pre-warmed Southern wash

buffer (1% SDS, 40 mM NaHPO$_4$ buffer pH 7.5, 1 mM EDTA) and exposed to phosphor

screen overnight.

Western blotting

Whole human cell extracts (WCE) were harvested by direct lysis in 1x Laemmli loading

buffer, denatured for 10 min at 95 ℃ and sonicated for 30 s using Bioruptor® Pico.

Samples were subjected to SDS-PAGE (7% or gradient gel) and transferred to

nitrocellulose membrane. Primary immunodetection was with antibodies targeting TOP2β

(Clone 40, BD Biosciences), TOP2α (ab52934, Abcam), or Ku80 (ab80592, Abcam).

Secondary immunodetection was with HRP-conjugated Rabbit anti-Mouse IgG

(ThermoFisher), prior to detection of peroxidase activity using ECL reagent and X-Ray film

(Scientific Laboratory Supplies Ltd).

Slot blotting

Samples were diluted 4-fold (500 uL total volume) in NaPO$_4$ buffer (25 mM, pH 6.5), and slot

blotted onto 0.2 µM nitrocellulose membrane (Amersham), using the Minifold I (Whatman)

manifold. The wells were washed twice with 750 µL NaPO$_4$ buffer. The membrane was then

blocked with 10% milk-TBST for 1 h at RT, prior to incubation overnight with anti-TOP2β

antibody (Clone 40, BD Biosciences) at 4 ℃. The membrane was washed 4 times with

TBST, incubated with HRP-conjugated Rabbit anti-Mouse IgG (ThermoFisher) for 1 h at RT,

washed 4 times with TBST, and incubated with ECL detection reagent for 1 min. X-Ray film

was used for detection.

Fluorescence-Assisted Cell Sorting (FACS)

Approximately 10 million RPE-1 cells were trypsinised, washed once in PBS and resuspended in 1.5 ml PBS. 3.5 ml ethanol was added dropwise to pellet, with vortexing. Cells were fixed for 1 hr at 4 °C, prior to centrifugation and aspiration of the supernatant. Cells were washed twice with PBS, prior to resuspension in 0.5 ml 0.25% Triton-X100-PBS for 15 min on ice. Cells were pelleted by centrifugation, supernatant was aspirated, and the pellet was resuspended in 0.5 ml TBS containing 10 ug/ml RNase A (Sigma) and 167 nM Sytox Green (ThermoFisher). After 30 min incubation in the dark at RT, the suspension was filtered through fine mesh into test tubes. DNA content in 50,000 cells was analysed using the Accuri C6 (BD Biosciences), with gating to exclude doublets and cell debris.

Immunofluorescence

Cells were seeded onto glass coverslips (Agar Scientific), incubated and treated according to the protocol outlined above. Cells were then fixed with 4% paraformaldehyde PBS for 10 min, washed three times with PBS, permeabilised with 0.2% Triton-X100-PBS for 10 min, blocked for 1 h with 10% FCS-PBS, incubated with primary antibodies targeting Phospho-Histone H2AX (S139) (JBW-301, Merck Millipore) and TOP2α (ab52934, Abcam), washed three times with PBS, incubated with secondary antibodies Alexa 488-conjugated Goat anti-mouse IgG (Fisher) and Alexa 647-conjugated Goat anti-Rabbit (Fisher), washed three times with PBS, washed once with distilled water, and mounted with VECTASHIELD containing DAPI (Vector Laboratories).

High-content microscopy

Automated wide-field microscopy was performed on an Olympus ScanR system (motorised IX83 microscope) with ScanR Image Acquisition and Analysis Software, 40x/0.6 (LUCPLFLN 40x PH) dry objectives and Hamamatsu ORCA-R2 digital CCD camera C10600. Numbers of anti-phospho-Histone H2AX (S139) foci (Alexa 488; FITC filter) were quantified in the nuclear region colocalising with DAPI, using Olympus ScanR Analysis software. TOP2α signal (Alexa 647; Cy5 filter) was also quantified in this region.

CC-seq: Enrichment of protein-linked DNA

1.5×10^7 RPE-1 cells were treated as described above, pelleted by centrifugation, washed once with 15 ml ice-cold PBS, and resuspended in three aliquots of 400 µL ice-cold PBS. 1×10^9 yeast cells were treated as described above, then spheroplasted in 1.5 ml spheroplasting buffer (1 M sorbitol, 50 mM NaHPO$_4$ buffer pH 7.2, 10 mM EDTA) containing

200 µg/ml Zymolyase 100T (AMS Biotech) and 1% β-mercaptoethanol (Sigma) for 20 min at

37 ˚C. 2 µL Protease Inhibitor Cocktail and 2 µL Pefabloc (Sigma) were added before

splitting into five aliquots of 400 µL. All subsequent steps of the protocol are the same for

yeast and human samples. 1 ml ice-cold ethanol was added to 400 µL cell suspensions in

microcentrifuge tubes, mixed, incubated for 10 min on ice, and pelleted by centrifugation.

The supernatant was thoroughly removed by aspiration, prior to addition of 200 µL 1x STE

buffer (2% SDS, 0.5M Tris pH 8.1, 10 mM EDTA, 0.05% bromophenol blue), cell disruption

using a pestle (VWR), addition of a further 400 µL 1x STE buffer, and incubation for 10 min

at 65 ˚C. Samples were cooled on ice and 500 µL Phenol-Chloroform-isoamyl alcohol

(25:24:1; Sigma) was added. The mixtures were emulsified by shaking and pipetting 5 times

with a 1 ml micropipette, prior to phase separation by centrifugation at 20,000 g for 20 min.

By minimising mechanical shearing of the lysate prior to phenol chloroform extraction,

peptides that are covalently linked to high molecular weight DNA segregate into the

aqueous phase. 500 µL of the aqueous phase was removed to a clean microcentrifuge tube

and nucleic acids were precipitated with 1 ml ice-cold ethanol, pelleted by centrifugation,

washed with ice-cold 70% ethanol, and dissolved in TE buffer overnight at 4 ˚C. Samples

were then incubated with 0.2 mg/ml RNase A (Sigma) for 1 h at 37 ˚C; nucleic acids were

precipitated with 1 ml ethanol, pelleted by centrifugation, washed twice with 70% ethanol

and dissolved in TE overnight. Aliquots were combined to a total of 1 ml and sonicated to

an average fragment size of 300-400 bp with Covaris (duty cycle: 10%, intensity/peak

power incidence: 75W, cycles/burst: 200, time: 15 min). 1 ml of sonicated sample was

added to 1.2 ml of binding buffer (10 mM Tris pH 8.1, 10 mM EDTA, 0.66 M NaCl, 0.22%

SDS, 0.44% N-Lauroylsarcosine sodium salt). Each sample was divided over several

Miniprep (QIAGEN) silica-fibre membrane spin columns, such that the total DNA loaded to

each was approximately 20 µg. The flowthrough was reapplied to the column to improve

binding. Columns were washed 6 times with 600 µL of TEN (10 mM Tris, 1 mM EDTA, 0.3 M

NaCl) per 1 min wash, prior to elution with 100 µL TES (10 mM Tris, 1 mM EDTA, 0.5%

SDS).

CC-seq: DNA end repair and adapter ligation

Eluted products were pooled to 500 µL in TES and incubated with 1 mg/ml Proteinase K

(Sigma) for 30 min at 60 ˚C, prior to overnight ethanol precipitation at -80 ˚C with 1.41 ml

ethanol, 0.2 mg/ml glycogen and 200 mM NaOAc. The DNA-glycogen precipitate was

pelleted by centrifugation at 20,000 g for 1 hr at 4 ˚C, washed once with 1.5 ml 70%

ethanol, and re-pelleted by centrifugation. The supernatant was aspirated and the pellet

was air-dried for 10 min at RT, prior to solubilisation in 52 µL 10 mM Tris-HCl. DNA

concentration was measured in a 2 µL sample with the Qubit (ThermoFisher) and High

Sensitivity reagents. The remaining 50 µL was used as input for one round of end repair

and adapter ligation with NEBNext Ultra II DNA Library Preparation kit (NEB), according to

manufacturer's instructions, except for the use of a custom P7 adapter (see table). The use

of custom adapters is to allow differentiation of the sheared end (P7 adapter) from the

Top2/Spo11 end (P5 adapter). After ligation of the P7 adapter, DNA was isolated with

AMPure XP beads (Beckman Coulter) according to manufacturer's instructions (beads:input

of 78:90) and eluted in 50 µL 10 mM Tris-HCl. Samples were diluted 2-fold with 50 µL Tdp2

reaction buffer (100 mM TrisOAc, 100 mM NaOAc, 2 mM MgOAc, 2 mM DTT, 200 µg/ml

BSA) and incubated with 3 µL of 10 µM recombinant human TDP2 (Hornyak et al., 2016;

Johnson et al., 2019), for 1 h at 37 ℃. DNA was isolated again with AMPure XP beads

(beads:input of 103:103) and eluted in 52 µL 10 mM Tris-HCl. Next, a second round of end

repair and adapter ligation was conducted, using adapter P5 (see table). After ligation of

the P5 adapter to the Top2/Spo11-cleaved end, DNA was isolated with AMPure XP beads

(beads:input of 78:90) and eluted in 17 µL 10 mM Tris-HCl.

CC-seq: PCR and size selection

DNA concentration was measured in 2 µL using the Qubit. The remaining 15 µL was used

as template for the PCR step of the NEBNext Ultra II PCR step using universal primer (P5

end) and indexed primers for multiplexing (P7 end), according to manufacturer's

instructions. PCR reactions were diluted with 50 µL 10 mM Tris-HCl, DNA was isolated with

AMPure XP beads (beads:input of 84:100) and eluted in 30 µL 1 mM Tris-HCl pH 8.1.

Samples were then subjected to 200-600 bp size selection using the BluePippin (Sage

Science), prior to quantification of molarity using the Bioanalyzer (Agilent).

CC-seq: NGS and data pipeline

Multiplexed library pools were sequenced on the Illumina MiSeq (Kit v3 - 150 cycles) or

Illumina NextSeq 500 (Kit v2 - 75 cycles), with paired-end read lengths of 75 or 42 bp,

respectively. Paired end reads that passed filter were aligned using bowtie2 (options: -X

1000 --no-discordant --very-sensitive --mp 5,1 --np 0), using MAPQ0 settings for yeast or

MAPQ10 settings for human experiments, then SAM files processed by the custom-built

Perl program termMapper that computes the coordinates of the protein-linked 5'-terminal

nucleotide. The reference genomes used in this study are hg19 (human), and Cer3H4L2 (*S.*

cerevisiae), which we generated by inclusion of the *his4*::*LEU2* and *leu2*::*hisG* loci into the

Cer3 yeast genome build. Yeast data sets were filtered to exclude long terminal repeats, retrotransposons, telomeres, and the rDNA. Human datasets were filtered to remove known ultra-high signal regions (Hoffman et al., 2013; ENCODE, 2012) and repeat regions (ENCODE, 2012). All subsequent analyses were performed in R (Version 3.4.3) using RStudio (Version 1.1.383), unless indicated otherwise.

CC-seq: Calibrated library generation

Calibration of Top2 CC-seq experiments was conducted to allow comparison of relative peak intensities in different yeast strains. This was achieved by spike-in of human DNA following the sonication stage of the protocol, which is a method that has been used to calibrate other sequencing methods (Hu et al., 2015; Grzybowski et al., 2015). S. cerevisiae and RPE-1 cells were exposed to etoposide and processed until just after sonication, exactly as according to the cell treatment and CC-seq protocols above. DNA concentration was quantified by Qubit, and then mixed at a molar ratio of human DNA:yeast DNA of 1:100. All subsequent stages of the protocol were identical, except read alignment, for which we used both hg19 and Cer3H4L2 builds successively. Cer3H4L2-aligned peak heights were corrected in each sample by multiplying by the reciprocal fraction of human reads in that sample.

CC-seq: Fine-scale mapping

Fine-scale (nucleotide resolution) maps of Spo11/Top2 CCs were produced as simple histograms over a specified region (Figures 1B, 2B, 3D, 4A, 5A, S1A, S3A, S5A and S5B). Dark red and blue line heights indicate numbers of 5'-terminal nucleotides detected at that position on the Watson and Crick strands, in units of HpM. Where indicated in the figure caption, smoothed data are also plotted as pale red and blue polygons, in addition to the unsmoothed nucleotide resolution data. This smoothing was either applied using a sliding Hann window of the indicated width, or using a custom smoothing function (VarX), as indicated in the figure caption.

CC-seq: Broad-scale mapping

Broad-scale maps of Top2 CCs were produced by binning nucleotide resolution data at 10 Kbp (Figure 3B) or 100 Kbp (Figure S4D) resolution. Binned data were either plotted directly (Figure S4D); or first scaled according to the estimated noise fraction (see below), smoothed with a 10-bin Hanning window, and median subtracted (Figure 3B).

CC-seq: Estimation of the noise fraction in Human datasets

The noise fraction in each sample was estimated using an adaptation of the previously

published NCIS method (Liang and Keleş, 2012). Briefly: the data for -VP16 and +VP16

samples were first binned at 10 Kbp resolution. Then the subpopulation of bins with the

lowest TOP2 CC signal was identified in each sample. The average signal density of this

subpopulation of bins, in each sample, was defined as the noise density (d_{-VP16} and d_{+VP16}).

Signal in the -VP16 sample was scaled by a normalisation factor equal to:

993
$$r = \frac{d_{+VP16}}{d_{-VP16}}$$

CC-seq: Quantification of Spo11 and Top2 activity in defined genomic regions

Yeast Spo11 hotspots were the same as defined previously (Pan et al., 2011). Yeast

intra/intergenomic regions were defined based on TSS and TTS coordinates reported on

the Saccharomyces Genome Database (https://www.yeastgenome.org). Human chromatin

compartments A and B were defined based on eigenvector analysis of previously published

100 kb resolution RPE-1 Hi-C data (Darrow et al., 2016) using the Juicer package

(Lieberman-Aiden et al., 2009; Durand et al., 2016). Human "CTCF" and "TSS" regions were

defined as 10 kb regions centred on the top quartile of expressed TSSs or the midpoint of

top quartile CTCF-bound CTCF-binding motifs, with overlaps merged using

GenomicFeatures::reduce. The "Both" region was defined as the intersection of the "TSS"

and "CTCF" regions using GenomicFeatures::intersect. This region was then excluded from

the individual "TSS" and "CTCF" regions. The "Neither" region was defined as regions of

hg19 which overlapped neither "TSS", nor "CTCF", nor "Both", using

GenomicFeatures::gaps. Nucleotide resolution data was tallied within defined regions, and

is expressed as an aggregate (bar plots) in which the signal is corrected using the average

signal in the "Neither" compartment, and/or as the distribution of region signal densities

(box-and-whisker plot), as indicated in figure captions.

1012

CC-seq: Correlation of Watson/Crick cleavage positions

Nucleotide resolution human TOP2 data was thresholded at 0.01 HpM. Nucleotide

resolution yeast Top2 and Spo11 data were not thresholded. Peak coordinates on the Crick

strand were offset over the range -100 to +100, relative to Watson coordinates. After each

offset, the data was filtered to include only sites with both Watson and Crick hits. The

Pearson correlation (r) between the Watson and Crick signal intensities in these n sites was

calculated. We also counted the fraction of reads found within these n sites, and

normalised this number over the -100 to +100 bp range. Data are expressed as R values

and/or normHpM values, as indicated in the figure captions.

1022

1023   **CC-seq: Aggregation of CC-seq data around loci of interest**

1024   Nucleotide resolution CC-seq data (line-plots) or binned CC-seq data (heatmaps) were

1025   aggregated within regions of specified size, centred on the loci of interest. The resulting

1026   sum total HpM was divided by numbers of loci to give a mean HpM per locus.

1027

CC-seq: Loci of interest used in this study

Yeast TSS and TTS coordinates were obtained from the Saccharomyces Genome

Database https://www.yeastgenome.org. These were stratified based on associated gene

length, expression level in vegetative SK1 (GSM907178, GSM907179, GSM907180), or

expression level in meiotic SK1 (GSM907176, GSM907177; (Dominissini et al., 2012)).

Human TSS and TTS coordinates were obtained from the UCSC hg19 knownGene

annotation, and stratified based on associated gene length, or expression level

(GSM1395252, GSM1395253, GSM1395254; (Ganem et al., 2014)). Occupied RPE-1 CTCF

motifs were identified as follows: The FIMO tool (Grant et al., 2011) and the CTCF Position

Weight Matrix (PWM) from the JASPAR database (Mathelier et al., 2014) were used to find

all significant hg19 CTCF motifs ($p < 1\times10^{-4}$). These were filtered to include only those

which overlapped positions of RPE-1 CTCF ChIP-seq peaks (GSM749673, GSM1022665;

(ENCODE, 2012)), and stratified based on this ChIP-seq data. RPE-1 loop anchor-

associated CTCF motifs were identified using the Juicer MotifFinder (Durand et al., 2016)

with the RPE-1 WT Hi-C looplist (GSE71831 (Darrow et al., 2016)) and CTCF ChIP-seq BED

file (GSM749673, GSM1022665; (ENCODE, 2012)) as input. Human TOP2 CC-seq peak

coordinates were identified by thresholding (0.05 HpM) of pooled +VP16 data presented.

1045

1046   **CC-seq: Spatial correlation of yeast Top2 and Spo11 signals**

1047   Nucleotide resolution Top2/Spo11 CC-seq data was binned at 50 bp resolution and the

1048   Pearson correlation was calculated between the sum HpM for all bins at a given inter-bin

1049   distance.

1050

1051

1052

CC-seq: Quantifying single and double-strand Top2 CCs

Nucleotide resolution Human and Yeast Top2 CC-seq data were first thresholded at 0.05 and 1 HpM, respectively. SSB sites were defined as those sites without a cognate on the opposite strand at the expected offset of 3 bp. DSBs were defined as cognate sites with a 3 bp offset. SSB and DSB percentages are expressed as a percentage of the total number of sites (SSB + DSB). A randomisation experiment was conducted in order to estimate expected numbers of SSBs and DSBs within the sample under a random model where signal is distributed independently on each strand. To achieve this, the amplitudes of Top2 CCs (HpM) were shuffled amongst the positions in the nucleotide resolution datasets, prior to thresholding and offset analysis as described above.

CC-seq: DNA sequence composition of single and double-strand Top2 CCs

Nucleotide resolution Human and Yeast Top2 CC-seq data were first thresholded at 0.05 and 1 HpM, respectively. The high frequency positions remaining were classified as part of a single or double-stranded Top2 CC based on absence or presence of a cognate cleavage at the expected W-C offset of 3 bp. For double-stranded Top2 CCs, a dyad axis coordinate was defined as the centrepoint between the Top2-linked nucleotides on the W and C strand (that is, the midpoint of the central two base pairs in the four base pair overhang). For single-stranded Top2 CCs, we used an imaginary dyad axis in the same relative position. DNA sequence was aggregated ±20 bp around these two classes of Top2 CCs. Statistical significance was determined using the one sample (goodness-of-fit) Chi-squared test, as described previously (Pommier et al., 1991).

CC-seq - MethylC Analysis

A publically-available RPE-1 reduced representation bisulphite sequencing (RRBS) dataset (two replicates) was used to define coordinates of nucleotides immediately 3'-relative to 919,817 unmethylated or 341,136 methylated Cs, based on <10% or >90% methylation. Next we aggregated RPE-1 CC-seq (0.01 HpM thresholded) hits on these loci, and divided by the number of loci to give a density (barplot). Statistical significance was determined by the Kolmogorov-Smirnov test.

Ideograms

Human chromosome ideograms were adapted from the open source ideogram package (https://eweitz.github.io/ideogram/).

Supplementary Figure Legends

Figure S1. CC-seq maps covalent Spo11-linked DNA breaks in *S. cerevisiae* meiosis with nucleotide accuracy (related to Figure 1)

A) Correlation of hotspot signals from CC-seq and oligo-seq.

B) Correlation of 500 bp binned Spo11 maps from two representative replicates.

C) Correlation of Spo11 cleavages on Watson and Crick strand when offset by 1 bp (boxed), relative to other offsets from -4 to +4 bp.

D) Spo11 breaks mapped by CC-seq (raw=black, or smoothed=red) anticorrelate with nucleosome occupancy measured by MNase-seq (blue).

E) Aggregation of Spo11 activity in a 10 Kbp window centred on orientated TSSs in *S. cerevisiae*. Three example TSSs are shown orientated in the 5'-3' direction (top). A heatmap of all TSSs in the *S. cerevisiae* genome, with 25 rows stratified by gene expression level in SK1 cells (middle). The colour scale indicates average Spo11 break density. Motifs are also stratified into 4 quartiles of gene expression in meiotic SK1 cells, and the average distribution of Spo11 activity in each quartile plotted (bottom).

Figure S2. CC-seq maps covalent Top2-linked DNA breaks in *S. cerevisiae* cycling cells with nucleotide accuracy (related to Figure 2).

A-C) Correlation of 1 Kbp binned Top2 CC maps from pdr1Δ (A), pdr1Δsae2Δ (B) and pdr1Δmre11Δ (C) cells treated with VP16.

D) Pairwise Pearson correlation values for 100 bp binned Top2 CC maps from all assayed conditions. Each condition is a pool of two biological replicates.

E) Nucleotide-resolution S. cerevisiae Top2 CC map of chromosome 3 for all assayed conditions. Each condition is a pool of two biological replicates.

F-H) The normalised number of Top2 CCs retained in the pdr1Δ (F), pdr1Δsae2Δ (G) and pdr1Δmre11Δ **(H)** cells treated ±VP16, after filtering to include only sites offset by the given number of base pairs. All data were normalised over a -100 to +100 bp window.

I-K) Pearson correlation (r) of Top2 CC-seq signal on Watson and Crick strands, offset by the indicated distance in pdr1Δ (I), pdr1Δsae2Δ (J) and pdr1Δmre11Δ (K) cells treated ±VP16.

Figure S3. Top2-linked DNA breaks in *S. cerevisiae* are not correlated with gene expression, anticorrelate with nucleosomes, and have biased nucleotide skews indicative of bent DNA (related to Figure 2)

A) Top2 CC mapped by CC-seq (raw=black, or smoothed=red) anticorrelate with nucleosome occupancy measured by MNase-seq (blue).

B) Aggregation of Top2 CCs in a 10 Kbp window centred on orientated transcription start sites (TSS) in pdr1Δ S. cerevisiae. Four example TSSs are shown orientated in the 5'-3' direction (top). Heatmap

of all TSSs in the S. cerevisiae genome, with 25 rows stratified by gene expression level in vegetative growth (middle). Colour scale indicates average Top2 CC density. Motifs are also stratified into four quartiles of gene expression, and the average distribution of Top2 CCs in each quartile plotted for untreated and +VP16 conditions (bottom).

C) The percentage of strong sites that are SSBs and DSBs in etoposide-treated *pdr1Δ*, *pdr1Δsae2Δ* and *pdr1Δmre11Δ* S. cerevisiae. Sites were thresholded at 1 HpM prior to sorting into DSB or SSB classes based on presence or absence of a 3 bp offset cognate (Obs.). As a control, the amplitudes of Top2 CCs (HpM) were randomised amongst the positions in the nucleotide resolution datasets, prior to thresholding and offset analysis as described above (Rand.)

D) Average nucleotide composition over a 200 bp window centred on the DSB dyad axis, in etoposide-treated *pdr1Δ* S. cerevisiae. The position of the flanking regions (F.R.) identified in Gyrase mapping experiments, and the flanking regions observed here for Top2 are indicated in blue and black, respectively.

E) Average nucleotide composition over a 120 bp window centred on the DSB dyad axis, in etoposide-treated *pdr1Δ S. cerevisiae*. The signal pattern is separated into A+T and G+C (top left and right), and into individual A, T, C and G plots (middle left, bottom left, middle right, and bottom right, respectively). The position of the flanking regions (F.R.) observed here for Top2 are indicated in black, and their ~10.5 bp periodicity is highlighted with black and red crosses.

Figure S4. CC-seq maps of TOP2-linked DNA breaks in Human cells are enriched by etoposide, and show high reproducibility and nucleotide accuracy (related to Figure 3)

A) Broad-scale *H. sapiens* TOP2 CC-seq maps in individual biological replicates of RPE-1 cells ±VP16. Raw data were binned at 100 Kbp prior to plotting. Each plot is offset on the y-axis by +0.3 HpB.

B) Replicate-to-replicate Pearson correlation values (r) for 10 Kbp binned TOP2 CC-seq maps of RPE-1 cells ±VP16.

C) Scatter plot of -VP16 and +VP16 TOP2 CC-seq maps binned at 10 Kbp resolution. Data were first scaled according to the estimated noise fraction (**Methods**), and are presented in a hexagonal-binned format, where the density of overplotting is indicated by the colour scale.

D) Violin plots of TOP2 CC-seq maps ±VP16 binned at 100 Kbp resolution. The inner black bar, black dot, and dotted horizontal line indicate the interquartile range, median, and expected mean Top2 CC density based on random distribution.

E) The normalised number of TOP2 CCs retained in the CC-seq maps in RPE-1 cells ±VP16 after filtering to include only sites offset by the given number of base pairs. Data were normalised over a -100 to +100 bp window.

Figure S5. CC-seq signal is TOP2-dependent

A) DNA content histograms of wild type (WT) and *TOP2B*$^{-/-}$ RPE-1 cells under asynchronous (10% FCS) and serum-deprived (0% FCS) conditions, as measured by FACS following propidium iodide (PI) staining. G1, S and G2 populations are clearly present under asynchronous growing conditions. A strong G1 arrest is observed in serum deprived conditions. Percentages of cells in each of the indicated regions (red dotted brackets) are given.

B) Western blots demonstrating the absence of TOP2β protein in serum deprived and asynchronous *TOP2B*$^{-/-}$ RPE-1 cells (left), and the absence of TOP2α in serum deprived wild type and *TOP2B*$^{-/-}$ RPE-1 cells (right). Ponceau S total protein loading is presented (Pon. S) for the left and right panels, and additionally a Ku80 loading control is included for the right panel.

C) Immunofluorescence experiment demonstrating induction of γ-H2AX foci (green) in asynchronous (Async.) and serum-deprived (Ser. Dep.) wild type (WT) and *TOP2B*$^{-/-}$ RPE-1 cells, all co-stained with DAPI (blue). Galleries of nine cells per condition were chosen randomly using Olympus ScanR Analysis software.

D) Quantification of (C). Numbers of γ-H2AX foci per cell were counted automatically using Olympus ScanR Analysis software. The mean ±SEM is reported for n= 3 biological replicate experiments.

E) Broad-scale *H. sapiens* TOP2 CC-seq maps in asynchronous and serum-deprived wild type (WT) and *TOP2B*$^{-/-}$ RPE-1 cells -VP16 (orange) and +VP16 (green). Raw hits on Watson and Crick strands were summed and smoothed according to local signal density (Fsize=501).

Figure S6. TOP2 CC-seq signal enrichment around CTCF and TSS sites, compared with END-seq DSB signal at TSSs

A) Fine-scale TOP2 CC-seq maps of *H. sapiens* CTCF-proximal loci in RPE-1 cells +VP16. Red and blue traces indicate TOP2-linked 5' DNA termini on the Watson and Crick strands, respectively. Pale shaded areas are the same data smoothed with a sliding 11 bp Hanning window. Red and Blue rectangles indicate the positions of CTCF motifs on the Watson and Crick strands respectively. The grey line indicates Hanning-smoothed sum of Watson and Crick TOP2 CCs.

B) Fine-scale mapping of TOP2 CCs surrounding three complex CTCF loci, processed as in (A).

C) Aggregation of TOP2 CCs in a 1 Kbp window centred on the subset of orientated CTCF motifs that can be assigned to a chromatin loop anchor in human RPE-1 cells (Darrow et al., 2016). Motifs are stratified into 4 quartiles of loop anchor interaction strength, and the average TOP2 CC distribution in each quartile plotted.

D) Fine-scale aggregation of TOP2 CCs (red) in a 800 bp window centred on TSSs in human RPE-1 cells, showing anticorrelation with aggregated MNase-seq signal (blue).

E) Aggregation of END-seq mapped DSBs in a 10 Kbp window centred on orientated TSSs in human MCF7 cells. Four example TSSs are shown orientated in the 5'-3' direction (top). A heatmap of all TSSs in the human genome, with 25 rows stratified by flanking gene expression level in MCF7 cells (middle). The colour scale indicates average END-seq DSB density. TSSs are also stratified into 4

1202 quartiles based on strength of flanking gene expression, and the average END-seq DSB distribution

1203 in each quartile plotted (bottom).

1204

Figure S7. TSS-proximal TOP2-linked DNA breaks in humans are correlated with gene length,

independently from gene expression level (related to Figure 5); TSS-proximal Top2-linked DNA

breaks in yeast are not strongly correlated with gene length, (related to Figure 2)

A) Aggregation of TOP2 CCs in a 10 Kbp window centred on orientated TSSs in human RPE-1 cells.

A heatmap of all TSSs in the human genome, with 25 rows stratified by gene length (top). The colour

scale indicates average TOP2 CC density. Motifs are also stratified into 4 quartiles of gene length,

and the average TOP2 CC distribution in each quartile plotted (bottom).

B) Aggregation of Top2 CCs in a 10 Kbp window centred on orientated transcription start sites (TSS)

in *pdr1Δ S. cerevisiae*. Heatmap of all TSSs in the *S. cerevisiae* genome, with 25 rows stratified by

gene length level (top). Colour scale indicates average Top2 CC density. Motifs are also stratified

into four quartiles of gene length, and the average distribution of Top2 CCs in each quartile plotted

(bottom).

C) Scatter plot of human gene length and RPE-1 gene expression, showing no correlation.

D) Scatter plot of *S. cerevisiae* gene length and gene expression during vegetative growth, showing

little correlation.

1220
1221
1222
1223

Supplementary Tables

1224

1225
1226

1227 **Table S1. DNA libraries used for mapping Spo11 activity**. The number of reads

1228 remaining after each stage of the data pipeline are indicated.

1229

Species	Strain/ Cell line	Condition	NGS Platform	Library code	Total Read Pairs	Mapped reads	Blacklist-filtered reads	% of Total
S. cerevisiae	*sae2Δ*	Meiosis	MiSeq	1	3512089	3376630	3177210	90.5
				2A	10101423	8796998	8564017	84.8
				3	4957477	4745192	4502308	90.8
				5	4012972	3705100	3559541	88.7
				7BC	4065600	3779390	3569290	87.8

1230
1231
1232
1233

1234 **Table S2. DNA libraries used for mapping Top2 activity.** The number of reads
1235 remaining after each stage of the data pipeline are indicated.
1236

Species	Strain/ Cell line	Condition	NGS Platform	Library code	Total Read Pairs	Mapped reads	De-duplicated reads	Blacklist-filtered reads	% of Total	Pooled Reads (Millions)
S. cerevisiae	*pdr1Δ*	-VP16	NextSeq 500	RA1	3741251	3542955	3179049	2746294	73.4	3.50
			MiSeq	RA8	1118483	879107	842290	751850	67.2	
S. cerevisiae	*pdr1Δ*	+VP16	NextSeq 500	RA2	3096257	3053356	2679303	2234941	72.2	3.93
			MiSeq	RA9	2262958	2071231	1958341	1690075	74.7	
S. cerevisiae	*pdr1Δ sae2Δ*	-VP16	NextSeq 500	RA4	3634284	3552818	3215523	2745668	75.5	3.79
			MiSeq	RA10	1420662	1214067	1181399	1046827	73.7	
S. cerevisiae	*pdr1Δ sae2Δ*	+VP16	NextSeq 500	RA5	2529368	2476714	1386551	1097282	43.4	3.54
			MiSeq	RA11	3226626	3063388	2824882	2444486	75.8	
S. cerevisiae	*pdr1Δ mre11Δ*	-VP16	NextSeq 500	RA6	5345587	5205411	4288662	3646842	68.2	4.76
			MiSeq	RA12	1519020	1326435	1280137	1109310	73.0	
S. cerevisiae	*pdr1Δ mre11Δ*	+VP16	NextSeq 500	RA7	3382546	3357168	2921333	2526342	74.7	5.54
			MiSeq	RA13	4133347	4032996	3540366	3010542	72.8	
Human	RPE-1 WT	-VP16	NextSeq 500	WG11	125203096	109503841	68845619	68639594	54.8	185.36
				WG15	125328792	110432447	46588296	46449378	37.1	
				WG23	89154181	77728045	70484665	70269334	78.8	
Human	RPE-1 WT	+VP16	NextSeq 500	WG12	110910655	95546007	55848200	55681939	50.2	276.97
				WG16	102804692	91872706	54312084	54164650	52.7	
				WG24	130816235	114086421	101162102	100857557	77.1	
				WG34	91829516	79751337	66458538	66270349	72.2	
Human	RPE-1 WT	-VP16 G1	MiSeq	WG13	2601642	2292863	2007177	1991088	76.5	4.27
				WG21	2610523	2298289	2295625	2277124	87.2	
Human	RPE-1 WT	+VP16 G1	MiSeq	WG14	3075620	2771819	2491674	2475350	80.5	5.23
				WG22	3139164	2779809	2772349	2751281	87.6	
Human	RPE-1 WT	-VP16 Async	MiSeq	WG15b*	3207507	2866283	2782876	2760652	86.1	5.35
				WG23	2959289	2616954	2608670	2586523	87.4	
Human	RPE-1 WT	+VP16 Async	MiSeq	WG16b*	3018163	2721770	2673273	2656580	88.0	6.30
				WG24	4165048	3682708	3668602	3639324	87.4	
Human	RPE-1 TOP2B-	-VP16 G1	MiSeq	WG17	2978044	2624262	2558332	2536752	85.2	4.69
				WG25	2460949	2181002	2173777	2155101	87.6	
Human	RPE-1 TOP2B-	+VP16 G1	MiSeq	WG18	2927745	2588870	2557532	2537085	86.7	5.10
				WG26	2922624	2596547	2587152	2565454	87.8	
Human	RPE-1 TOP2B-	-VP16 Async	MiSeq	WG19	3713068	3278164	3188049	3162895	85.2	7.04
				WG27	4413242	3943962	3910035	3877611	87.9	
Human	RPE-1 TOP2B-	+VP16 Async	MiSeq	WG20	3426100	3038601	2978047	2956076	86.3	8.13
				WG28	5878342	5251089	5211027	5169010	87.9	

1237
1238
1239

Table S3. *S. cerevisiae* strains used in this study. Strains MJ315 and M319 were used for Spo11 mapping; strains MJ429, MJ475 and MJ551 were used for Top2 mapping.

Strain	Number	Genotype
sae2Δ	MJ315	SK1: *ho::LYS2/', lys2/', ura3/', arg4-nsp/', leu2::hisG/', his4X::LEU2/', nuc1::LEU2/', sae2Δ::KanMX6/'*
sae2Δ spo11-Y135F	MJ319	SK1: *ho::LYS2/', lys2/', ura3/', arg4-nsp/', leu2::hisG/', his4X::LEU2/', nuc1::LEU2/', spo11(Y135F)-HA3His6::KanMX4/', sae2Δ::KanMX6/'*
pdr1Δ	MJ429	BY4741: *ura3Δ0, leu2Δ0, his3Δ1, met15Δ0, pdr1Δ::PDR1-DBD-CYC8::LEU2*
pdr1Δ sae2Δ	MJ475	BY4741: *ura3Δ0, leu2Δ0, his3Δ1, met15Δ0, pdr1Δ::PDR1-DBD-CYC8::LEU2, sae2Δ::KanMX6*
pdr1Δ mre11Δ	MJ551	BY4741: *ura3Δ0, leu2Δ0, his3Δ1, met15Δ0, pdr1Δ::PDR1-DBD-CYC8::LEU2, mre11Δ::KanMX4*

Table S4. Human cell lines used in this study.

Parent cell line	Clone number	Genotype
hTERT RPE-1	WT	WT
hTERT RPE-1	T2B/1	*TOP2B*

Table S5. The annealed structures of the custom P5 and P7 adapters. "T_i" indicates an inverted dT, linked via a 3'-3' phosphodiester. This inhibits unwanted ligation at this end of the adapter. Also note that the 5'-terminal moieties of each adapter (A for P5, G for P7) are nucleosides (3'-OH), which further inhibits ligation at this end

Adapter	Annealed Structure		
P5	blocked end	ACACTCTTTCCCTACACGACGCTCTTCCGATCT T_iTGTGAGAAAGGGATGTGCTGCGAGAAGGCTAGp	ligated end
P7	ligated end	pGATCGGAAGAGCACACGTCTGAACTCCAGTCACT$_i$ TCTAGCCTTCTCGTGTGCAGACTTGAGGTCAGTG	blocked end

Table S6. Publically available datasets used in this study.

Species/ Cell line	Dataset Type	Accession(s)	Reference
S. cerevisiae / SK1 (meiotic)	Gene expression microarray	GSM907178, GSM90719, GSM907180	(Dominissini et al., 2012)
S. cerevisiae / SK1 (vegetative)	Gene expression microarray	GSM907176, GSM907177	(Dominissini et al., 2012)
H. sapiens / RPE-1	Gene expression microarray	GSM1395252, GSM1395253, GSM1395254	(Ganem et al., 2014)
H. Sapiens / MCF7	Gene expression microarray	GSM1141244	(Nelson et al., 2013)
H. sapiens / RPE-1	Hi-C	GSE71831	(Darrow et al., 2016)
H. sapiens / RPE-1	bTMP-seq	GSM1062645, GSM1062646	(Naughton et al., 2013)
H. sapiens / MCF7	END-seq	GSM2635568	(Canela et al., 2017)
H. sapiens / RPE-1	CTCF ChIP-seq	GSM749673, GSM1022665	(ENCODE, 2012)
H. sapiens / RPE-1	H3K4Me3 ChIP-seq	GSM945271, GSM945271	(ENCODE, 2012)
H. sapiens / RPE-1	H3K27Ac ChIP-seq	GSM733771, GSM733718, GSM733755, GSM733691, GSM733656, GSM733674, GSM733646	(ENCODE, 2012)
H. sapiens / RPE-1	Methyl-RRBS	GSM683773, GSM683905	(ENCODE, 2012)
H. sapiens / K562	MNase-seq	GSM920557	(ENCODE, 2012)

Supplementary methods references

Canela, A., Maman, Y., Jung, S., Wong, N., Callen, E., Day, A., Kieffer-Kwon, K.-R., Pekowska, A., Zhang, H., Rao, S. S. P., Huang, S.-C., Mckinnon, P. J., Aplan, P. D., Pommier, Y., Aiden, E. L., Casellas, R., and Nussenzweig, A. (2017). Genome Organization Drives Chromosome Fragility. Cell *170*, 507-521.e18.

Darrow, E. M., Huntley, M. H., Dudchenko, O., Stamenova, E. K., Durand, N. C., Sun, Z., Huang, S. C., Sanborn, A. L., Machol, I., Shamim, M., Seberg, A. P., Lander, E. S., Chadwick, B. P., and Aiden, E. L. (2016). Deletion of DXZ4 on the human inactive X chromosome alters higher-order genome architecture. Proc Natl Acad Sci U S A *113*, E4504-12.

Dominissini, D., Moshitch-Moshkovitz, S., Schwartz, S., Salmon-Divon, M., Ungar, L., Osenberg, S., Cesarkas, K., Jacob-Hirsch, J., Amariglio, N., Kupiec, M., Sorek, R., and Rechavi, G. (2012). Topology of the human and mouse m6A RNA methylomes revealed by m6A-seq. Nature *485*, 201-206.

Durand, N. C., Shamim, M. S., Machol, I., Rao, S. S., Huntley, M. H., Lander, E. S., and Aiden, E. L. (2016). Juicer Provides a One-Click System for Analyzing Loop-Resolution Hi-C Experiments. Cell Syst *3*, 95-98.

ENCODE, P. C. (2012). An integrated encyclopedia of DNA elements in the human genome. Nature *489*, 57-74.

Ganem, N. J., Cornils, H., Chiu, S. Y., O'Rourke, K. P., Arnaud, J., Yimlamai, D., Théry, M., Camargo, F. D., and Pellman, D. (2014). Cytokinesis failure triggers hippo tumor suppressor pathway activation. Cell *158*, 833-848.

Grant, C. E., Bailey, T. L., and Noble, W. S. (2011). FIMO: scanning for occurrences of a given motif. Bioinformatics *27*, 1017-1018.

Grzybowski, A. T., Chen, Z., and Ruthenburg, A. J. (2015). Calibrating ChIP-Seq with Nucleosomal Internal Standards to Measure Histone Modification Density Genome Wide. Mol Cell *58*, 886-899.

Hoffman, M. M., Ernst, J., Wilder, S. P., Kundaje, A., Harris, R. S., Libbrecht, M., Giardine, B., Ellenbogen, P. M., Bilmes, J. A., Birney, E., Hardison, R. C., Dunham, I., Kellis, M., and Noble, W. S. (2013). Integrative annotation of chromatin elements from ENCODE data. Nucleic Acids Res *41*, 827-841.

Hornyak, P., Askwith, T., Walker, S., Komulainen, E., Paradowski, M., Pennicott, L. E., Bartlett, E. J., Brissett, N. C., Raoof, A., Watson, M., Jordan, A. M., Ogilvie, D. J., Ward, S. E., Atack, J. R., Pearl, L. H., Caldecott, K. W., and Oliver, A. W. (2016). Mode of action of DNA-competitive small molecule inhibitors of tyrosyl DNA phosphodiesterase 2. Biochem J *473*, 1869-1879.

Hsu, P. D., Scott, D. A., Weinstein, J. A., Ran, F. A., Konermann, S., Agarwala, V., Li, Y., Fine, E. J., Wu, X., Shalem, O., Cradick, T. J., Marraffini, L. A., Bao, G., and Zhang, F. (2013). DNA targeting specificity of RNA-guided Cas9 nucleases. Nat Biotechnol *31*, 827-832.

Hu, B., Petela, N., Kurze, A., Chan, K.-L., Chapard, C., and Nasmyth, K. (2015). Biological chromodynamics: a general method for measuring protein occupancy across the genome by calibrating ChIP-seq. Nucleic Acids Res

Johnson, D., Allison, R. M., Cannavo, E., Cejka, P., and Neale, M. (2019). Removal of Spo11 from meiotic DNA breaks in vitro but not in vivo by Tyrosyl DNA Phosphodiesterase 2. bioRxiv *http://dx.doi.org/10.1101/527333*,

Liang, K., and Keleş, S. (2012). Normalization of ChIP-seq data with control. BMC Bioinformatics *13*, 199.

Lieberman-Aiden, E., van Berkum, N. L., Williams, L., Imakaev, M., Ragoczy, T., Telling, A., Amit, I., Lajoie, B. R., Sabo, P. J., Dorschner, M. O., Sandstrom, R., Bernstein, B., Bender, M. A., Groudine, M., Gnirke, A., Stamatoyannopoulos, J., Mirny, L. A., Lander, E. S., and Dekker, J. (2009). Comprehensive mapping of long-range interactions reveals folding principles of the human genome. Science *326*, 289-293.

Mathelier, A., Zhao, X., Zhang, A. W., Parcy, F., Worsley-Hunt, R., Arenillas, D. J., Buchman, S., Chen, C. Y., Chou, A., Ienasescu, H., Lim, J., Shyr, C., Tan, G., Zhou, M., Lenhard, B., Sandelin, A.,

and Wasserman, W. W. (2014). JASPAR 2014: an extensively expanded and updated open-access
database of transcription factor binding profiles. Nucleic Acids Res *42*, D142-7.

Naughton, C., Avlonitis, N., Corless, S., Prendergast, J. G., Mati, I. K., Eijk, P. P., Cockroft, S. L.,
Bradley, M., Ylstra, B., and Gilbert, N. (2013). Transcription forms and remodels supercoiling domains
unfolding large-scale chromatin structures. Nat Struct Mol Biol *20*, 387-395.

Nelson, E. R., Wardell, S. E., Jasper, J. S., Park, S., Suchindran, S., Howe, M. K., Carver, N. J., Pillai,
R. V., Sullivan, P. M., Sondhi, V., Umetani, M., Geradts, J., and McDonnell, D. P. (2013). 27-
Hydroxycholesterol links hypercholesterolemia and breast cancer pathophysiology. Science *342*,
1094-1098.

Pan, J., Sasaki, M., Kniewel, R., Murakami, H., Blitzblau, H. G., Tischfield, S. E., Zhu, X., Neale, M.
J., Jasin, M., Socci, N. D., Hochwagen, A., and Keeney, S. (2011). A Hierarchical Combination of
Factors Shapes the Genome-wide Topography of Yeast Meiotic Recombination Initiation. Cell *144*,
719-731.

Pommier, Y., Capranico, G., Orr, A., and Kohn, K. W. (1991). Local base sequence preferences for
DNA cleavage by mammalian topoisomerase II in the presence of amsacrine or teniposide. Nucleic
Acids Res *19*, 5973-5980.

Stepanov, A., Nitiss, K. C., Neale, G., and Nitiss, J. L. (2008). Enhancing drug accumulation in
Saccharomyces cerevisiae by repression of pleiotropic drug resistance genes with chimeric
transcription repressors. Mol Pharmacol *74*, 423-431.

1332

1333

Figure 1

Figure 1. CC-seq maps covalent Spo11-linked DNA breaks in *S. cerevisiae* meiosis with nucleotide accuracy

A) Schematic of the CC-seq method.

B) Column-based enrichment of Spo11-linked DNA fragments detected by Southern blotting at the *his4*::*LEU2* recombination hotspot (**Methods**). Arrows indicate expected sizes of Spo11-DSBs.

C) Nucleotide resolution mapping of *S. cerevisiae* Spo11 hotspots by CC-seq or oligo-seq (Pan et al., 2011). Red and blue traces indicate Spo11-linked 5' DNA termini on the Watson and Crick strands, respectively. Grey arrows indicate positions of gene open reading frames.

D) Pearson correlation (r) of Spo11 CC-seq signal between Watson and Crick strands, offset by the indicated distances.

E) Average nucleotide composition over a 200 (top) and 30 bp (bottom) window centred on Spo11 breaks. Bases reported are for the top strand only.

HpM = Hits per million mapped reads per base pair.

Figure 2

Figure 2. CC-seq maps covalent Top2-linked DNA breaks in *S. cerevisiae* cycling cells with nucleotide accuracy
A) Serial dilution spot tests of VP16 tolerance for the indicated strains.
B) Nucleotide resolution mapping of *S. cerevisiae* Top2 CCs by CC-seq of the indicated strains after treatment for 4 hours with 1 mM VP16. Spo11-CC data is plotted for comparison. Top2 CC data were calibrated using a human DNA spike-in (**Methods**). Red and blue traces indicate CC-linked 5' DNA termini on the Watson and Crick strands, respectively. Grey arrows indicate positions of gene open reading frames. Lower panels show an expanded view of the region from 31.5 to 40 Kbp.
C) Quantification of Top2 and Spo11 CC signal stratified by genomic region. The genome was divided into intra and intergenic regions; the intergenic region was further divided into divergent, tandem and convergent based on orientation of flanking genes. Spo11 and Top2 activity mapped by CC-seq is expressed as box-and-whisker plots of density (upper and lower box limit: 3rd and 1st quartile; bar: median; upper and lower whisker: highest and lowest values within 1.5-fold of the interquartile range), or as the percentage of total mapped reads.
D) Local correlation of Top2 or Spo11 CC-seq signals. Top2 or Spo11 CC-seq data were binned at 50 bp resolution and the Pearson correlation calculated between bins of increasing separation.
E) Pearson correlation (r) of Top2 CC-seq signal on Watson and Crick strands, offset by the indicated distance.
HpM = Hits per million mapped reads per base pair.

Figure 3

Figure 3. CC-seq maps TOP2-linked DNA breaks in Human cells with nucleotide accuracy
A) Anti-TOP2β western slot blot of input, flow through, wash, and elution fractions. RPE-1 cells were treated or not with VP16, prior to processing according to **Figure 1A**.
B) Broad-scale maps of *H. sapiens* TOP2 CCs produced by CC-seq in RPE-1 cells ±VP16. Raw data were scaled, binned, smoothed and median subtracted prior to plotting (**Methods**). Chromatin compartments revealed by Hi-C eigenvector analysis (Darrow et al., 2016), and supercoiling revealed by bTMP ChIP-seq (Naughton et al., 2013) are shown for comparison.
C) Quantification of TOP2 CC in chromatin compartments A and B. Data are expressed as box-and-whisker plots of density as for **Figure 2C**.
D) Fine-scale mapping of *H. sapiens* TOP2 CCs by CC-seq in RPE-1 cells ±VP16. Red and blue traces indicate TOP2-linked 5' DNA termini on the Watson and Crick strands, respectively. Pale shaded areas are the same data smoothed according to local density (see **Methods**). RPE-1 CTCF and H3K4Me3 ChIP-seq data plus H3K27Ac ChIP-seq data overlaid from seven cell lines (ENCODE, 2012) is shown for comparison.
E) Medium-scale aggregate of END-seq mapped DSBs (Canela et al., 2017) surrounding nucleotide resolution CC-seq mapped TOP2 CCs.
F) Fine-scale strand-specific aggregate of END-seq mapped DSB ends (red and blue bars) and nucleotide resolution CC-seq mapped TOP2 CCs (grey bars) surrounding strong TOP2 CC sites.

Figure 4

Figure 4. CTCF-proximal TOP2 activity is tightly confined within nucleosome-depleted regions
A) Aggregation of TOP2 CCs in a 1 Kbp window centred on orientated CTCF motifs in human RPE-1 cells. Four example CTCF loci are shown orientated in the 5'-3' direction (top). A heatmap of all CTCF-motifs in the human genome, with 25 rows stratified by the strength of colocalising CTCF ChIP-seq peaks in RPE-1 cells (middle). The colour scale indicates average TOP2 CC density. Motifs are also stratified into 4 quartiles of CTCF-binding, and the average TOP2 CC distribution in each quartile plotted (bottom).
B) Aggregated TOP2 CC distribution (red line) in the highest quartile of CTCF-binding compared with the average MNase-seq signal (blue line).
C) Aggregated TOP2 CC distribution (black and red lines: single-nucleotide resolution and smoothed, respectively) and the average MNase-seq signal (blue line) surrounding strong TOP2 CC sites.
In (B) and (C) peaks in MNase-seq signal indicate inferred nucleosome positions (blue circles).

Figure 5

Figure 5. TSS-proximal TOP2 activity is strongly correlated with gene transcription
A) Aggregation of TOP2 CCs in a 10 Kbp window centred on orientated TSSs in human RPE-1 cells. Four example TSSs are shown orientated in the 5'-3' direction (top). A heatmap of all TSSs in the human genome, with 25 rows stratified by gene expression level in RPE-1 cells (middle). The colour scale indicates average TOP2 CC density. Motifs are also stratified into 4 quartiles of gene expression, and the average TOP2 CC distribution in each quartile plotted (bottom).
B) Comparison of CTCF-proximal and TSS-proximal TOP2 CC distributions in the highest quartile of CTCF-binding (pink) and gene-expression (blue), respectively.
C) Average TOP2 CC density in 10 Kbp regions centred on the highest quartile of TSS, CTCF, or regions where both features are present. Data are expressed as box-and-whisker plots of density as in **Figure 2C**. Statistical significance was determined using the KS test.
D) As in (C) but sum total TOP2 CCs found in these regions.

Figure 6

Figure 6. Local DNA sequence and methylation status direct the formation of Top2 CCs
A) The percentage of strong sites that are SSBs and DSBs in etoposide-treated *S. cerevisiae* and human cells.
B) Average nucleotide composition over a 30 bp window centred on the DSB or (inferred) SSB dyad axis, in etoposide-treated *S. cerevisiae* and human cells. Values reported are for the top strand only.
C) Heatmaps describing pairwise similarity of nucleotide composition patterns shown in (B). Rows 1-4 are the absolute differences between: yeast and human SSB patterns, yeast and human DSB patterns, yeast SSB and DSB patterns, and human SSB and DSB patterns, respectively.
D) The average number of TOP2 CCs at the +1 position relative to methylated and unmethylated cytosines in human RPE-1 cells, ±VP16. Statistical significance was determined using KS test.

Figure S1

Figure S1. CC-seq maps covalent Spo11-linked DNA breaks in *S. cerevisiae* meiosis with nucleotide accuracy (related to Figure 1)

A) Correlation of hotspot signals from CC-seq and oligo-seq.

B) Correlation of 500 bp binned Spo11 maps from two representative replicates.

C) Correlation of Spo11 cleavages on Watson and Crick strand when offset by 1 bp (boxed), relative to other offsets from -4 to +4 bp.

D) Spo11 breaks mapped by CC-seq (raw=black, or smoothed=red) anticorrelate with nucleosome occupancy measured by MNase-seq (blue).

E) Aggregation of Spo11 activity in a 10 Kbp window centred on orientated TSSs in *S. cerevisiae*. Three example TSSs are shown orientated in the 5'-3' direction (top). A heatmap of all TSSs in the *S. cerevisiae* genome, with 25 rows stratified by gene expression level in SK1 cells (middle). The colour scale indicates average Spo11 break density. Motifs are also stratified into 4 quartiles of gene expression in meiotic SK1 cells, and the average distribution of Spo11 activity in each quartile plotted (bottom).

Figure S2

Figure S2. CC-seq maps covalent Top2-linked DNA breaks in _S. cerevisiae_ cycling cells with nucleotide accuracy (related to Figure 2).

A-C) Correlation of 1 Kbp binned Top2 CC maps from _pdr1Δ_ (A), _pdr1Δsae2Δ_ (B) and _pdr1Δmre11Δ_ (C) cells treated with VP16.

D) Pairwise Pearson correlation values for 100 bp binned Top2 CC maps from all assayed conditions. Each condition is a pool of two biological replicates.

E) Nucleotide-resolution _S. cerevisiae_ Top2 CC map of chromosome 3 for all assayed conditions. Each condition is a pool of two biological replicates.

F-H) The normalised number of Top2 CCs retained in the _pdr1Δ_ (F), _pdr1Δsae2Δ_ (G) and _pdr1Δmre11Δ_ (H) cells treated ±VP16, after filtering to include only sites offset by the given number of base pairs. All data were normalised over a -100 to +100 bp window.

I-K) Pearson correlation (r) of Top2 CC-seq signal on Watson and Crick strands, offset by the indicated distance in _pdr1Δ_ (I), _pdr1Δsae2Δ_ (J) and _pdr1Δmre11Δ_ (K) cells treated ±VP16.

Figure S3

Figure S3. Top2-linked DNA breaks in *S. cerevisiae* are not correlated with gene expression, anticorrelate with nucleosomes, and have biased nucleotide skews indicative of bent DNA (related to Figure 2)

A) Top2 CC mapped by CC-seq (raw=black, or smoothed=red) anticorrelate with nucleosome occupancy measured by MNase-seq (blue).

B) Aggregation of Top2 CCs in a 10 Kbp window centred on orientated transcription start sites (TSS) in *pdr1Δ S. cerevisiae*. Four example TSSs are shown orientated in the 5'-3' direction (top). Heatmap of all TSSs in the *S. cerevisiae* genome, with 25 rows stratified by gene expression level in vegetative growth (middle). Colour scale indicates average Top2 CC density. Motifs are also stratified into four quartiles of gene expression, and the average distribution of Top2 CCs in each quartile plotted for untreated and +VP16 conditions (bottom).

C) The percentage of strong sites that are SSBs and DSBs in etoposide-treated *pdr1Δ*, *pdr1Δsae2Δ* and *pdr1Δmre11Δ S. cerevisiae*. Sites were thresholded at 1 HpM prior to sorting into DSB or SSB classes based on presence or absence of a 3 bp offset cognate (Obs.). As a control, the amplitudes of Top2 CCs (HpM) were randomised amongst the positions in the nucleotide resolution datasets, prior to thresholding and offset analysis as described above (Rand.)

D) Average nucleotide composition over a 200 bp window centred on the DSB dyad axis, in etoposide-treated *pdr1Δ S. cerevisiae*. The position of the flanking regions (F.R.) identified in Gyrase mapping experiments, and the flanking regions observed here for Top2 are indicated in blue and black, respectively.

E) Average nucleotide composition over a 120 bp window centred on the DSB dyad axis, in etoposide-treated *pdr1Δ S. cerevisiae*. The signal pattern is separated into A+T and G+C (top left and right), and into individual A, T, C and G plots (middle left, bottom left, middle right, and bottom right, respectively). The position of the flanking regions (F.R.) observed here for Top2 are indicated in black, and their ~10.5 bp periodicity is highlighted with black and red crosses.

Figure S4

Figure S4. CC-seq maps of TOP2-linked DNA breaks in Human cells are enriched by etoposide, and show high reproducibility and nucleotide accuracy (related to Figure 3)

A) Broad-scale *H. sapiens* TOP2 CC-seq maps in individual biological replicates of RPE-1 cells ±VP16. Raw data were binned at 100 Kbp prior to plotting. Each plot is offset on the y-axis by +0.3 HpB.

B) Replicate-to-replicate Pearson correlation values (r) for 10 Kbp binned TOP2 CC-seq maps of RPE-1 cells ±VP16.

C) Scatter plot of -VP16 and +VP16 TOP2 CC-seq maps binned at 10 Kbp resolution. Data were first scaled according to the estimated noise fraction (**Methods**), and are presented in a hexagonal-binned format, where the density of overplotting is indicated by the colour scale.

D) Violin plots of TOP2 CC-seq maps ±VP16 binned at 100 Kbp resolution. The inner black bar, black dot, and dotted horizontal line indicate the interquartile range, median, and expected mean Top2 CC density based on random distribution.

E) The normalised number of TOP2 CCs retained in the CC-seq maps in RPE-1 cells ±VP16 after filtering to include only sites offset by the given number of base pairs. Data were normalised over a -100 to +100 bp window.

Figure S5

Figure S5. CC-seq signal is TOP2-dependent

A) DNA content histograms of wild type (WT) and *TOP2B*-/- RPE-1 cells under asynchronous (10% FCS) and serum-deprived (0% FCS) conditions, as measured by FACS following propidium iodide (PI) staining. G1, S and G2 populations are clearly present under asynchronous growing conditions. A strong G1 arrest is observed in serum deprived conditions. Percentages of cells in each of the indicated regions (red dotted brackets) are given.

B) Western blots demonstrating the absence of TOP2β protein in serum deprived and asynchronous *TOP2B*-/- RPE-1 cells (left), and the absence of TOP2α in serum deprived wild type and *TOP2B*-/- RPE-1 cells (right). Ponceau S total protein loading is presented (Pon. S) for the left and right panels, and additionally a Ku80 loading control is included for the right panel.

C) Immunofluorescence experiment demonstrating induction of γ-H2AX foci (green) in asynchronous (Async.) and serum-deprived (Ser. Dep.) wild type (WT) and *TOP2B*-/- RPE-1 cells, all co-stained with DAPI (blue). Galleries of nine cells per condition were chosen randomly using Olympus ScanR Analysis software.

D) Quantification of (C). Numbers of γ-H2AX foci per cell were counted automatically using Olympus ScanR Analysis software. The mean ±SEM is reported for n= 3 biological replicate experiments.

E) Broad-scale *H. sapiens* TOP2 CC-seq maps in asynchronous and serum-deprived wild type (WT) and *TOP2B*-/- RPE-1 cells -VP16 (orange) and +VP16 (green). Raw hits on Watson and Crick strands were summed and smoothed according to local signal density (Fsize=501).

Figure S6

Figure S6. TOP2 CC-seq signal enrichment around CTCF and TSS sites, compared with END-seq DSB signal at TSSs

A) Fine-scale TOP2 CC-seq maps of *H. sapiens* CTCF-proximal loci in RPE-1 cells +VP16. Red and blue traces indicate TOP2-linked 5' DNA termini on the Watson and Crick strands, respectively. Pale shaded areas are the same data smoothed with a sliding 11 bp Hanning window. Red and Blue rectangles indicate the positions of CTCF motifs on the Watson and Crick strands respectively. The grey line indicates Hanning-smoothed sum of Watson and Crick TOP2 CCs.

B) Fine-scale mapping of TOP2 CCs surrounding three complex CTCF loci, processed as in (A).

C) Aggregation of TOP2 CCs in a 1 Kbp window centred on the subset of orientated CTCF motifs that can be assigned to a chromatin loop anchor in human RPE-1 cells (Darrow et al., 2016). Motifs are stratified into 4 quartiles of loop anchor interaction strength, and the average TOP2 CC distribution in each quartile plotted.

D) Fine-scale aggregation of TOP2 CCs (red) in a 800 bp window centred on TSSs in human RPE-1 cells, showing anticorrelation with aggregated MNase-seq signal (blue).

E) Aggregation of END-seq mapped DSBs in a 10 Kbp window centred on orientated TSSs in human MCF7 cells. Four example TSSs are shown orientated in the 5'-3' direction (top). A heatmap of all TSSs in the human genome, with 25 rows stratified by flanking gene expression level in MCF7 cells (middle). The colour scale indicates average END-seq DSB density. TSSs are also stratified into 4 quartiles based on strength of flanking gene expression, and the average END-seq DSB distribution in each quartile plotted (bottom).

Figure S7

Figure S7. TSS-proximal TOP2-linked DNA breaks in humans are correlated with gene length, independently from gene expression level (related to Figure 5); TSS-proximal Top2-linked DNA breaks in yeast are not strongly correlated with gene length, (related to Figure 2)

A) Aggregation of TOP2 CCs in a 10 Kbp window centred on orientated TSSs in human RPE-1 cells. A heatmap of all TSSs in the human genome, with 25 rows stratified by gene length (top). The colour scale indicates average TOP2 CC density. Motifs are also stratified into 4 quartiles of gene length, and the average TOP2 CC distribution in each quartile plotted (bottom).

B) Aggregation of Top2 CCs in a 10 Kbp window centred on orientated transcription start sites (TSS) in *pdr1Δ S. cerevisiae*. Heatmap of all TSSs in the *S. cerevisiae* genome, with 25 rows stratified by gene length level (top). Colour scale indicates average Top2 CC density. Motifs are also stratified into four quartiles of gene length, and the average distribution of Top2 CCs in each quartile plotted (bottom).

C) Scatter plot of human gene length and RPE-1 gene expression, showing no correlation.

D) Scatter plot of *S. cerevisiae* gene length and gene expression during vegetative growth, showing little correlation.