A nucleotide resolution map of Top2-linked DNA breaks in the yeast and human genome

William Gittens, Dominic J. Johnson, Rachal M. Allison, Tim J. Cooper, Holly Thomas, and Matthew J Neale

Genome Damage and Stability Centre, University of Sussex, UK, BN1 9RQ

Corresponding author: m.neale@sussex.ac.uk

Abstract

DNA topoisomerases are required to resolve DNA topological stress. Despite this essential role, abortive topoisomerase activity generates aberrant protein-linked DNA breaks, jeopardising genome stability. Here, to understand the genomic distribution and mechanisms underpinning topoisomerase-induced DNA breaks, we map Top2 DNA cleavage with strand-specific nucleotide resolution across the *S. cerevisiae* and human genomes—and use the meiotic Spo11 protein to validate the broad applicability of this method to explore the role of diverse topoisomerase family members. Our data characterises Mre11-dependent repair in yeast, and defines two strikingly different fractions of Top2 activity in humans: tightly localised CTCF-proximal and broadly distributed transcription-proximal. Moreover, the nucleotide resolution accuracy of our assay reveals the influence primary DNA sequence has upon Top2 cleavage—for the first time distinguishing canonical DNA double-strand breaks (DSBs) from a major population of DNA single-strand breaks (SSBs) induced by etoposide (VP16) *in vivo*.

Introduction

DNA topoisomerases are a broad and ubiquitous family of enzymes that tackle topological constraints to replication, transcription, the maintenance of genome structure and chromosome segregation in mitosis and meiosis (Pommier et al., 2016). Although the specific mechanisms by which this is accomplished vary considerably across the family, key aspects are shared: including single or double-strand DNA cleavage to form a transient covalent complex (CC), which allows alteration of the topology of the nucleic acid substrate prior to relegation (Wang, 2009). These processes are essential but carry with them a significant risk to genome stability because the CC may be stabilised as a permanent

protein-linked DNA break by several physiological factors; such as the proximity of other DNA lesions (Sabourin and Osheroff, 2000; Kingma et al., 1995; Khan et al., 2003; Vélez-Cruz et al., 2005; Deweese and Osheroff, 2009; Wang et al., 2000; Wang et al., 1999), the collision of transcription and replication complexes (Wu and Liu, 1997; Hsiang et al., 1989), denaturation of the topoisomerase (Liu et al., 1983), or by the binding of small molecules that inhibit religation (Reviewed in (Nitiss, 2009)). Topoisomerase-induced DNA strand breaks have been proposed to constitute a significant fraction of the total damage to genomic DNA per day, and have been linked to the genesis and development of various cancers (Lin et al., 2009; Haffner et al., 2010; Sasanuma et al., 2018), including a subset of therapy-related acute myeloid leukaemias (t-AML) caused by the use of Topoisomerase 2 (Top2) poisons in the chemotherapeutic treatment of primary cancers (Rowley and Olney, 2002; Cowell et al., 2012).

In *S. cerevisiae*, either Top1 or Top2 activity is sufficient to support transcription (Brill et al., 1987; Lotito et al., 2008; Bermejo et al., 2009). However, whilst *top1Δ* cells are viable, Top2 is essential for sister chromatid segregation (Baxter and Diffley, 2008; DiNardo et al., 1984; Holm et al., 1985; Uemura and Yanagida, 1984). In contrast to *S. cerevisiae*, all known vertebrate species encode two Top2 proteins (TOP2α and TOP2β). Interestingly, whilst TOP2α is essential for cellular proliferation—cells arrest in mitosis in its absence (Akimitsu et al., 2003)—TOP2β is not. Rather, TOP2β apparently plays an important role in promoting transcriptional programmes associated with neuronal development (Tiwari et al., 2012; Gómez-Herreros et al., 2014), a function that cannot be supported by TOP2α.

Accurate maps of the positions of topoisomerase-DNA covalent complexes (Top2 CCs) genome-wide and throughout the cell cycle provide insights into topological genome structural organisation, as well as providing the tools to study their repair should they become permanent DNA lesions. Whilst the dual catalytic sites present within the homodimeric Top2 enzyme suggest a primary role in DNA double-strand break (DSB) formation driven by a biological need for decatenation (Pommier et al., 2016), previous research has suggested that etoposide and other Top2 poisons may induce a population of DNA single-strand breaks (SSBs), due to independent inhibition of each active site (Wozniak and Ross, 1983; Long et al., 1986; Zwelling et al., 1981; Pommier et al., 1984; Bromberg et al., 2003). Yet, despite the use of etoposide in chemotherapy, and the different toxicity of these two classes of DNA lesion, the prevalence of Top2-SSBs remains unclear.

More recently, general DSB mapping techniques have been applied to map etoposide-
induced lesions (Canela et al., 2017; Crosetto et al., 2013; Yan et al., 2017). However, such
methods are only able to map DSB ends, which therefore excludes and obscures
etoposide-induced SSBs. Moreover, these methodologies lack specificity for protein-linked
DNA ends, and require nucleolytic processing to blunt 5' DNA termini as part of sample
preparation (Canela et al., 2017). When combined, these factors lead to loss of nucleotide
resolution at the site of Top2 cleavage.

To generate a more complete, and direct, picture of topoisomerase action across the
genome, we present here a technique for nucleotide resolution mapping of protein-linked
SSBs and DSBs (referred to collectively here as "Covalent Complexes"; CC), which we
demonstrate to be generally applicable in both yeast and human cell systems. We establish
with high confidence that the technique is able to map both the positions of Top2 CCs and
also CCs of DNA linked to the meiotic recombination protein Spo11, which is related to the
archaeal Topoisomerase VI (Bergerat et al., 1997). Furthermore, we provide insights into the
spatial distribution of human TOP2 CCs, including comparative analysis of their local
enrichment around transcription start sites (TSSs) and CTCF-binding motifs. We find that
TSS-proximal TOP2 CC levels are strongly correlated with transcription, a question that has
come under scrutiny recently (Canela et al., 2017). Finally, we present compelling evidence
that etoposide induces a majority of Top2-linked SSBs *in vivo*, corroborating previous
research and revealing, for the first time, that this is governed by primary DNA sequence at
the cleavage site.

Results

CC-seq enriches Spo11-linked DNA fragments
To elucidate the *in vivo* functions of topoisomerase-like enzymes, we set out to establish a
direct method (termed 'CC-seq') to enrich and map protein-DNA covalent complexes (CC)
genome-wide with nucleotide resolution. Silica fibre-based enrichment of CCs has been
used to study the adenovirus DNA terminal protein complex (Coombs and Pearson, 1978),
and in the original identification of the meiotic DSB-inducer Spo11 (Keeney and Kleckner,
1995; Keeney et al., 1997), an orthologue of archaeal Topoisomerase VI (Bergerat et al.,
1997). We first verified this enrichment principle using meiotic *sae2Δ* cells, in which Spo11-
linked DSBs are known to accumulate at defined loci due to abrogation of the nucleolytic
pathway that releases Spo11 (Keeney and Kleckner, 1995; Neale et al., 2005). To
demonstrate specific enrichment of protein-linked molecules, total genomic DNA from

meiotic *S. cerevisiae* cells was isolated in the absence of proteolysis, digested with *Pst*I

restriction enzyme, and isolated on glass-fibre spin columns (**Figure 1A, Methods**). We

used eluted material to assay a known Spo11-DSB hotspot by Southern blotting (**Figure**

1B). While DSB fragments are a minor fraction of input material (~10% of total), and were

absent in wash fractions, DSBs accounted for >99% of total eluted material, indicating

~1000-fold enrichment relative to non-protein-linked DNA (**Figure 1B**).

CC-seq maps known Spo11-DSB hotspots genome-wide with high reproducibility

To generate a genome-wide map of Spo11-DSBs, genomic DNA from meiotic *sae2Δ* cells

was sonicated to <400 bp in length, enriched upon the silica column, eluted, and ligated to

DNA adapters in a two-step procedure that utilised the known phosphotyrosyl-unlinking

activity of mammalian TDP2 to uncap the Spo11-bound end ('CC-seq'; **Figure 1A /**

Methods) (Cortes Ledesma et al., 2009; Johnson et al., 2019). Libraries were paired-end

sequenced and mapped to the *S. cerevisiae* reference genome (**Table S1**) alongside reads

from a previous mapping technique ('Spo11-oligo-seq') that relies on the isolation of

Spo11-linked oligonucleotides generated in wild-type cells during DSB repair (Pan et al.,

2011). CC-seq revealed sharp, localised peaks ('hotspots') in *SPO11*+ cells (**Figure 1C,**

middle) that visually (**Figure 1C**) and quantitatively (**Figure 1S1A**, r=0.82) correlate with

Spo11-oligo seq, and that are absent in the *spo11-Y135F* control strain (**Figure 1C,**

bottom) in which Spo11-DSBs do not form (Bergerat et al., 1997), demonstrating that

signal detected by CC-seq accurately reflects the distribution of Spo11 cleavages.

Moreover, CC-seq maps generated from independent biological samples were highly

correlated (**Figure S1B**; r = 0.98), demonstrating very high reproducibility. Finally, when

analysed at nucleotide resolution, CC-seq revealed high correlation (r = 0.85) between the

frequency of Watson-mapping and Crick-mapping cleavage sites, when offset by a single

bp (**Figure 1E and S1C**). This correlation is expected because of the 2 bp 5' overhang

generated at Spo11-DSBs *in vivo* (**Fig1D**; e.g. (Liu et al., 1995))—demonstrating the

nucleotide resolution accuracy of CC-seq, which is further supported by our observation of

nucleotide composition preference at the site of cleavage (**Figure 1E**).

CC-seq enriches and maps Top2-linked DNA fragments from *S. cerevisiae*

Confident in the specificity of CC-seq to map meiotic protein-linked DSBs with high

resolution and dynamic range, we next employed CC-seq to map within *S. cerevisiae* the

covalent complexes generated naturally by Topoisomerase 2 (Top2) that become stabilised

upon exposure to etoposide, and are thus a proxy for sites of Top2 catalytic activity

(Minocha and Long, 1984; Chen et al., 1984; Wu et al., 2011). Because *S. cerevisiae* is
relatively insensitive to etoposide, we utilised a strain background ('*pdr1Δ*') in which the
action of the major drug export pathways are downregulated due to modulation of *PDR1*
activity (Stepanov et al., 2008). As expected, the *pdr1Δ* background dramatically increased
cellular sensitivity to etoposide, which was further enhanced by mutation of *SAE2* and
MRE11 (**Figure 2A**), factors involved in the repair of covalent protein-linked DNA breaks
(Neale et al., 2005; Garcia et al., 2011; Anand et al., 2016; Cannavo and Cejka, 2014;
Cannavo et al., 2018; Hoa et al., 2016; Hartsuiker et al., 2009).

Next, we prepared sequencing libraries from mitotically growing etoposide-treated control
(*pdr1Δ*) *S. cerevisiae* cells, or cells that also harboured *sae2Δ* and *mre11Δ* mutations (**Table
S2**). A human DNA spike-in was included to enable calibration of relative signal between
strains (**Methods**). Libraries displayed high replicate-to-replicate reproducibility (r = 0.89,
0.95 and 0.97 for control, *sae2Δ* and *mre11Δ*, respectively; **Figures S2A-C**), therefore
replicates were pooled. In untreated control cells, Top2 CC signal was distributed
homogeneously across the genome, with few strong peaks (**Figure 2B**). Upon etoposide
treatment, sharp single-nucleotide peaks arose at similar locations across the genome in all
strains (**Figure 2B and S2D-E**), but with increased amplitude in the repair mutants—thereby
directly linking the MRX pathway to repair of Top2 CCs in *S. cerevisiae*, and corroborating
prior research in other organisms (Hartsuiker et al., 2009; Hoa et al., 2016; Nakamura et al.,
2010).

Global analyses revealed Top2 activity in *S. cerevisiae* to be relatively enriched in divergent
and tandem intergenic regions (IGRs) and depleted in intragenic and convergent IGRs
(**Figure 2B and C**), similar to, but less pronounced than, the patterns of Spo11 (**Figure 2B
and C**). Such connections to global gene organisation are likely driven by the need for
Spo11 and Top2 to interact with the DNA helix—an interpretation underpinned by an
anticorrelation with nucleosome occupancy (**Figure S1E and S3A**). Nevertheless, the fact
that *S. cerevisiae* Top2 CC signal is not correlated with proximal gene expression—neither
in untreated cells, nor following etoposide exposure (**Figure S3B**)—suggests that in the
gene-dense *S. cerevisiae* genome topological stress may be dealt with by Top2 at sites
dislocated from where it is generated. By contrast, Spo11 activity is positively correlated
with gene expression (**Figure S1D**), perhaps due to the influence transcription has upon the
generation of higher order chromatin loop structures (Schalbetter et al., 2018), which in turn
are known to pattern Spo11-DSBs (Sun et al., 2015).

Interestingly, Top2 CC signal remained weakly correlated over greater distances than

Spo11 (**Figure 2D**), suggesting localised 5-10 kb wide regions of enriched activity—much

larger than Spo11-DSB hotspots (generally <500 bp in size; **Figure 1C**; (Pan et al., 2011)).

Top2 activity was strongly correlated between Watson and Crick strands when offset by 3

bp (**Figure 2E and S2F-K**)—as expected for the 4 bp overhang generated by Top2-DSBs—

further demonstrating the nucleotide resolution accuracy of CC-seq. Together these data

describe how CC-seq reveals a strand-specific nucleotide resolution map of etoposide-

induced Top2 CCs.

CC-seq maps TOP2 CCs in the Human genome

Having demonstrated that CC-seq is applicable for mapping two divergent types of

topoisomerase-like covalent complexes in *S. cerevisiae*, we applied CC-seq to map TOP2

activity in a human cell system. Asynchronous, sub-confluent RPE-1 cells were treated with

etoposide (VP16) in the presence of MG132 proteasome inhibitor, in order to limit potential

proteolytic degradation of TOP2 CCs that might otherwise hamper enrichment (**Figure 1A**).

Slot-blotting and immunodetection with anti-TOP2β antibody demonstrated complete

recovery of input TOP2β-linked CCs within the column elution, with no TOP2β remaining in

the flow through or being removed by the washes (**Figure 3A**). Eluted material was used to

generate sequencing libraries from three replicate control (–VP16) and four replicate

etoposide-treated samples (+VP16), and high-depth paired-end sequencing reads (**Table**

S2) were aligned to the human genome (hg19; **Methods**).

Replicates displayed high correlation in the distribution of TOP2 CC signal at broad scale (r

values ≥ 0.79; **Figure S4A-B**), and so were pooled. Visual inspection revealed that -VP16

and +VP16 signals are spatially-correlated (**Figure 3B**), but that +VP16 signal intensity is

less uniform, with more dynamic range (**Figure S4C-D**). These differences are likely due to

higher signal-to-noise enabled by enrichment of TOP2 CC following VP16 treatment. Like

yeast Top2 libraries (**Figure 2E**), nucleotide resolution analysis of human CC-seq libraries

displayed a skew towards Watson-Crick read-pairs that are offset by 3 bp, as expected for

the 4 bp 5' overhang generated by TOP2 (**Figure S4E**). This skew is much greater in VP16-

treated samples (**Figure S4E**), supporting the view that +VP16 libraries have greater Top2

CC enrichment.

The human genome encodes two type-IIA topoisomerases, TOP2α and TOP2β
(**Introduction**). To demonstrate the specificity of CC-seq to map human TOP2 activity we
used CRISPR-Cas9 to generate a human RPE-1-derived cell line with homozygous
knockout mutations in the non-essential *TOP2B* gene, and arrested cells in G1 (**Figure S5A**)
when TOP2α is not expressed (Heck et al., 1988; Woessner et al., 1991; Negri et al., 1992).
TOP2α and TOP2β expression was undetectable in such G1-arrested cells (**Figure S5B**),
and importantly, VP16-induced ɣ-H2AX foci were reduced ~7-fold relative to wild type
control lines, indicating that most of the signal is TOP2β-dependent (**Figure S5C-D**).

Next, we applied CC-seq to asynchronous and G1-arrested wild type and *TOP2B$^{-/-}$* cells. In
G1, VP16 exposure induced localised regions of enriched TOP2 CC formation in wild type
cells that were largely diminished or absent in *TOP2B$^{-/-}$* cells (**Figure S5E**). By comparison,
TOP2β deletion did not prevent VP16-induced signal in asynchronous cells (**Figure S5E**), in
which TOP2α is still present. Together, these results indicate that CC-seq detects a mixture
of both TOP2α and TOP2β covalent complexes in wild type human cells depending on the
cell cycle phase.

Spatial distribution of Top2 activity in the human genome

The human genome is divided into a relatively gene-rich 'A' compartment and a relatively
gene-poor 'B' compartment, the 3D spatial segregation of which can be determined by HiC
(Lieberman-Aiden et al., 2009). In both untreated and VP16-treated asynchronous wild type
cells, TOP2 CC are enriched in the active A compartment (**Figure 3B and C**), consistent
with the role of TOP2 in facilitating transcription. TOP2 activity also correlated with regions
of negative DNA supercoiling (Naughton et al., 2013) (**Figure 3B**). These observations
provide a functional link between TOP2 activity, DNA topological stress, and large-scale
chromatin compartments.

We next focused our attention in more detail on the genome-wide pattern of TOP2 activity
enriched by VP16 treatment. Simple visual inspection (**Figure 3D**) suggested that TOP2 CC
signals are enriched around genome features previously identified as regions of high TOP2-
binding and etoposide-induced DNA strand breakage, including sites of CTCF-binding
(Uusküla-Reimand et al., 2016; Canela et al., 2017) and active transcription start sites
(TSSs) marked by H3K4Me3 and H3K27Ac (Baranello et al., 2014; Yang et al., 2015).

Genomic maps of etoposide-induced DNA breaks were recently generated using a related
methodology, 'END-seq' (Canela et al., 2017), that is not specific for protein-linked TOP2
CC. Aggregating END-seq data around strong CC-seq positions revealed concordant peak
signals at broad scale (Figure 3E), demonstrating a general agreement in the positions
reported by the two techniques. However, at finer scale, END-seq positions are offset by 1-
15 bp in the 3' direction relative to CC-seq positions on each strand (Figure 3F)—
something not observed when aggregating CC-seq data around the same loci (Figure 3F).
This important difference is likely to be explained by the different populations of DNA
breaks mapped by each technique: specific TOP2 cleavage positions (CC-seq) versus DNA
breaks that have undergone limited nucleolytic resection (END-seq).

To investigate the relationship between CTCF and TOP2 activity, we filtered CTCF motifs to
include only those that overlapped an RPE-1 CTCF ChIP-seq peak (Methods; (Akimitsu et
al., 2003)). After aligning Watson and Crick motifs in the same orientation, we frequently
observed a strong TOP2 CC peak upstream of the motif centre and accompanied by more
distal, weaker peaks on both sides of the motif (Figure 4A upper panel, and S6A). Loci
containing multiple CTCF-binding motifs display more complex TOP2 CC patterns (Figure
4A upper panel, and S6B) with enrichment on both sides of the double motif, as would be
expected from the aggregation of a heterogeneous population of CTCF and TOP2 activity
present across different cells.

We next stratified CTCF motifs into quantiles based on proximal CTCF binding intensity,
and aggregated CC-seq signals centred on these loci (Figure 4A, middle and bottom
panels). Most Top2 CC signal is concentrated in two peaks located ±~54 bp relative to the
centre of the CTCF motif, with the upstream peak ~2-fold more intense (Fig 4A, middle
and bottom panels). Remaining Top2 CC signal is focused in a flanking array of weaker
peaks. This signal pattern has been reported recently for VP16-induced DSBs (Canela et
al., 2017; Gothe et al., 2018), and has been suggested to result from a role for TOP2 in
facilitating genome organisation by cohesin-dependent loop extrusion operating in the
context of CTCF and other boundary elements. In support of this interpretation, total
aggregated TOP2 CC signal across these regions is positively correlated with CTCF
occupancy as measured by ChIP-seq (Figure 4A, middle and bottom); interestingly
however, TOP2 CCs were not correlated with interaction strength of associated loops
(Figure S6C), annotated previously by Hi-C (Darrow et al., 2016). The positions of TOP2 CC
surrounding CTCF motifs anticorrelate with the well-positioned nucleosomes at these sites

(**Figure 4B**). However, such anticorrelation with nucleosome positioning is not a unique

feature of CTCF loci, but is instead a general property of all TOP2 CC sites (**Figure 4C**),

consistent with maps of both Top2 CC (**Figure S3B**) and Spo11 (**Figure S1E;** (Pan et al.,

2011)) in *S. cerevisiae*.

In human cells, etoposide-induced TOP2 CCs are also enriched around gene TSSs (**Figure**

5), typically concentrated in broad peaks ~1-2 Kbp downstream and ~0.5-1 Kbp upstream

of the TSS (**Figure 5A**; (Yang et al., 2015)). Similar to elsewhere in the genome, in the region

immediately downstream of the TSS (0-400 bp), TOP2 CC signal anticorrelated with

nucleosome occupancy (**Figure S6D**). Stratifying TSSs based on gene expression

microarray data (Ganem et al., 2014) revealed a strong positive correlation between TSS-

proximal TOP2 CC enrichment and gene expression (**Figure 5A, middle and bottom**)—

something not identified when prior etoposide-induced DSB maps were compared to

nascent transcription as measured by GRO-Seq (Canela et al., 2017). Thus, while human

TOP2 activity is correlated with total transcription of the gene, it is not proportional to local

RNA polymerase activity when assayed at finer scale—suggesting a model where, as

postulated for *S. cerevisiae* (**Figure S3A**), superhelical stress is resolved at sites dislocated

from the process that generates them. Indeed, when analysed with an identical pipeline,

both TSS-proximal TOP2 CCs (**Figure 5A**) and etoposide-induced DSBs (Canela et al.,

2017) positively correlate with gene expression (**Figure S6E**), indicating that this correlation

is not due to technical differences in mapping methodologies. Based on the twin-domain

model of DNA supercoiling, topological stress is expected to increase not just with gene

expression level, but also with increasing gene length (Liu and Wang, 1999). In support of

this idea, we observe a strong positive correlation between human TOP2 CC enrichment

and gene length (**Figure S7A, top**) independent from gene expression (**Figure S7A,**

bottom). Interestingly, no such correlation with gene length is present in *S. cerevisiae*

(**Figure S7B**), further supporting a model whereby, in this compact gene-dense genome,

Top2 activity in the IGR is neither directly linked to expression nor length of the proximal

gene.

Notably, the increased density of etoposide-induced human TOP2 CCs at strongly

expressed genes is 2.7-fold greater than at strongly bound CTCF motifs (**Figure 5B**

and C). Thus, whilst strongly-bound CTCF motifs (14,924) greatly outnumber active genes

(4,194)—leading, globally, to more TOP2 activity in total at CTCF sites (**Figure 5D**)—our

data reinforce the important role of topoisomerase activity during transcription.

Local DNA sequence and methylation status direct the formation of Top2 CCs

Previous reports suggest that etoposide induces a majority of SSBs, rather than DSBs, due

to independent poisoning of each active site in the Top2 homodimer (Bromberg et al.,

2003). In our yeast and human datasets, we similarly observe more single-strand Top2 CCs

(90.1% and 94.2%, respectively) than double-strand Top2 CCs (9.9% and 5.8%,

respectively) following etoposide exposure (**Figure 6A**). Interestingly, *MRE11* deletion

increased the proportion of double-strand Top2 CCs (**Figure S3C** 12.9%, z-test p-value <

2.2 x 10^{-16}), consistent with the role of MRN/X in protein-linked DNA repair (Neale et al.,

2005; Garcia et al., 2011; Anand et al., 2016; Cannavo and Cejka, 2014; Cannavo et al.,

2018; Hoa et al., 2016; Hartsuiker et al., 2009).

Prior, fine-scale mapping of Top2 cleavage at specific substrates indicated that cleavage

patterns are influenced by local DNA sequence composition (Pommier et al., 1991;

Borgnetto et al., 1996; Strumberg et al., 1999; Wu et al., 2011). To investigate the generality

of such findings, and to precisely determine how DNA sequence influences Top2 site

preference across the yeast and human genome, we computed the nucleotide resolution

average base composition around Top2 CC SSB and DSB sites (**Fig 6B**)—something made

possible by the positional accuracy of our mapping technique. For DSBs, average DNA

sequence patterns are rotationally symmetrical around the dyad axis of cleavage (**Figure**

6B, left), consistent with the known homodimeric nature of eukaryotic Top2 (Pommier et

al., 2016). Base positions with notable skews are restricted to the region ±11 bp from the

dyad axis (p<10^{-4}, Chi-squared test), consistent with the DNA contacts made by Top2 (Lee

et al., 1989; Thomsen et al., 1990; Wu et al., 2011), yet with minor, species-specific

differences at a few positions (±2, ±7, ±9; **Figure 6C, top**), suggesting subtle differences in

the ScTop2 and HsTOP2 DNA binding surface. Positions -3 and +3 are the bases

immediately 5' to the scissile phosphodiester bonds cleaved by Top2 (**Figure 6B**). In the

averaged DSB sequence motif, strong preference for cytosine at these positions (69.0% of

DSBs) on the scissile strand (**Figure 6B, left)** agrees with *in vitro* studies of Top2

mechanism—attributed to favourable base-stacking interactions of etoposide with

guanosine at the -1 position on the non-scissile strand (Pommier et al., 1991; Strumberg et

al., 1999; Wu et al., 2011).

Importantly, whilst SSBs display similar overall symmetrical sequence features as DSBs,

SSBs differed substantially at position +3, where the strong preference for guanine on the

scissile strand (and paucity of cytosine on the non-scissile strand) is absent (**Figure 6B,**
right, and Figure 6C, bottom). Furthermore, the preference for cytosine at the -3 position
on the scissile strand increases to 88.1% for SSBs. This clear difference in sequence bias
provides strong evidence that such positions truly reflect sites of preferred VP16-induced
SSBs vs DSBs *in vivo*.

The human genome is subject to methylation of cytosine (meC) at CpG sites. Dramatically,
meC completely abolishes the formation of VP16-induced TOP2 CCs (**Figure 6D**),
suggesting this bulky group may interfere sterically with the VP16-Top2-DNA ternary
complex, modifying TOP2 catalysis. Collectively, these observations demonstrate that local
DNA sequence composition—including epigenetic marks—are major determinants of both
TOP2 CC abundance and the differential generation of SSBs and DSBs by VP16 *in vivo*.

Discussion

Topoisomerases are ubiquitous gatekeepers to the genetic material—facilitating changes to
DNA topology, nuclear organisation, and chromosome segregation that are essential for the
generation and propagation of all life on Earth. Yet, topoisomerase dysfunction can also
prove harmful, generating difficult to repair protein-linked DNA breaks that create risks to
genome stability. Thus, understanding how, where, and when topoisomerase activity takes
place is of great importance. Here, we demonstrate a robust, technically simple and
sensitive method to map sites of preferred type IIA topoisomerase activity across the
eukaryotic genomes of yeast and human cells at strand-specific nucleotide resolution. Our
methodology joins an extensive toolkit of complementary NGS solutions (Canela et al.,
2017; Crosetto et al., 2013; Yan et al., 2017), each with its own set of unique advantages.
However, many of these techniques do not achieve nucleotide resolution, either due to the
use of nucleolytic processing to blunt 5' DNA termini (Canela et al., 2017), and/or inability to
directly map DNA breaks covalently linked to protein (Canela et al., 2017; Crosetto et al.,
2013; Yan et al., 2017). Here, by utilising the tyrosyl phosphodiesterase activity of
recombinant Tdp2 to unblock 5'-DNA termini non-nucleolytically, we are able to accurately
map the position of not just Top2 CCs, but also related classes of topoisomerase-like
enzymes such as Spo11, which acts uniquely in meiotic cells. Importantly, such specificity
in our cleavage maps enable us to corroborate and extend—with completely independent
methodology—prior research findings from meiosis (Pan et al., 2011), the impact that VP16
has on SSB versus DSB formation—and for the first time demonstrate the influence that

primary DNA sequence composition and methylation status has on the process of Top2
catalysis *in vivo* in both yeast and human cells.

Top2 activity has long been associated with transcriptional activity (Yang et al., 2015; Brill
et al., 1987; Baranello et al., 2014; Yu et al., 2017). In addition, recent research has
identified etoposide-induced DSBs that colocalise with a subpopulation of chromatin-
bound TOP2 on the loop-external side of CTCF- and Rad21-bound CTCF motifs (Canela et
al., 2017; Uusküla-Reimand et al., 2016). TOP2 activity at these sites has been proposed to
facilitate genome organisation driven by chromatin loop extrusion, and thereby to constitute
a major fraction of the total nuclear TOP2 activity (Canela et al., 2017). Moreover, contrary
to expectations, the frequency of etoposide-induced DSBs was found to be locally
uncorrelated with nascent RNA levels (Canela et al., 2017)—generating some confusion in
the understanding of where TOP2 activity is most prevalent and most important. Here, our
independent CC-seq method and analyses permit us to revisit this question in explicit
detail, and clarify the role of TOP2 in these processes. Importantly, whilst Top2 activity is
certainly highly enriched at CTCF-bound genomic loci, the total amount of activity at any
given CTCF locus is many times lower than observed at transcriptionally active genes.
Nevertheless, it is critical to emphasise that because of the great abundance of CTCF-
bound sites compared to active genes, total aggregated TOP2 activity is globally similar for
CTCF-associated versus transcriptionally active regions of the genome.

Our data also permit us to investigate how other genomic features influence Top2 activity.
For example, at both TSS and CTCF-proximal regions—and indeed in the wider genome of
both yeast and human—Top2 cleavages are highly enriched in nucleosome-free DNA, in
support of prior findings made at a subset of genomic loci (Capranico et al., 1990; Udvardy
and Schedl, 1991; Käs and Laemmli, 1992). Notably, whilst yeast Top2 activity is also
enriched near TSSs, it is concentrated immediately (120-140 bp) upstream, unlike human
TOP2 activity, which is primarily concentrated in a broader peak 1-2 kb downstream, within
the gene body. Furthermore, unlike human TOP2, yeast Top2 activity is uncorrelated with
downstream gene expression. These differences may stem from the very different scale
and transcriptional organisation of the yeast and human genomes, and the strong
patterning of Top2 activity driven by accessibility to nucleosome-free intergenic promoter
regions in *S. cerevisiae*. In support of this interpretation, similar nucleosomal constraints
appear to substantially govern the activity of the related topoisomerase-like enzyme,
Spo11, during meiosis (Pan et al., 2011) (**Figure S1** herein).

Despite these differences in distribution of Top2 activity at this larger scale, the DNA

sequence preferences for yeast and human Top2 are similar, implying substantial

conservation of the DNA binding interface, which is supported by analysis of the published

crystal structure of VP16-TOP2-DNA (Wu et al., 2011). A nucleotide resolution genome-

wide map of *E. coli* gyrase activity was recently reported (Sutormin et al., 2018)), revealing a

sequence motif composed of a central 35 bp region containing the cleavage site, plus two

flanking 47 bp regions with ~10.5 bp periodicity centred ~±40 bp from the site of cleavage,

which are attributed to the known wrapping of proximal DNA around the gyrase C-terminal

domains (Reece and Maxwell, 1991). Strikingly, whilst we observe a core central motif for

Top2, there are no such flanking motifs over these ranges, in agreement with its lack of

DNA wrapping domains (**Figure S3D**). We do, however, observe two more-proximal

flanking regions centred ~±20 bp from the site of cleavage, each with ~10.5 bp periodicity

(**Figure S3E**), suggesting that Top2 also bends its DNA substrate—in agreement with a

recent biophysical study (Huang et al., 2017).

We have used CC-seq to probe MRX-dependent repair of Top2 CCs in *S. cerevisiae*,

corroborating and extending prior genetic experiments in this organism (Hamilton and

Maizels, 2010; Stepanov et al., 2008; Garcia et al., 2011), and others (Nakamura et al.,

2010; Hoa et al., 2016; Hartsuiker et al., 2009). Importantly, calibration of our signal using

an internal human standard enables the relative quantification of Top2 CCs, and thus

directly demonstrates for the first time that loss of Mre11 activity increases cellular levels of

Top2 CCs in *S. cerevisiae*. The advantage of the antigen-independent enrichment step is

three-fold. Firstly, the spike-in standard contains human TOP2 CCs, which are enriched to

a similar extent as the yeast Top2 CCs, and should therefore be less susceptible to

variation than background DNA. Secondly, the spike-in standard requires no genetic

epitope-tagging or reconstitution of recombinant protein-DNA complexes, unlike ChIP-seq

calibration methods (Hu et al., 2015; Grzybowski et al., 2015). Finally, we believe that this

calibration step could be performed using any other organism with a suitably divergent

genome. Thus, the capacity of CC-seq to quantify as well as map CCs between samples

makes this technique ideally suited to the future study of repair dynamics.

We have demonstrated that CC-seq maps CCs of both human TOP2α and TOP2β—

proteins that are conserved amongst all vertebrates and whose functional differences

remain a subject of interest within the field (Austin et al., 2018). Given the essential

requirement for TOP2α during mitosis (Akimitsu et al., 2003), and the apparent inability of TOP2β to compensate for its loss (Grue et al., 1998), it will be particularly interesting to determine the genomic distribution of TOP2α/TOP2β activities throughout the cell cycle, and to relate this to 3D chromatin conformation measured by Hi-C. Whilst not essential for cell proliferation, TOP2β plays an important role in promoting transcriptional programmes associated with neuronal development (Tiwari et al., 2012), and this function cannot be supported by TOP2α. Furthermore, loss of TDP2 function inhibits TOP2-dependent gene transcription and leads to neurological symptoms including intellectual disability, seizures and ataxia (Gómez-Herreros et al., 2014). Thus, accurate genomic maps of TOP2β CC formation in these tissues will help to identify those genes most pertinent to the development of neurodegeneration. Moreover, as demonstrated by our Spo11 maps, the antigen-independent, high-efficiency protein-DNA enrichment process makes CC-seq generally applicable for mapping not just Top2 activity in diverse organisms, but also for mapping similar types of topoisomerase-DNA covalent complexes, or even those less distinct DNA-protein crosslinks that arise upon ablation of the DNA-dependent protease SPRTN/Wss1 (Stingele et al., 2015).

Acknowledgements

We thank Keith Caldecott, Antony Oliver and Peter Hornyak for sharing recombinant TDP2, and Jon Nitiss for sharing the *pdr1Δ* strains.

Author Contributions

W.G., D.J. and M.J.N. conceived the project and developed the CC-seq methodology. W.G., D.J., H.T. and R.A. generated whole genome CC-seq libraries. W.G. performed all data processing and analysis, and all human cell work. D.J., H.T. and R.A. performed yeast sensitivity and meiotic DSB assays. T.J.C. developed the mapping pipeline and associated tools. W.G. and M.J.N. interpreted the observations and wrote the manuscript.

Competing interests

The authors declare no competing financial interests.

Funding statement

W.G., D.J., R.A., T.J.C., and M.J.N. were supported by an ERC Consolidator Grant (#311336), the BBSRC (#BB/M010279/1) and the Wellcome Trust (#200843/Z/16/Z).

Data availability

491 All strains and cell lines listed in Table S3 and S4 are available on request. Sequencing
492 reads are available via the NCBI Sequence Read Archive (accession numbers pending).

Methods

Yeast strains, culture methods and treatment

The *Saccharomyces cerevisiae* yeast strains used in this study are described in **Table 2** and
were derived using standard genetic techniques. Strains used for Spo11 mapping are
isogenic to the SK1 subtype and carry the *sae2Δ::kanMX* gene disruption allele. Strains
used for Top2 mapping are isogenic to the BY4741 subtype and carry the *pdr1DBD-CYC8*
drug sensitivity cassette (Stepanov et al., 2008). For Spo11 DSB mapping, cells were
induced to undergo synchronous meiosis as follows: Cells were grown overnight to
saturation in 4 ml YPD medium (1% yeast extract, 2% peptone, 2% glucose supplemented
with 0.5 mM adenine and 0.4 mM uracil) at 30 ℃, then diluted to OD_{600} 0.2 in 200 ml YPA
medium (1% yeast extract, 2% peptone, 1% potassium acetate) and grown vigorously for
15 h at 30 ℃. Cells were then washed with water, resuspended in 200 ml sporulation
medium (2% potassium acetate supplemented with diluted amino acids), incubated
vigorously for 6 h at 30 ℃ and harvested by centrifugation. For etoposide treatment, cells
were grown overnight to saturation in 4 ml YPD medium at 30 ℃, then diluted to OD_{600} 0.5
in 100 ml YPD and grown until OD_{600} 2. 50 ml cultures were then incubated for a further 4 h
in the presence of either 1 mM etoposide or 2% DMSO, before harvesting by
centrifugation.

Human cell lines, culture methods and treatment

The human cell lines used in this study are described (**Supplementary Table 3**). Human
hTERT RPE-1 cells were obtained from ATCC and cultured at 37 ℃, 5% CO_2 and 3% O_2 in
Dulbecco's Modified Eagle's Medium DMEM/F-12 (Sigma), supplemented with 10% Fetal
Calf Serum (FCS) and 90 Units/ml Penicillin-Streptomycin. For all experiments (CC-seq,
Slot Blot, WB, IF, FACs) with asynchronous cell populations, RPE-1 cells were seeded at a
density of (3.5×10^3 cells/cm^2l) and incubated for 72 h at 37 ℃, to ensure subconfluent log-
phase growth (~70% confluency) at the time of the experiment. For G1 cell populations, WT
and *TOP2B*$^{-/-}$ RPE-1 cells were seeded at a density of 5×10^3 cells/cm^2 and incubated for 48
h at 37 ℃ in DMEM/F12 containing 10% FCS, prior to a further 24 h incubation in

DMEM/F12 medium containing no FCS to ensure complete G1-phase arrest (verified by FACs, see below). For experiments with proteasome inhibitor and etoposide (CC-seq, Slot Blot), cells were preincubated with 5 µM MG132 (Sigma) for 90 min, trypsinised and incubated in suspension with 5 µM MG132 and 100 µM etoposide (Sigma) for 20 min at 37 ℃. For experiments with etoposide alone (IF), adherent cells were treated with 100 µM etoposide for 20 min at 37 ℃.

Generation of *TOP2B*^{-/-} RPE-1 cells

The oligonucleotides 5'-CACCGCCGCAGCCACCCGACT and 5'-AAACAGTCGGGTGGCTGCGGC (identified using Benchling; https://benchling.com) were annealed and cloned into pX330 following BbsI restriction, as described previously (Hsu et al., 2013). This SpCas9/ trugRNA co-expression plasmid was transiently expressed in RPE-1 cells to target the 17 bp target sequence GCCGCAGCCACCCGACT (TGG) within exon 1 of *TOP2B*. Single clones were trypsinised and passaged to isolated culture vessels prior to screening for absent protein by Western Blot (WB). All experiments involving *TOP2B*^{-/-} were conducted using the RPE-1 clone T2B/6, in which no TOP2β is detectable by WB or IF.

Spot tests of chronic etoposide sensitivity

Single colonies of each *S. cerevisiae* strain were incubated overnight in 4 ml YPD at 30 ℃ with shaking. 0.1 ml of this starter was used to inoculate 4 ml YPD, prior to incubation for 5 hours at 30 ℃. Cultures were diluted to make a stock with an OD_{600} of 2.0, then this was 10-fold serially diluted five times. Each dilution was spotted onto plates containing 0, 0.1, 0.3 or 1.0 mM VP16, prior to incubation for three days at 30 ℃. Plates were imaged on a 2400 Photo scanner (Epson).

Southern blotting of meiotic Spo11 DSBs

Approximately 2 µg of genomic DNA (isolated by non-proteolysing Phenol-Chloroform extraction, as described below) was digested at 37 °C overnight using *Pst*I restriction enzyme (NEB) in NEBuffer 3.1 (100 mM NaCl, 50 mM Tris Base·HCl pH 7.9, 10 mM $MgCl_2$, 100 µg ml-1 BSA). Additional *Pst*I was added for 4 hours before the addition of NEB purple loading dye to 1×. Digested samples were proteolysed using 1 mg ml-1 Proteinase K (Sigma) at 60 ℃ for 30 minutes, left to reach room temperature before 10 µg was loaded on a 0.7% 1× TAE agarose gel (40 mM Tris Base·HCl, 20 mM glacial acetic acid, 1 mM EDTA pH 8.0) containing 50 µg ml-1 ethidium bromide. DNA was separated in 1× TAE at 60 V for 18 hours. The gel was imaged using InGenius (Syngene) bioimaging system to check

migration and then exposed to 180 mJ/m^2 UV in the Stratalinker (Stratagene). The gel was

then soaked in three times its volume of denaturation solution (0.5 M NaOH, 1.5 M NaCl) for

30 minutes and then transferred to Zetaprobe (Bio-Rad) membrane by means of a vacuum

at 55 mBar for 2 hours. After transfer the membrane was washed in water ten times and

then cross-linked by exposing the membrane to 120 mJ/m^2 UV in the Stratalinker. The

membrane was incubated in 30 ml of hybridisation buffer (0.5 M NaHPO$_4$ buffer pH 7.5, 7%

SDS, 1 mM EDTA, 1% BSA) at 65 ˚C for 1 hour. The *MXR2* probe for looking at the

HIS4::LEU2 locus was created from 50 ng of template DNA, 0.1 ng of Lambda DNA (NEB)

digested with *BstE*II (NEB), and water. The mix was denatured at 100 °C for 5 minutes then

put on ice. High Prime (Roche) was added in addition to 0.5-3 mBq of α-^{32}P dCTP and

incubated at 37 °C for 15 minutes. 30 μl 1× TE was added and the probe spun through a G-

50 spin column (GE Healthcare) at 400 × *g* for 2 minutes. The probe was then denatured by

incubating at 100 °C for 5 minutes and then put on ice before being added to 20 ml

hybridisation mixture. The original 30 ml hybridisation buffer was discarded and the 20 ml

containing the probe was added to the membrane and incubated overnight at 65 ˚C. After

incubation, the membrane was washed five times with 100 ml pre-warmed Southern wash

buffer (1% SDS, 40 mM NaHPO$_4$ buffer pH 7.5, 1 mM EDTA) and exposed to phosphor

screen overnight.

Western blotting

Whole human cell extracts (WCE) were harvested by direct lysis in 1x Laemmli loading

buffer, denatured for 10 min at 95 ˚C and sonicated for 30 s using Bioruptor® Pico.

Samples were subjected to SDS-PAGE (7% or gradient gel) and transferred to

nitrocellulose membrane. Primary immunodetection was with antibodies targeting TOP2β

(Clone 40, BD Biosciences), TOP2α (ab52934, Abcam), or Ku80 (ab80592, Abcam).

Secondary immunodetection was with HRP-conjugated Rabbit anti-Mouse IgG

(ThermoFisher), prior to detection of peroxidase activity using ECL reagent and X-Ray film

(Scientific Laboratory Supplies Ltd).

Slot blotting

Samples were diluted 4-fold (500 uL total volume) in NaPO$_4$ buffer (25 mM, pH 6.5), and slot

blotted onto 0.2 μM nitrocellulose membrane (Amersham), using the Minifold I (Whatman)

manifold. The wells were washed twice with 750 μL NaPO$_4$ buffer. The membrane was then

blocked with 10% milk-TBST for 1 h at RT, prior to incubation overnight with anti-TOP2β

antibody (Clone 40, BD Biosciences) at 4 ˚C. The membrane was washed 4 times with

TBST, incubated with HRP-conjugated Rabbit anti-Mouse IgG (ThermoFisher) for 1 h at RT,

washed 4 times with TBST, and incubated with ECL detection reagent for 1 min. X-Ray film

was used for detection.

Fluorescence-Assisted Cell Sorting (FACS)

Approximately 10 million RPE-1 cells were trypsinised, washed once in PBS and

resuspended in 1.5 ml PBS. 3.5 ml ethanol was added dropwise to pellet, with vortexing.

Cells were fixed for 1 hr at 4 °C, prior to centrifugation and aspiration of the supernatant.

Cells were washed twice with PBS, prior to resuspension in 0.5 ml 0.25% Triton-X100-PBS

for 15 min on ice. Cells were pelleted by centrifugation, supernatant was aspirated, and the

pellet was resuspended in 0.5 ml TBS containing 10 ug/ml RNase A (Sigma) and 167 nM

Sytox Green (ThermoFisher). After 30 min incubation in the dark at RT, the suspension was

filtered through fine mesh into test tubes. DNA content in 50,000 cells was analysed using

the Accuri C6 (BD Biosciences), with gating to exclude doublets and cell debris.

Immunofluorescence

Cells were seeded onto glass coverslips (Agar Scientific), incubated and treated according

to the protocol outlined above. Cells were then fixed with 4% paraformaldehyde PBS for 10

min, washed three times with PBS, permeabilised with 0.2% Triton-X100-PBS for 10 min,

blocked for 1 h with 10% FCS-PBS, incubated with primary antibodies targeting Phospho-

Histone H2AX (S139) (JBW-301, Merck Millipore) and TOP2α (ab52934, Abcam), washed

three times with PBS, incubated with secondary antibodies Alexa 488-conjugated Goat

anti-mouse IgG (Fisher) and Alexa 647-conjugated Goat anti-Rabbit (Fisher), washed three

times with PBS, washed once with distilled water, and mounted with VECTASHIELD

containing DAPI (Vector Laboratories).

High-content microscopy

Automated wide-field microscopy was performed on an Olympus ScanR system (motorised

IX83 microscope) with ScanR Image Acquisition and Analysis Software, 40x/0.6

(LUCPLFLN 40x PH) dry objectives and Hamamatsu ORCA-R2 digital CCD camera

C10600. Numbers of anti-phospho-Histone H2AX (S139) foci (Alexa 488; FITC filter) were

quantified in the nuclear region colocalising with DAPI, using Olympus ScanR Analysis

software. TOP2α signal (Alexa 647; Cy5 filter) was also quantified in this region.

CC-seq: Enrichment of protein-linked DNA

1.5×10^7 RPE-1 cells were treated as described above, pelleted by centrifugation, washed

once with 15 ml ice-cold PBS, and resuspended in three aliquots of 400 µL ice-cold PBS.

1×10^9 yeast cells were treated as described above, then spheroplasted in 1.5 ml

spheroplasting buffer (1 M sorbitol, 50 mM NaHPO₄ buffer pH 7.2, 10 mM EDTA) containing

200 µg/ml Zymolyase 100T (AMS Biotech) and 1% β-mercaptoethanol (Sigma) for 20 min at

37 ℃. 2 µL Protease Inhibitor Cocktail and 2 µL Pefabloc (Sigma) were added before

splitting into five aliquots of 400 µL. All subsequent steps of the protocol are the same for

yeast and human samples. 1 ml ice-cold ethanol was added to 400 µL cell suspensions in

microcentrifuge tubes, mixed, incubated for 10 min on ice, and pelleted by centrifugation.

Supernatant was thoroughly removed by aspiration, prior to addition of 200 µL 1x STE

buffer (2% SDS, 0.5M Tris pH 8.1, 10 mM EDTA, 0.05% bromophenol blue), cell disruption

using a pestle (VWR), addition of a further 400 µL 1x STE buffer, and incubation for 10 min

at 65 ℃. Samples were cooled on ice and 500 µL Phenol-Chloroform-isoamyl alcohol

(25:24:1; Sigma) added. Mixtures were emulsified by shaking and pipetting 5 times with a 1

ml micropipette, prior to phase separation by centrifugation at 20,000 g for 20 min. By

minimising mechanical shearing of the lysate prior to phenol chloroform extraction,

peptides that are covalently linked to high molecular weight DNA partition into the aqueous

phase. 500 µL of the aqueous phase was removed to a clean microcentrifuge tube and

nucleic acids precipitated with 1 ml ice-cold ethanol, pelleted by centrifugation, washed

with ice-cold 70% ethanol, and dissolved in TE buffer overnight at 4 ℃. Samples were then

incubated with 0.2 mg/ml RNase A (Sigma) for 1 h at 37 ℃; nucleic acids were precipitated

with 1 ml ethanol, pelleted by centrifugation, washed twice with 70% ethanol and dissolved

in TE overnight. Aliquots were combined to a total of 1 ml and sonicated to an average

fragment size of 300-400 bp with Covaris (duty cycle: 10%, intensity/peak power incidence:

75W, cycles/burst: 200, time: 15 min). 1 ml of sonicated sample was added to 1.2 ml of

binding buffer (10 mM Tris pH 8.1, 10 mM EDTA, 0.66 M NaCl, 0.22% SDS, 0.44% N-

Lauroylsarcosine sodium salt). Each sample was divided over several Miniprep (QIAGEN)

silica-fibre membrane spin columns, such that the total DNA loaded to each was

approximately 20 µg. The flowthrough was reapplied to the column to improve binding.

Columns were washed 6 times with 600 µL of TEN (10 mM Tris, 1 mM EDTA, 0.3 M NaCl)

per 1 min wash, prior to elution with 100 µL TES (10 mM Tris, 1 mM EDTA, 0.5% SDS).

659   CC-seq: DNA end repair and adapter ligation

Eluted products were pooled to 500 µL in TES and incubated with 1 mg/ml Proteinase K

(Sigma) for 30 min at 60 ℃, prior to overnight ethanol precipitation at -80 ℃ with 1.41 ml

ethanol, 0.2 mg/ml glycogen and 200 mM NaOAc. The DNA-glycogen precipitate was

pelleted by centrifugation at 20,000 g for 1 hr at 4 ℃, washed once with 1.5 ml 70%

ethanol, and re-pelleted by centrifugation. The supernatant was aspirated and the pellet

was air-dried for 10 min at RT, prior to solubilisation in 52 µL 10 mM Tris-HCl. DNA

concentration was measured in a 2 µL sample with the Qubit (ThermoFisher) and High

Sensitivity reagents. The remaining 50 µL was used as input for one round of end repair

and adapter ligation with NEBNext Ultra II DNA Library Preparation kit (NEB), according to

manufacturer's instructions, except for the use of a custom P7 adapter (see table). The use

of custom adapters is to allow differentiation of the sheared end (P7 adapter) from the

Top2/Spo11 end (P5 adapter). After ligation of the P7 adapter, DNA was isolated with

AMPure XP beads (Beckman Coulter) according to manufacturer's instructions (beads:input

of 78:90) and eluted in 50 µL 10 mM Tris-HCl. Samples were diluted 2-fold with 50 µL Tdp2

reaction buffer (100 mM TrisOAc, 100 mM NaOAc, 2 mM MgOAc, 2 mM DTT, 200 µg/ml

BSA) and incubated with 3 µL of 10 µM recombinant human TDP2 (Hornyak et al., 2016;

Johnson et al., 2019), for 1 h at 37 ℃. DNA was isolated again with AMPure XP beads

(beads:input of 103:103) and eluted in 52 µL 10 mM Tris-HCl. Next, a second round of end

repair and adapter ligation was conducted, using adapter P5 (see table). After ligation of

the P5 adapter to the Top2/Spo11-cleaved end, DNA was isolated with AMPure XP beads

(beads:input of 78:90) and eluted in 17 µL 10 mM Tris-HCl.

CC-seq: PCR and size selection

DNA concentration was measured in 2 µL using the Qubit. The remaining 15 µL was used

as template for the PCR step of the NEBNext Ultra II PCR step using universal primer (P5

end) and indexed primers for multiplexing (P7 end), according to manufacturer's

instructions. PCR reactions were diluted with 50 µL 10 mM Tris-HCl, DNA was isolated with

AMPure XP beads (beads:input of 84:100) and eluted in 30 µL 1 mM Tris-HCl pH 8.1.

Samples were then subjected to 200-600 bp size selection using the BluePippin (Sage

Science), prior to quantification of molarity using the Bioanalyzer (Agilent).

CC-seq: NGS and data pipeline

Multiplexed library pools were sequenced on the Illumina MiSeq (Kit v3 - 150 cycles) or

Illumina NextSeq 500 (Kit v2 - 75 cycles), with paired-end read lengths of 75 or 42 bp,

respectively. Paired end reads that passed filter were aligned using bowtie2 (options: -X

1000 --no-discordant --very-sensitive --mp 5,1 --np 0), using MAPQ0 settings for yeast or

MAPQ10 settings for human experiments, then SAM files processed by the custom-built

Perl program termMapper that computes the coordinates of the protein-linked 5'-terminal

nucleotide. The reference genomes used in this study are hg19 (human), and Cer3H4L2 (*S.*

cerevisiae), which we generated by inclusion of the *his4*::*LEU2* and *leu2*::*hisG* loci into the

Cer3 yeast genome build. Yeast data sets were filtered to exclude long terminal repeats,

retrotransposons, telomeres, and the rDNA. Human datasets were filtered to remove known

ultra-high signal regions (Hoffman et al., 2013; ENCODE, 2012) and repeat regions

(ENCODE, 2012). All subsequent analyses were performed in R (Version 3.4.3) using

RStudio (Version 1.1.383), unless indicated otherwise.

CC-seq: Calibrated library generation

Calibration of Top2 CC-seq experiments was conducted to allow comparison of relative

peak intensities in different yeast strains. This was achieved by spike-in of human DNA

following the sonication stage of the protocol, which is a method that has been used to

calibrate other sequencing methods (Hu et al., 2015; Grzybowski et al., 2015). S. cerevisiae

and RPE-1 cells were exposed to etoposide and processed until just after sonication,

exactly as according to the cell treatment and CC-seq protocols above. DNA concentration

was quantified by Qubit, and then mixed at a molar ratio of human DNA:yeast DNA of

1:100. All subsequent stages of the protocol were identical, except read alignment, for

which we used both hg19 and Cer3H4L2 builds successively. Cer3H4L2-aligned peak

heights were corrected in each sample by multiplying by the reciprocal fraction of human

reads in that sample.

CC-seq: Fine-scale mapping

Fine-scale (nucleotide resolution) maps of Spo11/Top2 CCs were produced as simple

histograms over a specified region (Figures 1B, 2B, 3D, 4A, 5A, S1A, S3A, S5A and S5B).

Dark red and blue line heights indicate numbers of 5'-terminal nucleotides detected at that

position on the Watson and Crick strands, in units of HpM. Where indicated in the figure

caption, smoothed data are also plotted as pale red and blue polygons, in addition to the

unsmoothed nucleotide resolution data. This smoothing was either applied using a sliding

Hann window of the indicated width, or using a custom smoothing function (VarX), as

indicated in the figure caption.

CC-seq: Broad-scale mapping

Broad-scale maps of Top2 CCs were produced by binning nucleotide resolution data at 10

Kbp (Figure 3B) or 100 Kbp (Figure S4D) resolution. Binned data were either plotted

directly (**Figure S4D**); or first scaled according to the estimated noise fraction (see below),

smoothed with a 10-bin Hanning window, and median subtracted (**Figure 3B**).

CC-seq: Estimation of the noise fraction in Human datasets

The noise fraction in each sample was estimated using an adaptation of the previously

published NCIS method (Liang and Keleş, 2012). Briefly: the data for -VP16 and +VP16

samples were first binned at 10 Kbp resolution. Then the subpopulation of bins with the

lowest TOP2 CC signal was identified in each sample. The average signal density of this

subpopulation of bins, in each sample, was defined as the noise density (d_{-VP16} and d_{+VP16}).

Signal in the -VP16 sample was scaled by a normalisation factor equal to:

742
$$r = \frac{d_{+VP16}}{d_{-VP16}}$$

CC-seq: Quantification of Spo11 and Top2 activity in defined genomic regions

Yeast Spo11 hotspots were the same as defined previously (Pan et al., 2011). Yeast

intra/intergenomic regions were defined based on TSS and TTS coordinates reported on

the Saccharomyces Genome Database (https://www.yeastgenome.org). Human chromatin

compartments A and B were defined based on eigenvector analysis of previously published

100 kb resolution RPE-1 Hi-C data (Darrow et al., 2016) using the Juicer package

(Lieberman-Aiden et al., 2009; Durand et al., 2016). Human "CTCF" and "TSS" regions were

defined as 10 kb regions centred on the top quartile of expressed TSSs or the midpoint of

top quartile CTCF-bound CTCF-binding motifs, with overlaps merged using

GenomicFeatures::reduce. The "Both" region was defined as the intersection of the "TSS"

and "CTCF" regions using GenomicFeatures::intersect. This region was then excluded from

the individual "TSS" and "CTCF" regions. The "Neither" region was defined as regions of

hg19 which overlapped neither "TSS", nor "CTCF", nor "Both", using

GenomicFeatures::gaps. Nucleotide resolution data was tallied within defined regions, and

is expressed as an aggregate (bar plots) in which the signal is corrected using the average

signal in the "Neither" compartment, and/or as the distribution of region signal densities

(box-and-whisker plot), as indicated in figure captions.

CC-seq: Correlation of Watson/Crick cleavage positions

Nucleotide resolution human TOP2 data was thresholded at 0.01 HpM. Nucleotide

resolution yeast Top2 and Spo11 data were not thresholded. Peak coordinates on the Crick

strand were offset over the range -100 to +100, relative to Watson coordinates. After each

offset, the data was filtered to include only sites with both Watson and Crick hits. The

Pearson correlation (*r*) between the Watson and Crick signal intensities in these *n* sites was

calculated. We also counted the fraction of reads found within these *n* sites, and

normalised this number over the -100 to +100 bp range. Data are expressed as R values

and/or normHpM values, as indicated in the figure captions.

CC-seq: Aggregation of CC-seq data around loci of interest

Nucleotide resolution CC-seq data (line-plots) or binned CC-seq data (heatmaps) were

aggregated within regions of specified size, centred on the loci of interest. The resulting

sum total HpM was divided by numbers of loci to give a mean HpM per locus.

CC-seq: Loci of interest used in this study

Yeast TSS and TTS coordinates were obtained from the Saccharomyces Genome

Database https://www.yeastgenome.org. These were stratified based on associated gene

length, expression level in vegetative SK1 (GSM907178, GSM907179, GSM907180), or

expression level in meiotic SK1 (GSM907176, GSM907177; (Dominissini et al., 2012)).

Human TSS and TTS coordinates were obtained from the UCSC hg19 knownGene

annotation, and stratified based on associated gene length, or expression level

(GSM1395252, GSM1395253, GSM1395254; (Ganem et al., 2014)). Occupied RPE-1 CTCF

motifs were identified as follows: The FIMO tool (Grant et al., 2011) and the CTCF Position

Weight Matrix (PWM) from the JASPAR database (Mathelier et al., 2014) were used to find

all significant hg19 CTCF motifs (p < 1x10^{-4}). These were filtered to include only those

which overlapped positions of RPE-1 CTCF ChIP-seq peaks (GSM749673, GSM1022665;

(ENCODE, 2012)), and stratified based on this ChIP-seq data. RPE-1 loop anchor-

associated CTCF motifs were identified using the Juicer MotifFinder (Durand et al., 2016)

with the RPE-1 WT Hi-C looplist (GSE71831(Darrow et al., 2016)) and CTCF ChIP-seq BED

file (GSM749673, GSM1022665; (ENCODE, 2012)) as input. Human TOP2 CC-seq peak

coordinates were identified by thresholding (0.05 HpM) of pooled +VP16 data presented.

CC-seq: Spatial correlation of yeast Top2 and Spo11 signals

Nucleotide resolution Top2/Spo11 CC-seq data was binned at 50 bp resolution and the

Pearson correlation was calculated between the sum HpM for all bins at a given inter-bin

distance.

CC-seq: Quantifying single and double-strand Top2 CCs

Nucleotide resolution Human and Yeast Top2 CC-seq data were first thresholded at 0.05 and 1 HpM, respectively. SSB sites were defined as those sites without a cognate on the opposite strand at the expected offset of 3 bp. DSBs were defined as cognate sites with a 3 bp offset. SSB and DSB percentages are expressed as a percentage of the total number of sites (SSB + DSB). A randomisation experiment was conducted in order to estimate expected numbers of SSBs and DSBs within the sample under a random model where signal is distributed independently on each strand. To achieve this, the amplitudes of Top2 CCs (HpM) were shuffled amongst the positions in the nucleotide resolution datasets, prior to thresholding and offset analysis as described above.

CC-seq: DNA sequence composition of single and double-strand Top2 CCs

Nucleotide resolution Human and Yeast Top2 CC-seq data were first thresholded at 0.05 and 1 HpM, respectively. The high frequency positions remaining were classified as part of a single or double-stranded Top2 CC based on absence or presence of a cognate cleavage at the expected W-C offset of 3 bp. For double-stranded Top2 CCs, a dyad axis coordinate was defined as the centrepoint between the Top2-linked nucleotides on the W and C strand (that is, the midpoint of the central two base pairs in the four base pair overhang). For single-stranded Top2 CCs, we used an imaginary dyad axis in the same relative position. DNA sequence was aggregated ±20 bp around these two classes of Top2 CCs. Statistical significance was determined using the one sample (goodness-of-fit) Chi-squared test, as described previously (Pommier et al., 1991).

CC-seq - MethylC Analysis

A publically-available RPE-1 reduced representation bisulphite sequencing (RRBS) dataset (two replicates) was used to define coordinates of nucleotides immediately 3'-relative to 919,817 unmethylated or 341,136 methylated Cs, based on <10% or >90% methylation. Next we aggregated RPE-1 CC-seq (0.01 HpM thresholded) hits on these loci, and divided by the number of loci to give a density (barplot). Statistical significance was determined by the Kolmogorov-Smirnov test.

Ideograms

Human chromosome ideograms were adapted from the open source ideogram package (https://eweitz.github.io/ideogram/).

Figure Legends

Figure 1. CC-seq maps covalent Spo11-linked DNA breaks in *S. cerevisiae* meiosis with nucleotide accuracy

A) Schematic of the CC-seq method.

B) Column-based enrichment of Spo11-linked DNA fragments detected by Southern blotting at the *his4*::*LEU2* recombination hotspot (**Methods**). Arrows indicate expected sizes of Spo11-DSBs.

C) Nucleotide resolution mapping of *S. cerevisiae* Spo11 hotspots by CC-seq or oligo-seq (Pan et al., 2011). Red and blue traces indicate Spo11-linked 5' DNA termini on the Watson and Crick strands, respectively. Grey arrows indicate positions of gene open reading frames.

D) Pearson correlation (r) of Spo11 CC-seq signal between Watson and Crick strands, offset by the indicated distances.

E) Average nucleotide composition over a 200 (top) and 30 bp (bottom) window centred on Spo11 breaks. Bases reported are for the top strand only.

HpM = Hits per million mapped reads per base pair.

Figure 2. CC-seq maps covalent Top2-linked DNA breaks in *S. cerevisiae* cycling cells with nucleotide accuracy

A) Serial dilution spot tests of VP16 tolerance for the indicated strains.

B) Nucleotide resolution mapping of *S. cerevisiae* Top2 CCs by CC-seq of the indicated strains after treatment for 4 hours with 1 mM VP16. Spo11-CC data is plotted for comparison. Top2 CC data were calibrated using a human DNA spike-in (**Methods**). Red and blue traces indicate CC-linked 5' DNA termini on the Watson and Crick strands, respectively. Grey arrows indicate positions of gene open reading frames. Lower panels show an expanded view of the region from 31.5 to 40 Kbp.

C) Quantification of Top2 and Spo11 CC signal stratified by genomic region. The genome was divided into intra and intergenic regions; the intergenic region was further divided into divergent, tandem and convergent based on orientation of flanking genes. Spo11 and Top2 activity mapped by CC-seq is expressed as box-and-whisker plots of density (upper and lower box limit: 3rd and 1st quartile; bar: median; upper and lower whisker: highest and lowest values within 1.5-fold of the interquartile range), or as the percentage of total mapped reads.

D) Local correlation of Top2 or Spo11 CC-seq signals. Top2 or Spo11 CC-seq data were binned at 50 bp resolution and the Pearson correlation calculated between bins of increasing separation.

E) Pearson correlation (r) of Top2 CC-seq signal on Watson and Crick strands, offset by the indicated distance.

HpM = Hits per million mapped reads per base pair.

Figure 3. CC-seq maps TOP2-linked DNA breaks in Human cells with nucleotide accuracy

A) Anti-TOP2β western slot blot of input, flow through, wash, and elution fractions. RPE-1 cells were treated or not with VP16, prior to processing according to **Figure 1A**.

B) Broad-scale maps of *H. sapiens* TOP2 CCs produced by CC-seq in RPE-1 cells ±VP16. Raw data were scaled, binned, smoothed and median subtracted prior to plotting (**Methods**). Chromatin compartments revealed by Hi-C eigenvector analysis (Darrow et al., 2016), and supercoiling revealed by bTMP ChIP-seq (Naughton et al., 2013) are shown for comparison.

C) Quantification of TOP2 CC in chromatin compartments A and B. Data are expressed as box-and-whisker plots of density as for **Figure 2C**.

D) Fine-scale mapping of *H. sapiens* TOP2 CCs by CC-seq in RPE-1 cells ±VP16. Red and blue traces indicate TOP2-linked 5' DNA termini on the Watson and Crick strands, respectively. Pale shaded areas are the same data smoothed according to local density (see **Methods**). RPE-1 CTCF and H3K4Me3 ChIP-seq data plus H3K27Ac ChIP-seq data overlaid from seven cell lines (ENCODE, 2012) is shown for comparison.

885 **E)** Medium-scale aggregate of END-seq mapped DSBs (Canela et al., 2017) surrounding nucleotide
886 resolution CC-seq mapped TOP2 CCs.
887 **F)** Fine-scale strand-specific aggregate of END-seq mapped DSB ends (red and blue bars) and
888 nucleotide resolution CC-seq mapped TOP2 CCs (grey bars) surrounding strong TOP2 CC sites.
889

Figure 4. CTCF-proximal TOP2 activity is tightly confined within nucleosome-depleted regions
A) Aggregation of TOP2 CCs in a 1 Kbp window centred on orientated CTCF motifs in human RPE-1
cells. Four example CTCF loci are shown orientated in the 5'-3' direction (top). A heatmap of all
CTCF-motifs in the human genome, with 25 rows stratified by the strength of colocalising CTCF
ChIP-seq peaks in RPE-1 cells (middle). The colour scale indicates average TOP2 CC density. Motifs
are also stratified into 4 quartiles of CTCF-binding, and the average TOP2 CC distribution in each
quartile plotted (bottom).
B) Aggregated TOP2 CC distribution (red line) in the highest quartile of CTCF-binding compared with
the average MNase-seq signal (blue line).
C) Aggregated TOP2 CC distribution (black and red lines: single-nucleotide resolution and
smoothed, respectively) and the average MNase-seq signal (blue line) surrounding strong TOP2 CC
sites.
In (B) and (C) peaks in MNase-seq signal indicate inferred nucleosome positions (blue circles).
903

Figure 5. TSS-proximal TOP2 activity is strongly correlated with gene transcription
A) Aggregation of TOP2 CCs in a 10 Kbp window centred on orientated TSSs in human RPE-1 cells.
Four example TSSs are shown orientated in the 5'-3' direction (top). A heatmap of all TSSs in the
human genome, with 25 rows stratified by gene expression level in RPE-1 cells (middle). The colour
scale indicates average TOP2 CC density. Motifs are also stratified into 4 quartiles of gene
expression, and the average TOP2 CC distribution in each quartile plotted (bottom).
B) Comparison of CTCF-proximal and TSS-proximal TOP2 CC distributions in the highest quartile of
CTCF-binding (pink) and gene-expression (blue), respectively.
C) Average TOP2 CC density in 10 Kbp regions centred on the highest quartile of TSS, CTCF, or
regions where both features are present. Data are expressed as box-and-whisker plots of density as
in **Figure 2C**. Statistical significance was determined using the KS test.
D) As in (C) but sum total TOP2 CCs found in these regions.
916

Figure 6. Local DNA sequence and methylation status direct the formation of Top2 CCs
A) The percentage of strong sites that are SSBs and DSBs in etoposide-treated *S. cerevisiae* and
human cells.
B) Average nucleotide composition over a 30 bp window centred on the DSB or (inferred) SSB dyad
axis, in etoposide-treated *S. cerevisiae* and human cells. Values reported are for the top strand only.
C) Heatmaps describing pairwise similarity of nucleotide composition patterns shown in (B). Rows 1-
4 are the absolute differences between: yeast and human SSB patterns, yeast and human DSB
patterns, yeast SSB and DSB patterns, and human SSB and DSB patterns, respectively.
D) The average number of TOP2 CCs at the +1 position relative to methylated and unmethylated
cytosines in human RPE-1 cells, ±VP16. Statistical significance was determined using KS test.
927

Supplementary Figure Legends

Figure S1. CC-seq maps covalent Spo11-linked DNA breaks in *S. cerevisiae* meiosis with nucleotide accuracy (related to Figure 1)

A) Correlation of hotspot signals from CC-seq and oligo-seq.

B) Correlation of 500 bp binned Spo11 maps from two representative replicates.

C) Correlation of Spo11 cleavages on Watson and Crick strand when offset by 1 bp (boxed), relative to other offsets from -4 to +4 bp.

D) Spo11 breaks mapped by CC-seq (raw=black, or smoothed=red) anticorrelate with nucleosome occupancy measured by MNase-seq (blue).

E) Aggregation of Spo11 activity in a 10 Kbp window centred on orientated TSSs in *S. cerevisiae*. Three example TSSs are shown orientated in the 5'-3' direction (top). A heatmap of all TSSs in the *S. cerevisiae* genome, with 25 rows stratified by gene expression level in SK1 cells (middle). The colour scale indicates average Spo11 break density. Motifs are also stratified into 4 quartiles of gene expression in meiotic SK1 cells, and the average distribution of Spo11 activity in each quartile plotted (bottom).

Figure S2. CC-seq maps covalent Top2-linked DNA breaks in *S. cerevisiae* cycling cells with nucleotide accuracy (related to Figure 2).

A-C) Correlation of 1 Kbp binned Top2 CC maps from *pdr1Δ* (A), *pdr1Δsae2Δ* (B) and *pdr1Δmre11Δ* (C) cells treated with VP16.

D) Pairwise Pearson correlation values for 100 bp binned Top2 CC maps from all assayed conditions. Each condition is a pool of two biological replicates.

E) Nucleotide-resolution *S. cerevisiae* Top2 CC map of chromosome 3 for all assayed conditions. Each condition is a pool of two biological replicates.

F-H) The normalised number of Top2 CCs retained in the *pdr1Δ* (F), *pdr1Δsae2Δ* (G) and *pdr1Δmre11Δ* (H) cells treated ±VP16, after filtering to include only sites offset by the given number of base pairs. All data were normalised over a -100 to +100 bp window.

I-K) Pearson correlation (r) of Top2 CC-seq signal on Watson and Crick strands, offset by the indicated distance in *pdr1Δ* (I), *pdr1Δsae2Δ* (J) and *pdr1Δmre11Δ* (K) cells treated ±VP16.

Figure S3. Top2-linked DNA breaks in *S. cerevisiae* are not correlated with gene expression, anticorrelate with nucleosomes, and have biased nucleotide skews indicative of bent DNA (related to Figure 2)

A) Top2 CC mapped by CC-seq (raw=black, or smoothed=red) anticorrelate with nucleosome occupancy measured by MNase-seq (blue).

B) Aggregation of Top2 CCs in a 10 Kbp window centred on orientated transcription start sites (TSS) in *pdr1Δ S. cerevisiae*. Four example TSSs are shown orientated in the 5'-3' direction (top). Heatmap of all TSSs in the *S. cerevisiae* genome, with 25 rows stratified by gene expression level in vegetative growth (middle). Colour scale indicates average Top2 CC density. Motifs are also stratified into four quartiles of gene expression, and the average distribution of Top2 CCs in each quartile plotted for untreated and +VP16 conditions (bottom).

C) The percentage of strong sites that are SSBs and DSBs in etoposide-treated *pdr1Δ* and *pdr1Δmre11Δ S. cerevisiae*. Sites were thresholded at 1 HpM prior to sorting into DSB or SSB classes based on presence or absence of a 3 bp offset cognate (Obs.). As a control, the amplitudes of Top2 CCs (HpM) were randomised amongst the positions in the nucleotide resolution datasets, prior to thresholding and offset analysis as described above (Rand.)

D) Average nucleotide composition over a 200 bp window centred on the DSB dyad axis, in etoposide-treated *S. cerevisiae*. The position of the flanking regions (F.R.) identified in Gyrase

mapping experiments, and the flanking regions observed here for Top2 are indicated in blue and
black, respectively.
D) Average nucleotide composition over a 120 bp window centred on the DSB dyad axis, in
etoposide-treated *S. cerevisiae*. The signal pattern is separated into A+T and G+C (top left and right),
and into individual A, T, C and G plots (middle left, bottom left, middle right, and bottom right,
respectively). The position of the flanking regions (F.R.) observed here for Top2 are indicated in
black, and their ~10.5 bp periodicity is highlighted with black and red crosses.
986
Figure S4. CC-seq maps of TOP2-linked DNA breaks in Human cells are enriched by
etoposide, and show high reproducibility and nucleotide accuracy (related to Figure 3)
A) Broad-scale *H. sapiens* TOP2 CC-seq maps in individual biological replicates of RPE-1 cells
±VP16. Raw data were binned at 100 Kbp prior to plotting. Each plot is offset on the y-axis by +0.3
HpB.
B) Replicate-to-replicate Pearson correlation values (r) for 10 Kbp binned TOP2 CC-seq maps of
RPE-1 cells ±VP16.
C) Scatter plot of -VP16 and +VP16 TOP2 CC-seq maps binned at 10 Kbp resolution. Data were first
scaled according to the estimated noise fraction (**Methods**), and are presented in a hexagonal-
binned format, where the density of overplotting is indicated by the colour scale.
D) Violin plots of TOP2 CC-seq maps ±VP16 binned at 100 Kbp resolution. The inner black bar,
black dot, and dotted horizontal line indicate the interquartile range, median, and expected mean
Top2 CC density based on random distribution.
E) The normalised number of TOP2 CCs retained in the CC-seq maps in RPE-1 cells ±VP16 after
filtering to include only sites offset by the given number of base pairs. Data were normalised over a -
100 to +100 bp window.
1003
Figure S5. CC-seq signal is TOP2-dependent
A) DNA content histograms of wild type (WT) and *TOP2B*$^{-/-}$ RPE-1 cells under asynchronous (10%
FCS) and serum-deprived (0% FCS) conditions, as measured by FACS following propidium iodide
(PI) staining. G1, S and G2 populations are clearly present under asynchronous growing conditions.
A strong G1 arrest is observed in serum deprived conditions. Percentages of cells in each of the
indicated regions (red dotted brackets) are given.
B) Western blots demonstrating the absence of TOP2β protein in serum deprived and asynchronous
TOP2B$^{-/-}$ RPE-1 cells (left), and the absence of TOP2α in serum deprived wild type and *TOP2B*$^{-/-}$
RPE-1 cells (right). Ponceau S total protein loading is presented (Pon. S) for the left and right panels,
and additionally a Ku80 loading control is included for the right panel.
C) Immunofluorescence experiment demonstrating induction of γ-H2AX foci (green) in asynchronous
(Async.) and serum-deprived (Ser. Dep.) wild type (WT) and *TOP2B*$^{-/-}$ RPE-1 cells, all co-stained with
DAPI (blue). Galleries of nine cells per condition were chosen randomly using Olympus ScanR
Analysis software.
D) Quantification of (C). Numbers of γ-H2AX foci per cell were counted automatically using Olympus
ScanR Analysis software. The mean ±SEM is reported for n= 3 biological replicate experiments.
E) Broad-scale *H. sapiens* TOP2 CC-seq maps in asynchronous and serum-deprived wild type (WT)
and *TOP2B*$^{-/-}$ RPE-1 cells -VP16 (orange) and +VP16 (green). Raw hits on Watson and Crick strands
were summed and smoothed according to local signal density (Fsize=501).
1023
Figure S6. TOP2 CC-seq signal enrichment around CTCF and TSS sites, compared with END-
seq DSB signal at TSSs
A) Fine-scale TOP2 CC-seq maps of *H. sapiens* CTCF-proximal loci in RPE-1 cells +VP16. Red and
blue traces indicate TOP2-linked 5' DNA termini on the Watson and Crick strands, respectively. Pale
shaded areas are the same data smoothed with a sliding 11 bp Hanning window. Red and Blue

rectangles indicate the positions of CTCF motifs on the Watson and Crick strands respectively. The
grey line indicates Hanning-smoothed sum of Watson and Crick TOP2 CCs.
B) Fine-scale mapping of TOP2 CCs surrounding three complex CTCF loci, processed as in (A).
C) Aggregation of TOP2 CCs in a 1 Kbp window centred on the subset of orientated CTCF motifs
that can be assigned to a chromatin loop anchor in human RPE-1 cells (Darrow et al., 2016). Motifs
are stratified into 4 quartiles of loop anchor interaction strength, and the average TOP2 CC
distribution in each quartile plotted.
D) Fine-scale aggregation of TOP2 CCs (red) in a 800 bp window centred on TSSs in human RPE-1
cells, showing anticorrelation with aggregated MNase-seq signal (blue).
E) Aggregation of END-seq mapped DSBs in a 10 Kbp window centred on orientated TSSs in human
MCF7 cells. Four example TSSs are shown orientated in the 5'-3' direction (top). A heatmap of all
TSSs in the human genome, with 25 rows stratified by flanking gene expression level in MCF7 cells
(middle). The colour scale indicates average END-seq DSB density. TSSs are also stratified into 4
quartiles based on strength of flanking gene expression, and the average END-seq DSB distribution
in each quartile plotted (bottom).
1044
Figure S7. TSS-proximal TOP2-linked DNA breaks in humans are correlated with gene length,
independently from gene expression level (related to Figure 5); TSS-proximal Top2-linked DNA
breaks in yeast are not strongly correlated with gene length, (related to Figure 2)
A) Aggregation of TOP2 CCs in a 10 Kbp window centred on orientated TSSs in human RPE-1 cells.
A heatmap of all TSSs in the human genome, with 25 rows stratified by gene length (top). The colour
scale indicates average TOP2 CC density. Motifs are also stratified into 4 quartiles of gene length,
and the average TOP2 CC distribution in each quartile plotted (bottom).
B) Aggregation of Top2 CCs in a 10 Kbp window centred on orientated transcription start sites (TSS)
in *pdr1Δ S. cerevisiae*. Heatmap of all TSSs in the *S. cerevisiae* genome, with 25 rows stratified by
gene length level (top). Colour scale indicates average Top2 CC density. Motifs are also stratified
into four quartiles of gene length, and the average distribution of Top2 CCs in each quartile plotted
(bottom).
C) Scatter plot of human gene length and RPE-1 gene expression, showing no correlation.
D) Scatter plot of *S. cerevisiae* gene length and gene expression during vegetative growth, showing
little correlation.
1060
1061
1062

Supplementary Tables

Table S1. DNA libraries used for mapping Spo11 activity. The number of reads remaining after each stage of the data pipeline are indicated.

Species	Strain/ Cell line	Condition	NGS Platform	Library code	Total Read Pairs	Mapped reads	Blacklist-filtered reads	% of Total
S. cerevisiae	*sae2Δ*	Meiosis	MiSeq	1	3512089	3376630	3177210	90.5
				2A	10101423	8796998	8564017	84.8
				3	4957477	4745192	4502308	90.8
				5	4012972	3705100	3559541	88.7
				7BC	4065600	3779390	3569290	87.8

1073 **Table S2. DNA libraries used for mapping Top2 activity.** The number of reads
1074 remaining after each stage of the data pipeline are indicated.
1075

Species	Strain/ Cell line	Condition	NGS Platform	Library code	Total Read Pairs	Mapped reads	De-duplicated reads	Blacklist-filtered reads	% of Total	Pooled Reads (Millions)
S. cerevisiae	*pdr1Δ*	-VP16	NextSeq 500	RA1	3741251	3542955	3179049	2746294	73.4	3.50
			MiSeq	RA8	1118483	879107	842290	751850	67.2	
S. cerevisiae	*pdr1Δ*	+VP16	NextSeq 500	RA2	3096257	3053356	2679303	2234941	72.2	3.93
			MiSeq	RA9	2262958	2071231	1958341	1690075	74.7	
S. cerevisiae	*pdr1Δ sae2Δ*	-VP16	NextSeq 500	RA4	3634284	3552818	3215523	2745668	75.5	3.79
			MiSeq	RA10	1420662	1214067	1181399	1046827	73.7	
S. cerevisiae	*pdr1Δ sae2Δ*	+VP16	NextSeq 500	RA5	2529368	2476714	1386551	1097282	43.4	3.54
			MiSeq	RA11	3226626	3063388	2824882	2444486	75.8	
S. cerevisiae	*pdr1Δ mre11Δ*	-VP16	NextSeq 500	RA6	5345587	5205411	4288662	3646842	68.2	4.76
			MiSeq	RA12	1519020	1326435	1280137	1109310	73.0	
S. cerevisiae	*pdr1Δ mre11Δ*	+VP16	NextSeq 500	RA7	3382546	3357168	2921333	2526342	74.7	5.54
			MiSeq	RA13	4133347	4032996	3540366	3010542	72.8	
Human	RPE-1 WT	-VP16	NextSeq 500	WG11	125203096	109503841	68845619	68639594	54.8	185.36
				WG15	125328792	110432447	46588296	46449378	37.1	
				WG23	89154181	77728045	70484665	70269334	78.8	
Human	RPE-1 WT	+VP16	NextSeq 500	WG12	110910655	95546007	55848200	55681939	50.2	276.97
				WG16	102804692	91872706	54312084	54164650	52.7	
				WG24	130816235	114086421	101162102	100857557	77.1	
				WG34	91829516	79751337	66458538	66270349	72.2	
Human	RPE-1 WT	-VP16 G1	MiSeq	WG13	2601642	2292863	2007177	1991088	76.5	4.27
				WG21	2610523	2298289	2295625	2277124	87.2	
Human	RPE-1 WT	+VP16 G1	MiSeq	WG14	3075620	2771819	2491674	2475350	80.5	5.23
				WG22	3139164	2779809	2772349	2751281	87.6	
Human	RPE-1 WT	-VP16 Async	MiSeq	WG15b*	3207507	2866283	2782876	2760652	86.1	5.35
				WG23	2959289	2616954	2608670	2586523	87.4	
Human	RPE-1 WT	+VP16 Async	MiSeq	WG16b*	3018163	2721770	2673273	2656580	88.0	6.30
				WG24	4165048	3682708	3668602	3639324	87.4	
Human	RPE-1 TOP2B-	-VP16 G1	MiSeq	WG17	2978044	2624262	2558332	2536752	85.2	4.69
				WG25	2460949	2181002	2173777	2155101	87.6	
Human	RPE-1 TOP2B-	+VP16 G1	MiSeq	WG18	2927745	2588870	2557532	2537085	86.7	5.10
				WG26	2922624	2596547	2587152	2565454	87.8	
Human	RPE-1 TOP2B-	-VP16 Async	MiSeq	WG19	3713068	3278164	3188049	3162895	85.2	7.04
				WG27	4413242	3943962	3910035	3877611	87.9	
Human	RPE-1 TOP2B-	+VP16 Async	MiSeq	WG20	3426100	3038601	2978047	2956076	86.3	8.13
				WG28	5878342	5251089	5211027	5169010	87.9	

1076
1077
1078

1079 **Table S3. *S. cerevisiae* strains used in this study.** Strains MJ315 and M319 were
1080 used for Spo11 mapping; strains MJ429, MJ475 and MJ551 were used for Top2
1081 mapping.
1082

Strain	Number	Genotype
sae2Δ	MJ315	SK1: *ho::LYS2/', lys2/', ura3/', arg4-nsp/', leu2::hisG/', his4X::LEU2/', nuc1::LEU2/', sae2Δ::KanMX6/'*
sae2Δ spo11-Y135F	MJ319	SK1: *ho::LYS2/', lys2/', ura3/', arg4-nsp/', leu2::hisG/', his4X::LEU2/', nuc1::LEU2/', spo11(Y135F)-HA3His6::KanMX4/', sae2Δ::KanMX6/'*
pdr1Δ	MJ429	BY4741: *ura3Δ0, leu2Δ0, his3Δ1, met15Δ0, pdr1Δ::PDR1-DBD-CYC8::LEU2*
pdr1Δ sae2Δ	MJ475	BY4741: *ura3Δ0, leu2Δ0, his3Δ1, met15Δ0, pdr1Δ::PDR1-DBD-CYC8::LEU2, sae2Δ::KanMX6*
pdr1Δ mre11Δ	MJ551	BY4741: *ura3Δ0, leu2Δ0, his3Δ1, met15Δ0, pdr1Δ::PDR1-DBD-CYC8::LEU2, mre11Δ::KanMX4*

1083
1084
1085
1086 **Table S4. Human cell lines used in this study.**
1087

Parent cell line	Clone number	Genotype
hTERT RPE-1	WT	WT
hTERT RPE-1	T2B/1	*TOP2B*

1088
1089
1090
1091 **Table S5. The annealed structures of the custom P5 and P7 adapters.** "T_i" indicates
1092 an inverted dT, linked via a 3'-3' phosphodiester. This inhibits unwanted ligation at this
1093 end of the adapter. Also note that the 5'-terminal moieties of each adapter (A for P5, G
1094 for P7) are nucleosides (3'-OH), which further inhibits ligation at this end
1095

Adapter	Annealed Structure		
P5	blocked end	ACACTCTTTCCCTACACGACGCTCTTCCGATCT T_iTGTGAGAAAGGGATGTGCTGCGAGAAGGCTAGp	ligated end
P7	ligated end	pGATCGGAAGAGCACACGTCTGAACTCCAGTCACT$_i$ TCTAGCCTTCTCGTGTGCAGACTTGAGGTCAGTG	blocked end

1096
1097
1098 **Table S6. Publically available datasets used in this study.**
1099

Species/ Cell line	Dataset Type	Accession(s)	Reference
S. cerevisiae / SK1 (meiotic)	Gene expression microarray	GSM907178, GSM90719, GSM907180	(Dominissini et al., 2012)
S. cerevisiae / SK1 (vegetative)	Gene expression microarray	GSM907176, GSM907177	(Dominissini et al., 2012)
H. sapiens / RPE-1	Gene expression microarray	GSM1395252, GSM1395253, GSM1395254	(Ganem et al., 2014)
H. Sapiens / MCF7	Gene expression microarray	GSM1141244	(Nelson et al., 2013)
H. sapiens / RPE-1	Hi-C	GSE71831	(Darrow et al., 2016)
H. sapiens / RPE-1	bTMP-seq	GSM1062645, GSM1062646	(Naughton et al., 2013)
H. sapiens / MCF7	END-seq	GSM2635568	(Canela et al., 2017)
H. sapiens / RPE-1	CTCF ChIP-seq	GSM749673, GSM1022665	(ENCODE, 2012)
H. sapiens / RPE-1	H3K4Me3 ChIP-seq	GSM945271, GSM945271	(ENCODE, 2012)
H. sapiens / RPE-1	H3K27Ac ChIP-seq	GSM733771, GSM733718, GSM733755, GSM733691, GSM733656, GSM733674, GSM733646	(ENCODE, 2012)
H. sapiens / RPE-1	Methyl-RRBS	GSM683773, GSM683905	(ENCODE, 2012)
H. sapiens / K562	MNase-seq	GSM920557	(ENCODE, 2012)

References

Akimitsu, N., Adachi, N., Hirai, H., Hossain, M. S., Hamamoto, H., Kobayashi, M., Aratani, Y., Koyama, H., and Sekimizu, K. (2003). Enforced cytokinesis without complete nuclear division in embryonic cells depleting the activity of DNA topoisomerase IIalpha. Genes Cells *8*, 393-402.

Anand, R., Ranjha, L., Cannavo, E., and Cejka, P. (2016). Phosphorylated CtIP Functions as a Co-factor of the MRE11-RAD50-NBS1 Endonuclease in DNA End Resection. Mol Cell *64*, 940-950.

Austin, C. A., Lee, K. C., Swan, R. L., Khazeem, M. M., Manville, C. M., Cridland, P., Treumann, A., Porter, A., Morris, N. J., and Cowell, I. G. (2018). TOP2B: The First Thirty Years. Int J Mol Sci *19*,

Baranello, L., Kouzine, F., Wojtowicz, D., Cui, K., Przytycka, T. M., Zhao, K., and Levens, D. (2014). DNA break mapping reveals topoisomerase II activity genome-wide. Int J Mol Sci *15*, 13111-13122.

Baxter, J., and Diffley, J. F. (2008). Topoisomerase II inactivation prevents the completion of DNA replication in budding yeast. Mol Cell *30*, 790-802.

Bergerat, A., de Massy, B., Gadelle, D., Varoutas, P. C., Nicolas, A., and Forterre, P. (1997). An atypical topoisomerase II from Archaea with implications for meiotic recombination. Nature *386*, 414-417.

Bermejo, R., Capra, T., Gonzalez-Huici, V., Fachinetti, D., Cocito, A., Natoli, G., Katou, Y., Mori, H., Kurokawa, K., Shirahige, K., and Foiani, M. (2009). Genome-organizing factors Top2 and Hmo1 prevent chromosome fragility at sites of S phase transcription. Cell *138*, 870-884.

Borgnetto, M. E., Zunino, F., Tinelli, S., Kas, E., and Capranico, G. (1996). Drug-specific sites of topoisomerase II DNA cleavage in Drosophila chromatin: heterogeneous localization and reversibility. Cancer Res *56*, 1855-1862.

Brill, S. J., DiNardo, S., Voelkel-Meiman, K., and Sternglanz, R. (1987). Need for DNA topoisomerase activity as a swivel for DNA replication for transcription of ribosomal RNA. Nature *326*, 414-416.

Bromberg, K. D., Burgin, A. B., and Osheroff, N. (2003). A two-drug model for etoposide action against human topoisomerase IIalpha. J Biol Chem *278*, 7406-7412.

Canela, A., Maman, Y., Jung, S., Wong, N., Callen, E., Day, A., Kieffer-Kwon, K.-R., Pekowska, A., Zhang, H., Rao, S. S. P., Huang, S.-C., Mckinnon, P. J., Aplan, P. D., Pommier, Y., Aiden, E. L., Casellas, R., and Nussenzweig, A. (2017). Genome Organization Drives Chromosome Fragility. Cell *170*, 507-521.e18.

Cannavo, E., Johnson, D., Andres, S. N., Kissling, V. M., Reinert, J. K., Garcia, V., Erie, D. A., Hess, D., Thomä, N. H., Enchev, R. I., Peter, M., Williams, R. S., Neale, M. J., and Cejka, P. (2018). Regulatory control of DNA end resection by Sae2 phosphorylation. Nat Commun *9*, 4016.

Cannavo, E., and Cejka, P. (2014). Sae2 promotes dsDNA endonuclease activity within Mre11-Rad50-Xrs2 to resect DNA breaks. Nature *514*, 122-125.

Capranico, G., Jaxel, C., Roberge, M., Kohn, K. W., and Pommier, Y. (1990). Nucleosome positioning as a critical determinant for the DNA cleavage sites of mammalian DNA topoisomerase II in reconstituted simian virus 40 chromatin. Nucleic Acids Res *18*, 4553-4559.

Chen, G. L., Yang, L., Rowe, T. C., Halligan, B. D., Tewey, K. M., and Liu, L. F. (1984). Nonintercalative antitumor drugs interfere with the breakage-reunion reaction of mammalian DNA topoisomerase II. J Biol Chem *259*, 13560-13566.

Coombs, D. H., and Pearson, G. D. (1978). Filter-binding assay for covalent DNA-protein complexes: adenovirus DNA-terminal protein complex. Proc Natl Acad Sci U S A *75*, 5291-5295.

Cortes Ledesma, F., El Khamisy, S. F., Zuma, M. C., Osborn, K., and Caldecott, K. W. (2009). A human 5'-tyrosyl DNA phosphodiesterase that repairs topoisomerase-mediated DNA damage. Nature *461*, 674-678.

Cowell, I. G., Sondka, Z., Smith, K., Lee, K. C., Manville, C. M., Sidorczuk-Lesthuruge, M., Rance, H. A., Padget, K., Jackson, G. H., Adachi, N., and Austin, C. A. (2012). Model for MLL translocations in therapy-related leukemia involving topoisomerase IIβ-mediated DNA strand breaks and gene proximity. Proc Natl Acad Sci U S A *109*, 8989-8994.

Crosetto, N., Mitra, A., Silva, M. J., Bienko, M., Dojer, N., Wang, Q., Karaca, E., Chiarle, R., Skrzypczak, M., Ginalski, K., Pasero, P., Rowicka, M., and Dikic, I. (2013). Nucleotide-resolution DNA double-strand break mapping by next-generation sequencing. Nat Methods *10*, 361-365.

Darrow, E. M., Huntley, M. H., Dudchenko, O., Stamenova, E. K., Durand, N. C., Sun, Z., Huang, S. C., Sanborn, A. L., Machol, I., Shamim, M., Seberg, A. P., Lander, E. S., Chadwick, B. P., and Aiden, E. L. (2016). Deletion of DXZ4 on the human inactive X chromosome alters higher-order genome architecture. Proc Natl Acad Sci U S A *113*, E4504-12.

Deweese, J. E., and Osheroff, N. (2009). The DNA cleavage reaction of topoisomerase II: wolf in sheep's clothing. Nucleic Acids Res *37*, 738-748.

DiNardo, S., Voelkel, K., and Sternglanz, R. (1984). DNA topoisomerase II mutant of Saccharomyces cerevisiae: topoisomerase II is required for segregation of daughter molecules at the termination of DNA replication. Proc Natl Acad Sci U S A *81*, 2616-2620.

Dominissini, D., Moshitch-Moshkovitz, S., Schwartz, S., Salmon-Divon, M., Ungar, L., Osenberg, S., Cesarkas, K., Jacob-Hirsch, J., Amariglio, N., Kupiec, M., Sorek, R., and Rechavi, G. (2012). Topology of the human and mouse m6A RNA methylomes revealed by m6A-seq. Nature *485*, 201-206.

Durand, N. C., Shamim, M. S., Machol, I., Rao, S. S., Huntley, M. H., Lander, E. S., and Aiden, E. L. (2016). Juicer Provides a One-Click System for Analyzing Loop-Resolution Hi-C Experiments. Cell Syst *3*, 95-98.

ENCODE, P. C. (2012). An integrated encyclopedia of DNA elements in the human genome. Nature *489*, 57-74.

Ganem, N. J., Cornils, H., Chiu, S. Y., O'Rourke, K. P., Arnaud, J., Yimlamai, D., Théry, M., Camargo, F. D., and Pellman, D. (2014). Cytokinesis failure triggers hippo tumor suppressor pathway activation. Cell *158*, 833-848.

Garcia, V., Phelps, S. E. L., Gray, S., and Neale, M. J. (2011). Bidirectional resection of DNA double-strand breaks by Mre11 and Exo1. Nature *479*, 241-244.

Gómez-Herreros, F., Schuurs-Hoeijmakers, J. H., McCormack, M., Greally, M. T., Rulten, S., Romero-Granados, R., Counihan, T. J., Chaila, E., Conroy, J., Ennis, S., Delanty, N., Cortés-Ledesma, F., de Brouwer, A. P., Cavalleri, G. L., El-Khamisy, S. F., de Vries, B. B., and Caldecott, K. W. (2014). TDP2 protects transcription from abortive topoisomerase activity and is required for normal neural function. Nat Genet *46*, 516-521.

Gothe, H. J., Bouwman, B. A. M., Gusmao, E. G., Piccinno, R., Sayols, S., Drechsel, O., Petrosino, G., Minneker, V., Josipovic, N., Mizi, A., Nielsen, C. F., Wagner, E.-M., Takeda, S., Sasanuma, H., Hudson, D. F., Kindler, T., Baranello, L., Papantonis, A., Crosetto, N., and Roukos, V. (2018). Spatial chromosome folding and active transcription drive DNA fragility and formation of oncogenic MLL translocations. bioRxiv *http://dx.doi.org/10.1101/485763*,

Grant, C. E., Bailey, T. L., and Noble, W. S. (2011). FIMO: scanning for occurrences of a given motif. Bioinformatics *27*, 1017-1018.

Grue, P., Grässer, A., Sehested, M., Jensen, P. B., Uhse, A., Straub, T., Ness, W., and Boege, F. (1998). Essential mitotic functions of DNA topoisomerase IIalpha are not adopted by topoisomerase IIbeta in human H69 cells. J Biol Chem *273*, 33660-33666.

Grzybowski, A. T., Chen, Z., and Ruthenburg, A. J. (2015). Calibrating ChIP-Seq with Nucleosomal Internal Standards to Measure Histone Modification Density Genome Wide. Mol Cell *58*, 886-899.

Haffner, M. C., Aryee, M. J., Toubaji, A., Esopi, D. M., Albadine, R., Gurel, B., Isaacs, W. B., Bova, G. S., Liu, W., Xu, J., Meeker, A. K., Netto, G., De Marzo, A. M., Nelson, W. G., and Yegnasubramanian, S. (2010). Androgen-induced TOP2B-mediated double-strand breaks and prostate cancer gene rearrangements. Nat Genet *42*, 668-675.

Hamilton, N. K., and Maizels, N. (2010). MRE11 function in response to topoisomerase poisons is independent of its function in double-strand break repair in Saccharomyces cerevisiae. PLoS One *5*, e15387.

Hartsuiker, E., Neale, M. J., and Carr, A. M. (2009). Distinct requirements for the Rad32(Mre11) nuclease and Ctp1(CtIP) in the removal of covalently bound topoisomerase I and II from DNA. Mol Cell *33*, 117-123.

Heck, M. M., Hittelman, W. N., and Earnshaw, W. C. (1988). Differential expression of DNA topoisomerases I and II during the eukaryotic cell cycle. Proc Natl Acad Sci U S A *85*, 1086-1090.

Hoa, N. N., Shimizu, T., Zhou, Z. W., Wang, Z. Q., Deshpande, R. A., Paull, T. T., Akter, S., Tsuda, M., Furuta, R., Tsusui, K., Takeda, S., and Sasanuma, H. (2016). Mre11 Is Essential for the Removal of Lethal Topoisomerase 2 Covalent Cleavage Complexes. Mol Cell *64*, 580-592.

Hoffman, M. M., Ernst, J., Wilder, S. P., Kundaje, A., Harris, R. S., Libbrecht, M., Giardine, B., Ellenbogen, P. M., Bilmes, J. A., Birney, E., Hardison, R. C., Dunham, I., Kellis, M., and Noble, W. S. (2013). Integrative annotation of chromatin elements from ENCODE data. Nucleic Acids Res *41*, 827-841.

Holm, C., Goto, T., Wang, J. C., and Botstein, D. (1985). DNA topoisomerase II is required at the time of mitosis in yeast. Cell *41*, 553-563.

Hornyak, P., Askwith, T., Walker, S., Komulainen, E., Paradowski, M., Pennicott, L. E., Bartlett, E. J., Brissett, N. C., Raoof, A., Watson, M., Jordan, A. M., Ogilvie, D. J., Ward, S. E., Atack, J. R., Pearl, L. H., Caldecott, K. W., and Oliver, A. W. (2016). Mode of action of DNA-competitive small molecule inhibitors of tyrosyl DNA phosphodiesterase 2. Biochem J *473*, 1869-1879.

Hsiang, Y. H., Lihou, M. G., and Liu, L. F. (1989). Arrest of replication forks by drug-stabilized topoisomerase I-DNA cleavable complexes as a mechanism of cell killing by camptothecin. Cancer Res *49*, 5077-5082.

Hsu, P. D., Scott, D. A., Weinstein, J. A., Ran, F. A., Konermann, S., Agarwala, V., Li, Y., Fine, E. J., Wu, X., Shalem, O., Cradick, T. J., Marraffini, L. A., Bao, G., and Zhang, F. (2013). DNA targeting specificity of RNA-guided Cas9 nucleases. Nat Biotechnol *31*, 827-832.

Hu, B., Petela, N., Kurze, A., Chan, K.-L., Chapard, C., and Nasmyth, K. (2015). Biological chromodynamics: a general method for measuring protein occupancy across the genome by calibrating ChIP-seq. Nucleic Acids Res

Huang, W. C., Lee, C. Y., and Hsieh, T. S. (2017). Single-molecule Förster resonance energy transfer (FRET) analysis discloses the dynamics of the DNA-topoisomerase II (Top2) interaction in the presence of TOP2-targeting agents. J Biol Chem *292*, 12589-12598.

Johnson, D., Allison, R. M., Cannavo, E., Cejka, P., and Neale, M. (2019). Removal of Spo11 from meiotic DNA breaks in vitro but not in vivo by Tyrosyl DNA Phosphodiesterase 2. bioRxiv *http://dx.doi.org/10.1101/527333*,

Käs, E., and Laemmli, U. K. (1992). In vivo topoisomerase II cleavage of the Drosophila histone and satellite III repeats: DNA sequence and structural characteristics. EMBO J *11*, 705-716.

Keeney, S., Giroux, C. N., and Kleckner, N. (1997). Meiosis-specific DNA double-strand breaks are catalyzed by Spo11, a member of a widely conserved protein family. Cell *88*, 375-384.

Keeney, S., and Kleckner, N. (1995). Covalent protein-DNA complexes at the 5' strand termini of meiosis-specific double-strand breaks in yeast. Proc Natl Acad Sci U S A *92*, 11274-11278.

Khan, Q. A., Kohlhagen, G., Marshall, R., Austin, C. A., Kalena, G. P., Kroth, H., Sayer, J. M., Jerina, D. M., and Pommier, Y. (2003). Position-specific trapping of topoisomerase II by benzo[a]pyrene diol epoxide adducts: implications for interactions with intercalating anticancer agents. Proc Natl Acad Sci U S A *100*, 12498-12503.

Kingma, P. S., Corbett, A. H., Burcham, P. C., Marnett, L. J., and Osheroff, N. (1995). Abasic sites stimulate double-stranded DNA cleavage mediated by topoisomerase II. DNA lesions as endogenous topoisomerase II poisons. J Biol Chem *270*, 21441-21444.

Lee, M. P., Sander, M., and Hsieh, T. (1989). Nuclease protection by Drosophila DNA topoisomerase II. Enzyme/DNA contacts at the strong topoisomerase II cleavage sites. J Biol Chem *264*, 21779-21787.

Liang, K., and Keleş, S. (2012). Normalization of ChIP-seq data with control. BMC Bioinformatics *13*, 199.

Lieberman-Aiden, E., van Berkum, N. L., Williams, L., Imakaev, M., Ragoczy, T., Telling, A., Amit, I., Lajoie, B. R., Sabo, P. J., Dorschner, M. O., Sandstrom, R., Bernstein, B., Bender, M. A., Groudine, M., Gnirke, A., Stamatoyannopoulos, J., Mirny, L. A.,

Lander, E. S., and Dekker, J. (2009). Comprehensive mapping of long-range interactions reveals folding principles of the human genome. Science *326*, 289-293.

Lin, C., Yang, L., Tanasa, B., Hutt, K., Ju, B. G., Ohgi, K., Zhang, J., Rose, D. W., Fu, X. D., Glass, C. K., and Rosenfeld, M. G. (2009). Nuclear receptor-induced chromosomal proximity and DNA breaks underlie specific translocations in cancer. Cell *139*, 1069-1083.

Liu, J., Wu, T. C., and Lichten, M. (1995). The location and structure of double-strand DNA breaks induced during yeast meiosis: evidence for a covalently linked DNA-protein intermediate. EMBO J *14*, 4599-4608.

Liu, L. F., Rowe, T. C., Yang, L., Tewey, K. M., and Chen, G. L. (1983). Cleavage of DNA by mammalian DNA topoisomerase II. J Biol Chem *258*, 15365-15370.

Liu, Q., and Wang, J. C. (1999). Similarity in the catalysis of DNA breakage and rejoining by type IA and IIA DNA topoisomerases. Proc Natl Acad Sci U S A *96*, 881-886.

Long, B. H., Musial, S. T., and Brattain, M. G. (1986). DNA breakage in human lung carcinoma cells and nuclei that are naturally sensitive or resistant to etoposide and teniposide. Cancer Res *46*, 3809-3816.

Lotito, L., Russo, A., Chillemi, G., Bueno, S., Cavalieri, D., and Capranico, G. (2008). Global transcription regulation by DNA topoisomerase I in exponentially growing Saccharomyces cerevisiae cells: activation of telomere-proximal genes by TOP1 deletion. J Mol Biol *377*, 311-322.

Mathelier, A., Zhao, X., Zhang, A. W., Parcy, F., Worsley-Hunt, R., Arenillas, D. J., Buchman, S., Chen, C. Y., Chou, A., Ienasescu, H., Lim, J., Shyr, C., Tan, G., Zhou, M., Lenhard, B., Sandelin, A., and Wasserman, W. W. (2014). JASPAR 2014: an extensively expanded and updated open-access database of transcription factor binding profiles. Nucleic Acids Res *42*, D142-7.

Minocha, A., and Long, B. H. (1984). Inhibition of the DNA catenation activity of type II topoisomerase by VP16-213 and VM26. Biochem Biophys Res Commun *122*, 165-170.

Nakamura, K., Kogame, T., Oshiumi, H., Shinohara, A., Sumitomo, Y., Agama, K., Pommier, Y., Tsutsui, K. M., Tsutsui, K., Hartsuiker, E., Ogi, T., Takeda, S., and Taniguchi, Y. (2010). Collaborative action of Brca1 and CtIP in elimination of covalent modifications from double-strand breaks to facilitate subsequent break repair. PLoS Genet *6*, e1000828.

Naughton, C., Avlonitis, N., Corless, S., Prendergast, J. G., Mati, I. K., Eijk, P. P., Cockroft, S. L., Bradley, M., Ylstra, B., and Gilbert, N. (2013). Transcription forms and remodels supercoiling domains unfolding large-scale chromatin structures. Nat Struct Mol Biol *20*, 387-395.

Neale, M. J., Pan, J., and Keeney, S. (2005). Endonucleolytic processing of covalent protein-linked DNA double-strand breaks. Nature *436*, 1053-1057.

Negri, C., Chiesa, R., Cerino, A., Bestagno, M., Sala, C., Zini, N., Maraldi, N. M., and Astaldi Ricotti, G. C. (1992). Monoclonal antibodies to human DNA topoisomerase I and the two isoforms of DNA topoisomerase II: 170- and 180-kDa isozymes. Exp Cell Res *200*, 452-459.

Nelson, E. R., Wardell, S. E., Jasper, J. S., Park, S., Suchindran, S., Howe, M. K., Carver, N. J., Pillai, R. V., Sullivan, P. M., Sondhi, V., Umetani, M., Geradts, J., and McDonnell, D. P. (2013). 27-Hydroxycholesterol links hypercholesterolemia and breast cancer pathophysiology. Science *342*, 1094-1098.

Nitiss, J. L. (2009). Targeting DNA topoisomerase II in cancer chemotherapy. Nat Rev Cancer *9*, 338-350.

Pan, J., Sasaki, M., Kniewel, R., Murakami, H., Blitzblau, H. G., Tischfield, S. E., Zhu, X., Neale, M. J., Jasin, M., Socci, N. D., Hochwagen, A., and Keeney, S. (2011). A Hierarchical Combination of Factors Shapes the Genome-wide Topography of Yeast Meiotic Recombination Initiation. Cell *144*, 719-731.

Pommier, Y., Capranico, G., Orr, A., and Kohn, K. W. (1991). Local base sequence preferences for DNA cleavage by mammalian topoisomerase II in the presence of amsacrine or teniposide. Nucleic Acids Res *19*, 5973-5980.

Pommier, Y., Schwartz, R. E., Kohn, K. W., and Zwelling, L. A. (1984). Formation and rejoining of deoxyribonucleic acid double-strand breaks induced in isolated cell nuclei by antineoplastic intercalating agents. Biochemistry *23*, 3194-3201.

Pommier, Y., Sun, Y., Huang, S. N., and Nitiss, J. L. (2016). Roles of eukaryotic topoisomerases in transcription, replication and genomic stability. Nat Rev Mol Cell Biol *17*, 703-721.

Reece, R. J., and Maxwell, A. (1991). DNA gyrase: structure and function. Crit Rev Biochem Mol Biol *26*, 335-375.

Rowley, J. D., and Olney, H. J. (2002). International workshop on the relationship of prior therapy to balanced chromosome aberrations in therapy-related myelodysplastic syndromes and acute leukemia: overview report. Genes Chromosomes Cancer *33*, 331-345.

Sabourin, M., and Osheroff, N. (2000). Sensitivity of human type II topoisomerases to DNA damage: stimulation of enzyme-mediated DNA cleavage by abasic, oxidized and alkylated lesions. Nucleic Acids Res *28*, 1947-1954.

Sasanuma, H., Tsuda, M., Morimoto, S., Saha, L. K., Rahman, M. M., Kiyooka, Y., Fujiike, H., Cherniack, A. D., Itou, J., Callen Moreu, E., Toi, M., Nakada, S., Tanaka, H., Tsutsui, K., Yamada, S., Nussenzweig, A., and Takeda, S. (2018). BRCA1 ensures genome integrity by eliminating estrogen-induced pathological topoisomerase II-DNA complexes. Proc Natl Acad Sci U S A *115*, E10642-E10651.

Schalbetter, S. A., Fudenberg, G., Baxter, J., Pollard, K. S., and Neale, M. J. (2018). Principles of Meiotic Chromosome Assembly. bioRXiv *http://dx.doi.org/10.1101/442038*,

Stepanov, A., Nitiss, K. C., Neale, G., and Nitiss, J. L. (2008). Enhancing drug accumulation in Saccharomyces cerevisiae by repression of pleiotropic drug resistance genes with chimeric transcription repressors. Mol Pharmacol *74*, 423-431.

Stingele, J., Habermann, B., and Jentsch, S. (2015). DNA-protein crosslink repair: proteases as DNA repair enzymes. Trends Biochem Sci *40*, 67-71.

Strumberg, D., Nitiss, J. L., Dong, J., Kohn, K. W., and Pommier, Y. (1999). Molecular analysis of yeast and human type II topoisomerases. Enzyme-DNA and drug interactions. J Biol Chem *274*, 28246-28255.

Sun, X., Huang, L., Markowitz, T. E., Blitzblau, H. G., Chen, D., Klein, F., and Hochwagen, A. (2015). Transcription dynamically patterns the meiotic chromosome-axis interface. Elife *4*,

Sutormin, D., Rubanova, N., Logacheva, M., Ghilarov, D., and Severinov, K. (2018). Single-nucleotide-resolution mapping of DNA gyrase cleavage sites across the Escherichia coli genome. Nucleic Acids Res

Thomsen, B., Bendixen, C., Lund, K., Andersen, A. H., Sørensen, B. S., and Westergaard, O. (1990). Characterization of the interaction between topoisomerase II and DNA by transcriptional footprinting. J Mol Biol *215*, 237-244.

Tiwari, V. K., Burger, L., Nikoletopoulou, V., Deogracias, R., Thakurela, S., Wirbelauer, C., Kaut, J., Terranova, R., Hoerner, L., Mielke, C., Boege, F., Murr, R., Peters, A. H., Barde, Y. A., and Schübeler, D. (2012). Target genes of Topoisomerase IIβ regulate neuronal survival and are defined by their chromatin state. Proc Natl Acad Sci U S A *109*, E934-43.

Udvardy, A., and Schedl, P. (1991). Chromatin structure, not DNA sequence specificity, is the primary determinant of topoisomerase II sites of action in vivo. Mol Cell Biol *11*, 4973-4984.

Uemura, T., and Yanagida, M. (1984). Isolation of type I and II DNA topoisomerase mutants from fission yeast: single and double mutants show different phenotypes in cell growth and chromatin organization. EMBO J *3*, 1737-1744.

Uusküla-Reimand, L., Hou, H., Samavarchi-Tehrani, P., Rudan, M. V., Liang, M., Medina-Rivera, A., Mohammed, H., Schmidt, D., Schwalie, P., Young, E. J., Reimand, J., Hadjur, S., Gingras, A. C., and Wilson, M. D. (2016). Topoisomerase II beta interacts with cohesin and CTCF at topological domain borders. Genome Biol *17*, 182.

Vélez-Cruz, R., Riggins, J. N., Daniels, J. S., Cai, H., Guengerich, F. P., Marnett, L. J., and Osheroff, N. (2005). Exocyclic DNA lesions stimulate DNA cleavage mediated by human topoisomerase II alpha in vitro and in cultured cells. Biochemistry *44*, 3972-3981.

Wang, J. C. (2009). Untangling the double helix Cold Spring Harbor Laboratory Pr).

Wang, Y., Knudsen, B. R., Bjergbaek, L., Westergaard, O., and Andersen, A. H. (1999). Stimulated activity of human topoisomerases IIalpha and IIbeta on RNA-containing substrates. J Biol Chem *274*, 22839-22846.

Wang, Y., Thyssen, A., Westergaard, O., and Andersen, A. H. (2000). Position-specific effect of ribonucleotides on the cleavage activity of human topoisomerase II. Nucleic Acids Res *28*, 4815-4821.

Woessner, R. D., Mattern, M. R., Mirabelli, C. K., Johnson, R. K., and Drake, F. H. (1991). Proliferation- and cell cycle-dependent differences in expression of the 170 kilodalton and 180 kilodalton forms of topoisomerase II in NIH-3T3 cells. Cell Growth Differ *2*, 209-214.

Wozniak, A. J., and Ross, W. E. (1983). DNA damage as a basis for 4'-demethylepipodophyllotoxin-9-(4,6-O-ethylidene-beta-D-glucopyranoside) (etoposide) cytotoxicity. Cancer Res *43*, 120-124.

Wu, C. C., Li, T. K., Farh, L., Lin, L. Y., Lin, T. S., Yu, Y. J., Yen, T. J., Chiang, C. W., and Chan, N. L. (2011). Structural basis of type II topoisomerase inhibition by the anticancer drug etoposide. Science *333*, 459-462.

Wu, J., and Liu, L. F. (1997). Processing of topoisomerase I cleavable complexes into DNA damage by transcription. Nucleic Acids Res *25*, 4181-4186.

Yan, W. X., Mirzazadeh, R., Garnerone, S., Scott, D., Schneider, M. W., Kallas, T., Custodio, J., Wernersson, E., Li, Y., Gao, L., Federova, Y., Zetsche, B., Zhang, F., Bienko, M., and Crosetto, N. (2017). BLISS is a versatile and quantitative method for genome-wide profiling of DNA double-strand breaks. Nat Commun *8*, 15058.

Yang, F., Kemp, C. J., and Henikoff, S. (2015). Anthracyclines induce double-strand DNA breaks at active gene promoters. Mutat Res *773*, 9-15.

Yu, X., Davenport, J. W., Urtishak, K. A., Carillo, M. L., Gosai, S. J., Kolaris, C. P., Byl, J. A. W., Rappaport, E. F., Osheroff, N., Gregory, B. D., and Felix, C. A. (2017). Genome-wide TOP2A DNA cleavage is biased toward translocated and highly transcribed loci. Genome Res *27*, 1238-1249.

Zwelling, L. A., Michaels, S., Erickson, L. C., Ungerleider, R. S., Nichols, M., and Kohn, K. W. (1981). Protein-associated deoxyribonucleic acid strand breaks in L1210 cells treated with the deoxyribonucleic acid intercalating agents 4'-(9-acridinylamino) methanesulfon-m-anisidide and adriamycin. Biochemistry *20*, 6553-6563.

Figure 1

Figure 1. CC-seq maps covalent Spo11-linked DNA breaks in *S. cerevisiae* meiosis with nucleotide accuracy

A) Schematic of the CC-seq method.

B) Column-based enrichment of Spo11-linked DNA fragments detected by Southern blotting at the *his4*::*LEU2* recombination hotspot (**Methods**). Arrows indicate expected sizes of Spo11-DSBs.

C) Nucleotide resolution mapping of *S. cerevisiae* Spo11 hotspots by CC-seq or oligo-seq (Pan et al., 2011). Red and blue traces indicate Spo11-linked 5' DNA termini on the Watson and Crick strands, respectively. Grey arrows indicate positions of gene open reading frames.

D) Pearson correlation (r) of Spo11 CC-seq signal between Watson and Crick strands, offset by the indicated distances.

E) Average nucleotide composition over a 200 (top) and 30 bp (bottom) window centred on Spo11 breaks. Bases reported are for the top strand only.

HpM = Hits per million mapped reads per base pair.

Figure 2

Figure 2. CC-seq maps covalent Top2-linked DNA breaks in *S. cerevisiae* cycling cells with nucleotide accuracy

A) Serial dilution spot tests of VP16 tolerance for the indicated strains.

B) Nucleotide resolution mapping of *S. cerevisiae* Top2 CCs by CC-seq of the indicated strains after treatment for 4 hours with 1 mM VP16. Spo11-CC data is plotted for comparison. Top2 CC data were calibrated using a human DNA spike-in (**Methods**). Red and blue traces indicate CC-linked 5' DNA termini on the Watson and Crick strands, respectively. Grey arrows indicate positions of gene open reading frames. Lower panels show an expanded view of the region from 31.5 to 40 Kbp.

C) Quantification of Top2 and Spo11 CC signal stratified by genomic region. The genome was divided into intra and intergenic regions; the intergenic region was further divided into divergent, tandem and convergent based on orientation of flanking genes. Spo11 and Top2 activity mapped by CC-seq is expressed as box-and-whisker plots of density (upper and lower box limit: 3rd and 1st quartile; bar: median; upper and lower whisker: highest and lowest values within 1.5-fold of the interquartile range), or as the percentage of total mapped reads.

D) Local correlation of Top2 or Spo11 CC-seq signals. Top2 or Spo11 CC-seq data were binned at 50 bp resolution and the Pearson correlation calculated between bins of increasing separation.

E) Pearson correlation (r) of Top2 CC-seq signal on Watson and Crick strands, offset by the indicated distance.

HpM = Hits per million mapped reads per base pair.

Figure 3

Figure 3. CC-seq maps TOP2-linked DNA breaks in Human cells with nucleotide accuracy
A) Anti-TOP2β western slot blot of input, flow through, wash, and elution fractions. RPE-1 cells were treated or not with VP16, prior to processing according to **Figure 1A**.
B) Broad-scale maps of *H. sapiens* TOP2 CCs produced by CC-seq in RPE-1 cells ±VP16. Raw data were scaled, binned, smoothed and median subtracted prior to plotting (**Methods**). Chromatin compartments revealed by Hi-C eigenvector analysis (Darrow et al., 2016), and supercoiling revealed by bTMP ChIP-seq (Naughton et al., 2013) are shown for comparison.
C) Quantification of TOP2 CC in chromatin compartments A and B. Data are expressed as box-and-whisker plots of density as for **Figure 2C**.
D) Fine-scale mapping of *H. sapiens* TOP2 CCs by CC-seq in RPE-1 cells ±VP16. Red and blue traces indicate TOP2-linked 5' DNA termini on the Watson and Crick strands, respectively. Pale shaded areas are the same data smoothed according to local density (see **Methods**). RPE-1 CTCF and H3K4Me3 ChIP-seq data plus H3K27Ac ChIP-seq data overlaid from seven cell lines (ENCODE, 2012) is shown for comparison.
E) Medium-scale aggregate of END-seq mapped DSBs (Canela et al., 2017) surrounding nucleotide resolution CC-seq mapped TOP2 CCs.
F) Fine-scale strand-specific aggregate of END-seq mapped DSB ends (red and blue bars) and nucleotide resolution CC-seq mapped TOP2 CCs (grey bars) surrounding strong TOP2 CC sites.

Figure 4

Figure 4. CTCF-proximal TOP2 activity is tightly confined within nucleosome-depleted regions
A) Aggregation of TOP2 CCs in a 1 Kbp window centred on orientated CTCF motifs in human RPE-1 cells. Four example CTCF loci are shown orientated in the 5'-3' direction (top). A heatmap of all CTCF-motifs in the human genome, with 25 rows stratified by the strength of colocalising CTCF ChIP-seq peaks in RPE-1 cells (middle). The colour scale indicates average TOP2 CC density. Motifs are also stratified into 4 quartiles of CTCF-binding, and the average TOP2 CC distribution in each quartile plotted (bottom).
B) Aggregated TOP2 CC distribution (red line) in the highest quartile of CTCF-binding compared with the average MNase-seq signal (blue line).
C) Aggregated TOP2 CC distribution (black and red lines: single-nucleotide resolution and smoothed, respectively) and the average MNase-seq signal (blue line) surrounding strong TOP2 CC sites.
In (B) and (C) peaks in MNase-seq signal indicate inferred nucleosome positions (blue circles).

Figure 5

Figure 5. TSS-proximal TOP2 activity is strongly correlated with gene transcription
A) Aggregation of TOP2 CCs in a 10 Kbp window centred on orientated TSSs in human RPE-1 cells. Four example TSSs are shown orientated in the 5'-3' direction (top). A heatmap of all TSSs in the human genome, with 25 rows stratified by gene expression level in RPE-1 cells (middle). The colour scale indicates average TOP2 CC density. Motifs are also stratified into 4 quartiles of gene expression, and the average TOP2 CC distribution in each quartile plotted (bottom).
B) Comparison of CTCF-proximal and TSS-proximal TOP2 CC distributions in the highest quartile of CTCF-binding (pink) and gene-expression (blue), respectively.
C) Average TOP2 CC density in 10 Kbp regions centred on the highest quartile of TSS, CTCF, or regions where both features are present. Data are expressed as box-and-whisker plots of density as in **Figure 2C**. Statistical significance was determined using the KS test.
D) As in (C) but sum total TOP2 CCs found in these regions.

Figure 6

Figure 6. Local DNA sequence and methylation status direct the formation of Top2 CCs
A) The percentage of strong sites that are SSBs and DSBs in etoposide-treated *S. cerevisiae* and human cells.
B) Average nucleotide composition over a 30 bp window centred on the DSB or (inferred) SSB dyad axis, in etoposide-treated *S. cerevisiae* and human cells. Values reported are for the top strand only.
C) Heatmaps describing pairwise similarity of nucleotide composition patterns shown in (B). Rows 1-4 are the absolute differences between: yeast and human SSB patterns, yeast and human DSB patterns, yeast SSB and DSB patterns, and human SSB and DSB patterns, respectively.
D) The average number of TOP2 CCs at the +1 position relative to methylated and unmethylated cytosines in human RPE-1 cells, ±VP16. Statistical significance was determined using KS test.

Figure S1

Figure S1. CC-seq maps covalent Spo11-linked DNA breaks in *S. cerevisiae* meiosis with nucleotide accuracy (related to Figure 1)

A) Correlation of hotspot signals from CC-seq and oligo-seq.

B) Correlation of 500 bp binned Spo11 maps from two representative replicates.

C) Correlation of Spo11 cleavages on Watson and Crick strand when offset by 1 bp (boxed), relative to other offsets from -4 to +4 bp.

D) Spo11 breaks mapped by CC-seq (raw=black, or smoothed=red) anticorrelate with nucleosome occupancy measured by MNase-seq (blue).

E) Aggregation of Spo11 activity in a 10 Kbp window centred on orientated TSSs in *S. cerevisiae*. Three example TSSs are shown orientated in the 5'-3' direction (top). A heatmap of all TSSs in the *S. cerevisiae* genome, with 25 rows stratified by gene expression level in SK1 cells (middle). The colour scale indicates average Spo11 break density. Motifs are also stratified into 4 quartiles of gene expression in meiotic SK1 cells, and the average distribution of Spo11 activity in each quartile plotted (bottom).

Figure S2

Figure S2. CC-seq maps covalent Top2-linked DNA breaks in *S. cerevisiae* cycling cells with nucleotide accuracy (related to Figure 2).
A-C) Correlation of 1 Kbp binned Top2 CC maps from *pdr1Δ* (A), *pdr1Δsae2Δ* (B) and *pdr1Δmre11Δ* (C) cells treated with VP16.
D) Pairwise Pearson correlation values for 100 bp binned Top2 CC maps from all assayed conditions. Each condition is a pool of two biological replicates.
E) Nucleotide-resolution *S. cerevisiae* Top2 CC map of chromosome 3 for all assayed conditions. Each condition is a pool of two biological replicates.
F-H) The normalised number of Top2 CCs retained in the *pdr1Δ* (F), *pdr1Δsae2Δ* (G) and *pdr1Δmre11Δ* (H) cells treated ±VP16, after filtering to include only sites offset by the given number of base pairs. All data were normalised over a -100 to +100 bp window.
I-K) Pearson correlation (r) of Top2 CC-seq signal on Watson and Crick strands, offset by the indicated distance in *pdr1Δ* (I), *pdr1Δsae2Δ* (J) and *pdr1Δmre11Δ* (K) cells treated ±VP16.

Figure S3

Figure S3. Top2-linked DNA breaks in *S. cerevisiae* are not correlated with gene expression, anticorrelate with nucleosomes, and have biased nucleotide skews indicative of bent DNA (related to Figure 2)

A) Top2 CC mapped by CC-seq (raw=black, or smoothed=red) anticorrelate with nucleosome occupancy measured by MNase-seq (blue).

B) Aggregation of Top2 CCs in a 10 Kbp window centred on orientated transcription start sites (TSS) in *pdr1Δ S. cerevisiae*. Four example TSSs are shown orientated in the 5'-3' direction (top). Heatmap of all TSSs in the *S. cerevisiae* genome, with 25 rows stratified by gene expression level in vegetative growth (middle). Colour scale indicates average Top2 CC density. Motifs are also stratified into four quartiles of gene expression, and the average distribution of Top2 CCs in each quartile plotted for untreated and +VP16 conditions (bottom).

C) The percentage of strong sites that are SSBs and DSBs in etoposide-treated *pdr1Δ* and *pdr1Δmre11Δ S. cerevisiae*. Sites were thresholded at 1 HpM prior to sorting into DSB or SSB classes based on presence or absence of a 3 bp offset cognate (Obs.). As a control, the amplitudes of Top2 CCs (HpM) were randomised amongst the positions in the nucleotide resolution datasets, prior to thresholding and offset analysis as described above (Rand.)

D) Average nucleotide composition over a 200 bp window centred on the DSB dyad axis, in etoposide-treated *S. cerevisiae*. The position of the flanking regions (F.R.) identified in Gyrase mapping experiments, and the flanking regions observed here for Top2 are indicated in blue and black, respectively.

D) Average nucleotide composition over a 120 bp window centred on the DSB dyad axis, in etoposide-treated *S. cerevisiae*. The signal pattern is separated into A+T and G+C (top left and right), and into individual A, T, C and G plots (middle left, bottom left, middle right, and bottom right, respectively). The position of the flanking regions (F.R.) observed here for Top2 are indicated in black, and their ~10.5 bp periodicity is highlighted with black and red crosses.

Figure S4

Figure S4. CC-seq maps of TOP2-linked DNA breaks in Human cells are enriched by etoposide, and show high reproducibility and nucleotide accuracy (related to Figure 3)

A) Broad-scale *H. sapiens* TOP2 CC-seq maps in individual biological replicates of RPE-1 cells ±VP16. Raw data were binned at 100 Kbp prior to plotting. Each plot is offset on the y-axis by +0.3 HpB.

B) Replicate-to-replicate Pearson correlation values (r) for 10 Kbp binned TOP2 CC-seq maps of RPE-1 cells ±VP16.

C) Scatter plot of -VP16 and +VP16 TOP2 CC-seq maps binned at 10 Kbp resolution. Data were first scaled according to the estimated noise fraction (**Methods**), and are presented in a hexagonal-binned format, where the density of overplotting is indicated by the colour scale.

D) Violin plots of TOP2 CC-seq maps ±VP16 binned at 100 Kbp resolution. The inner black bar, black dot, and dotted horizontal line indicate the interquartile range, median, and expected mean Top2 CC density based on random distribution.

E) The normalised number of TOP2 CCs retained in the CC-seq maps in RPE-1 cells ±VP16 after filtering to include only sites offset by the given number of base pairs. Data were normalised over a -100 to +100 bp window.

Figure S5

Figure S5. CC-seq signal is TOP2-dependent

A) DNA content histograms of wild type (WT) and *TOP2B*[-/-] RPE-1 cells under asynchronous (10% FCS) and serum-deprived (0% FCS) conditions, as measured by FACS following propidium iodide (PI) staining. G1, S and G2 populations are clearly present under asynchronous growing conditions. A strong G1 arrest is observed in serum deprived conditions. Percentages of cells in each of the indicated regions (red dotted brackets) are given.

B) Western blots demonstrating the absence of TOP2β protein in serum deprived and asynchronous *TOP2B*[-/-] RPE-1 cells (left), and the absence of TOP2α in serum deprived wild type and *TOP2B*[-/-] RPE-1 cells (right). Ponceau S total protein loading is presented (Pon. S) for the left and right panels, and additionally a Ku80 loading control is included for the right panel.

C) Immunofluorescence experiment demonstrating induction of γ-H2AX foci (green) in asynchronous (Async.) and serum-deprived (Ser. Dep.) wild type (WT) and *TOP2B*[-/-] RPE-1 cells, all co-stained with DAPI (blue). Galleries of nine cells per condition were chosen randomly using Olympus ScanR Analysis software.

D) Quantification of (C). Numbers of γ-H2AX foci per cell were counted automatically using Olympus ScanR Analysis software. The mean ±SEM is reported for n= 3 biological replicate experiments.

E) Broad-scale *H. sapiens* TOP2 CC-seq maps in asynchronous and serum-deprived wild type (WT) and *TOP2B*[-/-] RPE-1 cells -VP16 (orange) and +VP16 (green). Raw hits on Watson and Crick strands were summed and smoothed according to local signal density (Fsize=501).

Figure S6

Figure S6. TOP2 CC-seq signal enrichment around CTCF and TSS sites, compared with END-seq DSB signal at TSSs

A) Fine-scale TOP2 CC-seq maps of *H. sapiens* CTCF-proximal loci in RPE-1 cells +VP16. Red and blue traces indicate TOP2-linked 5' DNA termini on the Watson and Crick strands, respectively. Pale shaded areas are the same data smoothed with a sliding 11 bp Hanning window. Red and Blue rectangles indicate the positions of CTCF motifs on the Watson and Crick strands respectively. The grey line indicates Hanning-smoothed sum of Watson and Crick TOP2 CCs.

B) Fine-scale mapping of TOP2 CCs surrounding three complex CTCF loci, processed as in (A).

C) Aggregation of TOP2 CCs in a 1 Kbp window centred on the subset of orientated CTCF motifs that can be assigned to a chromatin loop anchor in human RPE-1 cells (Darrow et al., 2016). Motifs are stratified into 4 quartiles of loop anchor interaction strength, and the average TOP2 CC distribution in each quartile plotted.

D) Fine-scale aggregation of TOP2 CCs (red) in a 800 bp window centred on TSSs in human RPE-1 cells, showing anticorrelation with aggregated MNase-seq signal (blue).

E) Aggregation of END-seq mapped DSBs in a 10 Kbp window centred on orientated TSSs in human MCF7 cells. Four example TSSs are shown orientated in the 5'-3' direction (top). A heatmap of all TSSs in the human genome, with 25 rows stratified by flanking gene expression level in MCF7 cells (middle). The colour scale indicates average END-seq DSB density. TSSs are also stratified into 4 quartiles based on strength of flanking gene expression, and the average END-seq DSB distribution in each quartile plotted (bottom).

Figure S7

Figure S7. TSS-proximal TOP2-linked DNA breaks in humans are correlated with gene length, independently from gene expression level (related to Figure 5); TSS-proximal Top2-linked DNA breaks in yeast are not strongly correlated with gene length, (related to Figure 2)

A) Aggregation of TOP2 CCs in a 10 Kbp window centred on orientated TSSs in human RPE-1 cells. A heatmap of all TSSs in the human genome, with 25 rows stratified by gene length (top). The colour scale indicates average TOP2 CC density. Motifs are also stratified into 4 quartiles of gene length, and the average TOP2 CC distribution in each quartile plotted (bottom).

B) Aggregation of Top2 CCs in a 10 Kbp window centred on orientated transcription start sites (TSS) in *pdr1Δ S. cerevisiae*. Heatmap of all TSSs in the *S. cerevisiae* genome, with 25 rows stratified by gene length level (top). Colour scale indicates average Top2 CC density. Motifs are also stratified into four quartiles of gene length, and the average distribution of Top2 CCs in each quartile plotted (bottom).

C) Scatter plot of human gene length and RPE-1 gene expression, showing no correlation.

D) Scatter plot of *S. cerevisiae* gene length and gene expression during vegetative growth, showing little correlation.