Loss of one Engrailed1 allele enhances induced α-synucleinopathy

Diptaman Chatterjee[a*] Daniel Saiz Sanchez[b,c*], Emmanuel Quansah[c], Nolwen L Rey[c], Sonia George[c],

Katelyn Becker[c], Zachary Madaj[d], Jennifer A Steiner[c], Jiyan Ma[c], Martha L Escobar Galvis[c], Jeffrey H

Kordower[a,c,e], Patrik Brundin[c]

[a]Department of Neurological Sciences, Rush University Medical Center, Chicago, IL, USA 60612

[b]Medical School, Ciudad Real, 13071 Castilla-La Mancha, Spain

[c]Center for Neurodegenerative Science, Van Andel Research Institute, Grand Rapids, MI, USA 49503

[d]Bioinformatics & Biostatistics Core, Van Andel Research Institute, Grand Rapids, MI, USA 49503

[e]Research Center for Brain Repair, Rush University Medical Center, Chicago, IL, USA 60612

* Shared first authors

Address for correspondence:

Patrik Brundin, M.D., Ph.D.

Center for Neurodegenerative Science

Van Andel Research Institute

333 Bostwick Ave., NE

Grand Rapids, MI 49503

USA

Telephone: (616) 234-5312

Fax: (616) 234-5183

Email: Patrik.Brundin@vai.org

Abstract

Background: Parkinson's disease (PD) is a synucleinopathy that has multiple neuropathological characteristics, with nigrostriatal dopamine system degeneration being a core feature. Current models of PD pathology typically fail to recapitulate several attributes of the pathogenic process and neuropathology. We aimed to define the effects of combining a mouse model exhibiting multiple PD-like changes with intrastriatal injections of α-synuclein (α-syn) pre-formed fibrils (PFFs) aggregates. We employed the heterozygous Engrailed 1 ($En1^{+/-}$) mouse that features several pathophysiological hallmarks of clinical PD. Objective: To test the hypothesis that the neuropathological changes in the $En1^{+/-}$ mice will promote formation of α-syn aggregates following intrastriatal injections of pathogenic human α-syn PFFs. Methods: We unilaterally injected PFFs into the striata of 1 month-old $En1^{+/-}$ and control wild-type mice and euthanized animals at 3 months for post-mortem analysis. Results: Using immunohistochemistry and unbiased stereology, we established that PFF-injected $En1^{+/-}$ mice exhibited a near-threefold increase in pS129-α-syn-positive neurons in the substantia nigra compared to PFF-injected wild-type mice. The PFF-injected $En1^{+/-}$ mice also displayed significant increases in pS129-α-syn-positive neurons in the amygdala and ventral tegmental area; regions of known PD pathology with projections to the striatum. Additionally, we observed amplified pS129-α-syn-positive aggregation in $En1^{+/-}$ mice in multiple cortical regions. Conclusions: Following intrastriatal injection of PFFs, absence of an $En1$ allele leads to additional aggregation of pathological α-syn, potentially due to $En1$-loss mediated nigrostriatal impairment. We propose that further development of this double-hit model could result in a PD mouse model that predicts which experimental therapies will be effective in PD.

Keywords

α-Synuclein; Parkinson's disease; engrailed; proteinopathy; misfolded proteins; propagation; mouse models; neurodegeneration

Introduction

Parkinson's disease (PD) is a synucleinopathy where degeneration of nigrostriatal dopamine

neurons is a core neuropathological feature. It is believed that mitochondrial deficits and aggregation of

misfolded α-synuclein (α-syn) are key components of the pathogenic process[1, 2]. Mitochondrial

insults, particularly complex I deficiencies, have long been suggested to contribute to the development

of PD pathology[3]. Toxins that cause complex I impairment, e.g. 1-methyl-4-phenyl-1,2,3,6-

tetrahydropyridine (MPTP), have been used to generate animal models of PD, but these typically cause

relatively acute degeneration of nigral dopaminergic neurons that is not representative of the

progressive pathology of PD. α-Syn pathology has in a few cases been observed in MPTP-treated

animals, but they are not a consistent feature of these models [4]. Furthermore, rotenone injections can

lead to nigral neuron degeneration, as well as some α-syn pathology, but reproducibility can be

problematic due to high rodent mortality [5, 6].

We [7] and others [8, 9] have previously characterized a PD-like neuropathology in the

Engrailed1 heterozygous mouse model (*En1+/-*) including mitochondrial deficits, autophagic

disturbances, axonal degeneration, neuroinflammation, diminished potassium-evoked dopamine release

in the dorsal striatum and progressive loss of midbrain dopamine neurons. Engrailed1 has been shown

to contribute to axonal maintenance and mitochondrial support in dopaminergic neurons into

adulthood and counteract deficits associated with experimentally induced neuronal insults (MPTP, 6-

OHDA, α-syn), both at the level of transcriptional regulation and translational output essential for

mitochondrial function [8-13]. The primary aim of the present study was to define the effect of

combining the complex neuropathology associated seen in *En1+/-* mice with the progressive

proteinopathy that follows intracerebral injection of α-syn fibrils [14]. Thus, we injected α-syn pre-

formed fibrils (PFFs) into the striatum of *En1+/-* mice. Eight weeks later we found significantly greater

numbers of phosphorylated serine-129 α-syn (p-S129)-positive inclusions in the substantia nigra (SN)

and ventral tegmental area (VTA), as well as in the basolateral amygdala (BLA) of male mice when

compared to wild-type (WT) mice injected with the same amount of PFFs. We believe this novel *En1*-PFF

mouse model represents a paradigm that mimics several aspects of cardinal PD pathologies. We believe

that it can prove to be a highly relevant model system in which one can test experimental therapeutics

aimed at slowing PD progression and obtain results that are highly predictive of clinical success.

Methods and materials

Animals

En1$^{+/-}$ mice were housed 4-5 per cage under a regular cycle (12 h light/12 h dark) and had access to food

and water *ad libitum*. The mice were treated in accordance with the Guide for the Care and Use of

Laboratory Animals (National Institutes of Health) and all procedures and experimental conditions were

undertaken with the prior approval of the Institutional Animal Use and Care Committee of Van Andel

Research Institute. The *En1*$^{+/-}$ mice were maintained on an OF1 genetic background as described in

Sonnier et al., 2007 [8-11].

Purification of recombinant alpha-synuclein and assembly of pre-formed fibrils

Recombinant human α-syn was purified similarly to a previously published protocol [15, 16]. Human α-

syn was used in this double-hit model with to provide a higher-fidelity platform for potential

downstream pre-clinical screening of α-syn targeting therapeutics. Briefly, the protein was expressed in

BL21 *E.coli* transfected with a pT7-7 expression plasmid expressing human α-syn. Once expressed, cells

were collected and stored at -80°C. For purification, cells were lysed by sonication and boiling, and

centrifuged to remove cell debris. The α-syn-containing supernatant was dialyzed overnight in 10 mM

Tris, pH 7.5, 50 mM NaCl, and 1 mM EDTA, using SnakeSkin Dialysis Tubing MWCO 7,000 (Thermo

Scientific). Chromatographic separation was performed using a Superdex 200 Column (GE Healthcare

Life Sciences) and a Hi-trap Q HP anion exchange column (GE Healthcare Life Sciences). Fractions

containing α-syn were identified by SDS-PAGE and Coomassie staining, and then dialyzed overnight into

PBS buffer (Life Sciences). A NanoDrop 2000 (Thermofisher) was used to determine the protein

concentration by OD280 reading and protein was concentrated to 5 mg/mL using a Vivaspin protein

concentrator spin column with a MWCO of 5kDa (GE Healthcare). Aliquots of 500 μL were stored at -

80°C until use. For amyloid fibril assembly, purified recombinant α-syn was thawed and subjected to

continuous shaking at 1,000 r.p.m at 37˚C in a Thermomixer (Eppendorf) for 7 days. Fibril morphology

was confirmed by imaging using a Tecnai G2 Spirit TWIN transmission electron microscope (FEI

Company) at 120kV and Thioflavin T (Sigma) fluorescence reading (data not shown). Fibrils were

aliquoted and frozen at -80˚C until use.

Stereotactic injections

Prior to injection, human α-syn PFFs were thawed and then sonicated at RT with a water-bath cup-horn

sonicator (Misonix XL2020, 50% power, 120 pulses 1 s ON, 1 s OFF). Post-sonication fibril morphology

was validated by TEM imaging protocol as described above. Prior to injection, 4-week-old mice (both

sexes) were anesthetized with an isoflurane/oxygen mixture and injected unilaterally with either 2 μL of

PFFs (5 μg/μl) or 2 μL of PBS as a control in the dorsal striatum (coordinates from bregma: AP: + 0.2 mm;

ML: -2.0 mm; DV: - 2.6 mm from dura) at a rate of 0.2 μL/min using a glass capillary attached to a 10 μL

Hamilton syringe. After injection, the capillary was left in place for 4 minutes before being slowly

removed.

Euthanasia

At 8 weeks post PFF-injection, mice were deeply anesthetized with sodium pentobarbital and perfused

transcardially with 0.9% saline followed by 4% paraformaldehyde in phosphate buffer. Brains were

collected and post-fixed for 24 hours in 4% PFA, and then stored at 4˚C in 30% sucrose in phosphate

buffer until sectioning.

Histology

Brains were then sectioned into 40 μm-thick free-floating coronal sections using a freezing microtome.

Brain sections were stored in cryoprotectant and quenched with sodium periodate. During the staining

protocols, sections were incubated with primary antibodies for tyrosine hydroxylase (mouse anti-TH,

ImmunoStar) at 1:1000 dilution and p-S129 α-syn (rabbit anti-pS129, Abcam [EP1536Y]) at 1:1000

dilution. Sections were incubated with respective anti-mouse or anti-rabbit biotinylated secondary

antibodies (Vector Laboratories) and conjugated with an ABC-HRP biotin/avidin complex kit (Vector

Laboratories). Staining was developed with 3,3'-diaminobenzidine and sections were mounted for

imaging and analysis. Representative images of pS129 pathology were acquired at 20x magnification

using a NIKON Eclipse Ni-U microscope.

pS129 α-synuclein scoring

The α-syn inclusions (cell body and neurites) were detected following pS129 staining of a whole series of

coronal sections (at 240 μm intervals) per animal. The stained slides were blind-coded prior to analysis.

We evaluated pS129 inclusions in 8–10 animals per experimental group. Using the 20x magnification on

a NIKON Eclipse Ni-U microscope, we evaluated all sections individually and assigned scores from 0 to 4

based on the relative levels of pS129-positive inclusions (such that 0 = no aggregates, 1 = sparse, 2 =

mild, 3 = dense, 4 = very dense), as previously described [17, 18]. The pathology depicted in the brain

regions in Fig. 5A are representative of the average score of animals in each experimental group. We

also used the average score values in each brain region per group to generate a heatmap (Fig. 5B) to

represent the entire dataset.

Stereology and Densitometry

To evaluate the statistical significance of the observed α-syn pathology, we counted α-syn inclusions

with unbiased stereological counting methods using the optical fractionator probe in StereoInvestigator

(v10.40, MBF Biosciences). Here, full series of brain sections (5-7 sections) spaced at 240 μm were

analyzed per animal and a total of five to seven animals per group (all n ≥ 5 per group) were included. To

count the pS129+ labeled cell bodies, contours of each region of interest (SN, VTA, BLA and motor

cortex) were drawn using a 4x objective. In the VTA, quantifications were accomplished with 60x

magnification using a counting frame of 90 μm X 90 μm and grid size of 180 μm (medio-lateral) and 180

μm (dorso-ventral), along with a guard zone of 2 μm and a dissector height of 15 μm. For each region of

interest, these quantification parameters were adjusted to assess at least 300 total cells per animal and

per side, with the error of coefficient ($m = 1$) being less than 0.1. Stereology on TH+ neurons in the SN

was performed on sections spaced 480 μm with a 70 μm X 70 μm counting frame, a grid size of 221 μm

X 221 μm, guard zone of 1 μm, and dissector height of 15 μm. Striatal TH+ optical density was measured

using Scion Image (v.163, Meyer Instruments) and imaged on an Olympus BH-2 microscope (Olympus).

Densitometric measurements were averaged from a minimum of 2 sections spaced 480 μm apart.

Statistics

All graphical data are presented as mean ± SEM. Stereological count data were analyzed using negative

binomial mixed-effects regression with zero-inflated adjustments in regions with negligible count data.

Densitometric measurements were assessed using a linear mixed-effects regression with a random

intercept for each animal subject. All statistical analyses were performed using GraphPad Prism (v7.03)

or R (v3.4.4). Significant findings and specific statistical analyses are listed in the figure legends.

Results

Study design

The study was performed utilizing the *En1*$^{+/-}$ (OF1 background) mouse line that features a knock-

in LacZ cassette in one allele of *En1* with WT levels of Engrailed 2 (Fig. 1A) [9]. The most recent

characterization of the *En1*$^{+/-}$ mouse model [7, 8, 19] revealed multiple indices of pathophysiology that

follow clinical features of PD. Critically, *En1*$^{+/-}$ animals exhibit downregulation of mitochondrial complex I

proteins, progressive loss of midbrain dopamine neurons (more pronounced in A9 than A10 neurons),

dysregulation of autophagic markers, axonal pathology and neuroinflammation (Fig. 1B) [7]. We tested

the hypothesis that the aforementioned nigrostriatal pathology in the *En1*$^{+/-}$ mouse would lead to an

exacerbation of α-syn aggregate formation in mice where the synucleinopathy is triggered by injection

of α-syn fibrils. Thus, we generated PFFs according to established protocols, and verified them by

transmission electron microscopy (Fig. 1C). To induce α-syn-mediated proteinopathy, we injected 4-

week old En1$^{+/-}$ mice with PFFs unilaterally into the striatum. After 2 months we euthanized the mice

and conducted neuropathological examinations of their brains using markers for α-syn pathology and

dopamine neuron survival (Fig. 1D, E).

α-Synuclein pathology is exacerbated following injections of fibrils in En1$^{+/-}$ *mice*

At eight weeks after unilateral, intrastriatal injection of PFFs into *En1*$^{+/-}$ and WT mice, we

assessed the numbers of α-syn-positive puncta in four regions with direct projections to the striatum –

the SN, BLA, motor cortex and VTA (Fig. 2, 3). We used unbiased stereology to count cell bodies

immunolabeled with p-S129-α-syn. The $En1^{+/-}$ mice injected with PFFs (*En1*-PFF mice) displayed a greater

than three-fold increase in the number of p-S129-α-syn-positive inclusions in the SN, ipsilateral to the

injection of PFFs, compared to WT mice injected with an identical amount of PFFs (WT-PFF mice) (1243 ±

262 cells and 406 ± 52 cells, respectively, p=0.009) (Fig. 2A, B). The numbers of p-S129-α-syn-positive

inclusions in the contralateral SN were negligible (Fig. 2B). We observed no Lewy-like α-syn pathology in

any PBS injected control animals. To assess the effects of α-syn pathology on nigrostriatal integrity, we

assessed the number of dopaminergic neurons in the SN (TH+) and the optical density of TH+ fibers in

the striatum. We observed a significant main effect in the loss of striatal fiber innervation caused by the

En1 genotype (p=0.001) as expected, but did not observe any added significant effect due to injection of

PFFs (Fig. 2D). At the level of the substantia nigra, however, the $En1^{+/-}$ animals (regardless of whether

they were injected with PBS or PFFs) did not exhibit significant loss of nigral TH-immunoreactive neurons

(Fig. 2C).

In the ipsilateral BLA, there was no significant difference in α-syn pathology between *En1*-PFF

and WT-PFF mice (2202 ± 476 cells and 1046 ± 208 cells, respectively, p = 0.428, Fig. 3A, B). We noted

that the variance was large, and when we analyzed the impact of biological sex we found a marked sex

difference, with male $En1^{+/-}$ mice displaying 359% increase (p=0.007, compared to WT-PFF mice) in

pS129+ deposition in the BLA ipsilateral to PFF injection and females exhibiting a non-significant trend

for a 14% increase (p=0.350) (Fig. 3B). In the VTA, similar to the SN, we observed a greater than 3-fold

increase in cells with α-syn pathology from the ipsilateral side of *En1*-PFF mice relative to WT mice (821

± 218 cells and 264 ± 34 cells, respectively, p = 0.033), with negligible pathology in the contralateral side

(Fig. 3C). In the motor cortex, we observed no statistically significant difference in α-syn pathology

between *En1*-PFF and WT-PFF mice (Fig. 3D).

To map long distance α-syn pathology propagation in the *En1*-PFF mice, we used a semi-

quantitative scale and determined a mean pathology score based upon p-S129-α-syn-positive perikarya

and neurites in all brain regions where α-syn inclusions were detectable. Enhanced pathology (both

ipsilateral and contralateral to PFF injection) was observed in *En1*-PFF mice across a number of cortical

regions, as depicted in Figure 4. The pathology scoring data is summarized in brain- and heat-maps (Fig.

5A, B). While there was less pathology in the olfactory bulb, we observed increased α-syn pathology in

the frontal association, orbital, motor, ectorhinal, somatosensory, perirhinal, auditory, and entorhinal

cortices of both *En1*-PFF and WT-PFF mice (Fig. 4), primarily ipsilateral to the intrastriatal PFF injection,

with more pronounced pathology observed in *En1*-PFF mice (Fig. 4, 5).

Discussion

We quantified p-S129-α-syn-positive pathology following injection of α-syn PFFs into the

striatum of mice and found that the absence of one *En1* allele led to significantly increased numbers of

cells displaying α-syn aggregates in the SN and VTA, as well as in the BLA of male mice. Qualitatively, we

also observed more p-S129-α-syn-positive staining in several cortical regions that project to the

striatum. We have previously investigated many facets of neuropathology in *En1*[+/-] mice and showed

that this mouse model exhibits several neuropathological features resembling those seen in PD, but

lacks α-syn aggregates [7]. Unusual features of the *En1*[+/-] mouse that are particularly pertinent to PD are

the protracted degeneration of nigral dopamine neurons over several months, with a later and less

pronounced loss of dopamine neurons in the VTA, as well as the early onset reductions in evoked striatal

dopamine release and a conspicuous axonal pathology reminiscent of the axonal dying back that is

believed to occur in PD [7, 20]. Notably, the first signs of axonopathy precede nigral degeneration in

En1[+/-] mice, adding to their relevance to PD where loss of dopaminergic terminals in the putamen also

has been shown to precede nigral cell loss [21].

There are many potential factors that could explain how heterozygous loss of *En1* exacerbates

pS129-α-syn pathology. Mitochondrial deficits in midbrain dopamine neurons that occur in *En1*[+/-] mice

may aggravate fibril recruitment of endogenous α-syn, and subsequent aggregation and propagation.

Deficiencies in mitochondrial complex I proteins, Ndufs 1 and 3, as well as critical mitochondrial

homeostatic mediator, LaminB2, are signature hallmarks in *En1* model pathophysiology [8, 22]. *En1*[+/-]

mice also feature robust oxidative stress, most likely attributed to signaling downstream of

mitochondrial impairment or to increases in LINE-1 expression and DNA damage [11, 12, 23]. Oxidative

damage in En1-expressing neurons may induce enhanced α-syn modification and accelerate

proteopathic seeding capacity.

Although the increase in p-S129-α-syn labeling was most prominent in the SN of the *En1*[+/-] mice,

a significant increase in p-S129-α-syn-positive inclusions was also apparent in the ipsilateral VTA. It is

possible that some of the injected PFF, which was targeted into the central striatum, may have diffused

into the ventral striatum leading to uptake of α-syn fibrils by terminals from dopaminergic neurons

located in the VTA. The deposition of p-S129-α-syn was not as pronounced in VTA neurons as in the SN,

possibly due to the anatomical coordinates of PFF injection into the striatum primarily leading to

targeting of central and dorsal striatal regions innervated by substantia nigra and not VTA. It is also

possible, however, that dopaminergic neurons in the SN are more impaired due to reduced *En1* levels

than those in the adjacent VTA dopamine neurons, which would be consistent with earlier reports of

greater dopamine neuron loss in the nigra compared to VTA in *En1*[+/-] mice [9, 24]. We did not observe

reduction in the number of TH-immunoreactive neurons in the SN of *En1*[+/-] mice; however, we detected

significant decreases in striatal fiber innervation which are known to precede SN neuron loss both in the

En1 heterozygous mouse and in clinical PD [7, 21]. Although SN neuron loss has been previously

reported at three months of age [9], genetic background is known to significantly impact the extent of

degeneration of nigral dopamine neurons [20], and subtle differences in such genetic susceptibility loci

between our present $En1^{+/-}$ colony and those used in earlier papers could potentially have affected the

results. Notably, the intrastriatal injections of PFFs did not cause any additional striatal degeneration or

accelerated nigral neuron loss. This is not surprising considering that our follow up after PFF injection

was only 2 months (mice aged 3 months at the time of sacrifice) and earlier work indicates that in WT

mice the loss of nigral TH-positive neurons does not become significant until between 3 and 6 months

after intrastriatal PFF injection, reaching 35% at 6 months [15]. We opted to inject the PFFs into $En1^{+/-}$

mice that were only one-month old, preceding the time when they exhibit extensive nigrostriatal axonal

degeneration, to ensure that the nigrostriatal neurons were still capable of uptake and transport of the

PFFs. Future studies with longer survival times are necessary to determine whether the PFF injections

cause nigral neuron death that is additive to the nigral cell death that occurs in $En1^{+/-}$ mice (estimated to

be around 40% at 6 months of age), or if it is the identically vulnerable subpopulation of nigral dopamine

neurons that dies in both paradigms.

Interestingly, we also observed increased α-syn deposition in the BLA of male $En1^{+/-}$ mice. It is

unlikely that the increased α-syn pathology in the BLA is primarily due to cell-autonomous deficits in BLA

neurons because there is no reported evidence of *En1* expression in the adult amygdala. Notably,

neurons in amygdala project to both SN and striatum [25]. Thus, the increased α-syn pathology could be

due to mitochondrial deficits in nigrostriatal dopamine neurons in $En1^{+/-}$ mice resulting in a greater

amount of aggregated α-syn seeds, or the neuroinflammatory environment induced by

neurodegeneration in the SN and striatum. While we observed a significant exacerbation of pathology in

the BLA of male $En1^{+/-}$ mice injected with PFFs, compared to WT mice similarly injected with PFFs, it is

unclear why female $En1^{+/-}$ mice did not show a similar increase in proteinopathy compared to WT female

mice. There are no previously reported sex differences in phenotypes of $En1^{+/-}$ mice, and it is unclear

that the impact of biological sex on the neuropathology of *En1*$^{+/-}$ mice has ever been addressed. It is

likely that due to the relatively short delay between PFF injection and analysis of neuropathology in our

study, female mice may develop equivalent levels of amygdala pathology latently. The divergence in

pathology may also be a result of intrinsic sex differences in amygdala networks and gene expression,

previously described in rodents, that may alter vulnerability to proteinopathy [26, 27]. Future studies

will require further investigation of sexual dimorphism pertaining to the *En1* allele loss and how it may

affect relevant brain structures.

Using stereological analysis, we found a non-significant trend for an increase in p-S129-α-syn-

positive aggregates in the ipsilateral motor cortex of the *En1*$^{+/-}$ mice. Bilaterally, in several other

neocortical brain regions which also project to the striatum and where we did not conduct formal

quantification of α-syn pathology, the overall visual impression was that there was more p-S129-α-syn

staining in the *En1*$^{+/-}$ mice (Fig. 4, 5 A, B). While we did not attempt to quantify pathology in all the

cortical regions and cannot judge whether there were robust effects of the absence of one *En1* allele, it

is possible our study was underpowered to detect differences in α-syn pathology between the *En1*$^{+/-}$ and

WT mice in the motor cortex. It is also likely that the precise location of the PFF injections in the

striatum, which receives topographically region-specific corticostriatal projections, contributed to a high

variability that masked a potential effect.

Taken together, we suggest that the *En1*-PFF mouse represents a potentially powerful PD mouse

model, with many molecular and neuropathological features resembling PD. Previous descriptions of the

En1$^{+/-}$ mouse have reported that it exhibits progressive, retrograde degeneration of dopamine neurons

focused in the SN and less marked in the VTA [7, 9] as well as reduced striatal levels of dopamine,

neuroinflammation, and changes in markers of autophagy [7]. In the current study, we found that the

additional hit of a unilateral intrastriatal PFF injection provides α-syn aggregate pathology which is

pertinent to PD pathology, and which was enhanced in *En1*$^{+/-}$ mice compared to wild type mice. The

combination of all these findings makes the *En1*-PFF mouse relevant for use in the testing of

experimental therapeutics. Further studies are needed with bilateral intrastriatal injections of PFFs and

long-term survival of animals to establish how severe the nigral degeneration and synucleinopathy can

become, and whether the mice develop a range of progressive motor and non-motor deficits that mimic

those seen in PD develop.

Acknowledgments

We acknowledge the Van Andel Research Institute and the many individuals and corporations that

financially support research into neurodegenerative disease at the Institute. D.S.S. was supported by a

José Castillejo Mobility Grant (CAS15/00201) awarded by the Ministry of Education, Culture and Sport

(Madrid, Spain). N.L.R.'s work is supported by the Peter C. and Emajean Cook Foundation. J.M. is

supported by National Institutes of Health grants (R21NS101676 and R01NS060729). J.H.K. is supported

by a Rush University center grant from the Parkinson's Disease Foundation. J.H.K. reports additional

support from the National Institutes of Health, the Michael J. Fox Foundation, and the Multiple System

Atrophy Coalition which are outside but relevant to the submitted work. P.B. is supported by grants

from the National Institutes of Health (1R01DC016519-01 and 5R21NS093993-02). J.M., P.B. and J.H.K.

are supported by an NIH IGNITE grant (1R21NS106078-01A1). P.B. reports additional grants from Office

of the Assistant Secretary of Defense for Health Affairs (Parkinson's Research Program, Award No.

W81XWH-17-1-0534), The Michael J Fox Foundation, National Institutes of Health, Cure Parkinson's

Trust, which are outside but relevant to the submitted work. We additionally acknowledge the technical

expertise of Lindsay Meyerdirk and Emily Schulz in their contributions to this study.

Conflicts of Interest

J.H.K. has received commercial support from Fujifilm-Cellular Dynamics. P.B. has received commercial

support as a consultant from Renovo Neural, Inc., Roche, and Teva Inc, Lundbeck A/S, AbbVie,

Neuroderm, Fujifilm-Cellular Dynamics, IOS Press Partners and CuraSen. P.B. has received commercial

support for grants/research from Renovo and Teva/Lundbeck. P.B. has ownership interests in Acousort

AB and is on the steering committee of the NILO-PD trial.

All other authors do not report conflicts of interest.

References

[1]   Ryan BJ, Hoek S, Fon EA, Wade-Martins R (2015) Mitochondrial dysfunction and mitophagy in Parkinson's: from familial to sporadic disease. *Trends Biochem Sci* **40**, 200-210.

[2]   Wong YC, Krainc D (2017) alpha-synuclein toxicity in neurodegeneration: mechanism and therapeutic strategies. *Nat Med* **23**, 1-13.

[3]   Schapira AH (2008) Mitochondria in the aetiology and pathogenesis of Parkinson's disease. *Lancet Neurol* **7**, 97-109.

[4]   Gibrat C, Saint-Pierre M, Bousquet M, Levesque D, Rouillard C, Cicchetti F (2009) Differences between subacute and chronic MPTP mice models: investigation of dopaminergic neuronal degeneration and alpha-synuclein inclusions. *J Neurochem* **109**, 1469-1482.

[5]   Cannon JR, Tapias V, Na HM, Honick AS, Drolet RE, Greenamyre JT (2009) A highly reproducible rotenone model of Parkinson's disease. *Neurobiol Dis* **34**, 279-290.

[6]   Jagmag SA, Tripathi N, Shukla SD, Maiti S, Khurana S (2015) Evaluation of Models of Parkinson's Disease. *Front Neurosci* **9**, 503.

[7]   Nordstroma U, Beauvais G, Ghosh A, Pulikkaparambil Sasidharan BC, Lundblad M, Fuchs J, Joshi RL, Lipton JW, Roholt A, Medicetty S, Feinstein TN, Steiner JA, Escobar Galvis ML, Prochiantz A, Brundin P (2015) Progressive nigrostriatal terminal dysfunction and degeneration in the engrailed1 heterozygous mouse model of Parkinson's disease. *Neurobiol Dis* **73**, 70-82.

[8]   Alvarez-Fischer D, Fuchs J, Castagner F, Stettler O, Massiani-Beaudoin O, Moya KL, Bouillot C, Oertel WH, Lombes A, Faigle W, Joshi RL, Hartmann A, Prochiantz A (2011) Engrailed protects mouse midbrain dopaminergic neurons against mitochondrial complex I insults. *Nat Neurosci* **14**, 1260-1266.

[9]   Sonnier L, Le Pen G, Hartmann A, Bizot JC, Trovero F, Krebs MO, Prochiantz A (2007) Progressive

loss of dopaminergic neurons in the ventral midbrain of adult mice heterozygote for Engrailed1.

J Neurosci **27**, 1063-1071.

[10]   Fuchs J, Stettler O, Alvarez-Fischer D, Prochiantz A, Moya KL, Joshi RL (2012) Engrailed signaling

in axon guidance and neuron survival. *Eur J Neurosci* **35**, 1837-1845.

[11]   Rekaik H, Blaudin de The FX, Fuchs J, Massiani-Beaudoin O, Prochiantz A, Joshi RL (2015)

Engrailed Homeoprotein Protects Mesencephalic Dopaminergic Neurons from Oxidative Stress.

Cell Rep **13**, 242-250.

[12]   Rekaik H, Blaudin de The FX, Prochiantz A, Fuchs J, Joshi RL (2015) Dissecting the role of

Engrailed in adult dopaminergic neurons--Insights into Parkinson disease pathogenesis. *FEBS*

Lett **589**, 3786-3794.

[13]   Simon HH, Thuret S, Alberi L (2004) Midbrain dopaminergic neurons: control of their cell fate by

the engrailed transcription factors. *Cell Tissue Res* **318**, 53-61.

[14]   Paumier KL, Luk KC, Manfredsson FP, Kanaan NM, Lipton JW, Collier TJ, Steece-Collier K, Kemp

CJ, Celano S, Schulz E, Sandoval IM, Fleming S, Dirr E, Polinski NK, Trojanowski JQ, Lee VM,

Sortwell CE (2015) Intrastriatal injection of pre-formed mouse alpha-synuclein fibrils into rats

triggers alpha-synuclein pathology and bilateral nigrostriatal degeneration. *Neurobiol Dis* **82**,

185-199.

[15]   Luk KC, Kehm V, Carroll J, Zhang B, O'Brien P, Trojanowski JQ, Lee VM (2012) Pathological alpha-

synuclein transmission initiates Parkinson-like neurodegeneration in nontransgenic mice.

Science **338**, 949-953.

[16]   Volpicelli-Daley LA, Kirik D, Stoyka LE, Standaert DG, Harms AS (2016) How can rAAV-alpha-

synuclein and the fibril alpha-synuclein models advance our understanding of Parkinson's

disease? *J Neurochem* **139 Suppl 1**, 131-155.

[17]   Rey NL, George S, Steiner JA, Madaj Z, Luk KC, Trojanowski JQ, Lee VM, Brundin P (2018) Spread

387   of aggregates after olfactory bulb injection of alpha-synuclein fibrils is associated with early

388   neuronal loss and is reduced long term. *Acta Neuropathol* **135**, 65-83.

389   [18]   Rey NL, Steiner JA, Maroof N, Luk KC, Madaj Z, Trojanowski JQ, Lee VM, Brundin P (2016)

390   Widespread transneuronal propagation of alpha-synucleinopathy triggered in olfactory bulb

391   mimics prodromal Parkinson's disease. *J Exp Med* **213**, 1759-1778.

392   [19]   Ghosh A, Tyson T, George S, Hildebrandt EN, Steiner JA, Madaj Z, Schulz E, Machiela E,

393   McDonald WG, Escobar Galvis ML, Kordower JH, Van Raamsdonk JM, Colca JR, Brundin P (2016)

394   Mitochondrial pyruvate carrier regulates autophagy, inflammation, and neurodegeneration in

395   experimental models of Parkinson's disease. *Sci Transl Med* **8**, 368ra174.

396   [20]   Kurowska Z, Jewett M, Brattas PL, Jimenez-Ferrer I, Kenez X, Bjorklund T, Nordstrom U, Brundin

397   P, Swanberg M (2016) Identification of Multiple QTLs Linked to Neuropathology in the Engrailed-

398   1 Heterozygous Mouse Model of Parkinson's Disease. *Sci Rep* **6**, 31701.

399   [21]   Kordower JH, Olanow CW, Dodiya HB, Chu Y, Beach TG, Adler CH, Halliday GM, Bartus RT (2013)

400   Disease duration and the integrity of the nigrostriatal system in Parkinson's disease. *Brain* **136**,

401   2419-2431.

402   [22]   Stettler O, Joshi RL, Wizenmann A, Reingruber J, Holcman D, Bouillot C, Castagner F, Prochiantz

403   A, Moya KL (2012) Engrailed homeoprotein recruits the adenosine A1 receptor to potentiate

404   ephrin A5 function in retinal growth cones. *Development* **139**, 215-224.

405   [23]   Blaudin de The FX, Rekaik H, Peze-Heidsieck E, Massiani-Beaudoin O, Joshi RL, Fuchs J,

406   Prochiantz A (2018) Engrailed homeoprotein blocks degeneration in adult dopaminergic neurons

407   through LINE-1 repression. *EMBO J* **37**.

408   [24]   Sgado P, Alberi L, Gherbassi D, Galasso SL, Ramakers GM, Alavian KN, Smidt MP, Dyck RH, Simon

409   HH (2006) Slow progressive degeneration of nigral dopaminergic neurons in postnatal Engrailed

410   mutant mice. *Proc Natl Acad Sci U S A* **103**, 15242-15247.

411   [25]   Lee HJ, Gallagher M, Holland PC (2010) The central amygdala projection to the substantia nigra

412   reflects prediction error information in appetitive conditioning. *Learn Mem* **17**, 531-538.

413   [26]   Cooke BM, Woolley CS (2005) Sexually dimorphic synaptic organization of the medial amygdala.

414   *J Neurosci* **25**, 10759-10767.

415   [27]   Lin LC, Lewis DA, Sibille E (2011) A human-mouse conserved sex bias in amygdala gene

416   expression related to circadian clock and energy metabolism. *Mol Brain* **4**, 18.

Figure Legends

Fig.1 – Inducing α-syn pathology in *En1*[+/-] mice. Engrailed1 expression is vital for CNS development and

maintenance, particularly for nigrostriatal projections. **A)** *En1*[+/-] mice feature a LacZ insertion cassette in

one allele and feature **B)** progressive "dying-back" nigrostriatal degeneration, dysfunctional mechanisms

in the autophagy-lysosomal pathway, and pathologic neuroinflammatory activation. *En1*[+/-] mice also

feature mitochondrial respiratory chain perturbations and motor dysfunction associated with PD clinical

pathology (Nordström *et. al.*, 2015; Sonnier *et. al.*, 2007) [7, 9]. **C)** Human pre-formed α-syn fibrils were

sonicated, validated by TEM, and injected into the dorsal striatum. **D)** PFF injection causes robust pS129

deposition at the site of injection. **E)** Both WT and *En1*[+/-] animals were unilaterally injected at one month

of age and sacrificed for post-mortem histology eight weeks following PFF injection.

Fig.2 – α-Syn pathology is preferentially exacerbated along the nigrostriatal tract. A)

Microphotographs of α-syn pathology (pS129-immunoreactive cell bodies and neurites) in the

substantia nigra of WT and *En1*[+/-] animals unilaterally injected with PFFs at one month of age. All

microphotograph data are representative images from one animal from each respective cohort. **B)**

Stereological estimates of pS129[+]-labeled cells in the substantia nigra ipsilateral and contralateral to the

site of PFF injection (N≥8). No α-syn pathology across any stereologically evaluated regions was

observed in PBS-treated animals. **C)** Stereological estimates of TH[+] labeled cells in the substantia nigra in

both WT and En1[+/-] mice injected unilaterally with either PBS or PFFs. **D)** Corresponding densitometric

quantification of TH-immunoreactive striatal integration (N≥5). Results are depicted as mean±SEM.

pS129+ and TH+ cell count data were analyzed by negative binomial mixed-effects regression with

separate analyses performed to compare pathology ipsilateral to injection and contralateral to injection.

Striatal TH[+] optical density data were analyzed by linear mixed-effects regression. Statistics: all *p<0.05*,

***p<0.01*. Scale bar: A - 250µm.

Fig.3 – Enhanced propagation of α-Syn pathology observed in extra-nigrostriatal brain regions of *En1*[+/-]

mice. A) Microphotographs of α-syn pathology in the basolatereal amygdala of WT and *En1*[+/-] animals

unilaterally injected with PFFs. All microphotograph data are representative images from one animal

from each respective male cohort. **B)** Corresponding stereological estimates of pS129[+] labeled cells in

the basolateral amygdala ipsilateral and contralateral to site of PFF injection (N≥8). The second

amygdala graph depicts pS129[+] labeled cells ipsilateral to site of PFF injection separated by sex of

animals. **C)** Stereological estimates of pS129[+] labeled cells in the ventral tegmental area (N≥5) and **D)**

motor cortex (N≥6). Results are depicted as mean±SEM. pS129+ count data were analyzed by negative

binomial mixed-effects regression with separate analyses performed to compare pathology ipsilateral to

injection and contralateral to injection. Statistics: all *p<0.05*, ***p<0.01*. Scale bar: A - 250µm.

Fig.4 – Striatal PFF injections induce widespread cortical α-Syn pathology in *En1*[+/-] mice.

Representative microphotographs of pS129 staining across cortical brain regions in both WT and *En1*[+/-]

mice. α-Syn pathology in cell bodies and neurites is detected across cortical regions with both direct and

indirect projections to/from the striatum. Pathology is observed both ipsilateral and contralateral to PFF

injection. Data are representative images from one respective animal of each cohort. Abbreviations: FrA

– Frontal Association, Orb – Orbital, M1/M2 – Motor, Ect – Ectorhinal, S1/S2 – Somatosensory, PRh –

Perirhinal, Aud – Auditory, Ent – Entorhinal. Scale bar: 500µm.

Fig.5 – Pathological propagation of α-Syn is enhanced throughout *En1*$^{+/-}$ neuraxis. A) Schematic

representation of brain regions depicting sites of observed pS129 cellular and neuritic deposition. **B)**

Heat map representing semi-quantititative pS129-pathological scores across numerous regions in both

WT and *En1*$^{+/-}$ mice injected with striatal PFFs (N=8). Regions were assessed both ipsilateral and

contralateral to PFF injection.

A

Engrailed1$^{+/+}$ | En1 → DA axon outgrowth / CNS Regionality / Morphogenesis & maintenance

LacZ

Engrailed1$^{+/-}$ | LacZ

B

Nigrostriatal Deficit	Striatal DA	▼
	DA Neurons in SN	▼
	DOPAC	▼
Altered Autophagy	p-mTOR	▲
	pS6	▲
	LC3b	▼
Inflammation	Iba1	▲
Mitochondrial Defects	Complex I	▼
Behavior	Motor Function	▼

C

200 nm   hPFF

D

pS129

E

Study Design

Intrastriatal hPFF Injection

Sacrifice
Post-Mortem Histology

Birth — 1 — 2 — 3

Age (Months)

A

Ipsilateral to PFF injection

WT   En1$^{+/-}$

Contralateral to PFF injection

WT   En1$^{+/-}$

B

Amygdala

pS129+ Labeled Cells

Ipsilateral to PFF inj.   Contralateral to PFF inj.

pS129+ Labeled Cells Ipsilateral to PFF Inj.

**

Male   Female

■ WT
■ En1$^{+/-}$

C

Ventral Tegmental Area

pS129+ Labeled Cells

*

Ipsilateral to PFF inj.   Contralateral to PFF inj.

D

Motor Cortex

pS129+ Labeled Cells

Ipsilateral to PFF inj.   Contralateral to PFF inj.

A.

WT   En1$^{+/-}$

hPFF Injection

B.

	Ipsilateral		Contralateral	
	WT	En1$^{+/-}$	WT	En1$^{+/-}$
Frontal association cortex				
Agranular insular cortex				
Lateral orbital cortex				
Temporal association cortex				
Medial orbital cortex				
Prelimbic cortex				
Cingulate cortex				
Motor cortex				
Ventral orbital cortex				
Granular/dysgranular insular cortex				
Ectorhinal cortex				
Somatosensory cortex				
Infralimbic cortex				
Striatum				
Nucleus accumbens				
Globus/Ventral pallidus				
Retrosplenial granular/dysgranular cx				
Basolateral amygdala				
Visual cortex; primary/secondary				
Perirhinal cortex				
Basomedial amygdala				
Piriform cortex				
Hippocampus				
Peduncular lateral hypothalamus				
Auditory cortex				
Parietal association cortex; lateral/medial				
Substantia nigra pars reticulata				
Substantia nigra pars compacta				
Ventral tegmental area, rostral				
Entorhinal cortex; Dorsal/Caudomedial				
Parafascicular thalamus/ventral posteromed thal				
Dorsal subiculum; near hippo				
Ventral/Intermedial nucleus lateral lemniscus				
Pontine reticular nucleus, oral				
Olfactory bulb				
Reticular nuclei				
Retromamillary nucleus				
Corpus calosum				
Dorsal raphe				
Retromamillary decuss				