Coupling adaptive molecular evolution to phylodynamics using fitness-dependent birth-death models

David A. Rasmussen[1,2*], Tanja Stadler[3,4]

1 Department of Entomology and Plant Pathology, North Carolina State University, Raleigh, North Carolina, USA
2 Bioinformatics Research Center, North Carolina State University, Raleigh, North Carolina, USA
3 Department of Biosystems Science and Engineering, ETH Zürich, Basel, Switzerland
4 Swiss Institute of Bioinformatics, Lausanne, Switzerland

* E-mail: drasmus@ncsu.edu

Abstract

Beneficial and deleterious mutations cause fitness to vary among individuals in a population, which natural selection can then act upon to drive adaptive evolution. Non-neutral mutations can likewise cause fitness to vary among lineages in a phylogeny and thereby shape its branching structure. While standard phylogenetic models do not allow mutations to feedback and shape tree topology, birth-death models can account for this feedback by letting the fitness of lineages vary depending on their type. To date though, these multi-type birth-death models have only been applied to cases where a lineage's fitness is determined by a single evolving character state and have not been extended to model sequence evolution across multiple sites. We introduce an extension of the multi-type birth-death model, the marginal fitness birth-death model, that tracks sequence evolution at multiple sites and how the fitness of a lineage depends on its genotype across all sites. This approach remains computationally tractable even for many evolving sites because it tracks the genotype of a lineage probabilistically in an approximate manner, and then marginalizes over all possible genotypes to determine the expected fitness of a lineage. Although approximate, we show that we can accurately estimate the fitness of a lineage and even site-specific mutational fitness effects from the branching pattern of a phylogeny. To demonstrate the power of this approach, we apply it to estimate the host population level fitness effects of mutations previously identified to increase the infectivity of Ebola virus in human cell lines during the 2013-16 epidemic in West Africa.

Introduction

The fitness effects of new mutations is a key determinant of a population's evolutionary potential to adapt over time. Studies exploring the distribution of fitness effects (DFE) of new mutations in a wide range of organisms have revealed that, while many mutations are neutral, a smaller but significant fraction have substantial effects on fitness [Sanjuán et al., 2004, Eyre-Walker and Keightley, 2007, Visher et al., 2016]. These findings have spurred interest in molecular evolutionary models that consider how non-neutral mutations shape sequence evolution and patterns of genetic diversity. Such models range in complexity from simple models assuming that selection operates uniformly across all sites [Muse and Gaut, 1994, Goldman and Yang, 1994, Yang and Nielsen, 2008] to parameter rich models with site-specific fitness effects [Halpern and Bruno, 1998, Lartillot and Philippe, 2004, Rodrigue et al., 2010]. While all of these models assume sequences evolve along an underlying phylogenetic tree representing their shared common ancestry, all also assume that the mutation process driving sequence evolution is independent of the other evolutionary processes giving rise to the tree. This independence assumption implies that mutations do not feedback and affect the fitness of lineages in the tree—such that lineages carrying highly

beneficial mutations are just as likely to survive and produce sampled descendants as lineages riddled with deleterious mutations.

While questionable in terms of biological realism, independence between the tree generating process and the mutation process allows for tractable statistical models. Assuming independence, the joint likelihood of a phylogenetic tree $\mathcal{T}$ and the sequence data $\mathcal{S}$ at the tips of the tree having evolved as observed can be factored into two distinct components:

$$L(\mathcal{S}, \mathcal{T}|\mu, \theta) = L(\mathcal{S}|\mathcal{T}, \mu)p(\mathcal{T}|\theta). \tag{1}$$

The likelihood of the sequence data $L(\mathcal{S}|\mathcal{T}, \mu)$ conditional on the tree and the mutational parameters μ can be computed efficiently for most continuous-time Markov models of sequence evolution [Felsenstein, 1981]. The probability density $p(\mathcal{T}|\theta)$ of the tree $\mathcal{T}$ given the parameters generating the tree θ can likewise be computed under widely used coalescent [Griffiths and Tavare, 1994, Pybus et al., 2000] or birth-death models [Rannala and Yang, 1996, Stadler, 2009]. In Bayesian phylogenetics, $p(\mathcal{T}|\theta)$ is normally thought of as the prior distribution over trees rather than a likelihood, because the tree itself is inferred from the sequence data.

The assumption of independence between the mutation and tree generating processes may be unproblematic in certain scenarios, such as if mutations are truly neutral or do not contribute to substantial fitness differences among lineages. A common argument invoked in defense of ignoring non-neutral mutations is that macroevoluntionary tree generating processes like speciation and extinction play out on longer timescales than the substitution process fixing or removing mutations within a population [Bustamante, 2005]. In this case, fitness variation drives the substitution process within a population but does not ultimately drive the formation of a phylogeny at the species level. But such separation-of-timescales arguments do not hold when segregating mutations contribute to substantial fitness variation between lineages in a phylogeny, such as for rapidly evolving microbes where several different mutant strains can co-circulate. In these cases, the tree generating and mutation processes occur on the same timescale, and the fitness effects of mutations can feedback and shape the branching structure of a phylogeny [Kaplan et al., 1988, Nicolaisen and Desai, 2012, Neher and Hallatschek, 2013]. Ignoring non-neutral evolution in this case may introduce biases into phylogenetic inference. But perhaps more importantly, fitness differences among lineages can be correlated with their ancestral genotypes, providing information about the molecular basis of adaptive evolution we would otherwise ignore.

We therefore explore an approach that couples molecular sequence evolution to the tree-generating process using multi-type birth-death (MTBD) models. Under this approach, mutations can directly impact the fitness of a lineage in the phylogeny by altering its birth or death rate. For a single evolving site or other character state, the joint likelihood of the phylogeny together with the observed tip states can be computed exactly under the MTBD model [Stadler and Bonhoeffer, 2013, Kühnert et al., 2016]. However, this approach is impractical for more than a few non-neutrally evolving sites due to the need to track all possible genotypes as separate types in the state space of the model. We therefore explore an approximate birth-death model that considers how mutations at multiple sites contribute to a lineage's overall fitness, without the need to track all possible genotypes in sequence space. This approach allows us to infer the fitness effects of individual mutations and the fitness of any particular lineage at any time (based on its inferred ancestral genotype) from the branching structure of a phylogeny. Because our approach is particularly relevant to rapidly adapting microbial pathogens, we apply it to Ebola virus and quantify the host population-level fitness effects of naturally occurring amino acid substitutions previously identified to increase viral infectivity in human cell lines [Diehl et al., 2016, Urbanowicz et al., 2016].

Methods

The MTBD at a single evolving site

At a single evolving site, the multi-type birth-death (MTBD) model of Stadler and Bonhoeffer [2013] can be used to compute the joint likelihood $L(S, \mathcal{T}|\mu, \theta)$ of the sequence or character state data $\mathcal{S}$ and phylogenetic tree $\mathcal{T}$ in a way that couples the mutation process with changes in fitness along a lineage. Let $D_n(t)$ represent the probability density that the subtree descending from lineage n evolved between time t and the present exactly as observed. Further, let $D_{n,i}(t)$ represent this probability density conditional on lineage n being in state i out of M possible states at time t. Here the state of a lineage refers to a particular allele or character state (e.g. nucleotide or amino acid) at a single site. We reserve the term genotype to refer to a particular configuration of states across multiple sites in a sequence.

The density $D_{n,i}(t)$ can be computed by solving a system of ordinary differential equations backwards in time from the present ($t = 0$) to time t:

$$
\begin{aligned}
\frac{d}{dt}D_{n,i}(t) = & -(\lambda_i + \sum_{j=1}^{M}\gamma_{i,j} + d_i)D_{n,i}(t) && \text{(a) no event} \\
& +2\lambda_i E_i(t)D_{n,i}(t) && \text{(b) birth of lineage with no sampled descendants} \\
& +\sum_{j=1}^{M}\gamma_{i,j}D_{n,j}(t) && \text{(c) mutation from } i \text{ to } j.
\end{aligned}
\tag{2}
$$

Here, λ_i is the birth rate and d_i is the death rate of lineages in state i, and thus reflect a lineage's fitness. Mutations between states i and j occur at a rate $\gamma_{i,j}$, independently of birth events. Each term in (2) describes how $D_{n,i}$ changes through time by accounting for all of the different events that could have occurred along the lineage. The first term (a) considers the change in probability density given that no birth, death or mutation event occurred. The second term (b) considers the probability of a birth event that went unobserved because one of the child lineages produced no sampled descendants (this event has probability $E_i(t)$, see below). The third term (c) reflects the probability that the lineage mutated from state i to j.

$E_i(t)$ represents the probability that a lineage in state i is not sampled and has no sampled descendants. This probability can be computed at any time t by solving a second set of ODEs:

$$
\begin{aligned}
\frac{d}{dt}E_i(t) = & (1 - s_i)d_i && \text{(a) death without sampling} \\
& -(\lambda_i + \sum_{j=1}^{M}\gamma_{i,j} + d_i)E_i(t) && \text{(b) no event} \\
& +\lambda_i E_i(t)^2 && \text{(c) birth, neither child has sampled descendants} \\
& +\sum_{j=1}^{M}\gamma_{i,j}E_j(t). && \text{(d) mutation from } i \text{ to } j
\end{aligned}
\tag{3}
$$

The first term (a) represents the probability that a lineage dies and is not sampled, where s_i is the probability that a lineage in state i is sampled upon dying. Terms (b-d) have similar interpretations as in (2).

At a tip lineage n, we initialize $D_{n,i}(t) = ds$ if the lineage was sampled upon death at time t. Alternatively, if n was sampled at the present time $t = 0$ before dying, then $D_{n,i}(t) = \rho$, where ρ is the probability that an individual was sampled. At a branching event, the probability density $D_{a,i}$ of the parent lineage a in state i giving rise to two descendent lineages n and m is updated as:

$$
D_{a,i} = 2\lambda_i D_{m,i}(t)D_{n,i}(t).
\tag{4}
$$

The factor of two enters because either lineage m or n could have given birth and we must consider both possible events.

At the root, we can compute the probability density of the entire tree by summing over all possible root states:

$$D_n = \sum_{i=1}^{M} q_i \frac{D_{n,i}(t_{root})}{1 - E_i(t_{root})}, \tag{5}$$

where q_i is the prior probability that the root is in state i at time t_{root}. D_n represents the probability that the entire tree and the tip states $\mathcal{S}$ evolved as exactly as observed. It is therefore equivalent to the joint likelihood $L(\mathcal{S}, \mathcal{T} | \mu, \theta)$ we seek where $\mu = \{\gamma\}$ and $\theta = \{\lambda, d, s\}$.

In theory, this approach could be extended to evolution at any number of sites as long as we track $D_{n,i}(t)$ for all possible genotypes i. Unfortunately, this approach has limited utility because the number of possible genotypes in sequence space scales exponentially with the number of sites L (i.e. 4^L possible genotypes for nucleotide sequences), making the MTBD model impractical for modeling evolution at more than a few sites.

The marginal fitness birth-death model

While the fitness of a lineage will generally depend on its genotype across multiple sites, tracking evolution in the space of all possible genotypes is, as just discussed, computationally infeasible. We therefore seek an approach that considers how mutations at multiple sites determine the fitness of a lineage without the need to track $D_{n,i}$ for all possible genotypes. In the approach described below, we therefore track molecular evolution at each site, computing the probability that each site occupies each state, and then approximate the probability of a lineage being in any particular genotype based on these site probabilities. To compute the expected fitness of a lineage, we can then sum, or marginalize, over the fitness of each genotype weighted by its approximate probability. We therefore refer to this approach as the marginal fitness birth-death (MFBD) model.

First, in order to couple a lineage's fitness with the birth-death process, we will assume that the birth rate λ_n of any lineage n scales according to the fitness f_g of its genotype:

$$\lambda_n = f_g \lambda_0, \tag{6}$$

where λ_0 is the base birth rate assigned to a particular reference genotype (e.g. the wildtype). A lineage's death rate can also be coupled to its fitness, but for simplicity we will assume a lineage's fitness is reflected only in its birth rate λ_n.

Let $\mathcal{G}$ be the set of all possible genotypes in sequence space and g_k be the state of genotype g at site k. To make it clear when we are considering evolution in genotype space rather than at a particular site, we will write the probability density $D_{n,i}$ as $D_{n,g}$ when i refers to a particular genotype. Furthermore, let $D_{n,k,i}$ be the probability density of the subtree descending from lineage n given that site k is in state i. By definition,

$$D_{n,k,i} = \sum_{\{g \in \mathcal{G} : g_k = i\}} D_{n,g}, \tag{7}$$

where the sum is over all genotypes in $\mathcal{G}$ with site k in state i.

We can derive a difference equation for $D_{n,k,i}$ from $D_{n,g}$ in a straightforward manner:

$$
\begin{aligned}
D_{n,k,i}(t + \Delta t) &= \sum_{\{g \in \mathcal{G}: g_k = i\}} D_{n,g}(t + \Delta t) \\
&= \sum_{\{g \in \mathcal{G}: g_k = i\}} \Big[\big(1 - (f_g \lambda_0 + \sum_{j=1}^{M} \sum_{\{g' \in \mathcal{G}: g'_k = j\}} \gamma_{g,g'} + d)\big) D_{n,g}(t) \Delta t \\
&\quad + 2 f_g \lambda_0 E_{n,g}(t) D_{n,g}(t) \Delta t \\
&\quad + \sum_{j=1}^{M} \sum_{\{g' \in \mathcal{G}: g'_k = j\}} \gamma_{g,g'} D_{n,g'}(t) \Delta t \Big].
\end{aligned}
\tag{8}
$$

Taking the limit as $\Delta t \to 0$, we get a new system of differential equations for $D_{n,k,i}(t)$:

$$
\begin{aligned}
\frac{d}{dt} D_{n,k,i}(t) = \sum_{\{g \in \mathcal{G}: g_k = i\}} \Big[&- (f_g \lambda_0 + \sum_{j=1}^{M} \sum_{\{g' \in \mathcal{G}: g'_k = j\}} \gamma_{g,g'} + d) D_{n,g}(t) \\
&+ 2 f_g \lambda_0 E_{n,g}(t) D_{n,g}(t) \\
&+ \sum_{j=1}^{M} \sum_{\{g' \in \mathcal{G}: g'_k = j\}} \gamma_{g,g'} D_{n,g'}(t) \Big]
\end{aligned}
\tag{9}
$$

Unfortunately, (9) would still require us to track $D_{n,g}(t)$ for all possible genotypes, precisely what we wish not to do. We show below that, if we can approximate f_g and $E_{n,g}$ for any given lineage, we can write (9) in terms of only $D_{n,k,i}$ (see (19)) and therefore do not need to track each genotype.

Approximating the fitness of a lineage

We begin by approximating the fitness f_n of a lineage n. Even if we do not know the exact genotype of a lineage at a particular time, we can compute the lineage's expected fitness by summing over the fitness of each genotype f_g weighted by the probability $\omega_{n,g}$ that lineage n is in genotype g:

$$
\mathbb{E}(f_n) = \sum_{g \in \mathcal{G}} f_g \omega_{n,g}.
\tag{10}
$$

The same logic can be extended to compute the expected marginal fitness $\mathbb{E}(f_{n,k,i})$ of a lineage n that at site k is in state i:

$$
\mathbb{E}(f_{n,k,i}) = \sum_{\{g \in \mathcal{G}: g_k = i\}} f_g \omega_{n,g}.
\tag{11}
$$

Computing $\mathbb{E}(f_{n,k,i})$ using (11) requires knowledge of the genotype probabilities $\omega_{n,g}$, which would again require us to track evolution in genotype space. We therefore introduce our major assumption: that we can approximate genotype probabilities using only the marginal site probabilities $\omega_{n,k,i}$ that site k is in state i. We describe how we compute $\omega_{n,k,i}$ below. For now, we make the approximation that

$$
\hat{\omega}_{n,g} = \prod_{k=1}^{L} \omega_{n,k,g_k}.
\tag{12}
$$

This approximation assumes that all sites evolve independently of one another, which is not generally true because mutations at different sites are linked together in genotypes with shared ancestral histories, creating correlations among sites that we ignore. Here though, our direct goal is not to compute these genotype probabilities, but to approximate them accurately enough to track the expected fitness of a lineage.

Using the approximate genotype probabilities $\hat{\omega}_{n,g}$, we can in turn approximate the expected marginal fitness of a lineage:

$$\hat{f}_{n,k,i} = \sum_{\{g \in \mathcal{G}: g_k = i\}} f_g \hat{\omega}_{n,g}. \tag{13}$$

If the fitness effects of each site act multiplicatively to determine the overall fitness of a lineage, we can compute $\hat{f}_{n,k,i}$ as:

$$\hat{f}_{n,k,i} = \sigma_{ki} \prod_{l=1, l \neq k}^{L} \sum_{j=1}^{M} \sigma_{lj} \omega_{n,l,j}, \tag{14}$$

where σ_{ki} is the fitness effect of site k being in state i. This formulation of $\hat{f}_{n,k,i}$ is useful if the number of sites L is large and the number of genotypes we need to sum over in (13) is therefore also extremely large.

Approximating the probability of no sampled descendants

The $E_{n,g}(t)$ term in (9) represents the probability that if lineage n gave birth to a new lineage at time t, the daughter lineage was not sampled and left no sampled descendants. $E_{n,g}(t)$ therefore necessarily depends on the fitness of unobserved lineages descending from n and how fitness along these lineages evolves through changes in their genotype. Because it is often easier track evolution in one dimensional fitness space rather than high-dimensional sequence space [Kepler and Perelson, 1993, Tsimring et al., 1996], we simplify this problem by tracking a proxy for $E_{n,g}(t)$ though fitness space.

Let E_u be the probability that a lineage with expected fitness u leaves no sampled descendants. While fitness can take on a continuous range of values, we track these probabilities only for a discrete set of points $\mathcal{V}$ in fitness space. We can track E_u for $u \in \mathcal{V}$ by modifying (3) to obtain:

$$\frac{d}{dt} E_u(t) = (1 - s_u) d_u - (\lambda_u + \sum_{v \in \mathcal{V}} \gamma_{u,v} + d_u) E_u(t) + \lambda_u E_u(t)^2 + \sum_{v \in \mathcal{V}} \gamma_{u,v} E_v(t). \tag{15}$$

We can then substitute $E_{n,g}(t)$ in (9) with E_u for the fitness value u closest to f_g or $\hat{f}_{n,k,i}$ in fitness space.

Tracking evolution in fitness space requires us to specify rates $\gamma_{u,v}$ for how lineages transition between fitness classes u and v. Let $\mathcal{G}_u$ be the set of genotypes with expected fitness closest to u out of all fitness values in $\mathcal{V}$. We approximate $\gamma_{u,v}$ as:

$$\gamma_{u,v} = \frac{1}{|\mathcal{G}_u|} \sum_{i \in \mathcal{G}_u} \sum_{j \in \mathcal{G}_v} \mu_{ij}, \tag{16}$$

where μ_{ij} is the mutation rate between genotypes i and j. In other words, we compute the average rate of transitions out of fitness class u into v by summing over all possible transitions between genotypes contained within each fitness class. Note that if each genotype falls in a unique fitness class such that $|\mathcal{G}_u| = 1$ for all $u \in \mathcal{V}$, then E_u is computed exactly.

Computing the marginal site densities $D_{n,k,i}$

Recall that (9) provided an exact way to track the marginal site densities $D_{n,k,i}$ based on the genotype densities $D_{n,g}$. To efficiently evaluate $D_{n,k,i}$ without the need to track $D_{n,g}$ for all genotypes, we apply

the three approximations made above. First, we approximate the genotype probabilities $\hat{\omega}_{n,g}$ based on the marginal site probabilities. Second, we marginalize over the fitness of each genotype (weighted by its genotype probability) to compute $\hat{f}_{n,k,i}$ and then substitute $\hat{f}_{n,k,i}$ for f_g for all genotypes where $g_k = i$ below. Third, we approximate $E_{n,g}$ by E_u for a single fitness value u closest to $\hat{f}_{n,k,i}$. Making these approximations in (9) leads to:

$$
\begin{aligned}
\frac{d}{dt} D_{n,k,i}(t) = \sum_{\{g \in \mathcal{G}: g_k = i\}} & \Big[- (\hat{f}_{n,k,i} \lambda_0 + \sum_{j=1}^{M} \sum_{\{g' \in \mathcal{G}: g'_k = j\}} \gamma_{g,g'} + d) D_{n,g}(t) \\
& + 2\hat{f}_{n,k,i} \lambda_0 E_u(t) D_{n,g}(t) \\
& + \sum_{j=1}^{M} \sum_{\{g' \in \mathcal{G}: g'_k = j\}} \gamma_{g,g'} D_{n,g'}(t) \Big]
\end{aligned}
\tag{17}
$$

Assuming that the mutation rate from i to j at site k does not depend on the genetic background, we can substitute $\sum_{j=1}^{M} \sum_{\{g' \in \mathcal{G}: g'_k = j\}} \gamma_{g,g'}$ with $\sum_{j=1}^{M} \gamma_{i,j}$, where $\gamma_{i,j}$ is the per site mutation rate. We can likewise substitute $\sum_{j=1}^{M} \sum_{\{g' \in \mathcal{G}: g'_k = j\}} \gamma_{g,g'} D_{n,g'}(t)$ with $\sum_{j=1}^{M} \gamma_{i,j} \sum_{\{g' \in \mathcal{G}: g'_k = j\}} D_{n,g'}(t)$. Making these substitutions and rearranging the sums in (17), we have:

$$
\begin{aligned}
\frac{d}{dt} D_{n,k,i}(t) = & - (\hat{f}_{n,k,i} \lambda_0 + \sum_{j=1}^{M} \gamma_{i,j} + d) \sum_{\{g \in \mathcal{G}: g_k = i\}} D_{n,g}(t) \\
& + 2\hat{f}_{n,k,i} \lambda_0 E_u(t) \sum_{\{g \in \mathcal{G}: g_k = i\}} D_{n,g}(t) \\
& + \sum_{j=1}^{M} \gamma_{i,j} \sum_{\{g' \in \mathcal{G}: g'_k = j\}} D_{n,g'}
\end{aligned}
\tag{18}
$$

Recalling that $D_{n,k,i} = \sum_{\{g \in \mathcal{G}: g_k = i\}} D_{n,g}$ (and by extension $D_{n,k,j} = \sum_{\{g \in \mathcal{G}: g_k = j\}} D_{n,g}$), we have:

$$
\begin{aligned}
\frac{d}{dt} D_{n,k,i}(t) = & - \left(\hat{f}_{n,k,i} \lambda_0 + \sum_{j=1}^{M} \gamma_{i,j} + d \right) D_{n,k,i}(t) \\
& + 2\hat{f}_{n,k,i} \lambda_0 E_u(t) D_{n,k,i}(t) \\
& + \sum_{j=1}^{M} \gamma_{i,j} D_{n,k,j}(t).
\end{aligned}
\tag{19}
$$

The significance of (19) is twofold. First, we can track sequence evolution at each site individually without tracking all genotypes. Second, given $\hat{f}_{n,k,i}$, we can track the overall fitness of a lineage by marginalizing over the fitness effects of all possible mutations at other sites. We can therefore track sequence evolution at each site while simultaneously taking into account the coupled fitness effects of mutations at all other sites on a lineage's fitness.

Computing $\hat{f}_{n,k,i}$ however still requires us to approximate the genotype probabilities using (12), which in turn requires the marginal site probabilities $\omega_{n,k,i}$. In our notation, $\omega_{n,k,i}$ represents the conditional probability $p(i|\mathcal{T}_n, \mathcal{S}_n)$ that lineage n is in particular state i, where $\mathcal{T}_n$ represents the subtree descending from n with tip sequences $\mathcal{S}_n$. $D_{n,k,i}$ represents the inverse conditional probability density $p(\mathcal{T}_n, \mathcal{S}_n|i)$. We can therefore apply Bayes theorem to compute $\omega_{n,k,i}$ given $D_{n,k,i}$:

$$
\omega_{n,k,i} = p(i|\mathcal{T}_n, \mathcal{S}_n) = \frac{p(\mathcal{T}_n, \mathcal{S}_n|i) q(i)}{\sum_{i}^{M} p(\mathcal{T}_n, \mathcal{S}_n|i) q(i)} = \frac{D_{n,k,i} q(i)}{\sum_{i}^{M} D_{n,k,i} q(i)}.
\tag{20}
$$

The $q(i)$ terms represent the prior probability that the lineage is in state i. Here we make a simplification in assuming that the tree ancestral and sister to lineage n has no information regarding $\omega_{n,k,i}$, and thus assume a uniform prior on $q(i) = 1/M$. The $q(i)$ terms therefore cancel above.

Because the fitness of a lineage depends on the state of all sites, we must solve (19) for all sites simultaneously as one coupled system of differential equations. This requires updating $D_{n,k,i}$ at each time step, which suggests the following iterative procedure.

At a tip n observed to be in genotype g, we initialize $\hat{f}_{n,k,i}$ as f_g if $g_k = i$ or else $\hat{f}_{n,k,i} = 0$, $\hat{D}_{n,k,i} = ds$ or ρ, and $\omega_{n,k,i} = 1$ if $g_k = i$, else $\omega_{n,k,i} = 0$. Then at each time step backwards through time from time t to time $t + \Delta t$, for each site and state we:

1. Update $D_{n,k,i}$ by numerically integrating (19) over time step Δt.

2. Update the marginal site probabilities $\omega_{n,k,i}$ using (20)

3. Update the expected marginal fitness values $\hat{f}_{n,k,i}$ using (13) or (14).

Computing the full joint likelihood

We can now compute the joint likelihood of the tree and sequence data if we track $D_{n,k,i}$ at each site back to the root. At the root, $D_{n,k,i}(t_{root})$ represents $p(\mathcal{T}, \mathcal{S}_k | \mu, \theta, i)$, the probability density of the entire tree $\mathcal{T}$ and the observed sequence data $\mathcal{S}_k$ as site k, conditional on site k being in state i at the root. To be precise, $D_{n,k,i}$ only approximates $p(\mathcal{T}, \mathcal{S}_k | \mu, \theta, i)$ because we computed $D_{n,k,i}$ using the expected marginal fitness of a lineage $\hat{f}_{n,k,i}$ based on approximate genotype probabilities. We therefore introduce an additional auxiliary variable $\mathcal{F}$ representing the entire set of expected fitness values $\hat{f}_{n,k,i}$ computed over all lineages, sites and states. Using this notation, $D_{n,k,i}(t_{root}) = p(\mathcal{T}, \mathcal{S}_k | \mu, \theta, \mathcal{F}, i)$. By summing over all possible root states at site k (and conditioning on survival), we can then compute:

$$p(\mathcal{T}, \mathcal{S}_k | \mu, \theta, \mathcal{F}) = \sum_{i=1}^{M} q(i) \frac{p(\mathcal{T}, \mathcal{S}_k | \mu, \theta, \mathcal{F}, i)}{1 - E_u(t_{root})} = \sum_{i=1}^{M} q(i) \frac{D_{n,k,i}(t_{root})}{1 - E_{n,k,i}(t_{root})}. \tag{21}$$

Likewise, we can compute the conditional probability density $p(\mathcal{S}_k | \mathcal{T}, \mu, \theta, \mathcal{F})$ of the sequence data at site k given the tree:

$$p(\mathcal{S}_k | \mathcal{T}, \mu, \theta, \mathcal{F}) = \frac{p(\mathcal{T}, \mathcal{S}_k | \mu, \theta, \mathcal{F})}{p(\mathcal{T} | \mu, \theta, \mathcal{F})}. \tag{22}$$

We already know $p(\mathcal{T}, \mathcal{S}_k | \mu, \theta, \mathcal{F})$ from above but now need the tree density $p(\mathcal{T} | \mu, \theta, \mathcal{F})$. This can easily be computed using a birth-death process where the birth rate of each lineage at any time t is always rescaled by its expected fitness $\hat{f}_n(t)$ contained within $\mathcal{F}$.

We can now compute the joint density $p(\mathcal{T}, \mathcal{S}_{1:L} | \mu, \theta)$ for all sites. Because each site is conditionally independent of all other sites given $\mathcal{F}$, we can factor $p(\mathcal{T}, \mathcal{S}_{1:L} | \mu, \theta, \mathcal{F})$ into a product of densities for $\mathcal{S}_k$ at each site and the density of the entire tree $\mathcal{T}$:

$$p(\mathcal{T}, \mathcal{S}_{1:L} | \mu, \theta, \mathcal{F}) = p(\mathcal{T} | \mu, \theta, \mathcal{F}) \prod_{k=1}^{L} p(\mathcal{S}_k | \mathcal{T}, \mu, \theta, \mathcal{F}). \tag{23}$$

We can thus approximate the joint likelihood of the sequence data and the phylogeny $p(\mathcal{T}, \mathcal{S}_{1:L} | \mu, \theta)$ as $p(\mathcal{T}, \mathcal{S}_{1:L} | \mu, \theta, \mathcal{F})$. This allows us to consider how selection shapes sequence evolution at each site while simultaneously considering how the fitness effects of mutations at multiple sites act together to shape the phylogeny. As (23) makes clear though, the goodness of our approximation depends on how well the fitness values in $\mathcal{F}$ are approximated, which in turn depends on how well we can approximate genotypes based on the marginal site probabilities. We explore the goodness of these approximations in the Results section.

Implementation

We first implemented the marginal fitness birth-death (MFBD) model in Matlab version R2017b. The Matlab implementation was used to test how well the MFBD model can approximate likelihoods and genotype probabilities relative to the exact multi-type birth death model tracking all possible genotypes for a simple model with only four genotypes. For statistical inference, the MFBD was implemented as an add-on package for BEAST 2 [Bouckaert et al., 2014] named Lumière, which extends the existing BDMM package for multi-type birth-death models [Kühnert et al., 2016]. The BEAST 2 implementation of Lumière allows for the joint posterior distribution of all model parameters and the phylogeny to be estimated from sequence data using Bayesian MCMC. Source code for Lumière and the Matlab implementation are freely available at `https://github.com/davidrasm/Lumiere`.

Simulations

To test the statistical performance of our approach, mock phylogenies and sequence data were simulated under a birth-death-mutation-sampling process using a variant of the Gillespie stochastic simulation algorithm [Gillespie, 2007] that recorded the ancestry of all individuals in the population. A binary sequence was associated with each lineage and allowed to mutate with a constant per-site mutation rate γ. Mutations could alter the fitness of a lineage by either increasing or decreasing its birth rate according to site-specific fitness effects. At death events, lineages were sampled with probability s, in which case they were included in the mock phylogeny. Code for these simulations is available at `https://github.com/davidrasm/Lumiere/sim/`.

Ebola analysis

We used the Lumière implementation of the MFBD model to estimate the fitness effects of amino acid mutations previously identified to increase the infectivity of Ebola virus in human cell lines [Diehl et al., 2016, Urbanowicz et al., 2016]. We reanalyzed a set of 1610 whole genome EBOV sequences sampled from Guinea, Sierra Leone and Liberia in 2014 to 2016. The sequence alignment along with the time-calibrated molecular phylogeny we used for our analysis were downloaded from `https://github.com/ebov/space-time/tree/master/Data`.

Urbanowicz et al. [2016] measured the fitness effects of 17 viral genotypes carrying 18 different amino acid mutations in either single, double or triple mutant backgrounds relative to the Makona genotype first sampled at the beginning of the epidemic. Because our methods cannot estimate fitness effects of mutations at very low frequencies, we only analyzed 9 of these mutations that were present in at least 10 of the 1610 viral samples. Preliminary analysis revealed that these mutations fall within 8 unique genetic backgrounds because of the way mutations are nested within other single or double mutant lineages in the phylogeny. Because the data of Urbanowicz et al. [2016] strongly suggest that epistatic interactions between mutations affect viral fitness, we estimated the genotypic fitness f_g of these 8 major genotypes rather than site-specific fitness effects σ. We therefore used the MFBD to track sequence evolution at each site, but used (13) to marginalize over these genotypes when approximating the fitness of a lineage.

We estimated the fitness of each genotype relative to the Makona genotype, assuming a uniform $[0, 2]$ prior distribution on these fitness values. For the other parameters in the model, we assumed a fixed death or removal rate d of 0.1667 per day based on earlier estimates [Gire et al., 2014, Stadler et al., 2014]. Sampling was modeled as occurring upon removal, with the sampling proportion s set to zero before March 2014, when the first sample was collected. After March 2014, we assumed a fixed sampling proportion of 0.056, reflecting the fact that the dataset included samples from 1610 individuals out of the 28,652 probable cases reported by the WHO [WHO, 2016]. Lastly, we assumed a constant amino acid mutation rate over all sites with an exponential prior on both the forward and backward mutation

rate with a mean rate of 2×10^{-3} per site per year. Our analysis of the Ebola data can be reproduced in Lumière with the XML input file available at `https://github.com/davidrasm/Lumiere/ebola/`.

Results

The four genotype model

We consider a simple model of molecular evolution in order to compare the marginal fitness birth-death (MFBD) model against the exact multi-type birth-death (MTBD) model tracking all genotypes. Specifically, we consider a binary evolving sequence of length $L = 2$ where all mutations are deleterious and carry a selective fitness cost σ. It is therefore possible to track the evolutionary dynamics of all four possible genotypes ($\mathcal{G} = \{00, 01, 10, 11\}$) under both models.

We first assume that the fitness effects of individual mutations act multiplicatively, such that the double mutant has fitness $(1 - \sigma)^2$. Figure 1A shows a phylogeny simulated under the four genotype model, colored according to the genotype of each lineage. We computed the joint likelihood that this tree and observed tip genotypes evolved under a range of different fitness values σ for both the exact MTBD and approximate MFBD models (Figure 1B). The likelihood profiles under both models peak around the true value of σ and closely match at lower values of σ, but begin to diverge at higher values. We reasoned that this divergence occurs because the approximate MFBD model does not directly track genotype probabilities, but only approximates them based on the marginal site probabilities. To test this idea, we computed another likelihood profile for a MFBD model where the exact genotype probabilities were provided by the MTBD model. This did not seem to reduce the divergence from the likelihoods computed under the exact model (Figure 1B, circles). We also compared the approximate genotype probabilities against the exact genotype probabilities. The approximate genotypes probabilities closely track the exact genotype probabilities when selection is weak to moderately strong ($\sigma = 0.5$; Figure 1C), but diverge as we increase σ and thereby the strength of purifying selection against the mutant genotypes (Figure 1D).

Overall though, the MFBD only introduces a small amount of error in approximating genotype probabilities, especially when we can compare the MFBD model against another approximation that tracks sequence evolution at each site completely independent of all other sites by setting the expected marginal fitness $\hat{f}_{n,k,i} = \sigma$ instead of using (14). Thus, this approximations completely ignores how the fitness of a lineage depends on mutations at other sites. Under this more naive approximation, the error in the genotype probabilities grows much more rapidly than under the MFBD approximation (Figure 1D).

Furthermore, the approximations made in the MFBD model turn out to not depend on the exact genetic architecture of fitness, i.e. how genotypes map to fitness. We can, for example, consider a variant of the four genotype model where each of the single mutant genotypes is neutral with $\sigma = 0$ but an epistatic interaction between the two sites causes the double mutant to be deleterious with some fitness cost ϵ. In effect, this situation represents a worst case scenario for the MFBD model because fitness depends entirely on the genotype of a lineage rather than on individual sites. For a phylogeny simulated under this model (Figure 1E), the likelihood curves largely agree but diverge as ϵ increases and selection against the double mutant becomes stronger (Figure 1F).

In the model with epistasis, the divergence in the likelihoods computed under the approximate and exact models clearly seems to arise from the error introduced by approximating genotypes probabilities based on the marginal site probabilities. If we condition on the true genotype probabilities, the likelihood profiles computed under the MFBD model closely match the exact MTBD model (Figure 1F, circles). As the strength of selection against the double mutant increases, so does the error introduced by approximating the genotype probabilities using the marginal site probabilities (Figure 1G). This occurs because selection acts at the level of genotypes, which induces additional correlations between sites that causes the genotype probabilities to deviate from those expected based on the marginal site probabilities.

Figure 1. Performance of the MFBD approximation for a model with four genotypes composed of two evolving sites. Mutations have multiplicative fitness effects in A-D and epistatic fitness effects in E-H. **(A & E)** Simulated phylogenies showing the genotype of each lineage through time. **(B & F)** Joint likelihood of the phylogeny and tip genotypes under different values of σ or ϵ computed using the exact MTBD model (solid line), the approximate MFBD (dashed line) and the MFBD where we have conditioned on the true genotype probabilities computed under the exact MTBD model (open circles). The vertical blue line marks the true parameter values. **(C & G)** The normalized probability of a single hypothetical lineage being in each genotype based on the MFBD approximation (solid line) versus the exact MTBD model (dotted line) with $\sigma = 0.5$ or $\epsilon = 0.5$. Note that the probabilities for genotypes 10 and 01 are identical. **(D & H)** The mean, time-integrated error in genotype probabilities introduced by the MFBD approximation with fitness effects coupled across sites (solid line) versus a naive approximation that ignores the fitness effects of other sites (dashed). All parameters besides σ and ϵ were fixed at $\lambda = 0.25$, $d = 0.05$, $s = 0.05$. The mutation rate γ was symmetric between forward and backwards mutations and fixed at 0.05.

Notably though, the error introduced by the MFBD approximation is still small and grows slowly with ϵ, whereas the error grows much more rapidly under the more naive approximation with fitness effects unlinked across sites (Figure 1H).

Estimating site-specific fitness effects

Next, we simulated phylogenies under a model where the fitness effect of the mutant allele at each site is drawn independently from a distribution of fitness effects (DFE) in order to test how well we can estimate site-specific fitness effects. Because there can be considerable uncertainty surrounding these fitness effects, we now estimate the posterior distribution of fitness effects using Bayesian MCMC. The accuracy and precision of the estimated fitness effects varies considerably across sites, as shown for a representative phylogeny with five evolving sites in Figure 2.

In order to better understand this variability, we simulated 100 phylogenies with randomly drawn fitness effects at either 2, 5 or 10 evolving sites. Overall, the estimated posterior median fitness effects are well correlated with their true values, although the strength of this correlation decreases as the number of sites increases (Figure 3A-C). Coverage of the 95% credible intervals on the other hand increased from 71.0 to 72.8 to 77.4%.

Figure 2. (A) A phylogeny simulated under a model with five evolving sites each with a random fitness effect. The lineages are colored according to the number of mutations they carry (blue = 0; yellow = 5). The distribution of fitness effects was assumed to be LogNormal with a mean of 0.85 and a standard deviation of 0.32. **(B)** Site-specific fitness effects estimated using the marginal fitness BD model. Red lines indicate the posterior median and 95% credible intervals. Blue lines mark the true fitness effect at each site.

While there is no systematic directional bias, fitness effects are underestimated for sites at which the mutant allele is at low frequency among sampled individuals and overestimated for sites where the mutant allele is at high frequencies. This however appears to be an intrinsic feature of estimating fitness effects from the branching structure of a phylogeny, as the same phenomena is observed under the exact MTBD model with two sites and 4 genotypes (Figure 3D), and the estimates made under the approximate MFBD

model are highly correlated with estimates made under the exact MTBD model (Figure 3E).

Figure 3. Inference of site-specific fitness effects from simulated phylogenies. (A-C)
Correlation between the true and estimated posterior median fitness effects for phylogenies simulated
with 2, 5 or 10 evolving sites. Results are aggregated over 100 simulated phylogenies, with each point
representing an estimate for a single site and phylogeny. The points are colored according to the
frequency of the mutant allele among sampled individuals in the phylogeny. **(D)** Fitness effects
estimated under the exact MTBD model tracking all four possible genotypes for the same two site
simulations as in A. **(E)** Correlation between the site-specific fitness effects estimated under the
approximate MFBD and exact MTBD for the two site simulations. **(F)** Error and uncertainty in
estimated site-specific fitness effects across all 2, 5, and 10 site simulations. Error was calculated as the
posterior median estimate minus the true fitness effect. Uncertainty was calculated as the standard
deviation of the posterior values sampled via MCMC. In all simulations, sites where the Effective
Sample Size of the MCMC samples was below 100 (less than 5% of all sites across simulations) were
discarded. The death rate was fixed at $d = 0.05$ but the birth, mutation and sampling rates were
randomly drawn for each simulation from a prior distribution: $\lambda \backsim$ Uniform(0.1,0.2); $\gamma \backsim$
Exponential(0.01); $s \backsim$ Uniform(0,1). Only the birth rate was jointly inferred with the site-specific
fitness effects.

Across all sites and simulations, accuracy decreased when the mutant allele at a given site was at low
or high frequencies, and there was considerably more uncertainty for sites where the mutant allele was
at very low frequencies (Figure 3F). Thus, while the MFBD model generally performs well at estimating
site-specific fitness effects, the accuracy and precision of these estimates varies greatly depending on the
frequency of a given mutation in a phylogeny.

Ebola virus adaptation to humans

The Ebola virus glycoprotein (GP) binds to cells during viral cell entry and is therefore thought to be
a key determinant of viral fitness in different hosts. Previously, Urbanowicz et al. [2016] analyzed a

large set of naturally occurring amino acid mutations in the GP isolated from patients during the 2013-16 epidemic in Western Africa. The effect of these GP mutations on fitness were then experimentally determined using infectivity assays in cell culture. Several mutant genotypes dramatically increased viral infectivity relative to the Makona genotype isolated during the earliest stages of the epidemic. However, the effect of these mutations on viral transmission and fitness at the host population level have not yet been determined. We therefore applied the MFBD model to a large dataset of 1610 Ebola virus (EBOV) genomes sampled during the 2013-16 epidemic to infer the population-level fitness effects of these GP mutations.

We analyzed 9 out of the 18 amino acid mutations analyzed by Urbanowicz et al. [2016] that were present in at least 10 of the 1610 viral samples. These 9 mutations fall in 8 different genetic backgrounds or genotypes (Figure 4). Because Urbanowicz et al. [2016] found evidence for epistatic interactions between several of these mutations, we estimated the fitness of these 8 genotypes rather than site-specific mutational fitness effects. Table 1 shows the relative fitness of these genotypes estimated at the population-level versus their fitness in cell culture.

Table 1. Estimated fitness of Ebola GP mutants relative to the Makona genotype

Genotype	Posterior median	95% CI	Effect in cell culture
Makona	1.00	-	Reference genotype
A82V	1.05	1.04-1.07	Increases infectivity 2X
P330S	0.98	0.82-1.14	Decreases infectivity
P330S+N107D+G480D	1.04	0.98-1.12	Increases infectivity >2X
A82V+R410S	1.09	1.00-1.18	No or small effect
A82V+R410S+K439E	1.14	1.01-1.26	Increases infectivity 2-3X
A82V+R29K	1.06	0.93-1.19	Increases infectivity 2-3X
A82V+T230A	1.03	0.93-1.11	Increases infectivity 2-3X
A82V+I371V	1.03	0.98-1.09	Increases infectivity 2-3X

Mapping the genotypes and fitness of lineages inferred under the MFBD model onto the phylogeny allows us to reconstruct the series of events by which EBOV adapted to humans (Figure 4). Shortly after the epidemic started in 2013, the A82V mutation occurred and gave rise to lineage B, which then spread to Sierra Leone, Liberia and Mali. Urbanowicz et al. [2016] found that the A82V mutation increases infectivity by 2-3 fold in cell culture. At the population-level, this mutation appears to have a less dramatic effect, increasing transmissibility by only 5% relative to the Makona genotype. The P330S mutation appears to have temporarily decreased the fitness of the main surviving clade in lineage A, although mutations N107D and G480D later rescue the fitness of this lineage, consistent with the findings of Urbanowicz et al. [2016]. Meanwhile, the R410S mutation occurred within lineage B but did not have an immediate effect on fitness. However, R410S appears to epistatically interact with mutation K439E, which occurs twice along the same lineage carrying the R410S mutation and in this genetic background increases infectivity 2-3 fold in cell culture. We estimate that the A82V+R410S+K439E genotype had the highest population-level fitness, but only increased fitness by 14% relative to the Makona genotype. Three other mutations, R29K, T230A and I371V, also occurred in the A82V genetic background, but were not estimated to have further increased the fitness of the A82V genotype. Overall, the population level fitness of all 8 genotypes agree with their fitness in cell culture in terms of the sign or direction of their effects, but these genotypes had much greater fitness relative to the Makona genotype in cell culture than at the population level.

Discussion

Many assumptions are made in phylogenetics to model molecular evolution in a statistically tractable way. Historically, one of the most biologically questionable yet pervasive of these assumptions is that sequences evolve neutrally along lineages, such the mutations do not feedback and alter the branching process shaping the phylogeny. Our marginal fitness birth-death (MFBD) model allows us to relax this core assumption in order to consider how non-neutral evolution at multiple sites affects sequence evolution, the fitness of lineages, and the overall branching structure of a phylogeny. While our approach is not exact in that it approximates genotype probabilities by assuming sites evolve independently when computing the marginal fitness of a lineage, we have shown that this approximation generally works well and only produces significant errors in rather extreme situations, such as the four genotype model with strong epistasis. While an earlier approach based on birth-death models allowed for lineage-specific fitness values to be inferred from the branching pattern of a phylogeny [Neher et al., 2014], this approach did not connect fitness back to the mutational process nor allow for the fitness effects of individual mutations or genotypes to be estimated. Using our approach, we demonstrated that the fitness effects of specific mutations can be estimated from simulated phylogenies under the MFBD with accuracy comparable to an exact multi-type birth death model. The MFBD model therefore provides a new, statistically powerful way of incorporating adaptive molecular evolution into phylodynamics.

The MFBD model allows us to exploit phylogenetic information about adaptive evolution that most methods for inferring selection from patterns in sequence data ignore. Currently, codon-substation models [Goldman and Yang, 1994, Muse and Gaut, 1994] and the related class of mutation-selection models [Yang and Nielsen, 2008] are by far the most widely used approach for inferring selection. These approaches rely on comparing sequence substitution patterns such as the dN/dS ratio of non-synonymous to synonymous substitutions across sites. These approaches can be very powerful when sequences from highly divergent taxa are available, such that enough time has elapsed for multiple substitutions at a single site to have accumulated between lineages. But on the shorter timescales relevant to evolution within a population, substitution patterns like the dN/dS ratio are relatively insensitive to selection pressures and may produce misleading inferences of selection [Kryazhimskiy and Plotkin, 2008]. For example, a highly beneficial non-synonymous mutation that occurs in a single lineage and then spreads through a population may produce a very low dN/dS ratio, indicative of purifying selection rather than adaptive evolution. In contrast, comparing the evolutionary dynamics of lineages with and without the mutation allows us to infer if that mutation confers a competitive advantage. Thus, considering the branching pattern of phylogenies provides additional information about molecular evolution not visible from substitution patterns in sequence data alone.

While new technologies increasingly allow researchers to quantify mutational fitness effects in vitro or even in vivo [Zanini and Neher, 2013, Thyagarajan and Bloom, 2014], how fitness measured in the lab translates to fitness in nature is largely unknown. This is especially pertinent for emerging pathogens whose epidemic potential often depends on new adaptive mutations [Antia et al., 2003, Longdon et al., 2014]. Phylodynamic approaches like the MFBD model that can quantify fitness at the host population level are therefore greatly needed, as they offer a means to assess the epidemiological significance of mutant lineages. Extrapolating from our experience with Ebola, where the population-level fitness effects of each mutant genotype we considered matched the sign of their effect in cell culture, we suspect that fitness measured in the lab will generally agree with fitness in nature. This seems reasonable, as mutations that increase replication or infectivity within hosts should generally promote transmissibility between hosts (e.g. Quinn et al. [2000], Fraser et al. [2007]). But at the same time, there is no reason to believe that transmission rates will increase linearly or even monotonically with increasing within-host growth rates. We therefore expect that the magnitude of fitness effects might often greatly differ across scales, as we found for the A82V glycoprotein mutation in Ebola. While A82V doubles infectivity in cell culture, we estimated that it only increases transmissibility at the population level by 5% (95% CI: 4-7%). Interestingly, Diehl et al. [2016] found that A82V only slightly increases viral titers in Ebola patients,

which is likely a much better proxy for transmissibility than cellular infectivity, lending support to our more moderate estimates at the population level.

In future applications, the MFBD could be applied to consider a much larger number of evolving sites since there is no need to track evolution in genotype space and computation time scales linearly with the number of sites. Based on our experience, tens or even hundreds of sites may be computationally feasible, depending on the number of sequences. However, with more sites the genetic background in which mutations occur grows in complexity due to the increased probability of mutations being linked to other mutations rather than occurring in isolated genetic backgrounds. This leads to strong correlations between fitness parameters at different sites in an increasingly high dimensional parameter space, making statistical inference challenging, especially using MCMC. Even worse, if two mutations always co-occur in the same genetic background, their individual effects on fitness will not be mutually identifiable. It may therefore not be possible to disentangle causative mutations from neutral passenger mutations. Spurious correlations may also arise due to additional sources of fitness variation beyond sequence changes. For example, if a mutation occurs along a lineage spreading through a higher fitness environment by chance, it will likely be inferred to increase fitness even if it is actually neutral. Thus, while the fitness of different lineages can be estimated under the MFBD, the fitness effects of individual mutations need to be interpreted carefully unless they occur in multiple genetic backgrounds and confounding sources of fitness variation are accounted for.

In spite of these shortcomings, we believe the MFBD model offers a powerful means to explore many questions not previously possible with strictly neutral phylodynamic models. Even if the fitness effects of individual mutations are not identifiable, it may still be possible to infer the distribution of fitness effects across sites, a key determinant of adaptive evolution that has only been explored in a few systems [Eyre-Walker and Keightley, 2007]. The MFBD model can also be used to compare the fitness of a mutation or lineage across different environments, such as in different hosts of a pathogen. Finally, the MFBD is not limited to exploring sequence evolution, as the model is generalizable to any discrete character state, including phenotypic or environmental characters. Thus, our model can be used to explore how multiple molecular and non-molecular characters interact to shape the overall fitness of lineages in a phylogeny.

Acknowledgments

We would like to thank Denise Kühnert for helpful advice on how to implement the MFBD model in BEAST 2.

References

R. Antia, R. R. Regoes, J. C. Koella, and C. T. Bergstrom. The role of evolution in the emergence of infectious diseases. *Nature*, 426(6967):658, 2003.

R. Bouckaert, J. Heled, D. Kühnert, T. Vaughan, C.-H. Wu, D. Xie, M. A. Suchard, A. Rambaut, and A. J. Drummond. BEAST 2: a software platform for Bayesian evolutionary analysis. *PLoS Computational Biology*, 10(4):e1003537, 2014.

C. D. Bustamante. *Population genetics of molecular evolution*. Springer, 2005.

W. E. Diehl, A. E. Lin, N. D. Grubaugh, L. M. Carvalho, K. Kim, P. P. Kyawe, S. M. McCauley, E. Donnard, A. Kucukural, P. McDonel, et al. Ebola virus glycoprotein with increased infectivity dominated the 2013–2016 epidemic. *Cell*, 167(4):1088–1098, 2016.

A. Eyre-Walker and P. D. Keightley. The distribution of fitness effects of new mutations. *Nature Reviews Genetics*, 8(8):610, 2007.

J. Felsenstein. Evolutionary trees from DNA sequences: a maximum likelihood approach. *Journal of Molecular Evolution*, 17(6):368–376, 1981.

C. Fraser, T. D. Hollingsworth, R. Chapman, F. de Wolf, and W. P. Hanage. Variation in HIV-1 set-point viral load: epidemiological analysis and an evolutionary hypothesis. *Proceedings of the National Academy of Sciences*, 104(44):17441–17446, 2007.

D. T. Gillespie. Stochastic simulation of chemical kinetics. *Annu. Rev. Phys. Chem.*, 58:35–55, 2007.

S. K. Gire, A. Goba, K. G. Andersen, R. S. Sealfon, D. J. Park, L. Kanneh, S. Jalloh, M. Momoh, M. Fullah, G. Dudas, et al. Genomic surveillance elucidates Ebola virus origin and transmission during the 2014 outbreak. *Science*, page 1259657, 2014.

N. Goldman and Z. Yang. A codon-based model of nucleotide substitution for protein-coding DNA sequences. *Molecular Biology and Evolution*, 11(5):725–736, 1994.

R. C. Griffiths and S. Tavare. Sampling theory for neutral alleles in a varying environment. *Phil. Trans. R. Soc. Lond. B*, 344(1310):403–410, 1994.

A. L. Halpern and W. J. Bruno. Evolutionary distances for protein-coding sequences: modeling site-specific residue frequencies. *Molecular Biology and Evolution*, 15(7):910–917, 1998.

N. L. Kaplan, T. Darden, and R. R. Hudson. The coalescent process in models with selection. *Genetics*, 120(3):819–829, 1988.

T. B. Kepler and A. S. Perelson. Cyclic re-entry of germinal center B cells and the efficiency of affinity maturation. *Immunology today*, 14(8):412–415, 1993.

S. Kryazhimskiy and J. B. Plotkin. The population genetics of dN/dS. *PLoS Genetics*, 4(12):e1000304, 2008.

D. Kühnert, T. Stadler, T. G. Vaughan, and A. J. Drummond. Phylodynamics with migration: a computational framework to quantify population structure from genomic data. *Molecular Biology and Evolution*, 33(8):2102–2116, 2016.

N. Lartillot and H. Philippe. A Bayesian mixture model for across-site heterogeneities in the amino-acid replacement process. *Molecular Biology and Evolution*, 21(6):1095–1109, 2004.

B. Longdon, M. A. Brockhurst, C. A. Russell, J. J. Welch, and F. M. Jiggins. The evolution and genetics of virus host shifts. *PLoS Pathogens*, 10(11):e1004395, 2014.

S. V. Muse and B. S. Gaut. A likelihood approach for comparing synonymous and nonsynonymous nucleotide substitution rates, with application to the chloroplast genome. *Molecular Biology and Evolution*, 11(5):715–724, 1994.

R. A. Neher and O. Hallatschek. Genealogies of rapidly adapting populations. *Proceedings of the National Academy of Sciences*, 110(2):437–442, 2013.

R. A. Neher, C. A. Russell, and B. I. Shraiman. Predicting evolution from the shape of genealogical trees. *Elife*, 3:e03568, 2014.

L. E. Nicolaisen and M. M. Desai. Distortions in genealogies due to purifying selection. *Molecular Biology and Evolution*, 29(11):3589–3600, 2012.

O. G. Pybus, A. Rambaut, and P. H. Harvey. An integrated framework for the inference of viral population history from reconstructed genealogies. *Genetics*, 155(3):1429–1437, 2000.

T. C. Quinn, M. J. Wawer, N. Sewankambo, D. Serwadda, C. Li, F. Wabwire-Mangen, M. O. Meehan, T. Lutalo, and R. H. Gray. Viral load and heterosexual transmission of human immunodeficiency virus type 1. *New England Journal of Medicine*, 342(13):921–929, 2000.

B. Rannala and Z. Yang. Probability distribution of molecular evolutionary trees: a new method of phylogenetic inference. *Journal of molecular evolution*, 43(3):304–311, 1996.

N. Rodrigue, H. Philippe, and N. Lartillot. Mutation-selection models of coding sequence evolution with site-heterogeneous amino acid fitness profiles. *Proceedings of the National Academy of Sciences*, page 200910915, 2010.

R. Sanjuán, A. Moya, and S. F. Elena. The distribution of fitness effects caused by single-nucleotide substitutions in an RNA virus. *Proceedings of the National Academy of Sciences*, 101(22):8396–8401, 2004.

T. Stadler. On incomplete sampling under birth–death models and connections to the sampling-based coalescent. *Journal of Theoretical Biology*, 261(1):58–66, 2009.

T. Stadler and S. Bonhoeffer. Uncovering epidemiological dynamics in heterogeneous host populations using phylogenetic methods. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, 368(1614):20120198, 2013.

T. Stadler, D. Kühnert, D. A. Rasmussen, and L. du Plessis. Insights into the early epidemic spread of Ebola in Sierra Leone provided by viral sequence data. *PLoS currents*, 6, 2014.

B. Thyagarajan and J. D. Bloom. The inherent mutational tolerance and antigenic evolvability of influenza hemagglutinin. *Elife*, 3:e03300, 2014.

L. S. Tsimring, H. Levine, and D. A. Kessler. Rna virus evolution via a fitness-space model. *Physical review letters*, 76(23):4440, 1996.

R. A. Urbanowicz, C. P. McClure, A. Sakuntabhai, A. A. Sall, G. Kobinger, M. A. Müller, E. C. Holmes, F. A. Rey, E. Simon-Loriere, and J. K. Ball. Human adaptation of Ebola virus during the West African outbreak. *Cell*, 167(4):1079–1087, 2016.

E. Visher, S. E. Whitefield, J. T. McCrone, W. Fitzsimmons, and A. S. Lauring. The mutational robustness of influenza A virus. *PLoS pathogens*, 12(8):e1005856, 2016.

WHO. Ebola situation reports, 2016. URL `http://apps.who.int/ebola/ebola-situation-reports`.

Z. Yang and R. Nielsen. Mutation-selection models of codon substitution and their use to estimate selective strengths on codon usage. *Molecular Biology and Evolution*, 25(3):568–579, 2008.

F. Zanini and R. A. Neher. Quantifying selection against synonymous mutations in HIV-1 env evolution. *Journal of Virology*, pages JVI–01529, 2013.

Figure 4. Relative fitness of Ebola virus genotypes circulating during the 2013-16 epidemic in Western Africa. Ancestral fitness values were reconstructed by first finding the probability of a lineage being in each possible genotype based on the marginal site probabilities computed using (20). Ancestral fitness values were then computed by averaging the posterior median fitness of all possible genotypes, weighted by the probability that the lineage was in each genotype. Fitness values are given relative to the Makona genotype isolated at the start of the epidemic. Clades are labeled according to their most probable genotype.