Perturbations of a Kinase-Pseudokinase Module Activate Plant Immunity

D. Patrick Bastedo[1], Derek Seto[1¶], Alexandre Martel[1¶], Madiha Khan[1¶], Inga Kireeva[2], Jianfeng Zhang[1], Wardah Masud[1], David Millar[1], Jee Yeon Lee[2], Amy Huei-Yi Lee[3,4], Yunchen Gong[2], André Santos-Severino[2], David S. Guttman[1,2&*], Darrell Desveaux[1,2&*]

[1] Department of Cell and Systems Biology, University of Toronto, Toronto, Ontario, Canada

[2] Centre for the Analysis of Genome Evolution & Function, University of Toronto, Toronto, Ontario, Canada

[3] Department of Microbiology and Immunology, University of British Columbia, Vancouver, British Columbia, Canada

[4] Department of Pharmaceutical Sciences, University of British Columbia, Vancouver, British Columbia, Canada

[¶] These authors contributed equally.

[&] These authors contributed equally.

[*] Corresponding authors

Email: darrell.desveaux@utoronto.ca (DD); david.guttman@utoronto.ca (DSG)

Author Contributions

DPB: Conceptualization, Investigation, Methodology, Validation, Formal Analysis, Software,

Visualization, Writing – Original Draft Preparation, Writing – Review & Editing

DS, AM, MK: Investigation, Methodology, Resources

IK: Investigation, Methodology

JZ: Methodology, Resources

WM, DM, JYL: Investigation

AHL: Methodology, Resources

YG: Data Curation

AS: Resources

DSG: Conceptualization, Data Curation, Funding Acquisition, Resources, Supervision, Writing

– Original Draft Preparation, Writing – Review & Editing

DD: Conceptualization, Methodology, Funding Acquisition, Resources, Supervision, Writing –

Original Draft Preparation, Writing – Review & Editing

Key Words

plant immunity, *Arabidopsis*, pseudokinase, ZED1, ZRK, kinase, PBL, NLR, ZAR1, *P. syringae*,

T3SS, T3E, effector, HopZ1a, YopJ

Short Title

Pseudokinase-kinase interactions in plant immunity

ABSTRACT

The *Pseudomonas syringae* acetyltransferase HopZ1a is delivered directly into host cells by the type III secretion system to promote bacterial growth. However, in the model plant host *Arabidopsis thaliana,* HopZ1a activity results in an effector-triggered immune response (ETI) that limits bacterial proliferation. HopZ1a-triggered immunity requires the nucleotide-binding, leucine-rich repeat domain (NLR) protein, ZAR1, and the ZED1 pseudokinase. Here we demonstrate that HopZ1a can acetylate members of a family of 'receptor-like cytoplasmic kinases' (RLCK family VII; also known as PBS1-like kinases, or PBLs) and promote their interaction with ZED1 and ZAR1 to form a ZAR1/ZED1/PBL ternary complex. Interactions between ZED1 and PBL kinases are determined by the pseudokinase features of ZED1, and mutants designed to restore ZED1 kinase motifs can (1) bind to PBLs, (2) recruit ZAR1, and (3) trigger immunity *in planta*, all independently of HopZ1a. Our results suggest that interactions between these two RLCK families are promoted by perturbations of structural features that distinguish active from inactive kinase domain conformations. We propose that effector-induced interactions between ZED1/ZRK pseudokinases (RLCK family XII) and PBL kinases (RLCK family VII) provide a sensitive mechanism for detecting perturbations of either kinase family and activating ZAR1-mediated ETI.

AUTHOR SUMMARY

All plants must ward off potentially infectious microbes, and those grown in large-scale crop operations are especially vulnerable to the rapid spread of disease by successful pathogens. Although many bacteria and fungi can supress plant immune responses by producing specialized virulence proteins called 'effectors', these effectors can also trigger immune responses that

render plants resistant to infection. We studied the molecular mechanisms underlying one such

effector-triggered immune response elicited by the bacterial effector HopZ1a in the model plant

host *Arabidopsis thaliana*. We have shown that HopZ1a promotes binding between a ZED1, a

'pseudokinase' required for HopZ1a-triggered immunity, and several 'true kinases' (known as

PBLs) that are likely targets of HopZ1a activity *in planta*. HopZ1a-induced ZED1-PBL

interactions also recruit ZAR1, an *Arabidopsis* 'resistance protein' previously implicated in

HopZ1a-triggered immunity. Importantly, ZED1 mutants that restore degenerate kinase motifs

can bridge interactions between PBLs and ZAR1 (independently of HopZ1a) and trigger

immunity *in planta*. Our results suggest that equilibria between active and inactive kinase

domain conformations regulate ZED1-PBL interactions and formation of ternary complexes with

ZAR1. Improved models describing molecular interactions between immunity determinants,

effectors and effector targets will inform efforts to exploit natural diversity for development of

crops with enhanced disease resistance.

INTRODUCTION

Both plants and animals use cell membrane-spanning receptor kinases to sense extracellular molecular patterns produced by invading pathogens. While agonists of these receptors can trigger signal cascades that result in protective host immune responses (pattern-triggered immunity, or PTI), Gram-negative bacteria equipped with a Type III secretion system (T3SS) can dampen such basal immune responses by delivery of effector proteins directly into host cells through the needle-like T3SS pilus. In plant cells, such T3SS-delivered effectors (T3Es) can be recognized by nucleotide-binding leucine-rich repeat (NLR) proteins, resulting in 'effector-triggered immunity' (ETI), which is often accompanied by a hypersensitive cell death response (HR) [1]. NLRs are members of the STAND (signal transduction ATPases with numerous domains) class of P-loop NTPases that also includes the NOD-like receptors of animal cells [2]. These proteins share common central nucleotide-binding and carboxy-terminal LRR domains, but are preceded by distinct classes of amino-terminal domains – plant NLRs have either 'coiled-coil' (CC) or 'Toll/Interleukin-1 Receptor' (TIR) domains at their amino-termini [2–5].

Plant NLRs become activated either by direct sensing of effectors, or by indirect sensing of effector-modified substrates [6]. Such effector-modified substrates can be virulence targets whose functions limit bacterial growth, or mimics of these virulence targets referred to as 'decoys' [7]. For example, the *Arabidopsis* kinase PBS1 (Avr<u>Pph</u><u>B</u>-<u>S</u>usceptible <u>1</u>), is cleaved by the *P. syringae* T3E HopAR1 (previously known as AvrPphB), resulting in activation of the *Arabidopsis* NLR RPS5 [8,9]. There are 45 PBS1-like kinases (PBL kinases, or PBLs) that together with PBS1 comprise receptor-like cytoplasmic kinase (RLCK) subfamily VII, and at least eight of these additional PBLs are also sensitive to cleavage by HopAR1, including BIK1

and PBL1, which are both involved in plant immunity [10]. PBS1 therefore appears to function

as a decoy that mimics HopAR1 virulence targets to activate RPS5-mediated ETI [10].

The *Arabidopsis* NLR ZAR1 (Hop<u>Z</u>-<u>A</u>ctivated <u>R</u>esistance <u>1</u>) belongs to the coiled-coil

class of plant NLRs and is required for ETI against at least three distinct T3Es: HopZ1a and

HopF2a from *Pseudomonas syringae*, and AvrAC from *Xanthomonas campestris* [11–13]. In

addition to ZAR1, recognition of all three T3Es requires members of RLCK family XII.

Members of this RLCK family lack at least one of several well-established and highly conserved

kinase motifs, and are therefore considered to encode pseudokinases [14–16]. Recognition of the

acetyltransferase HopZ1a requires the RLCK XII pseudokinase ZED1 (Hop<u>Z</u> <u>E</u>TI-<u>D</u>eficient <u>1</u>),

which is encoded in a gene cluster with seven additional ZED1-related pseudokinases (ZRKs).

ZRK1/RKS1 (hereafter ZRK1) and ZRK3 pseudokinases are required for recognition of AvrAC

and HopF2a, respectively [12–14]. ZED1, ZRK1 and ZRK3 have all been shown to interact with

ZAR1 in planta, although different roles in T3E recognition have been proposed [13,15,16].

ZED1 is acetylated by HopZ1a, but genetic ablation of *ZED1* does not appear to influence the

virulence of *P. syringae* [14]. As such, it has been proposed that ZED1 is a decoy substrate

monitored by ZAR1 to guard against the acetylation of other (kinase) substrates of HopZ1a [14–

16]. In contrast, ZRK1 has been proposed to function as an adaptor for ZAR1 by recruiting PBL2

(<u>PBS</u>1-<u>l</u>ike kinase 2) molecules uridylated by AvrAC to promote ETI [12]. ZRK3 is also thought

to act as a ZAR1 adaptor for an as-yet-unidentified kinase that is ADP-ribosylated by the T3E

HopF2a [13]. Overall, the ZED1/ZRK family of pseudokinases have expanded the T3E

recognition capacity of ZAR1 by directly sensing T3E modifications and/or serving as adaptors

that bridge interactions between the NLR and T3E-modfied plant kinases.

Highly-conserved kinase motifs were initially identified by comparative sequence

analysis [17,18], but since then, determination of the molecular structures of an ever-increasing

number of diverse kinase domains in the presence or absence of nucleotides, metal ions,

substrate peptides, and/or pharmacological inhibitors has provided rational explanations for the

functional importance of these motifs [19]. The kinase domain nucleotide-binding pocket is

formed by a cleft between a mostly β-stranded amino-terminal lobe and an α-helical carboxy-

terminal lobe. In the amino-terminal lobe, a glycine-rich motif in the loop connecting strands β1

and β2 ('G-loop') acts as a lid through backbone interactions with the β- and γ-phosphates of

bound ATP [20]. Appropriate positioning of ATP phosphates also requires metal ions (usually

Mg^{2+} or Mn^{2+}), a lysine from strand β3 that is stabilized by a salt-bridge with a glutamate from

helix αC, and an aspartate from the 'DFG motif'. The DFG motif also defines the beginning of

the 'activation loop', a conformationally flexible (and often disordered) region of 20-35 amino

acids that typically contains serine and/or threonine residue that can be phosphorylated (by

autophosphorylation or by upstream kinases). Phosphorylation of the activation loop can

stabilize an open conformation (i.e. a 'DFG-in' orientation, and a fully-assembled hydrophobic

'regulatory spine') that is associated with active kinases [21]. This open conformation allows

substrate peptides to approach the γ-phosphate (when appropriately-positioned, as described

above) for phospho-transfer by the aspartate from an 'HRD motif' in the catalytic segment.

These dynamic structural features play essential roles in regulating kinase activity as well as

modulating their interactions with other proteins [19,22]. ZED1 and the ZRKs have degeneracies

in nearly all of these canonical kinase motifs, and as such they may all represent pseudokinases

that perform T3E sensing functions similar to those proposed for ZED1, ZRK1 and ZRK3 [12–

14,16].

Early surveys of the complete kinase domain complements from organisms including

human, mouse, *Caenorhabditis elegans*, *Dictyostelium* and *Arabidopsis* found that

pseudokinases like ZED1 account for ~10% of the kinase domains in higher eukaryotes and 20%

of *Arabidopsis* RLCKs [23,24]. Rather than mere non-functional kinases resulting from relaxed

(or absent) selective constraints, there is instead a growing body of evidence indicating that

pseudokinases are important signaling molecules in spite of their predicted lack of catalytic

activity, often functioning as adaptors or scaffolds that modulate the activity of true kinase

partners [23,25–27]. In this report we show that the *P. syringae* T3E HopZ1a induces

interactions between ZED1 and PBL kinases and promotes the formation of a ZAR1/ZED1/PBL

ternary complex. The pseudokinase features of ZED1 mediate its conditional interactions with

PBL kinases, recruitment of ZAR1, and importantly, its regulation of ZAR1-mediated immunity.

Overall, our results support the hypothesis that ZED1-PBL (and more generally ZRK-PBL)

pseudokinase-kinase interplay provides a sensor for perturbations of host kinases by pathogen-

delivered effector proteins.

RESULTS

HopZ1a binds and acetylates PBL kinases

In order to assess whether PBL kinases are plausible targets of HopZ1a, we tested

whether HopZ1a can interact with members of this family. Using a yeast two-hybrid (Y2H)

screen (S1 Fig; interaction scheme A) with all 46 PBL kinases as prey and HopZ1a as bait (wild-

type or a C216A catalytic site mutant) we found strong binding between nine PBL kinases

(PBL21, PBL27, PBL8, PBL2, PBL3, PBL4, PBL18, PBL15 and PBL13) and HopZ1a[C216A], but

no interactions were observed with the wild-type effector, HopZ1a[wt] (Fig 1A).

We hypothesized that failure to observe binding between HopZ1a^{wt} and PBLs may reflect

efficient enzymatic activity and rapid substrate turnover (i.e. 'catch and release' catalysis). Such

interactions would be stabilized in the case of HopZ1a^{C216A} due to frustrated enzymatic activity.

To test whether HopZ1a can acetylate PBL kinases, we co-expressed FLAG-tagged derivatives

of both HopZ1a and PBS1 in yeast, prepared anti-FLAG immunoprecipitates from yeast cell

extracts and subjected these to liquid chromatography-tandem mass spectrometry (LC-MS/MS)

analysis. Although PBS1 was a weaker HopZ1a interactor, we chose to test its acetylation by

HopZ1a based on its established role in ETI. We observed three PBS1 peptides with mass

increases of 42 Da (corresponding to addition of an acetyl group) when PBS1 was co-expressed

with HopZ1a^{wt} but not with HopZ1a^{C216A}, indicating HopZ1a-mediated acetylation.

Fragmentation analysis of these three peptides established that the specific sites acetylated were

T32, S244, and S405 (Fig 1B; S3 Fig; S1 File). These results support our hypothesis that

HopZ1a can acetylate PBL kinases.

HopZ1a promotes binding between ZED1 pseudokinase and PBL kinases

Given the previously reported finding that AvrAC promotes interactions between PBL2

and ZRK1 [12], we sought to develop an assay to investigate whether HopZ1a can promote

interactions between ZED1 and PBL kinases. We devised a yeast three-hybrid (Y3H) assay

wherein interactions between a ZED1/ZRK bait protein and PBL preys were assessed in the

absence and presence of single-copy chromosomal integrations of a T3E of interest (S1 Fig,

interaction schemes A, B). As proof-of-principle we integrated *Xanthomonas* AvrAC at the yeast

ho locus and investigated its influence on ZRK1 interactions with PBL kinases. In the absence of

AvrAC expression, ZRK1 exhibited high-affinity binding to several PBLs (PBL21, PBL17,

PBL8 and PBL15) and moderate binding to several others, including PBL2 (S4 Fig). Binding

between ZRK1 and PBL2 was enhanced by the presence of AvrAC (as was binding between

ZRK1 and PBL3 / PBL29; S4 Fig), indicating that our Y3H system can be used to examine

modulation of ZRK/PBL interactions by T3Es.

We then integrated *hopZ1a* (wild-type or C216A alleles) at the yeast *ho* locus and

examined its influence on ZED1-PBL interactions. Only weak ZED1-PBL interactions were

observed in the absence of effector (Fig 2B; Fig 3B, column 1) or when HopZ1a^{C216A} was

expressed (Fig 2A; Fig 3B, column 3). Notably, expression of wild-type HopZ1a resulted in

enhanced interactions between ZED1 and 11 PBL kinases (PBL21, PBL22, PBL5, PBS1,

PBL27, PBL17, PBL8, PBL4, PBL9, PBL15, and PBL13), with the strongest interactions

occurring between ZED1 and PBL5 / PBL27 / PBL17 / PBL8 / PBL15 (Fig 2A; Fig 3B, column

2). These results indicate that HopZ1a activity can promote interactions between the ZED1

pseudokinase and PBL kinases.

In order to assess whether stimulation of ZED1-PBL binding was a property specific to

HopZ1a, we also created yeast strains with chromosomal integrations of genes encoding both

closely-related (HopZ1b; 65% amino acid identity to HopZ1a over 349 non-gapped sites) and

more distantly-related HopZ family members (HopZ2, HopZ3; 26% and 23% identical to

HopZ1a over 335 and 306 non-gapped sites, respectively). Interactions between ZED1 and PBLs

were assessed in the presence of each of these HopZ family effectors or with the unrelated

effector AvrAC (Fig 2B; S3 Fig; Fig 3B, columns 2-7). Like HopZ1a, HopZ1b promoted strong

interactions between ZED1 and PBL27 / PBL17 / PBL8, but in contrast, interactions between

ZED1 and PBL5 / PBL15 / PBL21 were weaker with HopZ1b than with HopZ1a, and HopZ1b

did not promote interactions between ZED1 and PBL 22 / PBS1 / PBL4 / PBL9 (Fig 2 and Fig

3B, columns 2, 4). Notably, the interactions between ZED1 and PBLs were unchanged by the

expression of HopZ2, HopZ3, or AvrAC (Fig 3B, columns 5-7; S5 Fig, panels A, B) although

integrated effectors are expressed to comparable levels (S5 Fig, panel C), demonstrating that

promotion of ZED1/PBL interactions is specific to HopZ1a (and to a lesser extent, HopZ1b).

Perturbation of ZED1 pseudokinase features modulates PBL interactions

The kinase domains of ZED1 and the ZRKs have diverged significantly from those of true

kinases, with degeneracies in one or more of the established kinase motifs (S6 Fig) [14]. They

also generally lack kinase activity under *in vitro* conditions that are sufficient for canonical

kinases to phosphorylate model substrates [14,28]. To determine the importance of ZED1

pseudokinase features for (HopZ1a-induced) binding to PBLs, we identified conspicuous ZED1

sequence elements based on their divergence from canonical kinase motifs and designed

mutations intended to 'restore' these degenerate motifs. We made a series of ZED1 variants with

various combinations of mutations at five different sites (Fig 3A), described below.

a) The glycine-rich motif ('GXGGFG') between strands β1 and β2 is unrecognizable in

211   ZED1 (S6 Fig, panel B). The triple mutant (S56G W58G F61G; hereafter referred to as

212   ZED1$^{'3xG'}$) restores a glycine-rich motif at this position.

213   **b)** The salt-bridge formed between the lysine from strand β3 and the glutamate from helix

214   αC is evident in the crystal structure of the *Arabidopsis* receptor kinase BRI1 (between

215   K911 and E927; S7 Fig, panel B) [29], however a homology-based structural model of

216   ZED1 indicates that the sidechains of the corresponding residues (K76 and D92) are too

217   far apart (S7 Fig, panel A). ZED1^{D92E} extends the acidic side chain at this position by a

218   single CH$_2$ group to restore an αC glutamate.

c) ZED1^{N173D} restores the active site aspartate of the HRD motif required for phospho-

transfer in active kinases.

d) In ZED1 the DFG motif glycine is replaced by a tryptophan (S6 Fig, panel B; S7 Fig,

panel A). ZED1^{W193G} restores the activation loop DFG motif.

e) ZED1 is unique, even among the ZRKs, in that its activation loop is completely devoid of

potential phospho-accepting serine or threonine residues (S6 Fig, panel B). We thus

created ZED1^{V212T}, since this position is an absolutely conserved threonine in all of the

other kinase sequences considered in our analysis, and its phosphorylated derivative has

been observed in a crystal structure of the *Arabidopsis* BAK1 kinase domain [30].

We tested the PBL-binding activity of each of these rationally-designed ZED1 mutations

(individually and in various combinations) both in the absence and presence of HopZ1a

expression using Y2H and Y3H assays. We examined the binding of ZED1 mutants to a

restricted subset of 15 PBLs that includes ten representatives with robust HopZ1a-induced

binding affinity to wild-type ZED1 as well as five with relatively weak HopZ1a-dependant

ZED1-binding activity (PBL24, PBL36, PBL18, PBL12, PBL11).

Our mutational analysis found that ZED1^{N173D} (HRD restoration), ZED1^{W193G} (DFG

restoration) and ZED1^{V212T} (activation loop restoration) all gained affinity for several PBL

kinases in the absence of HopZ1a, with ZED1^{N173D} and ZED1^{V212T} showing the strongest

interactions (S8 Fig and S9 Fig; Fig 3B, compare columns 12, 13, 18, 19 with columns 1, 10,

17). These mutations induced HopZ1a-independent ZED1 binding to PBL kinases showing both

strong (PBL21, PBL27, PBL15, PBL17, PBL4) and weak (PBL18) HopZ1a-dependent binding.

HopZ1a-independent binding between PBS1 and ZED1 was only observed in the context of the

ZED1^{N173D} mutation (i.e. HRD restoration; ZED1^{N173D} and ZED1$^{D92E\ N173D}$; S8 Fig and S9 Fig;

Fig 3B, columns 12, 14, 18). The triple mutant ZED1$^{\text{'3xG'}}$ (glycine loop restoration) disrupted

HopZ1a-induced PBL binding, and only moderate HopZ1a-independent binding was observed

(S9 Fig, panel B; Fig 3B, column 21). Combining ZED1 mutations differentially influenced

subsets of PBL interactions. For example, combining N173D with W193G (ZED1$^{\text{N173D W193G}}$; S8

Fig, panel B; Fig 3B, column 15) or with D92E and W193G (ZED1$^{\text{'EDG'}}$; S6 Fig, panel B; Fig

3B, column 16) resulted in a loss of PBS1 binding (observed with ZED1$^{\text{N173D}}$ and ZED1$^{\text{D92E}}$

$^{\text{N173D}}$; S8 Fig; Fig 3B, columns 12, 14, 18) but increased ZED1 binding affinity for PBL21 and

PBL24 (not observed with any of the three individual mutations; S8 Fig, panel A and S9 Fig,

panel A; Fig 3B, columns 11, 12, 13, 18). When we combined N173D with the restored glycine

loop (ZED1$^{\text{'3xG' N173D}}$), all binding to PBLs was abolished whether HopZ1a was present or not

(S9 Fig, panel B; Fig 3B, column 22). However, a subset of interactions was restored in the

context of a phospho-accepting activation loop in ZED1 (ZED1$^{\text{'3xG' V212T}}$ and ZED1$^{\text{'3xG' 'DT'}}$,

which adds both N173D and V212T; S9 Fig, panel B; Fig 3B, columns 23, 24). These results

suggest that the effects of ZED1 mutations differentially affect binding affinity for distinct PBLs,

indicating that PBLs differ in their abilities to interact with ZED1 through perturbation of

(pseudo)kinase features either through mutation or by T3E modification.

HopZ1a acetylation sites influence ZED1/PBL interactions

Based on the results of our ZED1 mutagenesis, we hypothesized that acetylation of one

or more sites on ZED1 might influence ZED1/ PBL binding by altering its pseudokinase

properties. We have previously described acetylation of ZED1 by HopZ1a at T125 and T177

[14], and subsequent experiments have indicated that S84 and T87 can also be acetylated by

HopZ1a (Fig 1C; S3 Fig; S1 File). We therefore made glutamine substitutions as potential

mimics of acetylated serine and/or threonine residues based on similarities in sidechain length

and branching structure (Fig 4A). We tested the PBL-binding capacity of three ZED1 glutamine

substitution mutants: ZED1^{T177Q}, which is proximal to the (non-catalytic) active site residue

N173, as well as ZED1^{T87Q} and ZED1^{S84Q}, which are both predicted to be part of a solvent-

exposed surface of helix αC (S7 Fig, panel A; S10 Fig, panel A). None of the ZED1 glutamine

substitution mutants (or any of the three possible pairwise combinations) resulted in HopZ1a-

independent binding to any of the PBLs tested (S10 Fig, panel B). In contrast, each of the

mutants bearing the T177Q substitution (ZED1^{T177Q}, ZED1$^{S84Q\ T177Q}$, ZED1$^{T87Q\ T177Q}$) lost all

capacity for HopZ1a-dependent binding (S10 Fig, panel B), indicating that ZED1^{T177Q} represents

a loss-of-function allele.

We also tested whether the PBS1 acetylation sites described above (Fig 1B; S3 Fig; S1

File) are important for its interaction with ZED1. Similar to our ZED1 findings, we found that

none of the PBS1 glutamine substitutions (alone or in combination) enable HopZ1a-independent

ZED1-binding activity (S10 Fig, panel C). Furthermore, as for ZED1 T177, each of the PBS1

mutants bearing a glutamine substitution of S244 lost capacity for HopZ1a-dependent binding to

ZED1 (S10 Fig, panel C). The combined insights from our mutagenesis of ZED1 and PBS1

acetylation sites suggest that glutamine does not mimic acetylated serine/threonine residues and

may instead act as a loss-of-function mutation by preventing acetylation and/or phosphorylation.

We also considered isoleucine substitutions as possible mimics of acetylated serine

and/or threonine residues based on similarities in hydrophobicity and sidechain branching (Fig

4A). We therefore tested glutamine and isoleucine (possible acetyl-mimics) as well as alanine

(loss-of-function) substitutions in PBL15, which was selected as a representative PBL based on

its strong HopZ1a-dependent binding to ZED1 (stronger than PBS1; Fig 2A). We mutated

PBL15 S260, an activation loop residue equivalent to PBS1 S244 (which is acetylated by

HopZ1a; Fig 1B). Of these three substitutions of PBL15 S260, only PBL15^{S260I} demonstrated

ZED1 binding in the absence of HopZ1a activity (Fig 4B). The strength of this interaction was

comparable to the ZED1-PBL15^{wt} binding induced by HopZ1a, and HopZ1a-induced ZED1

binding was abolished by the alanine and glutamine substitutions of S260. Overall, these results

suggest that acetylation of the activation loops of PBL kinases by HopZ1a is sufficient to

promote their interactions with ZED1 and that isoleucine substitutions can mimic acetylation of

serine (and possibly threonine) residues.

HopZ1a promotes the formation of a ZAR1-ZED1-PBL ternary complex

Having established that HopZ1a is able to stimulate protein-protein interactions between

ZED1 and PBLs, we wished to investigate whether HopZ1a can also promote formation of a

ternary complex between PBLs, ZED1, and ZAR1. For this purpose, we developed a yeast four-

hybrid system where ZAR1 baits and PBL kinase preys (11 with HopZ1a-dependent binding)

were expressed in the presence or absence of chromosomally-integrated *hopZ1a* (wild-type or

the C216A catalytic site mutant) and *ZED1* (S1 Fig, interaction scheme C). Since modulation of

inter-domain contacts is known to be important for the activation of plant NLRs [6,31], we

created various domain truncations of ZAR1 (panel A of both Fig 5 and S11 Fig) in order to

investigate domain-specific ZAR1 binding interactions with ZED1/PBLs that may be might be

modulated by HopZ1a activity. Full-length ZAR1 bait (ZAR1^{wt}) interacted with prey PBL

kinases (PBL5, and weaker interactions with PBL17, 4, 18, 15) when co-expressed with both

ZED1 and HopZ1a^{wt}, but not when co-expressed with ZED1 and HopZ1a^{C216A} (Fig 5B, compare

columns 4 and 6). These interactions were dependent on ZED1 expression (Fig 5B, compare

columns 3 and 4) and also required the ZAR1 LRR domain, since no interactions were observed

with the ZAR1$^{\Delta LRR}$ construct (Fig 5B, compare columns 4 and 12). Consistent with this finding,

the isolated LRR domain displayed weak HopZ1a^{wt}/ZED1-dependent PBL interactions, but the

CC and NB domains did not (S11 Fig, panel B, columns 4, 10, 16). Interestingly, ZAR1$^{\Delta CC}$

showed stronger HopZ1a/ZED1-dependent interactions with 9 of the 11 PBLs (Fig 5B, compare

columns 4 and 20), suggesting that the ZAR1 coiled-coil domain negatively regulates

ZAR1/ZED1/PBL complex formation.

Although we have demonstrated above that ZED1 mutants with restored kinase motifs

gain HopZ1a-independent binding affinity for PBL kinases, it is possible that these interactions

are formed by a binding interface that differs from that required for HopZ1a-dependent PBL-

ZED1-ZAR1 interactions. To address this, we integrated genes encoding ZED1^{wt} or ZED1$^{‘DT’}$

(ZED$^{N173D\ V212T}$) at the yeast *ho* locus to create a HopZ1a-independent Y3H system in which to

test possible PBL-ZED1-ZAR1 interactions. Expression of ZED1$^{‘DT’}$ allowed HopZ1a-

independent interactions between ZAR1^{wt} and PBL17 / PBL9, whereas ZED1^{wt} did not (Fig 5B,

compare columns 7 and 8). Moreover, in the presence of ZED1$^{‘DT’}$ (but not ZED1^{wt}), ZAR1$^{\Delta CC}$

interacts more robustly and with more PBLs (eight of the eleven tested; Fig 5B, compare

columns 7 and 8 with 23 and 24), indicating that the coiled coil domain negatively regulates

formation of the ZAR1/ZED1/PBL ternary complex in HopZ1a-independent contexts as well.

Importantly, ZAR1$^{\Delta LRR}$ did not interact with PBL kinases in the presence of ZED1$^{‘DT’}$ indicating

that the leucine-rich repeat domain is required for both HopZ1a-dependent and HopZ1a-

independent interactions.

ZED1 mutants with HopZ1a-independent PBL-binding promote HR in *Arabidopsis*

We hypothesized that ZED1/PBL interactions promote ZAR1-mediated immunity,

similar to the ZRK1/PBL2/ZAR1-mediated immunity triggered by AvrAC [12]. As such, mutant

ZED1 alleles that allow HopZ1a-independent interactions with PBL kinases should also promote

immunity in the absence of the effector. To test this, we transformed the *Arabidopsis* 'Col-0'

ecotype with four dexamethasone-inducible, HA epitope-tagged ZED1 alleles (ZED1^{wt},

ZED1^{N173D}, ZED1^{V212T}, and ZED1$^{'DT'}$), and examined the first generation (T1) for a

dexamethasone-induced tissue collapse (HR) similar to that observed in plants expressing the *P.*

syringae effector AvrRpt2 [32] (see column 2 in Fig 6AB, S12 Fig AB, and S13 Fig AB). We

established expression profiles for at least 8 individual T1 transformants from each ZED1 allele

(panel C in Fig 6, S12 Fig, S13 Fig) prior to testing for dexamethasone-inducible cell death

activity. These same plants (with one leaf removed), along with untransformed Col-0 and *zar1-1*

mutant [11] plants as controls, were then sprayed with dexamethasone and observed visually for

signs of dexamethasone-induced HR. As expected, untransformed wild-type *Arabidopsis* (Col-0)

and *zar1-1* mutant plants were both unaffected by dexamethasone treatment (columns 1 and 3 in

Fig 6AB, S12 Fig AB, and S13 Fig AB). *Arabidopsis* plants expressing ZED1^{V212T} and ZED1$^{'DT'}$

displayed tissue collapse similar to AvrRpt2-expressing plants 72 hours after dexamethasone

treatment (columns 4 and 5 in S13 Fig AB; columns 6 and 7 in Fig 6AB), whereas plants

expressing ZED1^{wt} and ZED1^{N173D} did not (columns 4, 5 in Fig 6AB; columns 4-7 in S12 Fig

AB). These results suggested that PBL-interacting ZED1 mutants can trigger HopZ1a-

independent immunity and led us to investigate whether this response is ZAR1-dependent.

ZED1 mutants trigger ZAR1-dependent HR in *Nicotiana benthamiana*

Nicotiana benthamiana has two ZAR1 homologues (*Nb*ZAR1 and *Nb*ZAR2) that are

~57% identical to *Arabidopsis* ZAR1 [33]. Despite lacking closely-related homologues of ZED1

and ZRKs (S14 Fig), ZAR1-dependent immune responses triggered by HopZ1a can be

recapitulated in *N. benthamiana* when complemented with *Arabidopsis* ZED1 [33]. Cell death

resulting from an HR is observed as early as 8 hours following induction of transgene expression

in *N. benthamiana* leaves co-transformed with ZED1 and HopZ1a, but not in leaves transformed

with either ZED1 or HopZ1a alone [33]. Since BLASTP searches demonstrate that *Nicotiana*

spp. do have highly-similar homologues of *Arabidopsis* PBLs (S14 Fig), we hypothesized that

our ZED1 mutants with HopZ1a-independent PBL binding may be sufficient to trigger ZAR1-

dependent HR in *N. benthamiana*. We therefore tested the ability of ZED1 mutants described

above to cause HR in *N. benthamiana* using transient transformations. As described previously

[33], macroscopic HR-like cell-death was apparent by 8-24 h post-induction of protein

expression in *N. benthamiana* tissue co-transformed with ZED1^{wt} and HopZ1a, but not in tissue

transformed with either ZED1^{wt} or HopZ1a alone (Fig 7; S15 Fig). In contrast, ZED1$^{‘DT’}$ and

ZED1^{V212T} were able to cause HR in *N. benthamiana* even in the absence of HopZ1a (Fig 7; S15

Fig). As observed in *Arabidopsis,* ZED1^{N173D} did not display HR-like collapse but did produce a

slow, late-onset yellowing of the infiltrated spot (Fig 7). Overexpression of wild-type ZED1 did

not induce HR despite similar levels of expression as ZED1$^{‘DT’}$ (S13 Fig, panel B).

We also tested ZAR1-dependence of this HopZ1a-independent cell death by subjecting

N. benthamiana plants to virus-induced gene silencing (VIGS) [33–35] two weeks prior to

transformation and induction of ZED1 expression. Although the HR-mediated cell death induced

by ZED1$^{‘DT’}$, ZED1^{V212T} and HopZ1a/ZED1^{wt} was unaffected in plants experiencing silencing of

the negative control *GUS* gene (Fig 7, left; S15 Fig), the HR induced by all of these alleles was

completely abolished in plants that received a *ZAR1* silencing construct (Fig 7, right).

DISCUSSION

In this study we have examined how kinase-pseudokinase interactions induced by the *P.*

syringae T3E HopZ1a contribute to the induction of plant immunity. We have shown that

HopZ1a can acetylate PBL (RLCK family VII) kinases and promote their interactions with the

ZED1 pseudokinase (RLCK family XII). In addition, we have also shown that ZED1 mutants

expected to adopt structural conformations more similar to those of active kinases gain HopZ1a-

independent PBL-binding activity and induce a ZAR1-dependent hypersensitive immune

response (HR) *in planta*. Overall our data suggest that ZED1-PBL interactions provide a sensor

that can monitor perturbations of either (or both) kinase families to trigger ZAR1-mediated ETI.

The switch-like structural features of kinases - including their propensity to conditionally interact

with one another - seem to have been exploited by the plant immune system to survey for

effector-induced perturbations of the plant kinome.

Models for ZAR1 activation by HopZ1a

According to our original decoy model of ZED1 function, acetylation of ZED1 should be

sufficient to activate ZAR1 and trigger immunity [14] (Fig 8, model A). Although the HopZ1a-

induced ZED1-PBL (and ZED1-PBL-ZAR1) interactions described in this study argue against a

strict decoy model and lend support to the adaptor model proposed by Wang *et al.* [12] (Fig 8,

model B), our experiments have not ruled out a role for effector-mediated modification of ZED1

in the activation of ZAR1. Indeed, HopZ1a can acetylate ZED1 at sites proximal to kinase motifs

of known functional importance (S7 Fig, panel A), and mutation of the ZED1 acetylation site

T177 is sufficient to abolish ZED1-PBL binding (S10 Fig, panel B). Furthermore, just as

HopZ1a activity can promote ZED1-PBL binding (Fig 2A), so too can ZED1 mutations that

restore degenerate pseudokinase motifs so that they more closely resemble those present in

catalytically-active 'true kinases' (Fig 3B; S8 Fig; S9 Fig). We speculate that these mutations

allow ZED1 to adopt an 'active-like' pseudokinase structure similar to one that may arise

following acetylation of ZED1 by HopZ1a. We therefore recognize an additional model for

activation of the PBL-ZRK-ZAR1 complex to recognize that effector-mediated modification of

either ZRKs or PBL kinases may be sufficient to activate ETI (Fig 8, models B, C). We

hypothesize that T3E-induced modifications of ZRKs promote an 'inactive' to 'active' structural

change that promotes interaction with 'active' PBL kinases to activate ZAR1 (Fig. 8, model C).

Conversely, PBL modification promotes an 'active' to 'inactive' structural change that promotes

interactions with 'inactive' ZRK pseudokinases that can (also) result in ZAR1 activation (Fig. 8,

model B). Clearly, future structural characterization of ZRK-PBL heterodimers (and of ZRK-

PBL-ZAR1 complexes) will be critical for providing experimental validation of the structural

speculations underlying our models portraying ZRK/PBL modules as molecular switches for

sensing kinase perturbations.

Since PBL kinases were not identified in our earlier forward genetic screen for loss of

HopZ1a-induced HR [14] (in contrast to the susceptibility of *Arabidopsis pbl2* mutants to

infections with AvrAC-expressing *X. campestris* [12]), we expect that there are multiple PBLs

capable of engaging ZED1/ZAR1 to activate HopZ1a-elicited ETI. As such, PBL kinases may be

individually dispensable for HopZ1a-induced ETI due to functional redundancy, or alternatively

one or more of these are essential for viability, precluding their recovery as loss-of-function

mutants.

It is likely that the PBL kinases with HopZ1a-dependent ZED1 binding include virulence

targets that are manipulated to promote *P. syringae* pathogenesis. Notably, a number of PBLs

have been implicated in plant immunity, including BIK1 [36–39], RIPK [40,41], PBL13 [42],

and PBL27 [43,44], and we have shown that these latter two participate in HopZ1a-dependent

interactions with ZED1. These and/or other PBLs represent likely virulence targets of HopZ1a in

the absence of ZAR1 recognition.

The ZED1 activation loop is implicated in PBL binding and immune activation *in planta*

Several lines of evidence suggest that (pseudo)kinase activation loops are specific targets

of HopZ1a (and other effectors). Although both ZED1^{N173D} and ZED1^{V212T} mutants acquired

strong HopZ1a-independent PBL binding activity, only the ZED1^{V212T} and ZED1$^{N173D\ V212T}$

mutants (with restored activation loops) were sufficient to induce immunity *in planta* (Fig 6; S13

Fig; Fig 7; S15 Fig). ZED1 is unique among the ZRKs in that it lacks any candidate phospho-

accepting residues in the activation loop (S6 Fig, panel B). This feature is not specific to the Col-

0 ecotype of *Arabidopsis*, since examination of 813 unique ZED1 sequences representing more

than 1000 distinct and globally-distributed ecotypes identified only four with potential

phospho/acetyl-accepting sites between the imperfect 'DFG' and 'APE' motifs that define the

activation loop (see Materials and Methods). The HopZ1a acetylatation target S244 of PBS1 (Fig

1B) provides further evidence for the functional importance the activation loop in kinase-

pseudokinase interactions. Glutamine substitution of this residue blocks HopZ1a-induced ZED1

binding activity, whereas an isoleucine substitution of the homologous activation loop position in

PBL15 (S260I) acts as a gain-of-function mutant that mimics HopZ1a-induced ZED1-PBL

binding (Fig 4B). These results suggest that kinase-pseudokinase (PBL-ZRK) interactions can be

promoted by activation loop-dependent structural changes in either of these protein families. An

interface between ZED1 and PBLs involving the sites implicated either by acetylation and/or by

mutational analyses may be similar to a recently reported structure describing a mutant IRAK4

kinase domain [45]. This mutant (bearing a catalytic site aspartate to asparagine substitution) was

crystalized as a dimer in the presence of the general kinase inhibitor staurosporine, and it also

gained the ability to dimerize in solution [45]. Remarkably, homology modeling that

superimposes ZED1 and PBS1 on the two monomers of the asymmetric IRAK4 homodimer

indicates that such a 'front-to-front' interface positions the acetylated PBL activation loop

residue (i.e. PBS1 S244 or PBL15 S260) in very close proximity to ZED1 residues implicated by

our mutational analyses (N173, T177, and V212; S16 Fig).

Kinase activation loops are also targeted by other T3Es. Cleavage of PBS1 by the T3E

HopAR1 occurs after the lysine (K243) immediately preceding S244 [8], a modification that is

perceived by the NLR RPS5 [9]. AvrAC uridylates serine and threonine residues in the

activation loops of PBL2, BIK1 and RIPK [46,47], similar to the acetylation of PBS1 by

HopZ1a. These conserved AvrAC uridylation sites are required for PBL2 interactions with

ZRK1/RKS1 [12], and are just two positions 'downstream' of PBS1 S244 (S6 Fig, panel B),

providing additional support for the functional significance of modifications in this region of the

activation loop. In human cells, YopJ (a HopZ-related effector from the human plague pathogen,

Yersinia pestis) acetylates the activation loops of MAP kinase kinases (MEK2, MAPKK6) and

both the α and β subunits of the IκB kinase (IKK) complex, resulting in inhibition of MAP

kinase and NF-κB signaling pathways [48,49]. It has long been established that the

phosphorylation status of activation loops has a role in influencing conformational changes that

both position catalytic components and regulate access of substrates to the catalytic site [50,51].

As such, kinase activation loops may represent attractive targets for invading pathogens, while

also providing the structural influence required to sense and transduce kinase domain

perturbations.

476　ZAR1 recruitment

Our yeast four-hybrid data indicate that ZAR1 forms a ternary complex with ZED1 and

PBL kinases that is stimulated by the presence of HopZ1a. Formation of this ternary complex is

dependent on ZED1 and is reminiscent of the ZAR1/ZRK1/PBL2 ternary complex likely formed

as a result of uridylation of PBL2 by AvrAC [12]. Notably, an intact ZAR1 LRR domain is

required for both ZAR1-ZRK1 interactions [12] and also for formation of the ZAR1-ZED1-PBL

ternary complex promoted by HopZ1a (Fig 5), consistent with co-immunoprecipitation

experiments in *N. benthamiana* with *Arabidopsis* ZED1 and ZAR1 truncation mutants [33].

These findings are distinguished from the PBS1/RPS5 model wherein PBS1 interacts with the

coiled-coil (CC) domain of RPS5 until cleavage of the PBS1 activation loop by HopAR1

abolishes this interaction, suggesting that in the case of RPS5, binding by PBS1 is required to

restrain an inherently active NLR [9]. Although direct interactions between ZED1 and the ZAR1

CC domain have been previously reported [14,33], our data suggest indicate that the CC domain

of ZAR1 can negatively regulate formation of a ZAR1/ZED1/PBL ternary complex (Fig 5B).

Overall our results support models suggesting that acetylation of ZED1 and/or of PBL kinases by

HopZ1a promotes formation of this ternary complex with ZAR1; this process requires the LRR

domain and is likely to be negatively-regulated by the ZAR1 CC domain (Fig 8, models B, C).

Ternary complex formation may occur by stabilizing a preformed ZAR1/ZED1 complex, as

proposed for the ZAR1/ZRK1/PBL2 interaction [12]. Alternatively, HopZ1a-stabilized

ZED1/PBL dimers may bind ZAR1 as a pre-formed unit to activate immunity, since we have

found that HopZ1a-induced ZED1/PBL interactions are ZAR1-independent (Fig 2A; Fig 4B; S8

Fig; S9 Fig; S10 Fig, panels B, C).

Functional diversity of HopZ alleles

HopZ1a is a member of a large and diverse family of bacterial effector proteins with

similarity to the YopJ effector from *Y. pestis*. YopJ-like effectors are produced by a variety of

pathogens of both plants and animals, and even among *P. syringae* strains there are five

recognized distinct lineages of HopZ effectors (HopZ1 through HopZ5) [52,53]. This inter-

allelic sequence variation confers significant functional differences, since only HopZ1a is

capable of inducing ETI in *Arabidopsis*. HopZ1b, the allele most closely-related to HopZ1a

(64% identity), can trigger a weak, HR-like tissue collapse, but this response is ZAR1-

independent and is only observed in ~25% of infiltrated leaves [11,54]. HopZ1b induced only a

subset of the PBL-ZED1 interactions promoted by HopZ1a, consistent with its inability to

activate robust ZAR1-dependent immunity. The PBL kinases that interact with ZED1

specifically (or more strongly) in the presence of HopZ1a than HopZ1b thus represent promising

candidates for key regulators of ZAR1 activation by HopZ1a. HopZ3, a more distantly-related

HopZ allele (23% identity to HopZ1a), does not activate ZAR1 but is able to interact with PBLs

(RIPK, PBS1, BIK1, and PBL1) and can also acetylate the activation loop of RIPK (S251, S252;

identical to the sites uridylylated by AvrAC) to dampen ETI mediated by the NLR RPM1 [55].

In our assays, however, co-expression with HopZ3 did not confer ZED1 binding activity upon

RIPK or any other PBL (S5 Fig). Overall, the promotion of most ZED1/PBL interactions is

specific to HopZ1a/HopZ1b, suggesting that other HopZ alleles may have adopted distinct

modification strategies to avoid recognition of their modified substrates by ZED1 and ZAR1 in

Arabidopsis.

Interestingly, a related HopZ-like T3E from *Xanthomonas perforans*, XopJ4/AvrXv4

(25.5% identical to HopZ1a across 337 non-gapped sites) triggers a ZAR1-dependent ETI in *N.*

benthamiana that is dependent on an RLCK XII protein named XOPJ4 IMMUNITY 2 (JIM2)

[56]. ZED1 is the *Arabidopsis* protein most closely-related to JIM2 (BLASTP E-value of 5e-73;

41% identity spanning 92% of the JIM2 query sequence) and notably, JIM2 has degenerate

kinase motifs (including a 'dead' HRD catalytic motif - 'YRI'), suggesting that NbZAR1 may

also use a pseudokinase/kinase module to detect perturbations of *N. benthamiana* signaling [56].

Concluding Statement

Kinases and pseudokinases possess switch-like structural features that are often

allosterically regulated by post-translational modifications (PTMs) to influence their activity (in

the case of active kinases) and/or alter their protein interaction profiles, resulting in

reorganization and redistribution of signaling network activities. The plant immune system

appears to have harnessed these switch-like features to detect kinase/pseudokinase perturbations

that are induced by pathogen-delivered effector proteins. Modifications of either the ZED1/ZRK

pseudokinases (RLCK XII) or of PBL kinases (RLCK VII) can promote interactions between

these two families, as well as subsequent/concomitant interactions with the NLR ZAR1 to

activate ETI. ZRK/PBL interactions are likely determined by a structural switch that is flipped

by effector-mediated PTMs (Fig 8), providing an array of discriminating sensors that can detect

diverse perturbations of the plant kinome (including both kinases and pseudokinases). Further

investigations of these intermolecular interactions, the ways in which they are influenced by

bacterial effectors, and the structural features required of these components for robust immunity

signaling will together provide valuable insights into the molecular mechanisms underlying

pathogen perception by plant immune systems.

MATERIALS AND METHODS

Construction of plasmids for yeast two-hybrid assays

PBL coding sequences were obtained from the ABRC (where available) as pENTR223 or

pUNI51 clones (Supplementary Table 1). Additional PBL coding sequences were synthesized by

General Biosystems, Inc. (USA) as Gateway^{TM}-compatible pDONR207 clones. Recombination

reactions using 'LR clonase' (Invitrogen) were used to shuttle coding sequences into a Gateway-

compatible derivative of the prey plasmid pJG4-5 (which encodes HA-tagged amino-terminal

fusions of the B42 activation domain with nuclear localization sequences; NLS-B42-HA-prey).

Donor plasmids carrying truncations of ZAR1 lacking the LRR (ΔLRR, nucleotides 1-

1545, amino acids 1-515) or coiled-coil (ΔCC, nucleotides 433-2556, amino acids 145-852)

domains, or carrying isolated individual domains (CC, nucleotides 1-432, amino acids 1-144;

NB, nucleotides 433-1545, amino acids 145-515; LRR, nucleotides 1546-2556, amino acids 516-

852) were prepared by PCR amplification and recombination into pDONR207 with BP clonase

(Invitrogen). Subsequent LR reactions were used to shuttle these constructs into the bait plasmid

pEG202 (which encodes an amino-terminal LexA DNA-binding domain; LexA-bait). Plasmids

were introduced into yeast strains according to standard LiAc/PEG methods [57,58] following

sequence verification by sequencing both DNA strands of the entire coding sequences and across

cloning junctions for each clone of interest, using Sanger sequencing services provided by the

Center for the Analysis of Genome Evolution and Function (CAGEF, University of Toronto).

Co-expression/immunoprecipitation of PBS1 and ZED1 with HopZ1a in yeast and sample

preparation

PBS1 was shuttled from pDONR207 into the centromere-based yeast plasmid pBA350V

[59] using LR clonase to create a galactose-dependent expression vector for production of PBS1-

FLAG in yeast. pBA350V::*PBS1-FLAG* was introduced into a derivative of yeast strain Y7092

with chromosomally-integrated *hopZ1a-FLAG* at the *ho* locus, as described previously [14,60].

FLAG-tagged proteins were expressed in yeast, and cell lysates were prepared and

probed with anti-FLAG-conjugated agarose resin (Sigma) as described previously [14,60].

FLAG-tagged proteins immunoprecipitated in this manner were eluted by incubating with 100

uL of FLAG peptide solution (150 ug/mL FLAG peptide in TBS) for one hour at 4°C. Eluted

material was then dried to a pellet under vacuum and stored at -80°C prior to subsequent mass

spectrometry analysis. Dried protein samples were re-solubilized in 50 mM ammonium

bicarbonate (pH 7.8) and then subjected to reduction with dithiothreitol at 56°C, alkylation with

iodoacetamide at room temperature, and overnight digestion with sequencing-grade trypsin

(Promega) at 37°C. This enzymatic reaction was terminated by addition of formic acid (to 3%),

and digestion products were purified and concentrated with Pierce C18 spin columns (Thermo

Fisher Scientific), then again dried to a pellet under vacuum. Peptide samples were then

solubilized in 0.1% formic acid prior to LC-MS/MS analyses.

LC-MS/MS analysis of immunoprecipitated proteins - chromatography and mass

spectrometry

Subsequent analytical separation was performed on a homemade, 75 μm i.d. column

(New Objective, Woburn, MA) gravity-packed with 10 cm of 100 Å, 5 μm Magic C18AQ

particles (Michrom, Auburn, CA). Peptide samples were loaded onto the analytical column using

a variable gradient with a flow rate of 300 nL/min. The gradient utilized two mobile phase

solutions: A - water/0.1% formic acid; and B - 80% acetonitrile/0.1% formic acid. Samples were

analyzed on a linear ion trap-Orbitrap hybrid analyzer outfitted with a nano-spray source and

EASY-nLC 1200 nano-LC system. The instrument method consisted of one MS full scan (400–

1400 *m/z*) in the Orbitrap mass analyzer, an automatic gain control target of 500,000 with a

maximum ion injection of 500 ms, one microscan, and a resolution of 60,000. Six data-

dependent MS/MS scans were performed in the linear ion trap using the three most intense ions

at 35% normalized collision energy. The MS and MS/MS scans were obtained in parallel

fashion. In MS/MS mode automatic gain control targets were 10,000 with a maximum ion

injection time of 100 ms. A minimum ion intensity of 1000 was required to trigger an MS/MS

spectrum. The dynamic exclusion was applied using an exclusion duration of 145 s. Each sample

was analyzed in triplicate. A spectral library was created using Proteome Discoverer 2.0

(Thermo Fisher Scientific).

Protein identification and database searches

Proteins were identified by searching all MS/MS spectra against a large database

composed of the complete proteome of *Saccharomyces cerevisiae* strain S288C (ATCC 204508;

UniProt proteome ID UP000002311) supplemented with sequences for *P. syringae* HopZ1a

(WP_011152901.1), and the *Arabidopsis* kinases PBS1 (NP_196820.1) or ZED1 (NP_567053.1)

using SEQUEST [61]. A fragment ion mass tolerance of 0.8 Da and a parent ion tolerance of 30

ppm were used. Up to two missed tryptic cleavages were allowed. Methionine oxidation

(+15.99492 Da), cysteine carbamidomethylation (+57.021465 Da), and acetylation (+42.01057

Da; for serines and threonines only) were set as variable modifications. Additional description of

these data is provided in S1 File.

614 Construction of yeast strains enabling three-hybrid and four-hybrid interaction analyses

Derivatives of plasmid pBA2262 used to generate yeast strains with chromosomally-

integrated copies of *hopZ1a^{wt}, hopZ1a^{C216A}, hopZ1b, hopZ2* and *hopZ3* have been described

previously [60]. Briefly, genes cloned into pBA2262 are under the control of the *GAL* promoter

(which is positively regulated by galactose), they are linked to the downstream *NAT^R* gene

(which provides resistance to nourseothricin, also known as clonNAT), and are flanked by 5' and

3' fragments of the *HO* gene (which encodes the homing endonuclease required for mating-type

switching in wild yeast but is dispensable and inactivated - i.e. *hoΔ* - in domesticated laboratory

strains). pBA2262 cannot replicate in yeast so pBA2262 derivatives are linearized by digestion

with the restriction enzyme NotI prior to transformation, and chromosomal integrants resulting

from double recombination events that replace the endogenous *hoΔ* locus with the gene of

interest are isolated by selection on YPDA plates (yeast extract, peptone, dextrose, adenine

sulfate) containing clonNAT at 100 µg/mL.

The *avrAC* coding sequence was amplified from pUC19-35S-*avrAC*-HA (plasmid

generously provided by Jian-Min Zhou; Chinese Academy of Sciences, Beijing) to first create

pDONR207-*avrAC* with BP clonase (Invitrogen), which in turn allowed subsequent creation of

pBA2262-*avrAC* with LR clonase (Invitrogen). Similarly, *ZED1^{wt}* and *ZED1^{N173D V212T}* were

shuttled from pDONR207-based plasmids to pBA2262 using LR clonase.

Assessing bait-prey interactions in yeast

The haploid yeast strain EGY48 ('alpha' mating type; i.e. *MAT* α) and derivative strains

bearing chromosomally-integrated additional genes (encoding bacterial effectors or alleles of the

Arabidopsis pseudokinase ZED1) were transformed with query bait plasmids (pEG202

derivatives; *HIS^+*), and transformants were isolated by selecting for prototrophy on Synthetic

Defined minimal media containing 2% glucose and lacking histidine (SD +Glc -His). Similarly,

haploid yeast strain RFY206 ('A' mating type; i.e. *MAT* **A**) carrying the *lacZ*-bearing reporter

plasmid pSH18-34 (*URA^+*) and a derivative strain bearing chromosomally-integrated *ZED1^{wt}*

were both transformed with prey plasmids (pJG4-5; *TRP^+*), and transformants were isolated by

selecting for prototrophy on SD glucose media lacking uracil and tryptophan (SD +Glc -Ura -

Trp). 12 x 8 arrays of bait (*MAT* α) and prey (*MAT* **A**) transformants were first arranged

manually on appropriate selective media, but all subsequent array manipulations were performed

using a 96-pin replicator. Diploid strains carrying bait, prey and reporter plasmids were created

by co-incubation of bait and prey arrays on YPDA media (8-18 h), followed by two selections on

SD glucose media lacking histidine, uracil and tryptophan (SD +Glc -His -Ura -Trp). Bait-prey

interactions were assessed on Synthetic Defined minimal media reporter plates containing 2 %

raffinose, 1 % galactose, sodium phosphate (0.05 *M*, pH=7.0), X-gal (10 mg/mL) and lacking

histidine, uracil and tryptophan (SD +Raf +Gal +X-gal -His -Ura -Trp). All yeast plates

contained 2 % agar and were incubated at 30^{o}C.

Acquisition of yeast plate images and image processing

Reporter plates were imaged from the bottom using a flat-bed scanner (Epson) against a

black felt background for contrast. The graphical summaries of the yeast interaction data shown

in Fig 1-3, Fig 5, S4 Fig, S5 Fig, S8 Fig, S9 Fig, and S11 Fig were prepared by finding the

average colour for an 81-pixel square (i.e. 9 pixels x 9 pixels) at the center of each colony.

Averaged pixel intensities from yeast interaction plates were extracted by defining array

positions with an interactive interface implemented in PYTHON (www.python.org) using

PROCESSING (www.processing.org; py.processing.org). Colours of text labels (white or black) for

the prey array interaction summaries accompanying yeast reporter plate images (and for the

leaves on the phylogenetic tree shown in Fig 3B) were determined according to a threshold based

on the relative intensities of the red, green and blue channels for the 9 x 9 averaged pixel:

specifically, the value of the ratio blue/(red + green) was evaluated at each PBL array position,

and positions where this ratio was ≥ 0.727 (relative interaction strength ≥ 0.455; i.e. mostly

blue; strong interactions) were assigned white labels, while all other positions were labeled in

black (S2 Fig). This value was used as a working threshold for distinguishing strong interactions

from the background since it bisects the path described by the total interaction dataset presented

in this study (S2 Fig; S15). Pixel plotting and analysis was implemented in PYTHON using

MATPLOTLIB [62]. Relevant scripts are available at https://github.com/DSGlab/Yeast-Array-

Analysis.

Creation of plant expression vectors

For construction of dexamethasone-inducible plant expression vectors, we cloned coding

sequences (lacking stop codons) for *hopZ1a* and for wild-type and mutant *ZED1* alleles into

pMAC14, which includes a dexamethasone-responsive promoter upstream of a multiple cloning

site, as well as linked genes encoding resistance to kanamycin and glufosinate (also known as

BASTA) [63]. *hopZ1a* and *ZED1* alleles were cloned between XhoI and StuI restriction sites

present in the pMAC14 multiple cloning site which places the coding sequence in frame with a

downstream, HA-tag-encoding sequence (and stop codon). *Agrobacterium* strain GV3101 was

transformed according to standard methods [64], and transformants were selected by screening

for plasmid-encoded kanamycin resistance.

Plant transformations and expression assays

We used the floral dip method [65] to generate germ-line transformants of *Arabidopsis*

using *Agrobacterium* strains transformed with pMAC14 derivatives. *Arabidopsis* transformants

were selected by germinating seeds on soil infused with 0.1% BASTA (v/v in water).

Transformed (BASTA-resistant) seedlings were then transplanted into fresh soil free of herbicide

along with control wild-type *Arabidopsis* (Col-0), third-generation (T3) AvrRpt2-expressing

plants, or *zar1-1* mutants. Transgene expression was induced by spraying each flat with ~ 30 mL

of 20 μM dexamethasone in ddH$_2$O with 0.01% surfactant Silwet-L77.

Leaves of *N. benthamiana* plants were locally transformed by pressure infiltration with

suspensions of pMAC14-transformed *Agrobacterium* strains using a needleless syringe applied

to the underside of the leaf. Transgene expression was induced by spraying whole leaves with 20

μM dexamethasone in water with 0.01% surfactant Silwet-L77 (8 - 24h following infiltrations).

Gene silencing was performed as described previously [33–35], using *Agrobacterium* strains

generously provided by Maël Baudin and Jennifer D. Lewis (University of California, Berkeley).

Preparation of protein extracts from plant tissue and immunoblot analysis

Expression of transgenes in *Arabidopsis* was assessed by punching three leaf cores from

a single leaf which were then floated on 20 µM dexamethasone solution (in ddH_2O) for ~18h.

Expression of transgenes in *N. benthamiana* was induced by spraying infiltrated plants with a

solution of 20 µM dexamethasone in ddH_2O with 0.01% Silwet L-77. For each transformation of

interest, tissue was harvested by punching three circular leaf cores 5 mm in diameter. These

tissue samples were then flash-frozen in liquid nitrogen before manual grinding with a mini-

pestle in a 1.5 mL Eppendorf tube. Powdered frozen plant tissue was then suspended in 100 µL

of Grinding Buffer (40 mM Tris pH=7.5, 150 mM NaCl, 1 mM EDTA, 5 mM DTT, 1% Triton

X-100, 0.1% sodium dodecyl sulfate) supplemented with a 1:500 dilution of plant protease

inhibitor cocktail P9599 (Sigma). 15 uL from each of these extracts was then resolved by

electrophoresis through 10 % polyacrylamide SDS-PAGE gels prior to protein transfer to

nitrocellulose membranes. Membranes were blocked with 5% powdered skim milk solution in

TBST (50 mM Tris-Cl, pH 7.5; 150 mM NaCl; 0.05% Tween-20). Primary (rat anti-HA, mouse

anti-LexA, and mouse anti-FLAG) and secondary antibodies (alkaline peroxidase-conjugated

goat anti-mouse) were both diluted 1:10,000 in TBST with 3% powdered milk.

Phylogenetic analysis of kinase domains

Sequences for 46 PBL kinases and 12 ZED1-related pseudokinases were retrieved from

TAIR (www.arabidopsis.org), aligned with Muscle [66], and trimmed to remove variable-length

amino-terminal and carboxy-terminal sequences flanking the conserved kinase domain. These

isolated kinase domain sequences were then supplemented with sequences from structurally-

characterized kinase domains of the *Arabidopsis* receptor kinases BRI1 (PDB: 5LPV) [29] and

BAK1 (PDB: 3UIM) [30], the tomato Pto kinase (PDB: 3HGK) [67], and the human interleukin-

1 receptor-associated kinase IRAK4 (PDB: 4U97) [45]. These 62 sequences were then realigned

with MUSCLE and any amino-terminal or carboxy-terminal sequences flanking the kinase domain

were again manually trimmed. The resulting alignment is provided as a supplementary file (S2

File) and was used as input for phylogenetic inference using PhyML [68,69]. Confidence in

nodes of the resulting phylogenetic tree was assessed using the approximate Likelihood Ratio

Test (aLRT) [70,71]. A representation of the resulting tree (rooted on IRAK4 and including only

those nodes with greater than 70% confidence) is presented in S6 Fig, panel A, and the input tree

is provided as a supplementary file (S3 File). A derivative of the same tree is also shown in Fig

3B and is the result of a pruning step to remove any non-PBL leaves. Kinase domain trees were

visualized using iTOL [72].

Analysis of ZED1 activation loop sequences from diverse *Arabidopsis* ecotypes

The *ZED1* sequences for available *Arabidopsis* ecotypes were derived from DNA SNP

information curated by the 1001 genomes project (http://1001genomes.org/) [73] and were

annotated using MAKER (http://www.yandell-lab.org/software/maker.html) [74]. 1121 translated

ZED1 sequences were collapsed into 813 unique sequences and aligned using MUSCLE [66] (S4

File). 25 of these unique sequences represent more than one *Arabidopsis* ecotype, while the

remaining 788 unique sequences are 'singletons' representing a single *Arabidopsis* ecotype (S5

File). Although confidence in translated ZED1 sequences depends on DNA sequence quality

(which varies among individual sequenced ecotypes), 92.5% of the unique ZED1 sequences

(mean un-gapped sequence length of 332.7 amino acids) had fewer than 38 positions translated

as an 'X' due to ambiguous DNA sequence information (i.e. $\leq 11.4\%$ of the mean sequence

length), and 76.3% of unique sequences had fewer than 23 positions translated as X (≤ 6.9% of

the mean sequence length) (S18 Fig, panels A, B, top left). A cropped alignment consisting of

only the activation loop sequences (mean un-gapped sequence length of 29.0 amino acids) had a

similar distribution of X position prevalence, with 89.0% of activation loop sequences containing

two or fewer X positions (≤ 6.9% of mean activation loop length) (S18 Fig, panels A, B, bottom

left). In contrast, the distributions of serine and threonine positions (S+T) were markedly

different when comparing the activation loop region with full-length protein sequences. While

88.6% of the unique *Arabidopsis* ecotype ZED1 sequences have at least 36 serine or threonine

positions (≥ 10.8% of the mean sequence length) across their full length (S18 Fig, panels A, B,

top right), the vast majority of ecotypes (99.5%) lack any serine or threonine residues when

considering activation loop sequences alone (S18 Fig, panels A, B, bottom right). Notably, the

four ecotypes with threonines in their activation loop sequences are all outliers with respect to

sequence quality, with either three or six ambiguous positions translated as X (sequence

ambiguity that is representative of ≤ 7% of all unique sequences; S18 Fig), so it is possible that

these rare apparent phospho/acetyl-accepting activations residues are in fact artefacts resulting

from poor sequence quality.

Molecular modeling

All three-dimensional protein structural models were prepared using the interface to

MODELLER [75] provided within UCSF CHIMERA [76]. Model templates were selected based on

high-scoring BLASTP hits in the PDB database. The ZED1 model presented in S7 Fig and S10

Fig was constructed using the multiple template option of MODELLER and is based on homology

with both *Solanum pimpinellifolium* (tomato) Pto kinase (PDB: 3HGK; 30% identity spanning

81% of query; E-value: 4e-31) [67] and the kinase domain of *Arabidopsis* receptor kinase BRI1

(PDB: 5LPV; 30% identity spanning 66% of the query; E-value: 5e-27) [29]. The PBS1 model in

S10 Fig was created much like the ZED1 model but using structures of BIK1 (PDB: 5TOS; 48%

identity spanning 70% of query; E-value: 1e-97) [39] and BRI1 (PDB: 5LPV; 44% identity

spanning 62% of query; E-value: 1e-71) [29] as model-building templates. Modeling was based

on two templates so as to generate a model reliably representing the entire kinase domain. For

the hypothetical asymmetric ZED1-PBS1 heterodimer shown in S16 Fig, IRAK4 (PDB: 4U97)

[45] was used as model template for both ZED1 (30% identity across 63% of query sequence; E-

value 7e-21) and PBS1 (42% identical across 62% of query sequence; E-value 8e-56) in addition

to the top-scoring BLASTP hits for ZED1 (Pto) and PBS1 (BIK1) described above. In this case,

however, activation loop residues from Pto and BIK1 were deleted from input alignments prior

to modeling of ZED1 and PBS1 so as to force modeled activation loops to be constrained only

by those from chains A and B from PDB entry 4U97 (IRAK4).

Plant imaging and image processing

Photographs were acquired with a Nikon D5200 DSLR camera. We have previously

described an ImageJ macro (PIDIQ, for <u>P</u>lant <u>I</u>mmunity and <u>D</u>isease <u>I</u>mage-based

<u>Q</u>uantification) that quantifies pixel areas corresponding to green or yellow-shaded regions of

input images [77]. Here we used a similar approach to replicate the behaviour of a factory-

installed camera filter that serendipitously produces images with pixels representing dead/dying

plant tissue and surrounding soil converted to grayscale, while those pixels representing healthy

Arabidopsis leaf tissue remain unchanged. We determined that this behavior can be replicated by

applying logic that outputs a full-colour, RGB pixel where the log-tranformed ratio between

pixel intensities for red and green channels ($\log_{10} \frac{red}{green}$) is ≤ -0.07, and a grayscale pixel where

this condition is not satisfied (S19 Fig). Pixel analysis and image processing were implemented

in PYTHON using MATPLOTLIB [62] and a script is available at https://github.com/DSGlab/Live-

Dead-Filter.

ACKNOWLEDGEMENTS

The authors wish to thank members of the Guttman and Desveaux labs for feedback and

critical discussion of these data, with special thanks to Timothy Lo for discovery of the

'Live/Dead' camera filter. We are also grateful to Maël Baudin and Jennifer D. Lewis

(University of California, Berkeley) for providing strains and advice for *ZAR1* gene-silencing,

and to Jian-Min Zhou (Chinese Academy of Sciences, Beijing) for providing pUC19-35S-

avrAC-HA.

REFERENCES

1. Jones JDG, Dangl JL. The plant immune system. Nature. 2006;444: 323–329. doi:10.1038/nature05286

2. Leipe DD, Koonin E V., Aravind L. STAND, a class of P-loop NTPases including animal and plant regulators of programmed cell death: Multiple, complex domain architectures, unusual phyletic patterns, and evolution by horizontal gene transfer. J Mol Biol. 2004;343: 1–28. doi:10.1016/j.jmb.2004.08.023

3. Duxbury Z, Ma Y, Furzer OJ, Huh SU, Cevik V, Jones JDG, et al. Pathogen perception by NLRs in plants and animals: Parallel worlds. BioEssays. 2016;38: 769–781. doi:10.1002/bies.201600046

4. Jones JDG, Vance RE, Dangl JL. Intracellular innate immune surveillance devices in plants and animals. Science (80-). 2016;354: aaf6395-aaf6395. doi:10.1126/science.aaf6395

5. Meunier E, Broz P. Evolutionary Convergence and Divergence in NLR Function and Structure. Trends Immunol. Elsevier Current Trends; 2017;38: 744–757. doi:10.1016/J.IT.2017.04.005

6. Qi D, Innes RW. Recent Advances in Plant NLR Structure, Function, Localization, and Signaling. Front Immunol. Frontiers; 2013;4: 348. doi:10.3389/fimmu.2013.00348

7. Khan M, Subramaniam R, Desveaux D. Of guards, decoys, baits and traps: Pathogen perception in plants by type III effector sensors. Curr Opin Microbiol. Elsevier Ltd; 2016;29: 49–55. doi:10.1016/j.mib.2015.10.006

8. Shao F, Golstein C, Ade J, Stoutemyer M, Dixon JE, Innes RW. Cleavage of Arabidopsis PBS1 by a Bacterial Type III Effector. Science (80-). 2003;301: 1230–1233. doi:10.1126/science.1085671

9. Ade J, DeYoung BJ, Golstein C, Innes RW. Indirect activation of a plant nucleotide binding site-leucine-rich repeat protein by a bacterial protease. Proc Natl Acad Sci U S A. 2007;104: 2531–2536. doi:10.1073/pnas.0608779104

10. Zhang J, Li W, Xiang T, Liu Z, Laluk K, Ding X, et al. Receptor-like cytoplasmic kinases integrate signaling from multiple plant immune receptors and are targeted by a Pseudomonas syringae effector. Cell Host Microbe. Elsevier Inc.; 2010;7: 290–301. doi:10.1016/j.chom.2010.03.007

11. Lewis JD, Wu R, Guttman DS, Desveaux D. Allele-specific virulence attenuation of the Pseudomonas syringae HopZ1a type III effector via the Arabidopsis ZAR1 resistance protein. PLoS Genet. 2010;6: 1–13. doi:10.1371/journal.pgen.1000894

12. Wang G, Roux B, Feng F, Guy E, Li L, Li N, et al. The Decoy Substrate of a Pathogen Effector and a Pseudokinase Specify Pathogen-Induced Modified-Self Recognition and Immunity in Plants. Cell Host Microbe. 2015;18: 285–295. doi:10.1016/j.chom.2015.08.004

13. Seto D, Koulena N, Lo T, Menna A, Guttman DS, Desveaux D. Expanded type III
effector recognition by the ZAR1 NLR protein using ZED1-related kinases. Nat Plants.
2017;3: 17027. doi:10.1038/nplants.2017.27

14. Lewis JD, Lee AH-Y, Hassan J a, Wan J, Hurley B, Jhingree JR, et al. The *Arabidopsis*
ZED1 pseudokinase is required for ZAR1-mediated immunity induced by the
Pseudomonas syringae type III effector HopZ1a. Proc Natl Acad Sci U S A. 2013;110:
18722–7. doi:10.1073/pnas.1315520110

15. Lewis JD, Lo T, Bastedo P, Guttman DS, Desveaux D. The rise of the undead:
pseudokinases as mediators of effector-triggered immunity. Plant Signal Behav. 2014;9:
e27563. doi:10.4161/psb.27563

16. Roux F, Noël L, Rivas S, Roby D. ZRK atypical kinases: Emerging signaling components
of plant immunity. New Phytol. 2014;203: 713–716. doi:10.1111/nph.12841

17. Hanks S, Quinn A, Hunter T. The protein kinase family: conserved features and deduced
phylogeny of the catalytic domains. Science (80-). 1988;241: 42–52.
doi:10.1126/science.3291115

18. Hanks SK, Hunter T. Protein kinases 6. The eukaryotic protein kinase superfamily: kinase
(catalytic) domain structure and classification. FASEB J. 1995;9: 576–96. doi:7768349

19. Kornev AP, Taylor SS. Dynamics-Driven Allostery in Protein Kinases. Trends Biochem
Sci. Elsevier Current Trends; 2015;40: 628–647. doi:10.1016/J.TIBS.2015.09.002

20. Kornev AP, Taylor SS. Defining the conserved internal architecture of a protein kinase.
Biochim Biophys Acta - Proteins Proteomics. Elsevier B.V.; 2010;1804: 440–444.
doi:10.1016/j.bbapap.2009.10.017

21. Taylor SS, Kornev AP. Protein kinases: Evolution of dynamic regulatory proteins. Trends
Biochem Sci. Elsevier Ltd; 2011;36: 65–77. doi:10.1016/j.tibs.2010.09.006

22. Lavoie H, Li JJ, Thevakumaran N, Therrien M, Sicheri F. Dimerization-induced allostery
in protein kinase regulation. Trends Biochem Sci. 2014;39: 475–486.
doi:10.1016/j.tibs.2014.08.004

23. Boudeau J, Miranda-Saavedra D, Barton GJ, Alessi DR. Emerging roles of pseudokinases.
Trends Cell Biol. 2006;16: 443–452. doi:10.1016/j.tcb.2006.07.003

24. Castells E, Casacuberta JM. Signalling through kinase-defective domains: The prevalence
of atypical receptor-like kinases in plants. Journal of Experimental Botany. 2007.
doi:10.1093/jxb/erm226

25. Eyers PA, Murphy JM. Dawn of the dead: protein pseudokinases signal new adventures in
cell biology. Biochem Soc Trans. 2013;41: 969–74. doi:10.1042/BST20130115

26. Kung JE, Jura N. Structural Basis for the Non-catalytic Functions of Protein Kinases.
Structure. Cell Press; 2016;24: 7–24. doi:10.1016/J.STR.2015.10.020

27. Müller R, Bleckmann A, Simon R. The receptor kinase CORYNE of Arabidopsis
transmits the stem cell-limiting signal CLAVATA3 independently of CLAVATA1. Plant

Cell. American Society of Plant Biologists; 2008;20: 934–46. doi:10.1105/tpc.107.057547

28.   Hurley BA. Development of a Proteomics Platform for Analysis of the Plant-pathogen
880   Interface (PhD Thesis) [Internet]. University of Toronto. 2015. Available:
881   http://hdl.handle.net/1807/70800

882   29.   Bojar D, Martinez J, Santiago J, Rybin V, Bayliss R, Hothorn M. Crystal structures of the
883   phosphorylated BRI1 kinase domain and implications for brassinosteroid signal initiation.
884   Plant J. 2014;78: 31–43. doi:10.1111/tpj.12445

885   30.   Yan L, Ma Y, Liu D, Wei X, Sun Y, Chen X, et al. Structural basis for the impact of
886   phosphorylation on the activation of plant receptor-like kinase BAK1. Cell Res. 2012;22:
887   1304–1308. doi:10.1038/cr.2012.74

888   31.   Sukarta OCA, Slootweg EJ, Goverse A. Structure-informed insights for NLR functioning
889   in plant immunity. Semin Cell Dev Biol. Elsevier Ltd; 2016;56: 134–149.
890   doi:10.1016/j.semcdb.2016.05.012

891   32.   McNellis TW, Mudgett MB, Li K, Aoyama T, Horvath D, Chua NH, et al.
892   Glucocorticoid-inducible expression of a bacterial avirulence gene in transgenic
893   Arabidopsis induces hypersensitive cell death. Plant J. 1998;14: 247–57.
894   doi:10.1046/j.1365-313X.1998.00106.x

895   33.   Baudin M, Hassan JA, Schreiber KJ, Lewis JD. Analysis of the ZAR1 Immune Complex
896   Reveals Determinants for Immunity and Molecular Interactions. Plant Physiol. 2017;174:
897   2038–2053. doi:10.1104/pp.17.00441

898   34.   Velásquez AC, Chakravarthy S, Martin GB. Virus-induced gene silencing (VIGS) in
899   Nicotiana benthamiana and tomato. J Vis Exp. MyJoVE Corporation; 2009;
900   doi:10.3791/1292

901   35.   Zhou B, Zeng L. Elucidating the role of highly homologous Nicotiana benthamiana
902   ubiquitin E2 gene family members in plant immunity through an improved virus-induced
903   gene silencing approach. Plant Methods. 2017; doi:10.1186/s13007-017-0210-6

904   36.   Lu D, Wu S, Gao X, Zhang Y, Shan L, He P. A receptor-like cytoplasmic kinase, BIK1,
905   associates with a flagellin receptor complex to initiate plant innate immunity. Proc Natl
906   Acad Sci U S A. 2010;107: 496–501. doi:10.1073/pnas.0909705107

907   37.   Lin W, Li B, Lu D, Chen S, Zhu N, He P, et al. Tyrosine phosphorylation of protein
908   kinase complex BAK1/BIK1 mediates Arabidopsis innate immunity. Proc Natl Acad Sci
909   U S A. National Academy of Sciences; 2014;111: 3632–7. doi:10.1073/pnas.1318817111

910   38.   Li L, Li M, Yu L, Zhou Z, Liang X, Liu Z, et al. The FLS2-associated kinase BIK1
911   directly phosphorylates the NADPH oxidase RbohD to control plant immunity. Cell Host
912   Microbe. Elsevier Inc.; 2014;15: 329–338. doi:10.1016/j.chom.2014.02.009

913   39.   Lal NK, Nagalakshmi U, Hurlburt NK, Flores R, Bak A, Sone P, et al. The Receptor-like
914   Cytoplasmic Kinase BIK1 Localizes to the Nucleus and Regulates Defense Hormone
915   Expression during Plant Innate Immunity. Cell Host Microbe. 2018;23: 485–497.e5.
916   doi:10.1016/j.chom.2018.03.010

40. Liu J, Elmore JM, Lin ZJD, Coaker G. A receptor-like cytoplasmic kinase phosphorylates
the host target RIN4, leading to the activation of a plant innate immune receptor. Cell
Host Microbe. Elsevier Inc.; 2011;9: 137–146. doi:10.1016/j.chom.2011.01.010

41. Xu N, Luo X, Li W, Wang Z, Liu J. The Bacterial Effector AvrB-Induced RIN4
Hyperphosphorylation Is Mediated by a Receptor-Like Cytoplasmic Kinase Complex in
Arabidopsis. Mol Plant-Microbe Interact. Molecular Plant-Microbe Interactions; 2017;30:
502–512. doi:10.1094/MPMI-01-17-0017-R

42. Lin Z-JD, Liebrand TWH, Yadeta KA, Coaker GL. PBL13 is a serine/threonine protein
kinase that negatively regulates Arabidopsis immune responses. Plant Physiol. 2015;169:
pp.01391.2015. doi:10.1104/pp.15.01391

43. Shinya T, Yamaguchi K, Desaki Y, Yamada K, Narisawa T, Kobayashi Y, et al. Selective
regulation of the chitin-induced defense response by the Arabidopsis receptor-like
cytoplasmic kinase PBL27. Plant J. Wiley/Blackwell (10.1111); 2014;79: 56–66.
doi:10.1111/tpj.12535

44. Yamada K, Yamaguchi K, Shirakawa T, Nakagami H, Mine A, Ishikawa K, et al. The
Arabidopsis CERK1-associated kinase PBL27 connects chitin perception to MAPK
activation. EMBO J. EMBO Press; 2016;35: 2468–2483. doi:10.15252/embj.201694248

45. Ferrao R, Zhou H, Shan Y, Liu Q, Li Q, Shaw DE, et al. IRAK4 Dimerization and trans-
Autophosphorylation Are Induced by Myddosome Assembly. Mol Cell. Elsevier Inc.;
2014;55: 891–903. doi:10.1016/j.molcel.2014.08.006

46. Feng F, Yang F, Rong W, Wu X, Zhang J, Chen S, et al. A Xanthomonas uridine 5′-
monophosphate transferase inhibits plant immune kinases. Nature. Nature Publishing
Group; 2012;485: 114–118. doi:10.1038/nature10962

47. Wang G, Roux B, Feng F, Guy E, Li L, Li N, et al. The decoy substrate of a pathogen
effector and a pseudokinase specify pathogen-induced modified-self recognition and
immunity in plants. Cell Host Microbe. 2015;18: 285–295. Available:
http://www.sciencedirect.com/science/article/pii/S1931312815003327

48. Mittal R, Peak-Chew S-Y, McMahon HT. Acetylation of MEK2 and I kappa B kinase
(IKK) activation loop residues by YopJ inhibits signaling. Proc Natl Acad Sci U S A.
2006;103: 18574–9. doi:10.1073/pnas.0608995103

49. Mukherjee S, Keitany G, Li Y, Wang Y, Ball HL, Goldsmith EJ, et al. Yersinia YopJ
acetylates and inhibits kinase activation by blocking phosphorylation. Science. 2006;312:
1211–4. doi:10.1126/science.1126867

50. Nolen B, Taylor S, Ghosh G. Regulation of protein kinases: Controlling activity through
activation segment conformation [Internet]. Molecular Cell Sep 10, 2004.
doi:10.1016/j.molcel.2004.08.024

51. Endicott JA, Noble MEM, Johnson LN. The Structural Basis for Control of Eukaryotic
Protein Kinases. Annu Rev Biochem. 2012;81: 587–613. doi:10.1146/annurev-biochem-
052410-090317

52. Ma K, Ma W. YopJ family effectors promote bacterial infection through a unique

acetyltransferase activity. Microbiol Mol Biol Rev. 2016;80: 1011–1027.
doi:10.1128/MMBR.00032-16.Address

53.   Jayaraman J, Choi S, Prokchorchik M, Choi DS, Spiandore A, Rikkerink EH, et al. A
bacterial acetyltransferase triggers immunity in Arabidopsis thaliana independent of
hypersensitive response. Sci Rep. 2017;7: 3557. doi:10.1038/s41598-017-03704-x

54.   Lewis JD, Abada W, Ma W, Guttman DS, Desveaux D. The HopZ family of
Pseudomonas syringae type III effectors require myristoylation for virulence and
avirulence functions in Arabidopsis thaliana. J Bacteriol. 2008;190: 2880–2891.
doi:10.1128/JB.01702-07

55.   Lee J, Manning AJ, Wolfgeher D, Jelenska J, Cavanaugh KA, Xu H, et al. Acetylation of
an NB-LRR Plant Immune-Effector Complex Suppresses Immunity. Cell Rep. The
Authors; 2015;13: 1670–1682. doi:10.1016/j.celrep.2015.10.029

56.   Schultink A, Qi T, Bally J, Staskawicz B. Using forward genetics in Nicotiana
benthamiana to uncover the immune signaling pathway mediating recognition of the
Xanthomonas perforans effector XopJ4. New Phytol. 2018; doi:10.1111/nph.15411

57.   Gietz RD, Schiestl RH. Quick and easy yeast transformation using the LiAc/SS carrier
DNA/PEG method. Nat Protoc. 2007;2: 35–37. doi:10.1038/nprot.2007.14

58.   Gietz RD, Schiestl RH. Frozen competent yeast cells that can be transformed with high
efficiency using the LiAc/SS carrier DNA/PEG method. Nat Protoc. 2007;2: 1–4.
doi:10.1038/nprot.2007.17

59.   Youn J-Y, Friesen H, Nguyen Ba AN, Liang W, Messier V, Cox MJ, et al. Functional
Analysis of Kinases and Transcription Factors in Saccharomyces cerevisiae Using an
Integrated Overexpression Library. G3 Genes, Genomes, Genet. 2017;
doi:10.1534/g3.116.038471

60.   Lee AH-Y, Bastedo DP, Youn J-Y, Lo T, Middleton MA, Kireeva I, et al. Identifying
Pseudomonas syringae Type III Secreted Effector Function via a Yeast Genomic Screen.
G3 Genes, Genomes, Genet. Genetics Society of America; 2018; g3.200877.2018.
doi:10.1534/g3.118.200877

61.   Eng J, McCormack A, Yates J. An approach to correlate tandem mass spectral data of
peptides with amino acid sequences in a protein database. J Am Soc Mass Spectrom.
1994;5: 976–89. doi:doi:10.1016/1044-0305(94)80016-2

62.   Hunter JD. Matplotlib: A 2D graphics environment. Comput Sci Eng. 2007;
doi:10.1109/MCSE.2007.55

63.   Aoyama T, Chua NH. A glucocorticoid-mediated transcriptional induction system in
transgenic plants. Plant J. 1997;11: 605–612. doi:10.1046/j.1365-313X.1997.11030605.x

64.   Höfgen R, Willmitzer L. Storage of competent cells for Agrobacterium transformation.
Nucleic Acids Res. Oxford University Press; 1988;16: 9877. Available:
http://www.ncbi.nlm.nih.gov/pubmed/3186459

65.   Clough SJ, Bent AF. Floral dip: A simplified method for Agrobacterium-mediated

transformation of Arabidopsis thaliana. Plant J. 1998;16: 735–743. doi:10.1046/j.1365-
313X.1998.00343.x

66. Edgar RC. MUSCLE: Multiple sequence alignment with high accuracy and high
throughput. Nucleic Acids Res. 2004;32: 1792–1797. doi:10.1093/nar/gkh340

67. Dong J, Xiao F, Fan F, Gu L, Cang H, Martin GB, et al. Crystal structure of the complex
between Pseudomonas effector AvrPtoB and the tomato Pto kinase reveals both a shared
and a unique interface compared with AvrPto-Pto. Plant Cell. 2009;21: 1846–1859.
doi:10.1105/tpc.109.066878

68. Guindon S, Gascuel O. A Simple, Fast, and Accurate Algorithm to Estimate Large
Phylogenies by Maximum Likelihood. Syst Biol. 2003;52: 696–704.
doi:10.1080/10635150390235520

69. Guindon S, Lethiec F, Duroux P, Gascuel O. PHYML Online - A web server for fast
maximum likelihood-based phylogenetic inference. Nucleic Acids Res. 2005;33.
doi:10.1093/nar/gki352

70. Anisimova M, Gascuel O. Approximate likelihood-ratio test for branches: A fast,
accurate, and powerful alternative. Syst Biol. 2006;55: 539–552.
doi:10.1080/10635150600755453

71. Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms
and methods to estimate maximum-likelihood phylogenies: Assessing the performance of
PhyML 3.0. Syst Biol. 2010;59: 307–321. doi:10.1093/sysbio/syq010

72. Letunic I, Bork P. Interactive tree of life (iTOL) v3: an online tool for the display and
annotation of phylogenetic and other trees. Nucleic Acids Res. 2016;
doi:10.1093/nar/gkw290

73. Weigel D, Mott R. The 1001 genomes project for Arabidopsis thaliana. Genome Biol.
2009;10: 107. doi:10.1186/gb-2009-10-5-107

74. Campbell MS, Holt C, Moore B, Yandell M. Genome Annotation and Curation Using
MAKER and MAKER-P. Curr Protoc Bioinforma. 2014;
doi:10.1002/0471250953.bi0411s48

75. Andrej Šali. Comparative protein modelling by satisfction of spatial restraints. J Mol
Biol. 1993;234: 779–815.

76. Meng EC, Pettersen EF, Couch GS, Huang CC, Ferrin TE. Tools for integrated sequence-
structure analysis with UCSF Chimera. BMC Bioinformatics. 2006;7: 339.
doi:10.1186/1471-2105-7-339

77. Laflamme B, Middleton M, Lo T, Desveaux D, Guttman DS. Image-Based Quantification
of Plant Immunity and Disease. Mol Plant-Microbe Interact. 2016; doi:10.1094/MPMI-07-
16-0129-TA

78. Zheng J, Trafny EA, Knighton DR, Xuong N, Taylor SS, Ten Eyck LF, et al. 2.2 Å
refined crystal structure of the catalytic subunit of cAMP-dependent protein kinase
complexed with MnATP and a peptide inhibitor. Acta Crystallogr Sect D Biol Crystallogr.

International Union of Crystallography; 1993;49: 362–365.
doi:10.1107/S0907444993000423

79.   Madhusudan PA, Xuong N-H, Taylor SS. Crystal structure of a transition state mimic of
the catalytic subunit of cAMP-dependent protein kinase. Nat Struct Biol. 2002;9: 273–
277. doi:10.1038/nsb780

80.   Akamine P, Madhusudan, Wu J, Xuong NH, Ten Eyck LF, Taylor SS. Dynamic features
of cAMP-dependent protein kinase revealed by apoenzyme crystal structure. J Mol Biol.
2003;327: 159–171. doi:10.1016/S0022-2836(02)01446-8

81.   Kornev AP, Taylor SS, Ten Eyck LF. A helix scaffold for the assembly of active protein
kinases. Proc Natl Acad Sci U S A. 2008;105: 14377–82. doi:10.1073/pnas.0807988105

1045

 FIGURE CAPTIONS

 Fig 1. HopZ1a binds to PBL kinases and acetylates PBS1.

 (A) Yeast two-hybrid interaction assay using wild-type or catalytically-inactive HopZ1a alleles

 as bait and an array of 46 PBL kinases as prey. Yeast colonies on an X-Gal reporter plate are

 shown on top. A grid describing the PBL array layout is shown below each half of the plate, with

 the colours of array positions representing the averaged colony colour for each bait-prey

 interaction. We used a 'relative interaction strength' metric derived from these averaged colony

 colours to discriminate strong interactions from the background (see S2 Fig; Materials and

 Methods); prey array positions with interaction strength ≥ 0.455 (i.e. strong interactions) were

 assigned white labels while other positions were assigned black labels). The interaction strength

 threshold is indicated in the colour-bar at right with a horizontal white line. Interaction scheme

 refers to S1 Fig. (B) LC-MS/MS identification of PBS1 residues acetylated by HopZ1a. The

 PBS1 protein sequence is indicated as horizontal bars for three experimental replicates in the

 presence of each HopZ1a allele (wild-type HopZ1a or the catalytically-inactive mutant,

 HopZ1a^{C216A}. Black bands indicate peptides that were reliably detected by the mass-spectrometry

 instruments, while peptides not detected with high confidence are indicated by grey shading. (C)

 LC-MS/MS identification of ZED1 residues acetylated by HopZ1a. Peptide detection and

 acetylation sites are presented as for panel B above. Note that in addition to S84, T87 and T177,

 S137 was also acetylated, although this modification was not consistently observed across

 multiple independent experiments.

 Fig 2. HopZ1a promotes binding between ZED1 and several PBL kinases.

 Y3H assays testing interactions between ZED1 and 46 PBLs in the presence of HopZ effector

 alleles are contrasted with a Y2H assay in the absence of effectors. (A) HopZ1a^{wt} (left) or

HopZ1a^{C216A} (center) were expressed from the chromosome (see Materials and Methods) with

ZED1 as bait and 46 individual PBLs as prey. The prey array layout shown at right represents

PBL interactions with ZED1 in the presence of HopZ1a^{wt}. The colours of the labels at each array

position (white or black) are determined by the relative interaction strength (see S2 Fig and

Materials and Methods). Interaction scheme refers to S1 Fig. (B) ZED1 bait / PBL prey

interactions in the absence of chromosomally expressed effector (left) or in the presence of

HopZ1b (center). The prey array layout shown at right represents PBL interactions with ZED1 in

the presence of HopZ1b. Interaction schemes refer to S1 Fig.

Fig 3. ZED1 mutagenesis targets, phylogenetic analysis of PBL kinase domains, and a

summary of ZED1-PBL binding interactions.

(A) A linear representation of the ZED1 sequence showing the positions of mutated sites (labeled

a-e, according to their descriptions in the text), and predicted secondary structure elements are

labeled according to homologous features present in the well-studied kinase, PKA [78–81]. β-

strands are coloured yellow, while α-helices are coloured dark blue, as for the three-dimensional

structural models of ZED1 presented in S7 Fig (panel A), S10 Fig (panel A) and S16 Fig. The

activation loop sequence is highlighted with an orange bracket. (B) *Left* – phylogenetic tree

showing relationships between the kinase domains of RLCK family VII (PBS1-like) kinases.

Labels for tree leaves are coloured according to yeast colony colours from the HopZ1a^{wt}-

dependent interactions presented in Fig 2A. *Right* - condensed graphical summary of the

Y2H/Y3H interaction data presented in Fig 2, S3 Fig, S2 Fig, S6 Fig and S7 Fig. Subsets of the

five ZED1 mutagenesis targets (sites a-e) that are altered in a given ZED1 allele are indicated at

the top of each column. Mutant ZED1 alleles whose expression causes induction of ETI *in planta*

are highlighted with red text. The colour-bar at lower left depicts the range of relative

interactions strengths, as defined in S2 Fig (see Materials and Methods).

Fig 4. Isoleucine substitution of a PBL activation loop phospho-accepting residue mimics

acetylation by HopZ1a.

(A) Comparison of the molecular structures of phospho-accepting residues L-threonine and L-

serine, their acetylated derivatives, candidate acetyl-mimetic residues L-isoleucine and L-

glutamine, and L-alanine, a presumed loss-of-function substitution that should block both

phosphorylation and acetylation. (B) Mutant alleles of PBL15 were tested in both Y2H (ZED1

bait / PBL15 prey) and Y3H contexts (ZED1 bait / PBL15 prey with HopZ1a expression) to

screen for substitutions that confer HopZ1a-independent ZED1 binding activity. Interaction

schemes refer to S1 Fig.

Fig 5. ZAR1-ZED1-PBL ternary complexes are formed in the presence either of HopZ1a

activity or of a mutant ZED1 allele.

(A) Schematic showing ZAR1 domain truncation boundaries. (B) Interactions between ZAR1

bait constructs and 11 PBL preys were assessed in the absence and presence of HopZ1a and/or

ZED1 alleles integrated at the *ho* locus of strains EGY48 (*MAT* α) or RFY206 (*MAT* **A**),

corresponding to Y2H (interaction scheme A), Y3H (interaction schemes B), and Y4H

(interaction scheme C) assays (see S1 Fig). The prey array layout shown at right represents PBL

interactions with ZAR1$^{\Delta CC}$ in the presence of both ZED1 and HopZ1a^{wt}. The colours of the

labels at each array position (white or black) are determined by the relative interaction strength

(see S2 Fig; Materials and Methods).

Fig 6. Dexamethasone-induced expression of ZED1$^{'DT'}$ (N173D V212T) causes whole-plant

HR in *Arabidopsis*.

BASTA-resistant *Arabidopsis* transformants bearing dexamethasone-inducible ZED1 alleles

(columns 4-7) were tested alongside control plants (columns 1-3; untransformed Col-0,

transgenic plants with dexamethasone-inducible AvrRpt2, and untransformed *zar1-1*).

Photographs of control plants and *Arabidopsis* T1 transformants are shown both before (A) and

~72 h after dexamethasone induction of transgenes (B). Panels A and B present full-colour RGB

images (top) as well as a filtered representation ('Live/Dead filter'; bottom) that converts

dead/dying plant tissue and surrounding soil to grayscale pixels while leaving pixels representing

healthy *Arabidopsis* tissue unchanged (see S19 Fig; Materials and Methods). Red numbered

circles indicate those plants for which transgene expression was detected and correspond to the

electrophoresis lane indices shown in panel C, below. Note that plants presented in this Figure

were grown on the same flat and so experienced identical conditions with respect to watering,

lighting, and dexamethasone sprays; whole-flat images were cropped to remove unrelated plants.

(C) Parallel assessment of transgene expression in tissue from the same plants shown in panels A

and B. HA-tagged transgene expression in T1 transformants is shown (top), and Ponceau S

staining of total protein transferred to the nitrocellulose membrane (bottom) demonstrates

consistent sample loading and protein transfer across all lanes.

Fig 7. Virus-induced gene silencing of *NbZAR1* expression abolishes HopZ1a-independent

HR triggered by ZED1 alleles.

Leaves from *N. benthamiana* plants were infected with Tobacco Rattle Virus-derived gene

silencing constructs targeting *GUS* (left) or *ZAR1* (right) two weeks prior to transformation by

localized pressure-infiltration of *Agrobacterium* cell suspensions delivering the indicated

dexamethasone-inducible transgenes. Images show infiltrated leaves ~48 h post-induction by

spray with dexamethasone. Note that the images showing infiltration spot #6 are from separate

leaves but from the same plants as the corresponding infiltration spots #1-5; leaf images were

cropped to remove additional unrelated infiltrations.

Fig 8. Model describing intermolecular interactions necessary for ZAR1-dependent

immune responses in *Arabidopsis*.

Left - An interdomain conformational change in ZAR1 and post-translational modification

(PTM) of ZRKs and/or PBLs are necessary for ternary complex formation and ZAR1-dependent

ETI induction. Active or 'active-like' kinase domain conformations are represented with rounded

rectangles, while inactive and 'inactive-like' kinase domain conformations are represented with

parallelograms. *Right* – Contrasting alternate models explaining activation of ZAR1-dependent

immunity in *Arabidopsis*. (A) In the decoy model, post-translationally-modified ZRKs are

presumed to adopt an 'active-like' conformation that is required to activate ZAR1 and trigger

ETI. (B) The adaptor model proposed by Wang et al. [12] suggests that post-translational

modification of PBLs results in conversion to an inactive conformation that is compatible for

high-affinity interactions with ZRKs (and ZAR1) to activate ETI. (C) An alternative adaptor

model suggested by our data implies that post-translational-modification of ZRK pseudokinases

may be sufficient to recruit PBLs for ZAR1 activation by inducing an 'active-like' pseudokinase

conformation. In other words, models B and C both allow heterodimer formation only when

kinase and pseudokinase have adopted similar structural conformations – inactivated kinases

bind to 'inactive-like' pseudokinases, but similarly, modified pseudokinases adopting 'active-

like' conformations as a result of post-translational modifications can also bind to active

(unmodified) PBLs to activate ZAR1.

 SUPPORTING INFORMATION

 S1 Fig. Yeast two-hybrid, three-hybrid and four-hybrid protein interaction schemes used

 to assess pairwise interactions between HopZ1a, ZED1/ZRKs, PBLs, and ZAR1.

(A) Y2H: Transformants of haploid yeast strain EGY48 (*MAT α*) carrying pEG202 bait fusion

plasmids (in this example, pEG202::*lexA^{DBD}-ZED1*) are mated with transformants of haploid

yeast strain RFY206 (*MAT* **A**) carrying pJG4-5 prey fusion plasmids (pJG4-5::*AD-PBL; AD =*

NLS-B42^{AD}-HA) and the reporter plasmid pSH18-34 (*lacZ*). (B) Y3H: Genes of interest (in this

example *hopZ1a*) are integrated at the *ho* locus of the haploid yeast strain EGY48 (*MAT α*;; see

Materials and Methods). These strains were subsequently transformed with pEG202 bait fusion

plasmids (in this example, pEG202::*lexA^{DBD}-ZED1*) and mated with transformants of haploid

yeast strain RFY206 (*MAT* **A**) carrying prey fusion plasmids (pJG4-5::*AD-PBL*) and the reporter

plasmid pSH18-34. (C) Y4H: Transformants of haploid yeast strain EGY48 (*MAT α*) with

integrated *hopZ1a* alleles and carrying pEG202 bait fusion plasmids (pEG202::*lexA^{DBD}-ZAR1*)

are mated with transformants of haploid yeast strain RFY206 (*MAT* **A**) with integrated *ZED1^{wt}*

and carrying prey fusion plasmids (pJG4-5::*AD-PBL*) and the reporter plasmid pSH18-34.

S2 Fig. Pixel analysis to define a 'relative interaction strength' metric.

(A) Averaged RGB pixel values for each of the yeast colonies presented in Fig 2, S4 Fig, S5 Fig,

S8 Fig, and S9 Fig (see Materials and Methods) are plotted as a function of the log-transformed

relative intensities of their blue (x-axis) and green (y-axis) channels. This plotting strategy results

in a smooth arc describing the transition between strong (mostly blue) and weak (yellowish

white) interactions, while spots representing array positions with no yeast colonies (agar plate

only) form a distinct cluster below. For comparison, alternate methods of plotting these same

data are presented in S17 Fig. The data were fitted to a quadratic function of the form $y =$

$a(x + b)^2 + cx + d$ (R^2 = 0.802; $a = -1.511, b = 0.67, c = 1.607, d = 0.456$; plotted as a red

line), and the point at which this curve reaches its maximum ($x = -0.138; \frac{blue}{red+green} = 0.727$)

was used a threshold for discriminating strong interactions from bait-prey pairs with only weak-

to-moderate interaction strength. Spots exceeding this threshold are shown with a white border

and are assigned white labels in the array layouts accompanying Fig 1A, Fig 2, Fig 5, S4 fig, S5

Fig, S8 Fig, S9 Fig, and S11 Fig. (B) Histogram of the x-coordinates of all yeast spots plotted in

panel A (excluding agar-only background spots). A consensus colour for the histogram bars

representing each of 85 bins was determined by finding the average red, green, and blue channel

intensities for all of the spots represented by a given bin. Bins exceeding the threshold described

in panel A are highlighted with a white border. *Inset* - a colour-bar derived from the histogram

data relating averaged pixel colours to a 'relative interaction strength' metric; the log-

transformed relative intensities of the blue channel for each spot ($-0.38 \leq x \leq 0.15$) were

scaled to values between 0 and 1. Empty bins (bins 76, 80, 81, 82, 83, and 84) were assigned the

colour of the immediately-preceding non-zero bin, and the resulting colour range was smoothed

by applying a five-bin sliding window average. The value corresponding to the white/black label

threshold established in panel A (0.455) is indicated with a white horizontal line.

S3 Fig. Peptides/modifications detected by LC-MS/MS at a more permissive threshold.

Peptides and acetylation sites from PBS1 (A) and ZED1 (B) are plotted as in Fig 1BC, except for

the additional inclusion of peptides that only surpass the 'Low' confidence threshold reported by

the Proteome Discoverer^{TM} software, as described in S1 File.

S4 Fig. AvrAC promotes interactions between ZRK1/RKS1 and PBL kinases.

Top - Binding of ZRK1/RKS1 to 46 PBLs was assessed in the absence (Y2H; left) or presence

(Y3H; right) of the *X. campestris* effector AvrAC; interaction schemes refer to S1 Fig. *Bottom* –

Prey array layouts showing the relative strength of interactions between ZRK1 and PBLs,

corresponding to the X-gal reporter plate above. Label colours (white or black) for each array

position are determined by the relative interaction strength (see S2 Fig; Materials and Methods).

Yeast spots and array positions corresponding to the positive control interaction between ZRK1

and PBL2 are highlighted with a red frame.

S5 Fig. PBL-ZED1 interactions in the presence of HopZ2, HopZ3, and AvrAC.

Y2H interactions between ZED1 and PBLs in the absence of effectors (panel B, left) are

compared with Y3H interactions in the presence of the indicated effectors (panel A; panel B,

middle). Prey array layouts at right indicate the relative strength of ZED-PBL interactions in the

presence of HopZ2 (A) and in the absence of effector (B). All array labels are black since the

relative interaction strengths are all below the threshold described in S2 Fig. Interaction schemes

refer to S1 Fig. (C) *Top* - Representative western blots showing consistent expression of both

bait (ZED1), prey (PBS1), and integrated effectors (HopZ alleles). *Bottom* – Nitrocellulose

membranes stained with Ponceau S demonstrate equal loading of yeast cell extracts and

consistent transfer from SDS-PAGE gels across all lanes.

S6 Fig. Degenerate kinase motifs in ZED1 and the ZED1-related kinases (ZRKs).

(A) Phylogenetic tree showing evolutionary relationships between the kinase domains of

ZED1/ZRKs (dark blue shading), PBLs (light blue shading), and sequences from four

structurally-characterized kinase domains - *Arabidopsis* BRI1 (PDB: 5LPV) [29] and BAK1

(PDB: 3UIM) [30], *Solanum pimpinellifolium* (tomato) Pto kinase (PDB: 3HGK) [67], and

human IRAK4 (PDB: 4U97) [45]. Note that PBL28 is an out-group compared to other PBLs and

may therefore not represent a 'true' PBL. (B) A subset of columns from the alignment used to

generate the tree shown in panel A are presented to highlight important kinase motifs present in

structurally-characterized kinase domains (top), ZED1/ZRKs (middle), and a representative

subset of PBLs (bottom). The numbers of omitted, non-gap positions between adjacent blocks of

consecutive columns are shown between square brackets for each sequence. Highlighted in blue

are exact matches to established kinase motifs. Serine and threonine residues between the 'DFG'

and 'APE' motifs that define the activation loop are highlighted in green. Serine/threonine

residues that have been observed in phosphorylated form in previously-described crystal

structures are highlighted in orange. PBS1 S244, a site acetylated by HopZ1a, and immediately

following the site cleaved by HopAR1, is highlighted in red. PBL2 S253 and T254, sites

uridylylated by the *X. campestris* effector AvrAC and required for AvrAC-induced interaction

with ZRK1 [12], are highlighted in purple. ZED1 residues targeted for mutagenesis in this study

are indicated in bold, black font.

S7 Fig. Active site residues in a homology-based model of pseudokinase ZED1 and in the

experimentally-determined structure of receptor kinase BRI1.

(A) Homology-based structural model (see Materials and Methods) of the ZED1 active site

showing the relative positions of the sites targeted for mutagenesis in this study. Blue residue

labels indicate positions that were mutated based on their established importance for kinase

function. Red residue labels (and all-red sidechains) indicate positions acetylated by HopZ1a

(S84, T87, T125, T177). The $\beta3$ lysine, K76 (labeled in grey), was not mutated but its sidechain

is shown for comparison to the salt-bridge-forming lysine (K911) from BRI1 (shown in panel B,

below). Note that in this model W58 and W193 would be expected to clash with the non-

hydrolyzable ATP analogue AMP-PNP and Mn^{2+} ions if directly superimposed from the

template BRI1 structure (shown in panel B). (B) An identical view of the kinase domain active

site showing equivalent positions on the template kinase domain structure of the *Arabidopsis*

brassinosteroid receptor kinase, BRI1 (PDB: 5LPV) [29]. Blue residue labels indicate positions

aligned with those that were mutated in ZED1 based on their established importance for kinase

function. Grey residue labels indicate positions equivalent to ZED1 sites that are acetylated by

HopZ1a, as well as the salt-bridge forming β3 lysine, K911. (Note that Q919 and R922 were

poorly resolved in the BRI1 structure, hence only the alpha-carbons of their sidechains are

visible.) The ATP analogue AMP-PNP is shown as a 'ball and stick' representation, and two

Mn^{2+} ions are shown as purple spheres. Images were prepared with UCSF CHIMERA [76].

S8 Fig. Restoration of degenerate kinase motifs allows promiscuous, HopZ1a-independent

ZED1-PBL interactions.

(A) Protein interaction assays to assess the binding affinities of wild-type ZED1 and ZED1

single mutants (sites 'b', 'c', and 'd' in Fig 3) for 15 PBLs in the absence (Y2H; S1 Fig, panel A)

or presence (Y3H; S1 Fig, panel B) of HopZ1a. The prey array layout at right indicates the

relative strength of PBL interactions with $ZED1^{N173D}$ in the absence of effector. The colours of

array labels at each position (black or white) are determined by the relative interaction strength,

as described in S2 Fig. (B) Double and triple mutant ZED1 alleles combining the mutations

shown in panel A (sites 'bc', 'bd', 'cd', and 'bcd' in Fig 3) were likewise tested against the same

array of 15 PBLs in both Y2H and Y3H contexts. The prey array layout shown at right indicates

the relative strength of PBL interactions with $ZED1^{N173D\ W193G}$ in the absence of effector.

Interaction schemes refer to S1 Fig.

S9 Fig. Restoration of a glycine-rich ZED1 'G-loop' blocks HopZ1a-dependent and

HopZ1a-independent PBL interactions.

(A) Protein interaction assays to assess the binding affinities of $ZED1^{wt}$, single mutants

($ZED1^{N173D}$, $ZED1^{V212T}$; sites 'c', 'e' in Fig 3), and a double mutant, ($ZED1^{"DT"}$; sites 'ce' in Fig

3) for 15 PBLs in Y2H and Y3H contexts. The prey array layout shown at right indicates the

relative strength of PBL interactions with ZED1^{V212T} in the absence of effector. The colours of

array labels at each position (black or white) is determined by the relative interaction strength, as

described in S2 Fig. (B) A ZED1 triple mutant ("3xG"; S56G W58G F61G) was tested against

the same 15 PBLs, either alone or combined with the single and double mutants shown in panel

A. The prey array layout shown at right indicates the relative strength of PBL interactions with

ZED1$^{\text{'3xG' V212T}}$ in the absence of effector. Interaction schemes refer to S1 Fig.

S10 Fig. Glutamine substitutions of HopZ1a acetylation sites block ZED1-PBL

interactions.

(A) Structural models of ZED1 and PBS1 showing the positions of HopZ1a-acetylated residues

as determined by LC-MS/MS. Note that acetylated PBS1 residues T32 and S405 do not form

part of the conserved kinase domain. Images were prepared with UCSF CHIMERA [76]. (B) Y2H

and Y3H assays for protein interactions between seven representative PBL preys and the

indicated ZED1 bait alleles (wild-type, and mutants with glutamine substitutions at acetylation

sites). (C) Y2H and Y3H assays for protein-protein interactions between wild-type ZED1 bait

and the indicated PBS1 prey alleles (wild-type, and mutants with glutamine substitutions at

acetylation sites). Interaction schemes refer to S1 Fig.

S11 Fig. Isolated ZAR1 domains are not sufficient for robust formation of HopZ1a-

dependent ternary complexes with ZED1 and PBLs.

(A) Schematic showing ZAR1 domain truncation boundaries. (B) Interactions between ZAR1

bait constructs and 11 PBL preys were assessed in the absence and presence of HopZ1a and/or

ZED1 alleles integrated at the *ho* locus of strain EGY48 (*MAT* α) or RFY206 (*MAT* **A**),

corresponding to Y2H (interaction scheme A), Y3H (interaction schemes B), and Y4H

(interaction scheme C) assays (S1 Fig). The prey array layout shown at right indicates the

relative strength of PBL interactions with ZAR1^{LRR} in the presence of both ZED1 and HopZ1a^{wt}.

All array labels are black since the relative interaction strengths are all below the threshold

described in S2 Fig.

S12 Fig. Dexamethasone-induced expression of ZED1^{wt} and ZED1^{N173D} alleles does not

induce immunity in *Arabidopsis*.

Control plants and BASTA-resistant *Arabidopsis* transformants bearing dexamethasone-

inducible ZED1 alleles (ZED1^{wt} and ZED1^{N173D}) were photographed before treatment (A) and 72

h post-dexamethasone spray (B) and screened for transgene expression (C), as described for

ZED1^{wt} and ZED1$^{'DT'}$ in Fig 6.

S13 Fig. Dexamethasone-induced expression of ZED1^{V212T} causes whole-plant HR in

***Arabidopsis*.**

Control plants and BASTA-resistant *Arabidopsis* transformants bearing dexamethasone-

inducible ZED1^{V212T} were photographed before treatment (A) and 72 h post-dexamethasone

spray (B) and screened for transgene expression (C), as described for ZED1^{wt} and ZED1$^{'DT'}$ in

Fig 6.

S14 Fig. BLASTP results showing sequences from *Nicotiana spp.* with similarity to

***Arabidopsis* PBL and ZRK query sequences.**

Plotted points represent sequences from *Nicotiana spp.* (*N. tabacum*, *N. attenuata*, *N. sylvestris*,

N. tomentosiformis) with regions homologous to the indicated *Arabidopsis* query sequences – 46

PBLs (blue) and 12 ZRKs (red). Up to 500 hits for each query are shown where the homologous

region spans at least 50% of the length of the query sequence and has an expectation score (E-

value) of less than 10e-20. Hits with E-values of zero were arbitrarily assigned values of 10e-225

prior to logarithmic transformation and plotting since the logarithm of zero is not defined.

S15 Fig. Expression of a ZED1 allele with promiscuous PBL binding activity activates the

Hypersensitive Response in *Nicotiana benthamiana*.

(A) A *N. benthamiana* leaf infiltrated with *Agrobacterium* strains delivering the indicated

(combinations of) transgenes, 26 h post-induction of expression by spray with dexamethasone.

(B) Protein lysates from leaf cores punched from an additional leaf, similarly infiltrated with the

same *Agrobacterium* cell suspensions, were assessed for expression of HA-tagged transgenes.

S16 Fig. Hypothetical 'front-to-front' pseudokinase-kinase heterodimer.

(A) Molecular models of ZED1 (dark blue helices, orange activation loop) and PBS1 (light blue

helices, pink activation loop) are superimposed over the asymmetric dimer of inactivated IRAK4

described by Ferrao *et al.* (PDB: 4U97) [45]. Note that the template IRAK4 structure is hidden

except for the co-crystalized inhibitor, staurosporine, which competes with ATP for binding to

kinase active sites. (B) A zoomed in view of the same hypothetical ZED1-PBS1 dimer shown in

panel A, rotated by 90 degrees to highlight a 'top-down' view of the predicted interface between

the ZED1 active site and the PBL activation loop acetylation site (S244 in PBS1). Images were

prepared with UCSF CHIMERA [76].

S17 Fig. Pixel analysis of spots representing the output from Y2H and Y3H interaction

assays.

Averaged pixels from the center of all yeast colonies presented in Fig 2, S4 Fig, S5 Fig, S8 Fig

and S9 Fig are plotted in three dimensions according to the absolute values of their red, green

and blue pixel channels (A), as log-transformed pairwise channel ratios (B), or as log-

transformed ratios of a given channel divided by the sum of the other two (C). Plotted points are

filled with the colour of the averaged pixel that they represent, while the colour of their borders

is determined by the relative intensity of the blue channel ($\frac{blue}{red+green}$), as described in S2 Fig

and Materials and Methods. Panels D, E, and F present two-dimensional plots of all possible

pairwise combinations of the three axes shown in panels A, B, and C, respectively. Note that the

right-most panel of panel F is comparable to the plot shown in S2 Fig, but with the x and y axes

flipped.

S18 Fig. Survey of the prevalence of ambiguous and possible phospho/acetyl-accepting

residues present in ZED1 sequences from diverse *Arabidopsis* ecotypes.

(A) Histograms showing the incidence of ambiguous (left) or potential phospho/acetyl-accepting

residues (right) in full-length ZED1 sequences (top) or in activation loop sequences only

(bottom). (B) The same data shown in panel A are plotted as cumulative proportions of the total

sample size. (C) Aligned ZED1 activation loop sequences for the Col-0 reference ecotype and

the four *Arabidopsis* ecotypes with apparent phospho/acetyl-accepting residues. Matches to

canonical DFG and APE motifs defining the activation loop are highlighted in blue, potential

phospho/acetyl-accepting residues are highlighted in green, and V212 from the reference Col-0

ecotype is highlighted in bold, black font, as in S6 Fig, panel B. Ambiguous activation loop

positions translated as 'X' are shown in bold, grey font.

S19 Fig. Demonstration of the pixel-processing logic used in transformations of *Arabidopsis*

plant images. Pixels from an input image (A) are subjected to threshold-masking of full-colour

RGB pixels (C) based on the ratio between red and green channel intensities (B). False-colour

representations of the absolute pixel intensities for individual red (D), green (E), and blue (F)

channels are shown below.

S1 Table. Plasmids generated and/or used for this study.

Plasmids are organized by type into spreadsheet tabs: 1A – Gateway-compatible 'entry vector'

clones; 1B – Y2H bait plasmids; 1C – Y2H prey plasmids; 1D – Yeast integration plasmids; 1E

– Plant expression plasmids.

S2 Table. Bait fusion plasmid transformants of yeast strain EGY48 (*MAT* α) and its

derivatives bearing chromosomal integrations of bacterial effectors or *Arabidopsis ZED1*.

S3 Table. Prey fusion plasmid transformants of yeast strain RFY206/pSH18-34 (*MAT* A)

and its derivatives bearing chromosomal integrations of *Arabidopsis ZED1*.

S4 Table. *Agrobacterium* strains used for transformation of *Arabidopsis* and *N.*

***benthamiana*.**

S1 File. Supplementary description of mass spectrometry analysis and results.

S2 File. Core kinase domain alignment used for phylogenetic inference of the trees shown

in Fig 3B and S4 Fig, panel A.

S3 File. Newick format phylogenetic tree describing relationships between kinase domain

sequences.

S4 File. Clustal-formatted Muscle alignment of 813 unique ZED1 protein sequences

representing 1121 distinct *Arabidopsis* ecotypes.

S5 File. List of *Arabidopsis* ecotypes corresponding to each of the unique ZED1 protein

sequence IDs in S4 File.

A

Bait: HopZ1a^wt   HopZ1a^C216A

Interaction scheme: A

Prey Array

B

Protein sequence	HopZ1a allele	Peptides/acetylation sites identified by LC-MS/MS	% coverage
PBS1	wt		32.7
PBS1	wt		32.5
PBS1	wt		31.1
PBS1	C216A		27.4
PBS1	C216A		41.2
PBS1	C216A		18.2

C

Protein sequence	HopZ1a allele	Peptides/acetylation sites identified by LC-MS/MS	% coverage
ZED1	wt		46.1
ZED1	C216A		46.4

A

ZED1 mutated sites:

a ('3xG')   b (D92)   c (N173)   d (W193)   e (V212)

N — β1 β2 β3 αC β4 β5 αD αE β6 β7 β8 β9 αF αG αH — C

Activation loop (191-219)

Conserved kinase domain (STKc_IRAK; cd14066) (55-330)

B

A

L-threonine → **O-acetyl-L-threonine**   L-isoleucine

HopZ1a

L-serine → **O-acetyl-L-serine**   L-glutamine

HopZ1a

L-alanine

B

A

ZAR1^{wt}   1 — [CC]—[NB]——[LRR]— 852

ZAR1^{ΔLRR}   1 — [CC]—[NB]— 515

ZAR1^{ΔCC}   145 —[NB]——[LRR]— 852

B

Bait:	ZAR1^{wt}	ZAR1^{ΔLRR}	ZAR1^{ΔCC}

ho ^{MAT α} :   -   HopZ1a^{wt}   HopZ1a^{C216A}   ZED1^{wt} ZED1^{DT}

ho ^{MAT A} :   - ZED1^{wt}   - ZED1^{wt}   - ZED1^{wt}   -   -

Prey Array

| PBL21 |
| PBL5 |
| PBS1 |
| PBL27 |
| PBL17 |
| PBL8 |
| PBL4 |
| PBL18 |
| PBL9 |
| PBL15 |
| PBL13 |

Relative interaction strength
1.0 — 0.75 — 0.50 — 0.25 — 0.0

x ZAR1^{ΔCC}
(Y4H with HopZ1a^{wt})

Interaction scheme:
A B B C B C B B | A B B C B C B B | A B B C B C B B

Column index:
1 2 3 4 5 6 7 8 | 9 10 11 12 13 14 15 16 | 17 18 19 20 21 22 23 24

Agrobacterium (co-)infiltrations

1. HopZ1a
2. **ZED1^{wt} + HopZ1a**
3. ZED1^{wt}
4. ZED1^{N173D}
5. **ZED1^{V212T}**
6. **ZED1$^{N173D\ V212T}$**

TRV1 + TRV2::***GUS***   TRV1 + TRV2::***ZAR1***

VIGS treatment
(two weeks prior)