Upscaling the surveillance of tick-borne pathogens in the French Caribbean islands

Mathilde Gondard [1,2]**, Sabine Delannoy** [3]**, Valérie Pinarello** [2,4]**, Rosalie Aprelon** [2,4]**, Elodie Devillers** [1]**, Clémence Galon** [1]**, Jennifer Pradel** [2,4]**, Muriel Vayssier-Taussat** [1]**, Emmanuel Albina** [2,4] **and Sara Moutailler** [1*]

[1] UMR BIPAR, Animal Health Laboratory, ANSES, INRA, Ecole Nationale Vétérinaire d'Alfort, Université Paris-Est, Maisons-Alfort, France; mathilde.gondard@gmail.com; elodie.devilliers@anses.fr; clemence.galon@anses.fr; Muriel.Vayssier@inra.fr; sara.moutailler@anses.fr

[2] CIRAD, UMR ASTRE, F-97170 Petit-Bourg, Guadeloupe, France; mathilde.gondard@gmail.com; valerie.pinarello@cirad.fr; rosalie.aprelon@cirad.fr; jennifer.pradel@cirad.fr; emmanuel.albina@cirad.fr

[3] IdentyPath Platform, Laboratory for Food Safety, ANSES, Maisons-Alfort, France. Sabine.delannoy@anses.fr

[4] ASTRE, Univ Montpellier, CIRAD, INRA, Montpellier, France; valerie.pinarello@cirad.fr; rosalie.aprelon@cirad.fr; jennifer.pradel@cirad.fr; emmanuel.albina@cirad.fr

* Correspondence: sara.moutailler@anses.fr; Tel.: +33 1 49 77 46 50

Please note that this paper has already been published online as a preprint on bioRxiv: https://doi.org/10.1101/532457

Abstract: Despite the high burden of vector-borne disease in (sub)-tropical areas, few information are available regarding the diversity of tick and tick-borne pathogens circulating in the Caribbean. Management and control of vector-borne disease require actual epidemiological data to better assess and anticipate the risk of (re)-emergence of tick-borne diseases in the region. To simplify and reduce the costs of such large-scale surveys, we implemented a high-throughput microfluidic real-time PCR system suitable for the screening of the main bacterial and parasitic genera involved in tick-borne disease and potentially circulating in the area. We used the new screening tool to perform an exploratory epidemiological study on 132 specimens of *Amblyomma variegatum* and 446 of *Rhipicephalus microplus* collected in Guadeloupe and Martinique. Not only the system was able to detect the main pathogens of the area– *Ehrlichia ruminantium*, *Rickettsia africae*, *Anaplasma marginale*, *Babesia bigemina*, *Babesia bovis*– but the system also provided evidence of unsuspected microorganisms in Caribbean ticks, belonging to the *Anaplasma*, *Ehrlichia*, *Borrelia* and *Leishmania* genera. Our study demonstrated how high-throughput microfluidic real-time PCR technology can assist large-scale epidemiological studies, providing a rapid overview of tick-borne pathogen and microorganism diversity, and opening up new research perspectives for the epidemiology of tick-borne pathogens.

Keywords: Tick; Bacteria; Parasites; Caribbean; Microfluidic real-time PCRs.

1. Introduction

Among hematophagous arthropods, ticks transmit the greatest variety of pathogens of public health and veterinary importance whose incidence is growing worldwide [1]. The French West Indies, including the islands of Guadeloupe and Martinique, are located in the heart of the Caribbean Neotropical zone, a cosmopolitan area characterized by a tropical climate, intercontinental trade and animal movements (legal and illegal trade as well as bird migration) that are favorable for the introduction and spread of ticks and tick-borne pathogens (TBPs) [2]. Yet, the epidemiological

situation of the Caribbean area with regard to the diversity of tick species and tick-borne diseases
(TBDs) is poorly documented [3].

Amblyomma variegatum, also known as the tropical bont tick (TBT) in the Caribbean, and
Rhipicephalus microplus (the "cattle tick") have been the two main tropical livestock pests since their
introduction in the Caribbean through imports of infested animals from Africa and Asia in the 18th-
19th centuries [4]. Both tick species are present in the French West Indies, where they are involved in
the transmission of TBPs of medical and veterinary importance [5-9].

R. microplus, a one-host tick highly specific to cattle, is mainly involved in the transmission of
Anaplasma marginale, *Babesia bovis* and *Babesia bigemina*, causing bovine anaplasmosis and babesiosis,
respectively. These endemic pathogens are responsible for important economical lost to farming
industries in the Caribbean and are still a sanitary threat [7,10].

A. variegatum is a three-host tick species, with immature stages that can parasitize a wide range
of hosts, including rodents, mongooses and birds, as well as an adult stage that is more specific to
cattle [11]. This tick species is mainly involved in *Ehrlichia ruminantium* transmission, the causative
agent of heartwater, a fatal ruminant ehrlichiosis. Although *A. variegatum* is present in both
Martinique (mainly in the south) and Guadeloupe (widespread), *E. ruminantium* has only been
reported in Guadeloupe [12]. In addition, *A. variegatum* ticks are also a vector of *Rickettsia africae*,
which is common in the Caribbean and can induce human rickettsiosis, called African tick-bite fever
[9,13,14]. African tick-bite fever remains a concern mainly for travelers. Indeed, despite high levels of
tick infection and seroprevalence in human and cattle sera, only two human cases of African tick-bite
fever have been reported to date, only in travelers returning from Guadeloupe [9,15]. Lastly, *A.*
variegatum is also involved in the epidemiology of *Theileria mutans* and *Theileria velifera*, two cattle
parasites with low and no virulence, respectively [6,8]. However, very few information are available
on the distribution and prevalence of these two apicomplexa in the Carribean.

Most of the epidemiological data available did not survey or determine the diversity of TBPs
circulating in the Carribean, since they were often limited to the detection of some well-known
pathogens, via serological studies in animals or humans, or on molecular biology testing (PCR, nested
PCR) [16,17]. Thus, regarding the lack of recent information, the limited extent of the epidemiological
data available, new insight into the epidemiology of ticks and TBPs was needed to better address the
prevalence and (re-)emergence of TBDs in the Caribbean.

In order to improve the surveillance ability of tick-borne pathogens in the Neotropical area, we
implemented a new large scale screening tool based on microfluidic real-time PCR approach.
Microfluidic real-time PCR system is based on the use of microfluidic chip allowing the performance
of up to 9216 individual PCR reactions per run, and thus the simultaneous detection of up to 96 target
into up to 96 samples. The recent development and use of a microfluidic real-time PCR system for
the rapid and concomitant detection of a large panel of TBPs in European ticks has paved the way
for promising and broader surveillance capacities [18-22]. Here, we adapted and designed a new
microfluidic real-time PCR system suited to the simultaneous screening of the main bacteria and
protozoans potentially transmitted by ticks in the Caribbean. Not only did the system enable the
direct detection of 49 bacterial and parasitic species, but it also enabled, within a single experiment,
broader capacities for the surveillance of potentially pathogenic microorganisms, by targeting the
main bacterial and protozoan genera involved in human and animal vector-borne diseases (one
protozoan phylum and eight bacterial and protozoan genera). In addition, the system enabled the
molecular identification of the three well-known tick species involved in TBDs in the Caribbean in
order to confirm the morphological tick species identification determined on the field. Finally, we
used the new high-throughput detection tool to conduct large-scale screening of TBPs in 132 *A.*
variegatum and 446 *R. microplus* specimens collected in Guadeloupe and Martinique. We
demonstrated the system's ability to detect well-known TBPs occurring in the French West Indies, as
well as unsuspected TBPs and potential new microorganisms. This new method can considerably
improve the ability to monitor emerging and non-emerging TBPs through large-scale surveys in the
Caribbean area.

2. Results

2.1 Implementation of the high-throughput microfluidic real-time PCR system for tick-borne pathogen
screening

The high-throughput microfluidic real-time PCR system developed for the screening of known
and potential TBPs in Caribbean ticks included 61 sets of primers and probes. Among them, 49
designs were developed for the detection of bacterial (n=32) and protozoan (n=17) species and
bacterial (n=5) and protozoan (n=3) genera/phyla (Table 1). Three sets of primers and probes were
developed for the molecular identification of the three tick species found in the Caribbean: *A.*
variegatum, *R. microplus* and *R. sanguineus* sensu lato (Table 1). Lastly, we developed a design
targeting a conserved region of the 16S rRNA genes in ticks, called "Tick spp.", used as a control for
DNA/RNA extraction (Table 1).

Table 1. List of primer/probe sets constituting the BioMark system, with the positive controls used
108   for their validation.

Microorganisms	Target	Design name	Sequence (5' à 3')	Length (bp)	Positive controls
Rickettsia spp.	gltA	Rick_spp_gltA_F	GTCGCAAATGTTCACGGTACTT	78	**, Culture of *R. slovaca*
		Rick_spp_gltA_R	TCTTCGTGCATTTCTTTCCATTG		
		Rick_spp_gltA_P	TGCAATAGCAAGAACCGTAGGCTGGATG		
Rickettsia massiliae *	ITS	Ri_ma_ITS_F	GTTATTGCATCACTAATGTTATACTG	128	Culture
		Ri_ma_ITS_R	GTTAATGTTGTTGCACGACTCAA		
		Ri_ma_ITS_P	TAGCCCCGCCACGATATCTAGCAAAAA		
Rickettsia rickettsii *	ITS	Ri_ri_ITS_F	TCTACTCACAAAGTTATCAGGTTAA	124	Plasmid
		Ri_ri_ITS_R	CCTACGATACTCAGCAAAATAATTT		
		Ri_ri_ITS_P	TCGCTGGATATCGTTGCAGGACTACAG		
Rickettsia conorii	sca1	Ri_co_sca1_F	GTAGATGCTTCATAGAATACTGC	88	Infected *Rhipicephalus sanguineus* s.l.
		Ri_co_sca1_R	CCAAATTTAGTCTACCTTGTGATC		
		Ri_co_sca1_P	TCCTCCTGACGTATTAAAAGAAGCTGAAGCT		
Rickettsia africae	sca1	Ri_af_sca1_F	GATACGACAAGTACCTCGCAG	122	Culture
		Ri_af_sca1_R	GGATTATATACTTTAGGTTCGTTAG		
		Ri_af_sca1_P	CAGATAGGAACAGTAATTGTAACGGAACCAG		
Rickettsia felis	orfB	Ri_fel_orfB_F	ACCCTTTTCGTAACGCTTTGC	163	Culture
		Ri_fel_orfB_R	TATACTTAATGCTGGGCTAAACC		
		Ri_fel_orfB_P	AGGGAAACCTGGACTCCATATTCAAAAGAG		
Rickettsia typhi	ompB	Ri_typ_ompB_F	CAGGTCATGGTATTACTGCTCA	133	Culture
		Ri_typ_ompB_R	GCAGCAGTAAAGTCTATTGATCC		
		Ri_typ_ompB_P	ACAAGCTGCTACTACAAAAAGTGCTCAAAATG		
Rickettsia prowazekii	gltA	Ri_pro_gltA_F	CAAGTATCGGTAAAGATGTAATCG	151	Plasmid
		Ri_pro_gltA_R	TATCCTCGATACCATAATATGCC		
		Ri_pro_gltA_P	ATATAAGTAGGGTATCTGCGGAAGCCGAT		

Borrelia spp. *	23S rRNA	Bo_bu_sl_23S_F	GAGTCTTAAAAGGGCGATTTAGT	73	**, Culture of *B. afzelii*, *B. garinii*, *B. valaisiana*, *B. lusitaniae*, *B. recurrentis*
		Bo_bu_sl-23S_R	CTTCAGCCTGGCCATAAATAG		
		Bo_bu_sl_23S_P	TAGATGTGGTAGACCCGAAGCCGAGT		
Borrelia burgdorferi sensu stricto	glpA	Bo_bu_glpA_F	GCAATTACAAGGGGGTATAAAGC	206	Culture
		Bo_bu_glpA_R	GGCGTGATAAGTGCACATTCG		
		Bo_bu_glpA_P	TTAATTAAACGGGGTGCATTCTTCTCAAGAATG		
Borrelia anserina	fla	Bor_ans_fla_F	GGAGCACAACAAGAGGGAG	76	Plasmid
		Bor_ans_fla_R	TTGGAGAATTAACCCCACCTG		
		Bor_ans_fla_P	TGCAAGCAACTCCAGCTCCAGTAGCT		
Borrelia lonestari	glpQ	Bor_lon_glpQ_F	GATCCAGAACTTGATACAACCAC	99	Infected *Amblyomma americanum*
		Bor_lon_glpQ_R	TTCATCTAGTGAGAAGTCAGTAG		
		Bor_lon_glpQ_P	AGTAATATCGTCCGTCTTCCCTAGCTCG		
Borrelia parkeri	gyrB	Bor_par_gyrB_F	GCAAAACGATTCAAAGTGAGTCC	184	Culture
		Bor_par_gyrB_R	CTCATTGCCTTTAAGAAACCACTT		
		Bor_par_gyrB_P	TTAAAACCAGCAACATGAGTTCCTCCTTCTC		
Borrelia bissettii *	rpoB	Bo_bi_rpoB_F	GCAACCAGTCAGCTTTCACAG	118	Plasmid
		Bo_bi_rpoB_R	CAAATCCTGCCCTATCCCTTG		
		Bo_bi_rpoB_P	AAAGTCCTCCCGGCCCAAGAGCATTAA		
Borrelia theileri	glpQ	Bo_th_glpQ_F	GTGCTAACAAAGGACAATATTCC	213	Plasmid
		Bo_th_glpQ_R	GGTTAGTGGAAAACGGTTAGGAT		
		Bo_th_glpQ_P	TATTATAATTCACGAGCCAGAGCTTGACAC		
Bartonella spp.	ssrA	Bart_spp_ssrA_F	CGTTATCGGGCTAAATGAGTAG	118	**, Culture of *B. quintana*
		Bart_spp_ssrA_R	ACCCCGCTTAAACCTGCGA		
		Bart_spp_ssrA_P	TTGCAAATGACAACTATGCGGAAGCACGTC		
Bartonella barcilliformis *	rpoB	Ba_ba_rpoB_F	GAAGAGTTTGTAGTTTGTCGTCA	105	Culture
		Ba_ba_rpoB_R	AGCAGCTACAGAAACCAACTG		
		Ba_ba_rpoB_P	TGCAGGTGAAGTTTTGATGGTGCCACG		
Bartonella henselae	ribC	Bar_he_ribC_F	GGGATGCGATTTAATAGTTCTAC	116	Culture
		Bar_he_ribC_R	CGCTTGTTGTTTTGATCCTCG		
		Bar_he_ribC_P	ACGTTATAGTAGCGAAAACTTAGAAATTGGTGC		
Bartonella vinsonii subsp. *berkhoffii*	ITS	Bar_vin_ITS_F	GGAATTGCTTAACCCACTGTTG	141	Culture
		Bar_vin_ITS_R	CCTTATTGATTTAGATCTGATGGG		
		Bar_vin_ITS_P2	AGAAACTCCCGCCTTTATGAGAGAAATCTCT		
	Icd	Co_bu_icd_F	AGGCCCGTCCGTTATTTTACG	74	Culture
		Co_bu_icd_R	CGGAAAATCACCATATTCACCTT		

Coxiella burnetii and Coxiella-like *	IS1111	Co_bu_icd_P	TTCAGGCGTTTTGACCGGGCTTGGC	86	Culture
		Co_bu_IS111_F	TGGAGGAGCGAACCATTGGT		
		Co_bu_IS111_R	CATACGGTTTGACGTGCTGC		
		Co_bu_IS111_P	ATCGGACGTTTATGGGGATGGGTATCC		
Francisella tularensis and Francisella-like endosymbionts *	tul4	Fr_tu_tul4_F	ACCCACAAGGAAGTGTAAGATTA	76	Culture
		Fr_tu_tul4_R	GTAATTGGGAAGCTTGTATCATG		
		Fr_tu_tul4_P	AATGGCAGGCTCCAGAAGGTTCTAAGT		
	fopA	Fr_tu_fopA_F	GGCAAATCTAGCAGGTCAAGC	91	Culture
		Fr_tu_fopA_R	CAACACTTGCTTGAACATTTCTAG		
		Fr_tu_fopA_P	AACAGGTGCTTGGGATGTGGGTGGTG		
Anaplasma spp.	16S rRNA	Ana_spp_16S_F	CTTAGGGTTGTAAAACTCTTTCAG	160	**
		Ana_spp_16S_R	CTTTAACTTACCAAACCGCCTAC		
		Ana_spp_16S_P	ATGCCCTTTACGCCCAATAATTCCGAACA		
Anaplasma marginale *	msp1b	An_ma_msp1_F	CAGGCTTCAAGCGTACAGTG	85	Experimentally infected bovine blood sample
		An_ma_msp1_R	GATATCTGTGCCTGGCCTTC		
		An_ma_msp1_P	ATGAAAGCCTGGAGATGTTAGACCGAG		
Anaplasma phagocytophilum *	msp2	An_ph_msp2_F	GCTATGGAAGGCAGTGTTGG	77	Infected *Ixodes* spp. tick
		An_ph_msp2_R	GTCTTGAAGCGCTCGTAACC		
		An_ph_msp2_P	AATCTCAAGCTCAACCCTGGCACCAC		
Anaplasma platys *	groEL	An_pla_groEL_F	TTCTGCCGATCCTTGAAAACG	75	Infected canine blood sample
		An_pla_groEL_R	CTTCTCCTTCTACATCCTCAG		
		An_pla_groEL_P	TTGCTAGATCCGGCAGGCCTCTGC		
Anaplasma bovis *	groEL	An_bo_groEL_F	GGGAGATAGTACACATCCTTG	73	Plasmid
		An_bo_groEL_R	CTGATAGCTACAGTTAAGCCC		
		An_bo_groEL_P	AGGTGCTGTTGGATGTACTGCTGGACC		
Anaplasma ovis *	msp4	An_ov_msp4_F	TCATTCGACATGCGTGAGTCA	92	Plasmid
		An_ov_msp4_r	TTTGCTGGCGCACTCACATC		
		An_ov_msp4_P	AGCAGAGAGACCTCGTATGTTAGAGGC		
Ehrlichia spp. *	16S rRNA	Neo_mik_16S_F	GCAACGCGAAAAACCTTACCA	98	**
		Neo_mik_16S_R	AGCCATGCAGCACCTGTGT		
		Neo_mik_16S_P	AAGGTCCAGCCAAACTGACTCTTCCG		
	gltA	Eh_ca_gltA_F	GACCAAGCAGTTGATAAAGATGG	136	Culture

		Eh_ca_gltA_R	CACTATAAGACAATCCATGATTAGG		
Ehrlichia canis		Eh_ca_gltA_P	ATTAAAACATCCTAAGATAGCAGTGGCTAAGG		
Ehrlichia chaffeensis *	dsb	Eh_ch_dsb_F	TATTGCTAATTACCCTCAAAAAGTC	117	Infected *Amblyomma americanum*
		Eh_ch_dsb_R	GAGCTATCCTCAAGTTCAGATTT		
		Eh_ch_dsb_P	ATTGACCTCCTAACTAGAGGGCAAGCA		
Ehrlichia ewingii *	dsb	Eh_ew_dsb_F	CAATACTTGGAGAAGCATCATTG	111	Infected *Amblyomma americanum*
		Eh_ew_dsb_R	TTGCTTATGGCTTAATGCTGCAT		
		Eh_ew_dsb_P	AAAGCAGTACGTGCAGCATTGGCTGTA		
Ehrlichia ruminantium	gltA	Eh_ru_gltA_F	CCAGAAAACTGATGGTGAGTTAG	116	Culture
		Eh_ru_gltA_R	AGCCTACATCAGCTTGAATGAAG		
		Eh_ru_gltA_P	AGTGTAAACTTGCTGTTGCTAAGGTAGCATG		
Panola Mountain Ehrlichia	gltA	Eh_PME_gltA_F	GCTAGTTATGAGTTAGAATGTAAAC	121	Infected *Amblyomma americanum*
		Eh_PME_gltA_R	TACTATAGGATAATCTTGAATCAGC		
		Eh_PME_gltA_P	TTGCTATCGCTAAAATTCCAAGTATGATTGCG		
Neoehrlichia mikurensis *	groEL	Neo_mik_groEL_F	AGAGACATCATTCGCATTTTGGA	96	Infected rodent blood sample
		Neo_mik_groEL_R	TTCCGGTGTACCATAAGGCTT		
		Neo_mik_groEL_P	AGATGCTGTTGGATGTACTGCTGGACC		
Aegyptianella pullorum	groEL	Ae_pul_groEL_F	AGCCAGTATTATCGCTCAAGG	168	Plasmid
		Ae_pul_groEL_R	GCCTCACGTGCCTTCATAAC		
		Ae_pul_groEL_P	TGCTTCTCAGTGTAACGACAGGGTTG		
Apicomplexa	18S rRNA	Apic_18S_F	TGAACGAGGAATGCCTAGTATG	104	**, Infected canine blood sample, with *B. canis rossi*, *B. canis canis*; Culture of *B. divergens*, *T. lestoquari*, *T. annulata*
		Apic_18S_R	CACCGGATCACTCGATCGG		
		Apic_18S_S	TAGGAGCGACGGGCGGTGTGTAC		
Babesia canis vogeli *	hsp70	Ba_vo_hsp70_F	TCACTGTGCCTGCGTACTTC	87	Infected canine blood sample
		Ba_vo_hsp70_R	TGATACGCATGACGTTGAGAC		
		Ba_vo_hsp70_P	AACGACTCCCAGCGCCAGGCCAC		
Babesia ovis *	18S rRNA	Ba_ov_18S_F	TCTGTGATGCCCTTAGATGTC	92	Plasmid
		Ba_ov_18S_R	GCTGGTTACCCGCGCCTT		
		Ba_ov_18S_P	TCGGAGCGGGGTCAACTCGATGCAT		
Babesia bigemina *		Ba_big_RNA18S_F	ATTCCGTTAACGAACGAGACC	99	Plasmid

	18S rRNA	Ba_big_RNA18S_R	TTCCCCCACGCTTGAAGCA		
		Ba_big_RNA18S_P	CAGGAGTCCCTCTAAGAAGCAAACGAG		
Babesia gibsoni	Rap1	Ba_gib_rap1_F	CTCTTGCTCATCATCTTTTCGG	130	Plasmid
		Ba_gib_rap1_R	TCAGCGTATCCATCCATTATATG		
		Ba_gib_rap1_S	TTTAATGCGTGCTACGTTGTACTTCCCAAAG		
Babesia caballi *	Rap1	Ba_cab_rap1_F	GTTGTTCGGCTGGGGCATC	94	Plasmid
		Ba_cab_rap1_R	CAGGCGACTGACGCTGTGT		
		Ba_cab_rap1_P	TCTGTCCCGATGTCAAGGGGCAGGT		
Babesia bovis *	CCTeta	Ba_bo_CCTeta_F	GCCAAGTAGTGGTAGACTGTA	100	Plasmid
		Ba_bo_CCTeta_R	GCTCCGTCATTGGTTATGGTA		
		Ba_bo_CCTeta_P	TAAAGACAACACTGGGTCCGCGTGG		
Babesia duncani *	ITS2	Ba_du_ITS_F	ATTTCCGTTTGCGAGAGTTGC	87	Plasmid
		Ba_du_ITS_R	AGGAAGCATCAAGTCATAACAAC		
		Ba_du_ITS_P	AACAAGAGGCCCCGAGATCAAGGCAA		
Babesia microti *	CCTeta	Bab_mi_CCTeta_F	ACAATGGATTTTCCCCAGCAAAA	145	Culture
		Bab_mi_CCTeta_R	GCGACATTTCGGCAACTTATATA		
		Bab_mi_CCTeta_P	TACTCTGGTGCAATGAGCGTATGGGTA		
Theileria parva *	18S rRNA	Th_pa_18S_F	GAGTATCAATTGGAGGGCAAG	173	Culture
		Th_pa_18S_R	CAGACAAAGCGAACTCCGTC		
		Th_pa_18S_P	AAATAAGCCACATGCAGAGACCCCGAA		
Theileria mutans	ITS	The_mu_ITS_F	CCTTATTAGGGGCTACCGTG	119	Plasmid
		The_mu_ITS_R	GTTTCAAATTTGAAGTAACCAAGTG		
		The_mu_ITS_P	ATCCGTGAAAAACGTGCCAAACTGGTTAC		
Theileria velifera	18S rRNA	The_ve_18S_F	TGTGGCTTATCTGGGTTCGC	151	Plasmid
		The_ve_18S_R	CCATTACTTTGGTACCTAAAACC		
		The_ve_18S_P	TTGCGTTCCCGGTGTTTTACTTTGAGAAAG		
Theileria equi	ema1	Th_eq_ema1_F4	CGGCAAGAAGCACACCTTC	167	Plasmid
		Th_eq_ema1_R4	TGCCATCGCCCTTGTAGAG		
		Th_eq_ema1_P4	AAGGCTCCAGGCAAGCGCGTCCT		
Cytauxzoon felis	ITS2	Cy_fel_ITS2_F	AAGATCCGAACGGAGTGAGG	119	Plasmid
		Cy_fel_ITS2_R	GTAGTCTCACCCAATTTCAGG		
		Cy_fel_ITS2_S	AAGTGTGGGATGTACCGACGTGTGAG		
Hepatozoon spp.	18S rRNA	Hepa_spp_18S_F	ATTGGCTTACCGTGGCAGTG	175	**
		Hepa_spp_18S_R	AAAGCATTTTAACTGCCTTGTATTG		
		Hepa_spp_18S_S	ACGGTTAACGGGGGATTAGGGTTCGAT		

Hepatozoon canis	18S rRNA	He_can_18S_F	TTCTAACAGTTTGAGAGAGGTAG	221	Infected canine blood sample
		He_can_18S_R	AGCAGACCGGTTACTTTTAGC		
		He_can_18S_S	AGAACTTCAACTACGAGCTTTTTAAC TGCAAC		
Hepatozoon americanum	18S rRNA	He_ame_18S_F2	GGTATCATTTTGGTGTGTTTTTAAC	159	Plasmid
		He_ame_18S_R2	CTTATTATTCCATGCTCCAGTATTC		
		He_ame_18S_P2	AAAAGCGTAAAAGCCTGCTAAAAAC ACTCTAC		
Leishmania spp.	hsp70	Leish_spp_hsp70_F	CGACCTGTTCCGCAGCAC	78	** and culture of *L. martiniquensis*
		Leish_spp_hsp70_R	TCGTGCACGGAGCGCTTG		
		Leish_spp_hsp70_S	TCCATCTTCGCGTCCTGCAGCACG		
Leishmania infantum	ITS	Le_inf_ITS_F	CGCACCGCCTATACAAAAGC	103	Culture
		Le_inf_ITS_R	GTTATGTGAGCCGTTATCCAC		
		Le_inf_ITS_S	ACACGCACCCACCCCGCCAAAAAC		
Rangelia vitalii	18S rRNA	Ra_vit_18S_F	TAACCGTGCTAATTGTAGGGC	92	Plasmid
		Ra_vit_18S_R	GAATCACCAAACCAAATGGAGG		
		Ra_vit_18S_S	TAATACACGTTCGAGGGCGCGTTTTG C		
Tick spp.	16S rRNA	Tick_spp_16S_F	AAATACTCTAGGGATAACAGCGT	99	**
		Tick_spp_16S_R	TCTTCATCAAACAAGTATCCTAATC		
		Tick_spp_16S_P	CAACATCGAGGTCGCAAACCATTTTG TCTA		
Amblyomma variegatum	ITS2	Amb_var_ITS2_F	GCCAGCCTCTGAAGTGACG	117	Tick extract (Guadeloupe)
		Amb_var_ITS2_R	TTCTGCGGTTTAAGCGACGC		
		Amb_var_ITS2_P	TCTTGCCACTCGACCCGTGCCTC		
Rhipicephalus microplus	ITS2	Rhi_mic_ITS2_F	GCTTAAGGCGTTCTCGTCG	144	Tick extract (Galapagos Islands)
		Rhi_mic_ITS2_R	CAAGGGCAGCCACGCAG		
		Rhi_mic_ITS2_P	TAGTCCGCCGTCGGTCTAAGTGCTTC		
Rhipicephalus sanguineus sensu lato	ITS2	Rhi_san_ITS2_F	TTGAACGCTACGGCAAAGCG	110	Tick extract (France)
		Rhi_san_ITS2_R	CCATCACCTCGGTGCAGTC		
		Rhi_san_ITS2_P	ACAAGGGCCGCTCGAAAGGCGAGA		

The detection ability of each design and the effect of pre-amplification on detection signals were
first checked by TaqMan real-time PCR on a LightCycler 480 apparatus using a range of dilutions of
positive controls (Table 1, Table S2). Three kinds of positive controls were used, including bacterial
or protozoan cultures when available, DNA from infected ticks or blood samples, and plasmidic
constructions as a last resort (Table 1). Except for the design targeting *Borrelia burgdorferi* sensu stricto,
which never succeeded in detecting the positive controls even after a pre-amplification step, the
remaining 60 designs targeting TBPs and tick species were able to detect their target with Ct values

between six and 38 (data not shown). Pre-amplification improved the quality of detection and was
therefore validated as part of the screening protocol (see Figure S1).
The relative specificity of the 61 designs was then evaluated using the BioMark system and a
total of 62 positive controls (Figure 1, Table S2).

Figure 1. BioMarkTM dynamic array system specificity test (96.96 chip). Each square corresponds to a
single real-time PCR reaction, where rows indicate the pathogen in the sample and columns represent the
target of the primer/probe set. Ct values for each reaction are represented by a color gradient; the color
scale is shown on the right y-axis. The darkest shades of blue and black squares are considered as negative
reactions with Ct > 30.

Forty-three primer/probe sets were able to specifically detect and amplify their target using a Ct
cut-off value of 30; they were then directly validated (Figure 1). The remaining designs were able to
detect and amplify their target, but they also gave positive results in outgroup controls. Interestingly,
two kinds of unsuspected signals were observed; some were related to cross-reactions with closely
related species and some to potential co-infections in controls corresponding to field samples (Figure
1). Thus, eight designs – *Rickettsia massiliae*, *Rickettsia conorii*, *Bartonella henselae*, *Bartonella bacilliformis*,
Babesia canis vogeli, *Babesia microti*, *Theileria parva*, *Hepatozoon americanum* – gave positive results in
outgroup controls, revealing cross-reactions with one to two closely related species (Figure 1).
Caution will be required when interpreting results obtained with these designs. Seven designs –
Rickettsia spp., *Rickettsia felis*, *Rickettsia africae*, *Apicomplexa*, *Babesia bigemina*, *Hepatozoon* spp.,
Hepatozoon canis – gave positive results in outgroup controls linked to potential co-infection in
controls corresponding to DNA from infected ticks or blood samples (Figure 1). As co-infections may

occur in natural tick or blood samples, these unexpected detections in biological samples were likely
due to the natural (co-)occurrence of microorganisms rather than to cross-reactions. Finally, the
Babesia ovis and *Rickettsia rickettsii* designs gave multiple cross-reactions with closely related species
or distant outgroups and thus were considered as non-specific and removed from the rest of the study
(Figure 1). More details on the relative specificity analysis of the designs are available in Appendix
A.
To conclude, with the exception of the sets of primers and probes targeting *Borrelia burgdorferi*
sensu stricto, *Babesia ovis* and *Rickettsia rickettsii* that were ultimately removed from the study, the 58
remaining designs were validated for the high-throughput screening of pathogens in Caribbean ticks,
taking into account the notified cross-reactions.

2.2 Large-scale TBP detection survey in ticks from Guadeloupe and Martinique

A total of 578 adult ticks were collected from cattle in Guadeloupe and Martinique. In total, 523
samples were tested using the BioMarkTM system developed in this study. The Molecular
identification of *Amblyomma variegatum* and *Rhipicephalus microplus* using the corresponding specific
designs were consistent with the morphological identification made after tick collection. The number
of positive ticks and the corresponding infection rates for each detected pathogen were calculated for
132 *A. variegatum* and 165 and 281 *R. microplus* specimens from Guadeloupe and Martinique,
respectively (Figure 2). As some of the *R. microplus* samples corresponded to pools of two to four
adult specimens, we reported the minimum and maximum infection rates (see Material and
methods).

	Guadeloupe		Martinique
	Positive *Amblyomma variegatum*, out of 132 (IR)	Positive *Rhipicephalus microplus*, out of 165 (IR min - max)	Positive *Rhipicephalus microplus*, out of 281 (IR min - max)
Anaplasma spp.	0	8 (4.8%) - 9 (5.5%)	112 (39.9%) - 116 (41.3%)
Anaplasma marginale	0	5 (3%) - 7 (4.2%)	111 (39.5%) - 116 (41.3%)
Ehrlichia spp.	11 (8.3%)	7 (4.2%) - 11(6.7%)	134 (47.7%) - 138 (49.1%)
Ehrlichia ruminantium	7 (5.3%)	0	0
Borrelia spp.	7 (5.3%)	1 (0.6%)	12 (4.3%)
Rickettsia spp.	130 (98.5%)	25 (15.2%) - 38 (23%)	0
Rickettsia africae	126 (95.5%)	0	0
Leishmania spp.	0	0	2 (0.7%)
Theileria mutans	2 (1.5%)	3 (1.8%) - 4 (2.4%)	4 (1.4%)
Theileria velifera	57 (43.2%)	39 (23.6%) - 52 (31.5%)	72 (25.6%) - 73 (26%)
Babesia bigemina	0	1 (0.6%) - 2 (1.2%)	35 (12.5%) - 36 (12.8%)
Babesia bovis	0	0	2 (0.7%)

Figure 2. Infection rates in ticks collected in Guadeloupe and Martinique. Number of positive *A.*
variegatum ticks (out of 132) and *R. microplus* ticks from Guadeloupe (out of 165) and Martinique (out
of 281). Dots on the map indicate the tick collection sites in Guadeloupe and Martinique. Yellow:
collection site of *R. microplus* ticks; Red: collection site of *A. variegatum* ticks; Blue: collection site of
both tick species, sometimes on the same animal. IR: Infection rate. As some *R. microplus* samples were
pooled, we have presented minimum and maximum tick infection rates.

Conventional PCRs/nested PCRs followed by amplicon sequencing were performed on several
tick samples to confirm some of the results of the newly designed BioMarkTM system (see Materials
and Methods section). Identity percentages of the sequences obtained with reference sequences
available in GenBank (NCBI) are presented in Table 2.

Table 2. Homology between the deposited sequences and reference sequences in GenBank.

Biomark Id	Sequence Name	T	S	An	L	Closest Homology	Id%	Reference
		3	1	MK049	24			
Rickettsia spp.	*Rickettsia africae* Tick208	0	4	851	8	*Rickettsia africae*	100	AF123706.1
Leishmania				MK049	27	Leishmania		
spp.	*Leishmania martiniquensis* Tick389	2	1	850	2	*martiniquensis*	100	AF303938.1
						Leishmania siamensis	100	GQ226033.1
		3		MK049	24			
Borrelia spp.	*Borrelia* sp. Tick7	0	1	846	5	*Borrelia anserina*	90	X75201.1
				MK049	32			
	Borrelia sp. Tick457		4	847	7	*Borrelia* sp. BR	100	EF141022.1
						Borrelia sp. strain Mo063b-flaB	100	KY070335.1
						Borrelia theileri	99	KF569936.1
				MK049	24	*Candidatus* Anaplasma		
Anaplasma spp.	*Anaplasma* sp. Tick314	2	2	845	5	boleense	100	KX987335.1
Anaplasma				MK049	24			
marginale	*Anaplasma* sp. Tick283	2	2	844	4	*Anaplasma marginale*	100	MH155593.1
						Anaplasma centrale	100	MF289482.1
						Anaplasma ovis	100	MG770440.1
						Anaplasma capra	100	MF000917.1
						Anaplasma phagocytophilum	100	DQ648489.1
				MK049	24			
Ehrlichia spp.	*Ehrlichia* sp. Tick428	2	2	849	6	*Ehrlichia* spp.	100	KY594915.1*
						Ehrlichia canis	99	KY594915.1
						Ehrlichia ewingii	99	U96436.1
						Ehrlichia chaffeensis	99	NR_074500.2
						Ehrlichia muris	99	KU535865.1
						Ehrlichia minasensis	99	NR_148800.1
Ehrlichia ruminantium	*Ehrlichia ruminantium* Tick116	1	1	MK049 848	20 7	*Erlichia ruminantium*	100	NR_074155.1
Babesia bigemina	*Babesia bigemina* Tick222	2	1	MK071 738	99	*Babesia bigemina*	100	KP710227.1
Babesia bovis	*Babesia bovis* Tick497	2	2	MK071 739	10 0	*Babesia bovis*	99	AB367921.1

2.2.1. Detection of known TBPs in Caribbean ticks

Seven TBPs known to circulate in the Caribbean were detected in ticks from Guadeloupe and
Martinique: *R. africae*, *E. ruminantium*, *An. marginale*, *B. bigemina*, *B. ovis*, *T. mutans* and *T. velifera*
(Figure 2).
Rickettsia spp. were only detected in ticks collected in Guadeloupe (Figure 2). *R. africae* was
identified in 95.6% of the *A. variegatum* samples (Figure 2). In contrast, *Rickettsia* spp. detected in 15.7-
23.5% of the *R. microplus* samples from Guadeloupe were not directly identified as *R. africae* with the
BioMarkTM system (Figure 2). Thus, 14 *A. variegatum* (6/14) and *R. microplus* (8/14) samples positive
for *Rickettsia* spp. were tested by nested PCR with primers targeting the ompB gene; this was followed
by sequencing. All the sequences recovered were identical and displayed 100% identity with *R.*
africae, confirming that the *Rickettsia* spp. detected in *R. microplus* from Guadeloupe corresponded
also to *R. africae*. (Table 2). The consensus sequence was deposited under the name *Rickettsia africae*
Tick208 (accession number MK049851).

E. ruminantium was identified in 5.1% of the *A. variegatum* ticks from Guadeloupe (Figure 2). We
confirmed the presence of *E. ruminantium* nucleic acids by testing one sample of *A. variegatum* by
conventional PCR targeting the 16S rRNA genes; this was followed by amplicon sequencing. The
sequence obtained displayed 100% sequence identity with *E. ruminantium* and was deposited under
the name *Ehrlichia ruminantium* Tick116 (accession number MK049848) (Table 2).
An. marginale was identified in *R. microplus* ticks from both islands, with infection rates reaching
3.6-4.8% and 39.5-41.3% of specimens from Guadeloupe and Martinique, respectively (Figure 2). We
confirmed the detection of *An. marginale* by testing two samples of *R. microplus* by conventional PCR
targeting the 16S rRNA genes; this was followed by amplicon sequencing. We obtained two identical
sequences, deposited under the name *Anaplasma* sp. Tick283 (accession number MK049844), which
displayed 100% sequence identity with *Anaplasma* spp. including *An. marginale* (Table 2).
B. bigemina was detected in 0.6-1.2% and 12.5-12.8% of the *R. microplus* ticks from Guadeloupe
and Martinique, respectively (Figure 2). *B. bovis* was only detected in ticks from Martinique, with an
infection rate of 0.7% in *R. microplus* samples (Figure 2). As conventional and nested PCR did not
succeed in detecting these parasites, we directly sequenced amplicons obtained with the *B. bigemina*
and *B. bovis* designs developed here, and corresponding sequences were identified (accession
numbers MK071738 and MK071739 respectively) (Table 2).
T. velifera and *T. mutans* were detected in both tick species and on both islands. *T. velifera* was
identified in 42.3% of the *A. variegatum* samples and in 24.1-31.9% and 25.6-26% of the *R. microplus*
samples from Guadeloupe and Martinique, respectively (Figure 2). Moreover, *T. mutans* was detected
in 1.5% of the *A. variegatum* samples and in 1.8-2.4% and 1.4% of the *R. microplus* samples from
Guadeloupe and Martinique, respectively (Figure 2). Unfortunately, neither conventional PCR nor
BioMark amplicon sequencing succeeded in confirming the BioMark results.

2.2.2. Detection of unexpected microorganisms in Caribbean ticks

Unexpected signals were obtained during the screening of microorganisms in ticks from
Guadeloupe and Martinique, including the first detection of untargeted species belonging to the
genera *Anaplasma*, *Ehrlichia*, *Borrelia* and *Leishmania* (Figure 2).
Ehrlichia spp. were detected in *R. microplus* ticks from both islands, with infection rates reaching
4.2-6.6% and 47.7-49.1% in Guadeloupe and Martinique, respectively (Figure 2). We tested two of the
Ehrlichia spp.-positive *R. microplus* samples by conventional PCR targeting the 16S rRNA genes in
order to identify the *Ehrlichia* spp. present in the Caribbean sample. We obtained two identical
sequences, deposited under the name *Ehrlichia* sp. Tick428 (accession number MK049849) (Table 2).
Phylogenetic and genetic distance analyses were performed using a portion of the 16S rRNA genes
of several *Ehrlichia* species (Figure 3). The *Ehrlichia* sp. Tick428 sequence was found within a cluster
including various uncharacterized *Ehrlichia* species detected in ticks from Asia and Africa (Figure 3).

Figure 3. Phylogenetic analysis of 16S rRNA sequences of *Ehrlichia* spp. Phylogenetic analysis of 16S rRNA sequences of *Ehrlichia* spp. using the maximum likelihood method based on the Tamura-Nei model. In the phylogenetic tree, GenBank sequences, species designations and strain names are given. The sequences investigated in the present study are marked with a black circle (*Ehrlichia* sp. Tick428, accession number MK049849) and a black diamond (*Ehrlichia ruminantium* Tick116, accession number MK049848). The tree with the highest log likelihood (-413.76) is shown. The percentage of trees in which the associated taxa clustered together is shown above the branches (bootstrap values). The analysis involved 25 nucleotide sequences. There were a total of 206 positions in the final dataset.

In addition, in around 50% and 18% of the *R. microplus* specimens positive for *Anaplasma* spp., none of the *Anaplasma* species targeted by the BioMarkTM system gave signals, suggesting the presence of an unexpected or new *Anaplasma* spp. (Figure 2). We tested two of the *Anaplasma* spp.-positive *R. microplus* samples by conventional PCR targeting the 16S rRNA genes. We obtained two identical sequences, deposited under the name *Anaplasma* sp. Tick314 (accession number MK049845) (Table 2). This sequence displayed 100% sequence identity with *Candidatus* Anaplasma boleense. Phylogenetic and genetic distance analyses were performed using a portion of the 16S rRNA genes of several *Anaplasma* species (Figure 4). The *Anaplasma* sp. Tick314 sequence was found in a cluster including *Candidatus* Anaplasma boleense, *Anaplasma platys* and *Anaplasma phagocytophilum*.

Figure 4. Phylogenetic analysis of 16S rRNA sequences of *Anaplasma* spp. Phylogenetic analysis of 16S rRNA sequences of *Anaplasma* spp. using the maximum likelihood method based on the Tamura-

Nei model. In the phylogenetic tree, GenBank sequences, species designations and strain names are
given. The sequences investigated in the present study are marked with a black triangle (*Anaplasma*
sp. Tick283, accession number MK049844) and a black square (*Anaplasma* sp. Tick314, accession
number MK049845). The tree with the highest log likelihood (-473.44) is shown. The percentage of
trees in which the associated taxa clustered together is shown above the branches (bootstrap values).
The analysis involved 12 nucleotide sequences. There were a total of 243 positions in the final dataset.

Borrelia spp. were detected in both tick species from both islands (Figure 2). Infection rates
reached 5.1% in *A. variegatum* and 0.6% and 4.3% in *R. microplus* from Guadeloupe and Martinique,
respectively (Figure 2). None of the specific targeted *Borrelia* species causing Lyme disease (*Borrelia*
burgdorferi sensu lato), or the *Borrelia* relapsing fever group, gave any positive results, suggesting the
occurrence of a new or unexpected *Borrelia* spp. in our samples (Figure 2). We tested 30 of the *Borrelia*
spp.-positive ticks by nested PCR targeting the flaB genes. Interestingly, we obtained two sequences
according to the tick species analyzed. The *Borrelia* sp. Tick7 (accession number MK049846) sequence
was recovered from one *A. variegatum* sample from Guadeloupe, and the *Borrelia* sp. Tick457 sequence
(accession number MK049847) was recovered from four *R. microplus* samples from Martinique (Table
2). Phylogenetic and genetic distance analyses were performed using a portion of the flaB gene of
several *Borrelia* species (Figure 5). Surprisingly, the *Borrelia* sp. Tick7 sequence recovered from the *A.*
variegatum sample, and found to be closely related to *Bo. anserina*, displayed an intermediate position,
sharing homology with both the relapsing fever and Lyme disease groups (Figure 5). Lastly, the
Borrelia sp. Tick457 sequence recovered from the *R. microplus* samples confirmed the previous
observations, forming a cluster with various relapsing fever *Borrelia* species encountered in hard
ticks, including *Bo. lonestari* and *Bo. theileri* (Figure 5).

Figure. 5. Phylogenetic analysis of flaB sequences of *Borrelia* spp. Phylogenetic analysis of flaB
sequences of *Borrelia* spp. using the maximum likelihood method based on the Tamura-Nei model. In
the phylogenetic tree, GenBank sequences, species designations and strain names are given. The
sequences investigated in the present study are marked with a black circle (*Borrelia* sp. Tick457,
accession number MK049847) and a black triangle (*Borrelia* sp. Tick7, accession number MK049846).
The Lyme disease and relapsing fever clades of *Borrelia* are marked. The tree with the highest log
likelihood (-963.24) is shown. The percentage of trees in which the associated taxa clustered together
is shown above the branches (bootstrap values). The analysis involved 16 nucleotide sequences. There
were a total of 245 positions in the final dataset.

Lastly, 0.7% of the *R. microplus* ticks from Martinique were positive for *Leishmania* spp. (Figure
2). We tested two of the *Leishmania* spp.-positive ticks by nested PCR targeting the small sub-unit
rRNA gene. We obtained one sequence from one sample, deposited under the name *Leishmania*

martiniquensis Tick389 (accession number MK049850) (Table 2). This sequence displayed 100%
identity with both the *Leishmania martiniquensis* and *Leishmania siamensis* sequences (Table 2).

2.2.3. Co-infections in ticks in Guadeloupe and Martinique

We analyzed the co-infections observed in *Amblyomma variegatum* (n=132 samples), *Rhipicephalus*
microplus collected in Guadeloupe (n=116 samples, including individual and pooled specimens) and
Martinique (n= 275 samples, including individual and pooled specimens). In Guadeloupe, almost all
of the *A. variegatum* samples (99.2%) were infected with at least one pathogen whereas only 56% of
the *R. microplus* samples were infected (Figure 6). In contrast, 81% of the *R. microplus* from Martinique
were infected with at least one pathogen (Figure 6). High and similar percentages of the two tick
species were infected with either one or two pathogens. The percentages drastically dropped for co-
infection with three pathogens, with less than 10% of the ticks infected. Respectively one and nine *A.*
variegatum and *R. microplus*, from Guadeloupe and Martinique, were co-infected with four pathogens
and one *R. microplus* from Martinique was found infected with five pathogens (Figure 6).
287

288
289 **Figure 6.** Co-infections detected in *Amblyomma variegatum* (n=132 samples) and *Rhipicephalus microplus*
290 collected in Guadeloupe (n=116 samples) and Martinique (n=275 samples).
291

A. variegatum from Guadeloupe were find heavily infected by *R.africae*, yet it did not seem to
affect the presence of other pathogen/microorganisms that were all find in co-infection with the
bacteria (Table A3). Interestingly, in *R. microplus* from Guadeloupe, most of the single-infection
reported corresponded to *R. africae* (12.9%) or *T. velifera* (21.6%) (Table A3). Positive association have
been identified between *T. velifera* and *T. mutans*, and *Anaplasma* spp. / *Borrelia* spp. (Table A5).
Finally, in *R. microplus* from Martinique, five positive associations have been detected, including *T.*
mutans / *T. velifera*, *T. mutans* / *Leishmania* spp., *T. mutans* / *Borrelia* spp., *T. velifera* / *B. bigemina*, *A.*
marginale/ Ehrlichia spp. (Table A6). The result of the co-occurrence test should be taken with
cautions and deserve further investigation regarding the few number of positive samples (Table A5-
A6). Nevertheless, no exclusion seemed to occur between the pathogens/microorganisms detected in
the two tick species from Guadeloupe and Martinique. More details on co-infections in ticks from
Guadeloupe and Martinique are available in Appendix B.

3. Discussion

In this study, a high-throughput microfluidic real-time PCR system based on the use of multiple
primers/probes was developed for large-scale surveys of bacteria and protozoans potentially
transmitted by ticks from the Caribbean area. The association of genus and species primer/probe
designs targeting TBPs improved the technology's screening capacity, enabling not only the
identification of infectious agents known to circulate in the studied area, but also the detection of

unsuspected TBPs and new microorganisms belonging to the main bacterial and protozoan
genera/phyla involved in TBDs worldwide. Nevertheless, as some endosymbiotic microorganisms
may belong to known TBP genera, such as *Rickettsia* and *Coxiella*, confirmatory tests are required
before suggesting the presence of a pathogenic microorganism [23-25]. When analyzing the
specificity of the microfluidic real-time PCR system, cross-reactions were observed for some designs
targeting closely related species; these must be taken into account when interpreting the results. Due
to high design constraints and a lack of available sequences in public databases, the improvement of
such cross-reacting oligonucleotides remains challenging. Here, the concomitant use of bacterial and
protozoan genera can assist in identifying non-specific signals. In addition to detecting
microorganisms, we developed sets of primers and probes enabling the molecular identification of
the three main tick species involved in TBDs in the Caribbean: *A. variegatum*, *R. microplus* and *R.*
sanguineus s.l. As the morphological identification of ticks collected in the field remains challenging,
molecular identification can be used to confirm the identification of the tick species analyzed.

We used the newly developed high-throughput microfluidic real-time PCR system to perform
an exploratory epidemiological study of TBPs and microorganisms potentially circulating in
Caribbean ticks. The analysis provided an overview of the diversity of microorganisms belonging to
the main bacterial and protozoan genera potentially transmitted by ticks. It enabled the detection
both of known TBPs of public and animal health importance in the area that require surveillance and
of unexpected microorganisms occurring in Caribbean ticks.

The four main pathogens responsible for ruminant diseases in the Caribbean – currently
classified as notifiable diseases by the World Organisation for Animal Health (OIE) – have been
detected by the microfluidic real-time PCR system. These are *E. ruminantium* in *A. variegatum*
specimens and *An. marginale*, *B. bigemina* and *B. bovis* in *R. microplus*.

Interestingly, the *E. ruminantium* infection rate in *A. variegatum* reported in our study was much
lower compared to in previous studies conducted between 2003 and 2005 in Guadeloupe (5.1% versus
36.7%) [12]. Although different study designs were used (different sampling strategies, study
periods, detection methods, etc.), which may explain this difference, it would be worth further
investigating whether the tick infection rate for *E. ruminantium* has decreased in Guadeloupe and
possibly assessing the epidemiological impact in terms of the incidence and prevalence of heartwater
in the ruminant population. These results are all the more surprising since systematic TBT
surveillance and control programs have been discontinued following the end of the POSEIDOM
eradication programs in 2006.

In this study, we have documented infection rates for *B. bigemina*, *B. bovis* and *An. marginale* in
the *R. microplus* vector tick in the French West Indies for the first time. Indeed, records of such
pathogens are mostly based on seroprevalence studies in cattle [7,8,10].

R. microplus ticks are both vectors and reservoirs of *B. bigemina* and *B. bovis*, transmitting the
parasites transovarially and trans-stadially [26,27]. As *R. microplus* ticks and cattle are both reservoirs
of infection, the infection rates reported here seemed quite low. The life cycle of *Babesia* spp. requires
complex interactions with its two hosts, which are the tick vector and the vertebrate host. The
efficiency of tick acquisition and of transovarial and trans-stadial transmission of *B. bovis* and *B.*
bigemina by *R. microplus*, involved in the long-term persistence of *Babesia* spp. in nature, is still poorly
understood and warrants further investigations [26,27].

Interestingly, *An. marginale* was detected in *R. microplus* from both islands, but the infection rate
reported in ticks from Guadeloupe seemed lower compared to in Martinique. The same trend had
been reported during previous seroprevalence studies [7,8,10]. Anaplasmosis can be transmitted by
vectors other than ticks, and some cattle breeds are known to be more susceptible than others to
Anaplasma infection [10]. The difference in *Anaplasma* infection rate in ticks between the two islands
may have been due to differences in the cattle populations. Indeed, there are mainly local Creole and
mixed European-Creole breeds in Guadeloupe. These are known to be more resistant to anaplasmosis
than Brahman and European breeds, which are the main breeds reared in Martinique [10]. In
addition, other factors, including differences in the population dynamics of alternate vectors such as
flies, may also have contributed to this difference.

Among the other known TBPs detected, we also found pathogens with low health impact in the
Caribbean – almost considered as endosymbionts – such as *R. africae*, *T. velifera* and *T. mutans* in their
A. variegatum vector and surprisingly in *R. microplus* ticks.

With almost all of the *A. variegatum* found to be infected, the *R. africae* infection rate was the
highest ever reported in the Caribbean [9,13,14,28]. As *A. variegatum* is both the vector and the
reservoir of the pathogen, with transovarial and trans-stadial transmission rates reaching 100%, this
high level of *R. africae* infection is not surprising per se [14,29]. Interestingly, the high *R. africae*
infection rate in vector ticks, associated with a very low number of African tick-bite fever cases in the
Caribbean, highlights the difficulty, in some cases, of clearly distinguishing between endosymbiosis
and pathogenicity [9,15]. The biological relationship between *R. africae* and *A. variegatum* as well as
the strain variety and virulence of *R. africae* in the Caribbean should be investigated in order to better
assess risks and guide prevention measures, especially for travelers [23,24,30]. The absence of direct
identification of *R. africae* in *R. microplus* ticks was probably due to lower sensitivity of the specific
target design compared to the genus target design. Indeed, *Rickettsia* spp.-positive *R. microplus*
samples displayed rather high Ct values, suggesting a low infection level that may have been below
the detection limit for *R. africae*. The unusual presence of *R. africae* in *R. microplus* ticks may have been
due to the co-occurrence of the two tick species, *R. microplus* and *A. variegatum*, on cattle. As the ticks
here were collected partially engorged, the presence of *R. africae* in *R. microplus* may have been due
to bacteria circulating in cattle blood picked up by engoring ticks, or to cross-contamination with *R.*
microplus ticks co-feeding next to infected *A. variegatum* [31,32].

This study provides the first update on the detection of *T. mutans* and *T. velifera* in Caribbean
ticks. Indeed, references to these parasites in the Caribbean are relatively old, and no prevalence
studies have been conducted since, whether in ticks or in cattle [5,6,33]. The low pathogenicity of
these piroplasms may explain the lack of diagnoses and the scarcity of information available on their
distribution and prevalence in the Caribbean. However, these parasite species may play an important
role in theileriosis management and protection, as chronically infected cattle can develop immunity
and heterologous protection against other pathogenic *Theileria* species, such as *Theileria parva* [34].
Unfortunately, we did not succeed in confirming these results by conventional or nested PCR,
suggesting either a level of infection below the detection threshold, or simply false signals.

Lastly, the high-throughput microfluidic real-time PCR system enabled the detection of
unexpected and/or potentially new microorganisms, leading to the recovery of nucleotide sequences
of *Anaplasma* spp., *Ehrlichia* spp., *Borrelia* spp. and *Leishmania* spp. in ticks collected in Guadeloupe
and Martinique.

The *Ehrlichia* sp. Tick428 sequence detected here formed a cluster with other uncharacterized
Ehrlichia species detected in ticks from Asia and Africa [13,35-39]. However, given the highly
conserved nature of the 16S rRNA genes, we could not more accurately define phylogenetic
relationships within the *Ehrlichia* species group. The *Anaplasma* sp. Tick314 sequence was identified
as *Candidatus* Anaplasma boleense, a bacterium described in ticks and mosquitoes in China [38,40].
No further information is available regarding the epidemiology of *Candidatus* Anaplasma boleense.
These observations highlight the need to set up characterization studies. Indeed, high-throughput
detection technologies can highlight the presence of DNA from potentially new microorganisms, but
it will still be necessary to isolate and characterize them in order to first confirm their existence and
then determine whether their presence in ticks poses a risk to public or animal health.

Here we provided the first report of *Borrelia* spp. in ticks from Guadeloupe and Martinique. Two
different sequences were recovered, according to the tick species analyzed. In *A. variegatum*, a
sequence named *Borrelia* sp. Tick7 was detected and was closely related to *Bo. anserina*, the agent of
avian spirochetosis. Both of them seemed to define an intermediate position between the relapsing
fever and Lyme disease groups. In contrast, the *Borrelia* sp. Tick457 sequence found in *R. microplus*
sample, clustered with uncharacterized *Borrelia* spp. described *R. microplus* specimens from
Madagascar and Brazil, such as *Borrelia* sp. strain Mo063b and *Borrelia* sp. BR, and with relapsing
fever Borrelia species encountered in hard ticks, including *Borrelia lonestari* and *Bo. theileri* [41,42].
Interestingly, the same observations had recently been made regarding *Borrelia* spp. found in *A.*

variegatum and *R. microplus* ticks from Ethiopia and Côte d'Ivoire [43,44]. As *A. variegatum* and *R.*
microplus were imported into the Caribbean from Africa during the time of the Atlantic triangular
trade, we may have detected bacteria probably characterized by an old introduction through infected
ticks and subsequent local evolution within their vector over a long period [4,45]. *Borrelia* spp. and
borreliosis case reports in the Caribbean are scarce and still being debated. In Cuba, one study
suggested the presence of antibodies to *Borrelia burgdorferi* sensu stricto in human sera associated
with clinical cases of Lyme disease-like syndrome [46,47]. However, the real specificity of these serum
antibodies has been questioned [48]. In the US Virgin Islands, seropositivity for *Borrelia hermsii* and
closely related species was reported in association with a human case of relapsing fever [49]. Lastly,
erythema migrans-like skin lesions and illness were reported in four Caribbean nationals [50].
Regarding the importance of *Borrelia* spp. for human and animal health, the characterization of these
potential new *Borrelia* species that seemed associated with tropical tick species requires further
investigation.
Lastly, *Leishmania* spp. were detected in *R. microplus* specimens from Martinique, and one
sequence was identified as *Leishmania martiniquensis* Tick389 (accession number MK049850). Studies
on *Leishmania* nomenclature have highlighted the fact that isolates of "*L. siamensis*" have never been
officially characterized and that therefore, this name should not be used [51-54]. Thus, since all the
sequences – except one – reported as "*L. siamensis*" in databases should be considered as synonyms
of *L. martiniquensis*, we assumed the occurrence of *L. martiniquensis* here. Parasites of the genus
Leishmania are usually transmitted by female phlebotomine sand flies (Diptera: Psychodidae:
Phlebotominae) and generally involve a wide variety of animal species, mainly including dogs and
canids in the epidemiological cycle. They are responsible for leishmaniasis, a zoonosis widespread in
tropical and sub-tropical areas [54]. *L. martiniquensis* belongs to the *L. enriettii* complex and has been
described in Martinique and Thailand, where it was responsible for both cutaneous and visceral
leishmaniosis [51,54-56]. *L. martiniquensis* is suspected to be endemic in Martinique [55]. Although
phlebotomines and rodents are present in Martinique, neither vectors nor reservoirs of this parasite
have yet been described [55]. Our study represents the first report of *L. martiniquensis* in *R. microplus*
ticks from the French West Indies. Although *Leishmania* spp. have been reported in ticks (*L. infantum*
in *R. sanguineus s.l.*, and *L. guyanensis* in *R. microplus* ticks in Peru, for example), the role of ticks in
Leishmania transmission is still being debated, and no evidence of vector capacity has been reported
yet [57-59]. Moreover, the finding of *Leishmania* spp. in a tick species that feeds mainly on cattle also
raises questions about the potential role of cattle in the epidemiology of leishmaniasis [60,61]. The
participation of ticks in *Leishmania* epidemiology warrants further investigation, especially since *R.*
microplus ticks could parasitize humans [62].
Surprisingly, co-infections with two or more TBPs were found in more than 50% of the infected
ticks, both for *A. variegatum* and *R. microplus* and on the two islands. In addition, we could not identify
any exclusion of infection between pathogens. These observations illustrate the efficiency of ticks as
reservoirs of multiple pathogens with no apparent significant effects on their life traits.
To conclude, although screening tools are useful for the discovery of pathogens in ticks, the
epidemiological significance of such results warrants further analysis. Detecting a microorganism's
DNA in ticks, especially in partially engorged ticks, does not necessarily mean that the ticks are
involved in the microorganism's life cycle; however, it provides useful information to supplement
vector competence studies [16]. Nevertheless, the detection of potentially new microorganisms in
ticks from the French West Indies has opened up new research perspectives for the future on the
epidemiology of TBPs in the Caribbean. A region-wide epidemiological survey on TBPs in ticks
collected in different countries and territories of the Caribbean area, organized in collaboration with
the Caribbean Animal Health Network (CaribVET) in order to strengthen our results, may be an
interesting way to supplement and strengthen some of this paper's findings.

4. Materials and Methods

4.1 Ticks collected in Guadeloupe and Martinique

The ticks used in this study were collected as part of two separate epidemiological surveys
conducted in Guadeloupe (between February 2014 and January 2015) and Martinique (between
February and March 2015), respectively. In Guadeloupe, adult ticks (any species, any sex) were
collected from 40 cattle originating from 22 different herds that were sampled in nine localities
situated in six different biotopes (urban area, dry coastal regions, valleys and hills, evergreen seasonal
forest, sub-mountainous rainforest, swamp forest). In Martinique, engorged females of *R. microplus*
only were collected from cattle in 29 farms participating in a study on acaricide resistance in ticks.
All the ticks were collected from cattle with the permission of farmers and cattle owners. The ticks
were morphologically identified at species level [63]. A total of 578 adult ticks were included in the
study: 132 *A. variegatum* and 165 *R. microplus* ticks from Guadeloupe and 281 *R. microplus* ticks from
Martinique (see maps, Figure 2). The GPS coordinates of the tick collection sites are available in Table
S1. All the ticks were partially engorged, and then stored at -80°C.

4.2 DNA extraction of ticks collected in Guadeloupe and Martinique

For 20 mg of tick, 1 ml of recently prepared PBS 1X was added to the sample. The ticks were
then washed by gently shaking for 2-3 min at 7 Hz/s in a TissueLyser (Qiagen, Germany). After
discarding the supernatant, the ticks were frozen at -80°C for 15-20 min. A steel ball was then added
and the samples were crushed twice for 2 min at 30 Hz/s with the TissueLyser (Qiagen, Germany).
450 µl of fresh PBS 1X were added to the samples. The samples were vortexed for 10 s and then
centrifuged for 2-3 min at 8000 g. Lastly, 20 µl of Proteinase K were added to 180 µl of crushed tick
sample and DNA was extracted using the NucleoSpin® 96 Virus Core Kit (Macherey-Nagel,
Germany) and the Biomek4000 automated platform (Beckman Coulter). This protocol enables the
simultaneous extraction of both DNA and RNA. Total nucleic acid per sample was eluted in 160 µl
of rehydration solution and stored at -80°C until further use. *A. variegatum* ticks were individually
extracted. *R. microplus* ticks were extracted both individually and in pools of two to four adult
specimens when they were too small to be treated individually.

4.3 Assay design

The list of pathogens to be monitored, the sets of primers and probes required for their detection,
as well as the targeted genes are shown in Table 1. Some of the oligonucleotides were specifically
designed for the purposes of this study; the others came from Michelet et al., 2014 [18]. The newly
developed oligonucleotides were validated for a range of dilutions of positive controls, including
cultures, plasmids and DNA samples (Table 1, Table S2), by real-time TaqMan PCR assays on a
LightCycler® 480 (LC480) (Roche Applied Science, Germany). Real-time PCR assays were performed
with LightCycler® 480 Probe Master Mix 1× (Roche Applied Science, Germany), using 200 nM of
primers and probes in a final volume of 12 µl, and 2 µl of control DNA were added. The thermal
cycling program was as follows: 95°C for 5 min, 45 cycles at 95°C for 10 s and 60°C for 15 s, and one
final cooling cycle at 40°C for 10 s.

4.4 Pre-amplification of DNA samples

All the DNA samples were subject to pre-amplification in order to enrich the pathogenic DNA
content compared with tick DNA. PerfeCTa® PreAmp SuperMix (Quanta Biosciences, Beverly, USA)
was used for DNA pre-amplification, following the manufacturer's instructions. All the primers were
pooled (except those targeting the tick species), with a final and equal concentration of 45 nM each.
The pre-amplification reaction was performed in a final volume of 5 µl containing 1 µl of PerfeCTa
PreAmp SuperMix (5X), 1.25 µl of pooled primer mix, 1.25 µl of DNA and 1.5 µl of Milli-Q water,
with one cycle at 95°C for 2 min and 14 cycles at 95°C for 10 s and 60°C for 3 min. At the end of the
cycling program, the reactions were 1:10 diluted. The pre-amplified DNA were stored at -20°C until
use.
510
4.5 High-throughput microfluidic real-time PCR

High-throughput microfluidic real-time PCR amplifications were performed using the
BioMark™ real-time PCR system (Fluidigm, USA) and 96.96 dynamic arrays (Fluidigm, USA),
enabling up to 9,216 individual reactions to be performed in one run [18]. Real-time PCRs were
performed using 6-carboxyfluorescein (6-FAM)- and Black Hole Quencher (BHQ1)-labeled TaqMan
probes with TaqMan Gene Expression Master Mix (Applied Biosystems, USA) following the
manufacturer's instructions. The cycling conditions were as follows: 2 min at 50°C and 10 min at
95°C, followed by 40 cycles of two-step amplification for 15 s at 95°C and 1 min at 60°C. The
BioMark™ real-time PCR system was used for data acquisition and the Fluidigm real-time PCR
analysis software for Ct value determination. Three kinds of controls per chip were used for
experiment validation: a negative water control to exclude contamination; a DNA extraction control,
corresponding to primers and probes targeting a portion of the 16S rRNA gene of ticks; and an
internal control, to check the presence of PCR inhibitors made of DNA from Escherichia coli strain
EDL933, added to each sample with specific primers and probes targeting the eae gene [64]. For the
relative specificity analysis of the newly designed Biomark system, DNA of 62 positive controls were
used as template (Table S2). Then, for the epidemiological survey of TBPs in Caribbean ticks, the 523
DNA samples of *A. variegatum* and *R. microplus* from Guadeloupe and Martinique were used as
template.

4.6 Infection rates for ticks from the French West Indies

Depending on the tick species and the island of origin, for each detected pathogen, infection
rates (the proportion of infected ticks divided by the total number of ticks analyzed) were calculated.
The majority of the samples were single specimens of ticks. When ticks were too small to be treated
individually, they were grouped into pools of two to four specimens. Thus, of the 523 samples
analyzed, 47 consisted of a pool of two to four tick specimens. The final estimation of infection rates
also includes the pools and is therefore expressed as the minimum (assuming at least one positive
tick in the pools) and maximum (assuming all positive ticks in the pools) proportions of infected ticks
out of the total number of ticks analyzed.

4.7 PCRs and sequencing for the confirmation of results

Conventional PCRs/nested PCRs using primers targeting different genes or regions than those
of the BioMark™ system were used to confirm the presence of pathogenic DNA in some field samples
and positive controls (Table 3). Amplicons were sequenced by Eurofins MWG Operon (BIOMNIS-
EUROFINS GENOMICS, France) and then assembled using BioEdit software (Ibis Biosciences,
Carlsbad, CA, USA). An online BLAST (Basic Local Alignment Search Tool) search was used to
compare the nucleotide sequences found in this study to reference sequences listed in GenBank
sequence databases (NCBI).

Table 3. Primers used to confirm the presence of pathogenic DNA in tick samples, and positive
controls.

Pathogen	Targeted gene	Primer name	Sequence (5' → 3')	Length (bp)	References
Rickettsia spp.	gltA	Rsfg877	GGGGGCCTGCTCACGGCGG	381	[65]
		Rsfg1258	ATTGCAAAAAGTACAGTGAACA		
	ompB	Rc.rompB.4362p	GTCAGCGTTACTTCTTCGATGC	475	[66]
		Rc.rompB.4,836n	CCGTACTCCATCTTAGCATCAG		
		Rc.rompB.4,496p	CCAATGGCAGGACTTAGCTACT	267	
		Rc.rompB.4,762n	AGGCTGGCTGATACACGGAGTAA		
Anaplasma/Ehrlichia spp.	16S rRNA	EHR16SD	GGTACCYACAGAAGAAGTCC	345	[67]
		EHR16SR	TAGCACTCATCGTTTACAGC		

Borrelia spp.	flaB	FlaB280F	GCAGTTCARTCAGGTAACGG	645	[68]
		FlaRL	GCAATCATAGCCATTGCAGATTGT		
		FlaB737F	GCATCAACTGTRGTTGTAACATTAACAGG	407	
		FlaLL	ACATATTCAGATGCAGACAGAGGT		
Leishmania spp.	SSU rRNA	R221	GGTTCCTTTCCTGATTTACG	603	[69]
		R332	GGCCGGTAAAGGCCGAATAG		
		R223	TCCATCGCAACCTCGGTT	358	
		R333	AAAGCGGGCGCGGTGCTG		

4.8 Phylogenetic sequence analysis

Alignments were performed using ClustalW [70]. Maximum likelihood trees were generated by 1,000
bootstrap repetitions based on the Tamura-Nei model [71] in MEGA7 [72]. The initial tree(s) for the
heuristic search were obtained automatically by applying neighbor-joining and BioNJ algorithms to
a matrix of pairwise distances estimated using the maximum composite likelihood (MCL) approach
and then selecting the topology with superior log likelihood value. The tree was drawn to scale, with
branch lengths measured in the number of substitutions per site. The codon positions included were
1st+2nd+3rd+Noncoding. All positions containing gaps and missing data were eliminated. Further
information is provided in the figure legends.

5. Conclusions

Our study demonstrated the high ability of microfluidic real-time PCR technology to provide a
rapid overview of the diversity of TBPs of veterinary and medical importance present in ticks from
the Caribbean. This innovative high-throughput tool is promising and could significantly improve
the surveillance and exploration of TBPs, enabling the rapid screening of multiple microorganisms
especially in regions where few epidemiological data are available and TBDs are numerous.

Supplementary Materials: The following are available online at www.mdpi.com/xxx/s1, Table S1: GPS
coordinates of the tick collection sites and number of ticks collected. A total of 578 adult ticks collected from
cattle from Guadeloupe and Martinique were used for the screening of tick-borne pathogens with the newly
implemented BioMark™ real-time PCR system, Table S2: List of the positive control samples used for the relative
specificity analysis of the Biomark system developped in this study, Figure S1: Improvement of detection signals
by pre-amplification. Test of primer/probe set sensitivity for a range of dilutions of positive controls by TaqMan
real-time PCR using LightCycler 480, before and after pre-amplification. Results of the sensitivity test of the
Leishmania infantum design using a *Leishmania infantum* culture, before (a) and after (c) pre-amplification; Results
of the sensitivity test of the *Rickettsia* spp. design using *Rickettsia conorii*-positive controls (extracted from an
infected *Rhipicephalus sanguineus* sensu lato tick), before (b) and after (d) pre-amplification.

Author Contributions: All authors have read and agree to the published version of the manuscript.
Conceptualization, E.A., J.P. and S.M.; methodology, M.G., E.D., V.P., R.A., C.G., S.D. and M.V.T.; formal
analysis, M.G.; investigation, M.G.; writing—original draft preparation, M.G.; writing—review and editing,
M.V.T., E.A., J.P. and S.M.; supervision, E.A., J.P. and S.M.; funding acquisition, M.V.T., E.A., J.P. and S.M.

Funding: This research was funded by grants from the French Agency for Food, Environmental and
Occupational Health & Safety (ANSES, CoVetLab grant), the French Agricultural Research Centre for
International Development (CIRAD) and the French National Institute for Agricultural Research (INRA,
PATHO-ID metaprogram). This study was partly supported by the MALIN project on "Surveillance, diagnosis,
control and impact of infectious diseases of humans, animals and plants in tropical islands", supported by the
European Union, in the framework of the European Regional Development Fund (ERDF), and the Regional
Council of Guadeloupe. The study used ticks collected in Guadeloupe and Martinique as part of previous
projects: the PathoID "Rodent and tick pathobiome" project funded by INRA and the ResisT project on
"Assessment of tick resistance to acaricides in the Caribbean – Development of strategies to improve surveillance
and control of tick-borne diseases in ruminants", funded by the Regional Cooperation Fund (FCR) of
Guadeloupe and Martinique. This work was performed in the framework of EurNegVec COST Action TD1303.

Acknowledgments: In this section you can acknowledge any support given which is not covered by the author
contribution or funding sections. This may include administrative and technical support, or donations in kind
(e.g., materials used for experiments).

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the
study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to
publish the results.

Appendix A

Analysis of the relative specificity of the 61 sets of primers and probe constituting the Biomark
system developed in this study.

In order to identify potential cross-reactions, we analyzed the relative specificity of the 61 sets
of primers and probe constituting the BioMark system used in this study using 62 positive control
samples including DNA from bacterial or parasitic cultures, or DNA from tick or blood samples
known to be infected, or plasmidic constructions (see Table S2).

Of the 61 designs, 42 designs were specific of their target. The Tick spp. design, used as a tick
nucleic acid extraction control, was able to detect *A. variegatum* and *R. sanguineus* s.l. samples as well
as the DNA of the *R. sanguineus* s.l. tick present in the *Rickettsia conorii* positive control as expected
(Table A1). However, the DNA of ticks from the *R. microplus* control sample and other positive
controls including tick DNA (such as the *Borrelia lonestari*, *Anaplasma phagocytophilum* controls, etc.)
were not detected (Table A1). The detection ability of this design was corrected by adding the Tick
spp. primers during the pre-amplification step; these had initially been excluded since the objective
was to enrich pathogenic DNA content compared to tick DNA (data not shown). Eight designs
displayed cross-reactions with one to two closely related species, and seven designs displayed
unexpected signals corresponding likely to the detection of unexpected co-infection in complex
control samples such as DNA extracted from ticks or blood samples (Table A1). Finally three design
were removed from the system, one design due to a lack of efficiency (no detection of the target), and
two designs were not specific, displaying multiple cross-reactions (Table A1).

Table A1. List of designs and their specificity using the BioMark system. CR: cross-reactions with
closely related species samples; CI: potential co-infections in control samples.

Design	Target detection	Specificity	Outgroup control samples
Rickettsia spp. gltA	YES	CI	*Borrelia lonestari* (Infected *A. americanum* ticks)
	YES	CI	*Ehrlichia chaffensis* (Infected *A. americanum* ticks)
	YES	CI	*Ehrlichia ewingii* (Infected *A. americanum* ticks)
	YES	CI	*Panola mountain Ehrlichia* (Infected *A. americanum* ticks)
	YES	CI	*Amblyomma variegatum* (Wild tick)
Rickettsia massiliae 23S-5S ITS	YES	CR	*Rickettsia slovaca* (Culture)
Rickettsia rickettsii 23S-5S ITS	YES	CR [1]	*Rickettsia slovaca* (Culture)
	YES		*Rickettsia conorii* (Infected *R. sanguineus* s.l. ticks)
	YES		*Rickettsia africae* (Culture)
	YES		*Amblyomma variegatum* (Wild tick)
Rickettsia conorii sca1	YES	CR	*Rickettsia slovaca* (Culture)
Rickettsia africae sca1	YES	CI	*Amblyomma variegatum* (Wild tick)
Rickettsia felis orfB	YES	CI	*Borrelia lonestari* (Infected *A. americanum* ticks)
	YES	CI	*Ehrlichia chaffensis* (Infected *A. americanum* ticks)
	YES	CI	*Ehrlichia ewingii* (Infected *A. americanum* ticks)
	YES	CI	*Panola mountain Ehrlichia* (Infected *A. americanum* ticks)
Rickettsia typhi ompB	YES	YES	

Rickettsia prowazekii gltA	*YES*	*YES*	
Borrelia spp. 23S rRNA	*YES*	*YES*	
Borrelia burgdorferi sensu stricto *glpA*	*NO* [1]		
Borrelia anserina fla	*YES*	*YES*	
Borrelia lonestari glpQ	*YES*	*YES*	
Borrelia parkeri gyrB	*YES*	*YES*	
Borrelia bissettii rpoB	*YES*	*YES*	
Borrelia theileri glpQ	*YES*	*YES*	
Bartonella spp. ssrA	*YES*	*YES*	
Bartonella bacilliformis rpoB	*YES*	*CR*	*Bartonella henselae* (Culture)
Bartonella henselae ribC	*YES*	*CR*	*Bartonella bacilliformis* (Culture)
Bartonella vinsonii subsp. berkhoffii ITS	*YES*	*YES*	
Coxiella burnetii icd	*YES*	*YES*	
Coxiella burnetii IS 1111	*YES*	*YES*	
Francisella tularensis tul4	*YES*	*YES*	
Francisella tularensis fopA	*YES*	*YES*	
Anaplasma spp. 16S rRNA	*YES*	*YES*	
Anaplasma marginale msp1b	*YES*	*YES*	
Anaplasma phagocytophilum msp2	*YES*	*YES*	
Anaplasma platys groEL	*YES*	*YES*	
Anaplasma bovis groEL	*YES*	*YES*	
Anaplasma ovis msp4	*YES*	*YES*	
Ehrlichia spp. 16S rRNA	*YES*	*YES*	
Ehrlichia canis gltA	*YES*	*YES*	
Ehrlichia chaffeensis dsb	*YES*	*YES*	
Ehrlichia ewingii dsb	*YES*	*YES*	
Ehrlichia ruminantium gltA	*YES*	*YES*	
Panola Mountain Ehrlichia gltA	*YES*	*YES*	
Neoehrlichia mikurensis groEL	*YES*	*YES*	
Aegyptianella pullorum groEL	*YES*	*YES*	
Apicomplexa 18S rRNA	*YES*	CI	*Borrelia lonestari* (Infected *A. americanum* ticks)
	YES	CI	*Anaplasma marginale* (Infected cow blood)
	YES	CI	*Panola mountain Ehrlichia* (Infected *A. americanum* ticks)
	YES	CI	*Neoehrlichia mikurensis* (Infected rodent blood)
	YES	CI	*Rhipicephalus sanguineus* s.l. (Wild tick)
Babesia canis vogeli hsp70	*YES*	*CR*	*Babesia canis canis* (Infected dog blood)
Babesia ovis 18S rRNA	*YES*	*CR* [1]	*Rickettsia massiliae* (Culture)
	YES		*Borrelia lonestari* (Infected *A. americanum* ticks)
	YES		*Anaplasma marginale* (Infected cow blood)
	YES		*Ehrlichia chaffensis* (Infected *A. americanum* ticks)
	YES		*Ehrlichia ewingii* (Infected *A. americanum* ticks)
	YES		*Panola mountain Ehrlichia* (Infected *A. americanum* ticks)
	YES		*Neoehrlichia mikurensis* (Infected rodent blood)

	YES		*Babesia divergens* (Culture)
	YES		*Babesia canis rossi* (Infected dog blood)
	YES		*Babesia canis canis* (Infected dog blood)
	YES		*Babesia canis vogeli* (Infected dog blood)
	YES		*Babesia microti* (Culture)
	YES		*Theileria annulata* (Culture)
	YES		*Theileria lestoquardi* (Culture)
	YES		*Theileria parva* (Culture)
	YES		*Hepatozoon canis* (Infected dog blood)
	YES		*Rhipicephalus sanguineus* s.l. (Wild tick)
Babesia bigemina 18S rRNA	*YES*	CI	*Anaplasma marginale* (Infected cow blood)
Babesia gibsoni Rap1	*YES*	*YES*	
Babesia caballi Rap1	*YES*	*YES*	
Babesia bovis CCTeta	*YES*	*YES*	
Babesia duncani ITS2	*YES*	*YES*	
Babesia microti CCTeta	*YES*	CR	*Babesia canis canis* (Infected dog blood)
Theileria parva 18S rRNA	*YES*	CR	*Theileria annulata* (Culture)
	YES	CR	*Theileria lestoquardi* (Culture)
Theileria mutans ITS	*YES*	*YES*	
Theileria velifera 18S rRNA	*YES*	*YES*	
Theileria equi ema1	*YES*	*YES*	
Cytauxzoon felis ITS2	*YES*	*YES*	
Hepatozoon spp 18S rRNA	*YES*	CI	*Neoehrlichia mikurensis* (Infected rodent blood)
	YES	CI	*Rhipicephalus sanguineus* s.l. (Wild tick)
Hepatozoon canis 18S rRNA	*YES*	CI	*Rhipicephalus sanguineus* s.l. (Wild tick)
Hepatozoon americanum 18S rRNA	*YES*	CR	*Hepatozoon canis* (Infected dog blood)
	YES	CI	*Rhipicephalus sanguineus* s.l. (Wild tick)
Leishmania spp hsp70	*YES*	*YES*	
Leishmania infantum ITS	*YES*	*YES*	
Rangelia vitalii 18S rRNA	*YES*	*YES*	
Tick spp 16S rRNA	*YES*	*YES* [2]	
Amblyomma variegatum ITS2	*YES*	*YES*	
Rhipicephalus microplus ITS2	*YES*	*YES*	
Rhipicephalus sanguineus s.l. ITS2	*YES*	*YES*	

[1] Designs removed from the screening analysis.

[2] Tick spp. primers required to be part of the pre-amplification mix in order to achieve correct detection signals.

Regarding the seven designs displaying unexpected signals, we decided to explore the hypothesis of potential co-infection in controls corresponding to DNA from ticks or blood samples using conventional PCR and amplicon sequencing (Table A1).

Both the *Rickettsia* spp. and *Rickettsia felis* designs gave positive results when testing the *Borrelia lonestari*, *Panola Mountain Ehrlichia*, *Ehrlichia ewingii* and *Ehrlichia chaffensis* controls. All these samples corresponded to DNA extracted from infected *Amblyomma americanum* collected in the field (USA). This result supports the fact that a *Rickettsia* species may have been present in these samples, whether it was *Rickettsia felis* or a closely related *Rickettsia* spp. infecting *Amblyomma americanum* ticks. When

testing the Panola Mountain Ehrlichia sample for *Rickettsia* spp. by conventional PCR targeting the
gltA gene [65], we obtained a 382 bp sequence (accession number MK049843) sharing 99% sequence
identity with *Rickettsia* spp. (MF511253.1) (Table A2). This result, in addition to natural co-infections
documented in *Amblyomma americanum* ticks, suggest that the detection of *Rickettsia* spp. – within the
four outgroup positive controls corresponding to *Amblyomma americanum* DNA – in this assay did
not correspond to cross-reactions [73,74]. Moreover, the *Amblyomma variegatum* sample,
corresponding to ticks collected from the field, was found to be positive for both *Rickettsia* spp. and
Rickettsia africae. As this sample originated from Guadeloupe, where *Rickettsia africae* circulates, this
result may have been in agreement with a natural infection [9].

Table A2. Taxonomic assignment of the sequences obtained after sequencing PCR products to
confirm the presence of co-infections in complex control samples corresponding to DNA extracted
from wild ticks or blood samples. AN: accession number, % I: percentage identity, % C: percentage
coverage.

Tested control sample	Tested for	AN	Length (bp)	Closest homology	% I	% C	AN
Panola mountain Ehrlichia (Infected *A. americanum* ticks)	*Rickettsia* spp.	MK049843	382	*Rickettsia* spp.	99	100	MF511253.1
Neoehrlichia mikurensis (Infected rodent blood)	*Hepatozoon* spp.	MK071735	169	*Hepatozoon* spp.	99	100	AB771515.1
Anaplasma marginale (Infected cow blood)	Apicomplexa	MK071737	104	*Babesia* spp., *Theileria* spp. (including *B. bigemina*)	100	99	MG604302.1
Panola mountain Ehrlichia (Infected *A. americanum* ticks)	Apicomplexa	MK071736	102	*Theileria* spp. (including *T. cervi*)	98	100	MH085203.1
Borrelia lonestari (Infected *A. americanum* ticks)	Apicomplexa						

639
Moreover, the *Rhipicephalus sanguineus* s.l. sample was positive with four designs targeting
parasites – Apicomplexa, *Hepatozoon* spp., *Hepatozoon canis* and *Hepatozoon americanum*. Since
Rhipicephalus sanguineus s.l. ticks are involved in the epidemiology of tick-borne parasites including
Hepatozoon spp., these results strongly suggest the occurrence of such parasites in the biological
sample [75]. Thus, these observations suggest that our designs were actually capable of detecting
pathogens present in naturally infected ticks rather than giving an unsuspected cross-reaction. The
Neoehrlichia mikurensis sample, corresponding to DNA extracted from rodent blood, was also found
to be positive for both Apicomplexa and *Hepatozoon* spp. The amplicon obtained from this sample
with the *Hepatozoon* spp. design was sequenced. The obtained 169 bp sequence (accession number
MK071735) displayed 99% sequence identity with the *Hepatozoon* spp. sequences (AB771515.1) (Table
A2). As rodents can be infected with *Hepatozoon* parasites, this result could also reflect a natural
infection [76]. The *Anaplasma marginale* sample, corresponding to a blood sample from an
experimentally infected cow, was also found to be positive for both Apicomplexa and *Babesia*
bigemina. The amplicon obtained from this sample with the Apicomplexa design was sequenced.
The obtained 104 bp sequence (accession number MK071737) displayed 99% sequence identity with
the Apicomplexan sequences, including *Babesia bigemina* (MG604302.1) (Table A2). As *Babesia*
bigemina and *Anaplasma marginale* are two cattle pathogens that have often evolved in the same region
and are transmitted by the same vector tick, co-infections with these two pathogens have already
been reported [77]. Thus, this cross-reaction may again have been a reflection of parasite co-infections.
Lastly, Apicomplexa also gave positive results in the *Borrelia lonestari* and *Panola Mountain Ehrlichia*
controls. As highlighted previously, these two controls corresponded to DNA extracted from *A.*
americanum ticks. The amplicons obtained from these two samples with the Apicomplexa design were
sequenced. The two obtained 102 bp sequences (accession number MK071736) were identical and

displayed 98% sequence identity with the *Theileria cervi* sequences (MH085203.1) (Table A2). As
Theileria cervi is a common deer pathogen found in *Amblyomma americanum*, the occurrence of this
parasite in these two control samples could explain the unexpected signals [78,79].

Appendix B

Analysis of the co-infections detected in *Amblyomma variegatum* and *Rhipicephalus microplus* ticks
collected in Guadeloupe and Martinique.
Here we reported the co-infections observed in *Amblyomma variegatum* (n=132 samples),
Rhipicephalus microplus collected in Guadeloupe (n=116 samples, including individual and pooled
specimens) and Martinique (n= 275 samples, including individual and pooled specimens) (Table A3).
We combined the results obtained with the *Rickettsia* spp. and the *R. africae* design, assuming that
only *R. africae* have been detected in the Caribbean samples analyzed here. In addition, *Anaplasma*
spp. results here correspond to sample only positive for *Anaplasma* spp. and not for *A. marginale*, as
well, *Ehrlichia* spp. results here correspond to sample only positive for *Ehrlichia* spp. and not for *E.*
ruminantium.

Table A3. Co-infection analysis in *A. variegatum* (n=132 samples) and *R. microplus* collected in
Guadeloupe (n=116 samples) and Martinique (n=275 samples).

		A. variegatum sample (percentage)	*R. microplus* sample from Guadeloupe (percentage)	*R. microplus* sample from Martinique (percentage)
	Total of sample	132	116	275
	Total of non-infected sample	1 (0.8%)	51 (44%)	52 (19%)
	Total of infected sample	131 (99.2%)	65 (56%)	223 (81%)
Single infections	Total	69 (52.3%)	47 (40.5%)	99 (36%)
	R. africae [1]	68 (51.5%)	15 (12.9%)	0
	Anaplasma spp. [2]		1 (0.9%)	10 (3.6%)
	A. marginale	0	2 (1.7%)	22 (8%)
	Ehrlichia spp. [3]	0	3 (2.6%)	43 (15.6%)
	Borrelia spp.	0	0	3 (1.1%)
	B. bigemina	0	1 (0.9%)	7 (2.5%)
	T. velifera	1 (0.8%)	25 (21.6%)	14 (5.1%)
Co-infections (2)	*Total*	53 (40.2%)	16 (13.8%)	88 (32%)
	R. africae [1] / *T. velifera*	47 (35.6%)	6 (5.2%)	0
	R. africae [1] / *Borrelia* spp.	3 (2.3%)	0	0
	R. africae [1] / *E. ruminantium*	3 (2.3%)	0	0
	Anaplasma spp. [2] / *Borrelia* spp.	0	1 (0.9%)	0
	Anaplasma spp. [2] / *T. velifera*	0	2 (1.7%)	3 (1.1%)
	Anaplasma spp. [2] / *Ehrlichia* spp. [3]	0	0	6 (2.2%)
	A. marginale / *T. velifera*	0	2 (1.7%)	10 (3.6%)
	A. marginale / *Borrelia* spp.	0	0	3 (1.1%)
	A. marginale / *Ehrlichia* spp. [3]	0	0	39 (14.29%)
	A. marginale / *B. bigemina*	0	0	4 (1.5%)

	Ehrlichia spp. [3] / *R. africae* [1]	0	2 (1.7%)	0
	Ehrlichia spp. [3] / *T. velifera*	0	1 (0.9%)	8 (2.9%)
	Ehrlichia spp. [3] / *Borrelia* spp.	0	0	3 (1.1%)
	Ehrlichia spp. [3] / *Leishmania* spp.	0	0	1 (0.4%)
	T. velifera / *B. bigemina*	0	0	10 (3.6%)
	T. velifera / *T. mutans*	0	2 (1.7%)	1 (0.4%)
Co-infections (3)	Total	8 (6.1%)	2 (1.7%)	26 (9.5%)
	R.africae [1] / *E. ruminantium* / *T. velifera*	3 (2.3%)	0	0
	R. africae [1] / *Borrelia* spp. / *T. velifera*	3 (2.3%)	0	0
	R. africae [1] / *T. velifera* / *T. mutans*	2 (1.5%)	1 (0.6%)	0
	R. africae [1] / *A. marginale* / *Ehrlichia* spp. [3]	0	1 (0.6%)	0
	A. marginale / *Ehrlichia* spp. [3] / *B. bigemina*	0	0	8 (2.9%)
	A. marginale / *Ehrlichia* spp. [3] / *B.bovis*	0	0	1 (0.4%)
	A. marginale / *Ehrlichia* spp. [3] / *T. velifera*	0	0	15 (5.5%)
	Anaplasma spp. [2] / *Ehrlichia* spp. [3] / *B. bigemina*	0	0	1 (0.4%)
	Anaplasma spp. [2] / *Ehrlichia* spp. [3] / *T.velifera*	0	0	1 (0.4%)
Co-infections (4)	Total	1 (0.8%)	0	9 (3.3%)
	R. africae [1] / *T. velifera* / *Borrelia* spp. / *E. ruminantium*	1 (0.8%)	0	0
	A. marginale / *Borrelia* spp. / *T. velifera* / *T. mutans*	0	0	1 (0.4%)
	A. marginale / *Ehrlichia* spp. [3] / *T. velifera* / *B. bigemina*	0	0	5 (1.8%)
	A. marginale / *Ehrlichia* spp. [3] / *T. velifera* / *B.bovis*	0	0	1 (0.4%)
	A. marginale / *Leishmania* spp. / *T. velifera* / *T. mutans*	0	0	1 (0.4%)
	Anaplasma spp. [2] / *Ehrlichia* spp. [3] / *Borrelia* spp. / *T. velifera*	0	0	1 (0.4%)
Co-infections (5)	Total	0	0	1 (0.4%)
	A. marginale / *Ehrlichia* spp. [3] / *Borrelia* spp. / *T. velifera* / *T. mutans*	0	0	1 (0.4%)

[1] Assuming that all the *Rickettsia* spp. found in *Rhipicephalus microplus* samples from Guadeloupe were *R. africae*.
[2] Sample positive for *Anaplasma* spp. and not for *A. marginale*.
[3] Sample positive for *Ehrlichia* spp. and not for *E. ruminantium*.

Almost all of the *A. variegatum* samples from Guadeloupe were infected with at least one
pathogen (99.2%) (Table A3). Interestingly, only 56% of the *R. microplus* samples from Guadeloupe
were infected with at least one pathogen, whereas this rate reach 81% of the *R. microplus* from
Martinique (Table A3). Most of the positive samples corresponded to single infection or co-infection
with two pathogens in both tick species. Then, less than 10% of the tick samples displayed co-
infections with three pathogens (Table A3). Finally, only one *A. variegatum* sample from Guadeloupe

and *nine R. microplus* samples from Martinique were co-infected with four pathogens and one *R.*
microplus from Martinique was found infected with five pathogens (Table A3).
The majority of the *A. variegatum* samples displayed single infections with *R. africae* (52%) or co-
infections with *R. africae* and *T. velifera* (36%) (Table A3). Nevertheless, no negative or positive
association have been detected between the microorganisms detected in *A. variegatum* samples (Table
A4). At least, the presence of *R. africae* do not seem to interfere with the presence of *T. velifera* (Table
A4).

Table A4. Co-occurrence matrix of the microorganisms detected in *A. variegatum* samples collected
in Guadeloupe (n=132 samples). No negative or positive association have been detected when
performing a co-occurrence test using the co-occur function (α=0.05) and R version 3.6.0 (2019-04-
26).

	E. ruminantium	*Borrelia* spp.	*R. africae*	*T. mutans*	*T. velifera*
E. ruminantium	7	1	7	0	4
Borrelia spp.		7	7	0	4
R. africae			130	2	56
T. mutans				2	2
T. velifera					57

Among the 116 *R. microplus* samples from Guadeloupe analyzed here, most of the positive
samples presented single-infection (40.5%), with *R. africae* (12.9%) or *T. velifera* (21.6%) (Table A3). As
R. microplus is not considered as a vector of both of these microorganisms, we made the hypothesis
of a possible contamination of this tick species via infected bovine blood still present in engorged
tick, and/or via co-feeding with infected *A. variegatum* ticks. Interestingly, if we remove *R. africae, T.*
velifera and *T. mutans* from the screening analysis, the percentage of infected *R. microplus* from
Guadeloupe dropped drastically to 13.8% (16/116 samples infected with at least one pathogen). This
observation is particularly surprising when comparing this rate to the 81% infected *R. microplus* from
Martinique (see below). When testing for co-occurrence linkage, two positive associations have been
detected between the microorganisms detected in *R. microplus* samples from Guadeloupe, including
T. velifera / T. mutans, and *Anaplasma* spp. / *Borrelia* spp. (Table A5). Co-infections and positive
association between *T. velifera* and *T. mutans* have already been reported in the literature, such as in
cattle sera from Uganda, and Kenya [80,81]. Regarding the few samples positive for *Anaplasma* spp.
and *Borrelia* spp., the result of the co-occurrence test should be taken with cautions and deserve
further investigation. Nevertheless, no exclusion seemed to occur between the
pathogens/microorganisms detected in *R. microplus* from Guadeloupe (Table A5).

Table A5. Co-occurrence matrix of the microorganisms detected in *R. microplus* samples collected in
Guadeloupe (n=116 samples). *: Positive association detected when performing a co-occurrence test
using the co-occur function (α=0.05) and R version 3.6.0 (2019-04-26).

	A. marginale	*Borrelia* spp.	*R. africae*	*T. mutans*	*T. velifera*	*B. bigemina*	*Anaplasma* spp.[1]	*Ehrlichia* spp.[2]
A. marginale	5	0	1	0	2	0	0	1
Borrelia spp.		1	0	0	0	0	1*	0
R. africae			25	1	7	0	0	3
T. mutans				3	3*	0	0	0
T. velifera					39	0	2	1
B. bigemina						1	0	0
Anaplasma spp.[1]							4	0
Ehrlichia spp.[2]								7

718   [1] Sample positive for *Anaplasma* spp. and not for *A. marginale*.

² Sample positive for *Ehrlichia* spp. and not for *E. ruminantium*.

Among the 275 *R. microplus* samples from Guadeloupe, most of the sample presented single-infection (36%) and co-infections with two pathogens (32%) (Table A1). Five positive associations have been detected between the microorganisms detected in *R. microplus* samples from Martinique (Table A6). *T. mutans* have been found in positive association with *T. velifera*, as previously observed in the same tick species from Guadeloupe, and with *Leishmania* spp. and *Borrelia* spp. In addition, *T. velifera* were found in positive association with *B. bigemina* and *Ehrlichia spp.* with *A. marginale.* Finally, no negative association have been reported between the pathogens/microorganisms detected in *R. microplus* from Martinique (Table A6).

Table A6. Co-occurrence matrix of the microorganisms detected in *R. microplus* samples collected in Martinique (n=275 samples). (*) Positive associations detected when performing a co-occurrence test using the co-occur function (α=0.05) and R version 3.6.0 (2019-04-26).

	A. marginale	*Borrelia* spp.	*Leishmania* spp.	*T. mutans*	*T. velifera*	*B. bigemina*	*B. bovis*	*Anaplasma* spp.[1]	*Ehrlichia* spp.[2]
A. marginale	111	5	1	3	34	17	2	0	**70***
Borrelia spp.		12	0	**2***	3	0	0	1	5
Leishmania spp.			2	**1***	1	0	0	0	1
T. mutans				4	**4***	0	0	0	1
T. velifera					72	**15***	1	5	32
B. bigemina						35	0	1	14
B. bovis							2	0	2
Anaplasma spp.1								22	9
Ehrlichia spp.2									134

¹ Sample positive for *Anaplasma* spp. and not for *A. marginale*.

² Sample positive for *Ehrlichia* spp. and not for *E. ruminantium*.

References

1. Dantas-Torres, F.; Chomel, B.B.; Otranto, D. Ticks and tick-borne diseases: a One Health perspective. *Trends Parasitol.* **2012**, *28*, 437–446. Doi: 10.1016/j.pt.2012.07.003

2. George, J.E.; Davey, R.B.; Pound, J.M. Introduced ticks and tick-borne diseases: the threat and approaches to eradication. *Vet. Clin. North Am. Food Anim. Pract.* **2002**, *18*, 401–416, vi.

3. Gondard, M.; Cabezas-Cruz, A.; Charles, R.A.; Vayssier-Taussat, M.; Albina, E.; Moutailler, S. Ticks and Tick-Borne Pathogens of the Caribbean: Current Understanding and Future Directions for More Comprehensive Surveillance. *Front Cell Infect Microbiol* **2017**, *7*, 490. Doi:10.3389/fcimb.2017.00490.

4. Barré, N.; Uilenberg, G. Spread of parasites transported with their hosts: case study of two species of cattle tick. *Rev. - Off. Int. Epizoot.* **2010**, *29*, 149–160, 135–147.

5. Morel, P.-C. Etudes sur les tiques du bétail en Guadeloupe et Martinique. II. Agents pathogènes transmis par les tiques. *Revue d'élevage et de médecine vétérinaire des pays tropicaux* **1967**, *20*, 291–299. Doi:10.19182/remvt.7508.

6. Uilenberg, G.; Camus, E.; Barré, N. [Existence of Theileria mutans and Theileria velifera (Sporozoa, Theileriidae) in Guadeloupe (French West Indies)]. *Rev Elev Med Vet Pays Trop* **1983**, *36*, 261–264.

7. Camus, E.; Barre, N. Vector situation of tick-borne diseases in the Caribbean islands. *Veterinary Parasitology* **1995**, *57*, 167–176. Doi: 10.1016/0304-4017(94)03118-G.

8. Alonso, M.; Arellano-Sota, C.; Cereser, V.H.; Cordoves, C.O.; Guglielmone, A.A.; Kessler, R.; Mangold,

751   A.J.; Nari, A.; Patarroyo, J.H.; Solari, M.A. Epidemiology of bovine anaplasmosis and babesiosis in Latin
752   America and the Caribbean. *Rev. - Off. Int. Epizoot.* **1992**, *11*, 713–733.

753 9. Parola, P.; Vestris, G.; Martinez, D.; Brochier, B.; Roux, V.; Raoult, D. Tick-borne rickettiosis in
754   Guadeloupe, the French West Indies: isolation of *Rickettsia africae* from *Amblyomma variegatum* ticks and
755   serosurvey in humans, cattle, and goats. *Am. J. Trop. Med. Hyg.* **1999**, *60*, 888–893.

756 10. Camus, E.; Montenegro-James, S. Bovine anaplasmosis and babesiosis in the Lesser Antilles: risk
757   assessment of an unstable epidemiologic situation. *Vet. Res.* **1994**, *25*, 313–317.

758 11. Barre, N.; Garris, G.I.; Borel, G.; Camus, E. Hosts and Population Dynamics of *Amblyomma variegatum*
759   (Acari: Ixodidae) on Guadeloupe, French West Indies. *J Med Entomol* **1988**, *25*, 111–115. Doi:
760   10.1093/jmedent/25.2.111.

761 12. Vachiéry, N.; Jeffery, H.; Pegram, R.; Aprelon, R.; Pinarello, V.; Kandassamy, R.L.Y.; Raliniaina, M.; Molia,
762   S.; Savage, H.; Alexander, R.; et al. *Amblyomma variegatum* ticks and heartwater on three Caribbean
763   Islands. *Ann. N. Y. Acad. Sci.* **2008**, *1149*, 191–195. Doi: 10.1196/annals.1428.081.

764 13. Parola, P.; Attali, J.; Raoult, D. First detection of *Rickettsia africae* on Martinique, in the French West Indies.
765   *Ann Trop Med Parasitol* **2003**, *97*, 535–537. Doi: 10.1179/0003498032225001382.

766 14. Kelly, P.; Lucas, H.; Beati, L.; Yowell, C.; Mahan, S.; Dame, J. *Rickettsia africae* in *Amblyomma variegatum*
767   and domestic ruminants on eight Caribbean islands. *J. Parasitol.* **2010**, *96*, 1086–1088. Doi: 10.1645/GE-
768   2552.1.

769 15. Raoult, D.; Fournier, P.E.; Fenollar, F.; Jensenius, M.; Prioe, T.; de Pina, J.J.; Caruso, G.; Jones, N.; Laferl,
770   H.; Rosenblatt, J.E.; et al. *Rickettsia africae*, a Tick-Borne Pathogen in Travelers to Sub-Saharan Africa.
771   *New England Journal of Medicine* **2001**, *344*, 1504–1510. Doi: 10.1056/NEJM200105173442003.

772 16. Estrada-Peña, A.; Gray, J.S.; Kahl, O.; Lane, R.S.; Nijhof, A.M. Research on the ecology of ticks and tick-
773   borne pathogens—methodological principles and caveats. *Front Cell Infect Microbiol* **2013**, *3*. Doi:
774   10.3389/fcimb.2013.00029.

775 17. Cabezas-Cruz, A.; Vayssier-Taussat, M.; Greub, G. Tick-borne pathogen detection: what's new? *Microbes
776   Infect.* **2018**. Doi: 10.1016/j.micinf.2017.12.015.

777 18. Michelet, L.; Delannoy, S.; Devillers, E.; Umhang, G.; Aspan, A.; Juremalm, M.; Chirico, J.; van der Wal,
778   F.J.; Sprong, H.; Boye Pihl, T.P.; et al. High-throughput screening of tick-borne pathogens in Europe.
779   *Front Cell Infect Microbiol* **2014**, *4*. Doi: 10.3389/fcimb.2014.00103.

780 19. Dupraz, M.; Toty, C.; Devillers, E.; Blanchon, T.; Elguero, E.; Vittecoq, M.; Moutailler, S.; McCoy, K.D.
781   Population structure of the soft tick *Ornithodoros maritimus* and its associated infectious agents within a
782   colony of its seabird host *Larus michahellis*. *Int J Parasitol Parasites Wildl* **2017**, *6*, 122–130. Doi:
783   10.1016/j.ijppaw.2017.05.001.

784 20. Gioia, G.V.; Vinueza, R.L.; Marsot, M.; Devillers, E.; Cruz, M.; Petit, E.; Boulouis, H.J.; Moutailler, S.;
785   Monroy, F.; Coello, M.A.; et al. Bovine anaplasmosis and tick-borne pathogens in cattle of the Galapagos
786   Islands. *Transbound Emerg Dis* **2018**, *65*, 1262–1271. Doi: 10.1111/tbed.12866.

787 21. Gondard, M.; Michelet, L.; Nisavanh, A.; Devillers, E.; Delannoy, S.; Fach, P.; Aspan, A.; Ullman, K.;
788   Chirico, J.; Hoffmann, B.; et al. Prevalence of tick-borne viruses in *Ixodes ricinus* assessed by high-
789   throughput real-time PCR. *Pathog Dis* **2018**, *76*. Doi: 10.1093/femspd/fty083.

790 22. Malmsten, J.; Dalin, A.-M.; Moutailler, S.; Devillers, E.; Gondard, M.; Felton, A. Vector-Borne Zoonotic
791   Pathogens in Eurasian Moose (Alces alces alces). *Vector Borne Zoonotic Dis.* **2018**. Doi:
792   10.1089/vbz.2018.2277.

793 23. Ahantarig, A.; Trinachartvanit, W.; Baimai, V.; Grubhoffer, L. Hard ticks and their bacterial

794   endosymbionts (or would be pathogens). *Folia Microbiol. (Praha)* **2013**, *58*, 419–428. Doi: 10.1007/s12223-
795   013-0222-1.

24. Duron, O.; Binetruy, F.; Noël, V.; Cremaschi, J.; McCoy, K.D.; Arnathau, C.; Plantard, O.; Goolsby, J.; Pérez
796   de León, A.A.; Heylen, D.J.A.; et al. Evolutionary changes in symbiont community structure in ticks.
797
798   *Mol. Ecol.* **2017**, *26*, 2905–2921. Doi: 10.1111/mec.14094.

25. Bonnet, S.; Jouglin, M.; Malandrin, L.; Becker, C.; Agoulon, A.; L'hostis, M.; Chauvin, A. Transstadial and
799
800   transovarial persistence *of Babesia divergens* DNA in *Ixodes ricinus* ticks fed on infected blood in a new
801   skin-feeding technique. *Parasitology* **2007**, *134*, 197–207. Doi: 10.1017/S0031182006001545.

26. Howell, J.M.; Ueti, M.W.; Palmer, G.H.; Scoles, G.A.; Knowles, D.P. Transovarial Transmission Efficiency
802
803   of *Babesia bovis* Tick Stages Acquired by *Rhipicephalus* (*Boophilus*) *microplus* during Acute Infection. *J Clin*
804   *Microbiol* **2007**, *45*, 426–431. Doi: 10.1128/JCM.01757-06.

27. Chauvin, A.; Moreau, E.; Bonnet, S.; Plantard, O.; Malandrin, L. *Babesia* and its hosts: adaptation to long-
805
806   lasting interactions as a way to achieve efficient transmission. *Vet. Res.* **2009**, *40*, 37. Doi:
807   10.1051/vetres/2009020.

28. Robinson, J.B.; Eremeeva, M.E.; Olson, P.E.; Thornton, S.A.; Medina, M.J.; Sumner, J.W.; Daschi, G.A. New
808
809   approaches to detection and identification of *Rickettsia africae* and *Ehrlichia ruminantium* in *Amblyomma*
810   *variegatum* (Acari: Ixodidae) ticks from the Caribbean. *J. Med. Entomol.* **2009**, *46*, 942–951.

29. Socolovschi, C.; Huynh, T.P.; Davoust, B.; Gomez, J.; Raoult, D.; Parola, P. Transovarial and trans-stadial
811
812   transmission of *Rickettsiae africae* in *Amblyomma variegatum* ticks. *Clin. Microbiol. Infect.* **2009**, *15 Suppl 2*,
813   317–318. Doi: 10.1111/j.1469-0691.2008.02278.x.

30. Maina, A.N.; Jiang, J.; Omulo, S.A.; Cutler, S.J.; Ade, F.; Ogola, E.; Feikin, D.R.; Njenga, M.K.; Cleaveland,
814
815   S.; Mpoke, S.; et al. High Prevalence of *Rickettsia africae* Variants in *Amblyomma variegatum* Ticks from
816   Domestic Mammals in Rural Western Kenya: Implications for Human Health. *Vector Borne Zoonotic Dis*
817   **2014**, *14*, 693–702. Doi: 10.1089/vbz.2014.1578.

31. Yssouf, A.; Socolovschi, C.; Kernif, T.; Temmam, S.; Lagadec, E.; Tortosa, P.; Parola, P. First molecular
818
819   detection of *Rickettsia africae* in ticks from the Union of the Comoros. *Parasites & Vectors* **2014**, *7*, 444. Doi:
820   10.1186/1756-3305-7-444.

32. Matsimbe, A.M.; Magaia, V.; Sanches, G.S.; Neves, L.; Noormahomed, E.; Antunes, S.; Domingos, A.
821
822   Molecular detection of pathogens in ticks infesting cattle in Nampula province, Mozambique. *Exp Appl*
823   *Acarol* **2017**, *73*, 91–102. Doi: 10.1007/s10493-017-0155-5.

33. Rodriguez O.N.; Espaine, L.; Rivas, A.; Rodriguez, P.. Epidemiology of cattle diseases caused by
824
825   haemoparasites in Cuba. Available online: http://www.cabi.org/isc/abstract/19902200923 (accessed on
826   Mar 25, 2017).

34. Woolhouse, M.E.J.; Thumbi, S.M.; Jennings, A.; Chase-Topping, M.; Callaby, R.; Kiara, H.; Oosthuizen,
827
828   M.C.; Mbole-Kariuki, M.N.; Conradie, I.; Handel, I.G.; et al. Co-infections determine patterns of
829   mortality in a population exposed to parasite infection. *Science Advances* **2015**, *1*, e1400026. Doi:
830   10.1126/sciadv.1400026.

35. Parola, P.; Inokuma, H.; Camicas, J.L.; Brouqui, P.; Raoult, D. Detection and identification of spotted fever
831
832   group Rickettsiae and Ehrlichiae in African ticks. *Emerging Infect. Dis.* **2001**, *7*, 1014–1017. Doi:
833   10.3201/eid0706.010616.

36. Inokuma, H.; Beppu, T.; Okuda, M.; Shimada, Y.; Sakata, Y. Detection of ehrlichial DNA in *Haemaphysalis*
834
835   ticks recovered from dogs in Japan that is closely related to a novel *Ehrlichia* sp. found in cattle ticks from
836   Tibet, Thailand, and Africa. *J. Clin. Microbiol.* **2004**, *42*, 1353–1355.

837   37.   Loftis, A.D.; Reeves, W.K.; Szumlas, D.E.; Abbassy, M.M.; Helmy, I.M.; Moriarity, J.R.; Dasch, G.A.
838   Rickettsial agents in Egyptian ticks collected from domestic animals. *Exp. Appl. Acarol.* **2006**, *40*, 67–81.
839   Doi: 10.1007/s10493-006-9025-2.

840   38.   Kang, Y.-J.; Diao, X.-N.; Zhao, G.-Y.; Chen, M.-H.; Xiong, Y.; Shi, M.; Fu, W.-M.; Guo, Y.-J.; Pan, B.; Chen,
841   X.-P.; et al. Extensive diversity of Rickettsiales bacteria in two species of ticks from China and the
842   evolution of the Rickettsiales. *BMC Evolutionary Biology* **2014**, *14*, 167. Doi: 10.1186/s12862-014-0167-2.

843   39.   Lu, M.; Tian, J.-H.; Yu, B.; Guo, W.-P.; Holmes, E.C.; Zhang, Y.-Z. Extensive diversity of rickettsiales
844   bacteria in ticks from Wuhan, China. *Ticks Tick Borne Dis* **2017**, *8*, 574–580. Doi:
845   10.1016/j.ttbdis.2017.03.006.

846   40.   Guo, W.-P.; Tian, J.-H.; Lin, X.-D.; Ni, X.-B.; Chen, X.-P.; Liao, Y.; Yang, S.-Y.; Dumler, J.S.; Holmes, E.C.;
847   Zhang, Y.-Z. Extensive genetic diversity of Rickettsiales bacteria in multiple mosquito species. *Sci Rep*
848   **2016**, *6*, 38770. Doi: 10.1038/srep38770.

849   41.   Yparraguirre, L.A.; Machado-Ferreira, E.; Ullmann, A.J.; Piesman, J.; Zeidner, N.S.; Soares, C.A.G. A hard
850   tick relapsing fever group spirochete in a Brazilian *Rhipicephalus* (*Boophilus*) *microplus*. *Vector Borne*
851   *Zoonotic Dis.* **2007**, *7*, 717–721. Doi: 10.1089/vbz.2007.0144.

852   42.   Hagen, R.M.; Frickmann, H.; Ehlers, J.; Krüger, A.; Margos, G.; Hizo-Teufel, C.; Fingerle, V.;
853   Rakotozandrindrainy, R.; Kalckreuth, V. von; Im, J.; et al. Presence of *Borrelia* spp. DNA in ticks, but
854   absence of *Borrelia* spp. and of *Leptospira* spp. DNA in blood of fever patients in Madagascar. *Acta Trop.*
855   **2018**, *177*, 127–134. Doi: 10.1016/j.actatropica.2017.10.002.

856   43.   Kumsa, B.; Socolovschi, C.; Raoult, D.; Parola, P. New Borrelia species detected in ixodid ticks in Oromia,
857   Ethiopia. *Ticks Tick Borne Dis* **2015**, *6*, 401–407. Doi: 10.1016/j.ttbdis.2015.03.009.

858   44.   Ehounoud, C.B.; Yao, K.P.; Dahmani, M.; Achi, Y.L.; Amanzougaghene, N.; Kacou N'Douba, A.;
859   N'Guessan, J.D.; Raoult, D.; Fenollar, F.; Mediannikov, O. Multiple Pathogens Including Potential New
860   Species in Tick Vectors in Côte d'Ivoire. *PLoS Negl Trop Dis* **2016**, *10*. Doi: 10.1371/journal.pntd.0004367.

861   45.   Estrada-Peña, A.; Álvarez-Jarreta, J.; Cabezas-Cruz, A. Reservoir and vector evolutionary pressures
862   shaped the adaptation of *Borrelia*. *Infection, Genetics and Evolution* **2018**. Doi: 10.1016/j.meegid.2018.03.023.

863   46.   Rodríguez, I.; Fernández, C.; Cinco, M.; Pedroso, R.; Fuentes, O. Do Antiborrelial Antibodies Suggest
864   Lyme Disease in Cuba? *Emerg Infect Dis* **2004**, *10*, 1698–1700. Doi: 10.3201/eid1009.031048.

865   47.   Rodríguez, I.; Fernández, C.; Sánchez, L.; Martínez, B.; Siegrist, H.H.; Lienhard, R. Prevalence of antibodies
866   to *Borrelia burgdorferi* sensu stricto in humans from a Cuban village. *Braz J Infect Dis* **2012**, *16*, 82–85. Doi:
867   http://dx.doi.org/10.1590/S1413-86702012000100015.

868   48.   Dessau, R.B. Infection due to *Borrelia burgdorferi* most likely does not occur in Cuba. *Braz J Infect Dis* **2012**,
869   *16*, 404; author reply 405-406. Doi: 10.1016/j.bjid.2012.06.007.

870   49.   Flanigan, T.P.; Schwan, T.G.; Armstrong, C.; Voris, V.; P, L.; Salata, R.A. Relapsing Fever in the US Virgin
871   Islands: A Previously Unrecognized Focus of Infection. *J Infect Dis* **1991**, *163*, 1391–1392. Doi:
872   10.1093/infdis/163.6.1391.

873   50.   Sharma, A.; Jaimungal, S.; Basdeo-Maharaj, K.; Rao, A.V.C.; Teelucksingh, S. Erythema Migrans–like
874   Illness among Caribbean Islanders. *Emerg Infect Dis* **2010**, *16*, 1615–1617. Doi: 10.3201/eid1610.100587.

875   51.   Pothirat, T.; Tantiworawit, A.; Chaiwarith, R.; Jariyapan, N.; Wannasan, A.; Siriyasatien, P.;
876   Supparatpinyo, K.; Bates, M.D.; Kwakye-Nuako, G.; Bates, P.A. First Isolation of *Leishmania* from
877   Northern Thailand: Case Report, Identification as *Leishmania martiniquensis* and Phylogenetic Position
878   within the *Leishmania enriettii* Complex. *PLoS Negl Trop Dis* **2014**, *8*. Doi: 10.1371/journal.pntd.0003339.

879   52.   Espinosa, O.A.; Serrano, M.G.; Camargo, E.P.; Teixeira, M.M.G.; Shaw, J.J. An appraisal of the taxonomy

and nomenclature of trypanosomatids presently classified as *Leishmania* and Endotrypanum. *Parasitology* **2018**, *145*, 430–442. Doi: 10.1017/S0031182016002092.

53. Leelayoova, S.; Siripattanapipong, S.; Manomat, J.; Piyaraj, P.; Tan-ariya, P.; Bualert, L.; Mungthin, M. Leishmaniasis in Thailand: A Review of Causative Agents and Situations. *Am J Trop Med Hyg* **2017**, *96*, 534–542. Doi: 10.4269/ajtmh.16-0604.

54. Steverding, D. The history of leishmaniasis. *Parasit Vectors* **2017**, *10*, 82. Doi : 10.1186/s13071-017-2028-5.

55. Desbois, N.; Pratlong, F.; Quist, D.; Dedet, J.-P. *Leishmania* (*Leishmania*) *martiniquensis* n. sp. (Kinetoplastida: Trypanosomatidae), description of the parasite responsible for cutaneous leishmaniasis in Martinique Island (French West Indies). *Parasite* **2014**, *21*, 12. Doi: 10.1051/parasite/2014011.

56. Liautaud, B.; Vignier, N.; Miossec, C.; Plumelle, Y.; Kone, M.; Delta, D.; Ravel, C.; Cabié, A.; Desbois, N. First case of visceral leishmaniasis caused by *Leishmania martiniquensis*. *Am. J. Trop. Med. Hyg.* **2015**, *92*, 317–319. Doi: 10.4269/ajtmh.14-0205.

57. Dantas-Torres, F. Ticks as vectors of *Leishmania* parasites. *Trends Parasitol.* **2011**, *27*, 155–159. Doi: 10.1016/j.pt.2010.12.006.

58. Rakhshanpour, A.; Malmasi, A.; Mohebali, M.; Nabian, S.; Mirhendi, H.; Zarei, Z.; Dalimi, A.; Mohammadiha, A.; Akhoundi, B.; Azarm, A. Transmission of *Leishmania infantum* by *Rhipicephalus sanguineus* (Acari: Ixodidae) in Dogs. *Iran J Parasitol* **2017**, *12*, 482–489.

59. Rojas-Jaimes, J.E.; Correa-Nuñez, G.H.; Rojas, N.; Cáceres-Rey, O. Detection of *Leishmania* (*V*) *guyanensis* in *Rhipicephalus* (*Boophilus*) *microplus* (Acari: Ixodidae) collected from *Pecari tajacu*. *Biomedica* **2017**, *37*, 208–214.

60. Alam, M.Z.; Rahman, M.M.; Akter, S.; Talukder, M.H.; Dey, A.R. An investigation about the possible role of cattle and goats as reservoir hosts for *Leishmania donovani* in Bangladesh. *J Vector Borne Dis* **2018**, *55*, 242–244. Doi: 10.4103/0972-9062.249484.

61. Paixão-Marques, M.D.S.; Alves-Martin, M.F.; Guiraldi, L.M.; Dos Santos, W.J.; de Lemos, F.A.; Sánchez, G.P.; Richini-Pereira, V.B.; Lucheis, S.B. First isolation of *Leishmania infantum* by blood culture in bovines from endemic area for canine visceral leishmaniasis. *Parasitology* **2019**, 1–3. Doi: 10.1017/S0031182019000088.

62. Lamattina, D.; Nava, S. Ticks infesting humans in Northern Misiones, Argentina. *Medicina (B Aires)* **2016**, *76*, 89–92.

63. Walker, A.R.; Bouattour, A.; Camicas, J.-L.; Estrada-Pena, A.; Horak, I.G.; Latif, A.A.; Pegram, R.G.; Preston, P.M. *Ticks of domestic animals in Africa: a guide to identification of species*; illustrée.; Bioscience Reports: Wisconsin University- Madison, **2003**; ISBN 0-9545173-0-X.

64. Nielsen, E.M.; Andersen, M.T. Detection and Characterization of Verocytotoxin-Producing *Escherichia coli* by Automated 5′ Nuclease PCR Assay. *J Clin Microbiol* **2003**, *41*, 2884–2893. Doi: 10.1128/JCM.41.7.2884-2893.2003.

65. Regnery, R.L.; Spruill, C.L.; Plikaytis, B.D. Genotypic identification of rickettsiae and estimation of intraspecies sequence divergence for portions of two rickettsial genes. *J. Bacteriol.* **1991**, *173*, 1576–1589.

66. Choi, Y.-J.; Lee, S.-H.; Park, K.-H.; Koh, Y.-S.; Lee, K.-H.; Baik, H.-S.; Choi, M.-S.; Kim, I.-S.; Jang, W.-J. Evaluation of PCR-Based Assay for Diagnosis of Spotted Fever Group Rickettsiosis in Human Serum Samples. *Clin Diagn Lab Immunol* **2005**, *12*, 759–763. Doi: 10.1128/CDLI.12.6.759-763.2005.

67. Hornok, S.; Földvári, G.; Elek, V.; Naranjo, V.; Farkas, R.; de la Fuente, J. Molecular identification of *Anaplasma marginale* and rickettsial endosymbionts in blood-sucking flies (Diptera: Tabanidae, Muscidae) and hard ticks (Acari: Ixodidae). *Vet. Parasitol.* **2008**, *154*, 354–359. Doi:

10.1016/j.vetpar.2008.03.019.

68. Loh, S.-M.; Gofton, A.W.; Lo, N.; Gillett, A.; Ryan, U.M.; Irwin, P.J.; Oskam, C.L. Novel *Borrelia* species

detected in echidna ticks, *Bothriocroton concolor*, in Australia. *Parasit Vectors* **2016**, *9*. Doi: 10.1186/s13071-

016-1627-x.

69. Albuquerque, A.; Campino, L.; Cardoso, L.; Cortes, S. Evaluation of four molecular methods to detect

Leishmania infection in dogs. *Parasit Vectors* **2017**, *10*, 57. Doi: 10.1186/s13071-017-2002-2.

70. Thompson, J.D.; Higgins, D.G.; Gibson, T.J. CLUSTAL W: improving the sensitivity of progressive

multiple sequence alignment through sequence weighting, position-specific gap penalties and weight

matrix choice. *Nucleic Acids Res.* **1994**, *22*, 4673–4680.

71. Tamura, K.; Nei, M. Estimation of the number of nucleotide substitutions in the control region of

mitochondrial DNA in humans and chimpanzees. *Mol. Biol. Evol.* **1993**, *10*, 512–526. Doi:

10.1093/oxfordjournals.molbev.a040023.

72. Kumar, S.; Stecher, G.; Tamura, K. MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for

Bigger Datasets. *Mol. Biol. Evol.* **2016**, *33*, 1870–1874. Doi: 10.1093/molbev/msw054.

73. Mixson, T.R.; Campbell, S.R.; Gill, J.S.; Ginsberg, H.S.; Reichard, M.V.; Schulze, T.L.; Dasch, G.A.

Prevalence of *Ehrlichia*, *Borrelia*, and Rickettsial Agents in *Amblyomma americanum* (Acari: Ixodidae)

Collected from Nine States. *Journal of Medical Entomology* **2006**, *43*, 1261–1268. Doi: 10.1603/0022-

2585(2006)43[1261:POEBAR]2.0.CO;2.

74. Trout Fryxell, R.T.; DeBruyn, J.M. The Microbiome of *Ehrlichia*-Infected and Uninfected Lone Star Ticks

(*Amblyomma americanum*). *PLoS One* **2016**, *11*. Doi: 10.1371/journal.pone.0146651.

75. Aktas, M.; Özübek, S. Transstadial Transmission of *Hepatozoon canis* by *Rhipicephalus sanguineus* (Acari:

Ixodidae) in Field Conditions. *J. Med. Entomol.* **2017**, *54*, 1044–1048. Doi: 10.1093/jme/tjx050.

76. Hamšíková, Z.; Silaghi, C.; Rudolf, I.; Venclíková, K.; Mahríková, L.; Slovák, M.; Mendel, J.; Blažejová, H.;

Berthová, L.; Kocianová, E.; et al. Molecular detection and phylogenetic analysis of *Hepatozoon* spp. in

questing *Ixodes ricinus* ticks and rodents from Slovakia and Czech Republic. *Parasitol Res* **2016**, *115*, 3897–

3904. Doi: 10.1007/s00436-016-5156-5.

77. Hailemariam, Z.; Krücken, J.; Baumann, M.; Ahmed, J.S.; Clausen, P.-H.; Nijhof, A.M. Molecular detection

of tick-borne pathogens in cattle from Southwestern Ethiopia. *PLoS One* **2017**, *12*. Doi:

10.1371/journal.pone.0188248.

78. Laird, J.S.; Kocan, A.A.; Kocan, K.M.; Presley, S.M.; Hair, J.A. Susceptibility of *Amblyomma americanum* to

natural and experimental infections with *Theileria cervi*. *J. Wildl. Dis.* **1988**, *24*, 679–683. Doi: 10.7589/0090-

3558-24.4.679.

79. Mans, B.J.; Pienaar, R.; Latif, A.A. A review of *Theileria* diagnostics and epidemiology. *International Journal*

for Parasitology: Parasites and Wildlife **2015**, *4*, 104–118. Doi: 10.1016/j.ijppaw.2014.12.006.

80. Njiiri, N.E.; Bronsvoort, B.M. deC; Collins, N.E.; Steyn, H.C.; Troskie, M.; Vorster, I.; Thumbi, S.M.; Sibeko,

K.P.; Jennings, A.; van Wyk, I.C.; et al. The epidemiology of tick-borne haemoparasites as determined

by the reverse line blot hybridization assay in an intensively studied cohort of calves in western Kenya.

Vet. Parasitol. **2015**, *210*, 69–76. Doi: 10.1016/j.vetpar.2015.02.020.

81. Oura, C. a. L.; Bishop, R.P.; Wampande, E.M.; Lubega, G.W.; Tait, A. Application of a reverse line blot

assay to the study of haemoparasites in cattle in Uganda. *Int. J. Parasitol.* **2004**, *34*, 603–613. Doi:

10.1016/j.ijpara.2003.12.012.

964