How many cubs can a mum raise? Maternal age and size influence litter size in polar

bears

Dorinda Marie Folio[1], Jon Aars[2], Olivier Gimenez[1], Andrew E. Derocher[3], Øystein Wiig[4],

Sarah Cubaynes[5]

[1]CEFE, CNRS, Univ Montpellier, Univ Paul Valéry Montpellier 3, EPHE, IRD, Montpellier,

France

[2]Norwegian Polar Institute, Fram Centre, Tromsø, Norway

[3]Department of Biological Sciences, University of Alberta, Edmonton, Canada

[4]Natural History Museum, University of Oslo, Oslo, Norway

[5]MMDN, INSERM U1198, EPHE/PSL & University of Montpellier, France

Abstract. Life history theory predicts that females' age and size affect the level of maternal

investment in current reproduction, balanced against future reproductive effort, maintenance

and survival. Using long-term (30 years) individual data on 231 female polar bears (*Ursus*

maritimus), we assessed age- and size-specific variation on litter size. Litter size varied with

maternal age, younger females had higher chances of losing a cub during their first months of

life. Results suggest an improvement of breeding abilities early in life due to experience with

subsequent reproductive senescence. Litter size increased with maternal size, indicating that

size may reflect individual quality. Heterogeneity was observed among the largest females,

suggesting that large size comes at a cost. Maternal traits and environmental conditions may

act together to influence reproductive success.

1. Introduction

Life history theory predicts that an optimal level of parental investment should maximize current reproductive success (RS) balanced against maintenance, survival, and future reproduction [1,2]. Among mammals, capital breeders are characterized by high maternal investment [10]. Lactation imposes high energetics demands on mothers [10] whereas energy is stored before breeding when foraging is constrained during reproduction [2]. Mothers' traits, namely age and body condition, should influence their ability to provide for their young, therefore influencing RS.

RS should increase with age due to an increase in maternal allocation to reproduction as residual reproductive value decreases ("terminal investment" hypothesis [1]). However, recent studies suggest a decline in RS with old age in wild vertebrates because of fewer resources to allocate to reproduction ("reproductive senescence" hypothesis [5]). An increase in RS has also been observed early in life due to increasing breeding abilities with experience ("constraint hypothesis" [6]). Moreover, irrespective of age, RS can vary with female body size and mass [3,4]. Larger size might benefit reproduction by improving or reflecting foraging abilities [7] and lactation [8]. RS might therefore increase until an optimal maternal size or age and then potentially decrease or level off because of maintenance costs exceeding the benefits associated with a higher size or experience of older females [3,9]. However, to date, the influence of maternal traits on reproductive outputs in mammalian capital breeders has received little attention [3].

We investigated the influence of maternal traits on RS in a capital breeder. Using long-term individual data on female polar bears (*Ursus maritimus*) of the Svalbard population, we assessed age- and size-specific variation on litter size that varies from one to

three young. High maternal investment is expected for polar bears, which live in an extreme

environment, rely only on stored fat reserves during pregnancy and for the first four months

of lactation and then care for and feed young during two and a half years [11]. We expected

litter size to increase until an optimal age and size, due to experience and individual quality,

and then decline for the largest and senescent individuals.

2. Materials and methods

(a) Data collection

We live-captured polar bears from 1992 to 2017 at Svalbard, Norway, from late March

to beginning of May – just after females have emerged from maternity dens with their cubs

[11] – using methods described in Stirling et al. 1989 [12]. The age of first reproduction for

Svalbard females is usually six years of age [11]. Age was estimated using a premolar tooth

extracted from sub-adult and adult bears (cubs were of known age based on size) [13]. Body

straight length (cm), hereafter size, was measured as the dorsal straight-line made from the tip

of the nose to the caudal end of the tail bone with bears laying sternally recumbent. Litter size

(one to three) was recorded upon capture as the number of cubs-of-the-year a mother had

reared to that time.

(b) Statistical analyses

We analysed litter size as a function of mother's traits using multinomial regression

[14] that extends standard logistic regression to more than two outcomes. Litters of one were

chosen as the reference category. Separate odd-ratios (OR) were determined for the relative

risk of a litter size of "two" *versus* "one"; and the relative risk of a litter size of "three" *versus*

"one" as a function of the covariates. Parameters α and γ respectively represent the OR of

twins and triplets, and will give an estimated probability of having one, two or three cubs as a

function of the covariates. We tested for linear and quadratic effects of maternal age and size.

Because fieldwork was spread over more than a month and that mortality rates for cubs within

their first year can be high [11], we considered capture date (as an ordinal date with 1^{st}

January being 1 in normal years and 0 in leap years) to account for a possible loss of young

between den emergence and observation. A yearly random effect was included on α and γ to

account for environmental variation. Twenty-seven models were fitted with a Bayesian

approach using Markov Chain Monte Carlo (MCMC) techniques in JAGS [15]. We used non-

informative normal prior distributions for the regression coefficients and a uniform prior

distribution for the standard deviation of the random effect. We ran two MCMC in parallel

with different initial values, 200,000 iterations each and an initial burn-in of 40,000 iterations.

One out of ten values were kept. We assessed convergence by visual inspection and by using

the Gelman and Rubin R-hat diagnostic (R-hat < 1.1 [16]). For model comparison, we used

the Deviance Information Criteria (DIC [17]) and considered the model with lowest DIC as

being best supported by the data. Model selection consisted of 5 steps. In step 1, we compared

different model structures for the intercept (same or different intercept on α and γ). In step 2

(respectively 3 and 4) we tested for the effect of maternal age (respectively maternal size and

capture date). Each tested variable could influence differently both OR (1: one common

coefficient, 2: two distinct ones, 3: one for α or 4: one for γ). In step 5, we compared models

with additive effects and interactions between the previously selected variables.

3. Results

The best model (model 5.2, Table 1) included an interaction between a quadratic effect

of age and capture date within the field season on the probability of having a litter of two over

one cub.

<u>Table 1:</u> List of all models considered with comparison based on Deviance Information

Criteria (DIC) values, with ΔDIC for the difference between the model with two intercepts

only and the model under investigation. Parameters α and γ represent the odds ratio (OR) of

twins and triplets, respectively, while # par. is for the number of model parameters. For each

step, the model best supported by the data is in bold.

Maternal size influenced both the probability of having two over one cub, and three

over one cub. Concerning age, passed mid-season (day 105), the probability of having twins

increased, then decreased, with age (Figure 1).

<u>Figure 1:</u> Estimated probability of having 1, 2 or 3 cubs as a function of capture date (in days

from March 21st) for (a) young (6 years old), (b) prime-aged (12 years old) and (c) old (18

years old) mothers and for a mean maternal size (194.8 cm). Predictions were obtained from

the best model (model 5.2 in Table 1). Solid lines are posterior means while dotted lines are

95% credible intervals.

Young females (5-9 years old (y.o.)) had a high probability of having twins shortly

after denning (day 90, $P(y=2) \approx 0.9$), but it declined within the field season, leading to a higher

probability of having just one cub alive towards the end of it (after day 125, $P(y=1) \approx 0.9$,

Figure 1a). Early in the season, prime-aged females (10-15 y.o.) also had a higher probability

of having two over one cub ($P(y=2)=0.7$ and $P(y=1)=0.3$, Figure 1b) while older females (>15

y.o.) had similar probabilities of having singletons or twins ($P(y=2) \approx P(y=1) \approx 0.5$, Figure 1c).

In contrast to the marked drop in spring litter size with time for young females, little variation

was observed for prime-aged ($0.7 < P(y=2) < 0.8$ and $0.2 < P(y=1) < 0.3$, Figure 1b) and old

females ($P(y=2) \approx P(y=1) \approx 0.5$, Figure 1c). Examining parameter estimates confirmed that the

interaction term between age and date was only significant for young mothers and not for

older ones (Table 2).

Table 2: Parameter estimates of the multinomial regression model best supported by the data

(see Table 1). Posterior mean, standard deviation (SD) and 95% credible intervals are

provided for the odds ratios (fixed effects) as well as the variance of the random effect year.

Rhat is the potential scale reduction factor (at convergence, Rhat < 1.2).

Litter size globally increased with the size of a mother (Figure 2). A higher probability

of having singletons was found for smaller females (size<183cm, $0.5 < P(y=1) < 0.8$,

$0.2 < P(y=2) < 0.5$, $P(y=3)=0$), of having twins for medium-sized females (190cm<size<200cm,

$P(y=1)=0.3$, $P(y=2)=0.7$, $P(y=3)=0$), and of having singletons or triplets for larger females

(size>210 cm, $P(y=1) \approx 0.5$, $0 < P(y=2) < 0.2$, $0.2 < P(y=3) < 0.5$). However, sample size was very

low for triplets (n=8).

Figure 2: Estimated probability of having 1, 2, or 3 cubs as a function of maternal size for a

mean value of mother's age (11.4 years) and capture date (day 105 ~ 16[th] of April).

Predictions were obtained from the best model (model 5.2 in Table 1). Solid lines are

posterior means while dotted lines are 95% credible intervals.

4. Discussion

Our results showed an influence of maternal traits on litter size, an index of

reproductive success, suggesting that a mothers' ability to invest in reproduction and care for

their offspring varied with their age and size. The quadratic pattern of variation in litter size

observed with age supported the hypothesis of a benefit of gaining experience early in life [6]

until 12 y.o. and of reproductive senescence [5] starting from 15 y.o. These results contradict

the terminal investment hypothesis [1], and support results on Canadian polar bears showing a

decrease in litter size and maternal body senescence after 16 years [18].

Under the experience hypothesis, an improvement of female's hunting skills might

explain the improvement of breeding abilities early in reproductive life [19]. Other studies on

mammals [3,18] suggest that an age-related increase in reproductive success could be linked

to mass gain and hence resources availability. Further supporting the experience hypothesis,

we found that younger females (< 9 y.o.) had about the same probability as prime-aged

females to produce two cubs, but higher chances to lose one during the capture spring season.

Under the senescence hypothesis, degradation of physiological functions with ageing

[20] might impair females' fat stores accumulation, causing a simultaneous decrease in

females' mass and reproductive outcomes [18]. Older mothers might therefore acquire less

energy, and might have higher energy allocation needs toward self-maintenance, reducing

energy allocated toward reproduction [1]. Reproductive senescence has been documented in

many wild populations for several reproductive parameters such as litter size [4,18], offspring

mass [3,5,18], and survival [4].

We showed that, on average, litter size increased with maternal size. The probability

of having triplets was only shown to depend of mother's size, although the sample size for

triplets was small. Large size might therefore be an index of individual quality, like in wolves

(*Canis lupus*) [4]. Other traits highlighting foraging capacities have been related to RS – e.g.

body mass and condition in bears and other species [3,18]. Among the largest females, the

chances of having a singleton or triplets were almost equal, while that of having twins was

low. This suggested heterogeneity among the largest females with one group of 'good quality

females' having triplets and another group of 'low quality females' having singletons. For the

latter group, larger size could be associated with a cost that may depend on other factors, such

as body condition and environmental quality. Large individuals occupying resource-poor

habitats or in a year of reduced resource availability might not have enough resources to

allocate to both their own maintenance and care for triplets. Heterogeneity in individual

quality may override reproductive cost [22], and costs may be restricted to resource-limited

contexts [23].

Overall, we found that litter size in polar bears increased with experience of mothers

early in life until a plateau, followed by a decrease for senescent females. Because population

growth mostly depends on female's RS, itself influenced by maternal traits, our findings

highlight the importance of accounting for individual heterogeneity to understand the species

response to environmental perturbations. Future research will aim at understanding the

determinants of female polar bears' reproductive tactics by accounting for environmental

conditions. Influence of climate variability has been shown to affect reproductive parameters

in several populations, including Svalbard [24].

Table 1.

Steps	Models	α twins OR	γ triplets OR	# par.	DIC	ΔDIC
1: Intercept	**1.1**	**Intercept**	**Intercept '**	**2**	**365.8**	**0**
	1.2	Intercept	Intercept	1	515.7	150.5
2: Age	**2.1**	**Age^2**	**β_0'**	**4**	**353.5**	**-12.3**
	2.2	Age^2	Age^2 '	6	356.5	-9.3
	2.3	Age^2	Age^2	4	359.4	-6.4
	2.4	β_0	Age'	3	364.7	-1.1
	2.5	Age	Age'	4	365.5	-0.3
	2.6	Age	β_0'	3	365.5	-0.3
	2.7	β_0	Age^2 '	4	366.4	0.6
	2.8	Age	Age	3	367	1.2
3: Size	**3.1**	**$Size^2$**	**Size'**	**5**	**360.3**	**-5.5**
	3.2	β_0	Size'	3	361.4	-4.4
	3.3	$Size^2$	$Size^2$ '	6	362	-3.8
	3.4	β_0	$Size^2$ '	4	362.1	-3.7
	3.5	$Size^2$	β_0'	4	362.9	-2.9
	3.6	Size	Size'	4	363.2	-2.6
	3.7	Size	β_0'	3	366.8	1
	3.8	$Size^2$	$Size^2$	4	367.5	1.7
	3.9	Size	Size	3	367.9	2.1
4: Date	**4.1**	**Date**	**β_0'**	**3**	**363.7**	**-2.1**
	4.2	Date	Date	3	364.6	-1.2
	4.3	Date	Date'	4	365.8	0
	4.4	β_0	Date'	3	367.4	1.6
5: Additive effects and interactions	5.1	$Age^2 + Date + Size^2$	Size'	8	346.9	0
	5.2	**Age^2 x Date + $Size^2$**	**Size'**	**10**	**342.8**	**-4.1**
	5.3	$Age^2 + Date * Size^2$	Size'	10	348.6	1.7
	5.4	Age^2 x $Size^2$ + Date	Size'	12	354.1	7.2

197 **Table 2.**
198

	Parameter	Mean	SD	95% credible interval	Rhat
			Fixed effects		
α	Intercept	1.399	0.275	[0.881; 1.962]	1.001
	Age (young)	-0.653	0.350	[-1.352; 0.023]	1.001
	Age (old)	-1.183	0.421	[-2.015; -0.356]	1.001
	Date	-0.047	0.207	[-0.451; 0.357]	1.001
	Size	-0.005	0.158	[-0.312; 0.312]	1.001
	$Size^2$	-0.279	0.112	[-0.505; -0.062]	1.001
	Age (young) x Date	-0.760	0.390	[-1.550; -0.017]	1.001
	Age (old) x Date	-0.149	0.404	[-0.946; 0.638]	1.001
γ	Intercept'	-2.634	0.486	[-3.671; -1.772]	1.001
	Size'	0.732	0.358	[0.056; 1.466]	1.001
			Year random effect		
	Variance	0.398	0.246	[0.024; 0.936]	1.001

References

1. Williams G. 1966 Natural selection, the costs of reproduction, and a refinement of Lack's

principle. *Am. Nat.* **100**, 687-690. (doi:10.1086/282461)

2. Stearns SC. 1992 *The evolution of life histories*. New York, NY: Oxford University Press.

3. Boltnev AI, York AE. 2001 Maternal investment in northern fur seals (*Callorhinus*

ursinus): interrelationships among mothers' age, size, parturition date, offspring size and sex

ratios. *J. Zool.* **254**, 219-228. (doi:10.1017/S0952836901000735)

4. Stahler DR, MacNulty DR, Wayne RK, VonHoldt B, Smith DW. 2013 The adaptive value

of morphological, behavioural and life-history traits in reproductive female wolves. *J. Anim.*

Ecol. **82,** 222-234. (doi:10.1111/j.1365-2656.2012.02039.x)

5. Nussey DH, Kruuk LEB, Donald A, Fowlie M, Clutton-Brock TH. 2006 The rate of

senescence in maternal performance increases with early-life fecundity in red deer. *Ecol.*

Lett. **9,** 1342-1350. (doi:10.1111/j.1461-0248.2006.00989.x)

6. Desprez M, Pradel R, Cam E, Monnat JY, Gimenez O. 2011 Now you see him, now you

don't: experience, not age, is related to reproduction in kittiwakes. *P. Roy. Soc. Lond. B Bio.*

278, 3060-3066. (doi:10.1098/rspb.2011.0189)

7. MacNulty DR, Smith DW, Mech LD, Eberly LE. 2009 Body size and predatory

performance in wolves: is bigger better? *J. Anim. Ecol.* **78**, 532-539. (doi:10.1111/j.1365-

2656.2008.01517.x)

8. Bowen WD, Iverson SJ, Boness DJ, Oftedal OT. 2001 Foraging effort, food intake and

lactation performance depend on maternal mass in a small phocid seal. *Funct. Ecol.* **15**, 325-

334. (doi:10.1046/j.1365-2435.2001.00530.x)

9. Houston AI, Stephens PA, Boyd IL, Harding KC, McNamara JM. 2006 Capital or income

breeding? A theoretical model of female reproductive strategies. *Behav. Ecol.* **18,** 241-250.

(doi:10.1093/beheco/arl080)

10. Langer P. 2008 The phases of maternal investment in eutherian mammals. *Zoology* **111,**

148-162. (doi:10.1016/j.zool.2007.06.007)

11. Wiig Ø. 1998 Survival and reproductive rates for polar bears at Svalbard. *Ursus* **10,** 25-

32. https://www.jstor.org/stable/3873105

12. Stirling I, Spencer C, Andriashek D. 1989 Immobilization of polar bears (*Ursus*

maritimus) with Telazol® in the Canadian Arctic. *J. Wildlife Dis.* **25,** 159-168.

(doi:10.7589/0090-3558-25.2.159)

13. Calvert W, Ramsay MA. 1998 Evaluation of age determination of polar bears by counts of

cementum growth layer groups. *Ursus.* **10,** 449-453. https://www.jstor.org/stable/3873156

14. Engel J. 1988 Polytomous logistic regression. *Stat. Neerl.* **42,** 233-252.

(doi:10.1111/j.1467-9574.1988.tb01238.x)

15. Plummer M. 2003 JAGS: A program for analysis of Bayesian graphical models using

Gibbs sampling. In *Proceedings of the 3rd international workshop on distributed statistical*

computing (Vol. 124, p. 125). Vienna.

16. Gelman A, Rubin DB. 1992 Inference from iterative simulation using multiple

sequences. *Stat. Sci.* **7,** 457-472. (doi:10.1214/ss/1177011136)

17. Spiegelhalter DJ, Best NG, Carlin BP, Van Der Linde A. 2002 Bayesian measures of

model complexity and fit. *J. Roy. Stat. Soc.* B. **64,** 583-639. (doi:10.1111/1467-9868.00353)

18. Derocher AE, Stirling I. 1994 Age-specific reproductive performance of female polar

bears (*Ursus maritimus*). *J. Zool.* **234,** 527-536. (doi:10.1111/j.1469-7998.1994.tb04863.x)

19. Derocher AE, Stirling I. 1992 The population dynamics of polar bears in western Hudson

Bay. *Wildlife 2001: Populations,* 1150-0059. (doi:10.1007/978-94-011-2868-1_88)

20. Mysterud A, Yoccoz NG, Stenseth NC, Langvatn R. 2001 Effects of age, sex and density

on body weight of Norwegian red deer: evidence of density–dependent senescence. *P. Roy.*

Soc. Lond. B Bio. **268,** 911-919. (doi:10.1098/rspb.2001.1585)

22. Hamel S, Cote SD, Gaillard JM, Festa-Bianchet M. 2009 Individual variation in

reproductive costs of reproduction: high-quality females always do better. *J. Anim. Ecol.* **78,**

143-151. (doi:10.1111/j.1365-2656.2008.01459.x)

23. Clutton-Brock TH, Stevenson IR, Marrow P, Maccoll AD, Houston AI, Mcnamara JM.

1996 Population Fluctuations, Reproductive Costs and Life-History Tactics in Female Soay

Sheep. *J. Anim. Ecol.* **65,** 675-689. (doi:10.2307/5667)

24. Derocher, AE 2005 Population ecology of polar bears at Svalbard, Norway. *Pop. Ecol.* **47**,

267-275. (doi: 10.1007/s10144-005-0231-2)

Acknowledgements. We thank Magnus Andersen for his participation in fieldwork, and Thor

Larsen for initiating the monitoring.

Authors' contributions. S.C., J.A. and O.G. conceived the analysis. J.A., AED and ØW

collected the data. D.F. carried out the analyses, and S.C., O.G., J.A. advised the analysis.

D.F. drafted the manuscript. All authors revised the manuscript critically, gave final approval

for publication and agreed to be accountable for all aspects of the work.

Data accessibility. Data are archived in the Dryad Digital Repository at

doi:10.5061/dryad.85qc15r.

Funding. WWF.

Competing interests. We have no competing interests.

Ethical statement. Animal handling methods were approved by the National Animal

Research Authority (http://www.mattilsynet.no/fdu).