Does data cleaning improve brain state classification?

Steven L. Meisler, Youssef Ezzyat, and Michael J. Kahana

Dept. of Psychology, University of Pennsylvania, Philadelphia PA 19104, USA

Corresponding Author:

Michael J. Kahana

Department of Psychology

University of Pennsylvania

425 S. University Avenue

Philadelphia, PA 19104

phone: 215.746.3501

fax: 215.746.3848

email: kahana@sas.upenn.edu

Abstract

Background: Neuroscientists routinely seek to identify and remove noisy or artifactual observations from their data. They do so guided by the belief that including such data would reduce power to detect effects of interest. Whereas standard methods can exclude certain well defined noise sources (e.g., line noise), most forms of noise do not easily separate from signals of interest. Here we ask how well methods routinely used to "clean" human electrophysiological recordings actually boost power to detect brain-behavior correlations.

New Method: This, to the authors' knowledge, is the first large-scale study of the impact of intracranial EEG preprocessing on brain state classification.

Results: We find that several commonly used data cleaning methods (automated methods based on statistical properties of the signal and manual methods based on expert review) reduce statistical power for both univariate and multivariate classification of successful memory encoding, a behavioral state with very well-characterized electrophysiological biomarkers.

Comparison with Existing Methods: By reallocating resources towards collecting more within-patient data instead of attempting to "clean" data, neuroscientists may see increases in the statistical power to detect physiological phenomena.

Conclusions: These findings highlight the challenge of partitioning signal and noise in the analysis of brain-behavior relations. They also prescribe increases in sample size and numbers of observations, rather than data cleaning, as the best approach to improving statistical power.

Keywords: intracranial EEG; physiology; noise removal; multivariate; episodic memory

Introduction

All measures of neural activity comprise mixtures of sources, some reflecting the physiological processes we seek to understand and others reflecting physiological or non-physiological sources that "contaminate" our signals of interest. In human neurophysiological recordings, nuisance signals may reflect electrical potentials produced outside of the brain (e.g., those produced by muscle or eye movements), electrical signals caused by sources within the brain other than those we seek to observe (e.g., epileptic activity contaminating signals related to normal cognition), or even sources outside of the body entirely, such as line noise or crosstalk between electrical channels within the recording system. To reduce the influence of these "nuisance" electrical signals, and thereby increase the statistical power to detect brain-behavior correlations, neurophysiologists employ a variety of "data cleaning" methods. Although the use of data cleaning methods is nearly ubiquitous in neuroscientific research, it remains unknown whether these methods actually increase the power to detect established effects.

In surveying the literature on intracranial EEG (iEEG) studies of human memory, a fairly restrictive domain, we identified six commonly used procedures designed to remove or attenuate putative noise sources prior to hypothesis testing (see Table 1). We compared methods based on statistical properties of the voltage timeseries (variance and kurtosis); annotating data based on suspected epileptic activity over channels or epochs; and monopolar vs. bipolar referencing. We applied each of these methods separately at both the electrode and epoch levels, and in each case we compared the effect of removing putatively noisy data with quantity-matched random data removal. Taking advantage of the large quantity of within-subject data, we further evaluated each method as a function of the quantity of data collected (# of electrodes and # of events) to determine whether the data quantity interacted with a particular method's effectiveness in increasing statistical power.

Because there is no current standard for preprocessing iEEG data, investigators are left to apply methods which they have used in the past or those used by others in the field without knowing which methods are likely to be effective. For methods based on statistical properties of the timeseries, which can be scripted and applied automatically, it is particularly easy for researchers to (consciously or not) evaluate multiple thresholds and methods and select the one that leads to the "cleanest" end result (Simmons, Nelson, & Simonsohn, 2011; Silberzahn et al., 2018). In the case of more labor intensive methods like identifying epileptic artifacts that require human annotation, subjectivity in criterion setting means that the rejected data will vary from one expert annotator to the next. Manual methods also have the obvious downside of requiring more time to carry out; to the extent that manual methods are effective at identifying and removing noise,

this may be worth the investment of time and effort. However if manual annotation does not, on average, have any benefit then it simply squanders the resources of an expert researcher.

We evaluated the aforementioned techniques based on their ability to resolve a well-established brain-behavior relation. Electrophysiological studies of memory encoding have consistently demonstrated that increases in broadband high-frequency activity, a correlate of neural activity (Rutishauser, Ross, Mamelak, & Schuman, 2010; Manning, Jacobs, Fried, & Kahana, 2009; Winawer et al., 2013), during encoding predicts subsequent recall, an effect seen across a broad network of brain regions including hippocampus, MTL, and lateral prefrontal cortex (Burke et al., 2014; Long, Burke, & Kahana, 2014; Long & Kahana, 2015; Sederberg et al., 2006). These broadly distributed patterns have also been incorporated into novel multivariate models for guiding closed-loop systems to detect memory encoding states (Ezzyat et al., 2018; Ezzyat & Rizzuto, 2018). We therefore applied our analyses to both a univariate and multivariate measure of memory encoding, in order to determine whether removing noise would differentially impact the two approaches.

Across univariate and multivariate measures we find that the majority of commonly used approaches to identifying and removing noisy epochs or channels do not increase statistical powder. For methods based on using a numeric threshold on statistical measures of the voltage timeseries (e.g. signal variance or kurtosis), liberal criteria ($z > 2.5$) do not increase statistical power compared to randomly removing an equivalent number of epochs or channels, while, as expected, using more conservative thresholds that exclude more data lead to reductions in statistical power. For methods based on using human manual annotation to identify epileptic channels or epochs, the overall effect is that removing data labeled "noisy" tends to decrease statistical power. We investigated whether the statistical or manual approaches might be more effective when applied to smaller datasets, however a within-subject parametric analysis of subsets of our dataset indicated this was not the case. The major exception to this pattern of null results occurred when comparing bipolar and monopolar referencing approaches, where bipolar referencing significantly outperformed monopolar referencing ins several analyses, particularly for the multivariate paradigm. The data show that the choice of referencing has a profound impact on iEEG analysis of the subsequent memory effect, while other commonly used noise removal approaches do not, and suggest that researchers will benefit from reallocating resources away from post-processing "noise" and toward collecting more within-subject data.

Table 1: Preprocessing approaches used in prior iEEG studies of human episodic memory.

	METHODS			
	Bipolar	Kurtosis	Standard Deviation	Epileptic Activity
Raghavachari et al., 2001				X
Caplan et al., 2001				X
Sederberg et al., 2003		X		
Ekstrom et al., 2005				X
Greco et al., 2007		X		
Sederberg et al., 2007		X		
van Vugt et al., 2009		X		
van Vugt et al., 2010		X		
Nolan et al., 2010			X	
Whitmer et al., 2010				
Dastjerdi et al., 2011				X
Burke et al., 2013	X			
Swann et al., 2013				X
Burke et al., 2014	X			
Lega et al., 2014		X		
Rangarajan et al., 2014			X	X
Zavala et al., 2014	X			
Greenberg et al., 2015	X	X		
Zhang and Jacobs, 2015		X		
Haque et al., 2015	X			
Kim et al., 2016				X
Vass et al., 2016				X
Piai et al., 2016	X			
Fonken et al., 2016			X	X
García-Cordero et al., 2017				X
Sheehan et al., 2018	X	X	X	X

Materials and Methods

Participants   We analyzed data from ninety-five neurosurgical patients with medication-resistant epilepsy who participated in at least three sessions of a verbal free recall memory task (described below) as part of an ongoing research collaboration coordinated by the University of Pennsylvania. As part of their clinical workup, subjects had electrodes implanted intracranially, either on the cortical surface or within the brain parenchyma–recordings from these electrodes during performance of the memory task were then used in the analyses described in this report. Subjects performed the memory task using a laptop at the bedside, and were tested at one of the following centers: Thomas Jefferson University Hospital (Philadelphia, PA), University of Texas Southwestern Medical Center (Dallas, TX), Emory University Hospital (Atlanta, GA), Dartmouth-Hitchcock Medical Center (Lebanon, NH), Hospital of the University of Pennsylvania (Philadelphia, PA), Mayo Clinic (Rochester, MN), and Columbia University Medical Center (New York, NY). The Institutional Review Board at each center approved the research protocol, and informed consent was obtained from each participant.

Free Recall Task   In a free recall task, a subject is presented with a list of words that he or she later tries to recall in any order (Glanzer, 1969). During the encoding phase, 12 nouns (http:// memory.psych.upenn.edu/WordPools) were presented one at a time on a computer screen in the subject's primary language (either English or Spanish). Each word remained on screen for 1600 milliseconds, followed by a blank inter-stimulus interval of between 750-1000 milliseconds (randomly drawn from a uniform distribution). Following the final word in the list, there was a 20 second distractor period, during which the subject used the laptop keyboard to answer arithmetic problems of the form A + B + C = ?, where A, B, and C were random integers between 0 and 9. After the delay ended, an 800 Hz auditory tone was played and a row of asterisks was displayed on the screen (simultaneously for 300 milliseconds), which signaled the beginning of the recall period (30 s). Subjects were instructed to recall as many words from the preceding list as could be remembered, in any order. Figure 1 illustrates the design of the task. Subjects completed up to 25 lists in a single session. Audio responses from the sessions were digitally recorded and manually parsed and annotated offline using Penn TotalRecall (http://memory.psych.upenn.edu/TotalRecall). There were two versions of the task: in one version the lists of words were constructed to minimize semantic relationships between the words; in the other version, four words from each of three semantic categories (e.g. DESK, CHAIR, TABLE, SOFA) were presented within the same list. Subjects could participate in one or both tasks across different sessions, however within a given session, only one task was used.

Figure 1: Schematic diagram of a single list presentation during free recall tasks. Reads from top to bottom.

Electrophysiological Recordings   iEEG data were gathered from both depth and subdural cortical surface electrodes. Subdural cortical surface leads were arranged in both strip and grid configurations. The types of electrodes used for recording varied across institutions based on clinician preference. Across the sites, the following electrode models were employed: PMT Depthalon (0.8 mm), AdTech Spencer RD (0.86 mm), AdTech Spencer SD (1.12 mm), AdTech Behnke-Fried (1.28 mm), and AdTech subdural strips and grids (2.3 mm).

iEEG data were recorded with one of the following clinical EEG interfaces, depending on institutional preference: Nihon Kohden EEG-1200, Natus XLTek EMU 128 or Grass Aura-LTM64. Electrode locations varied across subjects and were determined strictly by clinical monitoring needs. Depending on the amplifier and the preference of the clinical team, the signals were sampled at either 500, 1000, or 1600 Hz and were referenced to a common contact placed intracranially, on the scalp, or on the mastoid process. We notch filtered (Butterworth, 4th order, frequency range = center ± 2 Hz) all timeseries at 60, 120, and 180 Hz to reduce line noise.

General Procedure   The goal of this study was to use a large dataset to evaluate the effects of several methods for identifying and removing noise in iEEG recordings. We measured the effect of each method by determining the impact on the subsequent memory effect (SME), a well-characterized biomarker of successful memory encoding, during a free recall memory task. As described below, we separately assessed the effect on a univariate and multivariate measure of the SME, in order to answer the question of whether data removal had a differential impact on statistical power for uni- and multivariate measures. For both the univariate and multivariate analyses of the SME, we used spectral decomposition of the iEEG recordings to identify power in specific frequency bands; we describe the details in the next section. A schematic detailing our procedures for statistical thresholding and computing classifier metrics can be found in Figure S1.

Spectral Decomposition   We extracted all word encoding intervals from each subject's recordings (1600 ms following each word onset) and used wavelet convolution (wave number = 5) to spectrally decompose each epoch. We included a 1500 ms buffer period before and after each word encoding epoch before convolution, to mitigate edge artifacts, and subsequently discarded the buffer period before further analysis.

For the multivariate analysis, we used eight wavelets with center frequencies logarithmically-spaced between 3 and 175 Hz (3.0, 5.4, 9.6, 17.1, 30.6, 54.8, 97.9, 175.0). For the univariate analysis, we also used eight wavelets, but focused on the high frequency range (between 70 and 200 Hz; 70.0, 81.3, 94.5, 109.8, 127.5, 148.2, 172.1, 200.0). We then log-transformed and averaged the resulting powers over the 1600 ms

word encoding interval. We then normalized the powers (z-transform) across word encoding epochs, separately within each session, electrode and frequency (where relevant, the z-transformation was applied after applying one of the evaluated noise attenuation methods).

Multivariate Classification   The goal of the multivariate analysis was to attempt to use patterns of spectral power (across frequencies and electrodes) collected during encoding to classify individual word encoding epochs as recalled/not recalled. We used a logistic regression classifier implemented with Python's *scikit-learn* (Pedregosa et al., 2011) library, and using a balanced class weight to account for the difference in the base rate of recalled/not recalled trials. To reduce overfitting, we used L2 penalization (Hastie, Tibshirani, & Friedman, 2001) with penalty parameter $C = 2.4 \times 10^{-4}$, following protocols from previous free recall studies with a similar subject cohort (Ezzyat et al., 2017). We trained the classifier on feature matrices comprised of the z-scored powers (the input) and recall outcomes (binary -"recalled" or "not recalled"; the output) for all epochs with the exception of those corresponding to a single held-out list used to cross-validate the classifier. After being trained, we applied the model to the held-out data to generate estimated probabilities of recall for each held-out trial. We then iterated this procedure until all lists had been held-out once. We created receiver operating characteristic (ROC) curves for each subject using the cross-validated classifier probabilities and the true recalled/not recalled labels. An ROC curve plots the true positive rate against the false positive rate at various probability thresholds. We quantified model performance using area under the ROC curve (AUC), with an AUC of 0.5 representing chance (Hanley & McNeil, 1982).

Univariate Measure of SME   The goal of the univariate analyses was to compare spectral power in the high-frequency range (70–200 Hz) during the encoding period of recalled vs. not recalled words. Univariate high-frequency power during memory encoding across the brain is well-known to differentiate subsequently recalled and not recalled words (Burke et al., 2014). We averaged the z-scored powers over the word encoding period (0–1600 ms) separately for each epoch $\times$ frequency bin $\times$ electrode, and used Welch's t-test *across* epochs *within* each electrode to compare power for subsequently recalled vs. not recalled epochs (we employed Welch's test due to the possibility for different Ns and variances for the recalled/not recalled distributions). We then averaged the t-statistic across all electrodes to derive a summary measure of effect size for the univariate subsequent memory analysis.

Bipolar vs. Monopolar Referencing   Bipolar referencing involves referencing pairs of neighboring channels against each other, as opposed to monopolar referencing, which references against a single or average reference channel (Nunez & Srinivasan, 2006). Bipolar referencing is thought to minimize the impact of

reference channel noise on other channels (Burke et al., 2013), however a large-scale direct comparison of bipolar and monopolar reference schemes has not been reported. We therefore compared bipolar and monopolar referencing schemes to answer the question of whether there was any benefit, in terms of increasing statistical power by attenuating noise, to using bipolar referencing over monopolar referencing. Bipolar pairs were created for every group of two adjacent contacts on every depth, strip, and grid in a patient's montage. The timeseries for each bipolar pair is the difference between the signals in each electrode. We use the term "electrode" in this paper to denote either an individual channel or a bipolar pair, depending on the modality of the data of interest.

Bipolar referencing leads to a mismatch between the number of new bipolar referenced signals and the number of monopolar channels. To account for this, for any analysis comparing bipolar and monopolar referenced data, we calculated the average AUC (for multivariate analyses) and t-statistic (for univariate analyses) using 100 random samples from the full set of bipolar referenced recordings. We matched the size of each sample to the number of monopolar channels, thereby equating the number of features going into each of the bipolar and monopolar comparisons.

Electrode-Based Statistical Thresholding We evaluated the use of standard deviation and kurtosis thresholds for excluding channels from analysis. After collecting the timeseries data and separating them into epochs, we calculated the standard deviation and kurtosis of each epoch on each electrode. Within each channel and pair, we computed the average standard deviation and average kurtosis across epochs. Then, we z-scored these metrics separately across electrodes, and we evaluated the effect of excluding channels on the basis of standard deviation/kurtosis values at various thresholds in the distribution across channels for a given subject.

We excluded an electrode if its kurtosis or standard deviation z-score exceeded a given threshold. We evaluated the effect of thresholds at intervals of 0.5 between and including 0 and 3. As a benchmark for comparison, we also compared the threshold-based approach to randomly removing an equivalent number of channels at each threshold. We repeated the random removal 100 times, and the reported AUC/t-statistic for random removal represents the average across iterations. These results may be found in Figure 3. The next section describes the methods that eliminate epochs as opposed to electrodes.

Epoch-Based Statistical Thresholding Similar to the analysis described in the previous paragraph, we also tested the effect of excluding individual epochs based on their kurtosis and standard deviation. Within each epoch, we averaged the metric of interest (standard deviation, kurtosis) across electrodes. We pro-

ceeded to *z*-score these averages across epochs. To normalize the number of classifier features, we excluded an epoch only if its kurtosis or standard deviation exceed a given threshold for *both* monopolar and bipolar data. We randomly removed epochs similarly to the electrode randomization in the previous section. This process is also repeated 100 times, with the reported random AUC / *t*-statistic being the average of the 100 iterations. Epoch statistical thresholding results for kurtosis and variance may be found in Figure 6. While the previous sections described automated statistical thresholding, the next two sections detail removing data based on manual/semi-manual physician identification.

Neurologist-Based Electrode Exclusion   At each participating medical center, clinicians identified electrodes in regions associated with three types of abnormal neural activities: interictal spiking, seizure onsets, and lesions. For a subset of subjects, we also had information about contacts that may have been disconnected, saturated, or otherwise faulty. Electrode indications were stored in a subject-specific text file, although one did not exist for every subject in this experimental cohort. For those who had this information available ($N^* = 83$), this information was parsed to identify monopolar channels associated with the abnormal categories. Any bipolar pair that included at least one such channel was also grouped as belonging to the abnormal category associated with the channel(s). All possible combinations of categories were removed before performing spectral decomposition and classification. The resulting AUCs / *t*-statistics may be found in Figure 9.

Neurologist-Based Epoch Exclusion   Recently, a semi-automatic method has been evaluated to identify epochs with high frequency oscillatory (HFO) activity, a characteristic of epileptic EEG and iEEG (Waldman et al., 2018; Shimamoto et al., 2018; Weiss et al., 2018). The algorithm employed in the method annotates an epoch with one of the following labels that are then visually validated: 0) No HFO event detected, 1) Ripple superimposed in an interictal discharge, 2) Sharply contoured epileptiform spike, 3) Ripple on oscillation, 4) Fast ripple superimposed in an interictal discharge, 5) Fast ripple on oscillation. Following the guidance of the developers of the method, we removed as artifacts any epoch with an annotation of 1, 2, and/or 4. Then spectral decomposition and *t*-stat and AUC calculations proceeded similarly to other analyses. Only twenty-nine subjects had both at least 3 free recall sessions and HFO epochs to remove. To account for the smaller sample size, we lifted the restraint on minimum number of completed free-recall sessions, allowing for 70 subjects. Bipolar results for this analysis, including the resulting changes in *t*-stats and AUCs for classifiers built on both all electrodes and only those implicated in an HFO event, may be found in Figure 11 (monopolar results are in Figure S7). For the data subsampling method described in the next section, the

29 subjects with at least 3 sessions are used. These results may be found in Figures 12 and S8.

Data Subsampling   In addition to evaluating the effect of applying the preceding methods to a subject's full dataset, we were also interested in how the impact of the noise removal methods might differ for large and small within-subject datasets. To address this question, for all previously described preprocessing methods, we repeated the analyses using variously-sized subsets of data within subject. We subsampled deciles (10% intervals between 10% and 100%) of each subject's full set of electrodes or word lists were randomly sampled. This process was iterated 50 times per session and sampling percentage, and the reported result for the subject is the average across iterations and sessions. For methods that involved statistical thresholding, z-score thresholds below 2 were not tested since *post-hoc* analysis suggested they did not improve classifier performance. Since these analyses don't aim to contrast bipolar and monopolar performance, we did not match the number of features between complementary bipolar and monopolar datasets. Bipolar subsampling results may be found in the figures following each analytical paradigm's non-subsampled results. To preserve space, monopolar data subsampling results can be found in the Supplemental Information, but the trends closely resembled those for bipolar results.

Statistics   All results are presented as the mean $\pm$ standard error of the mean (SEM) unless noted otherwise. Any horizontal dashed lines in results figures represent baseline metrics present when no data are removed. The shaded region around them represents the SEM of the baseline.

Data Availability   Electrophysiological data may be found at http://memory.psych.upenn.edu/Electrophysiological_Data. The Python Time Series Analysis (PTSA) toolbox (https://github.com/pennmem/ptsa_new) was used to analyze the data. Custom Python scripts are available upon request from Steven Meisler (smeisler@mgh.harvard.edu).

Results

Figure 2 outlines the organization of the analyses conducted in the current study which span three levels: *Identification Method*, *Preprocessing Target*, and *Subsampled Data*. At the highest level (*Identification Method*), we evaluated noise preprocessing methods based on either (1) statistical thresholding of the voltage timeseries (standard deviation and kurtosis) or (2) manual review conducted by expert neurologists to identify data contaminated by epileptic activity. For each of these two broad approaches, at the next level (*Preprocessing Target*) we evaluated methods that targeted either (1) electrodes or (2) epochs. Finally, at

the lowest level (*Subsampled Data*) we evaluated the effect of each method on data of varying size by subsampling our within-subject datasets. We conducted our analyses with (1) no subsampling (full dataset); (2) subsampling electrodes; or (3) subsampling epochs. For the analyses that have different figures for bipolar and monopolar data, the monopolar results may be found in the Supplemental Information.

Electrode-Based Statistical Thresholding   As shown in Figure 3, removing electrodes based on their variance or kurtosis tends to diminish multivariate classifier performance. The AUC increases as the exclusion criteria becomes less strict. The average across subjects never reaches the AUC when no data are removed. At values marked by asterisks in the figure, the random removal (dotted lines) performed significantly better than the true exclusion (paired t-test; $P < .05$). At every threshold, the AUCs for bipolar pairs were significantly greater than the monopolar counterpart (paired t-test; $P < 1 \times 10^{-10}$). In the univariate measure, removing electrodes tended to outperform the random removal or baseline measure, though never significantly (paired t-test; $P > 0.1$). Changing the strictness of electrode exclusion did not produce as noticeable of a trend in univariate results than multivariate results.

Epoch-Based Statistical Thresholding   Similarly to the previous analysis, removing epochs had a negative impact on the AUCs, as depicted in Figure 6. Unlike the results from electrode removal, random removal never significantly out-performed true removal (paired t-test; $P < .05$). However, bipolar AUCs were again significantly higher than monopolar AUCs (paired t-test; $P < 1 \times 10^{-10}$). Similarly, in the univariate data, the t-statistics tended to increase as fewer epochs were excluded. The t-statistic never significantly exceeded the baseline.

Neurologist-Based Electrode Exclusion $N = 83$ subjects had neurologist-identified electrodes to exclude. Removing electrodes associated with abnormal neural activity did not improve classifier performance. As depicted in Figure 9, AUCs after removal never exceed the AUC when no electrodes are removed. Similar to the previous results, the bipolar AUCs significantly exceed those of monopolar data (paired t-test; $P < 1 \times 10^{-10}$). In the univariate measure, removing electrodes associated with interictal activity, lesions, or mechanical fault had slight improvements on performance, but none were significant (paired t-test; $P > .2$).

Neurologist-Based Epoch Exclusion   After lifting the restraint on minimum sessions completed, $N = 70$ subjects had HFO epochs to exclude. As depicted in Figure 11, the increase in the univariate metric was significant for classifiers trained *only* on electrodes that had HFO activity on them (1-sample t-test; $P < .05$). Besides that one case, removing these epochs did not have any significant effect.

Figure 2: Organization and location of results presented in the study. The supplemental figures depict monopolar results of the given paradigm.

Figure 3: Electrode-based statistical thresholding applied to the full dataset. The column and row labels denote the exclusion basis metric and classifier metric, respectively. * Indicates $P < 0.05$ compared to random. The dark red and blue dashed lines represent the respective values when no data are excluded. The shaded region surrounding those lines represent the SEM. SD: standard deviation, KUR: kurtosis. Lines were offset for visual clarity.

Figure 4: Electrode-based statistical thresholding applied to data subsampled at the electrode level. Each line corresponds to a different *z*-score threshold for statistical thresholding. SD: standard deviation, KUR: kurtosis. Lines were offset for visual clarity.

Figure 5: Electrode-based statistical thresholding applied to data subsampled at the list level. Each line corresponds to a different level of subsampling. SD: standard deviation, KUR: kurtosis. Lines were offset for visual clarity.

Figure 6: Epoch-based statistical thresholding applied to the full dataset. Dotted lines represent random removal. At no threshold were the random and non-random AUCs significantly different. SD: standard deviation, KUR: kurtosis. Lines were offset for visual clarity.

Figure 7: Epoch-based statistical thresholding applied to data subsampled at the electrode level. Each line corresponds to a different level of subsampling. SD: standard deviation, KUR: kurtosis. Lines were offset for visual clarity.

Figure 8: Epoch-based statistical thresholding applied to data subsampled at the list level. Each line corresponds to a different level of subsampling. SD: standard deviation, KUR: kurtosis. Lines were offset for visual clarity.

Figure 9: Neurologist-based electrode exclusion applied to the full dataset. S: Seizure onset, I: Interictal spiking, L: Lesion, B: Bad (loose/faulty contact).

Figure 10: Neurologist-based electrode exclusion applied to data subsampled at the electrode and list levels (N = 83). N-ID: neurologist-identified. Lines were offset for visual clarity.

Figure 11: Neurologist-based epoch exclusion applied to the full bipolar dataset and restricted to electrodes that had epochs removed. **A)**: Histograms of the changes in classifier metrics after removing HFO epochs for classifiers trained on all electrodes (top row) and only electrodes implicated in an HFO event (bottom row). Blue line denotes average of distribution. * denotes average greater than 0 (1-sample t-test; $P < .05$). **B)**: AUC and t-stats for all bipolar data (red) and those from HFO-implicated electrodes (orange). Error bars are standard error of the difference pairs of data (pre- and post- epoch removal). * denotes significant difference (paired t-test; $P < .05$).

Figure 12: Neurologist-based epoch exclusion subsampled at the electrode and list levels (N = 29). HFO: High-frequency oscillations. Lines were offset for visual clarity.

Discussion

Neuroscientists regularly identify and exclude data through both automatic and manual means, with the underlying assumption that doing so will increase the statistical power of detecting physiological phenomena. In this study, we systematically evaluated the impact of several preprocessing methods commonly employed in iEEG studies of human memory with the goal of determining if the approaches are effective at increasing statistical power at both a univariate and multivariate level. The methods we analyzed included bipolar rereferencing as well as removing both electrodes and epochs through automatic statistical thresholding and manual identification. In our cohort of 95 subjects, we also tested the efficacy of preprocessing while varying the amount of within-patient data, both in terms of the number of electrodes and session length.

We found that, with the exception of one approach, the preprocessing methods that involved data (e.g. epoch and electrode) exclusion, did not improve statistical power. In fact, at the most lenient thresholds for data exclusion, the statistical power was often significantly lower than when no preprocessing is performed. In some cases, primarily in the multivariate paradigm, the targeted data removal decreased statistical power more than the complementary random exclusion did. When varying the amount of within-patient data through subsampling, the data exclusion methods also failed to significantly increase statistical power. It is possible that the features that were targeted by the preprocessing, such as epileptic spikes, were so transient that their confounding impacts were nullified when the power spectrums were averaged across the entire encoding period.

Although our results strongly argue that commonly used data cleaning methods more often lead to reduced statistical power, these methods may nonetheless be valuable in situations where the artifact or noise is correlated with the behavior of interest. For example, one could imagine a scenario in which confounding EMG activity from muscle contractions (e.g. due to nervousness or concentration) could be correlated with memory performance. In such a situation, a researcher would certainly want to eliminate contamination from such artifacts in order to maximize the purity of the signal that actually comes from the brain. However, being able to do this this relies on the assumption that a researcher has a good model of the artifact and has strong evidence that the artifact is correlated with the behavior or neurophysiological signal of interest. If the artifact is not strongly related to the behavior, and if the ability to distinguish artifact from signal is poor, then removing artifacts is more likely to remove valid signal than artifactual signal, making it unlikely that conclusions drawn from the "cleaned" data will be representative of the true relationship between physiology and behavior. In the case of bipolar recordings from intracranially implanted electrodes,

one would not expect to see significant EMG and EOG artifacts. In situations in which the characteristics of the artifact are known (e.g. 50/60 Hz line noise), data cleaning is advantageous. However, our results suggest that for cases in which the distinction between signal and noise is less obvious, the researcher will be better off collecting and analyzing as much data as possible.

A second reason to reconsider data cleaning is because it offers researchers additional parameters with which to tune analyses. Because there is no widely accepted approach for intracranial studies of human memory that researchers can adopt *a priori*, there is a greater opportunity for a study to lead to distorted inferences about brain-behavior relationships that will not generalize. Opportunities to collect data from iEEG electrodes are usually scarce, (Parvizi & Kastner, 2018) making the recordings themselves highly valuable–ideally researchers would make use of as much of the collected data as possible. Our results suggest that investigators should be empowered to analyze as much of their data as possible, and that a fruitful use of resources would be to collect of large numbers of within-subject observations.

Generally, whether it be through targeted preprocessing, random exclusion, or subsampling, removing epochs tended to have a larger impact on statistical power than removing electrodes, and this difference was more pronounced in the univariate data. The reported *t*-stat was the average of tests across individual electrodes, so removing electrodes did not change the statistical power of the tests, which is determined by the number of epochs. Rather, it just limited the number of *t*-statistics to average across. Surprisingly, although removing "noisy" data failed to improve statistical power, bipolar rereferencing increased statistical power, even after controlling for the number of classification features. Although the improvement was modest, it was statistically significant in the multivariate analyses. This effect could be due to the utility of bipolar referencing in cancelling out artifacts on reference electrodes. This also suggests that local as compared with global spectral changes better characterize variability in mnemonic processing.

Another way to view our results would be that the classifiers, particularly the multivariate logistic regression, were robust to the noise that data exclusion targeted (Tomioka, Aihara, & Müller, 2007). This is consistent with previous work showing that L2-penalized logistic regression classifiers generalized to situations in which electrodes were excluded (Hammon & de Sa, 2007). Logistic regression also proved to be reliable in single-trial EEG classification (Tomioka et al., 2007) and cursor movement (Penny, Roberts, Curran, & Stokes, 2000) for a BCI. Similar robustness has also been observed for classifiers in other domains (Ryali, Supekar, Abrams, & Menon, 2010; Bootkrajang & Kabán, 2012; Carroll & Pederson, 1993; Komarek & Moore, 2003).

Conclusion

This study tested the impact of various preprocessing methods on multivariate and univariate classification of detecting subsequent memory effects. Across all tests that involved removing data based on automatic statistical thresholding or manual identification, exclusion rarely improved classifier performance. In fact, in many cases, removing data significantly diminished performance. Bipolar rereferencing, however, improved classifier metrics, particularly in the multivariate paradigm. These results were consistent across varying amounts of within-patient data. The study suggests that researchers should be deliberate in deciding their preprocessing pipeline, especially if it involves excluding data, and suggests that experimenter resources may be better deployed in the service of collecting larger within-subject datasets.

Conflicts of Interest

Dr. Michael Kahana has co-founded a company, Nia Therapeutics, LLC ("Nia"), intended to develop and commercialize brain stimulation therapies for memory restoration. He holds more than 5% equity interest in Nia. The views, opinions, and/or findings contained in this material are those of the authors and should not be interpreted as representing the official views or policies of the Department of Defense or the U.S. Government.

Acknowledgments

The authors would like to acknowledge Blackrock Microsystems' generosity in providing neural recording equipment. This work was supported by the DARPA Restoring Active Memory (RAM) program (Cooperative Agreement N66001-14-2-4032), as well as National Institutes of Health grant MH55687 and T32NS091006. This study would not be possible without all of the patients who volunteered their time to participate in the study.

References

Bootkrajang, J., & Kabán, A. (2012). Label-noise robust logistic regression and its applications. In *Joint european conference on machine learning and knowledge discovery in databases* (pp. 143–158).

Burke, J. F., Long, N. M., Zaghloul, K. A., Sharan, A. D., Sperling, M. R., & Kahana, M. J. (2014). Human intracranial high-frequency activity maps episodic memory formation in space and time. *NeuroImage*, *85*, 834–843.

Burke, J. F., Zaghloul, K. A., Jacobs, J., Williams, R. B., Sperling, M. R., Sharan, A. D., & Kahana, M. J. (2013). Synchronous and asynchronous theta and gamma activity during episodic memory formation. *Journal of Neuroscience*, *33*(1), 292–304.

Caplan, J. B., Madsen, J. R., Raghavachari, S., & Kahana, M. J. (2001). Distinct patterns of brain oscillations underlie two basic parameters of human maze learning. *Journal of Neurophysiology*, *86*, 368–380.

Carroll, R. J., & Pederson, S. (1993). On robustness in the logistic regression model. *Journal of the Royal Statistical Society. Series B (Methodological)*, 693–706.

Dastjerdi, M., Foster, B. L., Nasrullah, S., Rauschecker, A. M., Dougherty, R. F., Townsend, J. D., ... others (2011). Differential electrophysiological response during rest, self-referential, and non–self-referential tasks in human posteromedial cortex. *Proceedings of the National Academy of Sciences*, *108*(7), 3023–3028.

Ekstrom, A. D., Caplan, J., Ho, E., Shattuck, K., Fried, I., & Kahana, M. (2005). Human hippocampal theta activity during virtual navigation. *Hippocampus*, *15*, 881–889.

Ezzyat, Y., Kragel, J. E., Burke, J. F., Levy, D. F., Lyalenko, A., Wanda, P., ... Kahana, M. J. (2017). Direct brain stimulation modulates encoding states and memory performance in humans. *Current Biology*, *27*(9), 1251–1258.

Ezzyat, Y., & Rizzuto, D. S. (2018). Direct brain stimulation during episodic memory. *Current Opinion in Biomedical Engineering*, *8*, 78-83.

Ezzyat, Y., Wanda, P., Levy, D. F., Kadel, A., Aka, A., Pedisich, I., ... Kahana, M. J. (2018). Closed-loop stimulation of temporal cortex rescues functional networks and improves memory. *Nature Communications*, *9*(1), 365. doi: 10.1038/s41467-017-02753-0

Fonken, Y. M., Rieger, J. W., Tzvi, E., Crone, N. E., Chang, E., Parvizi, J., ... Krämer, U. M. (2016). Frontal and motor cortex contributions to response inhibition: evidence from electrocorticography. *Journal of neurophysiology*, *115*(4), 2224–2236.

Foster, B. L., Kaveh, A., Dastjerdi, M., Miller, K. J., & Parvizi, J. (2013). Human retrosplenial cortex

displays transient theta phase locking with medial temporal cortex prior to activation during autobiographical memory retrieval. *The Journal of Neuroscience*, *33*(25), 10439–10446.

García-Cordero, I., Esteves, S., Mikulan, E. P., Hesse, E., Baglivo, F. H., Silva, W., ... others (2017). Attention, in and out: scalp-level and intracranial eeg correlates of interoception and exteroception. *Frontiers in neuroscience*, *11*, 411.

Glanzer, M. (1969). Distance between related words in free recall: Trace of the STS. *Journal of Verbal Learning and Verbal Behavior*, *8*, 105-111.

Greco, A., Mammone, N., Morabito, F., & Versaci, M. (2007). Semi-automatic artifact rejection procedure based on kurtosis, Renyi's entropy and independent component scalp maps. *International Journal of Medical, Health, Biomedical and Pharmaceutical Engeneering*, *1*, 466–470.

Greenberg, J. A., Burke, J. F., Haque, R., Sharan, A. D., Litt, B., Baltuch, G. H., ... Zaghloul, K. A. (2015). Decreases in theta and increases in high frequency activity underlie associative memory encoding. *NeuroImage*, *114*, 257–263.

Hammon, P. S., & de Sa, V. R. (2007). Preprocessing and meta-classification for brain-computer interfaces. *IEEE Transactions on Biomedical Engineering*, *54*(3), 518–525.

Hanley, J. A., & McNeil, B. J. (1982). The meaning and use of the area under a receiver operating characteristic (roc) curve. *Radiology*, *143*(1), 29–36.

Haque, R. U., Wittig, J. H., Damera, S. R., Inati, S. K., & Zaghloul, K. A. (2015). Cortical low-frequency power and progressive phase synchrony precede successful memory encoding. *The Journal of Neuroscience*, *35*(40), 13577–13586.

Hastie, T., Tibshirani, R., & Friedman, J. (2001). *The elements of statistical learning*. New York: Springer-Verlag.

Kim, K., Ekstrom, A. D., & Tandon, N. (2016). A network approach for modulating memory processes via direct and indirect brain stimulation: Toward a causal approach for the neural basis of memory. *Neurobiology of Learning and Memory*.

Komarek, P., & Moore, A. W. (2003). Fast robust logistic regression for large sparse datasets with binary outputs. In *Aistats*.

Lega, B., Burke, J. F., Jacobs, J., & Kahana, M. J. (2015). Slow theta-to-gamma phase amplitude coupling in human hippocampus supports the formation of new episodic memories. *Cerebral Cortex*, *26*(1), 268-278.

Lega, B., Kahana, M. J., Jaggi, J. L., Baltuch, G. H., & Zaghloul, K. A. (2011). Neuronal and oscillatory activity during reward processing in the human ventral striatum. *NeuroReport*, *22*(16), 795-800.

Long, N. M., Burke, J. F., & Kahana, M. J. (2014). Subsequent memory effect in intracranial and scalp EEG. *NeuroImage*, *84*, 488–494. doi: 10.1016/j.neuroimage.2013.08.052

Long, N. M., & Kahana, M. J. (2015). Successful memory formation is driven by contextual encoding in the core memory network. *NeuroImage*, *119*, 332-337.

Manning, J. R., Jacobs, J., Fried, I., & Kahana, M. J. (2009). Broadband shifts in local field potential power spectra are correlated with single-neuron spiking in humans. *Journal of Neuroscience*, *29*(43), 13613–13620.

Nolan, H., Whelan, R., & Reilly, R. (2010). Faster: fully automated statistical thresholding for eeg artifact rejection. *Journal of neuroscience methods*, *192*(1), 152–162.

Nunez, P. L., & Srinivasan, R. (2006). *Electric fields of the brain*. New York: Oxford University Press.

Parvizi, J., & Kastner, S. (2018). Promises and limitations of human intracranial electroencephalography. *Nature neuroscience*, 1.

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., . . . Duchesnay, E. (2011). Scikit-learn: Machine learning in Python. *Journal of Machine Learning Research*, *12*, 2825–2830.

Penny, W. D., Roberts, S. J., Curran, E. A., & Stokes, M. J. (2000). Eeg-based communication: a pattern recognition approach. *IEEE transactions on Rehabilitation Engineering*, *8*(2), 214–215.

Piai, V., Anderson, K. L., Lin, J. J., Dewar, C., Parvizi, J., Dronkers, N. F., & Knight, R. T. (2016). Direct brain recordings reveal hippocampal rhythm underpinnings of language processing. *Proceedings of the National Academy of Sciences*, *113*(40), 11366–11371.

Raghavachari, S., Kahana, M. J., Rizzuto, D. S., Caplan, J. B., Kirschen, M. P., Bourgeois, B., . . . Lisman, J. E. (2001). Gating of human theta oscillations by a working memory task. *Journal of Neuroscience*, *21*(9), 3175-3183.

Rangarajan, V., Hermes, D., Foster, B. L., Weiner, K. S., Jacques, C., Grill-Spector, K., & Parvizi, J. (2014). Electrical stimulation of the left and right human fusiform gyrus causes different effects in conscious face perception. *Journal of Neuroscience*, *34*(38), 12828–12836.

Rutishauser, U., Ross, I., Mamelak, A., & Schuman, E. (2010). Human memory strength is predicted by theta-frequency phase-locking of single neurons. *Nature*, *464*(7290), 903–907.

Ryali, S., Supekar, K., Abrams, D. A., & Menon, V. (2010). Sparse logistic regression for whole-brain classification of fmri data. *NeuroImage*, *51*(2), 752–764.

Sederberg, P. B., Gauthier, L. V., Terushkin, V., Miller, J. F., Barnathan, J. A., & Kahana, M. J. (2006). Oscillatory correlates of the primacy effect in episodic memory. *NeuroImage*, *32*(3), 1422–1431. doi: 10.1016/j.neuroimage.2006.04.223

Sederberg, P. B., Kahana, M. J., Howard, M. W., Donner, E. J., & Madsen, J. R. (2003). Theta and gamma oscillations during encoding predict subsequent recall. *Journal of Neuroscience*, *23*(34), 10809–10814.

Sederberg, P. B., Schulze-Bonhage, A., Madsen, J. R., Bromfield, E. B., Litt, B., Brandt, A., & Kahana, M. J. (2007). Gamma oscillations distinguish true from false memories. *Psychological Science*, *18*(11), 927–932.

Sheehan, T. C., Sreekumar, V., Inati, S. K., & Zaghloul, K. A. (2018). Signal complexity of human intracranial eeg tracks successful associative memory formation across individuals. *Journal of Neuroscience*, 2389–17.

Shimamoto, S., Waldman, Z. J., Orosz, I., Song, I., Bragin, A., Fried, I., ... others (2018). Utilization of independent component analysis for accurate pathological ripple detection in intracranial eeg recordings recorded extra-and intra-operatively. *Clinical Neurophysiology*, *129*(1), 296–307.

Silberzahn, R., Uhlmann, E. L., Martin, D. P., Anselmi, P., Aust, F., Awtrey, E., ... others (2018). Many analysts, one data set: Making transparent how variations in analytic choices affect results. *Advances in Methods and Practices in Psychological Science*, *1*(3), 337–356.

Simmons, J. P., Nelson, L. D., & Simonsohn, U. (2011). False-positive psychology: Undisclosed flexibility in data collection and analysis allows presenting anything as significant. *Psychological science*, *22*(11), 1359–1366.

Swann, N. C., Tandon, N., Pieters, T. A., & Aron, A. R. (2012). Intracranial electroencephalography reveals different temporal profiles for dorsal-and ventro-lateral prefrontal cortex in preparing to stop action. *Cerebral Cortex*, *23*(10), 2479–2488.

Tomioka, R., Aihara, K., & Müller, K.-R. (2007). Logistic regression for single trial eeg classification. In *Advances in neural information processing systems* (pp. 1377–1384).

van Vugt, M. K., Schulze-Bonhage, A., Litt, B., Brandt, A., & Kahana, M. J. (2010). Hippocampal gamma oscillations increase with memory load. *Journal of Neuroscience*, *30*(7), 2694–2699.

van Vugt, M. K., Schulze-Bonhage, A., Sekuler, R., Litt, B., Brandt, A., Baltuch, G., & Kahana, M. J. (2009). Intracranial electroencephalography reveals two distinct similarity effects during item recognition. *Brain Research*, *1299*, 33–44. doi: 10.1016/j.brainres.2009.07.016

Vass, L. K., Copara, M. S., Seyal, M., Shahlaie, K., Farias, S. T., Shen, P. Y., & Ekstrom, A. D. (2016). Oscillations go the distance: Low-frequency human hippocampal oscillations code spatial distance in the absence of sensory cues during teleportation. *Neuron*, *89*(6), 1180–1186.

Waldman, Z. J., Shimamoto, S., Song, I., Orosz, I., Bragin, A., Fried, I., ... Weiss, S. A. (2018). A method

for the topographical identification and quantification of high frequency oscillations in intracranial electroencephalography recordings. *Clinical Neurophysiology*, *129*(1), 308–318.

Weiss, S. A., Berry, B., Chervoneva, I., Waldman, Z., Guba, J., Bower, M., ... others (2018). Visually validated semi-automatic high-frequency oscillation detection aides the delineation of epileptogenic regions during intra-operative electrocorticography. *Clinical Neurophysiology*, *129*(10), 2089–2098.

Whitmer, D., Worrell, G., Stead, M., Lee, I. K., & Makeig, S. (2010). Utility of independent component analysis for interpretation of intracranial eeg. *Frontiers in human neuroscience*, *4*, 184.

Winawer, J., Kay, K. N., Foster, B. L., Rauschecker, A. M., Parvizi, J., & Wandell, B. A. (2013). Asynchronous broadband signals are the principal source of the bold response in human visual cortex. *Current Biology*, *23*(13), 1145–1153.

Zavala, B. A., Tan, H., Little, S., Ashkan, K., Hariz, M., Foltynie, T., ... Brown, P. (2014). Midline frontal cortex low-frequency activity drives subthalamic nucleus oscillations during conflict. *Journal of Neuroscience*, *34*(21), 7322–7333.

Zhang, H., & Jacobs, J. (2015). Traveling theta waves in the human hippocampus. *The Journal of Neuroscience*, *35*(36), 12477–12487.