1 Impoverished auditory cues fail to engage
2 brain networks controlling spatial selective
3 attention

Yuqi Deng[1], Inyong Choi[2], Barbara Shinn-Cunningham[1, 3], Robert Baumgartner[1, 4, *]

[1] Biomedical Engineering, Boston University, Boston, MA 02215, USA

[2] Communication Sciences & Disorders, University of Iowa, Iowa City, IA 52242, USA

[3] Neuroscience Institute, Carnegie Mellon University, Pittsburgh, PA 15213 USA

[4] Acoustics Research Institute, Austrian Academy of Sciences, Vienna, Austria

[*] Corresponding Author: robert.baumgartner@oeaw.ac.at

Author contributions: R.B., B.S.-C., Y.D., and I.C. designed research; Y.D. and R.B. performed research;

Y.D. and R.B. analyzed data; and Y.D., R.B., B.S.-C., and I.C. wrote the paper.

Abstract

Spatial selective attention enables listeners to process a signal of interest in natural settings. However,

most past studies on auditory spatial attention used impoverished spatial cues: presenting competing

sounds to different ears, using only interaural differences in time (ITDs) and/or intensity (IIDs), or using

non-individualized head-related transfer functions (HRTFs). Here we tested the hypothesis that

impoverished spatial cues impair spatial auditory attention by only weakly engaging relevant cortical

networks. Eighteen normal-hearing listeners reported the content of one of two competing syllable

streams simulated at roughly +30 ˚ and -30˚ azimuth. The competing streams consisted of syllables from

two different-sex talkers. Spatialization was based on natural spatial cues (individualized HRTFs),

individualized IIDs, or generic ITDs. We measured behavioral performance as well as

electroencephalographic markers of selective attention. Behaviorally, subjects recalled target streams

most accurately with natural cues. Neurally, spatial attention significantly modulated early evoked

sensory response magnitudes only for natural cues, not in conditions using only ITDs or IIDs. Consistent

with this, parietal oscillatory power in the alpha band (8-14 Hz; associated with filtering out distracting

events from unattended directions) showed significantly less attentional modulation with isolated

spatial cues than with natural cues. Our findings support the hypothesis that spatial selective attention

networks are only partially engaged by impoverished spatial auditory cues. These results not only

suggest that studies using unnatural spatial cues underestimate the neural effects of spatial auditory

attention, they also illustrate the importance of preserving natural spatial cues in assistive listening

devices to support robust attentional control.

Keywords: auditory spatial selective attention, head-related transfer functions, electroencephalography

Introduction

Spatial hearing is crucial to selectively attend to sounds of interest in everyday social settings. The

remarkable ability of normal-hearing listeners to focus on a sound source within a complex acoustic

scene is often referred to as "the cocktail party phenomenon," and has a rich history (Cherry, 1953).

Nevertheless, the mechanisms controlling spatial selective attention are still poorly understood.

Acoustically, in everyday situations, the two ears provide the listener with a listener-specific

combination of spatial cues that include interaural time and intensity differences (ITDs and IIDs,

respectively), as well as spectral cues caused by acoustical filtering of the pinnae (Blauert, 1997).

Together, these cues, captured by individualized head-related transfer functions (HRTFs), allow the brain

to create a clear, punctate internal representation of the location of sound sources in the environment

(Majdak et al., 2019; Middlebrooks, 2015).

When only isolated or impoverished spatial cues are present, auditory localization performance

degrades and the natural perception of external auditory objects may even collapse into the listener's

head (e.g., Baumgartner et al., 2017; Callan et al., 2013a; Hartmann and Wittenberg, 1996).

Nevertheless, isolated ITDs and IIDs still create a strong sense of lateralization within the head;

moreover, even such highly impoverished spatial cues can be used to achieve spatial release from

speech-on-speech masking, behaviorally (Culling et al., 2004; Ellinger et al., 2017; Glyde et al., 2013;

Kidd et al., 2010b; Loiselle et al., 2016). The relative importance of ITDs and IIDs in spatial release from

masking remains unclear, with past studies reporting conflicting results when directly comparing

different binaural conditions (Ellinger et al., 2017; Glyde et al., 2013; Higgins et al., 2017; Shinn-

Cunningham et al., 2005).

Previous electroencephalography (EEG) and magnetoencephalography (MEG) studies have

demonstrated that rich spatial cues in sound stimuli lead to different cortical activity compared to using

isolated cues during sound localization (Callan et al., 2013a; Leino et al., 2007; Palomäki et al., 2005) and

auditory motion processing (Getzmann and Lewald, 2010). However, the apparently minor behavioral

consequences of using unnatural, non-individualized spatial cues on spatial release from masking,

combined with the ease of implementing studies with simple, non-individualized spatial cues, led to

their wide usage in auditory neuroscience studies (Cusack et al., 2001; Dahmen et al., 2010; Dai et al.,

2018; Itoh et al., 2000; Kong et al., 2014; Sach et al., 2000). Indeed, in the auditory neuroscience

literature, many studies did not even present true binaural signals, but instead studied "spatial"

attention by using dichotic signals, with one sound presented monaurally to one ear, and a competing

sound presented monaurally to the other ear (e.g., Ahveninen et al., 2011; Alho et al., 1999; Wöstmann

et al., 2016). These studies implicitly assumed that because listeners were able to use impoverished

spatial cues to listen to one sound from a particular (relative) direction, the cognitive networks

responsible for controlling spatial attention must be engaged just as they are when listening to rich,

natural spatial cues. Nonetheless, it is unclear whether and how engagement of higher-order cognitive

processes such as deployment of selective attention is affected by the use of unnatural or impoverished

spatial cues.

Modulation of neural signatures, such as event-related potentials (ERPs) and induced oscillatory activity,

is often taken as evidence of effective attentional control (Herrmann and Knight, 2001; Siegel et al.,

2012). In particular, auditory spatial attention is known to modulate early sensory ERPs in the N1 time

range (processing latencies of 100 to 150 ms) (Choi et al., 2013; Röder et al., 1999), whereas modulation

of P1 ERPs (50 to 100 ms) has only recently been demonstrated in a free field experiment (Giuliano et

al., 2014). Induced alpha oscillation (8 to 14 Hz) has been hypothesized to function as an information

gating mechanism (Klimesch et al., 2007). During auditory spatial attention, parietal alpha power often

decreases in the contralateral hemisphere of attended stimuli and/or increases in the ipsilateral

hemisphere (Banerjee et al., 2011; Lim et al., 2015; Wöstmann et al., 2016). These neural modulations

constitute objective metrics of the efficacy of attentional control.

Here, we test listeners in a selective attention paradigm with simultaneous, spatially separated talkers.

We use the aforementioned EEG measures to compare both perceptual ability and the neural signatures

of attentional control for simulations with impoverished vs. natural spatial cues. Eighteen subjects

performed an auditory spatial attention task with two competing streams located at roughly +30 ˚ and -

30˚ azimuth (Figure 1). On every trial, listeners were cued by an auditory cue to attend to either the left

or right stream and report the content of the cued stream. The competing streams consisted of syllables

(/ba/, /da/ or /ga/) from two different-sex talkers. Sound stimuli (including the cuing sound) were

spatialized using three different levels of naturalness and richness: 1) generic ITDs only, 2) individualized

IIDs, or 3) individualized HRTFs containing all of the naturally occurring spatial cues a listener

experiences in the everyday world. We show that behavioral performance is better when listeners hear

natural, individualized spatial cues than when they hear impoverished cues. Importantly, only natural

spatial cues yield significant attentional modulation of P1 amplitudes. Moreover, induced alpha activity

is less robust and poorly lateralized with isolated spatial cues compared to rich, natural spatial cues.

Fig. 1. *Auditory spatial attention task with two competing streams was used to assess the consequence of impoverished auditory spatial cues on neural proxies of attention control. An auditory cue was presented first from the location of the upcoming target stream, processed by the same spatialization scheme as the upcoming mixture. Following the cue, the competing streams began, one from around +30° the other from around -30° azimuth. Listeners were asked to recall the syllable sequence presented from the cued side. The first syllables of both streams were temporally aligned; however, the latter two syllables in the competing streams were staggered, enabling us to isolate neural responses to each. Feedback was provided after every trial.*

Results

Natural spatial cues facilitate behavioral performance. We compared the percentage of correctly recalling each syllable of the target stream across the three spatialization conditions (Figure 2). Paired t-tests were performed on the logit transformed percentages. For the first syllable, where the target sound and the distractor sound overlapped in time, subjects were least accurate in discriminating the target sound in the ITD condition compared to the IID condition ($t_{(17)}$ = 5.15, *P* < 0.001) and HRTF condition ($t_{(17)}$ = 7.09, *P* < 0.001). However, no statistically significant difference was observed between IID and HRTF conditions for the first syllable ($t_{(17)}$ = 1.4, *P* = 0.18). For the second and the third syllable, where target and distractor streams occurred staggered in time, subjects performed significantly better in the HRTF condition than in both the ITD condition (2nd syllable: $t_{(17)}$ = 2.78, *P* = 0.013; 3rd syllable:

$t_{(17)}$ = 2.68, P = 0.016) and the IID condition (2nd syllable: $t_{(17)}$ = 3.15, P < 0.01; 3rd syllable: $t_{(17)}$ = 2.15,

P = 0.046). There was no significant difference between the ITD and IID conditions for the two staggered

syllables (2nd syllable: $t_{(17)}$ = 0.49, P = 0.63; 3rd syllable: $t_{(17)}$ = 1.74, P = 0.099).

Fig. 2. *Listeners' (N = 18) recall performance was evaluated for every syllable and different*

spatialization conditions. Sounds were spatialized either based on generic ITDs, individualized IIDs,

or the natural combination of ITDs, IIDs, and spectral cues in individualized HRTFs. Behavioral

advantages of having more consistent spatial information were significant but small. * P < .05;

** P < .001; *** P < .0001

Impoverished spatial cues affect attentional modulation of ERPs. Figure 3A shows the ERPs evoked by

the onset of the second syllable of the attended target sound and the unattended distractor sound,

aligning the onsets of the target and distractor syllables to 0 s to allow direct comparison. Stimulus

onsets elicited a fronto-central positivity (P1) between 50 to 100 ms followed by a negativity (N1)

between 100 to 180 ms (Figure 3A-B). The amplitudes of these two components were extracted and the

difference between attended stimuli (target sound) and unattended stimuli (distractor sound) was

calculated in order to quantify attentional modulation for both the P1 and N1 components (Figure 3C).

A.

B.

C.

Fig. 3. *P1 amplitudes were only modulated by the direction of attention in the HRTF condition,*

whereas N1 amplitudes were modulated equivalently across spatialization conditions (N = 18). **A.**

ERP waveforms at fronto-central electrodes were compared between the attended target stream

and the unattended distractor stream for every spatialization condition. The P1 time range was

defined between 50 ms and 100 ms, and the N1 time range was defined between 100 ms and

180 ms. **B.** *Most topographies of both ERP components show maxima at the fronto-central sites*

(black dots) used for evaluation. **C.** *The modulation strength of ERP components was assessed by*

the amplitude differences between attended and unattended streams.

P1 responses to attended stimuli were significantly larger than to unattended stimuli in the HRTF

condition ($t_{(17)}$ = 3.12, *P* < 0.01). However, no significant attentional modulation was found in the ITD

($t_{(17)}$ = 0.50, *P* = 0.62) and IID conditions ($t_{(17)}$ = 0.06, *P* = 0.95). Consistent with this, a direct comparison

of the amount of modulation (difference in ERP magnitudes for attended versus unattended sounds)

found that attentional modulation was significantly larger in the HRTF condition than in the ITD

($t_{(17)}$ = 2.48, *P* = 0.024) and IID conditions ($t_{(17)}$ = 2.29, *P* = 0.035); however, modulation did not differ

significantly between the ITD and IID conditions ($t_{(17)}$ = 0.27, *P* = 0.79) (Figure 3C).

In all three spatialization conditions, the N1 amplitude was modulated by spatial attention. An attended

sound elicited larger N1 amplitudes than when a sound was ignored in all three conditions (ITD:

$t_{(17)}$ = 3.01, *P* < 0.01; IID: $t_{(17)}$ = 4.12, *P* < 0.001; HRTF: $t_{(17)}$ = 3.56, *P* < 0.01). The magnitude of this N1

modulation did not differ significantly across the three spatialization conditions (ITD vs IID: $t_{(17)}$ = 0.32,

P = 0.76; ITD vs HRTF: $t_{(17)}$ = 0.21, *P* = 0.84; HRTF vs IID: $t_{(17)}$ = 0.13, *P* = 0.90) (Figure 3C).

Alpha oscillation power shows less attentional modulation with impoverished spatial cues. To

investigate the effect of spatialization on attentional control, we analyzed the power in alpha oscillations

during the attentional preparation period (-800 ms to 0 ms), a time period in which listeners know

where to orient spatial attention based on the preceding acoustic cue, but before the sound mixture of

competing streams begins. We averaged the power in alpha across all trials for each spatialization

condition, regardless of where spatial attention was focused, to get a measure of the total engagement

of alpha activity. We then compared relative power for different attentional directions. Averaged across

direction of attentional focus, we calculated the time courses of alpha global field power (GFP, Figure

4A) and compared within-subject differences of the temporal average within the preparatory time

period across spatialization conditions (Figure 4B). We found significantly larger alpha GFP differences in

the HRTF condition than in the other two conditions (HRTF vs ITD: $t_{(17)}$ = 2.75, P = 0.014; HRTF vs IID:

$t_{(17)}$ = 2.63, P = 0.017). No significant difference was found between the ITD and IID conditions

($t_{(17)}$ = 0.021, P = 0.98).

Fig. 4. *Within-subject differences in alpha-band GFP are larger in the HRTF condition, especially*

during the preparatory time window (after the sound cue but before the first syllables of the

*competing streams). **A.** Waveforms of the average (± SEM) GFP differences are shown during the*

*baseline period, preparatory phase, and stimulus phase with stream competition. **B.** The temporal*

average of the preparatory alpha GFP difference is larger for the HRTF condition.

We next explored whether the lateralization of alpha with the spatial focus of attention was greater

with more natural spatial cues than with impoverished cues. To this end, for each spatialization

condition we calculated the alpha power "attentional modulation index" (AMI), defined as the alpha

power difference between attend left and attend right trials divided by the overall alpha power

(Wöstmann et al., 2016). Calculated in this way, the AMI is expected to be positive in left and negative in

right parietal channels. Scalp topographies of the resulting AMI are shown in Figure 5. The alpha AMI

was then averaged within the left and the right parietal channels (see Methods). Statistical comparisons

between left- and right-hemisphere AMIs were made within conditions to quantify lateralization of

alpha power. Statistically significant hemispheric differences were found in the HRTF condition only

($t_{(17)}$ = 3.09, P < 0.01), not in the ITD ($t_{(17)}$ = 1.29, P = 0.21) or IID conditions ($t_{(17)}$ = 0.15, P = 0.88). To

compare alpha lateralization directly across conditions, the hemispheric difference in alpha AMI was

calculated by subtracting average right-hemisphere AMI from the average left-hemisphere AMI. Direct

comparison of conditions shows that alpha lateralization was significantly greater in the HRTF condition

than in the ITD condition ($t_{(17)}$ = 2.18, P = 0.043); alpha lateralization in the IID condition did not differ

significantly from that of either the HRTF condition ($t_{(17)}$ = 1.82, P = 0.086) or the ITD condition

($t_{(17)}$ = 0.72, P = 0.48).

Fig. 5. *Attentional modulation of alpha activity was lateralized to the hemisphere ipsilateral to the*

target stream only in the HRTF condition. AMI topographies and hemispheric averages are shown

*for every spatialization condition (N = 18). ** P < .001*

Impoverished spatial cues lead to worse behavioral performance, smaller P1 modulation, reduced

modulation of preparatory alpha GFP, and reduced lateralization of alpha power with attentional focus,

confirming our hypothesis that impoverished spatial cues impaired engagement of spatial attention. We

then explored relationships between these metrics to see if there were ordered relationships in the

individual measures of performance and neural signatures of attentional control. To investigate the

relationship between evoked response modulation and alpha oscillatory activities, we first calculated

the regression slope relating P1 amplitude to preparatory alpha GFP for each subject, and then

performed a paired t-test on the coefficients obtained. No consistent relationship between alpha GFP

and P1 amplitudes was observed ($t_{(17)}$ = 0.90, P = 0.38). Correlation analysis was also conducted

comparing behavioral accuracy to P1 modulation, defined as the attended P1 amplitude minus

unattended P1 amplitude. No consistent relationships between P1 modulation and behavioral

performance were observed for any syllable (1^{st} syllable: $t_{(17)}$ = 0.54, P = 0.59; 2^{nd} syllable: $t_{(17)}$ = 0.31, P

= 0.76; 3^{rd} syllable: $t_{(17)}$ = 0.69, P = 0.50). Similarly, we did not observe consistent relationships between

alpha AMI lateralization and response accuracy for any syllable (1^{st} syllable: $t_{(17)}$ = 0.19, P = 0.85; 2^{nd}

syllable: $t_{(17)}$ = 1.39, P = 0.18; 3^{rd} syllable: $t_{(17)}$ = 0.11 , P = 0.91). Thus, although there were significant

differences in engagement of attention across spatial conditions as measured both behaviorally and

neurally, the individual subject differences in these metrics were not closely related.

Discussion

Behaviorally, we found that impoverished spatial cues impair performance on an auditory spatial

attention task in a multi-talker scene. We used objective electrophysiological measures to assess

whether the naturalness and richness of spatial cues also impacts how strongly auditory spatial

attention modulates brain responses. We found that impoverished spatial cues reduce the strength of

the evoked and induced neural signatures of attentional control. Specifically, evoked P1 amplitudes and

induced alpha oscillatory power showed less attentional modulation for sound stimuli with

impoverished spatial cues compared to when spatial cues were tailored to recreate the natural, rich

experience of individual listeners.

Impoverished spatial cues result in less neural modulation during selective attention. We investigated

attentional modulation of four established neural signatures of selective attention: evoked P1 and N1

amplitudes and induced power and lateralization of alpha oscillation. While attentional modulation of

N1 amplitude was observed in all conditions, attentional modulation of the earlier P1 amplitude was not

observed or was significantly weaker in the impoverished cue conditions compared to the natural cue

condition. Similarly, we found less preparatory alpha power activity in the impoverished spatial cue

conditions than in the natural cue condition, reflected by two indexes quantifying the amount of spatial

variability of alpha power: alpha GFP (Figure 4) and AMI (Figure 5). In the ITD and IID conditions,

although there was a hint of preparatory alpha lateralization over parietal sensors, the amount of

lateralization was significantly smaller than in the HRTF condition and did not reach statistical

significance. Preparatory alpha activity during spatial attention tasks has been well documented to form

a specific lateralization pattern in both vision and audition (Banerjee et al., 2011; Kelly, 2006; Sauseng et

al., 2005; Worden et al., 2000), which is thought to be evidence of a preparatory information-gating

mechanism (Foxe and Snyder, 2011; Jensen and Mazaheri, 2010; Klimesch, 2012; Klimesch et al., 2007).

In vision, alpha lateralization has been observed to increase with the laterality of attention focus (Rihs et

al., 2007a; Samaha et al., 2016), reflecting an inhibition pattern topographically specific to attention

focus. Moreover, evidence for active top-down control of the phase of alpha oscillation during visual

spatial attention suggests that alpha oscillatory activity represents active engagement and

disengagement of the attentional network (Samaha et al., 2015). In addition, a previous somatosensory

study has revealed that the alpha lateralization is positively correlated to pre-stimulus cue reliability,

further suggesting that alpha lateralization reflects the top-down control in order to optimize the

processing of upcoming stimuli (Haegens et al., 2011). Although relatively few studies have investigated

alpha activity in audition, studies suggest that alpha control mechanisms are supra-modal rather than

sensory specific (33). In the current experiment, a pre-stimulus auditory cue directed listeners where to

focus attention in an upcoming sound mixture. The cue was spatialized using the same auditory features

used to spatialize the stream mixture. Our results thus suggest that compared to stimuli with natural

spatial cues, stimuli featuring only ITDs or only IIDs are less reliable in directing attentional focus,

producing weaker engagement of spatial attention and reduced attentional modulation of neural

responses.

Consistent with the idea that impoverished spatial cues lead to weaker engagement of spatial attention,

we found that the P1 ERP component was modulated by attention only with natural spatial cues, not

with impoverished cues; this result is consistent with a weak spatial representation failing to engage

attentional modulation of early sensory responses (Figure 3). Our finding that attentional focus leads to

a modulation of P1 amplitude for natural spatial cues is consistent with previous reports of effects of

attention on the P1 amplitude observed in previous spatial attention studies across sensory modalities

[auditory: (Giuliano et al., 2014); visual: (Hillyard and Anllo-Vento, 1998; Hopfinger et al., 2004)]. Past

studies agree that P1 modulation reflects an early sensory inhibition mechanism related to suppression

of task-irrelevant stimuli. Although debates remain as to whether P1 modulation results from bottom-up

sensory gain control (Hillyard and Anllo-Vento, 1998; Luck, 1995; Slagter et al., 2016) or for some top-

down inhibitory process (Freunberger et al., 2008; Klimesch, 2011a), it is generally accepted in visual

spatial studies that greater P1 amplitude modulation is associated with greater inhibition of to-be-

ignored stimuli (Couperus and Mangun, 2010; Hillyard and Anllo-Vento, 1998; Klimesch, 2012).

Interestingly, attentional modulation of auditory P1 has been found to be positively correlated to visual

working memory capacity, a result that was used to suggest that stronger P1 modulation reflects better

attentional control of the flow of sensory information into working memory (Fukuda and Vogel, 2009;

Giuliano et al., 2014). Our result is consistent with the hypothesis that P1 modulation directly reflects

attentional control. Specifically, impoverished spatial cues likely produce a "muddy" representation of

auditory space that supports only imprecise, poorly focused top-down spatial attention. The resulting

lack of control and specificity of spatial auditory attention results in early P1 responses that are

unmodulated by attentional focus.

N1 modulation is well documented as a neural index of attentional control (Choi et al., 2013; Hillyard

and Anllo-Vento, 1998; Stevens et al., 2008; Wyart et al., 2012). The attentional modulation of N1 is

thought to reflect attentional facilitation rather than inhibition (Couperus and Mangun, 2010;

Marzecová et al., 2018; Slagter et al., 2016). In contrast to preparatory alpha and P1, we found that the

later N1 evoked response was modulated similarly, regardless of the richness and naturalness of spatial

cues.

Due to the robustness and relatively large amount of modulation, changes in auditory N1 amplitude

have been used as a biomarker and a primary feature for classification of attentional focus ((Blankertz et

al., 2011; Schreuder et al., 2011); see also recent work on decoding attentional focus for running speech

using the correlation between neural responses and the power envelope of the speech streams: (Chait

et al., 2010; Mesgarani and Chang, 2012; Rimmele et al., 2015)). However, there is little known about

how N1 amplitudes reflects the processing of different spatial cues during auditory spatial attention.

Previous studies have revealed different N1 topographies during ITD and IID processing, leading to the

conclusion that ITD and IID are processed by different neural populations in the auditory cortex

(Johnson and Hautus, 2010; Tardif et al., 2006; Ungan et al., 2001). However, debates remain about

whether this difference in topography depends on perceived laterality, instead of different neural

populations specialized for processing different spatial cues. Results from a more recent study show that

auditory N1 modulation does not differ across spatial cue conditions, indicating integrated processing of

sound locations in auditory cortex regardless of cues (Salminen et al., 2015). In the current study, N1

modulation did not differ across the three spatialization conditions. Thus, our results support the idea

that the same cortical neural population is responsible for processing different binaural spatial cues.

Behavioral disadvantages associated with impoverished spatial cues are modest and depend on sound

stimulus characteristics. Despite the influence of spatial cue richness on neural metrics, our behavioral

results showed only small (albeit significant) behavioral differences between impoverished spatial cues

and natural, individualized spatial cues (Figure 2). In line with previous studies that observed greater

spatial release from masking with combined spatial cues compared to with isolated cues (Culling et al.,

2004; Ellinger et al., 2017), accuracy was best in the HRTF condition. The small accuracy improvement

over using impoverished cues is seen consistently across subjects. In the first syllable where the target

and distractor streams overlap in time, the HRTF condition yielded an increase of 13% points in accuracy

over the ITD condition, but is comparable to performance in the IID condition. In the two staggered

syllables, accuracy in the HRTF condition is greater than in the ITD and IID conditions by only about 6%

and 1% point, respectively. These differences in behavioral performance across syllables suggest that

the characteristics of sound stimuli influence the difficulty of the task and may influence the behavioral

advantages of having richer, more robust spatial cues (Kidd et al., 2010a). Concordantly, a previous

study with complex tone stimuli has shown much larger differences in behavioral performance of up to

20% points (Schröger, 1996), whereas studies presenting speech stimuli in a multi-talker environment

found no behavioral advantage of having combined cues compared to impoverished cues (Glyde et al.,

2013). These behavioral discrepancies, in combination with our neural findings, indicate that behavioral

performance alone is not a sensitive metric for determining whether cortical networks controlling

spatial selective attention are fully engaged.

Summary

Our results indicate that although impoverished spatial cues can support spatial segregation of speech in

a multi-talker environment, they do not fully engage the brain networks controlling spatial attention and

lead to weak attentional control. Previous auditory studies have provided evidence that impoverished

spatial cues do not evoke the same neural processing mechanisms as natural cue combinations during

localization tasks with single sounds (Callan et al., 2013b; Getzmann and Lewald, 2010; Leino et al.,

2007; Palomäki et al., 2005). The current study extends these findings, demonstrating that the efficacy

of higher-level cognitive processing, such as deployment of auditory selective attention, also depends on

the naturalness of spatial cues. Poor attentional control was reflected in limited modulation of neural

biomarkers of attentional processes. These findings suggest that the many past auditory attention

studies using impoverished spatial cues may have underestimated the robust changes in cortical activity

associated with deployment of spatial auditory attention in natural settings. Although impoverished

auditory spatial cues can allow listeners to deploy spatial attention effectively enough to perform well in

simple acoustic scenes, noisy, complex listening environments like those encountered in everyday

environments pose greater challenges to attentional processing. In natural settings, spatial attention

may fail unless attentional control networks are fully engaged. Thus, these results demonstrate the

importance of preserving rich, natural spatial cues in hearing aids and other assistive listening devices.

Materials and Methods

Subjects. Twenty-one paid volunteers and one author within the age of 18-42 years (M = 22.9, SD = 5.5;

12 females, 10 males) participated in this study. None of the subjects had audiometric thresholds

greater than 20 dB for frequencies from 250 Hz to 8 kHz. All participants gave informed consent as

approved by the Boston University Institutional Review Board. Two subjects were withdrawn from the

study due to the inability to perform the task (percentage of correct response less than 30% after

training), and two subjects were removed during EEG data preprocessing due to excessive artifacts.

Therefore 18 subjects remained for further analysis (N = 18).

Stimuli and Procedure. The sound stimuli consisted of consonant-vowel syllables (/ba/, /da/, & /ga/),

each 0.4 s in duration. These syllables were recorded from three talkers that naturally differed in

fundamental frequency (F0). The auditory cue was a single syllable /ba/ spoken by a low-pitch male

voice (F0 = 91 Hz, estimated by Praat software) (Boersma, 2001). The following target and distractor

streams both consisted of three syllables randomly chosen out of the set of three syllables (with

replacement). The target stream was spoken by either a female (F0 = 189 Hz) or a high-pitch male talker

(F0 = 125 Hz), and the distractor stream was spoken by the other talker different than the target stream.

The first syllable of the target and distractor sound overlapped in time, while the latter two syllables

were separated by 200 ms, onset to onset (Figure 1). To avoid engagement of temporal attention rather

than spatial attention, the assignment of the target stream being leading or lagging was equally

distributed across trials. In the leading stream, the onsets of all three syllables were separated by

400 ms; in the lagging stream, the onsets of the first and the second syllable were separated by 600 ms,

whereas those of the second and the third syllable were separated by 400 ms. All sound stimuli were

presented at a sound pressure level of approximately 75 dB.

Cue and stimuli were presented via earphones (ER-2, Etymotic Research, Inc.) and spatialized to

approximately ±30° azimuth (0° elevation). Three different spatialization conditions were used: HRTF,

IID, and ITD. In the HRTF condition, individualized HRTFs, providing natural combinations of ITDs, IIDs,

and spectral cues, were used. In the IID condition, ITDs were removed from the individualized HRTFs by

computing minimum-phase representations of the filters via removing the non-causal part of the

cepstrum (Oppenheim and Schafer, 1975). Hence, the IID and HRTF conditions provided the same

monaural magnitude spectra and thus the same energetic advantage of the ear ipsilateral to the target.

In the ITD condition, spatialization was based on simply delaying the signal presented to the

contralateral ear by 300 μs, thus providing no energetic advantage to the ipsilateral ear.

The measurement procedure for individualized HRTFs was identical to that described in (Baumgartner et

al., 2017). In short, acoustical transfer characteristics between loudspeakers positioned at the target

angles 1.5 m away from the subject's head and miniature microphones inside the subject's blocked ear

canals were measured inside a semi-anechoic chamber. Room reflections were removed from the

resulting impulse responses via temporal windowing (0.5-ms cosine ramps) to a duration of 3 ms.

Subjects performed a spatial attention task in a Posner paradigm (Figure 1) (Posner, 1980). Sound

spatialization was realized by one of the three spatialization conditions fixed within trials but pseudo-

randomized across trials. Subjects were instructed to fixate their gaze on a dot at the center of the

screen at the beginning of each trial. The fixation dot lasted for 1.2 s before an auditory cue was

presented. The auditory cue came from either left or right indicating where the target sound would

come from. A target sound started 0.8 s later from the cued location. At the same time a distractor

sound started from the opposite location of the target sound. Subjects were asked to report the syllable

sequence of the target sound by pressing a keyboard after the sounds finished and a response cue was

shown. Feedback about whether or not they correctly reported the syllables was given at the end of

every trial.

Each subject performed 450 randomized trials of this task, divided into 9 blocks each consisting of 50

trials. In total, every subject performed 150 trials for each of the three sound spatialization conditions

(75 trials attending left and 75 trials attending right). Prior to the test sessions, all participants received a

practice session to get familiarized with the task. Participants with a percentage of correct response

lower than 30% after 3 blocks of training (50 trials per block) were excluded from the study.

EEG Acquisition and Preprocessing. 32-channel scalp EEG data was recorded (Activetwo system with

Activeview acquisition software, Biosemi B.V.) in a sound proof booth (Eckel Industries, Inc.) while

subjects were performing the task. Two additional reference electrodes were placed on the earlobes.

Horizontal eye movements were recorded by two electrooculography (EOG) electrodes placed on the

outer canthi of each eye. Vertical eye movement was recorded by one EOG electrode placed below the

right eye. The timing of stimulus was controlled by Matlab (Mathworks) with Psychtoolbox (extension 3)

(Brainard, 1997; Delorme and Makeig, 2004). EEG preprocessing was conducted with Eeglab toolbox in

Matlab.

EEG data were corrected against the average of the two reference channels. Bad channels were marked

by manual selection during recording and automatically detected based on joint probability measures of

Eeglab (Delorme et al., 2007). EEG signals were then down-sampled to 256 Hz and epochs containing

responses to individual trials were extracted. Each epoch was baseline corrected against 100 ms prior to

the cue onset by removing the mean of the baseline period from the whole trial. ICA artifact rejection

was performed with Eeglab to remove components of eye movements, blinks, and muscle artifacts

(Delorme and Makeig, 2004). The maximum number of independent components rejected for each

subject was five. After ICA rejection, bad channels were removed and interpolated. Trials with a

maximum absolute value over 80 μV were rejected (Delorme et al., 2007). Two subjects with excessive

artifacts were removed from further EEG analysis because less than 50% of trials remained after

thresholding. For the rest of the 18 subjects, at least about two thirds of the trials (minimum was 48 out

of 75 trials) remained for each condition after artifact rejection. Trial numbers were equalized within

and across subjects by randomly selecting the minimum number of available trials (N = 48) for each

condition across the whole recording session.

Data analysis. Behavioral performance was quantified by the percentage of correct responses for every

one of the three syllables in the target stream and each spatialization condition. Behavioral results were

collapsed across the attend-left and attend-right trials. The percentages of correct response were then

normalized by logit transformation before parametric statistical testing was performed on the resulting

data.

ERP responses were evaluated for the second syllable of the target sound and distractor sound,

respectively. The reason we looked at the second syllable only is that 1) the first syllable of the target

and distractor aligned in time and therefore the ERPs were inseparable, and 2) the ERP amplitude in

response to the third syllable was small, and therefore more contaminated by noise. The preprocessed

data (details see EEG preprocessing) was bandpass filtered from 0.5 to 20 Hz by a finite impulse

response filter with Kaiser window design (β = 7.2, n = 1178). Data from four fronto-central channels (Cz,

Fz, FC1, and FC2) were averaged to get the auditory ERP response. We picked these four channels a

priori because auditory ERP responses in sensor space are largest in the fronto-central area of the scalp.

To quantify the amplitudes of ERP components, the maximum value within the window of 50 to 100 ms

after the second syllable onset was taken to be the P1 amplitude; the minimum value within the window

of 100 to 180 ms after the second syllable onset was calculated to be the N1 amplitude. The values

extracted from the selected windows were calculated for each channel and plotted onto a 2D scalp map

to generate topography plots. The values of the ERP components from the four selected channels were

then averaged and compared across different spatialization conditions.

To get the amplitude of alpha oscillation, the preprocessed EEG data was bandpass filtered to the alpha

range (8 to 14 Hz) before a Hilbert transform was applied. The magnitude of the resulting data was

taken as the extracted alpha power envelope. To get induced alpha power, the alpha power was

calculated for single trials first and then averaged across trials (Snyder and Large, 2005). The time course

of alpha power was baseline corrected against 700 ms before the auditory cue onset. GFP (Murray et al.,

2008; Skrandies, 1990) constitutes the spatial standard deviation across all scalp electrodes; it has been

used as a measurement to quantify the amount of alpha variation across the scalp (Lim et al., 2015). We

calculated the time courses of alpha GFP by taking the standard deviation of all electrodes. To quantify

the degree of alpha lateralization based on direction of attention, we calculated the Attentional

Modulation Index (AMI) of alpha power, defined as the alpha power difference between attended left

and attended right trials divided by the overall alpha power (Wöstmann et al., 2016). The AMI of alpha

was calculated for each time point, yielding the time course of AMI for each spatialization condition. We

then averaged the alpha AMI of each spatialization condition over the 800 ms immediately before

stimulus onset (-800 ms to 0 ms, re: onset). This is the period where subjects have been cued to orient

their spatial attention in preparation for the target sound, but before the speech streams begin. Scalp

topographies of the preparatory alpha AMI were plotted for each condition. Hemispheric lateralization

of alpha AMI was further compared across spatialization conditions and evaluated as the difference

between the left hemisphere and the right hemisphere.

For hypothesis testing, we compared group level differences between conditions with paired t-tests.

Due to the relatively small number of subjects (N=18), the Lilliefors test was performed prior to

statistical testing to check normality of the data. Data was considered normally distributed at P>0.05.

Data availability. All data and code used to conduct this study and analyze the results will be provided

via Zenodo immediately after acceptance of this manuscript for publication.

Acknowledgements

433   We thank Ashvini Meltoke for collecting part of the data, Gerald Kidd Jr. for providing the facilities for

434   individualized HRTF measurements, and Virginia Best for fruitful discussions. This work was supported

435   by the Austrian Science Fund (grant J3803-N30 to R.B.) and NICD (grant DC013825 to B.S.-C.).

References

Ahveninen, J., Hämäläinen, M., Jääskeläinen, I. P., Ahlfors, S. P., Huang, S., Lin, F.-H., Raij, T., et al. (**2011**). "Attention-driven auditory cortex short-term plasticity helps segregate relevant sounds from noise," *Proc. Natl. Acad. Sci.*, **108**, 4182–4187, doi: 10.1073/pnas.1016134108.

Alho, K., Medvedev, S. V., Pakhomov, S. V., Roudas, M. S., Tervaniemi, M., Reinikainen, K., Zeffiro, T., et al. (**1999**). "Selective tuning of the left and right auditory cortices during spatially directed attention," *Cogn. Brain Res.*, **7**, 335–341, doi: 10.1016/S0926-6410(98)00036-6.

Banerjee, S., Snyder, A. C., Molholm, S., and Foxe, J. J. (**2011**). "Oscillatory Alpha-Band Mechanisms and the Deployment of Spatial Attention to Anticipated Auditory and Visual Target Locations: Supramodal or Sensory-Specific Control Mechanisms?," *J. Neurosci.*, **31**, 9923–9932, doi: 10.1523/JNEUROSCI.4660-10.2011.

Baumgartner, R., Reed, D. K., Tóth, B., Best, V., Majdak, P., Colburn, H. S., and Shinn-Cunningham, B. (**2017**). "Asymmetries in behavioral and neural responses to spectral cues demonstrate the generality of auditory looming bias," *Proc. Natl. Acad. Sci.*, **114**, 9743–9748, doi: 10.1073/pnas.1703247114.

Blankertz, B., Lemm, S., Treder, M., Haufe, S., and Müller, K.-R. (**2011**). "Single-trial analysis and classification of ERP components — A tutorial," *NeuroImage*, **56**, 814–825, doi: 10.1016/J.NEUROIMAGE.2010.06.048.

Blauert, J. (**1997**). *Spatial hearing. The Psychophysics of Human Sound Localization*, (MIT-Press, Cambridge, MA), pp. 497.

Boersma, P. (**2001**). "Praat, a system for doing phonetics by computer," *Glot Int.*, **5**, 341–345.

Brainard, D. H. (**1997**). "The psychophysics toolbox," *Spat. Vis.*, **10**, 433–436.

Callan, A., Callan, D. E., and Ando, H. (**2013**). "Neural correlates of sound externalization," *NeuroImage*, **66**, 22–27, doi: 10.1016/J.NEUROIMAGE.2012.10.057.

Callan, A., Callan, D. E., and Ando, H. (**2013**). "Neural correlates of sound externalization," *NeuroImage*, **66**, 22–27, doi: 10.1016/j.neuroimage.2012.10.057.

Chait, M., de Cheveigné, A., Poeppel, D., and Simon, J. Z. (**2010**). "Neural dynamics of attending and ignoring in human auditory cortex," *Neuropsychologia*, **48**, 3262–3271, doi: 10.1016/J.NEUROPSYCHOLOGIA.2010.07.007.

Cherry, E. C. (**1953**). "Some Experiments on the Recognition of Speech, with One and with Two Ears," *J. Acoust. Soc. Am.*, **25**, 975–979, doi: 10.1121/1.1907229.

Choi, I., Rajaram, S., Varghese, L. A., and Shinn-Cunningham, B. G. (**2013**). "Quantifying attentional
modulation of auditory-evoked cortical responses from single-trial electroencephalography,"
Front. Hum. Neurosci., **7**, 115, doi: 10.3389/fnhum.2013.00115.

Couperus, J. W. and Mangun, G. R. (**2010**). "Signal enhancement and suppression during visual–spatial
selective attention," *Brain Res.*, **1359**, 155–177, doi: 10.1016/J.BRAINRES.2010.08.076.

Culling, J. F., Hawley, M. L., and Litovsky, R. Y. (**2004**). "The role of head-induced interaural time and
level differences in the speech reception threshold for multiple interfering sound sources," *J.
Acoust. Soc. Am.*, **116**, 1057–1065, doi: 10.1121/1.1772396.

Cusack, R., Carlyon, R. P., and Robertson, I. H. (**2001**). "Auditory Midline and Spatial Discrimination in
Patients with Unilateral Neglect," *Cortex*, **37**, 706–709, doi: 10.1016/S0010-9452(08)70620-8.

Dahmen, J. C., Keating, P., Nodal, F. R., Schulz, A. L., and King, A. J. (**2010**). "Adaptation to Stimulus
Statistics in the Perception and Neural Representation of Auditory Space," *Neuron*, **66**, 937–948,
doi: 10.1016/J.NEURON.2010.05.018.

Dai, L., Best, V., and Shinn-Cunningham, B. G. (**2018**). "Sensorineural hearing loss degrades behavioral
and physiological measures of human spatial selective auditory attention," *Proc. Natl. Acad. Sci.*,
115, E3286--E3295, doi: 10.1073/pnas.1721226115.

Delorme, A. and Makeig, S. (**2004**). "EEGLAB: an open source toolbox for analysis of single-trial EEG
dynamics including independent component analysis," *J. Neurosci. Methods*, **134**, 9–21, doi:
10.1016/j.jneumeth.2003.10.009.

Delorme, A., Sejnowski, T., and Makeig, S. (**2007**). "Enhanced detection of artifacts in EEG data using
higher-order statistics and independent component analysis," *NeuroImage*, **34**, 1443–1449, doi:
10.1016/j.neuroimage.2006.11.004.

Ellinger, R. L., Jakien, K. M., and Gallun, F. J. (**2017**). "The role of interaural differences on speech
intelligibility in complex multi-talker environments," *J. Acoust. Soc. Am.*, **141**, EL170–EL176, doi:
10.1121/1.4976113.

Foxe, J. J. and Snyder, A. C. (**2011**). "The role of alpha-band brain oscillations as a sensory suppression
mechanism during selective attention," *Front. Psychol.*, **2**, 154, doi: 10.3389/fpsyg.2011.00154.

Freunberger, R., Höller, Y., Griesmayr, B., Gruber, W., Sauseng, P., and Klimesch, W. (**2008**). "Functional
similarities between the P1 component and alpha oscillations," *Eur. J. Neurosci.*, **27**, 2330–2340,
doi: 10.1111/j.1460-9568.2008.06190.x.

Fukuda, K. and Vogel, E. K. (**2009**). "Human Variation in Overriding Attentional Capture," *J. Neurosci.*, **29**,
8726–8733, doi: 10.1523/JNEUROSCI.2145-09.2009.

Getzmann, S. and Lewald, J. (**2010**). "Effects of natural versus artificial spatial cues on

electrophysiological correlates of auditory motion," *Hear. Res.*, **259**, 44–54, doi:

10.1016/j.heares.2009.09.021.

Giuliano, R. J., Karns, C. M., Neville, H. J., and Hillyard, S. A. (**2014**). "Early Auditory Evoked Potential Is

Modulated by Selective Attention and Related to Individual Differences in Visual Working

Memory Capacity," *J. Cogn. Neurosci.*, doi: 10.1162/jocn_a_00684.

Glyde, H., Buchholz, J. M., Dillon, H., Cameron, S., and Hickson, L. (**2013**). "The importance of interaural

time differences and level differences in spatial release from masking," *J. Acoust. Soc. Am.*, **134**,

EL147–EL152, doi: 10.1121/1.4812441.

Haegens, S., Luther, L., and Jensen, O. (**2011**). "Somatosensory Anticipatory Alpha Activity Increases to

Suppress Distracting Input," *J. Cogn. Neurosci.*, **24**, 677–685, doi: 10.1162/jocn_a_00164.

Hartmann, W. M. and Wittenberg, A. (**1996**). "On the externalization of sound images," *J Acoust Soc Am*,

99, 3678–3688.

Herrmann, C. S. and Knight, R. T. (**2001**). "Mechanisms of human attention: event-related potentials and

oscillations," *Neurosci. Biobehav. Rev.*, **25**, 465–476, doi: 10.1016/S0149-7634(01)00027-6.

Higgins, N. C., McLaughlin, S. A., Rinne, T., and Stecker, G. C. (**2017**). "Evidence for cue-independent

spatial representation in the human auditory cortex during active listening," *Proc. Natl. Acad.*

Sci., **114**, E7602--E7611, doi: 10.1073/pnas.1707522114.

Hillyard, S. A. and Anllo-Vento, L. (**1998**). "Event-related brain potentials in the study of visual selective

attention," *Proc. Natl. Acad. Sci.*, **95**, 781–787, doi: 10.1073/pnas.95.3.781.

Hopfinger, J. B., Luck, S. J., and Hillyard, S. A. (**2004**). "Selective Attention: Electrophysiological and

Neuromagnetic Studies," in *The cognitive neurosciences, 3rd ed*, (MIT Press, Cambridge, MA,

US), pp. 561–574.

Itoh, K., Yumoto, M., Uno, A., Kurauchi, T., and Kaga, K. (**2000**). "Temporal stream of cortical

representation for auditory spatial localization in human hemispheres," *Neurosci. Lett.*, **292**,

215–219, doi: 10.1016/S0304-3940(00)01465-8.

Jensen, O. and Mazaheri, A. (**2010**). "Shaping Functional Architecture by Oscillatory Alpha Activity:

Gating by Inhibition," *Front. Hum. Neurosci.*, **4**, 186, doi: 10.3389/fnhum.2010.00186.

Johnson, B. W. and Hautus, M. J. (**2010**). "Processing of binaural spatial information in human auditory

cortex: Neuromagnetic responses to interaural timing and level differences," *Neuropsychologia*,

48, 2610–2619, doi: 10.1016/J.NEUROPSYCHOLOGIA.2010.05.008.

Kelly, S. P. (**2006**). "Increases in Alpha Oscillatory Power Reflect an Active Retinotopic Mechanism for
Distracter Suppression During Sustained Visuospatial Attention," *J. Neurophysiol.*, **95**, 3844–
3851, doi: 10.1152/jn.01234.2005.

Kidd, G., Mason, C. R., Best, V., and Marrone, N. (**2010**). "Stimulus factors influencing spatial release
from speech-on-speech masking," *J. Acoust. Soc. Am.*, **128**, 1965–1978, doi: 10.1121/1.3478781.

Klimesch, W. (**2011**). "Evoked alpha and early access to the knowledge system: The P1 inhibition timing
hypothesis," *Brain Res.*, **1408**, 52–71, doi: 10.1016/j.brainres.2011.06.003.

Klimesch, W. (**2012**). "Alpha-band oscillations, attention, and controlled access to stored information,"
Trends Cogn. Sci., **16**, 606–617, doi: 10.1016/J.TICS.2012.10.007.

Klimesch, W., Sauseng, P., and Hanslmayr, S. (**2007**). "EEG alpha oscillations: The inhibition-timing
hypothesis," *Brain Res. Rev.*, **53**, 63–88, doi: 10.1016/j.brainresrev.2006.06.003.

Kong, L., Michalka, S. W., Rosen, M. L., Sheremata, S. L., Swisher, J. D., Shinn-Cunningham, B. G., and
Somers, D. C. (**2014**). "Auditory Spatial Attention Representations in the Human Cerebral
Cortex," *Cereb. Cortex*, **24**, 773–784, doi: 10.1093/cercor/bhs359.

Leino, S., May, P. J. C., Alku, P., Liikkanen, L. A., and Tiitinen, H. (**2007**). "The contribution of high
frequencies to human brain activity underlying horizontal localization of natural spatial sounds,"
BMC Neurosci., **8**, 78, doi: 10.1186/1471-2202-8-78.

Lim, S.-J., Wöstmann, M., and Obleser, J. (**2015**). "Selective Attention to Auditory Memory Neurally
Enhances Perceptual Precision," *J. Neurosci.*, **35**, 16094–16104, doi: 10.1523/JNEUROSCI.2674-
15.2015.

Loiselle, L. H., Dorman, M. F., Yost, W. A., Cook, S. J., and Gifford, R. H. (**2016**). "Using ILD or ITD Cues for
Sound Source Localization and Speech Understanding in a Complex Listening Environment by
Listeners With Bilateral and With Hearing-Preservation Cochlear Implants," *J. Speech Lang. Hear.*
Res. JSLHR, **59**, 810–818, doi: 10.1044/2015_JSLHR-H-14-0355.

Luck, S. J. (**1995**). "Multiple mechanisms of visual-spatial attention: recent evidence from human
electrophysiology," *Behav. Brain Res.*, **71**, 113–123, doi: 10.1016/0166-4328(95)00041-0.

Majdak, P., Baumgartner, R., and Jenny, C. (**2019**). "Formation of three-dimensional auditory space,"
*ArXiv:*1901.03990 Q-Bio (13 January 2019).

Marzecová, A., Schettino, A., Widmann, A., SanMiguel, I., Kotz, S. A., and Schröger, E. (**2018**).
"Attentional gain is modulated by probabilistic feature expectations in a spatial cueing task: ERP
evidence," *Sci. Rep.*, **8**, doi: 10.1038/s41598-017-18347-1.

Mesgarani, N. and Chang, E. F. (**2012**). "Selective cortical representation of attended speaker in multi-
562   talker speech perception," *Nature*, **485**, 233–236, doi: 10.1038/nature11020.

563   Middlebrooks, J. C. (**2015**). "Sound localization," *Handb. Clin. Neurol.*, *The Human Auditory System*, **129**,
564   99–116, doi: 10.1016/B978-0-444-62630-1.00006-8.

565   Murray, M. M., Brunet, D., and Michel, C. M. (**2008**). "Topographic ERP Analyses: A Step-by-Step Tutorial
566   Review," *Brain Topogr.*, **20**, 249–264, doi: 10.1007/s10548-008-0054-5.

567   Oppenheim, A. V. and Schafer, R. W. (**1975**). *Digital Signal Processing*, (Prentice-Hall, Englewood Cliffs,
568   NJ).

569   Palomäki, K. J., Tiitinen, H., Mäkinen, V., May, P. J. C., and Alku, P. (**2005**). "Spatial processing in human
570   auditory cortex: The effects of 3D, ITD, and ILD stimulation techniques," *Cogn. Brain Res.*, **24**,
571   364–379, doi: 10.1016/j.cogbrainres.2005.02.013.

572   Posner, M. I. (**1980**). "Orienting of attention," *Q. J. Exp. Psychol.*, **32**, 3–25, doi:
573   10.1080/00335558008248231.

574   Rihs, T. A., Michel, C. M., and Thut, G. (**2007**). "Mechanisms of selective inhibition in visual spatial
575   attention are indexed by α-band EEG synchronization," *Eur. J. Neurosci.*, **25**, 603–610, doi:
576   10.1111/j.1460-9568.2007.05278.x.

577   Rihs, T. A., Michel, C. M., and Thut, G. (**2009**). "A bias for posterior α-band power suppression versus
578   enhancement during shifting versus maintenance of spatial attention," *NeuroImage*, **44**, 190–
579   199, doi: 10.1016/j.neuroimage.2008.08.022.

580   Rimmele, J. M., Zion Golumbic, E., Schröger, E., and Poeppel, D. (**2015**). "The effects of selective
581   attention and speech acoustics on neural speech-tracking in a multi-talker scene," *Cortex*, **68**,
582   144–154, doi: 10.1016/J.CORTEX.2014.12.014.

583   Röder, B., Teder-Sälejärvi, W., Sterr, A., Rösler, F., Hillyard, S. A., and Neville, H. J. (**1999**). "Improved
584   auditory spatial tuning in blind humans," *Nature*, **400**, 162–166, doi: 10.1038/22106.

585   Sach, A. J., Hill, N. I., and Bailey, P. J. (**2000**). "Auditory spatial attention using interaural time
586   differences," *J. Exp. Psychol. Hum. Percept. Perform.*, **26**, 717–729, doi: 10.1037//0096-
587   1523.26.2.717.

588   Salminen, N. H., Takanen, M., Santala, O., Lamminsalo, J., Altoè, A., and Pulkki, V. (**2015**). "Integrated
589   processing of spatial cues in human auditory cortex," *Hear. Res.*, **327**, 143–152, doi:
590   10.1016/J.HEARES.2015.06.006.

Samaha, J., Bauer, P., Cimaroli, S., and Postle, B. R. (**2015**). "Top-down control of the phase of alpha-

band oscillations as a mechanism for temporal prediction," *Proc. Natl. Acad. Sci. U. S. A.*, **112**,

8439–8444, doi: 10.1073/pnas.1503686112.

Samaha, J., Sprague, T. C., and Postle, B. R. (**2016**). "Decoding and Reconstructing the Focus of Spatial

Attention from the Topography of Alpha-band Oscillations," *J. Cogn. Neurosci.*, 1–8, doi:

10.1162/jocn_a_00955.

Sauseng, P., Klimesch, W., Stadler, W., Schabus, M., Doppelmayr, M., Hanslmayr, S., Gruber, W. R., et al.

(**2005**). "A shift of visual spatial attention is selectively associated with human EEG alpha

activity," *Eur. J. Neurosci.*, **22**, 2917–2926, doi: 10.1111/j.1460-9568.2005.04482.x.

Schreuder, M., Rost, T., and Tangermann, M. (**2011**). "Listen, You are Writing! Speeding up Online

Spelling with a Dynamic Auditory BCI," *Front. Neurosci.*, **5**, 112, doi: 10.3389/fnins.2011.00112.

Schröger, E. (**1996**). "Interaural time and level differences: Integrated or separated processing?," *Hear.*

Res., **96**, 191–198, doi: 10.1016/0378-5955(96)00066-4.

Shinn-Cunningham, B. G., Kopco, N., and Martin, T. J. (**2005**). "Localizing nearby sound sources in a

classroom: Binaural room impulse responses," *J. Acoust. Soc. Am.*, **117**, 3100–3115, doi:

10.1121/1.1872572.

Siegel, M., Donner, T. H., and Engel, A. K. (**2012**). "Spectral fingerprints of large-scale neuronal

interactions," *Nat. Rev. Neurosci.*, **13**, 121–134, doi: 10.1038/nrn3137.

Skrandies, W. (**1990**). "Global field power and topographic similarity," *Brain Topogr.*, **3**, 137–141.

Slagter, H. A., Prinssen, S., Reteig, L. C., and Mazaheri, A. (**2016**). "Facilitation and inhibition in attention:

Functional dissociation of pre-stimulus alpha activity, P1, and N1 components," *NeuroImage*,

125, 25–35, doi: 10.1016/j.neuroimage.2015.09.058.

Snyder, J. S. and Large, E. W. (**2005**). "Gamma-band activity reflects the metric structure of rhythmic

tone sequences," *Brain Res. Cogn. Brain Res.*, **24**, 117–126, doi:

10.1016/j.cogbrainres.2004.12.014.

Stevens, C., Fanning, J., Coch, D., Sanders, L., and Neville, H. (**2008**). "Neural mechanisms of selective

auditory attention are enhanced by computerized training: Electrophysiological evidence from

language-impaired and typically developing children," *Brain Res.*, **1205**, 55–69, doi:

10.1016/J.BRAINRES.2007.10.108.

Tardif, E., Murray, M. M., Meylan, R., Spierer, L., and Clarke, S. (**2006**). "The spatio-temporal brain

dynamics of processing and integrating sound localization cues in humans," *Brain Res.*, **1092**,

161–176, doi: 10.1016/J.BRAINRES.2006.03.095.

Ungan, P., Yagcioglu, S., and Goksoy, C. (**2001**). "Differences between the N1 waves of the responses to

interaural time and intensity disparities: scalp topography and dipole sources," *Clin.*

Neurophysiol., **112**, 485–498, doi: 10.1016/S1388-2457(00)00550-2.

Worden, M. S., Foxe, J. J., Wang, N., and Simpson, G. V. (**2000**). "Anticipatory biasing of visuospatial

attention indexed by retinotopically specific alpha-band electroencephalography increases over

occipital cortex," *J. Neurosci.*, **20**, RC63, doi: 10.1523/JNEUROSCI.20-06-j0002.2000.

Wöstmann, M., Herrmann, B., Maess, B., and Obleser, J. (**2016**). "Spatiotemporal dynamics of auditory

attention synchronize with speech," *Proc. Natl. Acad. Sci.*, **113**, 3873–3878, doi:

10.1073/pnas.1523357113.

Wyart, V., Dehaene, S., and Tallon-Baudry, C. (**2012**). "Early dissociation between neural signatures of

endogenous spatial attention and perceptual awareness during visual masking," *Front. Hum.*

Neurosci., **6**, 16, doi: 10.3389/fnhum.2012.00016.