Impoverished auditory cues limit engagement of brain networks controlling spatial selective attention

Authors: Yuqi Deng[1], Inyong Choi[2], Barbara Shinn-Cunningham[1,3], Robert Baumgartner[1,4*]

Affiliations:

[1] Biomedical Engineering, Boston University, Boston, MA 02215, USA

[2] Communication Sciences & Disorders, University of Iowa, Iowa City, IA 52242, USA

[3] Neuroscience Institute, Carnegie Mellon University, Pittsburgh, PA 15213 USA

[4] Acoustics Research Institute, Austrian Academy of Sciences, 1040 Vienna, Austria

[*] Corresponding Author: robert.baumgartner@oeaw.ac.at

Author contributions: R.B., B.S.-C., Y.D., and I.C. designed research; Y.D. and R.B. performed research; Y.D. and R.B. analyzed data; and Y.D., R.B., B.S.-C., and I.C. wrote the paper.

The authors have no competing interests to declare.

Abstract

Spatial selective attention enables listeners to process a signal of interest in natural settings. However,

most past studies on auditory spatial attention used impoverished spatial cues: presenting competing

sounds to different ears, using only interaural differences in time (ITDs) and/or intensity (IIDs), or using

non-individualized head-related transfer functions (HRTFs). Here we tested the hypothesis that

impoverished spatial cues impair spatial auditory attention by only weakly engaging relevant cortical

networks. Eighteen normal-hearing listeners reported the content of one of two competing syllable

streams simulated at roughly +30 ˚ and -30˚ azimuth. The competing streams consisted of syllables from

two different-sex talkers. Spatialization was based on natural spatial cues (individualized HRTFs),

individualized IIDs, or generic ITDs. We measured behavioral performance as well as

electroencephalographic markers of selective attention. Behaviorally, subjects recalled target streams

most accurately with natural cues. Neurally, spatial attention significantly modulated early evoked

sensory response magnitudes only for natural cues, not in conditions using only ITDs or IIDs. Consistent

with this, parietal oscillatory power in the alpha band (8-14 Hz; associated with filtering out distracting

events from unattended directions) showed significantly less attentional modulation with isolated spatial

cues than with natural cues. Our findings support the hypothesis that spatial selective attention networks

are only partially engaged by impoverished spatial auditory cues. These results not only suggest that

studies using unnatural spatial cues underestimate the neural effects of spatial auditory attention, they also

illustrate the importance of preserving natural spatial cues in assistive listening devices to support robust

attentional control.

Keywords: auditory spatial selective attention, head-related transfer functions, electroencephalography

1 Introduction

Spatial hearing is crucial to selectively attend to sounds of interest in everyday social settings. The remarkable ability of normal-hearing listeners to focus on a sound source within a complex acoustic scene is often referred to as "the cocktail party phenomenon," and has a rich history (Cherry, 1953). Nevertheless, the mechanisms controlling spatial selective attention are still poorly understood. Acoustically, in everyday situations, the two ears provide the listener with a listener-specific combination of spatial cues that include interaural time and intensity differences (ITDs and IIDs, respectively), as well as spectral cues caused by acoustical filtering of the pinnae (Blauert, 1997a). Together, these cues, captured by individualized head-related transfer functions (HRTFs), allow the brain to create a clear, punctate internal representation of the location of sound sources in the environment (Majdak et al., 2018; Middlebrooks, 2015).

When only isolated or impoverished spatial cues are present, auditory localization performance degrades and the natural perception of external auditory objects may even collapse into the listener's head (Baumgartner et al., 2017; Callan et al., 2013; Cubick et al., 2018; Hartmann and Wittenberg, 1996). Nevertheless, degraded or isolated ITDs and IIDs still create a strong sense of lateralization within the head; moreover, even highly impoverished spatial cues can be used to achieve spatial release from speech-on-speech masking, behaviorally (Cubick et al., 2018; Culling et al., 2004; Ellinger et al., 2017; Glyde et al., 2013; Kidd et al., 2010; Loiselle et al., 2016). The relative importance of ITDs and IIDs in spatial release from masking remains unclear, with past studies reporting conflicting results when directly comparing different binaural conditions (Ellinger et al., 2017; Glyde et al., 2013; Higgins et al., 2017; Shinn-Cunningham et al., 2005). More importantly, it is a puzzle as to why realistic and degraded spatial cues yield at best small behavioral differences in masking release even though spatial perception is clearly degraded when cues are impoverished (e.g., Cubick et al., 2018).

Previous electroencephalography (EEG) and magnetoencephalography (MEG) studies have demonstrated that rich spatial cues in sound stimuli lead to different cortical activity compared to using isolated cues

during sound localization (Callan et al., 2013; Leino et al., 2007; Palomäki et al., 2005) and auditory

motion processing (Getzmann and Lewald, 2010). However, the apparently minor behavioral

consequences of using unnatural, non-individualized spatial cues on spatial release from masking,

combined with the ease of implementing studies with simple, non-individualized spatial cues, led to their

wide usage in auditory neuroscience studies (Cusack et al., 2001; Dahmen et al., 2010; Dai et al., 2018;

Itoh et al., 2000; Kong et al., 2014; Sach et al., 2000). Indeed, in the auditory neuroscience literature,

many studies did not even present true binaural signals, but instead studied "spatial" attention by using

dichotic signals, with one sound presented monaurally to one ear and a competing sound presented

monaurally to the other ear (Ahveninen et al., 2011; Alho et al., 1999b; Das et al., 2016; Wöstmann et al.,

2016). These studies implicitly assumed that because listeners were able to use impoverished spatial cues

to listen to one sound from a particular (relative) direction, the cognitive networks responsible for

controlling spatial attention must be engaged just as they are when listening to rich, natural spatial cues.

Nonetheless, it is unclear whether and how engagement of higher-order cognitive processes such as

deployment of selective attention is affected by the use of unnatural or impoverished spatial cues.

Modulation of neural signatures, such as event-related potentials (ERPs) and induced oscillatory activity,

is often taken as evidence of effective attentional control (Herrmann and Knight, 2001; Siegel et al.,

2012). In particular, auditory spatial attention is known to modulate early sensory ERPs in the N1 time

range (processing latencies of 100 to 150 ms; see Choi et al., 2013; Röder et al., 1999), whereas

modulation of P1 ERPs (50 to 100 ms) has only recently been demonstrated in a free field experiment

(Giuliano et al., 2014). Induced alpha oscillation (8 to 14 Hz) has been hypothesized to function as an

information gating mechanism (Klimesch et al., 2007). During auditory spatial attention, parietal alpha

power often decreases in the contralateral hemisphere of attended stimuli and/or increases in the

ipsilateral hemisphere (Banerjee et al., 2011; Lim et al., 2015; Wöstmann et al., 2016). These neural

modulations constitute objective metrics of the efficacy of attentional control.

Here, we test listeners in a selective attention paradigm with simultaneous, spatially separated talkers. We

use the aforementioned EEG measures to compare both perceptual ability and the neural signatures of

attentional control for simulations with impoverished vs. natural spatial cues. Eighteen subjects performed

an auditory spatial attention task with two competing streams located at roughly +30 ˚ and -30˚ azimuth

(Figure 1). On every trial, listeners were cued by an auditory cue to attend to either the left or right stream

and report the content of the cued stream. The competing streams consisted of syllables (/ba/, /da/ or /ga/)

from two different-sex talkers. Sound stimuli (including the cuing sound) were spatialized using three

different levels of naturalness and richness: 1) generic ITDs only, 2) individualized IIDs, or 3)

individualized HRTFs containing all of the naturally occurring spatial cues a listener experiences in the

everyday world. We show that behavioral performance is better when listeners hear natural,

individualized spatial cues than when they hear impoverished cues. Importantly, only natural spatial cues

yield significant attentional modulation of P1 amplitudes. Moreover, induced alpha activity is less robust

and poorly lateralized with isolated spatial cues compared to rich, natural spatial cues.

2 Materials and Methods

2.1 Subjects

Twenty-one paid volunteers and one author within the age of 18-42 years (M = 22.9, SD = 5.5;

12 females, 10 males) participated in this study. None of the subjects had audiometric thresholds greater

than 20 dB for frequencies from 250 Hz to 8 kHz. All participants gave informed consent as approved by

the Boston University Institutional Review Board. Two subjects were withdrawn from the study due to

the inability to perform the task (percentage of correct response less than 30% after training), and two

subjects were removed during EEG data preprocessing due to excessive artifacts. Therefore 18 subjects

remained for further analysis (N = 18).

2.2 Stimuli and Procedure

The sound stimuli consisted of consonant-vowel syllables (/ba/, /da/, & /ga/), each 0.4 s in duration. These

syllables were recorded from three talkers that naturally differed in fundamental frequency (F0). Details

on stimulus are provided in Stimulus Presentation. Cue and stimuli were presented via earphones (ER-2,

Etymotic Research, Inc.) and spatialized to approximately ±30° azimuth (0° elevation). Three different

spatialization conditions were used: HRTF, IID, and ITD. In the HRTF condition, individualized HRTFs,

providing natural combinations of ITDs, IIDs, and spectral cues, were used (See Individual HRTF

Measurement for measurement methods). In the IID condition, ITDs were removed from the

individualized HRTFs by computing minimum-phase representations of the filters via removing the non-

causal part of the cepstrum. Hence, the IID and HRTF conditions provided the same monaural magnitude

spectra and thus the same energetic advantage of the ear ipsilateral to the target while differing in spatial

perception. In the ITD condition, spatialization was based on simply delaying the signal presented to the

contralateral ear by 300 μs, thus providing no energetic advantage to the ipsilateral ear. This spatialization

method was tested due to its popularity in auditory neuroscience.

The auditory cue was a single syllable /ba/ spoken by a low-pitch male voice (F0 = 91 Hz, estimated by

Praat software) (Boersma, 2001). The following target and distractor streams both consisted of three

syllables randomly chosen out of the set of three syllables (with replacement). The target stream was

spoken by either a female (F0 = 189 Hz) or a high-pitch male talker (F0 = 125 Hz), and the distractor

stream was spoken by the other talker different than the target stream. The first syllable of the target and

distractor sound overlapped in time, while the latter two syllables were separated by 200 ms, onset to

onset (Figure 1). To avoid engagement of temporal attention rather than spatial attention, the assignment

of the target stream being leading or lagging was equally distributed across trials. In the leading stream,

the onsets of all three syllables were separated by 400 ms; in the lagging stream, the onsets of the first and

the second syllable were separated by 600 ms, whereas those of the second and the third syllable were

separated by 400 ms. All sound stimuli were presented at a sound pressure level of approximately 75 dB.

Figure 1

Fig. 1. *Auditory spatial attention task with two competing streams was used to assess the*

consequence of impoverished auditory spatial cues on neural proxies of attention control. An

auditory cue was presented first from the location of the upcoming target stream, processed by the

same spatialization scheme as the upcoming mixture. Following the cue, the competing streams

began, one from around +30° the other from around -30° azimuth. Listeners were asked to recall

the syllable sequence presented from the cued side. The first syllables of both streams were

temporally aligned; however, the latter two syllables in the competing streams were staggered,

enabling us to isolate neural responses to each. Feedback was provided after every trial.

2.3 Task

Subjects performed a spatial attention task in a Posner paradigm (Figure 1) (Posner et al., 1980). Sound

spatialization was realized by one of the three spatialization conditions fixed within trials but pseudo-

randomized across trials. Subjects were instructed to fixate their gaze on a dot at the center of the screen

at the beginning of each trial. The fixation dot lasted for 1.2 s before an auditory cue was presented. The

auditory cue came from either left or right indicating where the target sound would come from. A target

sound started 0.8 s later from the cued location. At the same time a distractor sound started from the

opposite location of the target sound. Subjects were asked to report the syllable sequence of the target

sound by pressing a keyboard after the sounds finished and a response cue was shown. Feedback about

whether or not they correctly reported the syllables was given at the end of every trial.

Each subject performed 450 randomized trials of this task, divided into 9 blocks each consisting of 50

trials. In total, every subject performed 150 trials for each of the three sound spatialization conditions

(75 trials attending left and 75 trials attending right). Prior to the test sessions, all participants received a

practice session to get familiarized with the task. Participants with a percentage of correct response lower

than 30% after 3 blocks of training (50 trials per block) were excluded from the study.

2.4 EEG Acquisition and Preprocessing

32-channel scalp EEG data was recorded (Activetwo system with Activeview acquisition software,

Biosemi B.V.) in a sound proof booth (Eckel Industries, Inc.) while subjects were performing the task.

Two additional reference electrodes were placed on the earlobes. Horizontal eye movements were

recorded by two electrooculography (EOG) electrodes placed on the outer canthi of each eye. Vertical eye

movement was recorded by one EOG electrode placed below the right eye. The timing of stimulus was

controlled by Matlab (Mathworks) with Psychtoolbox (extension 3) (Brainard, 1997).

EEG preprocessing was conducted in Matlab with Eeglab toolbox (Delorme and Makeig, 2004). EEG

data were corrected against the average of the two reference channels. Bad channels were marked by

manual selection during recording and automatically detected based on joint probability measures of

Eeglab. EEG signals were then down-sampled to 256 Hz and epochs containing responses to individual

trials were extracted. Each epoch was baseline corrected against 100 ms prior to the cue onset by

removing the mean of the baseline period from the whole trial. ICA artifact rejection was performed with

Eeglab to remove components of eye movements, blinks, and muscle artifacts. The maximum number of

independent components rejected for each subject was five. After ICA rejection, bad channels were

removed and interpolated. Trials with a maximum absolute value over 80 µV were rejected (Delorme et

al., 2007). Two subjects with excessive artifacts were removed from further EEG analysis because less

than 50% of trials remained after thresholding. For the rest of the 18 subjects, at least about two thirds of

the trials (minimum was 48 out of 75 trials) remained for each condition after artifact rejection. Trial

numbers were equalized within and across subjects by randomly selecting the minimum number of

available trials (N = 48) for each condition across the whole recording session.

2.5 Data analysis

Behavioral performance was quantified by the percentage of correct responses for every one of the three

syllables in the target stream and each spatialization condition. Behavioral results were collapsed across

the attend-left and attend-right trials. The percentages of correct response were then normalized by logit

transformation before parametric statistical testing was performed on the resulting data.

ERP responses were evaluated for the second syllable of the target sound and distractor sound,

respectively. The reason we looked at the second syllable only is that 1) the first syllable of the target and

distractor aligned in time and therefore the ERPs were inseparable, and 2) the ERP amplitude in response

to the third syllable was small, and therefore more contaminated by noise. ERP components were then

extracted from the time series data. The preprocessed data (details see EEG Preprocessing Procedures)

was bandpass filtered from 0.5 to 20 Hz by a finite impulse response filter with Kaiser window design (β

= 7.2, n = 1178). Data from four fronto-central channels (Cz, Fz, FC1, and FC2) were averaged to get the

auditory ERP response. We picked these four channels a priori because auditory ERP responses in sensor

space are largest in the fronto-central area of the scalp. To quantify the amplitudes of ERP components,

the maximum value within the window of 50 to 100 ms after the second syllable onset was taken to be the

P1 amplitude; the minimum value within the window of 100 to 180 ms after the second syllable onset was

calculated to be the N1 amplitude. The values extracted from the selected windows were calculated for

each channel and plotted onto a 2D scalp map to generate topography plots. The values of the ERP

components from the four selected channels were then averaged and compared across different

spatialization conditions.

To get the amplitude of alpha oscillation, the preprocessed EEG data was bandpass filtered to the alpha

range (8 to 14 Hz) before a Hilbert transform was applied. The magnitude of the resulting data was taken

as the extracted alpha power envelope. To get induced alpha power, the alpha power was calculated for

single trials first and then averaged across trials (Snyder and Large, 2005). The time course of alpha

power was baseline corrected against 700 ms before the auditory cue onset. GFP (Murray et al., 2008;

Skrandies, 1990) constitutes the spatial standard deviation across all scalp electrodes; it has been used as a

measurement to quantify the amount of alpha variation across the scalp (Lim et al., 2015). We calculated

the time courses of alpha GFP by taking the standard deviation of all electrodes. To quantify the degree of

alpha modulation based on direction of attention, we calculated the Attentional Modulation Index (AMI)

of alpha power, defined as the alpha power difference between attended left and attended right trials

divided by the overall alpha power (Wöstmann et al., 2016). The AMI of alpha was calculated for each

time point, yielding the time course of AMI for each spatialization condition. We then averaged the alpha

AMI of each spatialization condition over the 800 ms immediately before stimulus onset (-800 ms to

0 ms, re: onset). This is the period where subjects have been cued to orient their spatial attention in

preparation for the target sound, but before the speech streams begin. Scalp topographies of the

preparatory alpha AMI were plotted for each condition. Hemispheric lateralization of alpha AMI was

further compared across spatialization conditions and evaluated as the difference between the left

hemisphere and the right hemisphere. Calculated in this way, the AMI is expected to be positive in left

and negative in right parietal channels.

For testing the significance of different means across conditions, we conducted repeated measures

ANOVAs followed by post-hoc analyses for all significant main effects and interactions using Fisher's

least significant difference procedure. We separately tested whether condition means differed

significantly from zero using Bonferroni-corrected t-tests (P_{adj}). The Lilliefors test was performed prior to

statistical testing to check normality of the data. Data was considered normally distributed at $P > 0.05$.

Prior to statistical analysis of behavioral performance, the percentages of correctly reported syllable were

logit transformed in order to obtain normally distributed data.

3 Results

3.1 Natural spatial cues facilitate behavioral performance

Percentages of correctly recalling each syllable of the target stream differed across the three spatialization

conditions (Figure 2; 1st syllable: $F_{(2,34)} = 25.25$, $P < 0.001$; 2nd syllable: $F_{(2,34)} = 6.27$, $P = 0.005$; 3rd

syllable: $F_{(2,34)} = 5.60$, $P = 0.008$). For the first syllable, where the target and distractor sounds overlapped

in time, subjects were least accurate in the ITD condition compared to the IID condition ($t_{(34)} = 5.31$,

$P < 0.001$) and HRTF condition ($t_{(34)} = 6.74$, $P < 0.001$). However, no statistically significant difference

was observed between IID and HRTF conditions for that syllable ($t_{(34)} = 1.43$, $P = 0.16$). For the second

and the third syllable, where target and distractor streams occurred staggered in time, subjects performed

better in the HRTF condition than in both the ITD condition (2nd syllable: $t_{(34)} = 3.27$, $P = 0.002$;

3rd syllable: $t_{(34)} = 3.33$, $P = 0.002$) and the IID condition (2nd syllable: $t_{(34)} = 2.81$, $P = 0.008$; 3rd syllable:

$t_{(34)} = 1.94$, $P = 0.06$). There was no significant difference between the ITD and IID conditions for the two

staggered syllables (2nd syllable: $t_{(34)} = 1.41$, $P = 0.17$; 3rd syllable: $t_{(34)} = 1.39$, $P = 0.17$).

Figure 2

Fig. 2. *Listeners' (N = 18) recall performance was evaluated for every syllable and different*

spatialization conditions. Sounds were spatialized either based on generic ITDs,

individualized IIDs, or the natural combination of ITDs, IIDs, and spectral cues in

individualized HRTFs. Behavioral advantages of having more consistent spatial information

*were significant but small. * P < .05; ** P < .001; *** P < .0001*

3.2 Impoverished spatial cues affect attentional modulation of ERPs

Figure 3A shows the ERPs evoked by the onset of the second syllable of the attended target sound and the

unattended distractor sound, aligning the onsets of the target and distractor syllables to 0 s to allow direct

comparison. Stimulus onsets elicited a fronto-central positivity (P1) between 50 to 100 ms followed by a

negativity (N1) between 100 to 180 ms (Figure 3A-B). The amplitudes of these two components were

extracted and the difference between attended stimuli (target sound) and unattended stimuli (distractor

sound) was calculated in order to quantify attentional modulation for both the P1 and N1 components

(Figure 3C).

Figure 3

A.

B.

C.

Fig. 3. *P1 amplitudes were only modulated by the direction of attention in the HRTF condition,*

whereas N1 amplitudes were modulated equivalently across spatialization conditions (N = 18).

A. *ERP waveforms at fronto-central electrodes were compared between the attended target stream*

and the unattended distractor stream for every spatialization condition. The P1 time range was

defined between 50 ms and 100 ms, and the N1 time range was defined between 100 ms and 180 ms.

B. *Most topographies of both ERP components show maxima at the fronto-central sites (black dots)*

used for evaluation. C. The modulation strength of ERP components was assessed by the amplitude

*differences between attended and unattended streams. * P < .05; ** P < .01*

We tested whether P1 responses were significantly larger to attended stimuli than to unattended stimuli in

each of the three conditions. Only the HRTF condition showed a significant P1 modulation ($t_{(17)}$ = 3.12,

P_{adj} = 0.017); no significant attentional modulation was found in either the ITD ($t_{(17)}$ = 0.50, P_{adj} = 1) or

IID conditions ($t_{(17)}$ = 0.06, P_{adj} = 1). Across conditions we found a statistically significant main effect of

spatial cue on P1 amplitude modulation ($F_{(2,34)}$ = 3.34, P = 0.047). Attentional modulation was

significantly larger in the HRTF condition than in the ITD ($t_{(34)}$ = 2.38, P = 0.023) and IID conditions

($t_{(34)}$ = 2.07, P = 0.046); however, modulation did not differ significantly between the ITD and IID

conditions ($t_{(34)}$ = 0.31, P = 0.76) (Figure 3C).

In all three spatialization conditions, the N1 amplitude was modulated significantly by spatial attention,

that is, attended sounds evoked larger N1 amplitudes than unattended sounds (ITD: $t_{(17)}$ = 3.01,

P_{adj} = 0.024; IID: $t_{(17)}$ = 4.12, P_{adj} = 0.002; HRTF: $t_{(17)}$ = 3.56, P_{adj} = 0.007). Across the three spatialization

conditions the magnitude of N1 modulation did not differ significantly ($F_{(2,34)}$ = 0.060, P = 0.94;

Figure 3C).

3.3 Alpha oscillation power shows less attentional modulation with impoverished spatial cues

To investigate the effect of spatialization on attentional control, we analyzed the power in alpha

oscillations during the attentional preparation period (-800 ms to 0 ms), a time period in which listeners

knew where to orient spatial attention based on the preceding acoustic cue, but before the sound mixture

of competing streams began. We averaged the power in alpha across all trials for each spatialization

condition, regardless of where spatial attention was focused, to get a measure of the total engagement of

alpha activity. We then compared relative power for different attentional directions. On average across

directions of attentional focus, we calculated the time courses of alpha global field power (GFP, Figure

4A) and compared within-subject differences of the temporal average within the preparatory time period

across spatialization conditions (Figure 4B). Alpha GFP was not significantly modulated in either the ITD

or ILD conditions (ITD: $t_{(17)} = 0.44$, $P_{adj} = 1$; ILD: $t_{(17)} = 0.43$, $P_{adj} = 1$), while in the HRTF condition, the

GFP tended to be greater than zero (HRTF: $t_{(17)} = 2.56$, $P_{adj} = 0.061$). In a direct comparison,

spatialization conditions differed significantly in alpha GFP ($F_{(2,34)} = 5.26$, $P = 0.010$). In particular, alpha

GFP in the HRTF condition was significantly larger than in the other two conditions (HRTF vs ITD:

$t_{(34)} = 2.80$, $P = 0.008$; HRTF vs IID: $t_{(34)} = 2.82$, $P = 0.008$). No significant difference was found between

the ITD and IID conditions ($t_{(34)} = 0.019$, $P = 0.99$).

Figure 4

Fig. 4. *Within-subject differences in alpha-band GFP are larger in the HRTF condition, especially*

during the preparatory time window (after the sound cue but before the first syllables of the

competing streams). *A.* *Waveforms of the average (± SEM) GFP differences are shown during the*

baseline period, preparatory phase, and stimulus phase with stream competition. *B.* *The temporal*

average of the preparatory alpha GFP difference is larger for the HRTF condition. *** P < .01*

We next assessed the lateralization of alpha power with the spatial focus of attention by comparing AMI

differences across hemispheres (Figure 5). In general, the scalp topographies of AMIs show the expected

hemispheric differences. However, statistically significant hemispheric differences were found only in the

296   HRTF condition ($t_{(17)}$ = 3.09, P_{adj} = 0.020), not in the ITD ($t_{(17)}$ = 1.29, P_{adj} = 0.64) and the IID condition

297   ($t_{(17)}$ = 0.15, P_{adj} = 1). A direct comparison of these hemisphere differences across conditions revealed a

298   trend in which the HRTF condition had larger differences in AMI across hemispheres ($F_{(2,34)}$ = 2.98,

299   P = 0.064).

Figure 5

Fig. 5. *Attentional modulation of alpha activity was lateralized to the hemisphere ipsilateral to the*

target stream only in the HRTF condition. AMI topographies and hemispheric averages are shown

*for every spatialization condition (N = 18). * P < .05*

In summary, impoverished spatial cues lead to worse behavioral performance, smaller P1 modulation,

reduced modulation of preparatory alpha power GFP, and reduced lateralization of alpha power with

attentional focus, confirming our hypothesis that impoverished spatial cues impaired engagement of

spatial attention.

3.4 Relationships between Attentional Modulation Metrics

Given all these consistent effects of modulation metrics, we explored, post hoc, whether there were

ordered relationships in the individual measures of performance and neural signatures of attentional

control, including P1 modulation, preparatory alpha GFP, and alpha power lateralization. To investigate

the relationship between evoked response modulation and alpha oscillatory activities, we first calculated

the regression slope relating P1 amplitude to preparatory alpha GFP for each subject, and then performed

a paired t-test on the coefficients obtained. No consistent relationship between alpha GFP and P1

amplitudes was observed ($t_{(17)} = 0.90$, $P = 0.38$). Correlation analysis was also conducted comparing

behavioral accuracy to P1 modulation, defined as the attended P1 amplitude minus unattended P1

amplitude. No consistent relationships between P1 modulation and behavioral performance were observed

for any syllable (1st syllable: $t_{(17)} = 0.54$, $P = 0.59$; 2nd syllable: $t_{(17)} = 0.31$, $P = 0.76$; 3rd syllable:

$t_{(17)} = 0.69$, $P = 0.50$). Similarly, we did not observe consistent relationships between alpha AMI

lateralization and response accuracy for any syllable (1st syllable: $t_{(17)} = 0.19$, $P = 0.85$; 2nd syllable:

$t_{(17)} = 1.39$, $P = 0.18$; 3rd syllable: $t_{(17)} = 0.11$, $P = 0.91$). Thus, although there were significant

differences in engagement of attention across spatial conditions as measured both behaviorally and

neurally, the individual subject differences in these metrics were not closely related.

4 Discussion

Behaviorally, we found that impoverished spatial cues impair performance on an auditory spatial attention task in a multi-talker scene. We used objective electrophysiological measures to assess whether the naturalness and richness of spatial cues also impacts how strongly auditory spatial attention modulates brain responses. We found that impoverished spatial cues reduce the strength of the evoked and induced neural signatures of attentional control. Specifically, evoked P1 amplitudes and induced alpha oscillatory power showed less attentional modulation for sound stimuli with impoverished spatial cues compared to when spatial cues were tailored to recreate the natural, rich experience of individual listeners.

4.1 Impoverished spatial cues result in less neural modulation during selective attention

We investigated attentional modulation of four established neural signatures of selective attention: evoked P1 and N1 amplitudes and induced power and lateralization of alpha oscillation. While attentional modulation of N1 amplitude was observed in all conditions, attentional modulation of the earlier P1 amplitude was not observed or was significantly weaker in the impoverished cue conditions compared to the natural cue condition. Similarly, we found less preparatory alpha power activity in the impoverished spatial cue conditions than in the natural cue condition, reflected by two indexes quantifying the amount of spatial variability of alpha power: alpha GFP (Figure 4) and AMI (Figure 5). In the ITD and IID conditions, although there was a hint of preparatory alpha lateralization over parietal sensors, the amount of lateralization was significantly smaller than in the HRTF condition and did not reach statistical significance. Preparatory alpha activity during spatial attention tasks has been well documented to form a specific lateralization pattern in both vision and audition (Banerjee et al., 2011; Kelly, 2006; Sauseng et al., 2005; Worden et al., 2018), which is thought to be evidence of a preparatory information-gating mechanism (Foxe and Snyder, 2011; Jensen and Mazaheri, 2010; Klimesch, 2012; Klimesch et al., 2007). In vision, alpha lateralization has been observed to increase with the laterality of attention focus (Rihs et al., 2007; Samaha et al., 2015), reflecting an inhibition pattern topographically specific to attention focus. Moreover, evidence for active top-down control of the phase of alpha oscillation during visual spatial

attention suggests that alpha oscillatory activity represents active engagement and disengagement of the

attentional network (Samaha et al., 2016). In addition, a previous somatosensory study has revealed that

the alpha lateralization is positively correlated to pre-stimulus cue reliability, further suggesting that alpha

lateralization reflects the top-down control in order to optimize the processing of upcoming stimuli

(Haegens et al., 2011). Although relatively few studies have investigated alpha activity in audition,

studies suggest that alpha control mechanisms are supra-modal rather than sensory specific (Banerjee et

al., 2011). In the current experiment, a pre-stimulus auditory cue directed listeners where to focus

attention in an upcoming sound mixture. The cue was spatialized using the same auditory features used to

spatialize the stream mixture. Our results thus suggest that compared to stimuli with natural spatial cues,

stimuli featuring only ITDs or only IIDs are less reliable in directing attentional focus, producing weaker

engagement of spatial attention and reduced attentional modulation of neural responses.

Consistent with the idea that impoverished spatial cues lead to weaker engagement of spatial attention, we

found that the P1 ERP component was modulated by attention only with natural spatial cues, not with

impoverished cues; this result is consistent with a weak spatial representation failing to engage attentional

modulation of early sensory responses (Figure 3). Our finding that attentional focus leads to a modulation

of P1 amplitude for natural spatial cues is consistent with previous reports of effects of attention on the P1

amplitude observed in previous spatial attention studies across sensory modalities [auditory: (Giuliano et

al., 2014); visual: (Hillyard and Anllo-Vento, 1998; Hopfinger et al., 2004)]. Past studies agree that P1

modulation reflects an early sensory inhibition mechanism related to suppression of task-irrelevant

stimuli. Although debates remain as to whether P1 modulation results from bottom-up sensory gain

control (Hillyard and Anllo-Vento, 1998; Luck, 1995; Slagter et al., 2016) or for some top-down

inhibitory process (Freunberger et al., 2008; Klimesch, 2011), it is generally accepted in visual spatial

studies that greater P1 amplitude modulation is associated with greater inhibition of to-be-ignored stimuli

(Couperus and Mangun, 2010; Hillyard and Anllo-Vento, 1998; Klimesch, 2012). Interestingly,

attentional modulation of auditory P1 has been found to be positively correlated to visual working

memory capacity, a result that was used to suggest that stronger P1 modulation reflects better attentional

control of the flow of sensory information into working memory (Fukuda and Vogel, 2009; Giuliano et

al., 2014). Our result is consistent with the hypothesis that P1 modulation directly reflects attentional

control. Specifically, impoverished spatial cues likely produce a "muddy" representation of auditory

space that supports only imprecise, poorly focused top-down spatial attention. The resulting lack of

control and specificity of spatial auditory attention results in early P1 responses that are unmodulated by

attentional focus.

N1 modulation is well documented as a neural index of attentional control (Choi et al., 2013; Hillyard et

al., 1998; Stevens et al., 2008; Wyart et al., 2012). The attentional modulation of N1 is thought to reflect

attentional facilitation rather than inhibition (Couperus and Mangun, 2010; Marzecová et al., 2018;

Slagter et al., 2016). In contrast to preparatory alpha and P1, we found that the later N1 evoked response

was modulated similarly, regardless of the richness and naturalness of spatial cues.

Due to the robustness and relatively large amount of modulation, changes in auditory N1 amplitude have

been used as a biomarker and a primary feature for classification of attentional focus (Blankertz et al.,

2011; Schreuder et al., 2011); see also recent work on decoding attentional focus for running speech using

the correlation between neural responses and the power envelope of the speech streams: (Chait et al.,

2010; Mesgarani and Chang, 2012; Rimmele et al., 2015). However, there is little known about how N1

amplitudes reflects the processing of different spatial cues during auditory spatial attention. Previous

studies have revealed different N1 topographies during ITD and IID processing, leading to the conclusion

that ITD and IID are processed by different neural populations in the auditory cortex (Johnson and

Hautus, 2010; Tardif et al., 2006; Ungan et al., 2001). However, debates remain about whether this

difference in topography depends on perceived laterality, instead of different neural populations

specialized for processing different spatial cues. Results from a more recent study show that auditory N1

modulation does not differ across spatial cue conditions, indicating integrated processing of sound

locations in auditory cortex regardless of cues (Salminen et al., 2015). In the current study, N1

modulation did not differ across the three spatialization conditions. Thus, our results support the idea that

the same cortical neural population is responsible for processing different binaural spatial cues.

4.2 Behavioral disadvantages associated with impoverished spatial cues are modest and depend

on sound stimulus characteristics

Despite the influence of spatial cue richness on neural metrics, our behavioral results showed only small

(albeit significant) behavioral differences between impoverished spatial cues and natural, individualized

spatial cues (Figure 2). In line with previous studies that observed greater spatial release from masking

with combined spatial cues compared to with isolated cues (Culling et al., 2004; Ellinger et al., 2017),

accuracy was best in the HRTF condition. The small accuracy improvement over using impoverished

cues is seen consistently across subjects. In the first syllable where the target and distractor streams

overlap in time, the HRTF condition yielded a 13% increase in accuracy over the ITD condition, but is

comparable to performance in the IID condition. In the two staggered syllables, accuracy in the HRTF

condition is greater than in the ITD and IID conditions by only about 6% and 1%, respectively. These

differences in behavioral performance across syllables suggest that the characteristics of sound stimuli

influence the difficulty of the task and may influence the behavioral advantages of having richer, more

robust spatial cues (Kidd et al., 2010). Concordantly, a previous study with complex tone stimuli has

shown much larger differences in behavioral performance of up to 20% (Schröger, 1996), whereas studies

presenting speech stimuli in a multi-talker environment found no behavioral advantage of having

combined cues compared to impoverished cues (Glyde et al., 2013). These behavioral discrepancies, in

combination with our neural findings, indicate that behavioral performance alone is not a sensitive metric

for determining whether cortical networks controlling spatial selective attention are fully engaged.

Conclusions

Our results indicate that although impoverished spatial cues can support spatial segregation of speech in a multi-talker environment, they do not fully engage the brain networks controlling spatial attention and lead to weak attentional control. Previous auditory studies have provided evidence that impoverished spatial cues do not evoke the same neural processing mechanisms as natural cue combinations during localization tasks with single sounds (Callan et al., 2013; Getzmann and Lewald, 2010; Leino et al., 2007; Palomäki et al., 2005). The current study extends these findings, demonstrating that the efficacy of higher-level cognitive processing, such as deployment of auditory selective attention, also depends on the naturalness of spatial cues. Poor attentional control was reflected in limited modulation of neural biomarkers of attentional processes. These findings suggest that the many past auditory attention studies using impoverished spatial cues may have underestimated the robust changes in cortical activity associated with deployment of spatial auditory attention in natural settings. Although impoverished auditory spatial cues can allow listeners to deploy spatial attention effectively enough to perform well in simple acoustic scenes, noisy, complex listening environments like those encountered in everyday environments pose greater challenges to attentional processing. In natural settings, spatial attention may fail unless attentional control networks are fully engaged. Thus, these results demonstrate the importance of preserving rich, natural spatial cues in hearing aids and other assistive listening devices.

Acknowledgements

We thank Ashvini Meltoke for collecting part of the data, Gerald Kidd Jr. for providing the facilities for individualized HRTF measurements, and Virginia Best for fruitful discussions. This work was supported by the Austrian Science Fund (grant J3803-N30 to R.B.) and NICD (grant DC013825 to B.S.-C.).

References

Ahveninen, J., Hämäläinen, M., Jääskeläinen, I.P., Ahlfors, S.P., Huang, S., Lin, F.-H., Raij, T., Sams, M., Vasios, C.E., Belliveau, J.W., 2011. Attention-driven auditory cortex short-term plasticity helps

segregate relevant sounds from noise. Proceedings of the National Academy of Sciences 108, 4182–4187.
https://doi.org/10.1073/pnas.1016134108

Alho, K., Medvedev, S.V., Pakhomov, S.V., Roudas, M.S., Tervaniemi, M., Reinikainen, K., Zeffiro, T.,
Näätänen, R., 1999a. Selective tuning of the left and right auditory cortices during spatially directed
attention. Cognitive Brain Research 7, 335–341. https://doi.org/10.1016/S0926-6410(98)00036-6

Banerjee, S., Snyder, A.C., Molholm, S., Foxe, J.J., 2011. Oscillatory Alpha-Band Mechanisms and the
Deployment of Spatial Attention to Anticipated Auditory and Visual Target Locations: Supramodal or
Sensory-Specific Control Mechanisms? Journal of Neuroscience 31, 9923–9932.
https://doi.org/10.1523/jneurosci.4660-10.2011

Baumgartner, R., Reed, D.K., Tóth, B., Best, V., Majdak, P., Colburn, H.S., Shinn-Cunningham, B.,
2017. Asymmetries in behavioral and neural responses to spectral cues demonstrate the generality of
auditory looming bias. Proceedings of the National Academy of Sciences 114, 9743–9748.
https://doi.org/10.1073/pnas.1703247114

Blankertz, B., Lemm, S., Treder, M., Haufe, S., Müller, K.-R., 2011. Single-trial analysis and
classification of ERP components — A tutorial. NeuroImage 56, 814–825.
https://doi.org/10.1016/J.NEUROIMAGE.2010.06.048

Blauert, J., 1997b. Spatial hearing. The Psychophysics of Human Sound Localization, 2nd edition. ed.
MIT-Press, Cambridge, MA.

Boersma, P., 2001. Praat, a system for doing phonetics by computer. Glot International 5, 341–345.

Brainard, D.H., 1997. The psychophysics toolbox. Spatial vision 10, 433–436.

Callan, A., Callan, D.E., Ando, H., 2013. Neural correlates of sound externalization. NeuroImage 66, 22–
27. https://doi.org/10.1016/j.neuroimage.2012.10.057

Chait, M., de Cheveigné, A., Poeppel, D., Simon, J.Z., 2010. Neural dynamics of attending and ignoring
in human auditory cortex. Neuropsychologia 48, 3262–3271.
https://doi.org/10.1016/J.NEUROPSYCHOLOGIA.2010.07.007

Cherry, E.C., 1953. Some Experiments on the Recognition of Speech, with One and with Two Ears. The
Journal of the Acoustical Society of America 25, 975–979. https://doi.org/10.1121/1.1907229

Choi, I., Rajaram, S., Varghese, L.A., Shinn-Cunningham, B.G., 2013. Quantifying attentional
modulation of auditory-evoked cortical responses from single-trial electroencephalography. Frontiers in
Human Neuroscience 7, 115. https://doi.org/10.3389/fnhum.2013.00115

Couperus, J.W., Mangun, G.R., 2010. Signal enhancement and suppression during visual–spatial selective
attention. Brain Research 1359, 155–177. https://doi.org/10.1016/j.brainres.2010.08.076

Cubick, J., Buchholz, J.M., Best, V., Lavandier, M., Dau, T., 2018. Listening through hearing aids affects
spatial perception and speech intelligibility in normal-hearing listeners. The Journal of the Acoustical
Society of America 144, 2896–2905. https://doi.org/10.1121/1.5078582

Culling, J.F., Hawley, M.L., Litovsky, R.Y., 2004. The role of head-induced interaural time and level
differences in the speech reception threshold for multiple interfering sound sources. The Journal of the
Acoustical Society of America 116, 1057–1065. https://doi.org/10.1121/1.1772396

Cusack, R., Carlyon, R.P., Robertson, I.H., 2001. Auditory Midline and Spatial Discrimination in Patients
with Unilateral Neglect. Cortex 37, 706–709. https://doi.org/10.1016/S0010-9452(08)70620-8

Dahmen, J.C., Keating, P., Nodal, F.R., Schulz, A.L., King, A.J., 2010. Adaptation to Stimulus Statistics
in the Perception and Neural Representation of Auditory Space. Neuron 66, 937–948.
https://doi.org/10.1016/J.NEURON.2010.05.018

Dai, L., Best, V., Shinn-Cunningham, B.G., 2018. Sensorineural hearing loss degrades behavioral and
physiological measures of human spatial selective auditory attention. Proceedings of the National
Academy of Sciences 201721226. https://doi.org/10.1073/pnas.1721226115

Das, N., Biesmans, W., Bertrand, A., Francart, T., 2016. The effect of head-related filtering and ear-
specific decoding bias on auditory attention detection. Journal of Neural Engineering 13, 056014.
https://doi.org/10.1088/1741-2560/13/5/056014

Delorme, A., Makeig, S., 2004. EEGLAB: an open source toolbox for analysis of single-trial EEG
dynamics including independent component analysis. Journal of Neuroscience Methods 134, 9–21.
https://doi.org/10.1016/j.jneumeth.2003.10.009

Delorme, A., Sejnowski, T., Makeig, S., 2007. Enhanced detection of artifacts in EEG data using higher-
order statistics and independent component analysis. NeuroImage 34, 1443–1449.
https://doi.org/10.1016/j.neuroimage.2006.11.004

Ellinger, R.L., Jakien, K.M., Gallun, F.J., 2017. The role of interaural differences on speech intelligibility
in complex multi-talker environmentsa). The Journal of the Acoustical Society of America 141, EL170–
EL176. https://doi.org/10.1121/1.4976113

Foxe, J.J., Snyder, A.C., 2011. The Role of Alpha-Band Brain Oscillations as a Sensory Suppression
Mechanism during Selective Attention. Frontiers in Psychology 2, 154.
https://doi.org/10.3389/fpsyg.2011.00154

Freunberger, R., Höller, Y., Griesmayr, B., Gruber, W., Sauseng, P., Klimesch, W., 2008. Functional
similarities between the P1 component and alpha oscillations. European Journal of Neuroscience 27,
2330–2340. https://doi.org/10.1111/j.1460-9568.2008.06190.x

Fukuda, K., Vogel, E.K., 2009. Human Variation in Overriding Attentional Capture. Journal of
Neuroscience 29, 8726–8733. https://doi.org/10.1523/jneurosci.2145-09.2009

Getzmann, S., Lewald, J., 2010. Effects of natural versus artificial spatial cues on electrophysiological
correlates of auditory motion. Hearing Research 259, 44–54. https://doi.org/10.1016/j.heares.2009.09.021

Giuliano, R.J., Karns, C.M., Neville, H.J., Hillyard, S.A., 2014. Early Auditory Evoked Potential Is
Modulated by Selective Attention and Related to Individual Differences in Visual Working Memory
Capacity. Journal of Cognitive Neuroscience. https://doi.org/10.1162/jocn_a_00684

Glyde, H., Buchholz, J.M., Dillon, H., Cameron, S., Hickson, L., 2013. The importance of interaural time
differences and level differences in spatial release from masking. The Journal of the Acoustical Society of
America 134, EL147–EL152. https://doi.org/10.1121/1.4812441

Haegens, S., Handel, B.F., Jensen, O., 2011. Top-Down Controlled Alpha Band Activity in
Somatosensory Areas Determines Behavioral Performance in a Discrimination Task. Journal of
Neuroscience 31, 5197–5204. https://doi.org/10.1523/jneurosci.5199-10.2011

Hartmann, W.M., Wittenberg, A., 1996. On the externalization of sound images. J Acoust Soc Am 99,
3678–88.

Herrmann, C.S., Knight, R.T., 2001. Mechanisms of human attention: event-related potentials and
oscillations. Neuroscience & Biobehavioral Reviews 25, 465–476. https://doi.org/10.1016/S0149-
7634(01)00027-6

Higgins, N.C., McLaughlin, S.A., Rinne, T., Stecker, G.C., 2017. Evidence for cue-independent spatial
representation in the human auditory cortex during active listening. Proceedings of the National Academy
of Sciences 114, E7602--E7611. https://doi.org/10.1073/pnas.1707522114

Hillyard, S.A., Anllo-Vento, L., 1998. Event-related brain potentials in the study of visual selective
attention. Proceedings of the National Academy of Sciences 95, 781–787.
https://doi.org/10.1073/pnas.95.3.781

Hillyard, S.A., Vogel, E.K., Luck, S.J., 1998. Sensory gain control (amplification) as a mechanism of
selective attention: electrophysiological and neuroimaging evidence. Philosophical transactions of the
Royal Society of London. Series B, Biological sciences 353, 1257–70.
https://doi.org/10.1098/rstb.1998.0281

Hopfinger, J., Luck, S., Hillyard, S., 2004. Selective attention: Electrophysiological and neuromagnetic
studies. The cognitive neurosciences III 561–574.

Itoh, K., Yumoto, M., Uno, A., Kurauchi, T., Kaga, K., 2000. Temporal stream of cortical representation
for auditory spatial localization in human hemispheres. Neuroscience Letters 292, 215–219.
https://doi.org/10.1016/S0304-3940(00)01465-8

Jensen, O., Mazaheri, A., 2010. Shaping Functional Architecture by Oscillatory Alpha Activity: Gating
by Inhibition. Frontiers in Human Neuroscience 4, 186. https://doi.org/10.3389/fnhum.2010.00186

Johnson, B.W., Hautus, M.J., 2010. Processing of binaural spatial information in human auditory cortex:
Neuromagnetic responses to interaural timing and level differences. Neuropsychologia 48, 2610–2619.
https://doi.org/10.1016/J.NEUROPSYCHOLOGIA.2010.05.008

Kelly, S.P., 2006. Increases in Alpha Oscillatory Power Reflect an Active Retinotopic Mechanism for
Distracter Suppression During Sustained Visuospatial Attention. Journal of Neurophysiology 95, 3844–
3851. https://doi.org/10.1152/jn.01234.2005

Kidd, G., Mason, C.R., Best, V., Marrone, N., Marrone, N., 2010. Stimulus factors influencing spatial
release from speech-on-speech masking. The Journal of the Acoustical Society of America 128, 1965–78.
https://doi.org/10.1121/1.3478781

Klimesch, W., 2012. Alpha-band oscillations, attention, and controlled access to stored information.
Trends in Cognitive Sciences 16, 606–617. https://doi.org/10.1016/J.TICS.2012.10.007

Klimesch, W., 2011. Evoked alpha and early access to the knowledge system: The P1 inhibition timing
hypothesis. Brain Research. https://doi.org/10.1016/j.brainres.2011.06.003

Klimesch, W., Sauseng, P., Hanslmayr, S., 2007. EEG alpha oscillations: The inhibition-timing
hypothesis. Brain Research Reviews 53, 63–88. https://doi.org/10.1016/j.brainresrev.2006.06.003

Kong, L., Michalka, S.W., Rosen, M.L., Sheremata, S.L., Swisher, J.D., Shinn-Cunningham, B.G.,
Somers, D.C., 2014. Auditory Spatial Attention Representations in the Human Cerebral Cortex. Cerebral

Cortex 24, 773–784. https://doi.org/10.1093/cercor/bhs359

Leino, S., May, P.J.C., Alku, P., Liikkanen, L.A., Tiitinen, H., 2007. The contribution of high frequencies to human brain activity underlying horizontal localization of natural spatial sounds. BMC neuroscience 8, 78. https://doi.org/10.1186/1471-2202-8-78

Lim, S.-J., Wöstmann, M., Obleser, J., 2015. Selective Attention to Auditory Memory Neurally Enhances Perceptual Precision. Journal of Neuroscience 35, 16094–16104. https://doi.org/10.1523/JNEUROSCI.2674-15.2015

Loiselle, L.H., Dorman, M.F., Yost, W.A., Cook, S.J., Gifford, R.H., 2016. Using ILD or ITD Cues for Sound Source Localization and Speech Understanding in a Complex Listening Environment by Listeners With Bilateral and With Hearing-Preservation Cochlear Implants. Journal of speech, language, and hearing research : JSLHR 59, 810–8. https://doi.org/10.1044/2015_JSLHR-H-14-0355

Luck, S.J., 1995. Multiple mechanisms of visual-spatial attention: recent evidence from human electrophysiology. Behavioural Brain Research 71, 113–123. https://doi.org/10.1016/0166-4328(95)00041-0

Majdak, P., Baumgartner, R., Jenny, C., 2019. Formation of three-dimensional auditory space. arXiv:1901.03990 [q-bio].

Marzecová, A., Schettino, A., Widmann, A., SanMiguel, I., Kotz, S.A., Schröger, E., 2018. Attentional gain is modulated by probabilistic feature expectations in a spatial cueing task: ERP evidence. Scientific Reports 8. https://doi.org/10.1038/s41598-017-18347-1

Mesgarani, N., Chang, E.F., 2012. Selective cortical representation of attended speaker in multi-talker speech perception. Nature 485, 233–236. https://doi.org/10.1038/nature11020

Middlebrooks, J.C., 2015. Sound localization. Handbook of Clinical Neurology, The Human Auditory System 129, 99–116. https://doi.org/10.1016/B978-0-444-62630-1.00006-8

Murray, M.M., Brunet, D., Michel, C.M., 2008. Topographic ERP Analyses: A Step-by-Step Tutorial Review. Brain Topography 20, 249–264. https://doi.org/10.1007/s10548-008-0054-5

Palomäki, K.J., Tiitinen, H., Mäkinen, V., May, P.J.C., Alku, P., 2005. Spatial processing in human auditory cortex: The effects of 3D, ITD, and ILD stimulation techniques. Cognitive Brain Research 24, 364–379. https://doi.org/10.1016/j.cogbrainres.2005.02.013

Posner, M.I., Snyder, C.R., Davidson, B.J., 1980. Attention and the detection of signals. Journal of Experimental Psychology: General 109, 160–174. https://doi.org/10.1037/0096-3445.109.2.160

Rihs, T.A., Michel, C.M., Thut, G., 2007. Mechanisms of selective inhibition in visual spatial attention are indexed by ?-band EEG synchronization. European Journal of Neuroscience 25, 603–610. https://doi.org/10.1111/j.1460-9568.2007.05278.x

Rimmele, J.M., Zion Golumbic, E., Schröger, E., Poeppel, D., 2015. The effects of selective attention and speech acoustics on neural speech-tracking in a multi-talker scene. Cortex 68, 144–154. https://doi.org/10.1016/J.CORTEX.2014.12.014

Röder, B., Teder-Sälejärvi, W., Sterr, A., Rösler, F., Hillyard, S.A., Neville, H.J., 1999. Improved auditory spatial tuning in blind humans. Nature 400, 162–166. https://doi.org/10.1038/22106

Sach, A.J., Hill, N.I., Bailey, P.J., 2000. Auditory spatial attention using interaural time differences.
Journal of Experimental Psychology: Human Perception and Performance 26, 717–729.
https://doi.org/10.1037/0096-1523.26.2.717

Salminen, N.H., Takanen, M., Santala, O., Lamminsalo, J., Altoè, A., Pulkki, V., 2015. Integrated
processing of spatial cues in human auditory cortex. Hearing Research 327, 143–152.
https://doi.org/10.1016/J.HEARES.2015.06.006

Samaha, J., Bauer, P., Cimaroli, S., Postle, B.R., 2015. Top-down control of the phase of alpha-band
oscillations as a mechanism for temporal prediction. Proceedings of the National Academy of Sciences of
the United States of America 112, 8439–8444. https://doi.org/10.1073/pnas.1503686112

Samaha, J., Sprague, T.C., Postle, B.R., 2016. Decoding and Reconstructing the Focus of Spatial
Attention from the Topography of Alpha-band Oscillations. Journal of Cognitive Neuroscience 28, 1090–
1097. https://doi.org/10.1162/jocn_a_00955

Sauseng, P., Klimesch, W., Doppelmayr, M., Pecherstorfer, T., Freunberger, R., Hanslmayr, S., 2005.
EEG alpha synchronization and functional coupling during top-down processing in a working memory
task. Human Brain Mapping 26, 148–155. https://doi.org/10.1002/hbm.20150

Schreuder, M., Rost, T., Tangermann, M., 2011. Listen, You are Writing! Speeding up Online Spelling
with a Dynamic Auditory BCI. Frontiers in Neuroscience 5, 112.
https://doi.org/10.3389/fnins.2011.00112

Schröger, E., 1996. Interaural time and level differences: Integrated or separated processing? Hearing
Research 96, 191–198. https://doi.org/10.1016/0378-5955(96)00066-4

Shinn-Cunningham, B.G., Kopco, N., Martin, T.J., 2005. Localizing nearby sound sources in a classroom:
Binaural room impulse responses. The Journal of the Acoustical Society of America 117, 3100–3115.
https://doi.org/10.1121/1.1872572

Siegel, M., Donner, T.H., Engel, A.K., 2012. Spectral fingerprints of large-scale neuronal interactions.
Nature Reviews Neuroscience 13, 121–134. https://doi.org/10.1038/nrn3137

Skrandies, W., 1990. Global field power and topographic similarity. Brain Topography 3, 137–141.

Snyder, J.S., Large, E.W., 2005. Gamma-band activity reflects the metric structure of rhythmic tone
sequences. Cognitive Brain Research 24, 117–126. https://doi.org/10.1016/j.cogbrainres.2004.12.014

Stevens, C., Fanning, J., Coch, D., Sanders, L., Neville, H., 2008. Neural mechanisms of selective
auditory attention are enhanced by computerized training: Electrophysiological evidence from language-
impaired and typically developing children. Brain Research 1205, 55–69.
https://doi.org/10.1016/J.BRAINRES.2007.10.108

Tardif, E., Murray, M.M., Meylan, R., Spierer, L., Clarke, S., 2006. The spatio-temporal brain dynamics
of processing and integrating sound localization cues in humans. Brain Research 1092, 161–176.
https://doi.org/10.1016/J.BRAINRES.2006.03.095

Ungan, P., Yagcioglu, S., Goksoy, C., 2001. Differences between the N1 waves of the responses to
interaural time and intensity disparities: scalp topography and dipole sources. Clinical Neurophysiology
112, 485–498. https://doi.org/10.1016/S1388-2457(00)00550-2

Worden, M.S., Foxe, J.J., Wang, N., Simpson, G. V., 2018. Anticipatory Biasing of Visuospatial

Attention Indexed by Retinotopically Specific α-Bank Electroencephalography Increases over Occipital
Cortex. The Journal of Neuroscience 20, RC63–RC63. https://doi.org/10.1523/jneurosci.20-06-
j0002.2000

Wöstmann, M., Herrmann, B., Maess, B., Obleser, J., 2016. Spatiotemporal dynamics of auditory
attention synchronize with speech. Proceedings of the National Academy of Sciences 113, 3873–3878.
https://doi.org/10.1073/pnas.1523357113

Wyart, V., Dehaene, S., Tallon-Baudry, C., 2012. Early dissociation between neural signatures of
endogenous spatial attention and perceptual awareness during visual masking. Frontiers in Human
Neuroscience 6, 16. https://doi.org/10.3389/fnhum.2012.00016