Extinction-resistant attention to long-term conditioned threat is indexed by selective

visuocortical alpha suppression in humans

Christian Panitz[a] (christian.panitz@staff.uni-marburg.de),

Andreas Keil[b] (akeil@ufl.edu),

&

Erik M. Mueller[a] (erik.mueller@staff.uni-marburg.de)

[a]University of Marburg, Department of Psychology, Gutenbergstr. 18, 35032 Marburg, Germany

[b]University of Florida, Center for the Study of Emotion and Attention, 3063 Longleaf Road,

Gainesville, FL 32608, USA

This study was supported by a grant of the German Research Foundation to Erik M. Mueller

[DFG MU3535/2-1]. None of the authors has financial interests or other potential conflicts of interest.

Correspondence concerning this article should be addressed to Christian Panitz, Department of

Psychology, University of Marburg, Gutenbergstr. 18, 35032 Marburg, Germany.

E-mail: christian.panitz@staff.uni-marburg.de

Abstract

While ERP studies have shown heightened early visual attention to conditioned threat, it is unknown whether this attentional prioritization is sustained throughout later processing stages and whether it is robust to extinction. To investigate sustained visual attention, we assessed visuocortical alpha suppression in response to conditioned and extinguished threat. $N = 87$ participants underwent a two-day threat conditioning paradigm with acquisition and extinction on one day and a critical recall test one day later. EEG time-frequency analyses revealed that, on Day 2, threat-conditioned vs. safety cues evoked stronger occipital alpha power suppression from 600 to 1200 ms. Notably, this suppression was resistant to extinction. The present study showed for the first time that threat conditioning enhances sustained modulation of visuocortical attention to threat in the long term. The long-term stability and extinction resistance of alpha suppression suggests a crucial role of visuocortical attention mechanisms in the maintenance of learned fears.

Keywords: threat; long-term conditioning; long-term extinction; alpha; visual attention

Introduction

Learning to predict threat and safety based on environmental cues is fundamental for adaptive

behavior. A well-established paradigm for elucidating mechanisms of associative cue-outcome

learning in the laboratory is classical threat conditioning (also referred to as fear conditioning or

aversive conditioning) where conditioned stimuli are threat cues (CS+) when they signal an upcoming

aversive event (i.e., an unconditioned stimulus; US) and safety cues (CS-) when they signal the

absence of an aversive event (LeDoux, 2014; Lonsdorf et al., 2017; Rescorla & Wagner, 1972). In

threat conditioning, participants acquire a conditioned defensive response in response to the CS+,

which is expected to diminish again upon extinction training, i.e., repeated presentation of CS+

without US (Myers & Davis, 2007; Quirk & Mueller, 2008).

Importantly, both conditioning and extinction memories need to be consolidated, retained, and

recalled in future situations to allow adaptive responding in the long term. Successful short-term

learning – i.e., acquisition and extinction of a conditioned response within an experimental session – is

considered necessary but not sufficient for successful long-term learning. Such long-term learning

effects are indicated by the successful recall of previously acquired and extinguished contingencies in

a delayed test session (Myers & Davis, 2007; Quirk & Mueller, 2008). Importantly, acquisition and

extinction learning form separate memory traces and individuals may recall previously acquired threat

memories without recalling extinction memories. In other words, they may show long-term extinction

resistance, which is assumed to be key in the maintenance of both non-pathological and pathological

fears (Duits et al., 2015; Lissek et al., 2005; Milad & Quirk, 2012). In this case, despite repeated

omission of aversive events (i.e., extinction), individuals keep showing robust conditioned threat

responses over time.

Conditioned threat responses have different adaptive functions and may manifest at various

levels of central and autonomic physiology. Among these, increased attentive processing promotes

capture of threat-relevant information, improving chances of successful defensive responding

(Bradley, 2009; Lang & Bradley, 2010). Supporting this notion, several studies using visual evoked

brain potentials in humans have found evidence for selectively heightened visual attention when

viewing threat cues (Bublatzky & Schupp, 2012; Gruss, Langaee, & Keil, 2016; Kastner-Dorn,

Andreatta, Pauli, & Wieser, 2018; Michalowski, Pané-Farré, Löw, & Hamm, 2015; Muench, Westermann, Pizzagalli, Hofmann, & Mueller, 2016; Wieser, Reicherts, Juravle, & von Leupoldt, 2016). However, because visual evoked potentials to threat generally reflect early brain activity (i.e. < 500 ms; Miskovic & Keil, 2012) it is unknown, whether enhanced visual attention is sustained throughout later visual processing stages of conditioned threat. Moreover, it is unclear, if heightened attention displays long-term resistance to extinction. These are clinically relevant questions given that enduring prioritization of visual threat processing would presumably interfere with the reduction of fear through extinction and exposure therapy. The present study addresses these open questions using a robust electrophysiological marker of sustained visual attention, suppression of visuocortical oscillatory activity in the alpha range.

Alpha oscillations (e.g., 8-12 Hz; Bartsch, Hamuni, Miskovic, Lang, & Keil, 2015; de Cesarei & Codispoti, 2011) indicate active inhibition processes in neural populations (Klimesch, Sauseng, & Hanslmayr, 2007; Pfurtscheller, 2001). Consequentially, stimulus-evoked posterior alpha suppression to visual stimuli has been associated with disinhibition and increased excitability of early visual areas in the occipital cortex (Green et al., 2017; Iemi & Busch, 2018; Iemi, Chaumon, Crouzet, & Busch, 2017; Klimesch, Fellinger, & Freunberger, 2011; Scheeringa, Koopmans, van Mourik, Jensen, & Norris, 2016) in response to increased attentional demands (Ben-Simon et al., 2013; Cole & Ray, 1985; Foxe & Snyder, 2011; Frey, Ruhnau, & Weisz, 2015; Klimesch, 2012; Ray & Cole, 1985). In addition, posterior alpha power suppression has been found stronger in response to aversive vs. neutral/appetitive pictures(de Cesarei & Codispoti, 2011; Huster, Stevens, Gerlach, & Rist, 2009; Vagnoni, Lourenco, & Longo, 2015; but also see Uusberg, Uibo, Kreegipuu, & Allik, 2013), taken to reflect prioritized visual processing of threat information (Bradley, 2009; Lang & Bradley, 2010).

In the present study, we estimated current source density of alpha-band oscillatory activity specifically at scalp sites consistent with visuocortical sources, to investigate (a) the extent to which heightened visuocortical alpha suppression indexes selective visual attention to threat cues one day after initial conditioning, and (b) if conditioned alpha suppression is extinguished in the long term. For this purpose, we employed a two-day differential threat conditioning paradigm with threat acquisition

1   and extinction on one day and a critical recall test one day later, comparing alpha power changes in

2   response to previously conditioned and extinguished as well as non-extinguished CS.

6

Results

Conditioning and extinction effects during Day 2 recall test

The ANOVA showed a significant main effect of Contingency ($F(1, 86) = 9.85$, $p = .002$, $\eta_p^2 = .103$) as CS+ were followed by a stronger suppression of alpha power compared to CS- (also see Figure 2). Meanwhile, the main effect Extinction ($F(1, 86) = 0.18$, $p = .669$, $\eta_p^2 = .002$) and the Contingency x Extinction interaction ($F(1, 86) = 1.29$, $p = .259$, $\eta_p^2 = .015$) were not significant.

In line with the frequentist ANOVA, Bayesian ANOVA provided strongest evidence for a main effect of Contingency in the absence of other effects ($BF_{10} = 26.0$, all other models: $BF_{10} < 3.3$). In line with this pattern, Bayesian inclusion factors provided support for the inclusion of the main effect of Contingency ($BF_{Incl} = 25.9$) and against the inclusion of the main effect of Extinction ($BF_{Incl} = 1 / 8.1$) or the Contingency x Extinction interaction ($BF_{Incl} = 1 / 4.0$). Figure 3 shows time-frequency plots and topographic mapping of the Contingency effect on alpha power.

Figure 1. **Conditioning effects on Day 2 alpha power.** Mean alpha power (relative to baseline) at occipital sites in the time window of 500 – 1200 ms. Light bars represent CS presented during Day 1 extinction, dark bars represent CS not presented during Day 1 extinction. Error bars indicate SEM based on within-subject variance. **$p < .01$ for the main effect of Contingency.

Figure 2. **Main effect of Contingency on occipital alpha power. (a)** and **(b)** Time-frequency plots for CS+ and CS-, respectively. Power values are % change relative to baseline (-400 to -200 ms) and averaged across Oz, POz, O1, and O2. **(c)** Time-frequency plot of the difference between CS+ and CS-, across Oz, POz, O1, and O2; the white rectangle indicates time (500 to 1200 ms) and frequency (8.1 to 11.9 Hz) windows for statistical analyses. **(d)** Topography of the difference between CS+ and CS- in the a priori defined time window (500 to 1200 ms). White dots depict the electrodes used for statistical analyses.

Time course of Contingency effect

As suggested by permutation *t*-tests, viewing CS+ compared to CS- prompted significantly lower alpha power in the time window from 630 to 1230 ms post-CS (Figure 4), largely converging with our a priori window (500 to 1200 ms).

Figure 3. **Time course of Contingency effect on mean occipital alpha power.** Alpha power (relative to baseline) for collapsed CS+ and CS-, respectively, averaged across occipital sites. The gray-shaded area indicates significant differences between CS+ and CS- as determined by permutation testing ($p < .05$, two-sided).

Discussion

In the present study, we investigated the role of heightened selective attention to threat cues in

long-term threat conditioning and extinction. For this purpose, participants underwent a differential

threat conditioning paradigm with acquisition and extinction on Day 1 and a test phase on Day 2 that

allowed to assess long-term recall of extinguished and non-extinguished threat-conditioned responses.

We found that, one day after threat conditioning, CS evoked a suppression of alpha power at occipital

sites between 600 and 1200 ms, which was more pronounced for threat cues (i.e., CS+) compared to

safety cues (i.e., CS-). Importantly, this conditioning effect was unaffected by Day 1 extinction

training.

In the present sample, we observed robust alpha power suppression at occipital electrodes that

was stronger for threat-conditioned CS+ than for CS-. This is the first time that occipital alpha power

suppression is shown in response to conditioned threat cues. The present effects converge with

previously shown alpha suppression to naturally threatening stimuli (de Cesarei & Codispoti, 2011;

Huster et al., 2009; Vagnoni et al., 2015) and can be interpreted as stronger disinhibition of early

visuocortical areas facilitating attentional prioritization of behaviorally more relevant (i.e., threat) cues

(Foxe & Snyder, 2011; Frey et al., 2015). They are also consistent with previous ERP research

showing threat-related prioritization of visuocortical processing (Bublatzky & Schupp, 2012; Gruss et

al., 2016; Kastner-Dorn et al., 2018; Michalowski et al., 2015; Mueller & Pizzagalli, 2016; Muench et

al., 2016; Thigpen, Bartsch, & Keil, 2017; Wieser et al., 2016). Prioritization of threat is highly

important for the successful choice of adaptive responding (Bradley, 2009; Bradley, Keil, & Lang,

2012; Lang & Bradley, 2010), given limited capacities for information processing in the brain

(Broadbent, 1965; Chun, Golomb, & Turk-Browne, 2011; Lavie, 2005).

Threat-potentiated alpha suppression was found in the time window from about 600 to 1200

ms post-CS. Therefore, this alpha suppression likely indicates *sustained* attention allocation to the

threat cue via recruitment of visuocortical resources (de Cesarei & Codispoti, 2011). This process may

be initiated after initial stimulus recognition and affective stimulus categorization as presumably

indicated by early event-related potential (Miskovic & Keil, 2012; Schupp, Flaisch, Stockburger, &

Junghöfer, 2006) and may reflect a more thorough analysis of threat cues. Taken together, the present

results suggest occipital alpha as a promising new marker of sustained visuocortical attention modulation in response to conditioned threat cues.

Importantly, the stronger alpha suppression in response to CS+ was evident one day after initial learning, indicating successful long-term recall of selectively increased attention allocation to threat cues. To our knowledge, this is the first study to show such long-term threat conditioning effects on selective visual attention, stable across timely spaced situations. The long-term stability of the present alpha suppression suggests selective attention modulation in threat as a central feature of stable fear. While there is some evidence that selective attention to threatening information contributes to the long-term maintenance of fear (Richards, Benson, Donnelly, & Hadwin, 2014; Van Bockstaele et al., 2014), the neural underpinnings of this link are not well understood. The present study suggests a mechanism in which differential activation of visuocortical areas is a temporally stable threat-conditioned response and underlies increased attentional processing of threat cues across time. Increased attentional threat processing may consolidate fearful behavior via increased levels of state fear (and other negative affects; MacLeod, Rutherford, Campbell, Ebsworthy, & Holker, 2002; Van Bockstaele et al., 2014) and by a reduced ability to disengage from threat cues at the cost of other potentially relevant stimuli (Fox et al., 2002; Georgiou et al., 2005; Rinck, Reinecke, Ellwart, Heuer, & Becker, 2005) – e.g., concurrent safety cues or cues for successful coping. Moreover, attentional biases may interact with negative expectancy, memory, and interpretation biases related to threat (Aue & Okon-Singer, 2015). Future studies may use occipital alpha to investigate these and other potential mechanisms in order to elucidate attentional mechanisms of long-term fear maintenance.

In addition to long-term threat recall, we observed long-term extinction resistance of threat-conditioned alpha suppression. In other words, the initially threat-conditioned CS+ continued to evoke conditioned threat responses on Day 2, regardless of Day 1 fear extinction. The conclusion of extinction resistance was backed by Bayesian analyses, which favored models excluding the extinction factor. Note that the present result pattern of extinction-resistant alpha suppression mirrors the pattern of long-term extinction-resistance of (a) rapid processing enhancements of threat-conditioned faces (Mueller & Pizzagalli, 2016) and (b) cardiac deceleration in the present sample (Panitz et al., 2018) as well as in previous studies (Panitz, Hermann, & Mueller, 2015; Sperl, Panitz, Hermann, & Mueller,

2016). Threat-evoked cardiac deceleration has been suggested to also reflect attentional processes, namely orienting in the face of imminent harm (Bastos et al., 2016; Löw, Weymar, & Hamm, 2015; Roelofs, 2017; Wendt, Löw, Weymar, Lotze, & Hamm, 2017). On the other hand, the present sample showed reduced differential skin conductance responses for extinguished (CS+E vs. CS-E) compared to non-extinguished (CS+N vs. CS-N) threat cues (Panitz et al., 2018), which indicates successful extinction recall of CS-US contingency awareness (Hamm & Vaitl, 1996). This suggests that attention-related processes in general, and occipital alpha suppression in particular, could be more extinction-resistant than other conditioned responses and may occur in a better-safe-than-sorry fashion. In other words, the cost of increased attentional engagement to invalid threat cues (i.e., false alarms) may be judged as significantly smaller than the cost for overlooking valid threat cues – even after repeated omission of harmful events.

As discussed earlier, alpha-related attentional processes may be crucial in the long-term maintenance of fears. The observed extinction resistance in the present study suggests that former threat cues continue drawing on attentional resources even in the absence of contingent reinforcement by harmful outcomes, i.e., even when cues are not followed by negative consequences anymore after initial learning. Moreover, occipital alpha suppression may prove useful for investigating the influence of threat-focused attention and attentional biases on the effectiveness of exposure therapy (Barry, Vervliet, & Hermans, 2015; Mogg & Bradley, 2018) .

The limitations of the current study should be addressed. First, using the current paradigm, it cannot be ruled out that CS+ evoked stronger alpha suppression than CS- due to partial reinforcement and changing contingencies across phases, making predictions of US occurrence more difficult in CS+ vs. CS- and motivating participants to process CS+ more thoroughly. This, however, is unlikely to explain the present effects given that threat-depicting pictures, with no learning history, also evoked stronger alpha suppression in previous studies (de Cesarei & Codispoti, 2011; Güntekin & Basar, 2007; Huster et al., 2009; Vagnoni et al., 2015). Second, we only used male participants in order to investigate conditioning and extinction without the influence of known sex differences (Lebron-Milad et al., 2012; Merz et al., 2012). As there also may be gender differences in visuocortical threat

processing (Pintzinger, Pfabigan, Pfau, Kryspin-Exner, & Lamm, 2017; Sass et al., 2010), the present results should be replicated in females.

In the present study, we could show that conditioned visual threat cues evoke enhanced alpha suppression at occipital sites. Moreover, we showed for the first time that increased attention allocation to conditioned threat cues via sustained recruitment of early visuocortical areas is long-term stable and resistant to extinction. Future studies may use occipital alpha power to further elucidate mechanisms of visual attention in the development, maintenance, and extinction of fears.

Materials and Methods

Sample

We analyzed data of a sample of $N = 87$ healthy, male participants (mean age: 23.7, SD: 3.85,

range: 18-34), described in more detail elsewhere (Panitz et al., 2018). Participants were compensated

65 € (ca. 75 US$) for two experimental sessions (total 7 hours on two subsequent days). The study

was conducted in accordance to the Declaration of Helsinki and was approved by the ethics committee

of the German Psychological Association (DGPs).

Conditioning and extinction procedures

Paradigm. We employed a two-day differential threat conditioning and extinction paradigm

(Mueller, Panitz, Hermann, & Pizzagalli, 2014; Panitz et al., 2018; also see Figure 1). Briefly, on Day

1, participants underwent CS habituation, followed by an acquisition phase. Two CS (CS+E, CS+N)

were paired with a US in 21 out of 45 trials (46.6 %), the other two CS (CS-E, CS-N) were never

paired with the US (also 45 trials per CS). The extinction phase started exactly three hours after the

end of acquisition and consisted of 40 CS+E and CS-E presentations ('E' standing for *presented*

during extinction phase). CS+N, CS-N, and US were not presented during the extinction phase ('N'

standing for *not presented during extinction phase)*. Approximately 24 h after the extinction,

participants returned for a Day 2 recall test which included 60 trials of each CS. No US were presented

on Day 2.

Stimuli and trial structure. We used four different male faces with neutral expression from the

Karolinska Directed Emotional Faces set (Lundqvist, Flykt, & Öhman, 1998) as CS (pictures:

AM10NES, AM13NES, AM31NES, BM08NES; also see Figure 1). Assignment of face stimuli to the

different CS types was permutated and balanced across participants. The US was a 1 s white noise

burst at 95 dB(A) delivered by a room speaker as we had previously shown that noise bursts are

particularly well suited for threat conditioning with many trials (Sperl et al., 2016). In every trial, a

fixation cross (1 s duration) was presented before participants saw the CS for 4 s. In paired trials the

CS co-terminated with the US for 1 s. A black screen (jittered duration, 6-8 s) was presented between

trials.

	Day 1			Day 2 (24 h later)
	Habituation	**Acquisition**	**Extinction**	**Recall Test**
	5 CS+E	45 CS+E **+ 21 US**	40 CS+E	60 CS+E
	5 CS-E	45 CS-E	40 CS-E	60 CS-E
	5 CS+N	45 CS+N **+ 21 US**		60 CS+N
	5 CS-N	45 CS-N		60 CS-N

Figure 4. **Conditioning and extinction paradigm**. Face stimuli and number of presentations in the two-day differential threat conditioning and extinction paradigm. The US was only presented during acquisition, indicated by the speaker symbol. Assignment of different faces to CS type was permutated across participants. CS+E = extinguished CS+, CS+N = non-extinguished CS+, CS-E = CS- presented during extinction phase, CS-N = CS- not presented during extinction phase. Stimuli were presented in color. KDEF stimuli IDs from top to bottom: AM10NES, AM13NES, AM31NES, BM08NES. Figure from Panitz et al. (2018).

EEG recording and preprocessing.

64-channel EEG was recorded with a QuickAmp72 amplifier and actiCAP active electrodes (Brain Products, Gilching, Germany) at 1000 Hz, with a 200 Hz online lowpass-filter and referenced against the average. EEG processing was performed in BrainVision Analyzer 2 (Brain Products, Gilching, Germany). The EEG was downsampled to 500 Hz, highpass-filtered (-3 dB at 0.1 Hz, 24 dB/oct., zero-phase IIR Butterworth filter) and notch-filtered (50 Hz, 5 Hz bandwidth, 16th order). Extended Infomax ICA (Lee, Girolami, & Sejnowski, 1999) was applied to the continuous data. Critical independent components reflective of horizontal and vertical eye movements, blinks, and cardiac artifacts were identified and removed by an experienced rater. To increase signal stationarity required for ICA, large EEG artifacts were removed manually, the signal was 0.5 Hz highpass-filtered for ICA only and the resulting weights were subsequently applied to the 0.1 Hz filtered data (Winkler, Debener, Muller, & Tangermann, 2015). Segments with remaining artifacts were removed manually and channels with excessive amounts of bad data were interpolated (spline interpolation). Continuous

EEG data were lowpass-filtered (30 Hz, filter specifications identical to highpass filter). After correcting marker latencies for monitor delay (33 ms), data were segmented relative to CS onset (-600 to 2000 ms).

To estimate current source density and improve the spatial specificity of the voltage map, the surface Laplacian was computed using the algorithm of Perrin, Pernier, Bertrand, and Echallier (1989) as implemented in BrainVision Analyzer 2 (spline order: 4, Legendre polynomial: 10, lambda: 1e-5). In order to facilitate comparison with previous studies, we also analyzed the average-referenced scalp data. The results converged with the present results in the surface Laplacian data and are provided in the Supplement.

Wavelet analyses.

Wavelet analyses were conducted in MATLAB 2013a (MathWorks, Natick, MA, USA) using established custom scripts (Keil, Stolarova, Moratti, & Ray, 2007). First, EEG segments in the time domain were baseline-corrected (subtraction of the mean from -600 to -500 ms) and tapered with a cosine square window (20 samples rise/fall). Complex Morlet wavelets were applied with variable bandwidths (Morlet parameter = f/σ_f = 12) to compute power for frequency bands from 3.8 to 30.4 Hz in linear steps of 0.38 Hz. Power in each frequency band was baseline-corrected by dividing the signal by the mean amplitude between -400 ms and -200 ms relative to CS onset. Mean power of all discrete frequencies from 8.1 Hz to 11.9 Hz (e.g., Bartsch et al., 2015; de Cesarei & Codispoti, 2011) was used for statistical analyses on alpha.

Statistical analyses

In order to assess the influence of Day 1 learning on occipital alpha during Day 2, we computed mean alpha power across Oz and adjacent electrodes (Oz, POz, O1, and O2) in the time window of 500 – 1200 ms after the CS. The resulting values were entered into an ANOVA with the within-subject factors Contingency (CS+ vs. CS-) and Extinction (extinguished vs. non-extinguished) using the *aov* function in R (R Core Team, 2018). In addition to null-hypothesis testing, we conducted Bayesian analyses as implemented in the *anovaBF* function in the BayesFactor package (Morey &

Rouder, 2018) for R. We computed Bayes factors (100,000 iterations) for four different models (main

effect Contingency only, main effect Extinction only, additive model of Contingency + Extinction

main effects, complete model including both main effects and the interaction term; cf. Rouder et al.,

2016) compared to the null model (BF_{10}). All models had equal prior probabilities and included a

random effect term to account for between-subject variance. In a second step, we computed Bayesian

inclusion factors (BF_{Incl}) for each effect (i.e., Contingency, Extinction, and Contingency x Extinction)

that indicate whether models including this effect are more likely to explain the data than matched

models without this effect. More precisely, all models containing the effect of interest – but no higher-

order interactions of this effect – were compared to matched models stripped of the effect (BAWS

factor suggested by Sebastiaan Mathôt; also implemented in JASP 0.9, JASP Team, 2018). As an

example, for the Contingency effect, the models *Contingency only* and *Contingency + Extinction* were

compared to the null model and the *Extinction only* model.

After using a predefined time window (500 to 1200 ms; Chien et al., 2018; Vagnoni et al.,

2015) for hypothesis testing, we conducted follow-up analyses on the exact time window of the

Contingency effect based on the present data. We used permutation-controlled *t*-tests (adapted from

the t_{max} approach from Blair & Karniski, 1993), comparing alpha power (averaged across Oz, POz,

O1, O2) at each time sample between both CS+ and both CS- (i.e., Contingency contrast). First, we

randomly permutated the CS+ and CS- condition in each participant 1,000 times and computed *t*-

values for each of the 1,300 time samples. Then, the tails of each permutation's *t*-value distribution

were determined and stored in a t_{min} and t_{max} distribution, respectively, each having 1,000 values

corresponding to the 1,000 permutations. Finally, 2.5^{th} and 97.5^{th} percentiles from the resulting

distribution of t_{min} and t_{max} values were used as critical *t*-values to compare empirical *t*-values against

(i.e., $\alpha = .05$, two-sided; t_{crit}: CS+ < CS-: -2.59; CS+ > CS-: 2.64).

References

Aue, T., & Okon-Singer, H. (2015). Expectancy biases in fear and anxiety and their link to biases in attention. *Clinical Psychology Review*, *42*, 83–95. doi: 10.1016/j.cpr.2015.08.005

Barry, T. J., Vervliet, B., & Hermans, D. (2015). An integrative review of attention biases and their contribution to treatment for anxiety disorders. *Frontiers in Psychology*, *6*, 968. doi: 10.3389/fpsyg.2015.00968

Bartsch, F., Hamuni, G., Miskovic, V., Lang, P. J., & Keil, A. (2015). Oscillatory brain activity in the alpha range is modulated by the content of word-prompted mental imagery. *Psychophysiology*, *52*, 727–735. doi: 10.1111/psyp.12405

Bastos, A. F., Vieira, A. S., Oliveira, J. M., Oliveira, L., Pereira, M. G., Figueira, I., … Volchan, E. (2016). Stop or move: Defensive strategies in humans. *Behavioural Brain Research*, *302*, 252–262. doi: 10.1016/j.bbr.2016.01.043

Ben-Simon, E., Podlipsky, I., Okon-Singer, H., Gruberger, M., Cvetkovic, D., Intrator, N., & Hendler, T. (2013). The dark side of the alpha rhythm: FMRI evidence for induced alpha modulation during complete darkness. *European Journal of Neuroscience*, *37*, 795–803. doi: 10.1111/ejn.12083

Blair, R. C., & Karniski, W. (1993). An alternative method for significance testing of waveform difference potentials. *Psychophysiology*, *30*, 518–524. doi: 10.1111/j.1469-8986.1993.tb02075.x

Bradley, M. M. (2009). Natural selective attention: Orienting and emotion. *Psychophysiology*, *46*, 1–11. doi: 10.1111/j.1469-8986.2008.00702.x

Bradley, M. M., Keil, A., & Lang, P. J. (2012). Orienting and emotional perception: facilitation, attenuation, and interference. *Forntiers in Psychology*, *3*, 493. doi: 10.3389/fpsyg.2012.00493

Broadbent, D. E. (1965). Information Processing in the Nervous System. *Science*, *150*, 457–462. doi: 10.1126/science.150.3695.457

Bublatzky, F., & Schupp, H. T. (2012). Pictures cueing threat: Brain dynamics in viewing explicitly instructed danger cues. *Social Cognitive and Affective Neuroscience*, *7*, 611–622. doi: 10.1093/scan/nsr032

Chien, J. H., Colloca, L., Korzeniewska, A., Cheng, J. J., Campbell, C. M., Hillis, A. E., & Lenz, F. A.

(2018). Oscillatory EEG Activity Induced by Conditioning Stimuli During Fear Conditioning Reflects Salience and Valence of these Stimuli More than Expectancy. *Neuroscience*, *346*, 81–93. doi: 10.1016/j.neuroscience.2016.12.047

Chun, M. M., Golomb, J. D., & Turk-Browne, N. B. (2011). A Taxonomy of External and Internal Attention. *Annual Review of Psychology*, *62*, 73–101. doi: 10.1146/annurev.psych.093008.100427

Cole, H. W., & Ray, W. J. (1985). EEG correlates of emotional tasks related to attentional demands. *International Journal of Psychophysiology*, *3*, 33–41. doi: 10.1016/0167-8760(85)90017-0

de Cesarei, A., & Codispoti, M. (2011). Affective modulation of the LPP and α-ERD during picture viewing. *Psychophysiology*, *48*, 1397–1404. doi: 10.1111/j.1469-8986.2011.01204.x

Duits, P., Cath, D. C., Lissek, S., Hox, J. J., Hamm, A. O., Engelhard, I. M., … Baas, J. M. P. (2015). Updated meta-analysis of classical fear conditioning in the anxiety disorders. *Depression and Anxiety*, *32*, 239–253. doi: 10.1002/da.22353

Fox, E., Russo, R., Dutton, K., Fox, E., Russo, R., & Dutton, K. (2002). Attentional bias for threat: Evidence for delayed disengagement from emotional faces. *Cognition and Emotion*, *16*, 355–379. doi: 10.1080/02699930143000527

Foxe, J. J., & Snyder, A. C. (2011). The role of alpha-band brain oscillations as a sensory suppression mechanism during selective attention. *Frontiers in Psychology*, *2*, 154. doi: 10.3389/fpsyg.2011.00154

Frey, J. N., Ruhnau, P., & Weisz, N. (2015). Not so different after all: The same oscillatory processes support different types of attention. *Brain Research*, *1626*, 183–197. doi: 10.1016/j.brainres.2015.02.017

Georgiou, G. A., Bleakley, C., Hayward, J., Russo, R., Dutton, K., Eltiti, S., & Fox, E. (2005). Focusing on fear: Attentional disengagement from emotional faces. *Visual Cognition*, *12*, 145–158. doi: 10.1080/13506280444000076

Green, J. J., Boehler, C. N., Roberts, K. C., Chen, L.-C., Krebs, R. M., Song, A. W., & Woldorff, M. G. (2017). Cortical and Subcortical Coordination of Visual Spatial Attention Revealed by Simultaneous EEG–fMRI Recording. *The Journal of Neuroscience*, *37*, 7803–7810. doi:

10.1523/JNEUROSCI.0326-17.2017

Gruss, L. F., Langaee, T., & Keil, A. (2016). The role of the COMT val158met polymorphism in mediating aversive learning in visual cortex. *NeuroImage*, *125*, 633–642. doi: 10.1016/j.neuroimage.2015.11.003

Güntekin, B., & Basar, E. (2007). Emotional face expressions are differentiated with brain oscillations. *International Journal of Psychophysiology*, *64*, 91–100. doi: 10.1016/j.ijpsycho.2006.07.003

Hamm, A. O., & Vaitl, D. (1996). Affective learning: Awareness and aversion. *Psychophysiology*, *33*, 698–710. doi: 10.1111/j.1469-8986.1996.tb02366.x

Huster, R. J., Stevens, S., Gerlach, A. L., & Rist, F. (2009). A spectralanalytic approach to emotional responses evoked through picture presentation. *International Journal of Psychophysiology*, *72*, 212–216. doi: 10.1016/j.ijpsycho.2008.12.009

Iemi, L., & Busch, N. A. (2018). Moment-to-Moment Fluctuations in Neuronal Excitability Bias Subjective Perception Rather than Strategic Decision-Making. *ENeuro*, *5*, 1–13. doi: 10.1523/ENEURO.0430-17.2018

Iemi, L., Chaumon, M., Crouzet, S. M., & Busch, N. A. (2017). Spontaneous Neural Oscillations Bias Perception by Modulating Baseline Excitability. *The Journal of Neuroscience*, *37*, 807–819. doi: 10.1523/JNEUROSCI.1432-16.2017

JASP Team. (2018). JASP (Version 0.9)[Computer Software].

Kastner-Dorn, A. K., Andreatta, M., Pauli, P., & Wieser, M. J. (2018). Hypervigilance during anxiety and selective attention during fear: Using steady-state visual evoked potentials (ssVEPs) to disentangle attention mechanisms during predictable and unpredictable threat. *Cortex*, *106*, 120–131. doi: 10.1016/j.cortex.2018.05.008

Keil, A., Stolarova, M., Moratti, S., & Ray, W. J. (2007). Adaptation in human visual cortex as a mechanism for rapid discrimination of aversive stimuli. *NeuroImage*, *36*, 472–479. doi: 10.1016/j.neuroimage.2007.02.048

Klimesch, W. (2012). Alpha-band oscillations, attention, and controlled access to stored information. *Trends in Cognitive Sciences*, *16*, 606–617. doi: 10.1016/j.tics.2012.10.007

Klimesch, W., Fellinger, R., & Freunberger, R. (2011). Alpha oscillations and early stages of visual encoding. *Frontiers in Psychology*, *2*, 118. doi: 10.3389/fpsyg.2011.00118

Klimesch, W., Sauseng, P., & Hanslmayr, S. (2007). EEG alpha oscillations: The inhibition-timing hypothesis. *Brain Research Reviews*, *53*, 63–88. doi: 10.1016/j.brainresrev.2006.06.003

Lang, P. J., & Bradley, M. M. (2010). Emotion and the motivational brain. *Biological Psychology*, *84*, 437–450. doi: 10.1016/j.biopsycho.2009.10.007

Lavie, N. (2005). Distracted and confused?: Selective attention under load. *Trends in Cognitive Sciences*, *9*, 75–82. doi: 10.1016/j.tics.2004.12.004

Lebron-Milad, K., Abbs, B., Milad, M. R., Linnman, C., Rougemount-Bücking, A., Zeidan, M. A., … Goldstein, J. M. (2012). Sex differences in the neurobiology of fear conditioning and extinction: a preliminary fMRI study of shared sex differences with stress-arousal circuitry. *Biology of Mood & Anxiety Disorders*, *2*, 7. doi: 10.1186/2045-5380-2-7

LeDoux, J. E. (2014). Coming to terms with fear. *Proceedings of the National Academy of Sciences*, *111*, 2871–2878. doi: 10.1073/pnas.1400335111

Lee, T. W., Girolami, M., & Sejnowski, T. J. (1999). Independent component analysis using an extended infomax algorithm for mixed subgaussian and supergaussian sources. *Neural Computation*, *11*, 417–441. doi: 10.1162/089976699300016719

Lissek, S., Powers, A. S., McClure, E. B., Phelps, E. A., Woldehawariat, G., Grillon, C., & Pine, D. S. (2005). Classical fear conditioning in the anxiety disorders: A meta-analysis. *Behaviour Research and Therapy*, *43*, 1391–1424. doi: 10.1016/j.brat.2004.10.007

Lonsdorf, T. B., Menz, M. M., Andreatta, M., Fullana, M. A., Golkar, A., Haaker, J., … Merz, C. J. (2017). Don't fear 'fear conditioning': Methodological considerations for the design and analysis of studies on human fear acquisition, extinction, and return of fear. *Neuroscience and Biobehavioral Reviews*, *77*, 247–285. doi: 10.1016/j.neubiorev.2017.02.026

Löw, A., Weymar, M., & Hamm, A. O. (2015). When Threat Is Near, Get Out of Here: Dynamics of Defensive Behavior During Freezing and Active Avoidance. *Psychological Science*, *26*, 1706–1716. doi: 10.1177/0956797615597332

Lundqvist, D., Flykt, A., & Öhman, A. (1998). *The Karolinska Directed Emotional Faces - KDEF.*

CD ROM from the Department of Clinical Neuroscience, Psychology Section, Karolinska Institutet.

MacLeod, C., Rutherford, E., Campbell, L., Ebsworthy, G., & Holker, L. (2002). Selective Attention and Emotional Vulnerability: Assessing the Causal Basis of Their Association Through the Experimental Manipulation of Attentional Bias. *Journal of Abnormal Psychology*, *111*, 107–123. doi: 10.1037//0021-843X.111.1.107

Merz, C. J., Tabbert, K., Schweckendiek, J., Klucken, T., Vaitl, D., Stark, R., & Wolf, O. T. (2012). Neuronal correlates of extinction learning are modulated by sex hormones. *Social Cognitive and Affective Neuroscience*, *7*, 819–830. doi: 10.1093/scan/nsr063

Michalowski, J. M., Pané-Farré, C. A., Löw, A., & Hamm, A. O. (2015). Brain dynamics of visual attention during anticipation and encoding of threat- and safe-cues in spider-phobic individuals, *10*, 1177–1186. doi: doi:10.1093/scan/nsv002

Milad, M. R., & Quirk, G. J. (2012). Fear Extinction as a Model for Translational Neuroscience: Ten Years of Progress. *Annual Review of Psychology*, *63*, 129–151. doi: 10.1146/annurev.psych.121208.131631

Miskovic, V., & Keil, A. (2012). Acquired fears reflected in cortical sensory processing: A review of electrophysiological studies of human classical conditioning. *Psychophysiology*, *49*, 1230–1241. doi: 10.1111/j.1469-8986.2012.01398.x

Mogg, K., & Bradley, B. P. (2018). Anxiety and Threat-Related Attention: Cognitive-Motivational Framework and Treatment. *Trends in Cognitive Sciences*, *22*, 225–240. doi: 10.1016/j.tics.2018.01.001

Morey, R. D., & Rouder, J. N. (2018). BayesFactor: Computation of Bayes Factors for Common Designs.

Mueller, E. M., Panitz, C., Hermann, C., & Pizzagalli, D. A. (2014). Prefrontal Oscillations during Recall of Conditioned and Extinguished Fear in Humans. *Journal of Neuroscience*, *34*, 7059–7066. doi: 10.1523/JNEUROSCI.3427-13.2014

Mueller, E. M., & Pizzagalli, D. A. (2016). One-year-old fear memories rapidly activate human fusiform gyrus. *Social Cognitive and Affective Neuroscience*, *11*, 308–316. doi:

10.1093/scan/nsv122

Muench, H. M., Westermann, S., Pizzagalli, D. A., Hofmann, S. G., & Mueller, E. M. (2016). Self-relevant threat contexts enhance early processing of fear-conditioned faces. *Biological Psychology*, *121*, 194–202. doi: 10.1016/j.biopsycho.2016.07.017

Myers, K. M., & Davis, M. (2007). Mechanisms of fear extinction. *Molecular Psychiatry*, *12*, 120–150. doi: 10.1038/sj.mp.4001939

Panitz, C., Hermann, C., & Mueller, E. M. (2015). Conditioned and extinguished fear modulate functional corticocardiac coupling in humans. *Psychophysiology*, *52*, 1351–1360. doi: 10.1111/psyp.12498

Panitz, C., Sperl, M. F. J., Hennig, J., Klucken, T., Hermann, C., & Mueller, E. M. (2018). Fearfulness, neuroticism/anxiety, and COMT Val158Met in long-term fear conditioning and extinction. *Neurobiology of Learning and Memory*, *155*, 7–20. doi: 10.1016/j.nlm.2018.06.001

Perrin, F., Pernier, J., Bertrand, O., & Echallier, J. F. (1989). Spherical splines for scalp potential and current density mapping. *Electroencephalography and Clinical Neurophysiology*, *72*, 184–187. doi: 10.1016/0013-4694(89)90180-6

Pfurtscheller, G. (2001). Functional brain imaging based on ERD / ERS. *Vision Research*, *41*, 1257–1260. doi: 10.1016/S0042-6989(00)00235-2

Pintzinger, N. M., Pfabigan, D. M., Pfau, L., Kryspin-Exner, I., & Lamm, C. (2017). Temperament differentially influences early information processing in men and women: Preliminary electrophysiological evidence of attentional biases in healthy individuals. *Biological Psychology*, *122*, 69–79. doi: 10.1016/j.biopsycho.2016.07.007

Quirk, G. J., & Mueller, D. (2008). Neural mechanisms of extinction learning and retrieval. *Neuropsychopharmacology*, *33*, 56–72. doi: 10.1038/sj.npp.1301555

R Core Team. (2018). R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing.

Ray, W. J., & Cole, H. W. (1985). EEG alpha activity reflects attentional demands, and beta activity reflects emotional and cognitive processes. *Science*, *228*, 750–752. doi: 10.1126/science.3992243

Rescorla, R. A., & Wagner, A. R. (1972). A theory of Pavlovian conditioning: Variations in the

effectiveness of reinforcement and nonreinforcement. In A. H. Black & W. F. Prokasy (Eds.), *Classical conditioning II: Current research and theory* (pp. 64–99). New York: Appleton-Century-Crofts.

Richards, H. J., Benson, V., Donnelly, N., & Hadwin, J. A. (2014). Exploring the function of selective attention and hypervigilance for threat in anxiety. *Clinical Psychology Review*, *34*, 1–13. doi: 10.1016/j.cpr.2013.10.006

Rinck, M., Reinecke, A., Ellwart, T., Heuer, K., & Becker, E. S. (2005). Speeded Detection and Increased Distraction in Fear of Spiders: Evidence From Eye Movements. *Journal of Abnormal Psychology*, *114*, 235–248. doi: 10.1037/0021-843X.114.2.235

Roelofs, K. (2017). Freeze for action: Neurobiological mechanisms in animal and human freezing. *Philosophical Transactions of the Royal Society B*, *372*, 20160206. doi: 10.1098/rstb.2016.0206

Rouder, J. N., Engelhardt, C. R., McCabe, S., & Morey, R. D. (2016). Model comparison in ANOVA. *Psychonomic Bulletin and Review*, *23*, 1779–1786. doi: 10.3758/s13423-016-1026-5

Sass, S. M., Heller, W., Stewart, J. L., Levin Silton, R., Edgar, J. C., Fisher, J. E., & Miller, G. A. (2010). Time course of attentional bias in anxiety: Emotion and gender specificity. *Psychophysiology*, *47*, 247–259. doi: 10.1111/j.1469-8986.2009.00926.x

Scheeringa, R., Koopmans, P. J., van Mourik, T., Jensen, O., & Norris, D. G. (2016). The relationship between oscillatory EEG activity and the laminar-specific BOLD signal. *Proceedings of the National Academy of Sciences*, *113*, 6761–6766. doi: 10.1073/pnas.1522577113

Schupp, H. T., Flaisch, T., Stockburger, J., & Junghöfer, M. (2006). Emotion and attention: event-related brain potential studies. *Progress in Brain Research*, *156*, 31–51. doi: 10.1016/S0079-6123(06)56002-9

Sperl, M. F., Panitz, C., Hermann, C., & Mueller, E. M. (2016). A pragmatic comparison of noise burst and electric shock unconditioned stimuli for fear conditioning research with many trials. *Psychophysiology*, *53*. doi: 10.1111/psyp.12677

Thigpen, N. N., Bartsch, F., & Keil, A. (2017). The malleability of emotional perception: Short-term plasticity in retinotropic neurons accompanies the formation of perceptual biases to threat. *Journal of Experimental Psychology: General*, *146*, 464–471. doi: 10.1037/xge0000283.The

Uusberg, A., Uibo, H., Kreegipuu, K., & Allik, J. (2013). EEG alpha and cortical inhibition in affective attention. *International Journal of Psychophysiology*, *89*, 26–36. doi: 10.1016/j.ijpsycho.2013.04.020

Vagnoni, E., Lourenco, S. F., & Longo, M. R. (2015). Threat modulates neural responses to looming visual stimuli. *European Journal of Neuroscience*, *42*, 2190–2202. doi: 10.1111/ejn.12998

Van Bockstaele, B., Verschuere, B., Tibboel, H., Houwer, J. De, Crombez, G., & Koster, E. H. W. (2014). A Review of Current Evidence for the Causal Impact of Attentional Bias on Fear and Anxiety. *Psychological Bulletin*, *140*, 682–721. doi: 10.1037/a0034834

Wendt, J., Löw, A., Weymar, M., Lotze, M., & Hamm, A. O. (2017). Active avoidance and attentive freezing in the face of approaching threat. *NeuroImage*, *158*, 196–204. doi: 10.1016/j.neuroimage.2017.06.054

Wieser, M. J., Reicherts, P., Juravle, G., & von Leupoldt, A. (2016). Attention mechanisms during predictable and unpredictable threat — A steady-state visual evoked potential approach. *NeuroImage*, *139*, 167–175. doi: 10.1016/j.neuroimage.2016.06.026

Winkler, I., Debener, S., Muller, K. R., & Tangermann, M. (2015). On the influence of high-pass filtering on ICA-based artifact reduction in EEG-ERP. *Proceedings of the Annual International Conference of the IEEE Engineering in Medicine and Biology Society, EMBS*, *2015–Novem*, 4101–4105. doi: 10.1109/EMBC.2015.7319296

Supplementary Material

In order to facilitate comparison with de Cesarei & Codispoti (2011) and Vagnoni et al. (2015), we

conducted analyses on posterior alpha power using average-referenced EEG all parieto-occipital sites

(here: P7, P5, P3, P1, Pz, P2, P4, P6, P8, PO9, PO7, PO3, POz, PO4, PO8, PO10, O1, Oz, O2). All

other parameters (e.g., time window, wavelet parameters) are identical to the main analysis of the

present manuscript.

Results on Day 2 recall test

In line with the main analyses on CSD data, the ANOVA showed a significantly stronger

alpha suppression to CS+ vs CS- (main effect of Contingency: $F(1, 86) = 15.3$, $p < .001$, $\eta_p^2 = .151$;

Supplementary Figure 1). The main effect Extinction ($F(1, 86) = 0.88$, $p = .350$, $\eta_p^2 = .010$) and the

Contingency x Extinction interaction ($F(1, 86) = 0.95$, $p = .333$, $\eta_p^2 = .011$) were not significant.

In line with the frequentist ANOVA, Bayesian ANOVA provided strongest evidence for a

main effect of Contingency in the absence of other effects ($BF_{10} = 90.1$, all other models: $BF_{10} < 16.4$).

In line with this pattern, Bayesian inclusion factors provided support for the inclusion of the main

effect of Contingency ($BF_{Incl} = 90.6$) and against the inclusion of the main effect of Extinction (BF_{Incl}

$= 1 / 5.5$) or the Contingency x Extinction interaction ($BF_{Incl} = 1 / 3.9$). Supplementary Figure 2 shows

time-frequency plots and topographic mapping of the Contingency effect on alpha power on average-

referenced EEG.

Supplementary Figure 1. **Conditioning effects on Day 2 alpha power.** Mean alpha power (relative to baseline)
at P-, PO-, and O-electrodes in the time window of 500 – 1200 ms. Light bars represent CS presented during
Day 1 extinction, dark bars represent CS not presented during Day 1 extinction. Error bars indicate SEM based
on within-subject variance. ***p < .01 for the main effect of Contingency.

Supplementary Figure 2. **Main effect of Contingency on occipital alpha power.** (a) and (b) Time-frequency
plots for CS+ and CS-, respectively. Power values are % change relative to baseline (-400 to -200 ms) and
averaged across all P-, PO-, and O-electrodes. (c) Time-frequency plot of the difference between CS+ and CS-,
across all P-, PO-, and O-electrodes; the white rectangle indicates time (500 to 1200 ms) and frequency (8.1 to
11.9 Hz) windows for statistical analyses. (d) Topography of the difference between CS+ and CS- in the a priori
defined time window (500 to 1200 ms). White dots depict the electrodes used for statistical analyses.
9

10