Loss of centromere function drives karyotype evolution in closely related *Malassezia*
species

Sundar Ram Sankaranarayanan[1], Giuseppe Ianiri[2,$], Marco A. Coelho[2], Md. Hashim Reza[1],
Bhagya C. Thimmappa[1,#], Promit Ganguly[1], Rakesh Netha Vadnala[3], Sheng Sun[2], Rahul
Siddharthan[3], Christian Tellgren-Roth[4], Thomas L Dawson Jr.[5,6], Joseph Heitman[2,*], and
Kaustuv Sanyal[1,*]

[1]Molecular Mycology Laboratory, Molecular Biology and Genetics Unit, Jawaharlal Nehru
Center for Advanced Scientific Research, Jakkur P.O, Bengaluru- 560064

[2]Department of Molecular Genetics and Microbiology, Duke University Medical Center,
Durham, NC 27710, USA

[3]The Institute of Mathematical Sciences/HBNI, Taramani, Chennai 600113, India

[4]National Genomics Infrastructure, Science for Life Laboratory, Department of Immunology,
Genetics and Pathology, Uppsala University, 75108 Uppsala, Sweden

[5]Skin Research Institute Singapore, Agency for Science, Technology, and Research (A*STAR),
138648, Singapore

[6]Medical University of South Carolina, School of Pharmacy, Department of Drug Discovery,
29425, USA

[$]Present address: Department of Agricultural, Environmental and Food Sciences, University of
Molise, 86100, Campobasso, Italy

[#]Present address: Department of Biochemistry, Robert-Cedergren Centre for Bioinformatics and
Genomics, University of Montreal, 2900 Edouard-Montpetit, Montreal, H3T1J4, QC, Canada

E-mail addresses of authors

SRS (sundar_ram@jncasr.ac.in), GI (giuseppe.ianiri@unimol.it), MHR
(hashimreza@jncasr.ac.in), BCT (bhagyathimmappa@gmail.com), PG
(promitganguly1@gmail.com), MAC (marco.dias.coelho@duke.edu), RNV
(rakeshnetha@imsc.res.in), SS (sheng.sun@duke.edu), RS (rsidd@imsc.res.in), CTR
(christian.tellgren@igp.uu.se), TLD (thomas_dawson@sris.a-star.edu.sg)

*Corresponding authors

Kaustuv Sanyal (sanyal@jncasr.ac.in)

Joseph Heitman (heitm001@duke.edu)

Classification

Biological Sciences, Genetics

Keywords: Chromosome fusion, skin microbe, double-strand breaks, kinetochore, dicentric.

Abstract

Genomic rearrangements associated with speciation often result in chromosome number

variation among closely related species. *Malassezia* species show variable karyotypes ranging

between 6 and 9 chromosomes. Here, we experimentally identified all 8 centromeres in *M.*

sympodialis as 3 to 5 kb long kinetochore-bound regions spanning an AT-rich core and depleted

of the canonical histone H3. Centromeres of similar sequence features were identified as CENP-

A-rich regions in *Malassezia furfur* with 7 chromosomes, and histone H3 depleted regions in

Malassezia slooffiae and *Malassezia globosa* with 9 chromosomes each. Analysis of synteny

conservation across centromeres with newly generated chromosome-level genome assemblies

suggests two distinct mechanisms of chromosome number reduction from an inferred 9-

chromosome ancestral state: (a) chromosome breakage followed by loss of centromere DNA

and (b) centromere inactivation accompanied by changes in DNA sequence following

chromosome-chromosome fusion. We propose AT-rich centromeres drive karyotype diversity in

the *Malassezia* species complex through breakage and inactivation.

Introduction

Centromeres are the genomic loci on which the kinetochore, a multi-subunit complex, assembles to facilitate high fidelity chromosome segregation. The centromere-specific histone H3 variant CENP-A is the epigenetic hallmark of centromeres, as it replaces canonical histone H3 in the nucleosomes to make specialized centromeric chromatin that acts as the foundation to recruit other kinetochore proteins. A remarkable diversity in the organization of centromere DNA sequences has been observed to accomplish this conserved role (Roy and Sanyal, 2011, Yadav et al., 2018b).

The smallest known centromeres are the point centromeres present in budding yeasts of the family *Saccharomycetaceae* that span <200 bp in length (Clarke and Carbon, 1980, Gordon et al., 2011, Kobayashi et al., 2015). These centromeres are organized into conserved DNA elements I, II, and III that are recognized by a cognate kinetochore protein complex called the CBF3 complex, making them genetically defined centromeres. Small regional centromeres, identified in several *Candida* species, form the second category (Sanyal et al., 2004, Padmanabhan et al., 2008, Kapoor et al., 2015, Chatterjee et al., 2016) and have a 2 to 5 kb region enriched by kinetochore proteins. These centromeres can either have unique DNA sequences or a homogenized core that is flanked by inverted repeats. The third type of centromere structure is the large regional centromere, often repetitive in sequence and spanning over 15 kb. Large regional centromeres can be transposon enriched as in *Cryptococcus* species or organized into repeat structures around a central core as in *Schizosaccharomyces pombe* (Chikashige et al., 1989, Clarke and Baum, 1990, Sun et al., 2017, Yadav et al., 2018b).

While the organization of DNA elements is variable, a majority of known centromeres share AT-richness as a common feature. Examples include the CDEII of point centromeres, central core sequences in *S. pombe*, and centromeres of *Neurospora crassa*, *Magnaporthe oryzae*, *Plasmodium falciparum*, and diatoms (Fitzgerald-Hayes et al., 1982, Iwanaga et al., 2010, Rhind et al., 2011, Kapoor et al., 2015, Diner et al., 2017, Yadav et al., 2019). Even the recently described mosaic centromere structure observed in *Mucor circinelloides* that lost CENP-A comprises an AT-rich kinetochore-bound core region (Navarro-Mendoza et al., 2019). While suppression of recombination around centromeres has been correlated with reduced GC content (Lynch et al., 2010), the genetic underpinning of how an AT-rich DNA region favors

kinetochore assembly remains unclear. Ironically, AT-rich sequences have been shown to be

fragile sites of a chromosome (Zhang and Freudenreich, 2007).

Several lines of evidence suggest that centromeres are species-specific and are among the

most rapidly evolving genomic regions, even between closely related species (Bensasson et al.,

2008, Padmanabhan et al., 2008, Rhind et al., 2011, Roy and Sanyal, 2011). This is accompanied

by the concomitant evolution of CENP-A and the associated kinetochore proteins (Talbert et al.,

2004). Functional incompatibilities between centromeres result in uniparental genome

elimination in interspecies hybrids (Ravi and Chan, 2010, Sanei et al., 2011). The divergent

nature of centromeres is proposed to be a driving force for speciation (Henikoff et al., 2001,

Malik and Henikoff, 2009).

Asexual organisms, by virtue of inter- and intra-chromosomal rearrangements, diversify

into species clusters that are distinct in genotype and morphology (Barraclough et al., 2003).

These genotypic differences include changes in both chromosomal organization and number.

Centromere function is directly related to karyotype stabilization following a change in

chromosome number. Rearrangements in the form of telomere-telomere fusions and nested

chromosome insertions (NCIs), wherein an entire donor chromosome is 'inserted' into or near

the centromere of a non-homologous recipient chromosome, are among the major sources of

chromosome number reduction (Lysak, 2014). Such events often result in the formation of

dicentric chromosomes that are subsequently stabilized by breakage-fusion-bridge cycles

(McClintock, 1941) or via inactivation of one centromere through different mechanisms (Han et

al., 2009, Sato et al., 2012). Well known examples of telomere-telomere fusions include the

formation of extant human chromosome 2 by fusion of two ancestral chromosomes (Yunis and

Prakash, 1982, Jdo et al., 1991), the reduction in karyotype observed within the members of the

Saccharomycotina such as *Candida glabrata*, *Vanderwaltozyma polyspora*, *Kluyveromyces*

lactis, and *Zygosaccharomyces rouxii* (Gordon et al., 2011), and the exchange of chromosomal

arms seen in plants and fungi (Schubert and Lysak, 2011, Sun et al., 2017). NCIs have

predominantly shaped karyotype evolution in grasses (Murat et al., 2010). Chromosome number

reduction by centromere loss has also been reported (Gordon et al., 2011).

To investigate if centromere breakage can be a natural source of karyotype diversity in

closely related species, we sought to identify centromeres in a group of *Malassezia* yeast species

that exhibit chromosome number variation. *Malassezia* species are lipid-dependent basidiomycetous fungi that are naturally found as part of the animal skin microbiome (Theelen et al., 2018). The *Malassezia* genus presently includes 18 species divided into three clades – A, B, and C. These species also have unusually compact genomes of less than 9 Mb, organized into 6 to 9 chromosomes as revealed by electrophoretic karyotyping of some of these species (Boekhout and Bosboom, 1994, Boekhout et al., 1998, Wu et al., 2015). Fungemia-associated species like *Malassezia furfur* belong to Clade A. Clade B includes common inhabitants of human skin that are phylogenetically clustered into two subgroups namely Clade B1 that contains *Malassezia globosa* and *Malassezia restricta* and Clade B2 that contains *Malassezia sympodialis* and related species. Clade C includes *Malassezia slooffiae* and *Malassezia cuniculi* which diverged earlier from a *Malassezia* common ancestor (Wu et al., 2015, Lorch et al., 2018).

Besides humans, *Malassezia* species have been detected on the skin of animals. For example, *M. slooffiae* was isolated from cows and goats, *M. equina* from horses, *M. brasiliensis* and *M. psittaci* from parrots, and a cold-tolerant species *M. vesperilionis* isolated from bats (Lorch et al., 2018, Theelen et al., 2018). Additionally, culture-independent studies of fungi from environmental samples showed that the *Malassezia* species closely realted to those found on human skin were also detected in diverse niches such as deep-sea vents, soil invertebrates, hydrothermal vents, corals, and Antarctic soils (Amend, 2014). More than ten *Malassezia* species have been detected as a part of the human skin microbiome (Findley and Grice, 2014). The human skin commensals such as *M. globosa, M. restricta,* and *M. sympodialis* have been associated with dermatological conditions such as dandruff/seborrheic dermatitis, atopic dermatitis, and folliculitis (Theelen et al., 2018). Recent reports implicate *M. restricta* in conditions such as Crohn's disease and *M. globosa* in the progression of pancreatic cancer (pancreatic ductal adenocarcinoma) (Aykut et al., 2019, Limon et al., 2019). Elevated levels of *Malassezia* species and the resulting inflammatory host response have been implicated in both of these disease states. The nature of genomic rearrangements in each species may influence its ability to adapt and cause disease in a specific host niche. Thus, studying the mechanisms of karyotype evolution is an important step towards understanding the evolution of the *Malassezia* species complex.

Kinetochore proteins serve as useful tools in the identification of centromeres due to their centromere-exclusive localization. CENP-A replaces histone H3 in the centromeric nucleosomes. This has been shown as a reduction in histone H3 levels at the centromeres in *C. lusitaniae* (Kapoor et al., 2015) as well as in a human neocentromere (Lo et al., 2001). These CENP-A nucleosomes act as a foundation to recruit CENP-C, the KMN (KNL1C-MIS12C-NDC80C) network, and other kinetochore proteins (Musacchio and Desai, 2017).

In this study, we experimentally validated all of the eight centromeres of *M. sympodialis* using the Mtw1 protein (Mis12 in *S. pombe*), a subunit of the KMN complex, as the kinetochore marker. The Mis12 complex proteins are evolutionarily conserved outer kinetochore proteins that link the chromatin-associated inner kinetochore proteins to the microtubule-associated outer kinetochore proteins. Members of the Mis12 complex localize to centromeres in other organisms (Goshima et al., 1999, Goshima et al., 2003, Westermann et al., 2003, Roy et al., 2011). Recent studies suggest that the protein domains associated with the Mis12 complex members are exclusive to kinetochore proteins and are not detected in any other proteins making them attractive tools to identify centromere sequences (Tromer et al., 2019). Using the features of centromeres of *M. sympodialis* and newly generated chromosome-level genome assemblies, we predicted centromeres in related *Malassezia* species carrying seven, eight, or nine chromosomes and experimentally validated the centromere identity in representative species of each karyotype, belonging to different *Malassezia* clades. We employed gene synteny conservation across these centromeres to understand the transition between these states from an inferred ancestral state of 9 chromosomes. Based on our results, we propose that centromere loss by two distinct mechanisms drives karyotype diversity.

Results

Chromosome number varies in the *Malassezia* species complex

Previous reports based on pulsed-field gel electrophoresis (PFGE) have suggested that chromosome number varies within the *Malassezia* species complex. The early diverged species *M. slooffiae* of Clade C was reported to have 9 chromosomes (Boekhout et al., 1998). Clade B *Malassezia* species are reported to have 9 (*M. globosa* and *M. restricta*), 8 (*M. sympodialis*), or 6 chromosomes (*M. pachydermatis*). Among the Clade A species, *M. obtusa* and *M. furfur* CBS14141 were reported to have 7 chromosomes each (Boekhout and Bosboom, 1994, Zhu et

al., 2017). A high-quality reference genome is a prerequisite to understanding the rearrangements

associated with chromosome number variation. Additionally, this will also assist in resolving

ambiguities in PFGE-based estimates of chromosome number when similar-sized chromosomes

are present. Complete genome assemblies were not available for many of the species with

reported numbers of chromosomes. To obtain better-assembled reference genomes, we

sequenced the genomes of *M. slooffiae* and *M. globosa* as representatives of the 9-chromosome

state, and *M. furfur* as a representative of the 7-chromosome state using the PacBio SMRT

sequencing technology (Figure 1, Figure 1-figure supplement 1).

The *M. globosa* genome was completely assembled into 9 contigs with telomeres on both

ends (BioSample accession SAMN10720087). We validated these by assigning each band on the

pulsed-field gel with the sizes from the genome assembly and further confirmed these by

chromoblot analysis following PFGE. This analysis shows that chromosome 5 contains the

rDNA locus and migrates higher than the expected size of 902 kb, as a diffuse ensemble of

different sizes along with chromosome 3 (Figure 1A). The assembled genome of *M. slooffiae*

has 14 contigs of which 9 contigs have telomeres on both ends, indicative of 9 chromosomes

(BioSample accession SAMN10720088). Each of the 9 contigs could be assigned to the bands

observed in the pulsed-field gel (Figure 1B). For *M. furfur*, the final genome assembly consisted

of 7 contigs with telomeres on both ends and matched the expected chromosome sizes obtained

from an earlier PFGE analysis of CBS14141 (Figure 1C). The complete genome assembly of *M.*

sympodialis reported earlier is distributed into 8 chromosomes with telomere repeats on both

ends (Figure 1D), and serves as a representative of an 8-chromosome state in this study.

Changes in chromosome number are always associated with birth or loss of centromeres

to stabilize the karyotype in organisms with monocentric chromosomes. To understand the

transitions between these different karyotypic states observed in the *Malassezia* species complex,

we sought to experimentally validate centromeres in species representative of each karyotype.

Kinetochores cluster and localize to the nuclear periphery in *M. sympodialis*

Organisms with point centromeres possess Ndc10, Cep3, and Ctf13 of the CBF3

complex, a cognate protein complex specific to point centromeres. None of these point

centromere-specific proteins could be detected in *M. sympodialis*. However, we could detect

homologs of CENP-A, CENP-C, and most of the outer kinetochore proteins in the genome of *M.*

sympodialis (Figure 2A and Figure 2-figure supplement 1). We functionally expressed an N-

terminally GFP-tagged Mtw1 protein (Protein ID: SHO76526.1) from its native promoter and the

expression of the fusion protein was confirmed by western blotting (Figure 2B). Upon staining

with anti-GFP antibodies and DAPI (4′,6-diamidino-2-phenylindole), we could detect punctate

localization of Mtw1 at the nuclear periphery (Figure 2C) consistent with the clustered

kinetochore localization observed in other yeasts (Goshima et al., 1999, Euskirchen, 2002, Roy

et al., 2011). Live-cell images of MSY001 (GFP-*MTW1*) cells revealed that the kinetochores

(GFP-Mtw1) remained clustered throughout the cell cycle, starting from unbudded G1 cells in

interphase to large budded cells in mitosis (Figure 2D).

Mtw1 is localized to a single region at the GC minima of each *M. sympodialis* chromosome

Having identified Mtw1 as an authentic kinetochore protein, we performed ChIP-

sequencing using the GFP-Mtw1 expressing strain of *M. sympodialis* (MSY001). Mapping the

reads to the reference genome of *M. sympodialis* strain ATCC42132 (Zhu et al., 2017) identified

one significantly enriched locus on each of the eight chromosomes (Figure 3A). The length of

the Mtw1-enriched centromere regions identified from the ChIP-seq analysis ranges between

3167 bp and 5143 bp with an average length of 4165 bp (Table S1). However, the region of

maximum Mtw1 enrichment on each chromosome (based on the number of sequenced reads

aligned) mapped to the intergenic region harboring the GC trough (approximately 1 kb long),

which was previously predicted to be the centromeres of *M. sympodialis* (Figure 3B) (Zhu et al.,

2017). The regions of Mtw1 enrichment span beyond the core centromeres and include active

genes located proximal to these troughs (Figure 3B, Figure 3-figure supplement 1A). However,

these ORFs do not show consensus features such as the orientation of transcription or functional

classification. We validated this enrichment by ChIP-qPCR analysis with primers homologous to

the predicted centromeres compared to a control region distant from the centromere (Figure 3-

figure supplement 1B).

Histone H3 is depleted at the core centromere with active genes at the pericentric regions in

M. sympodialis

The presence of CENP-A nucleosomes at the kinetochore should result in reduced

histone H3 enrichment at the centromeres when compared to a non-centromeric locus. To test

this, we performed ChIP with anti-histone H3 antibodies and analyzed the immunoprecipitated

(IP) DNA by qPCR. As compared to a control ORF region unlinked to the centromere (190 kb

away from *CEN1*), the pericentric regions flanking the core centromere showed a marginal

reduction in histone H3 enrichment that was further reduced at the core that maps to the GC

trough with the highest enrichment of the kinetochore protein Mtw1. That the core centromere

region showing the maximum depletion of histone H3 coincided with the regions most enriched

with Mtw1 further supports that histone H3, in these regions, is possibly replaced by its

centromere-specific variant CENP-A (Figure 3C).

The short regional centromeres of *M. sympodialis* are enriched with a 12 bp long AT-rich

consensus sequence motif

To understand the features of *M. sympodialis* centromeres, we analyzed the centromere

DNA sequences for the presence of consensus motifs or structures such as inverted repeats.

PhyloGibbs-MP (Siddharthan et al., 2005, Siddharthan, 2008) predicted a 12 bp long AT-rich

motif common to all of the centromere sequences of *M. sympodialis* (Figure 3D). We swept the

Position Weight Matrix (PWM) from the PhyloGibbs-MP output across each chromosome of *M.*

sympodialis and counted the number of motif predictions in sliding 500 bp window, sliding by

100 bp at a time. Sites with log-likelihood-ratio (LLR) of >7.5 were counted as motif predictions.

The LLR is the natural logarithm of the ratio of the likelihood of the 12 bp substring arising as a

sample from the PWM, to the likelihood of it being generic "background". In each case, the

global peak coincides with the centromere (Figure 3-figure supplement 2A). In each

chromosome, the centromere region shows between 7 and 13 motif matches, while no other 500

bp window shows more than 3 matches. This suggests that the AT-rich motif is more enriched at

the centromeres than at any other region in the *M. sympodialis* genome (Figure 3-figure

supplement 2A). To ensure that this is not an artifact of the GC-poor nature of the centromere,

we repeated the analysis with a synthetic shuffled PWM, created by scrambling the order of the

columns of the original PWM (that is, scrambling the positions in the motif while keeping the

corresponding weight vectors the same). This shuffled motif showed more matches in the

centromeres than what is seen in the non-centromeric genomic sequence, but significantly fewer

than the authentic centromeric sequences in most chromosomes (Figure 3-figure supplement 2B).

A dot-plot analysis was performed to detect the presence of any direct or inverted-repeat

structure associated with the centromeres in *M. sympodialis*. Analysis of all of the centromere

sequences along with 5 kb flanking sequences using SyMap confirmed the lack of direct/inverted

repeat structures (Figure 3-figure supplement 2C).

In the absence of any centromere exclusive DNA sequence, the unique and distinguishing

features of centromere regions in *M. sympodialis* are an AT-rich core region of <1 kb (average

AT content of 78% as compared to the genome average of 41.5%) enriched with the 12 bp motif

(Figure 3-figure supplement 3) in a kinetochore protein-bound region of 3 to 5 kb. As expected,

the kinetochore bound region contains a reduced level of histone H3.

Centromeres in *M. furfur, M. slooffiae,* and *M. globosa* map to chromosomal GC minima

Using the unique centromere features identified in *M. sympodialis*, we predicted one

centromere locus on each of the 7 *M. furfur* chromosomes and these all map to chromosomal GC

troughs (Figure 4-figure supplement 1A, Table S2). Similar to *M. furfur*, we predicted the

centromeres in *M. slooffiae, M. globosa,* and *M. restricta*, each of which contains 9

chromosomes (Figure 4-figure supplement 1B-D, Table S2). Each of the predicted centromeres

showed enrichment of the 12 bp AT-rich motif identified with *M. sympodialis* centromere

sequences as compared to other regions in the genomes (Figure 4-figure supplement 2).

To experimentally validate the centromeric loci in *M. furfur*, we functionally expressed

the centromeric histone H3 variant CENP-A with a 3xFLAG tag at the C-terminus (Figure 4A).

We performed ChIP in strain MF001 (CENP-A-3xFLAG) and analyzed immunoprecipitated

DNA by qPCR using primers specific to each of the seven predicted centromeres and a

centromere-unlinked control locus 1.3 Mb away from *CEN1*. Enrichment of CENP-A at all

seven centromeres over the control locus confirmed that the regions predicted are indeed

centromeres in *M. furfur* CBS14141 (Figure 4B, C).

Given the lack of genetic manipulation methods for *M. slooffiae* and *M. globosa*, we

tested the enrichment of histone H3 at the predicted centromeres in these two species. All of the

nine centromeric loci in these two species showed a reduced histone H3 level when compared to

a control locus unlinked to centromeres (Figure 4D, E). Furthermore, upon analyzing the

enrichment profile at one centromere (*CEN1*) in *M. slooffiae*, we observed a reduction in the

enrichment levels of histone H3 at the GC troughs as compared to the flanking regions (Figure

4F). In the case of *M. globosa*, the regions spanning a centromere (*CEN2*) also depicted a similar

reduction in the histone H3 levels (Figure 4G). Taken together, the significant reduction in the

histone H3 levels at the predicted centromeres, indicative of the presence of CENP-A, suggests

that these putative centromere regions are indeed *bona fide* centromeres in these species.

Centromere loss by breakage resulted in chromosome number reduction in *M. sympodialis*

Synteny of genes across centromeres is largely conserved in closely related species

(Byrne and Wolfe, 2005, Padmanabhan et al., 2008, Yadav et al., 2018b). To understand the

transition between different chromosome number states, we analyzed the conservation of gene

synteny across centromeres in these species. By mapping gene synteny at the centromeres of *M.*

globosa and *M. slooffiae* (each carries 9 chromosomes), compared with *M. sympodialis*

(containing 8 chromosomes), we found complete gene synteny conservation in 8 of the 9

centromeres (Figure 5A, B). Thus, syntenic regions of all 8 *M. sympodialis* centromeres are

present in the genomes of *M. globosa* and *M. slooffiae*. In the case of *M. restricta*, 7 putative

centromeres are completely syntenic with *M. sympodialis* centromeres and one centromere

retained partial gene synteny (Table S3). However, no gene synteny conservation was observed

at the centromeres of Chr2 in *M. globosa,* Chr5 in *M. slooffiae,* or Chr8 in *M. restricta* (Table

S3), indicative of loss of a centromere during the transition from the 9-chromosome state to the

8-chromosome state.

The GC trough corresponding to *MgCEN2/MslCEN5* is flanked by genes that map to

MsyChr2 on one arm and MsyChr4 on the other (Figure 5C, D). The centromere region in each

of MgChr2 and MslChr5 marks a synteny breakpoint showing no homologous region in the *M.*

sympodialis genome indicative of a loss of this centromere DNA sequence. We also observed

that the genes flanking the breakpoint are conserved in *M. sympodialis*, suggesting that the

specific intergenic region was involved (Figure 5E, F). Evidence for internalization of telomere

adjacent ORFs or the presence of interstitial telomere repeats indicative of telomere-telomere

fusions was not detected in the *M. sympodialis* genome. These observations strongly support our

hypothesis that breakage of *MgCEN2/MslCEN5* (or the orthologous ancestral *CEN*) and fusion of

the two acentric arms to other chromosomes resulted in the chromosome number reduction

observed between these species.

Centromere inactivation by sequence divergence and loss of AT-richness resulted in chromosome number reduction in *M. furfur*

To understand the basis of the change in chromosome number from 9 to 7 in *Malassezia* species, we compared the synteny of ORFs flanking the *M. slooffiae* or *M. globosa* centromeres with that of *M. furfur*. Of the 9 centromeres in *M. slooffiae*, three centromeres belonged to conserved gene synteny blocks and four other centromeres showed partial gene synteny conservation in *M. furfur* (Figure 6A, Table S3). A similar pattern of gene synteny conservation was observed between *M. globosa* and *M. furfur* (Figure 6B, Table S3). The genes flanking the remaining two centromeres (*CEN8* and *CEN9*) in *M. slooffiae* were present in conserved gene synteny blocks in the two arms of MfChr3 (Figure 6C). However, the regions corresponding to *CEN8* and *CEN9* in *M. slooffiae* appear to have evolved to decreased AT-richness in *M. furfur*. A similar centromere inactivation mechanism was observed when *CEN8* and *CEN9* of *M. globosa* were compared to the corresponding syntenic regions in *M. furfur* (Figure 6D). These results are suggestive of centromere inactivation by changes in the centromeric DNA sequence in this species (Figure 6-figure supplement 1).

The common ancestral *Malassezia* species contained 9 chromosomes

To trace the ancestral karyotype in *Malassezia*, we predicted centromeres and inferred chromosome numbers in other species of clades A and B based on GC troughs and gene synteny. We identified putative centromeres in *M. dermatis* and *M. nana* in Clade B because of their relatively better-assembled genomes distributed in 18 and 13 scaffolds respectively. Of these, we could predict 8 centromeric regions marked by GC troughs that were also enriched with the 12 bp motif in each species (Figure 7-figure supplement 1A-B, Figure 7-figure supplement 2A-B and Table S2). Furthermore, in both of these species, the 8 putative centromeres shared complete gene synteny conservation with the regions spanning *M. sympodialis* centromeres, indicating that their common ancestor had 8 chromosomes (green circle in Figure 7A, Figure 7-figure supplement 3 and Table S3). To map the common ancestor in Clade B *Malassezia* species, we analyzed regions flanking centromeres of Chr2 of *M. globosa* and Chr8 of *M. restricta,* both of which mapped to the gene synteny breakpoint of the genome of the *Malassezia* species with 8 chromosomes suggesting their common ancestor, named as Ancestor B (Anc. B), also had 9 chromosomes (Figure 7A, Figure 7-figure supplement 3). Based on our centromere predictions

in Clade B species and synteny analysis, we propose that centromere breakage would have

occurred in the common ancestor of *M. sympodialis*, *M. nana,* and *M. dermatis* after divergence

from the common ancestor of *M. globosa* and *M. restricta* which retained a 9-chromosome

configuration (Figure 7A, B).

As mentioned earlier, *M. furfur* and *M. obtusa* of Clade A contain 7 chromosomes each

(Boekhout and Bosboom, 1994, Boekhout et al., 1998). To further understand the karyotype

variations within this clade, we predicted the chromosome number in *Malassezia vespertilionis*

and *Malassezia japonica* as their genomes are relatively well assembled (Sugita et al., 2003,

Lorch et al., 2018). In both of these species, we could predict 9 GC troughs indicative of

centromeres of 9 chromosomes (Figure 7-figure supplement 1C-D, Table S2). In the case of *M.

japonica*, all of the predicted centromeres (except the centromere of scaffold 7) showed

enrichment of the 12 bp motif (Figure 7-figure supplement 2C). However, the 12 bp motif was

found to be enriched in all of the predicted centromeres of *M. vespertilionis* (Figure 7-figure

supplement 2D). The presence of species with 9 chromosomes in Clade A suggests that the

ancestral state in this clade, named Ancestor A (Anc. A) also contained 9 chromosomes (Figure

7A).

We identified 9 centromeres in *M. slooffiae*, the only species in Clade C with a well-

assembled genome. The presence of species with 9 chromosomes in each of the three clades of

Malassezia species, the conservation of gene synteny across orthologous centromeres, and the

similar centromere features shared by all 9 species analyzed in this study suggest that *Malassezia*

species diverged from a common ancestor that had 9 chromosomes with short regional

centromeres enriched with the 12 bp AT-rich DNA sequence motif.

Discussion

In this study, we experimentally validated the chromosome number in *M. slooffiae* and

M. globosa by PFGE analysis. We sequenced and assembled the genomes of *M. slooffiae*, *M.

globosa*, and *M. furfur* and compared each one with the genome of *M. sympodialis* to understand

the karyotype differences observed in members of the *Malassezia* species complex. These

species represent each of the three major clades of *Malassezia* species with chromosome

numbers ranging from 7 to 9. Because centromere loss or gain directly influences the

chromosome number of a given species, we experimentally identified the centromeres of these

representative species to understand the mechanisms of karyotype diversity. Kinetochore
proteins are useful tools in identifying the centromeres of an organism. Localization of the
evolutionarily conserved kinetochore protein Mtw1 tagged with GFP suggested that kinetochores
are clustered throughout the cell cycle in *M. sympodialis*. ChIP-sequencing analysis identified
short regional (< 5 kb long) centromeres in *M. sympodialis* that are depleted of histone H3, and
are enriched with an AT-rich sequence motif. Identification of centromeres in *M. slooffiae*, *M.*
globosa, and *M. furfur* further suggested that centromere properties are conserved across these
Malassezia species. By predicting putative centromere in five other species along with four
species with experimentally mapped centromeres described above across three clades of
Malassezia, we concluded that an AT-rich centromere core of < 1 kb in length enriched with the
12 bp sequence motif is a potential signature of centromeres in nine *Malassezia* species analyzed
in this study. Comparative genomics analysis revealed two major mechanisms of centromere
inactivation resulting in karyotype change. The presence of a 9-chromosome state in each of the
three clades along with conserved centromere features and conserved gene synteny across these
centromeres, helped us to infer that the ancestral *Malassezia* species had 9 chromosomes with
short regional centromeres with an AT-rich core enriched with the 12 bp sequence motif.

Centromeres in the *Malassezia* species complex represent the first example of short
regional centromeres in basidiomycetes. Centromeres reported in other basidiomycetes, such as
those in *Cryptococcus* and *Ustilago* species, are of the large regional type (Yadav et al., 2018b).
Malassezia species analyzed in this study have a significantly smaller genome (< 9 Mb) as
compared to other basidiomycetes and lack RNAi machinery. The occurrence of short regional
centromeres in RNAi-deficient *Malassezia* species is in line with a previous finding wherein a
reduction in centromere size was observed in RNAi-deficient basidiomycete species as compared
to their RNAi-proficient relatives (Yadav et al., 2018b). With the presence of clustered
kinetochores across cell cycle stages, and the absence of key genes encoding the RNAi
machinery, these *Malassezia* species resemble ascomycetes such as many of the CTG clade
species with short regional centromeres (Sanyal et al., 2004, Nakayashiki et al., 2006,
Padmanabhan et al., 2008, Kapoor et al., 2015, Chatterjee et al., 2016, Yadav et al., 2018a),
rather than the basidiomycetes with large regional centromeres. By combining these features, we
conclude that the genome size and the presence of complete RNAi machinery could be the
determinants of the centromere type of a species irrespective of the phylum it belongs to.

Based on the binding pattern of the kinetochore protein across the *M. sympodialis*

genome, the 3 to 5 kb long region can be divided into two domains: (a) an AT-rich *CEN* core that

maps to the intergenic region containing the GC trough showing maximum kinetochore binding

(< 1 kb) and (b) the regions flanking the core that show basal levels of kinetochore protein

binding. We observed conservation of the 12 bp AT-rich motif in the centromere core across the

nine *Malassezia* species. It should also be noted that the 12 bp motif is significantly enriched at

the centromeres but not exclusive to the centromeres as it is detected across the chromosomes.

We did not observe any orientation bias of these motifs at the centromeres. The functional

significance of the frequent occurrence of this motif at centromeres is unknown. It will be

intriguing to test the roles played by this motif, the core, and the flanking sequences in

centromere function. Testing these domains for centromere function *in vivo* by making

centromeric plasmids in various *Malassezia* species is challenging at present due to technical

limitations. Other than *M. sympodialis*, *M. pachydermatis,* and *M. furfur*, no other *Malassezia*

species have been successfully transformed (Ianiri et al., 2016, Celis et al., 2017, Ianiri et al.,

2017b). Moreover, in all of these cases, the genetic manipulations are performed by

Agrobacterium-mediated transconjugation, which is not amenable for the introduction of circular

plasmids.

The centromeres in *M. sympodialis* contain transcribed ORFs, as documented in

centromeres of rice, maize, and *Zymoseptoria tritici* (Nagaki et al., 2004, Wang et al., 2014). In

contrast to these cases, our read count analysis did not reveal any significant difference in the

transcription (RPKM values) of centromere associated ORFs to that elsewhere in the genome of

M. sympodialis. We posit that the 12 bp AT-rich motif sequences could facilitate the

transcription of these genes by recruiting transcription factors with a possible role in kinetochore

assembly. Binding of Cbf1 at CDEI in *S. cerevisiae* centromeres and Ams2 at the central core

sequences of *S. pombe* centromeres are classic examples of transcription factors facilitating

kinetochore stability in fungal systems (Hemmerich et al., 2000, Chen et al., 2003). A fine

regulation of transcription by Cbf1, Ste12, and Htz1 and the resulting low-level cenRNAs have

been implicated in proper chromosome segregation in budding yeast (Ohkuni and Kitagawa,

2011, Ling and Yuen, 2019). These studies reinforce the role of transcription in centromere

function irrespective of centromere structure.

In this study, we report three high-quality chromosome-level genome assemblies and

identified centromeres in nine *Malassezia* species, representing all of the three *Malassezia* clades

with differing numbers of chromosomes. This will serve as a rich resource for comparative

genomics in the context of niche adaptation and speciation. Analysis of gene synteny

conservation across centromeres using these genomes revealed breakage at the centromere as one

of the mechanisms resulting in a karyotype change between closely related species - those with 9

chromosomes, such as *M. slooffiae* and *M. globosa,* and those with 8 chromosomes like *M.*

sympodialis. Gene synteny breakpoints adjacent to the centromeres have been reported in *C.*

tropicalis that has seven chromosomes - one less than *C. albicans* (Chatterjee et al., 2016).

Centromere loss by breakage was proposed to have reduced the *Ashbya gossypii* karyotype by

one when compared to the pre-whole genome duplication ancestor (Gordon et al., 2011).

Breakpoints of conserved gene synteny between mammalian and chicken chromosomes were

also mapped to the centromeres (International Chicken Genome Sequencing, 2004). Similar

consequences in the karyotype have been reported in cases where centromeres were

experimentally excised. Besides neocentromere formation, survival by fusion of acentric

chromosome arms has been shown in *S. pombe* (Ishii et al., 2008). Such fusions are detected

upon deletion of centromeres in another basidiomycete, *C. deuterogattii* (Schotanus and

Heitman, 2019). By comparing the ancestral state (*M. slooffiae*) and other *Malassezia* species

with either the same number or fewer chromosomes, we observed gene synteny breaks adjacent

to centromeres (indicated by partial synteny conservation) apart from the break observed at

MgCEN2 or *MslCEN5*. Is this suggestive of a fragile nature of *Malassezia* centromeres? We

advance the following hypothesis to explain the observed breaks at centromeres.

Studies of the common fragile sites in the human genome suggest different forms of

replication stress as a major source of instability and subsequent breakage at these sites

(Helmrich et al., 2011, Letessier et al., 2011, Ozeri-Galai et al., 2011). The resolution of the

resulting replication fork stall has been shown to be critical for the stability of these fragile sites

(Schwartz et al., 2005). Studies of the human fragile site FRA16D show that the AT-rich DNA

(Flex1) results in fork stalling as a consequence of cruciform or secondary structure formation

(Zhang and Freudenreich, 2007). Centromeres are natural replication fork stall sites in the

genome (Greenfeder and Newlon, 1992, Smith et al., 1995, Mitra et al., 2014). The AT-rich core

centromere sequence in *M. globosa* is also predicted to form secondary structures (Figure 7-

figure supplement 4), which can be facilitated by the inherent replication fork stall at the

centromeres. Whenever these secondary structures are unresolved and the fork restart fails, DSBs

can occur at the centromeres. Chromosomal breakage and aneuploidy resulting from such defects

are known in cancers (Kops et al., 2005). In mammals, centromeric DSBs are repaired efficiently

compared to regions elsewhere in the genome, largely due to the presence of several homology

tracts in the form of repetitive DNA sequences and the stiffness provided by the inherent

heterochromatic state to facilitate ligation (Rief and Lobrich, 2002). *Malassezia* species are

haploid in nature and lack typical pericentric heterochromatin marks. While the efficiency of

centromeric DSB repair in the absence of long tracts of homologous sequences is not known in

this species complex, we propose that the AT-rich core sequences, by virtue of secondary

structure formation during DNA replication, could occasionally undergo DNA breakage at the

centromere in *Malassezia* species.

The second mechanism of chromosome number reduction based on our comparative

analyses of the *M. furfur* genome with genomes of *M. slooffiae* or *M. globosa* involves the

inactivation of centromeres in the process of transition from a 9-chromosome state to a 7-

chromosome state. Centromere inactivation occurs in cases involving the fusion of centric

chromosomal fragments, stabilizing the fusion product and generating a functionally

monocentric chromosome, as seen in case of the origin of human Chr2 from the shared ancestor

with the great apes (Yunis and Prakash, 1982, Jdo et al., 1991). A larger proportion of known

centromere inactivation events were shown to be mediated by epigenetic modification wherein

inactivated centromeres are enriched with marks such as H3K9me2/3, H3K27me2/3, or DNA

methylation, emphasizing the role of heterochromatin in this process (Zhang et al., 2010, Koo et

al., 2011, Sato et al., 2012). Deletion of the centromere sequence corresponding to kinetochore

binding has also been reported as an alternate mechanism, albeit less frequent, in both humans

and in yeasts (Stimpson et al., 2010, Gordon et al., 2011, Sato et al., 2012). Unlike the above two

modes, we observed divergence in the sequences corresponding to the inactivated centromeres

(*CEN8* and *CEN9* of both *M. slooffiae* and *M. globosa*) in the arms of *M. furfur* Chr3, resulting

in the loss of AT-richness of these centromere core regions (Figure 7C). This is also suggestive

of a functional role for AT-rich DNA in centromere function in these species.

A change in chromosome number between two closely related species like *C. albicans* and *C. tropicalis* is associated with a change in centromere structure, between unique short regional centromeres in *C. albicans* that are epigenetically regulated, as compared to a genetically defined homogenized inverted repeat-associated centromere in *C. tropicalis* (Chatterjee et al., 2016). Strikingly, in both the transitions described above for *Malassezia* species, we did not observe any change in the centromere structure. The emergence of evolutionarily new centromeres as seen in the case of primates was not detected in *Malassezia* species (Rocchi et al., 2009, Kalitsis and Choo, 2012). This is particularly striking in the absence of conservation of any specific centromere-exclusive DNA sequence. This suggests that a strong driving force helps maintain the highly conserved centromere properties in closely related *Malassezia* species descended from their common ancestor, even after extensive chromosomal rearrangements involving centromeres that might have driven speciation. Furthermore, centromere inactivation/loss of centromere function seems to be a conserved theme mediating chromosome number variation from unicellular yeast species to metazoans, including primates.

Materials and methods

The reagents, strains, plasmids, and primers used in this study are listed in the *SI appendix* (Tables S4-S7). *Malassezia* strains were grown on modified Dixon's media (Malt extract 36 g/L, Desiccated Ox-bile 20 g/L, Tween40 10 mL/L, Peptone 6 g/L, Glycerol 2 mL/L, Oleic acid 2.89 mL/L). *M. sympodialis, M. furfur* strains were grown at 30°C. Cultures of *M. globosa* and *M. slooffiae* were grown at 32°C.

Construction of the *M. sympodialis* strain expressing GFP-Mtw1

The allele for N-terminal tagging of Mtw1 with GFP was prepared by gap repair in the *Saccharomyces cerevisiae* BY4741 strain (Eckert-Boulet et al., 2012). Briefly, a 1.6 kb fragment consisting of the upstream and promoter sequence of the *MTW1* gene and a 1.6 kb fragment having the *MTW1* ORF (Protein ID: SHO76526) along with the downstream sequence was amplified from *M. sympodialis* genomic DNA. The GFP ORF (without the stop codon) and NAT were amplified from plasmids pVY7 and pAIM1 respectively (Yadav et al., 2018b). *S. cerevisiae* was transformed with all four fragments and the linearized plasmid pGI3 (digested with KpnI and BamHI) and the epitope-tagged allele were assembled in an ordered way by gap repair. Total DNA was isolated from *S. cerevisiae* and *E. coli* DH5α strain was transformed. The pGFP-Mtw1

526 construct was screened by restriction digestion and further confirmed by sequencing. The pGFP-
527 Mtw1 construct was used to transform *M. sympodialis* strain ATCC42132 by *Agrobacterium*
528 *tumefaciens*-mediated transconjugation (Ianiri et al., 2016, Ianiri et al., 2017a).

Construction of the *M. furfur* strain expressing CENP-A FLAG

The allele for C-terminal tagging of CENP-A with a 3xFLAG epitope tag was prepared
by gap repair in the *Saccharomyces cerevisiae* BY4741 strain (Eckert-Boulet et al., 2012).
Briefly, a 1 kb fragment consisting of the upstream and promoter sequence of the CENP-A gene
of *M. furfur* including the ORF (CENP-A ORF coordinates in Chr1: 1,453,468- 1,453,921) and 1
kb fragment containing the sequence downstream of CENP-A ORF were amplified from *M.*
furfur genomic DNA. The 3xFLAG tag was introduced in the reverse primer annealing to the
CENP-A ORF. The NAT marker was amplified from plasmid pAIM1 as above. *S. cerevisiae*
was transformed with all three fragments and plasmid pGI3 (digested with KpnI and BamHI) and
the epitope-tagged allele was assembled in an ordered way by gap repair. Total DNA was
isolated from *S. cerevisiae* and *E. coli* DH5α strain was transformed. The resulting pMF1
construct was screened by restriction digestion and further confirmed by sequencing. The pMF1
construct was used to transform *M. furfur* strain CBS14141 by *Agrobacterium tumefaciens*-
mediated transconjugation (Ianiri et al., 2016, Ianiri et al., 2017a) to obtain the epitope-tagged
strain MF001.

Microscopic imaging of live cells and processing

The GFP-Mtw1 strain was inoculated to 1% v/v from a saturated starter culture
grown in mDixon medium. After growth for 6 h at 30°C/ 150 rpm, these cells were pelleted
at 4,000 rpm and washed 3 times with 1x phosphate-buffered saline (PBS) and the cell
suspension was placed on a clean glass slide. A coverslip was placed on the spot and sealed
prior to imaging. The images were acquired at room temperature using a laser scanning
inverted confocal microscope LSM 880-Airyscan (ZEISS, Plan Apochromat 63x, NA oil 1.4)
equipped with highly sensitive photo-detectors. The filters used were GFP/FITC 488
excitation and GFP/FITC 500/550 band pass, long pass for emission. Z-stack images were
taken at every 0.3 μm and processed using ZEISS Zen software or ImageJ. All of the images
were processed post-acquisition with minimal adjustments to levels and linear contrast until
the signals were highlighted.

Preparation of spheroplasts

Cells were grown on mDixon's medium and washed with water by centrifugation at 4000 rpm for 5 min. Cells were resuspended in 10 mL of 5% (v/v) 2-mercaptoethanol solution in water and incubated at 30°C/ 150 rpm for 45 min. The cells were pelleted, washed, and resuspended in 3 mL spheroplasting buffer (40 mM Citric acid, 120 mM Na_2HPO_4 and 1.2 M Sorbitol) for every 1.5×10^9 cells. Cell clumps were dissociated by mild sonication for 30 s using the medium intensity setting in a Bioruptor (Diagenode). Lysing enzymes from *Trichoderma harzianum* (Sigma) and Zymolyase-20T (MP Biomedicals) were added at 20 mg/mL and 100 µg/mL respectively. The spheroplasting suspension was incubated at 30°C/ 65 rpm for 6 to 8 h. The suspension was examined under a microscope to estimate the proportion of spheroplasts. Spheroplasts were washed with ice-cold 1xPBS and used as per the experimental design (adapted from (Boekhout, 2003)).

Indirect Immunofluorescence

The GFP-Mtw1 strain was inoculated to 1% (v/v) from a saturated starter culture grown in mDixon medium. After growth for 6 h, the cells were fixed by the addition of formaldehyde to a final concentration of 3.7% for 1 h. Post-fixing, the cells were washed with water and taken for preparation of spheroplasts (described above). Spheroplasts were washed with ice-cold 1xPBS and resuspended in ice-cold 1xPBS to a cell density suitable for microscopy. Slides for microscopy were washed with water and coated with poly L-Lysine (15 µL of 10 mg/mL solution per well) for 5 min at room temperature. The solution was aspirated and washed once with water. The cell suspension was added to each well (15-20 µL) and allowed to stand at room temperature for 5 min. The cell suspension was aspirated and the slides were washed once with water to remove unbound cells. The slides were fixed in ice-cold methanol for 6 min followed by treatment with ice-cold acetone for 30 s. Post fixing, blocking solution (2% non-fat skim milk in 1xPBS) was added to each well, incubated at room temperature for 30 min. After this, the blocking solution was aspirated and primary antibodies were added (mouse anti-GFP antibodies [Sigma] at 1:100 dilution). After incubation for 1 h at room temperature, each slide was washed 8 times with 1xPBS giving a 2 min incubation for every wash. Secondary antibody solution (goat anti-mouse-AlexaFluor488 [Invitrogen] at 1:500 dilution) was added to each well and incubated for 1 h

in the dark at room temperature. Post-incubation, slides were washed as described above.

Mounting medium (DAPI at 100 ng/mL in 70% glycerol) was added, incubated for 5 min and

aspirated out. Slides were sealed with a clean coverslip and proceeded for imaging. The

images were acquired at room temperature using an inverted fluorescence microscope

(ZEISS Axio Observer, Plan Apochromat 100x, NA oil 1.4). Z- stack images were taken at

every 0.3 μm and processed using ZEISS Zen software/ ImageJ.

Chromatin immunoprecipitation (ChIP)

The ChIP protocol was adapted from the one implemented for *C. neoformans* (Yadav

et al., 2018b). Logarithmically grown cells were fixed with formaldehyde at a final

concentration of 1% for 30 min (for Mtw1 ChIP) and 15 min (for CENP-A, histone H3, and

histone H4 ChIP) respectively. The reaction was quenched by the addition of glycine to a

final concentration of 0.135 M. Cells were pelleted and processed for spheroplasting as

described above. Spheroplasts were washed once sequentially using 10 mL of the following

ice-cold buffers: 1xPBS, Buffer-I (0.25% Triton X-100, 10 mM EDTA, 0.5 mM EGTA, 10

mM Na-HEPES pH=6.5), Buffer-II (200 mM NaCl, 1 mM EDTA, 0.5 mM EGTA, 10 mM

Na-HEPES pH=6.5). The pellet after the final wash was resuspended in 1 mL lysis buffer (50

mM HEPES pH=7.4, 1% Triton X-100, 140 mM NaCl, 0.1% Na-deoxycholate, 1 mM

EDTA) for every 1.5×10^9 cells. Protease inhibitor cocktail was added to 1x final

concentration.

The resuspended spheroplasts were sonicated with a Bioruptor (Diagenode) using 30

s ON/OFF pulse at high-intensity mode with intermittent incubation on ice to obtain

chromatin fragments in the size range of 100-300 bp. The lysate was cleared after sonication

by centrifugation at 13,000 rpm for 10 min at 4ºC. The input DNA fraction was separated at

this step ($1/10^{th}$ volume of lysate) and processed for de-crosslinking by the addition of 400

μL elution buffer (0.1 M NaHCO₃, 1% SDS) per 100 μL lysate (processing for de-

crosslinking mentioned below). The remaining lysate was split equally and processed as IP

and control samples. For GFP-Mtw1 ChIP, 20 μL GFP-trap beads and blocked agarose beads

respectively were used for IP and control. For CENP-A-3xFLAG ChIP, 20 μL anti-FLAG

affinity gel and blocked agarose beads respectively were used for IP and control. In the case

of histone H3 or histone H4 ChIP, 5 μL of antibodies were used per IP fraction along with 20

µL Protein-A sepharose beads. Samples were rotated for 6 h at 4ºC. Post incubation, samples

were sequentially washed as follows: twice with 1 mL low salt wash buffer (0.1% SDS, 1%

Triton X-100, 2 mM EDTA, 20 mM Tris pH=8.0, 150 mM NaCl), twice with 1 mL high salt

wash buffer (0.1% SDS, 1% Triton X-100, 2 mM EDTA, 20 mM Tris pH=8.0, 500 mM

NaCl), once with 1 mL LiCl wash buffer (0.25 M LiCl, 1% NP-40, 1% Na-deoxycholate, 1

mM EDTA, 10 mM Tris pH=8.0) and twice with 1 mL 1xTE (10 mM Tris pH=8.0, 1 mM

EDTA). Samples were rotated in a rotaspin for 5 min at room temperature for every wash (15

min in case of histone H3 or histone H4 ChIP). After washes, DNA was eluted from the

beads twice using 250 µL elution buffer. The samples for elution were incubated at 65ºC for

5 min, rotated for 15 min and then collected by centrifugation.

Samples were decrosslinked by the addition of 20 µL 5 M NaCl and incubation at

65ºC for 6 h. Following this, samples were deproteinized by the addition of 10 µL 0.5 M

EDTA, 20 µL 1 M Tris pH=6.8, 2 µL Proteinase K (20 mg/L) and incubation at 45ºC for 2 h.

After incubation, samples were treated with an equal volume of phenol-chloroform-isoamyl

alcohol (25:24:1) mix, and the aqueous phase was extracted by centrifugation. DNA was

precipitated by addition of $1/10^{th}$ volume of 3 M Na-acetate, 1 µL glycogen (20 mg/mL), 1

mL absolute ethanol and incubation at -20ºC for at least 12 h. Finally, the samples were

harvested by centrifugation at 13,000 rpm for 45 min at 4ºC followed by washing the pellet

once with ice-cold 70% ethanol. Air-dried pellets were then resuspended in 20 µL sterile

MilliQ water with 10 µg/mL RNAse. Samples were either processed for library preparation

for ChIP-sequencing or analyzed by qPCR with the primers listed in Table S7.

Analysis of sequencing data

GFP-Mtw1 ChIP sequencing was performed at the Clevergene Biocorp. Pvt. Ltd.,

Bengaluru, India. A total of 46,704,720 and 63,524,912 150 bp paired-end reads were

obtained for IP and Input samples, respectively. The reads were mapped to *M. sympodialis*

ATCC42132 genome using Geneious 9.0 (http://www.geneious.com/) with default

conditions. Each read was allowed to map only once randomly anywhere in the genome. The

alignments were exported to BAM files, sorted, and visualized using Integrative Genomics

Viewer (IGV, Broad Institute). The images from IGV were imported into Adobe Photoshop

(Adobe) and scaled for representation purposes. RNA-sequencing data (E-MTAB-4589)

from a previous study was downloaded from the ArrayExpress website, sorted, and

visualized using IGV. GC-content was calculated using Geneious 9.0 with a sliding window

size of 250 bp. The data was exported as wig files and further visualized using IGV.

Western Blotting

Protein lysates for western blot were prepared by the TCA method. One mL overnight

grown cultures were harvested, washed, and resuspended in 400 μL of 12.5% ice-cold TCA

solution. The suspension was vortexed briefly and stored at – 20°C for 4 to 6 h. The suspension

was thawed on ice, pelleted at 14,000 rpm for 10 min, and washed twice with 350 μL of 80%

Acetone (ice-cold). The washed pellets were air-dried completely and resuspended in the desired

volume of lysis buffer (0.1 N NaOH+1% SDS). Samples were separated in 12% polyacrylamide

gels and transferred onto nitrocellulose membranes. For probing, mouse anti-GFP antibody

(Roche) and the HRP conjugated goat anti-mouse secondary antibody (Bangalore Genei), were

used at 1:3000 and 1:5000 dilution respectively in 2.5% skim milk powder in 1xPBS. The blots

were developed using Chemiluminescence Ultra substrate (BioRad) and imaged using the

VersaDoc imaging system (BioRad).

PFGE analysis for *M. globosa* and *M. slooffiae*

For CHEF analysis of *M. globosa* (CBS7966) and *M. slooffiae* (CBS7956), the cells were

grown on solid mDixon medium and then collected and resuspended in 1xPBS. CHEF plugs

were prepared as described in previous studies (Sun et al., 2014, Sun et al., 2017). Chromosomes

were separated in 1% Megabase certified agarose gels made with 0.5×TBE, using a BioRad

CHEF-DR II System running at 3.2 V/cm with linear ramping switching time from 90 to 360 s

for 120 h in 0.5xTBE at 14°C. The gel was stained with EtBr and visualized under UV.

For the chromoblot analyses of *M. globosa* (CBS7966), the gel from the CHEF

analysis was first transferred to a membrane, and the resulting chromoblots were then

hybridized sequentially with four probes from the chromosomes 3, 4, 5, and 6 of the

CBS7966 genome assembly, respectively (see the Table S7 for the primer information), as

described in previous studies (Findley et al., 2012, Yadav et al., 2018b).

M. globosa genome assembly

Sequence reads were assembled with HGAP3 included in SMRTPortal v2.3 (PacBio, Menlo Park, CA, USA) and default parameters except for the genome size set to 9 Mb. Assembly completeness was evaluated by checking for telomeric repeats. Non-telomeric contig-ends were aligned to other contigs using BLAST and unique overlaps used to build complete chromosomes. Short telomere ends were extended using uniquely mapping reads longer than 10 kb and repolishing of the assembly using the resequencing pipeline in SMRTPortal v2.3. The assembly resulted in 19 contigs, with a total length of 9.2 Mb. 17 long and 1 short telomere could be identified. 6 contigs had telomeres on both the 5'- and the 3'-end and thus represent full-length chromosomes (Chr1, 2, 3, 6, 7, and 8). 6 contigs had only one telomere and 7 contigs had no telomeric sequence. 2 contigs without telomeres were from the mitochondrion and 2 were from the ribosomal repeats. Chromosome 5 was constructed from 2 contigs ending in ribosomal rDNA repeats. The assembly contains 6 copies of the repeat, but read coverage suggests a length of 30-40 repeat units that cannot be resolved with the available read-length. The remaining contigs were used to build chromosomes 4 and 9 that share highly similar 5'-ends. The two ends can be distinguished by two microsatellite expansions. Chromosome 4 had a very short 3'-telomere from the default assembly, but the raw data contained a uniquely mapping read that extended several repeat units past the assembly end. After polishing the reference, all 9 chromosomes had clear 5'- and 3'- telomeres.

M. slooffiae genome assembly

Sequence reads were assembled using HGAP3 included in SMRTPortal v2.3 (PacBio, Menlo Park, CA, USA) with default parameters. This resulted in an assembly with 14 scaffolds with telomeric repeats at both ends in 9 contigs. Of the remaining five contigs, three of them could be assigned to mitochondrial DNA based on BLAST analysis with *M. globosa*. The remaining two contigs of sized 5.8 kb and 2.3 kb respectively did not show BLAST hits against *M. globosa* or *M. sympodialis* genomes.

M. furfur genome assembly

Sequence reads generated from CBS14141 were assembled using HGAP3 included in SMRTPortal v2.3 (PacBio, Menlo Park, CA, USA) with default parameters. This resulted in an

703   assembly with 8 scaffolds of which the nuclear genome was organized in 7 scaffolds having

704   telomere repeats at both ends.

Gene synteny analysis

For gene synteny conservation across the centromeres, the analysis was performed by

BLAST as follows. The genomes for *M. restricta*, *M. nana,* and *M. dermatis* were downloaded

from the NCBI genomes portal. The PacBio assembled genomes of *M. globosa* and *M. slooffiae*

were used for synteny analysis. Synteny analysis was done in the context of ORFs flanking the

centromeres of *M. sympodialis*. The protein sequences for each of these ORFs served as the

query in BLAST analysis against the genome of other species. The local database for each

genome was set up in the Geneious software for this analysis. The percentage identity values for

each ORFs are mentioned in the boxes in Figure 7- figure supplement 3. Additionally, synteny

analyses between *M. globosa* and *M. sympodialis* were conducted with megablast (word size: 28)

and plotted together with GC content (calculated as the deviation from the genomic mean, in

non-overlapping 1 kb windows), using Circos (v0.69-6) (Krzywinski et al., 2009). Additional

whole-genome alignments were conducted with Satsuma (Grabherr et al., 2010), with default

parameters. The linear synteny comparisons shown in Figures 5 and 6 were generated with the

Python application EasyFig (Sullivan et al., 2011).

Species phylogeny

To reconstruct the phylogenetic relationship among the 9 *Malassezia* species selected,

orthologs were identified using the bidirectional best-hit (BDBH), COGtriangles (v2.1), and

OrthoMCL (v1.4) algorithms implemented in the GET_HOMOLOGUES software package

(Contreras-Moreira and Vinuesa, 2013). The proteome of *M. sympodialis* ATCC42132 was used

as a reference. A phylogeny was inferred from a set of 738 protein sequences as follows.

Individual proteins were aligned using MAFFT v7.310 (L-INS-i strategy) and poorly aligned

regions were trimmed with TrimAl (-gappyout). The resulting alignments were concatenated to

obtain a final supermatrix consisting of a total of 441,200 amino acid sites (159,270 parsimony-

informative). This sequence was input to IQ-TREE v1.6.5 (Nguyen et al., 2015) and a maximum

likelihood phylogeny was estimated using the LG+F+R4 amino acid model of substitution.

Branch support values were obtained from 10,000 replicates of both ultrafast bootstrap

approximation (UFBoot) and the nonparametric variant of the approximate likelihood ratio test

(SH-aLRT) implemented in IQ-TREE. The best likelihood tree was graphically visualized with

iTOL v4.3.3 (Letunic and Bork, 2019).

Data availability

Acknowledgments

We thank Clevergene Biocorp Pvt. Ltd., Bengaluru for generating the Mtw1 ChIP-

sequencing data. S.R.S is a Senior Research Fellow supported by intramural funding from

JNCASR. M.H.R is a National Postdoctoral Fellow (PDF/2016/002858), supported by the

Science and Engineering Research Board (SERB), Department of Science and Technology

(DST), Government of India. K.S is a Tata Innovation Fellow (grant number

BT/HRT/35/01/03/2017) and is supported by a grant for Life Science Research, Education and

Training (BT/INF/22/SP27679/2018) of Department of Biotechnology, Govt. of India and

intramural funding from JNCASR. Studies were supported in part by NIH/NIAID R37 award

AI39115-21 and R01 award AI50113-15 to J.H. J.H is co-director and fellow of CIFAR program

Fungal Kingdom: Threats & opportunities. T.L.D acknowledges the A* STAR Industry

alignment fund H18/01a0/016, Asian Skin Microbiome Program. We thank the members of K.S

lab and J.H lab for valuable discussions and comments during bi-weekly Skype meetings.

Author contributions

K.S and J.H conceived the study and secured funding. S.S performed PFGE and

chromoblot analysis of *M. slooffiae* and *M. globosa*. M.H.R generated the constructs for tagging

MsyMtw1 and MfCENP-A. Transformation of *Malassezia* strains and PCR confirmation was

done by G.I. S.R.S performed western blot, microscopy, ChIP-qPCR analysis, and prepared the

samples for ChIP-sequencing. P.G analyzed the ChIP-seq data and RNA-seq data for *M.*

sympodialis. S.R.S, B.C.T, M.H.R, and M.A.C performed the synteny conservation analysis.

M.A.C performed genome comparisons, phylogenetic analysis, and prepared the figures. R.N.V

and R.S generated the GC/GC3 plots and performed motif analysis reported in this study. T.L.D

and C.T.R provided the genome assemblies of *M. globosa*, *M. slooffiae*, and *M. furfur*. S.R.S and

K.S wrote the manuscript with help from all of the authors. All of the authors revised and

approved the final version.

Declaration of Interests

The authors declare no competing interests.

References

AMEND, A. 2014. From dandruff to deep-sea vents: *Malassezia*-like fungi are ecologically hyper-diverse.
PLoS Pathog, 10**,** e1004277.
AYKUT, B., PUSHALKAR, S., CHEN, R., LI, Q., ABENGOZAR, R., KIM, J. I., SHADALOEY, S. A., WU, D., PREISS,
P., VERMA, N., GUO, Y., SAXENA, A., VARDHAN, M., DISKIN, B., WANG, W., LEINWAND, J., KURZ,
E., KOCHEN ROSSI, J. A., HUNDEYIN, M., ZAMBRINIS, C., LI, X., SAXENA, D. & MILLER, G. 2019.
The fungal mycobiome promotes pancreatic oncogenesis via activation of MBL. *Nature,* 574**,**
264-67.
BARRACLOUGH, T. G., BIRKY, C. W., JR. & BURT, A. 2003. Diversification in sexual and asexual organisms.
Evolution, 57**,** 2166-72.
BENSASSON, D., ZAROWIECKI, M., BURT, A. & KOUFOPANOU, V. 2008. Rapid evolution of yeast
centromeres in the absence of drive. *Genetics,* 178**,** 2161-67.
BOEKHOUT, T. & BOSBOOM, R. W. 1994. Karyotyping of *Malassezia* yeasts - taxonomic and
epidemiologic implications. *Systematic and Applied Microbiology,* 17**,** 146-53.
BOEKHOUT, T., KAMP, M. & GUEHO, E. 1998. Molecular typing of *Malassezia* species with PFGE and
RAPD. *Med Mycol,* 36**,** 365-72.
BOEKHOUT, T. A. N., E.S., 2003. *Pulsed Field Gel Electrophoresis (PFGE) of Yeasts*.
BYRNE, K. P. & WOLFE, K. H. 2005. The Yeast Gene Order Browser: combining curated homology and
syntenic context reveals gene fate in polyploid species. *Genome Res,* 15**,** 1456-61.
CELIS, A. M., VOS, A. M., TRIANA, S., MEDINA, C. A., ESCOBAR, N., RESTREPO, S., WOSTEN, H. A. & DE
COCK, H. 2017. Highly efficient transformation system for *Malassezia furfur* and *Malassezia*
pachydermatis using *Agrobacterium tumefaciens*-mediated transformation. *J Microbiol*
Methods, 134**,** 1-6.
CHATTERJEE, G., SANKARANARAYANAN, S. R., GUIN, K., THATTIKOTA, Y., PADMANABHAN, S.,
SIDDHARTHAN, R. & SANYAL, K. 2016. Repeat-associated fission yeast-like regional centromeres
in the ascomycetous budding yeast *Candida tropicalis*. *PLoS Genet,* 12**,** e1005839.
CHEN, E. S., SAITOH, S., YANAGIDA, M. & TAKAHASHI, K. 2003. A cell cycle-regulated GATA factor
promotes centromeric localization of CENP-A in fission yeast. *Mol Cell,* 11**,** 175-87.
CHIKASHIGE, Y., KINOSHITA, N., NAKASEKO, Y., MATSUMOTO, T., MURAKAMI, S., NIWA, O. & YANAGIDA,
M. 1989. Composite motifs and repeat symmetry in *S. pombe* centromeres: direct analysis by
integration of NotI restriction sites. *Cell,* 57**,** 739-51.
CLARKE, L. & BAUM, M. P. 1990. Functional analysis of a centromere from fission yeast: a role for
centromere-specific repeated DNA sequences. *Mol Cell Biol,* 10**,** 1863-72.
CLARKE, L. & CARBON, J. 1980. Isolation of a yeast centromere and construction of functional small
circular chromosomes. *Nature,* 287**,** 504-9.
CONTRERAS-MOREIRA, B. & VINUESA, P. 2013. GET_HOMOLOGUES, a versatile software package for
scalable and robust microbial pangenome analysis. *Appl Environ Microbiol,* 79**,** 7696-701.

DINER, R. E., NODDINGS, C. M., LIAN, N. C., KANG, A. K., MCQUAID, J. B., JABLANOVIC, J., ESPINOZA, J. L.,
NGUYEN, N. A., ANZELMATTI, M. A., JR., JANSSON, J., BIELINSKI, V. A., KARAS, B. J., DUPONT, C.
L., ALLEN, A. E. & WEYMAN, P. D. 2017. Diatom centromeres suggest a mechanism for nuclear
DNA acquisition. *Proc Natl Acad Sci U S A,* 114**,** 6015-24.
ECKERT-BOULET, N., PEDERSEN, M. L., KROGH, B. O. & LISBY, M. 2012. Optimization of ordered plasmid
assembly by gap repair in *Saccharomyces cerevisiae*. *Yeast,* 29**,** 323-34.
EUSKIRCHEN, G. M. 2002. Nnf1p, Dsn1p, Mtw1p, and Nsl1p: a new group of proteins important for
chromosome segregation in *Saccharomyces cerevisiae*. *Eukaryot Cell,* 1**,** 229-40.
FINDLEY, K. & GRICE, E. A. 2014. The skin microbiome: a focus on pathogens and their association with
skin disease. *PLoS Pathog,* 10**,** e1004436.
FINDLEY, K., SUN, S., FRASER, J. A., HSUEH, Y. P., AVERETTE, A. F., LI, W., DIETRICH, F. S. & HEITMAN, J.
2012. Discovery of a modified tetrapolar sexual cycle in *Cryptococcus amylolentus* and the
evolution of *MAT* in the *Cryptococcus* species complex. *PLoS Genet,* 8**,** e1002528.
FITZGERALD-HAYES, M., CLARKE, L. & CARBON, J. 1982. Nucleotide sequence comparisons and
functional analysis of yeast centromere DNAs. *Cell,* 29**,** 235-44.
GORDON, J. L., BYRNE, K. P. & WOLFE, K. H. 2011. Mechanisms of chromosome number evolution in
yeast. *PLoS Genet,* 7**,** e1002190.
GOSHIMA, G., KIYOMITSU, T., YODA, K. & YANAGIDA, M. 2003. Human centromere chromatin protein
hMis12, essential for equal segregation, is independent of CENP-A loading pathway. *J Cell Biol,*
160**,** 25-39.
GOSHIMA, G., SAITOH, S. & YANAGIDA, M. 1999. Proper metaphase spindle length is determined by
centromere proteins Mis12 and Mis6 required for faithful chromosome segregation. *Genes Dev,*
13**,** 1664-77.
GRABHERR, M. G., RUSSELL, P., MEYER, M., MAUCELI, E., ALFOLDI, J., DI PALMA, F. & LINDBLAD-TOH, K.
2010. Genome-wide synteny through highly sensitive sequence alignment: Satsuma.
Bioinformatics, 26**,** 1145-51.
GREENFEDER, S. A. & NEWLON, C. S. 1992. Replication forks pause at yeast centromeres. *Mol Cell Biol,*
12**,** 4056-66.
HAN, F., GAO, Z. & BIRCHLER, J. A. 2009. Reactivation of an inactive centromere reveals epigenetic and
structural components for centromere specification in maize. *Plant Cell,* 21**,** 1929-39.
HELMRICH, A., BALLARINO, M. & TORA, L. 2011. Collisions between replication and transcription
complexes cause common fragile site instability at the longest human genes. *Mol Cell,* 44**,** 966-
77.
HEMMERICH, P., STOYAN, T., WIELAND, G., KOCH, M., LECHNER, J. & DIEKMANN, S. 2000. Interaction of
yeast kinetochore proteins with centromere-protein/transcription factor Cbf1. *Proc Natl Acad
Sci U S A,* 97**,** 12583-8.
HENIKOFF, S., AHMAD, K. & MALIK, H. S. 2001. The centromere paradox: stable inheritance with rapidly
evolving DNA. *Science,* 293**,** 1098-102.
IANIRI, G., APPLEN CLANCEY, S., LEE, S. C. & HEITMAN, J. 2017a. FKBP12-dependent inhibition of
calcineurin mediates immunosuppressive antifungal drug action in *Malassezia*. *mBio,* 8**,** e01752-
17.
IANIRI, G., AVERETTE, A. F., KINGSBURY, J. M., HEITMAN, J. & IDNURM, A. 2016. Gene function analysis
in the ubiquitous human commensal and pathogen *Malassezia* genus. *mBio,* 7 , e01853-16.
IANIRI, G., BOYCE, K. J. & IDNURM, A. 2017b. Isolation of conditional mutations in genes essential for
viability of *Cryptococcus neoformans*. *Curr Genet,* 63**,** 519-30.
INTERNATIONAL CHICKEN GENOME SEQUENCING, C. 2004. Sequence and comparative analysis of the
chicken genome provide unique perspectives on vertebrate evolution. *Nature,* 432**,** 695-716.

ISHII, K., OGIYAMA, Y., CHIKASHIGE, Y., SOEJIMA, S., MASUDA, F., KAKUMA, T., HIRAOKA, Y. &
TAKAHASHI, K. 2008. Heterochromatin integrity affects chromosome reorganization after
centromere dysfunction. *Science,* 321**,** 1088-91.
IWANAGA, S., KHAN, S. M., KANEKO, I., CHRISTODOULOU, Z., NEWBOLD, C., YUDA, M., JANSE, C. J. &
WATERS, A. P. 2010. Functional identification of the Plasmodium centromere and generation of
a Plasmodium artificial chromosome. *Cell Host Microbe,* 7**,** 245-55.
JDO, J., BALDINI, A., WARD, D. C., REEDERS, S. T. & WELLS, R. A. 1991. Origin of human chromosome 2:
an ancestral telomere-telomere fusion. *Proc Natl Acad Sci U S A,* 88**,** 9051-55.
KALITSIS, P. & CHOO, K. H. 2012. The evolutionary life cycle of the resilient centromere. *Chromosoma,*
121**,** 327-40.
KAPOOR, S., ZHU, L., FROYD, C., LIU, T. & RUSCHE, L. N. 2015. Regional centromeres in the yeast *Candida*
lusitaniae lack pericentromeric heterochromatin. *Proc Natl Acad Sci U S A,* 112**,** 12139-44.
KOBAYASHI, N., SUZUKI, Y., SCHOENFELD, L. W., MULLER, C. A., NIEDUSZYNSKI, C., WOLFE, K. H. &
TANAKA, T. U. 2015. Discovery of an unconventional centromere in budding yeast redefines
evolution of point centromeres. *Curr Biol,* 25**,** 2026-33.
KOO, D. H., HAN, F., BIRCHLER, J. A. & JIANG, J. 2011. Distinct DNA methylation patterns associated with
active and inactive centromeres of the maize B chromosome. *Genome Res,* 21**,** 908-14.
KOPS, G. J., WEAVER, B. A. & CLEVELAND, D. W. 2005. On the road to cancer: aneuploidy and the mitotic
checkpoint. *Nat Rev Cancer,* 5**,** 773-85.
KRZYWINSKI, M., SCHEIN, J., BIROL, I., CONNORS, J., GASCOYNE, R., HORSMAN, D., JONES, S. J. &
MARRA, M. A. 2009. Circos: an information aesthetic for comparative genomics. *Genome Res,*
19**,** 1639-45.
LETESSIER, A., MILLOT, G. A., KOUNDRIOUKOFF, S., LACHAGES, A. M., VOGT, N., HANSEN, R. S., MALFOY,
B., BRISON, O. & DEBATISSE, M. 2011. Cell-type-specific replication initiation programs set
fragility of the FRA3B fragile site. *Nature,* 470**,** 120-23.
LETUNIC, I. & BORK, P. 2019. Interactive Tree Of Life (iTOL) v4: recent updates and new developments.
Nucleic Acids Res, 47**,** W256-W259.
LIMON, J. J., TANG, J., LI, D., WOLF, A. J., MICHELSEN, K. S., FUNARI, V., GARGUS, M., NGUYEN, C.,
SHARMA, P., MAYMI, V. I., ILIEV, I. D., SKALSKI, J. H., BROWN, J., LANDERS, C., BORNEMAN, J.,
BRAUN, J., TARGAN, S. R., MCGOVERN, D. P. B. & UNDERHILL, D. M. 2019. *Malassezia* is
associated with crohn's disease and exacerbates colitis in mouse models. *Cell Host Microbe,* 25**,**
377-88 e6.
LING, Y. H. & YUEN, K. W. Y. 2019. Point centromere activity requires an optimal level of centromeric
noncoding RNA. *Proc Natl Acad Sci U S A,* 116**,** 6270-79.
LO, A. W., CRAIG, J. M., SAFFERY, R., KALITSIS, P., IRVINE, D. V., EARLE, E., MAGLIANO, D. J. & CHOO, K.
H. 2001. A 330 kb CENP-A binding domain and altered replication timing at a human
neocentromere. *EMBO J,* 20**,** 2087-96.
LORCH, J. M., PALMER, J. M., VANDERWOLF, K. J., SCHMIDT, K. Z., VERANT, M. L., WELLER, T. J. &
BLEHERT, D. S. 2018. *Malassezia vespertilionis* sp. nov.: a new cold-tolerant species of yeast
isolated from bats. *Persoonia-Molecular Phylogeny and Evolution of Fungi,* 41**,** 56-70.
LYNCH, D. B., LOGUE, M. E., BUTLER, G. & WOLFE, K. H. 2010. Chromosomal G + C content evolution in
yeasts: systematic interspecies differences, and GC-poor troughs at centromeres. *Genome Biol*
Evol, 2**,** 572-83.
LYSAK, M. A. 2014. Live and let die: centromere loss during evolution of plant chromosomes. *New*
Phytologist, 203**,** 1082-89.
MALIK, H. S. & HENIKOFF, S. 2009. Major evolutionary transitions in centromere complexity. *Cell,* 138**,**
1067-82.
MCCLINTOCK, B. 1941. The stability of broken ends of chromosomes in *Zea mays*. *Genetics,* 26**,** 234-82.

MITRA, S., GOMEZ-RAJA, J., LARRIBA, G., DUBEY, D. D. & SANYAL, K. 2014. Rad51-Rad52 mediated
maintenance of centromeric chromatin in *Candida albicans*. *PLoS Genet,* 10**,** e1004344.
MURAT, F., XU, J. H., TANNIER, E., ABROUK, M., GUILHOT, N., PONT, C., MESSING, J. & SALSE, J. 2010.
Ancestral grass karyotype reconstruction unravels new mechanisms of genome shuffling as a
source of plant evolution. *Genome Res,* 20**,** 1545-57.
MUSACCHIO, A. & DESAI, A. 2017. A molecular view of kinetochore assembly and function. *Biology*
(Basel), 6**,** 5-54.
NAGAKI, K., CHENG, Z., OUYANG, S., TALBERT, P. B., KIM, M., JONES, K. M., HENIKOFF, S., BUELL, C. R. &
JIANG, J. 2004. Sequencing of a rice centromere uncovers active genes. *Nat Genet,* 36**,** 138-45.
NAKAYASHIKI, H., KADOTANI, N. & MAYAMA, S. 2006. Evolution and diversification of RNA silencing
proteins in fungi. *J Mol Evol,* 63**,** 127-35.
NAVARRO-MENDOZA, M. I., PEREZ-ARQUES, C., PANCHAL, S., NICOLAS, F. E., MONDO, S. J., GANGULY,
P., PANGILINAN, J., GRIGORIEV, I. V., HEITMAN, J., SANYAL, K. & GARRE, V. 2019. Early diverging
fungus *Mucor circinelloides* lacks centromeric histone CENP-A and displays a mosaic of point and
regional centromeres. *Curr Biol,* 22**,** 3791-802.
NGUYEN, L. T., SCHMIDT, H. A., VON HAESELER, A. & MINH, B. Q. 2015. IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol,* 32**,** 268-74.
OHKUNI, K. & KITAGAWA, K. 2011. Endogenous transcription at the centromere facilitates centromere
activity in budding yeast. *Curr Biol,* 21**,** 1695-703.
OZERI-GALAI, E., LEBOFSKY, R., RAHAT, A., BESTER, A. C., BENSIMON, A. & KEREM, B. 2011. Failure of
origin activation in response to fork stalling leads to chromosomal instability at fragile sites. *Mol*
Cell, 43**,** 122-31.
PADMANABHAN, S., THAKUR, J., SIDDHARTHAN, R. & SANYAL, K. 2008. Rapid evolution of Cse4p-rich
centromeric DNA sequences in closely related pathogenic yeasts, *Candida albicans* and *Candida*
dubliniensis. *Proc Natl Acad Sci U S A,* 105**,** 19797-802.
RAVI, M. & CHAN, S. W. 2010. Haploid plants produced by centromere-mediated genome elimination.
Nature, 464**,** 615-18.
RHIND, N., CHEN, Z., YASSOUR, M., THOMPSON, D. A., HAAS, B. J., HABIB, N., WAPINSKI, I., ROY, S., LIN,
M. F., HEIMAN, D. I., YOUNG, S. K., FURUYA, K., GUO, Y., PIDOUX, A., CHEN, H. M., ROBBERTSE,
B., GOLDBERG, J. M., AOKI, K., BAYNE, E. H., BERLIN, A. M., DESJARDINS, C. A., DOBBS, E., DUKAJ,
L., FAN, L., FITZGERALD, M. G., FRENCH, C., GUJJA, S., HANSEN, K., KEIFENHEIM, D., LEVIN, J. Z.,
MOSHER, R. A., MULLER, C. A., PFIFFNER, J., PRIEST, M., RUSS, C., SMIALOWSKA, A., SWOBODA,
P., SYKES, S. M., VAUGHN, M., VENGROVA, S., YODER, R., ZENG, Q., ALLSHIRE, R., BAULCOMBE,
D., BIRREN, B. W., BROWN, W., EKWALL, K., KELLIS, M., LEATHERWOOD, J., LEVIN, H.,
MARGALIT, H., MARTIENSSEN, R., NIEDUSZYNSKI, C. A., SPATAFORA, J. W., FRIEDMAN, N.,
DALGAARD, J. Z., BAUMANN, P., NIKI, H., REGEV, A. & NUSBAUM, C. 2011. Comparative
functional genomics of the fission yeasts. *Science,* 332**,** 930-36.
RIEF, N. & LOBRICH, M. 2002. Efficient rejoining of radiation-induced DNA double-strand breaks in
centromeric DNA of human cells. *J Biol Chem,* 277**,** 20572-82.
ROCCHI, M., STANYON, R. & ARCHIDIACONO, N. 2009. Evolutionary new centromeres in primates. *Prog*
Mol Subcell Biol, 48**,** 103-52.
ROY, B., BURRACK, L. S., LONE, M. A., BERMAN, J. & SANYAL, K. 2011. CaMtw1, a member of the
evolutionarily conserved Mis12 kinetochore protein family, is required for efficient inner
kinetochore assembly in the pathogenic yeast *Candida albicans*. *Mol Microbiol,* 80**,** 14-32.
ROY, B. & SANYAL, K. 2011. Diversity in requirement of genetic and epigenetic factors for centromere
function in fungi. *Eukaryot Cell,* 10**,** 1384-95.

SANEI, M., PICKERING, R., KUMKE, K., NASUDA, S. & HOUBEN, A. 2011. Loss of centromeric histone H3
(CENH3) from centromeres precedes uniparental chromosome elimination in interspecific barley
hybrids. *Proc Natl Acad Sci U S A,* 108**,** 498-505.
SANYAL, K., BAUM, M. & CARBON, J. 2004. Centromeric DNA sequences in the pathogenic yeast *Candida*
albicans are all different and unique. *Proc Natl Acad Sci U S A,* 101**,** 11374-79.
SATO, H., MASUDA, F., TAKAYAMA, Y., TAKAHASHI, K. & SAITOH, S. 2012. Epigenetic inactivation and
subsequent heterochromatinization of a centromere stabilize dicentric chromosomes. *Curr Biol,*
22**,** 658-67.
SCHOTANUS, K. & HEITMAN, J. 2019. Centromere deletion in *Cryptococcus deuterogattii* leads to
neocentromere formation and chromosome fusions. *bioRxiv***,** 526962.
SCHUBERT, I. & LYSAK, M. A. 2011. Interpretation of karyotype evolution should consider chromosome
structural constraints. *Trends Genet,* 27**,** 207-16.
SCHWARTZ, M., ZLOTORYNSKI, E., GOLDBERG, M., OZERI, E., RAHAT, A., LE SAGE, C., CHEN, B. P., CHEN,
D. J., AGAMI, R. & KEREM, B. 2005. Homologous recombination and nonhomologous end-joining
repair pathways regulate fragile site stability. *Genes Dev,* 19**,** 2715-26.
SIDDHARTHAN, R. 2008. PhyloGibbs-MP: module prediction and discriminative motif-finding by Gibbs
sampling. *PLoS Comput Biol,* 4**,** e1000156.
SIDDHARTHAN, R., SIGGIA, E. D. & VAN NIMWEGEN, E. 2005. PhyloGibbs: a Gibbs sampling motif finder
that incorporates phylogeny. *PLoS Comput Biol,* 1**,** e67.
SMITH, J. G., CADDLE, M. S., BULBOACA, G. H., WOHLGEMUTH, J. G., BAUM, M., CLARKE, L. & CALOS, M.
P. 1995. Replication of centromere II of *Schizosaccharomyces pombe*. *Mol Cell Biol,* 15**,** 5165-72.
STIMPSON, K. M., SONG, I. Y., JAUCH, A., HOLTGREVE-GREZ, H., HAYDEN, K. E., BRIDGER, J. M. &
SULLIVAN, B. A. 2010. Telomere disruption results in non-random formation of de novo dicentric
chromosomes involving acrocentric human chromosomes. *PLoS Genet,* 6, **,** e1001061.
SUGITA, T., TAKASHIMA, M., KODAMA, M., TSUBOI, R. & NISHIKAWA, A. 2003. Description of a new
yeast species, *Malassezia japonica*, and its detection in patients with atopic dermatitis and
healthy subjects. *J Clin Microbiol,* 41**,** 4695-99.
SULLIVAN, M. J., PETTY, N. K. & BEATSON, S. A. 2011. Easyfig: a genome comparison visualizer.
Bioinformatics, 27**,** 1009-10.
SUN, S., BILLMYRE, R. B., MIECZKOWSKI, P. A. & HEITMAN, J. 2014. Unisexual reproduction drives
meiotic recombination and phenotypic and karyotypic plasticity in *Cryptococcus neoformans*.
PLoS Genet, 10**,** e1004849.
SUN, S., YADAV, V., BILLMYRE, R. B., CUOMO, C. A., NOWROUSIAN, M., WANG, L., SOUCIET, J. L.,
BOEKHOUT, T., PORCEL, B., WINCKER, P., GRANEK, J. A., SANYAL, K. & HEITMAN, J. 2017. Fungal
genome and mating system transitions facilitated by chromosomal translocations involving
intercentromeric recombination. *PLoS Biol,* 15**,** e2002527.
TALBERT, P. B., BRYSON, T. D. & HENIKOFF, S. 2004. Adaptive evolution of centromere proteins in plants
and animals. *J Biol,* 3**,** 18.1-18.17.
THEELEN, B., CAFARCHIA, C., GAITANIS, G., BASSUKAS, I. D., BOEKHOUT, T. & DAWSON, T. L., JR. 2018.
Malassezia ecology, pathophysiology, and treatment. *Med Mycol,* 56**,** S10-25.
TROMER, E. C., VAN HOOFF, J. J. E., KOPS, G. & SNEL, B. 2019. Mosaic origin of the eukaryotic
kinetochore. *Proc Natl Acad Sci U S A,* 116**,** 12873-82.
WANG, K., WU, Y., ZHANG, W., DAWE, R. K. & JIANG, J. 2014. Maize centromeres expand and adopt a
uniform size in the genetic background of oat. *Genome Res,* 24**,** 107-16.
WESTERMANN, S., CHEESEMAN, I. M., ANDERSON, S., YATES, J. R., 3RD, DRUBIN, D. G. & BARNES, G.
2003. Architecture of the budding yeast kinetochore reveals a conserved molecular core. *J Cell*
Biol, 163**,** 215-22.

WU, G., ZHAO, H., LI, C., RAJAPAKSE, M. P., WONG, W. C., XU, J., SAUNDERS, C. W., REEDER, N. L.,
REILMAN, R. A., SCHEYNIUS, A., SUN, S., BILLMYRE, B. R., LI, W., AVERETTE, A. F., MIECZKOWSKI,
P., HEITMAN, J., THEELEN, B., SCHRODER, M. S., DE SESSIONS, P. F., BUTLER, G., MAURER-
STROH, S., BOEKHOUT, T., NAGARAJAN, N. & DAWSON, T. L., JR. 2015. Genus-wide comparative
genomics of *Malassezia* delineates its phylogeny, physiology, and niche adaptation on human
skin. *PLoS Genet,* 11**,** e1005614.
YADAV, V., SREEKUMAR, L., GUIN, K. & SANYAL, K. 2018a. Five pillars of centromeric chromatin in fungal
pathogens. *PLoS Pathog,* 14**,** e1007150.
YADAV, V., SUN, S., BILLMYRE, R. B., THIMMAPPA, B. C., SHEA, T., LINTNER, R., BAKKEREN, G., CUOMO,
C. A., HEITMAN, J. & SANYAL, K. 2018b. RNAi is a critical determinant of centromere evolution in
closely related fungi. *Proc Natl Acad Sci U S A,* 115**,** 3108-13.
YADAV, V., YANG, F., REZA, M. H., LIU, S., VALENT, B., SANYAL, K. & NAQVI, N. I. 2019. Cellular dynamics
and genomic identity of centromeres in cereal blast fungus. *mBio,* 10**,** e01581-19.
YUNIS, J. J. & PRAKASH, O. 1982. The origin of man: a chromosomal pictorial legacy. *Science,* 215**,** 1525-
30.
ZHANG, H. & FREUDENREICH, C. H. 2007. An AT-rich sequence in human common fragile site FRA16D
causes fork stalling and chromosome breakage in *S. cerevisiae*. *Mol Cell,* 27**,** 367-79.
ZHANG, W., FRIEBE, B., GILL, B. S. & JIANG, J. 2010. Centromere inactivation and epigenetic
modifications of a plant chromosome with three functional centromeres. *Chromosoma,* 119**,**
553-63.
ZHU, Y., ENGSTROM, P. G., TELLGREN-ROTH, C., BAUDO, C. D., KENNELL, J. C., SUN, S., BILLMYRE, R. B.,
SCHRODER, M. S., ANDERSSON, A., HOLM, T., SIGURGEIRSSON, B., WU, G.,
SANKARANARAYANAN, S. R., SIDDHARTHAN, R., SANYAL, K., LUNDEBERG, J., NYSTEDT, B.,
BOEKHOUT, T., DAWSON, T. L., JR., HEITMAN, J., SCHEYNIUS, A. & LEHTIO, J. 2017.
Proteogenomics produces comprehensive and highly accurate protein-coding gene annotation
in a complete genome assembly of *Malassezia sympodialis*. *Nucleic Acids Res,* 45**,** 2629-43.

1018

1019

   Figures and figure supplements

1021

1022   **Figure 1**

1023

Figure 1. Genome assembly and karyotype diversity in representative *Malassezia* species. The

genomes of *M. globosa* (A), *M. slooffiae* (B), and *M. furfur* (C) were sequenced and assembled in this

study, while the genome assembly of *M. sympodialis* (D) was reported earlier (Zhu et. al., 2015) and is

shown for comparison. In each panel, bar plots represent the assembled chromosomes of the indicated

Malassezia species, with the telomeres and the ribosomal DNA (rDNA) marked as dark grey and blue

bars, respectively. Telomere-repeat motifs are shown at the 5'- and 3'-ends of the Chr1 in each species.

Electrophoretic karyotypes of *M. globosa* (*Mg*) and *M. sloffiiae* (*Msl*), are shown in (A) and (B)

respectively, with chromosome sizes estimated from the genome assembly. Chromosomes of

Saccharomyces cerevisiae (*Sc*) served as size markers. The chromosome containing the rDNA (marked

with an "r"), in *M. globosa,* co-migrates with Chr3. This was assessed by chromoblot hybridization using

unique sequences from Chr3, Chr4, Chr5, and Chr6 as probes (regions indicated by red arrowheads).

Chromosomes of similar size (denoted in red) migrate together in the gel and appear as a doublet band

(i.e. MgChr3-MgChr5, MslChr2-MslChr3, and MslChr5-MslChr6).

1037

	M. globosa	M. slooffiae	M. furfur
Strain	CBS7966	CBS7956	CBS14141
Sequencing platform	PacBio	PacBio	PacBio
No. of reads	122,622	92,622	57,425
Coverage	90X	140X	64X
Size (bp)	9,119,538	8,938,001	8,311,773
GC content (%)	52.09	66.31	64.9
Assembly status			
Total contigs	10	14	8
Nuclear genome	**Complete** 9 contigs telomere on both ends	**Complete** 9 contigs telomere on both ends	**Complete** 7 contigs telomere on both ends
Gaps	none	none	none
Assembly polished with Illumina reads	No	No	No

1038

1039 **Figure 1-figure supplement 1.** Statistics of the genome assemblies of *M. globosa*, *M. slooffiae,* and *M.*

1040 *furfur* generated in this study.

1041

1042   **Figure 2**

1043

Figure 2. Kinetochores cluster and localize at the nuclear periphery in *M. sympodialis*.

(A) Schematic of the kinetochore organization of *M. sympodialis*. Gray boxes indicate proteins absent in

M. sympodialis. The outer kinetochore protein Mtw1 (a component of Mis12C) served as the kinetochore

marker in the present study. (B) Line diagram representation of Mtw1 tagged with GFP at the N-

terminus. Immunoblot analysis of whole cell lysates prepared from the untagged *M. sympodialis* strain

(ATCC42132) and GFP-Mtw1 expressing cells (MSY001) probed with anti-GFP antibodies and anti-

PSTAIRE antibodies. PSTAIRE served as a loading control. Relative intensity values normalized to

PSTAIRE are indicated below each lane. (C) Logarithmically grown MSY001 cells expressing GFP-

Mtw1 were fixed and stained with DAPI (blue) and anti-GFP antibodies (pseudo-colored in red). Scale

bar, 2.5 µm. (D) Cell cycle stage-specific localization dynamics of GFP-Mtw1. Scale bar, 2.5 µm.

1054

1055

1056

1057

1058

1059

1060

1061

1062   **Figure 2- figure supplement 1**

	Sub complex	Query	Protein ID (SHOxxxxx.x)	% identity	*E*-value	Query coverage
Outer kinetochore	**Dam1 complex**	Ask1	79565.1	27.4	4.84e-22	65.64
		Dad1	75772.1	47.1	4.44e-01	27.36
		Dad2*	79986.1	40	9.9e-23	82.89
		Dad3	77770.1	42.5	3.62e-10	78.75
		Dad4	79701.1	39.1	1.24e-09	95.83
		Dam1	79667.1	43.7	6.44e-08	45.45
		Duo1	76611.1	27.4	1.19e-04	17.63
		Spc19	n.d.			
		Spc34	n.d.			
		Hsk3*	77065.1	41.9	1.6e-15	92.68
	Ndc80 complex	Ndc80	79901.1	25.6	1.15e-36	80.33
		Nuf2	78898.1	25.8	7.64e-14	74.59
		Spc24	n.d.			
		Spc25	78263.1	25.8	1.1e-07	42.18
	Mis12 complex	Mis12/Mtw1	76526.1	25.1	5.67e-09	70.91
		Dsn1	78762.1	24.4	3.94e-08	24.05
		Nnf1*	77715.1	40.8	2.10E-19	35.9
		Nsl1*	79581.1	28.3	1.68E-06	42
	KNL1	Spc105	75936.1	19.9	5.36e-11	40.59
Inner kinetochore	**Constitutive Centromere Associated Network (CCAN)**	Cnn1/Wip1/Mhf1/Mhf2	n.d.			
		Mcm16/Mcm22/Ctf3	n.d.			
		Okp1/Ame1/Ctf19/Mcm21	n.d.			
		Chl4/Iml3	n.d.			
	CENP-C	Mif2	79930.1	35.8	5.77e-29	22.67
	CENP-A	Cse4	76408.1	70.8	2.77e-46	66.67
Point CEN specific complex	**CBF3 complex**	Ndc10	n.d.			
		Cep3	n.d.			
		Ctf13	n.d.			

1063

Figure 2- figure supplement 1

Identification of kinetochore proteins in *M. sympodialis* by BLAST.

Putative homologs of the indicated proteins in *M. sympodialis* were identified by BLAST analysis using

corresponding protein sequences of *C. neoformans* as the query. Asterisk (*) indicates cases where

corresponding protein sequences from *Ustilago maydis* were used as query. To detect proteins of the

CBF3 complex, protein sequences of Ndc10, Cep3, and Ctf13 from *S. cerevisiae* were used as query.

'n.d.' indicates homolog not detected.

1071

1072   **Figure 3**

1073

1074

1075

Figure 3. Localization of Mtw1to single-peaks identifies centromeres on each of the eight

chromosomes of *M. sympodialis.* (A) GFP-Mtw1 ChIP-seq reads were mapped to each chromosome.

The *x*-axis indicates chromosomal coordinates (in kb) and the *y*-axis indicates read depth. "Input", reads

from total DNA; "IP," reads from immunoprecipitated sample; *CEN,* Mtw1-enriched regions derived by

subtracting input reads from those of the IP sample (peak values 0-2500). Additional peaks observed in

both IP and input tracks on Chr5 are from the rDNA locus. (B) A 13 kb-window of Mtw1 enrichment

profile (*CEN*, represented as peaks and heat-map in two different tracks, red) plotted along with the GC

content (%GC, blue) and regions of transcription (RNA-seq, green). Numbers mentioned in the topmost

track in every panel indicate chromosomal coordinates (in kb). The scales in the *y*-axis are as follows:

CEN (0-8000), %GC (0-75), RNA-seq reads (0-1500). Gray arrows in each panel indicate predicted ORFs

based on RNA-seq data with arrowheads pointing the direction of transcription of the corresponding gene,

also marked as '+' and '-' in the axis label. (C) Fold difference in Mtw1 and histone H3 enrichment at

CEN8 as compared to a non-centromeric control region (190 kb away on the right of *CEN1*) by qPCR

analysis. Schematic of a 4 kb region of Chr8 with *CEN8* core (yellow box) is depicted below the graph.

Blue lines indicate regions assayed by PCR: core-region corresponding to the GC trough, L1 and R1- 750

bp away from the core, L2- 1500 bp away from the core, and non-centromeric control region (190 kb

away from centromere in Chr1). The *x*-axis indicates regions across the *CEN8* probed by PCR and *y*-axis

indicates fold difference in the enrichment of Mtw1 and histone H3 as compared to the control region.

Error bars indicate standard deviation (SD). (D) Logo of the consensus DNA sequence identified from *M.*

sympodialis centromeres, graphically represented with the size of the base corresponding with the

frequency of occurrence.

1097

1098 **Figure 3- figure supplement 1**

1099

1100

Figure 3- figure supplement 1

Mtw1-enriched regions in *M. sympodialis* contain transcriptionally active genes.

(A) Normalized read counts (RPKM) for transcripts from ORFs flanking the centromere as compared

with the genome average RPKM value calculated from RNA-seq data of *M. sympodialis* (ATCC42132).

The *x*-axis represents ORFs labeled L1 - L5 and R1 - R5, with L1 and R1 being proximal to the GC

trough of the centromere. Genome average considered for comparison is labeled as global. The *y*-axis

represents Log_2 RPKM values. (B) ChIP-qPCR assays validating the enrichment of Mtw1 at the

centromeres identified by ChIP-sequencing analysis. The *x*-axis indicates individual *CEN* regions assayed

with primers mapping to the chromosomal GC troughs (see Table S7 for primer sequences) and the *y*-axis

indicates enrichment of Mtw1 over a centromere unlinked control region as 'normalized % input

immunoprecipitated'. Error bars indicate standard deviation (SD).

1112

Figure 3- figure supplement 2

1114

1115

1116

Figure 3- figure supplement 2

Sequence features of centromeres in *M. sympodialis*.

(A) The genome of *M. sympodialis* was scanned for matches to the 12 bp AT-rich motif using a 500 bp

sliding window. Hit counts (*y*-axis) were plotted against the chromosomal coordinates (*x*-axis, in kb). Red

asterisks near the line corresponding to maximum enrichment in every chromosome mark the regions

predicted as centromeres. (B) Comparison of the maximum motif counts in a 500 bp window in each *M.*

sympodialis chromosome (*y*-axis) estimated with the PWM of the 12 bp motif (blue bars, labelled real)

and a scrambled version of the matrix (orange bars, labelled scrambled). The *x*-axis indicates individual

chromosomes in *M. sympodialis* (C) Dot-plot generated by plotting sequences that include 5 kb flanking

the Mtw1-enriched regions in *M. sympodialis* against themselves is depicted. The coordinates of regions

used in *M. sympodialis* are *CEN1*: 781716-791716, *CEN2*: 350852-360852, *CEN3*: 232777-242777,

CEN4: 413320-423320, *CEN5*: 120251-130251, *CEN6*: 97796-107796, *CEN7*: 426611-436611 and

CEN8: 19905-29905.

1130

1131

MsyChr1

Mtw1 enriched region (784,833 to 788,599)

MsyCEN1 core (786,541 to 787,061)

5'- [DNA sequence figure — MsyChr1 centromere region, reading 5' to 3'] -3'

1132

1133

Figure 3- figure supplement 3. Enrichment of the 12 bp motif at the centromere core in *M.*
***sympodialis*.**

The sequences corresponding to the Mtw1 enriched regions in Chr1 (with the AT-rich core region

highlighted in pink) are taken as a representative of *M. sympodialis* centromeres. The sequences

corresponding to the 12 bp motif shown in figure 3D are highlighted. Red and blue highlights indicate the

forward and reverse orientation of motif occurrence.

Figure 4

1141

Figure 4. Centromeres in *M. furfur*, *M. slooffiae,* and *M. globosa* map to global GC troughs on each

chromosome. (A) Schematic of epitope tagging of CENP-A with 3xFLAG at the C-terminus is shown.

(Bottom) Immunoblot analysis using whole cell lysates prepared from the untagged wild-type *M. furfur*

(CBS14141) cells and CENP-A-3xFLAG expressing cells (MF001) probed with anti-FLAG antibodies

and anti-PSTAIRE antibodies. PSTAIRE served as the loading control. (B) Abundance of CENP-A at

each of the predicted *M. furfur* centromeres by qPCR analysis of DNA immunoprecipitated with anti-

FLAG affinity gel in MF001 cells expressing CENP-A-3xFLAG. The *x*-axis indicates individual *CEN*

regions assayed with primers homologous to the GC troughs on each chromosome that were predicted as

centromeres (see Table S7 for primer sequences). The non-centromeric control maps to a region 1.3 Mb

away from predicted *CEN1*. The *y*-axis indicates enrichment of CENP-A estimated as the percentage of input immunoprecipitated. (C) Abundance of CENP-A across *MfCEN7* by ChIP-qPCR analysis in MF001 cells. Schematic representation of a 2 kb region is shown below the graph. Yellow bar indicates the centromere core of *CEN7* corresponding to the GC trough. Blue bars indicate regions analyzed by qPCR: L1 and R1- 750 bp away from the centromere core. 'Control' refers to a region 1.3 Mb away from *CEN1*. The CENP-A enrichment is plotted along the *y*-axis as the percentage of input immunoprecipitated. (D, E) Comparison of relative abundance of histone H3 compared to histone H4 at the predicted centromeres to a non-centromeric control locus in *M. slooffiae* and *M. globosa* respectively. Enrichment was estimated as the percentage of input immunoprecipitated with histone H3 and histone H4 antibodies and their ratio, as indicated in *y*-axis, is plotted as relative enrichment. The *x*-axis indicates centromeres in each species and these were assayed with primers homologous to GC troughs in each chromosome that were predicted as the centromeres (see Table S7 for primer sequences). The control region unlinked to the centromere corresponds to a locus 630 kb away from *CEN1* in *M. slooffiae* and 416 kb away from *CEN2* in *M. globosa* respectively. (F) The relative abundance of histone H3 compared to histone H4 across *MslCEN1* by qPCR analysis of the DNA immunoprecipitated using histone H3 and histone H4 antibodies. Schematic of *MslCEN1* locus is shown below the graph. Yellow bar indicates the *CEN1* core region corresponding to the GC trough. Blue bars indicate regions analyzed by qPCR: L1 and L2 map to regions 750 bp and 1.5 kb to the left of the *CEN1* core; R1 maps to a region 750 bp to the right of the *CEN1* core. The control region corresponds to a locus 630 kb away from the *CEN1* core. The ratio of enrichment of histone H3 to that of histone H4 is plotted as the relative enrichment in *y*-axis. (G) The relative abundance of histone H3 compared to histone H4 across *MgCEN2* by qPCR analysis of the DNA immunoprecipitated using histone H3 and histone H4 antibodies. Schematic of a 5 kb region containing *MgCEN2* is shown below the graph. Yellow bar indicates the *CEN2* core region corresponding to the GC trough. Blue bars indicate regions analyzed by qPCR: L1 and L2 indicate regions 750 bp and 1.5 kb to the left of the *CEN2* core; R1 and R2 indicate regions 750 bp and 1.5 kb to the right of the *CEN2* core. The control region corresponds to a locus 416 kb away from the *CEN2* core. The ratio of enrichment of histone H3 to that of histone H4 is plotted as the relative enrichment in the *y*-axis. Error bars in figures B-G indicate standard deviation (SD); Statistical significance was tested by one-way ANOVA: * significant at $P < 0.05$, *** significant at $P < 0.001$.

1183   **Figure 4- figure supplement 1**

1184

1185

Figure 4- figure supplement 1

Putative centromeres of *M. furfur*, *M. slooffiae*, *M. globosa*, and *M. restricta* map to global GC

troughs in each chromosome. (A-D) Graphs indicating GC content (red lines) and GC3 content (black

lines) of each chromosome of *M. furfur*, *M. slooffiae, M. globosa,* and *M. restricta.* Position of putative

centromeres mapping to chromosomal GC minima are marked in blue. The *x*-axis indicates chromosomal

coordinates in Mb.

1192

1193   **Figure 4- figure supplement 2**

1194

Figure 4- figure supplement 2

The 12 bp AT-rich motif is enriched at the putative centromeres of *M. furfur, M. slooffiae, M. globosa,* and *M. restricta*. (A-D) The genomes of *M. furfur, M. slooffiae, M. globosa,* and *M. restricta* were scanned for matches to the 12 bp AT-rich motif using a 500 bp sliding window. Hit counts (y-axis) were plotted against the chromosomal coordinates (x-axis, in kb) for each of the above species. Red asterisks near the line corresponding to maximum enrichment in every chromosome or scaffold mark the regions predicted as centromeres in each species.

1206   **Figure 5**

1207

1208

Figure 5. *MgCEN2* and *MslCEN5* map to a synteny breakpoint in *M. sympodialis*. (A, B) Circos plots

depicting the conserved gene synteny blocks between *M. globosa* and *M. sympodialis* chromosomes and

M. slooffiae and *M. sympodialis* chromosomes. Tracks from outside to inside represent positions of

centromeres and telomeres, GC content (plotted as blue and red lines indicating GC content below or

above genome average, calculated in 0.4 kb non-overlapping windows), and colored connectors indicate

regions of conserved gene synteny between the two species. (C, D) Linear chromosome plots depicting

syntenic regions between Chr2 of *M. globosa* and Chr5 of *M. slooffiae* respectively with chromosomes of

M. sympodialis. GC content (in %) is shown as red/blue lines above each chromosome. Labels in black

circles mark the gene synteny breakpoints. Synteny breakpoint at *MgCEN2* and *MslCEN5* is marked as

III. The regions on MsyChr2 and MsyChr4 where the homologs of ORFs flanking the breakpoint are

located, are marked II and IV respectively. Labels I and V indicate gene synteny conservation on the

other side of the fusion points II and IV on MsyChr2 and MsyChr4 as compared to *M. globosa* and *M.*

slooffiae chromosomes respectively. (E, F) Zoomed-in image of the gene synteny breakpoint at *MgCEN2*

and *MslCEN5* representing the conservation of genes flanking these centromeres in *M. sympodialis*

chromosomes at the ORF level.

1224

1225   **Figure 6**

1226

1227

Figure 6. Inactivation of *CEN8* and *CEN9* of *M. slooffiae* and *M. globosa* in MfChr3 resulted in

chromosome number reduction in *M. furfur*. (A, B) Circos plots depicting the conserved gene synteny

blocks between the *M. slooffiae* and *M. globosa* chromosomes respectively with *M. furfur* chromosomes.

Tracks from outside to inside represent positions of centromeres and telomeres, GC content (plotted as

blue and red lines indicating GC content below or above genome average, calculated in 0.4 kb non-

overlapping windows), and colored connectors indicate regions of conserved gene synteny between the

two species. (C) Linear chromosome plot depicting the syntenic regions between chromosome 3 of *M.*

furfur and chromosomes 3, 4, 7, 8, and 9 of *M. sloofffiae.* GC content (in %) is shown as red/blue lines

above each chromosome. Regions corresponding to *MslCEN8* and *MslCEN9* in MfChr3 are marked I and

II respectively. (D) Linear chromosome plot depicting the gene synteny conservation between

chromosome 3 of *M. furfur* and chromosomes 1, 4, 8, and 9 of *M. globosa*. Regions corresponding to

MgCEN9 and *MgCEN8* in MfChr3 are marked I and II respectively.

1240

1241 **Figure 6- figure supplement 1**

1242

Figure 6- figure supplement 1

Loss of AT-content in regions corresponding to *CEN8* and *CEN9* of *M. sloofffiae* and *M. globosa* in

MfChr3.

(A, B) Zoomed-in view of regions marked I and II in figure 6B, corresponding to inactivated *MslCEN8*

and *MslCEN9* in *M. furfur* along with the regions of gene synteny is represented. GC content (in %) is

shown as red/blue line diagram above each chromosome. (C, D) Zoomed-in view of the regions marked I

and II in figure 6D, corresponding to inactivated *MgCEN9* and *MgCEN8* in *M. furfur* along with the

regions of gene synteny is represented. GC content (in %) is shown as red/blue lines above each

chromosome.

1252

1253

1254 **Figure 7**

1255

Figure 7. Karyotype evolution by loss of centromere function in *Malassezia* species.

(A) Phylogenetic relationships between the *Malassezia* species analyzed in this study are represented and

their chromosome numbers shown. Species representing each clade are color coded based on previous

reports (Theelen et al., 2018). The chromosome numbers for *M. slooffiae* and *M. globosa* are based on

results from this study. In the case of *M. sympodialis, M. restricta,* and *M. furfur*, the chromosome

numbers are based on previous reports (Zhu et al., 2017, Senczek et al., 1999, Boekhout and Bosboom,

1994). For *M. dermatis, M. nana*, *M. vespertilionis,* and *M. japonica,* the number of chromosomes were

estimated from the predicted number of centromeres. The node corresponding to the ancestral state of

Clade A and Clade B are labelled 'Anc. A' and 'Anc. B' respectively. Green and red circles indicate the

origins of karyotypes with 8 and 7 chromosomes respectively from an ancestral state of 9 chromosomes.

(B) Schematic of the centromere loss by breakage and the resulting reduction in chromosome number as

observed in *M. sympodialis* (represented as the current state). A karyotype with 9 chromosomes (as

shown for *M. globosa*) is depicted as the ancestral state. (C) Proposed model of centromere inactivation

observed in *M. furfur* as a consequence of fusion of AnChr8 and AnChr9 to the AnMfChr3 equivalent,

resulting in a 7-chromosome configuration. The fusion product corresponding to extant MfChr3 is

represented as the current state.

1272

1273

1274

1275

1276

1277

Figure 7- figure supplement 1

Putative centromeres of *M. dermatis*, *M. nana*, *M. vespertilionis,* and *M. japonica* map to global GC

troughs in each chromosome. (A-D) Graphs indicating GC content (red lines) and GC3 content (black

lines) of each chromosome of *M. dermatis*, *M. nana*, *M. vespertilionis,* and *M. japonica.* Position of

putative centromeres mapping to chromosomal GC minima are marked in blue. The *x*-axis indicates

chromosomal coordinates in Mb.

1284

1285 **Figure 7-figure supplement 2**

1286

1287   **Figure 7- figure supplement 3**

1288

1289

1290

Figure 7- figure supplement 2

The 12 bp AT-rich motif is enriched at the putative centromeres of *M. dermatis, M. nana, M.*

***japonica,* and *M. vespertilionis*.** The genomes of *M. dermatis, M. nana, M. japonica,* and *M.*

vespertilionis were scanned for matches to the 12 bp AT-rich motif using a 500 bp sliding window. Hit

counts (*y*-axis) were plotted against the chromosomal coordinates (*x*-axis, in kb) for each of the above

species. Red asterisks near the line corresponding to maximum enrichment in every chromosome or

scaffold mark the regions predicted as centromeres in each species.

1298

Figure 7- figure supplement 3

All of the predicted centromeres of *M. sympodialis*, *M. nana,* and *M. dermatis* belong to conserved

gene synteny blocks when compared with species containing 9 chromosomes. Gene order and synteny

of ORFs flanking the centromeres of 6 *Malassezia* species representative of clade B and C analyzed in

this study by BLAST analysis using the protein sequences from *M. sympodialis* as query. Species

included are as follows: *Mna- M. nana*, *Mde- M. dermatis*, *Msy- M. sympodialis*, *Mgl- M. globosa*, *Mre-*

M. restricta, and *Msl- M. slooffiae*. Chromosome/scaffold numbers are indicated at the start of every

track. Boxes represent ORFs with the numbers inside them indicating percentage identity from BLAST

analysis. Broken lines marked with filled circle towards the ends indicate gene synteny breakpoints. Red

arrows indicate inverted orientation of the scaffold/chromosome.

1309

MgCEN2   MgCEN3

1310

1311

1312   **Figure 7- figure supplement 4**

1313   ***MgCEN2* and *MgCEN3* are predicted to form secondary structures.**

1314   The secondary structure of the centromere cores of *MgCEN2* and *MgCEN3* predicted by the ViennaRNA,

1315   generated using Geneious 9.0 (Lorenz et al., 2011). Blue and red circles mark the 5'- and the 3'- DNA ends

1316   respectively.

1317