Erosion of phylogenetic diversity in Neotropical bat assemblages: findings from a whole-

ecosystem fragmentation experiment

Sabhrina G. Aninta[1,4], Ricardo Rocha[2,3,5], Adrià López-Baucells[2,3,6], Christoph F. J. Meyer[2,3,7]

[1]Munich Graduate School for Evolution, Ecology and Systematics (EES), Faculty of Biology,

Ludwig-Maximilians-Universität München, Germany

[2]Centre for Ecology, Evolution and Environmental Changes – cE3c, Faculty of Sciences,

University of Lisbon, 1749-016 Lisbon, Portugal

[3]Biological Dynamics of Forest Fragments Project, National Institute for Amazonian

Research and Smithsonian Tropical Research Institute, 69011-970, Manaus, Brazil

[4]Tambora Muda Indonesia, Jalan Samiaji 3 Nomor 10 Bantarjati, Perumahan Indraprasta

Bogor Utara, 16153 Bogor, Indonesia

[5]Conservation Science Group, Department of Zoology, University of Cambridge, CB2 3EJ

Cambridge, United Kingdom

[6]Granollers Museum of Natural Sciences, Granollers 08402, Catalonia, Spain

[7]Ecosystems and Environment Research Centre (EERC), School of Environment and Life

Sciences, University of Salford, Salford M5 4WT, United Kingdom

Corresponding author: Sabhrina G. Aninta, sagitaninta[at]gmail.com

───────────────

Our study highlights the erosion of phylogenetic diversity of bat assemblages associated with habitat fragmentation and degradation in the world's largest and longest-running whole-ecosystem fragmentation experiment. This work advances our understanding of the effects of habitat modification on bats as key ecosystem service providers in tropical ecosystems.

Author Contributions: RR and AL conducted the fieldwork, SGA and CFJM formulated the idea, SGA performed the analysis, and SGA, RR, AL, and CFJM wrote the manuscript.

Abstract

The traditional focus on taxonomic diversity metrics for investigating species responses to

habitat loss and fragmentation has limited our understanding on how biodiversity is impacted

by habitat modification. This is particularly true for taxonomic groups such as bats which

exhibit species-specific responses. Here, we investigate phylogenetic alpha and beta diversity

of Neotropical bat assemblages across two environmental gradients, one in habitat quality and

one in habitat amount. We surveyed bats in 39 sites located across a whole-ecosystem

fragmentation experiment in the Brazilian Amazon, representing a gradient of habitat quality

(interior-edge-matrix, hereafter IEM) in both continuous forest and forest fragments of

different sizes (1, 10, and 100 ha; forest size gradient). For each habitat category, we

quantified alpha and beta phylogenetic diversity, then used linear models and cluster analysis

to explore how forest area and IEM gradient affect phylogenetic diversity. We found that the

secondary forest matrix harboured significantly lower total evolutionary history compared to

the fragment interiors, especially the 1 ha fragments, containing bat assemblages with more

closely related species. Forest fragments ≥10 ha had levels of phylogenetic richness similar to

continuous forest, suggesting that large fragments retain considerable levels of evolutionary

history. The edge and matrix adjacent to large fragments tend to have closely related lineages

nonetheless, suggesting phylogenetic homogenization in these IEM gradient categories. Thus,

despite the high mobility of bats, fragmentation still induces considerable levels of erosion of

phylogenetic diversity, suggesting that the various evolutionary history might not be able to

persist in present-day human-modified landscapes.

Keywords: evolutionary history, environmental gradients, Amazon, forest fragments,

secondary forests

Introduction

Humans have fundamentally changed the face of the Earth, with negative side-effects for biodiversity across all major biomes. Tropical forests are among the biomes impacted most heavily given the large footprint of pervasive land use changes which have resulted in widespread landscape fragmentation and loss of habitat for many species (Austin et al. 2017; Barlow et al. 2018). These land use changes have been documented to have detrimental effects on the richness, abundance, and composition of many tropical taxa (Alroy 2017), leading to a pattern of extensive defaunation, with cascading effects on ecosystem functioning (Young et al. 2016).

Idiosyncratic responses of species to fragmentation are ubiquitous, rendering assemblage-level inferences regarding fragmentation effects generally difficult (Ewers and Didham 2006; Fahrig 2017). This is mainly because the treatment of species as equal entities by neglecting their unique evolutionary history, functional roles in the ecosystem, and their association with each other within the community (Pellens and Grandcolas 2016), paints an incomplete picture of the effects of habitat fragmentation. Therefore, recent studies assessing the effect of habitat fragmentation on tropical taxa have started to incorporate evolutionary information using phylogenetic diversity metrics in addition to species richness (Frishkoff et al. 2014; Santos et al. 2014; Cisneros et al. 2015, 2016; Aguirre et al. 2016; Frank et al. 2017). By doing so, these studies were able to uncover patterns previously undetected by studies with a sole focus on the taxonomic dimension of biodiversity. For example, the random pattern of plant species absence across the phylogenetic tree in a fragmented landscape, or phylogenetic over-dispersion, can potentially be linked to low phylogenetic conservatism of key life-history traits (Santos et al. 2014), suggesting that the negative effect of habitat fragmentation depends on the evolutionary history of the taxa in question. However, studies on mobile species such as birds and bats showed non-random patterns of absence across their phylogeny in various

types of disturbed habitats (Riedinger et al. 2013; Frishkoff et al. 2014; Frank et al. 2017).

This pattern, often referred to as phylogenetic clustering, whereby species within an

assemblage are more closely related than expected by chance, indicates a strong effect of

habitat filtering. Gaining better insights into the extent to which phylogenetic diversity of

assemblages is eroded as a result of habitat fragmentation therefore is critical to improve our

general understanding of biodiversity persistence in human-modified landscapes.

Despite their mobility, bats (Chiroptera) are among the many animal groups that are

demonstrably affected by habitat loss and fragmentation (Meyer et al. 2016; Alroy 2017).

Notwithstanding increased research effort devoted over recent years to better understand how

bats respond to habitat fragmentation, studies at the assemblage level, typically comparing

species richness, diversity, and assemblage composition between forest fragments and

continuous forest, showed inconsistent results and highlighted the need for more research

focusing on the functional and phylogenetic biodiversity dimensions (Meyer et al. 2016). Bats

are a good model group to study the effect of habitat fragmentation on phylogenetic diversity

given their high species richness, functional diversity, and key roles in ecosystem functioning

(Kunz et al. 2011). Studies employing a phylogenetic approach to investigate bat responses

towards habitat disturbance (Cisneros et al. 2015; Frank et al. 2017; Presley et al. 2018) have

been made possible by the availability of phylogenetic trees of all extant bat species (Jones et

al. 2002, 2005; Shi and Rabosky 2015) that can be used to calculate phylogenetic diversity

metrics.

Of the few studies that have investigated bat phylogenetic diversity in fragmented landscapes,

none has assessed responses across the entire gradient in habitat quality typically encountered,

formed by the interiors (I) and edges (E) of continuous forest and forest fragments, as well as

the intervening matrix (M), or the IEM gradient (Rocha et al. 2017). Explicit consideration of

the full IEM gradient, however, is important to better understand the extent of environmental

filtering that usually is regarded as the cause of phylogenetic clustering in disturbed habitats

(Riedinger et al. 2013; Frank et al. 2017; Presley et al. 2018) as species persistence in

fragmented landscapes may be differentially affected by this gradient in habitat quality

(Ferreira et al. 2017). The observed phylogenetic richness and structure in each habitat that

comprises the IEM gradient can give an indication about the amount of evolutionary history

retained by the constituent habitat elements of a fragmented landscape (Cisneros et al. 2015).

Moreover, exploring which habitats share similar evolutionary history or harbour lineages that

are more closely related compared to other habitats may give insights into the evolution of

habitat preferences (Graham and Fine 2008).

To elucidate how habitat fragmentation affects the evolutionary dimension of bat diversity,

we investigated the changes in phylogenetic alpha and beta diversity of Amazonian bat

assemblages across two environmental gradients, one in habitat quality (IEM gradient) and

one in habitat amount (forest size: continuous forest; fragments of 1, 10, 100 ha), in the

experimentally fragmented landscape of the Biological Dynamics of Forest Fragments Project

(BDFFP), the world's largest and longest-running experimental study of habitat fragmentation

(Haddad et al. 2015). For phylogenetic alpha diversity, we expected assemblages in the

secondary forest matrix to retain the least total evolutionary history due to selection of bat

lineages that are best adapted to it (Rocha et al. 2018), followed by edges, while forest

interiors were predicted to harbour the most total evolutionary history due to greatest resource

availability (Ries and Sisk 2004). Accordingly, we predicted phylogenetic diversity to

increase with forest size, and phylogenetic clustering to be associated with smaller forest

fragments and strongest in the matrix. Finally, we expected that habitat filtering will leave

each habitat of the IEM gradient with its own set of unique assemblages so that phylogenetic

beta diversity will show that the same category of IEM gradient will contain similar lineages

and a similar amount of total evolutionary as a consequence of low phylogenetic turnover.

Methods

Study Area. The BDFFP spans ~1,000 km^2 and is located approximately 80 km north of

Manaus, Brazil (S2°30', W60°). The area contains different-sized forest fragments separated

by 80-650 m from the surrounding continuous forest that serves as experimental control

(Laurance et al. 2018). Following abandonment of the cattle pastures which initially

surrounded the experimentally isolated fragments after their creation in the early 1980s,

Vismia- and *Cecropia*-dominated secondary forest developed in the matrix (Mesquita et al.

2015).

The forest at the BDFFP is a typical non-flooded forest of the Amazon basin (De Oliveira and

Mori 1999), with approximately 280 species of trees (dbh >10 cm) per hectare (Laurance et

al. 2010). The area has a relatively flat topography (80-160 m), with nutrient-poor soils.

Rainfall ranges from 1,900 to 3,500 mm annually, with a moderately strong dry season from

June to October (Laurance et al. 2018).

Bat Sampling. Bats were sampled with ground-level mist nets in eight primary forest

fragments —three of 1 ha, three of 10 ha and two of 100 ha—and in nine control sites in three

areas of continuous forest. Fragments and controls were sampled in the interior, at the edges,

and in the secondary forest matrix, resulting in a total of 39 sites (for continuous forest, only

three edge and matrix sites were sampled; cf. Rocha et al. 2017). Distances between interior

and edge sites of continuous forest and fragments were respectively 1,118 ± 488 and 245 ±

208 m (mean ± SD). Matrix sites were located ca. 100 m away from the border between

primary and secondary forest. Each sampling site was visited eight times over a 2-year period

(August 2011 – June 2013). See (Rocha et al. 2017) for a more detailed description.

Phylogenetic Information. Analysis was restricted to the family Phyllostomidae, which can

be adequately sampled with ground-level mist nets (Kalko 1998). The phylogenetic

information of phyllostomid species present at the BDFFP, hereafter referred to as the local

phylogeny, was extracted from the most recent species-level phylogeny of bats (Shi and

Rabosky 2015) using R package `picante` (Kembel et al. 2010). We chose this particular tree

as it covered more of the phyllostomid species that occur at the BDFFP compared to another

frequently used bat phylogenetic tree published by (Jones et al. 2002, 2005). The supertree

was downloaded from TreeBASE and pruned to obtain the local phylogeny. The branch

lengths of the local phylogeny represent the divergence time of the species in millions of

years. As we used distance-based phylogenetic diversity metrics, the phylogenetic pairwise

distance matrix was extracted from the local phylogeny using the `cophenetic.phylo()`

function from R package `ape` (Paradis et al. 2004).

Species that occur in the study area but were not present in the pruned tree were substituted

by their congeners, following (Cisneros et al. 2016). Only for two out of 43 captured

phyllostomid species was this the case, i.e. *Artibeus gnomus* and *A. cinereus*, which hence

were both represented by their closest congener, *A. glaucus* (Redondo et al. 2008).

Measuring Phylogenetic Diversity

Alpha Diversity. We explored variation in phylogenetic richness and structure within

assemblages separately across the IEM gradient of 1, 10, and 100 ha fragments and

continuous forest (CF) using Faith's phylogenetic diversity (Faith 1992) and mean pairwise

distance (Clarke and Warwick 1998; Webb 2000), hereafter referred to as PD and MPD

respectively. As PD is sensitive to sample size (Vamosi et al. 2009), we rarefied the observed

values of PD when comparing different assemblages using the R package 'BAT'(Cardoso et

al. 2015).

To assess the effect of phylogenetic information *per se* on PD, we eliminated any potential

effect of species richness by calculating SES_{PD} (standardized effect size of PD) under a

'richness' null model (Swenson 2014). Similarly, we quantified phylogenetic structure *per se*

by calculating SES_{MPD} (standardized effect size of MPD) using the tip-shuffling null model

(Swenson 2014). However, as the local phylogeny consisted of closely related phyllostomids

with quite a balanced topology, MPD could underestimate phylogenetic clustering in the

terminal part of the phylogeny. We therefore also calculated the mean nearest taxon distance

(MNTD) and computed SES_{MNTD} using the tip-shuffling null model as it is more sensitive

than MPD for detecting phylogenetic clustering in the terminal part of the local phylogeny

(Tucker et al. 2017). For SES_{MPD} and SES_{MNTD}, high quantiles ($p > 0.95$) indicate a

phylogenetically over-dispersed assemblage whereas low quantiles ($p < 0.05$) indicate a

phylogenetically clustered assemblage (Swenson 2014).

Beta Diversity. To explore between-assemblage variation in phylogenetic richness and

structure in relation to the habitat quality (IEM) and size gradients, we calculated COMDIST

and phylogenetic beta diversity ($P\beta_{total}$), respectively. COMDIST is the mean pairwise

phylogenetic distance (MPD) between species from different assemblages (Swenson 2011)

which could help reveal which habitats along the gradient contain closely related lineages. To

calculate COMDIST, we used R package 'picante' (Kembel et al. 2010). Phylogenetic beta

diversity was partitioned into its richness ($P\beta_{rich}$) and replacement ($P\beta_{repl}$) components to

capture, respectively, the difference in shared total branch lengths between assemblages and

the uniqueness of each assemblage based on the evolutionary lineages present (Cardoso et al.

2014). To calculate $P\beta_{total}$ and its partitions, we used R package 'BAT' (Cardoso et al. 2015).

Phylogenetic alpha and beta diversity across the IEM and forest size gradient

Alpha Diversity. We modelled phylogenetic diversity in relation to the IEM and forest size

gradient to understand how the two affect the evolutionary dimension of biodiversity. While

in principle the six camp locations (Colosso, Porto Alegre, Dimona, Cabo Frio, Florestal and

Km 41) should be incorporated in the statistical model as a random effect to account for

potential spatial autocorrelation of the response variables (Bolker et al. 2009), we found its

effect to be negligible in practice after we fit linear mixed models using the `lme4` R package

(Bates et al. 2014) (see Table S1). We therefore dropped the block effect, leaving us with a

simple linear model to assess significance of the fixed effects. The linear models were fitted

using the R package 'stats' (R Core Team 2017), with IEM gradient (interior, edge, or

matrix) and forest size (CF, 100 ha, 10 ha, or 1 ha) as explanatory variables, comparing 12

types of assemblages in total. Both additive and interactive effects for IEM gradient and forest

size along with the null models were evaluated based on the small sample size version of

Akaike's Information Criterion (AICc) (Burnham and Anderson 2002) as implemented in R

package 'MuMIn' (Barton 2016). Along with AICc, we also calculated Akaike weights (w_i)

and Akaike differences (Δ_i) to gauge relative variable importance. Models with $\Delta_i \leq 2$ were

considered to have substantial support (Burnham and Anderson 2002). Significant effects

were further evaluated via multiple comparison tests with Tukey contrasts (adjusted *P* values

reported in Table S3) using the R package 'multcomp' (Hothorn et al. 2008).

Beta Diversity. To detect any implicit spatial structure in $P\beta_{total}$, $P\beta_{rich}$, $P\beta_{repl}$, and

COMDIST, we visualized these metrics through UPGMA clustering (Borcard et al. 2011)

using the 'hclust' function in R (R Core Team 2017). When applied to $P\beta_{rich}$, UPGMA will

cluster assemblages with similar amount of phylogenetic richness whereas $P\beta_{repl}$ will cluster

assemblages with similar lineages. For COMDIST, UPGMA will cluster closely related

assemblages. A Mantel test (Mantel 1967) was conducted using the R package 'vegan'

(Oksanen et al. 2017) to detect any linear (Pearson correlation) or non-linear (Spearman's

rank correlation) spatial structure in $P\beta_{total}$, $P\beta_{rich}$, $P\beta_{repl}$, and COMDIST.

Results

Comparison within assemblages: phylogenetic diversity is lowest in the smallest fragments

Rarefied PD showed a decreasing trend from continuous forest and fragment interiors to

forest edges and matrix, with the matrix sites adjacent to the 1 ha fragments harbouring the

lowest PD overall (Fig. 1). In agreement with this, the model including IEM gradient and

forest size as additive predictors of PD received overwhelming support ($w_i = 0.99$) compared

to the other candidate models (Table 1). The decrease in PD from interior towards edge and

matrix was significant ($P < 0.001$ for both), with edges experiencing larger decreases in PD

compared to the matrix (Table S2). However, multiple comparison tests did not support any

significant differences in PD between edges and matrix, irrespective of forest size (Table S3).

Also, there was no significant loss in PD from larger towards smaller fragments except for 1

ha fragments ($P < 0.001$, Table S2). Multiple comparison tests confirmed the significantly

lower phylogenetic richness in 1 ha fragments relative to larger fragments and continuous

forest across the entire IEM gradient (Table S3). According to the local phylogeny, most of

the old species lineages, i.e. those with long tree branches, do not appear in the 1 ha fragments

(Fig. S1).

When the effect of species richness on PD was accounted for (SES_{PD}), only the IEM gradient

emerged as the best-supported predictor in the model ($w_i = 0.78$, Table 1). SES_{PD} for forest

fragments showed lower phylogenetic richness than expected given the number of species

present, particularly for fragment edges and the matrix (Fig 2A). SES_{PD} was also significantly

lower in the matrix than in forest interiors, particularly for the 1 ha fragments (Fig 2A).

In line with PD, observed MPD of phyllostomid assemblages in most habitat types was lower

than expected under the simulated null communities, particularly for the interiors of 1 ha

fragments, fragment edges, and the matrix around the larger (10 and 100 ha) fragments (Fig.

2B). Only the assemblages in the matrix surrounding 1 ha fragments, although characterized

as relatively phylogenetically over-dispersed compared to the other assemblages based on

SES_{MPD}, showed terminal clustering based on its significantly lower SES_{MNTD} (Fig. 2C).

Patterns of MPD were best explained by a model that included only the IEM gradient as

explanatory variable ($w_i = 0.33$, Table 1). There was, however, substantial model selection

uncertainty, with the null, size, and IEM + size models also being supported as plausible

($\Delta AIC_c < 2$, Table 1). According to the single variable models IEM gradient and size (Table

S2), MPD was significantly reduced in the matrix ($P = 0.03$) and 1 ha fragments ($P = 0.04$),

respectively, suggesting that the matrix around 1 ha fragments contains more closely related

lineages as indicated by its significantly lower SES_{MNTD}. For SES_{MPD}, the effect of the IEM

gradient was even more tenuous ($w_i = 0.36$) as the null model received the strongest support

($w_i = 0.57$, Table 1).

Comparison between assemblages: no clear pattern of lineage replacement

Dendrograms based on the total evolutionary history shared between assemblages ($P\beta_{total}$,

$P\beta_{rich}$, and $P\beta_{repl}$) were substantially different from the one based on relatedness between

lineages within assemblages (COMDIST). UPGMA clustering based on total phylogenetic

beta diversity ($P\beta_{total}$) suggested that the interiors of continuous forest and forest fragments

harbour similar amounts of phylogenetic richness, except for the 1 ha fragments (Fig 3A),

further confirming the phylogenetic erosion of 1 ha fragments. The similarity in total

phylogenetic richness between the interiors of continuous forest and larger forest fragments

(10 and 100 ha) was maintained after $P\beta_{total}$ was partitioned into $P\beta_{rich}$ and $P\beta_{repl}$. For $P\beta_{rich}$,

however, the interior sites of the larger fragments clustered together, unlike $P\beta_{total}$ which

grouped the interior of continuous forest closer together with that of 100 ha fragments (Fig

3B). For $P\beta_{repl}$, the interior sites were completely dispersed (Fig 3C), suggesting that the

difference in $P\beta_{total}$ compared to $P\beta_{rich}$ is caused by different lineages contained within the

interiors. The position of the IEM habitat categories in the $P\beta_{repl}$ dendrogram are unlikely due

to their geographic proximity as there was no significant relationship between $P\beta_{repl}$ and

geographic distance based on the Mantel test (Pearson's Mantel statistic r = 0.031, P = 0.299,

Table S4).

UPGMA clustering of COMDIST revealed that the investigated assemblages were closely

related and were not clustered according to either the same IEM gradient or the same forest

size categories (Fig. 4). The sites within the interior of 1 ha fragment are more closely related

to the matrix and edges of forest fragments and continuous forest. The position of the interior

sites of 1 ha fragments distant from those of larger fragments and continuous forest,

underscores the distinct composition of bat lineages in the assemblages of 1 ha fragments.

When the local phylogeny was annotated with information on species' presence across the

two environmental gradients, different lineages were present within the same IEM category

except for the interior of continuous forest and the larger forest fragments (Fig. S1). The

interiors of 1 ha fragments were distinct from other interior sites whereas there was no clear

pattern of lineage distribution at forest edges and in the matrix adjoining continuous forest

and fragments. This is also unlikely due to spatial effects as there was no significant

relationship between COMDIST and geographic distance (Pearson's Mantel statistic r =

0.055, P = 0.230, Table S4).

Discussion

Using metrics of phylogenetic alpha and beta diversity, we showed that there was not only a disproportionate reduction of phylogenetic richness in the smallest forest fragments (1 ha) but also phylogenetic homogenization of assemblages at forest edges and in the matrix as assemblages in these categories of the IEM gradient contained closely related bat lineages. Whereas the two environmental gradients investigated in this study explained quite well the observed variation in total phylogenetic richness (PD), they could not adequately account for how closely related lineages clustered in small forest fragments (MPD). Additionally, the results of the clustering analysis based on phylogenetic beta diversity metrics were counter to our predictions since the resulting dendrograms did not clearly cluster assemblages from the same category of habitat quality and forest size together.

Environmental filtering is strongest in the matrix surrounding small forest fragments

Across the full disturbance gradient of interior-edge-matrix, phylogenetic richness was curtailed in both edge and matrix habitats, especially in the case of the 1 ha fragments. Thus, habitat fragmentation at the BDFFP does not only reduce bat taxonomic and functional diversity in the small forest fragments, edges, and matrix as we have previously shown (Farneda et al. 2015; Rocha et al. 2017), but also negatively affects the total evolutionary history preserved. However, the magnitude of loss of phylogenetic richness at fragment edges and in the adjacent matrix relative to the interior of continuous forest was only slight, probably due to the low structural contrast between matrix (tall secondary forest) and edges during the sampling period (Rocha et al. 2017). In accordance with other studies comparing phylogenetic richness between various human-modified habitats of different levels of forest/vegetation complexity (Riedinger et al. 2013; Frishkoff et al. 2014; Edwards et al. 2015, 2017; Frank et al. 2017; Martins et al. 2017; Pereira et al. 2018), the observed decrease in total evolutionary history in a structurally less complex habitat such as the secondary forest

matrix is not surprising. Phylogenetic diversity is often associated with canopy cover (Frank

et al. 2017; Martins et al. 2017), which decreased along our IEM gradient (cf Table S2, Rocha

et al. 2017). This phenomenon is often attributed to environmental filtering, whereby species

with particular characteristics are not well adapted to and thus unable to persist in all parts of

an environmental gradient (Cisneros et al. 2016; Frank et al. 2017).

Through the use of SES_{MPD} and SES_{MNTD}, we further detected that edge and matrix

assemblages were phylogenetically clustered, whereby the matrix of 1 ha fragments was

characterized by especially closely related lineages, pointing towards stronger environmental

filtering compared to the edge. Matrix sites surrounding 1 ha fragments are impoverished with

respect to lineages with long evolutionary history, i.e. *Lampronycteris brachyotis* and several

Micronycteris species (Fig S1), and contain lineages that are phylogenetically clustered in the

terminal branches. Nevertheless, it is important to note that matrix sites adjacent to the larger

fragments (10 ha and 100 ha) did not differ much from their edges in terms of phylogenetic

richness and structure, indicating that the habitat quality gradient investigated in this study

represents a strong environmental filter in both large (≥10 ha) and small (1 ha) forest

fragments.

Our model selection results, however, did not provide strong support for the environmental

gradient investigated in this study to explain the observed phylogenetic clustering, which was

weak as the values of SES_{MPD} and and SES_{MNTD} did not deviate much from the 5[th] and 95[th]

quantiles of the null model. This is potentially due to the influence of landscape-level

attributes which encompass wider environmental gradients (Rocha et al. 2017; Tinoco et al.

2018) and are responsible for the ability of the large fragments to retain more distantly related

lineages than the small forest fragments. The low structural contrast between secondary

regrowth in the matrix and the forest interiors during the sampling period could also attenuate

any strong phylogenetic structure of bat assemblages along the interior-edge-matrix gradient;

(Patrick and Stevens 2016) for instance found that environmental variables have a weaker

effect upon phylogenetic structure when assemblages experience less harsh environmental

conditions. Additionally, the environmental filtering that is responsible for phylogenetic

clustering could possibly result from non-linear responses (Smith and Lundholm 2010; Stegen

and Hurlbert 2011) that we were unable to detected using linear models. Stronger support of

an effect of the IEM gradient in explaining phylogenetic structure may have been received if

we had considered abundance-based metrics and quantitative variables such as seasonal

temperature (Stevens and Gavilanez 2015) or elevation (Cisneros et al. 2014). Nonetheless,

our findings still confirm our expectations that the low resource availability in the matrix

apparently selects for frugivorous bat lineages that have traits favourable by the existing

environmental filter (Farneda et al. 2015).

There is a possibility that the phylogenetic clustering documented in small fragments also

reflects biotic filtering through competitive exclusion of species with low competitive ability

in addition to abiotic filtering through selection by the IEM gradient (Mayfield and Levine

2010). However, we consider this unlikely to be the case for phyllostomid bats which are

known to have a high diversification rate compared to other bat families (Monteiro and

Nogueira 2011; Santana et al. 2012). The wide array of feeding strategies exhibited by

phyllostomids likely eliminates the need for strong competition between species as pre-

requisite of inferring biotic filtering. For bats at the BDFFP in particular, traits associated with

fragmentation sensitivity such as trophic level, body mass, and wing morphology were

phylogenetically conserved (Farneda et al. 2015) so that the observed clustering could be

related to environmental filtering.

By investigating two measures of phylogenetic alpha diversity simultaneously (PD and

MPD), we showed that assemblages that have a relatively lower amount of phylogenetic

richness (PD) compared to those of the interior of continuous forest also are made up of more

closely related lineages as shown by the relative increase in phylogenetic clustering (MPD).

This is consistent with Frank et al (2017) and Riedinger et al (2013) who pointed out the

tendency of closely-related bat species to respond similarly to habitat disturbance due to the

similarity in traits that are evolutionarily conserved.

Variation in phylogenetic diversity between assemblages is not related to habitat quality

Phylogenetic beta diversity metrics further corroborate the weak effect of habitat quality on

between-assemblage differences in phylogenetic richness and lineage composition. There was

no clear-cut pattern with regard to the relatedness and phylogenetic richness of bat

assemblages in relation to IEM and forest size gradients. Except for 1 ha fragments, the

interiors of the forest fragments and continuous forest contain a large number of distantly

related species and harbour a similar amount of total evolutionary history, but not the same

bat lineages. Larger forest fragments (≥10 ha) may therefore act as important repositories for

preserving the total evolutionary history of bat assemblages in fragmented landscapes. In

contrast, the marked changes in phylogenetic richness and structure of 1 ha fragment interiors

and similarity to their edges in terms of preserved evolutionary history, suggests that patterns

of phylogenetic diversity are fundamentally driven by the pervasive negative edge effects that

commonly plague fragments of this size (Santos et al. 2010; Laurance et al. 2018). We

probably did not detect any spatially implicit structure between different habitat categories of

the IEM gradient due to the predominant effect of dispersal ability in the absence of

contrasting environmental gradients (Moreno and Halffter 2001).

The distribution of edge and matrix sites associated with the larger forest fragments and

continuous forest across the dendrograms are possibly the result of phylogenetic

homogenization. Assemblages of these two categories of the IEM gradient did not differ

markedly in phylogenetic richness and structure according to the phylogenetic alpha diversity

metrics, and UPGMA clustering further confirmed that they are closely related to each other

(Fig. 4). Thus, despite the ability of large forest fragments to retain amounts of total

evolutionary history similar to the interior of continuous forest, the level of forest degradation

at the edges and in the matrix still has a negative effect on the phylogenetic diversity of bats at

the BDFFP.

New insights gained from a focus on the phylogenetic biodiversity dimension

A focus on the phylogenetic dimension of biodiversity has allowed us to complement our

previous analyses that approached bat responses to landscape fragmentation at the BDFFP

from taxonomic and functional perspectives (Rocha et al. 2017; Farneda et al. 2018). The

reduction of total evolutionary history in the smallest fragments, particularly in the

surrounding matrix, was apparently selective and resulted in assemblages comprised of

closely related bat lineages with similar traits due to environmental filtering. This adds more

significance to the term "biodiversity loss" in fragmented landscapes as losing a lineage of

species usually means losing a unique evolutionary history which may be irreplaceable for a

particular habitat and disrupt ecosystem processes therein. The non-significant decrease of

phylogenetic richness in larger fragments (10 ha and 100 ha) compared to continuous forest

indicates that fragments of that size still serve as important refuges for the overall lineage

pool in the BDFFP landscape. Although the habitat quality and size gradients investigated in

this study could not entirely explain the pattern of phylogenetic beta diversity, cluster analysis

showed their constituent categories to harbour distinct assemblages that are probably shaped

by a specific assembly mechanism.

Any environmental filtering detected in this study, however, may not be generalizable to other

fragmented landscapes. The BDFFP as an experimentally controlled fragmented landscape is

a best-case scenario as levels of disturbance are reduced in relation to most real-world

landscapes (Laurance et al. 2018). Additionally, forest fragments are surrounded by a benign

matrix dominated by forest regrowth (Mesquita et al. 2015), and which is largely free from

human disturbances which could interact additively or synergistically with fragmentation

(Laurance et al. 2018). The matrix was dominated by tall secondary forest regrowth at the

time of study (Rocha et al. 2017), resulting in a rather homogenous makeup of the overall

landscape which could promote random placement in bat assemblages (Morlon et al. 2011).

Nonetheless, the most challenging part in investigating environmental filtering using

phylogenetic diversity metrics is deciding which aspect of the environment actually matters

for the taxa in question and on which ecological and evolutionary scale to better understand

the evolutionary consequences of habitat fragmentation.

Phylogenetic diversity summarizes species characteristics that are not covered by taxon and

functional diversity (Flynn et al. 2011; Cisneros et al. 2014), especially the ecological

redundancies and complementarity of organisms (Cadotte et al. 2013). Our choice of

incidence-based metrics, however, was limited to assess among-species variation and could

not account for within-species variation that is better captured by abundance-based metrics.

Abundance-based phylogenetic diversity metrics are more sensitive to environmental

gradients (Stevens and Gavilanez 2015) and could correct for stochasticity in the detection of

rare taxa or sample effects that have been observed for various taxa at BDFFP (Laurance et al.

2018).

Our study adds to a growing body of evidence suggesting biotic homogenization in human-

modified landscapes is widespread not only with regard to the taxonomic facet of biodiversity

but also its evolutionary dimension (La Sorte et al. 2018; Park and Razafindratsima 2019).

Despite the limitation of our methods, we showed fragmentation to result in the loss of

species lineages and phylogenetic homogenization of assemblages in degraded edge and

matrix habitats. This could further affect ecosystem stability as communities where species

are evenly and distantly related to one another are more stable compared to communities

where phylogenetic relationships are more clumped (Cadotte et al. 2012). The conservation of

large forest fragments and the improvement of habitat quality thus should be priority in

managing fragmented forest landscapes to conserve species with longer evolutionary history.

Acknowledgements

We are in debt to the many volunteers, students and field assistants that helped us during

fieldwork and the coordination team of the BDFFP. We also thank Paulo E.D. Bobrowiec for

logistic support and Dirk Metzler for his feedback in the statistical analysis.

Funding: The Portuguese Foundation for Science and Technology provided funds to C.F.J.M.

(PTDC/BIA-BIC/111184/2009), R.R. (SFRH/BD/80488/2011) and A.L.-B.

(PD/BD/52597/2014). S.G.A was supported by the Indonesia Endowment Fund for Education

(LPDP). This research was conducted under ICMBio permit (26877-2) and constitutes

publication number XXX in the BDFFP technical series.

Conflict of Interest: The authors declare that they have no conflict of interest.

References

Aguirre LF, Montaño-Centellas FA, Gavilanez MM, Stevens RD (2016) Taxonomic and

Phylogenetic Determinants of Functional Composition of Bolivian Bat Assemblages.

PLoS One 11:e0158170. doi: 10.1371/journal.pone.0158170

Alroy J (2017) Effects of habitat disturbance on tropical forest biodiversity. PNAS 114:6056–

6061. doi: 10.1073/pnas.1611855114

Austin KG, González-Roglich M, Schaffer-Smith D, et al (2017) Trends in size of tropical

deforestation events signal increasing dominance of industrial-scale drivers. Environ

Res Lett 12:79601. doi: 10.1088/1748-9326/aa7760

Barlow J, França F, Gardner TA, et al (2018) The future of hyperdiverse tropical ecosystems.

Nature 559:517. doi: 10.1038/s41586-018-0301-1

Barton K (2016) MuMIn: Multi-Model Inference.

Bates D, Mächler M, Bolker B, Walker S (2014) Fitting linear mixed-effects models using

lme4. arXiv preprint arXiv:14065823

Bolker BM, Brooks ME, Clark CJ, et al (2009) Generalized linear mixed models: a practical

guide for ecology and evolution. Trends Ecol Evol 24:127–135. doi:

10.1016/j.tree.2008.10.008

Borcard D, Gillet F, Legendre P (2011) Numerical Ecology with R. Springer New York, New

York, NY

Burnham KP, Anderson DR (2002) Model selection and multimodel inference: a practical

information-theoretic approach, 2nd ed. Springer, New York

Cadotte MW, Albert CH, Walker SC (2013) The ecology of differences: assessing community

assembly with trait and evolutionary distances. Ecol Lett 16:1234–1244. doi:

10.1111/ele.12161

Cadotte MW, Dinnage R, Tilman D (2012) Phylogenetic diversity promotes ecosystem

stability. Ecology 93:S223–S233

Cardoso P, Rigal F, Carvalho JC (2015) BAT - Biodiversity Assessment Tools, an R package

for the measurement and estimation of alpha and beta taxon, phylogenetic and

functional diversity. Methods in Ecology and Evolution 6:232–236. doi:

10.1111/2041-210X.12310

Cardoso P, Rigal F, Carvalho JC, et al (2014) Partitioning taxon, phylogenetic and functional

481   beta diversity into replacement and richness difference components. J Biogeogr

482   41:749–761. doi: 10.1111/jbi.12239

483   Cisneros LM, Burgio KR, Dreiss LM, et al (2014) Multiple dimensions of bat biodiversity

484   along an extensive tropical elevational gradient. J Anim Ecol 83:1124–1136. doi:

485   10.1111/1365-2656.12201

486   Cisneros LM, Fagan ME, Willig MR (2015) Effects of human-modified landscapes on

487   taxonomic, functional and phylogenetic dimensions of bat biodiversity. Divers Distrib

488   21:523–533. doi: 10.1111/ddi.12277

489   Cisneros LM, Fagan ME, Willig MR (2016) Environmental and spatial drivers of taxonomic,

490   functional, and phylogenetic characteristics of bat communities in human-modified

491   landscapes. PeerJ 4:e2551. doi: 10.7717/peerj.2551

492   Clarke KR, Warwick RM (1998) A taxonomic distinctness index and its statistical properties.

493   Journal of Applied Ecology 35:523–531

494   De Oliveira AA, Mori SA (1999) A central Amazonian terra firme forest. I. High tree species

495   richness on poor soils. Biodivers Conserv 8:1219–1244

496   Edwards DP, Gilroy JJ, Thomas GH, et al (2015) Land-Sparing Agriculture Best Protects

497   Avian Phylogenetic Diversity. Current Biology 25:2384–2391. doi:

498   10.1016/j.cub.2015.07.063

499   Edwards DP, Massam MR, Haugaasen T, Gilroy JJ (2017) Tropical secondary forest

500   regeneration conserves high levels of avian phylogenetic diversity. Biological

501   Conservation 209:432–439. doi: 10.1016/j.biocon.2017.03.006

502   Ewers RM, Didham RK (2006) Confounding factors in the detection of species responses to

503   habitat fragmentation. Biol Rev 81:117. doi: 10.1017/S1464793105006949

504   Fahrig L (2017) Ecological responses to habitat fragmentation per se. Annu Rev Ecol Evol

505   Syst 48:1–23. doi: 10.1146/annurev-ecolsys-110316- 022612

506   Faith DP (1992) Conservation evaluation and phylogenetic diversity. Biological Conservation

507   61:1–10

508   Farneda FZ, Rocha R, López-Baucells A, et al (2015) Trait-related responses to habitat

509   fragmentation in Amazonian bats. J Appl Ecol 52:1381–1391. doi: 10.1111/1365-

510   2664.12490

511   Farneda FZ, Rocha R, López-Baucells A, et al (2018) Functional recovery of Amazonian bat

512   assemblages following secondary forest succession. Biol Conserv 218:192–199. doi:

513   10.1016/j.biocon.2017.12.036

514   Ferreira DF, Rocha R, López-Baucells A, et al (2017) Season-modulated responses of

515   Neotropical bats to forest fragmentation. Ecol Evol 7:4059–4071. doi:

516   10.1002/ece3.3005

517   Flynn DF, Mirotchnick N, Jain M, et al (2011) Functional and phylogenetic diversity as

518   predictors of biodiversity–ecosystem-function relationships. Ecology 92:1573–1581

519   Frank HK, Frishkoff LO, Mendenhall CD, et al (2017) Phylogeny, Traits, and Biodiversity of

520   a Neotropical Bat Assemblage: Close Relatives Show Similar Responses to Local

521   Deforestation. Am Nat 190:0–0. doi: 10.1086/692534

522   Frishkoff LO, Karp DS, M'Gonigle LK, et al (2014) Loss of avian phylogenetic diversity in

523   neotropical agricultural systems. Science 345:1343–1346

524   Graham CH, Fine PVA (2008) Phylogenetic beta diversity: linking ecological and

525   evolutionary processes across space in time. Ecol Lett 11:1265–1277. doi:

526   10.1111/j.1461-0248.2008.01256.x

Haddad NM, Brudvig LA, Clobert J, et al (2015) Habitat fragmentation and its lasting impact

on Earth's ecosystems. Science Advances 1:e1500052–e1500052. doi:

10.1126/sciadv.1500052

Hothorn T, Bretz F, Westfall P (2008) Simultaneous inference in general parametric models.

Biom J 50:346–363

Jones KE, Bininda-Emonds ORP, Gittleman JL (2005) Bats, clocks, and rocks: diversification

patterns in Chiroptera. Evolution 59:2243. doi: 10.1554/04-635.1

Jones KE, Purvis A, MacLarnon A, et al (2002) A phylogenetic supertree of the bats

(Mammalia: Chiroptera). Biol Rev 77:223–259

Kalko EKV (1998) Organisation and diversity of tropical bat communities through space and

time. Zoology 101:281–297

Kembel SW, Cowan PD, Helmus MR, et al (2010) Picante: R tools for integrating

phylogenies and ecology. Bioinformatics 26:1463–1464. doi:

10.1093/bioinformatics/btq166

Kunz TH, Braun de Torrez E, Bauer D, et al (2011) Ecosystem services provided by bats.

Annals of the New York Academy of Sciences 1223:1–38. doi: 10.1111/j.1749-

6632.2011.06004.x

La Sorte FA, Lepczyk CA, Aronson MFJ, et al (2018) The phylogenetic and functional

diversity of regional breeding bird assemblages is reduced and constricted through

urbanization. Divers Distrib 24:928–938. doi: 10.1111/ddi.12738

Laurance SGW, Laurance WF, Andrade A, et al (2010) Influence of soils and topography on

Amazonian tree diversity: a landscape-scale study. J Veg Sci 21:96–106. doi:

10.1111/j.1654-1103.2009.01122.x

Laurance WF, Camargo JLC, Fearnside PM, et al (2018) An Amazonian rainforest and its

fragments as a laboratory of global change. Biol Rev 93:223–247. doi:

10.1111/brv.12343

Mantel N (1967) The detection of disease clustering and a generalized regression approach.

Cancer Res 27:209–220

Martins ACM, Willig MR, Presley SJ, Marinho-Filho J (2017) Effects of forest height and

vertical complexity on abundance and biodiversity of bats in Amazonia. For Ecol

Manage 391:427–435. doi: 10.1016/j.foreco.2017.02.039

Mayfield MM, Levine JM (2010) Opposing effects of competitive exclusion on the

phylogenetic structure of communities: Phylogeny and coexistence. Ecol Lett

13:1085–1093. doi: 10.1111/j.1461-0248.2010.01509.x

Mesquita R de CG, Massoca PE dos S, Jakovac CC, et al (2015) Amazon Rain Forest

Succession: Stochasticity or Land-Use Legacy? BioScience 65:849–861. doi:

10.1093/biosci/biv108

Meyer CFJ, Struebig MJ, Willig MR (2016) Responses of Tropical Bats to Habitat

Fragmentation, Logging, and Deforestation. In: Voigt CC, Kingston T (eds) Bats in

the Anthropocene: Conservation of Bats in a Changing World. Springer International

Publishing, Cham, pp 63–103

Monteiro LR, Nogueira MR (2011) Evolutionary patterns and processes in the radiation of

phyllostomid bats. BMC Evol Biol 11:137

Moreno CE, Halffter G (2001) Spatial and temporal analysis of α, β and γ diversities of bats

in a fragmented landscape. Biodivers Conserv 10:367–382

Morlon H, Schwilk DW, Bryant JA, et al (2011) Spatial patterns of phylogenetic diversity.

Ecol Lett 14:141–149. doi: 10.1111/j.1461-0248.2010.01563.x

Oksanen J, Blanchet FG, Friendly M, et al (2017) vegan: Community Ecology Package

Paradis E, Claude J, Strimmer K (2004) APE: Analyses of Phylogenetics and Evolution in R

language. Bioinformatics 20:289–290. doi: 10.1093/bioinformatics/btg412

Park DS, Razafindratsima OH (2019) Anthropogenic threats can have cascading

homogenizing effects on the phylogenetic and functional diversity of tropical

ecosystems. Ecography 42:148–161. doi: 10.1111/ecog.03825

Patrick LE, Stevens RD (2016) Phylogenetic community structure of North American desert

bats: influence of environment at multiple spatial and taxonomic scales. J Anim Ecol

85:1118–1130. doi: 10.1111/1365-2656.12529

Pellens R, Grandcolas P (2016) Phylogenetics and Conservation Biology: Drawing a Path into

the Diversity of Life. In: Biodiversity Conservation and Phylogenetic Systematics:

Preserving our evolutionary heritage in an extinction crisis. Springer Open, pp 1–15

Pereira MJR, Fonseca C, Aguiar LMS (2018) Loss of multiple dimensions of bat diversity

under land-use intensification in the Brazilian Cerrado. HYSTRIX 29:25–32. doi:

10.4404/hystrix–00020-2017

Presley SJ, Cisneros LM, Higgins CL, et al (2018) Phylogenetic and functional

underdispersion in Neotropical phyllostomid bat communities. Biotropica 50:135–

145. doi: 10.1111/btp.12501

R Core Team (2017) R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing, Vienna, Austria

Redondo RAF, Brina LPS, Silva RF, et al (2008) Molecular systematics of the genus Artibeus

(Chiroptera: Phyllostomidae). Mol Phylogenet Evol 49:44–58. doi:

10.1016/j.ympev.2008.07.001

Riedinger V, Müller J, Stadler J, et al (2013) Assemblages of bats are phylogenetically

clustered on a regional scale. Basic Appl Ecol 14:74–80. doi:

10.1016/j.baae.2012.11.006

Ries L, Sisk TD (2004) A predictive model of edge effects. Ecology 85:2917–2926

Rocha R, López-Baucells A, Farneda FZ, et al (2017) Consequences of a large-scale
    fragmentation experiment for Neotropical bats: disentangling the relative importance
    of local and landscape-scale effects. Landscape Ecol 32:31–45. doi: 10.1007/s10980-
    016-0425-3

Rocha R, Ovaskainen O, López-Baucells A, et al (2018) Secondary forest regeneration
    benefits old-growth specialist bats in a fragmented tropical landscape. Sci Rep 8:3819.
    doi: 10.1038/s41598-018-21999-2

Santana SE, Grosse IR, Dumont ER (2012) Dietary hardness, loading behavior, and the
    evolution of skull form in bats. Evolution 66:2587–2598. doi: 10.1111/j.1558-
    5646.2012.01615.x

Santos BA, Arroyo-Rodríguez V, Moreno CE, Tabarelli M (2010) Edge-Related Loss of Tree
    Phylogenetic Diversity in the Severely Fragmented Brazilian Atlantic Forest. PLoS
    ONE 5:e12625. doi: 10.1371/journal.pone.0012625

Santos BA, Tabarelli M, Melo FPL, et al (2014) Phylogenetic Impoverishment of Amazonian
    Tree Communities in an Experimentally Fragmented Forest Landscape. PLoS ONE
    9:e113109. doi: 10.1371/journal.pone.0113109

Shi JJ, Rabosky DL (2015) Speciation dynamics during the global radiation of extant bats.
    Evolution 69:1528–1545. doi: 10.1111/evo.12681

Smith TW, Lundholm JT (2010) Variation partitioning as a tool to distinguish between niche
    and neutral processes. Ecography 33:648–655. doi: 10.1111/j.1600-
    0587.2009.06105.x

Stegen JC, Hurlbert AH (2011) Inferring Ecological Processes from Taxonomic, Phylogenetic
    and Functional Trait β-Diversity. PLoS ONE 6:e20906. doi:
    10.1371/journal.pone.0020906

Stevens RD, Gavilanez MM (2015) Dimensionality of community structure: phylogenetic,

morphological and functional perspectives along biodiversity and environmental

gradients. Ecography 38:861–875. doi: 10.1111/ecog.00847

Swenson NG (2014) Functional and Phylogenetic Ecology in R. Springer New York, New

York, NY

Swenson NG (2011) Phylogenetic Beta Diversity Metrics, Trait Evolution and Inferring the

Functional Beta Diversity of Communities. PLoS ONE 6:e21264. doi:

10.1371/journal.pone.0021264

Tinoco BA, Santillán VE, Graham CH (2018) Land use change has stronger effects on

functional diversity than taxonomic diversity in tropical Andean hummingbirds. Ecol

Evol 8:3478–3490. doi: 10.1002/ece3.3813

Tucker CM, Cadotte MW, Carvalho SB, et al (2017) A guide to phylogenetic metrics for

conservation, community ecology and macroecology: A guide to phylogenetic metrics

for ecology. Biol Rev 92:698–715. doi: 10.1111/brv.12252

Vamosi SM, Heard SB, Vamosi JC, Webb CO (2009) Emerging patterns in the comparative

analysis of phylogenetic community structure. Mol Ecol 18:572–592. doi:

10.1111/j.1365-294X.2008.04001.x

Webb CO (2000) Exploring the phylogenetic structure of ecological communities: an

example for rain forest trees. Am Nat 156:145–155

Young HS, McCauley DJ, Galetti M, Dirzo R (2016) Patterns, Causes, and Consequences of

Anthropocene Defaunation. Annu Rev Ecol Evol Syst 47:333–358. doi:

10.1146/annurev-ecolsys-112414-054142

Fig. 1 Comparison of rarefied phylogenetic richness of bat assemblages across gradients of

habitat quality (IEM) and habitat amount (forest size) at the Biological Dynamics of Forest

Fragmentation Project, Brazil. Horizontal bars indicate the median of rarefied values, the

mean of observed values for each habitat category is depicted as 'x'. Asterisks denote

significant differences relative to continuous forest interiors (***$P<0.001$; **$P<0.01$; *P

<0.05).

Fig. 2 Standardized effect size (dots) of (A) Faith's phylogenetic diversity (PD), (B) mean

pairwise distance (MPD), and (C) mean nearest taxon distance (MNTD) along with 5^{th} and

95^{th} quantiles (dashed lines) of the simulated null communities (box and whisker plots). In the

box plots, values of observed PD, MPD, and MNTD overlaid on the simulated null

communities are indicated by a black diamond.

Fig. 3 Unweighted Pair Group Method with Arithmetic Mean (UPGMA) clustering of bat

total phylogenetic beta richness ($P\beta_{tot}$) (A) in the Biological Dynamics of Forest Fragments

Project landscape, partitioned into its richness ($P\beta_{rich}$) (B) and replacement ($P\beta_{repl}$) (C)

components.

Fig. 4 Unweighted Pair Group Method with Arithmetic Mean (UPGMA) clustering of

COMDIST metric representing the divergence aspects of phylogenetic beta diversity of bats

in the Biological Dynamics of Forest Fragments Project landscape.

Table 1. Results of Akaike Information Criterion (AIC)-based model selection assessing support for an effect of habitat quality (IEM) and habitat amount (size) on phylogenetic richness (PD and SES_{PD}) and divergence (MPD and SES_{MPD}) of bat assemblages at the Biological Dynamics of Forest Fragments Project, Brazil. For each model, the log-likelihood (Log-L), number of estimable parameters (K), sample-size adjusted AIC (AICc), Akaike differences (Δ_i), and Akaike weights (w_i) are presented.

Response variable	Model	K	logLik	AICc	Δ_i	w_i
PD	IEM + size	7	-206.78	431.20	0	0.99
	IEM * size	13	-199.86	440.30	9.09	0.01
	IEM	4	-220.36	449.90	18.73	0
	size	5	-220.46	452.70	21.56	0
	null	2	-230.45	465.20	34.06	0
SES_{PD}	IEM	4	-50.78	110.74	0	0.78
	null	2	-54.95	114.23	3.49	0.14
	IEM + size	7	-49.02	115.65	4.92	0.07
	size	5	-53.49	118.79	8.06	0.01
	IEM * size	13	-45.56	131.69	20.95	0

MPD	IEM	4	-88.24	185.66	0	0.33
	null	2	-90.99	186.31	0.65	0.24
	size	5	-87.28	186.37	0.71	0.23
	IEM + size	7	-84.54	186.70	1.04	0.20
	IEM * size	13	-78.14	196.83	11.17	0
SES$_{MPD}$	null	2	-55.84	116.01	0	0.57
	IEM	4	-53.89	116.96	0.95	0.36
	size	5	-54.54	120.90	4.89	0.05
	IEM + size	7	-52.43	122.46	6.46	0.02
	IEM * size	13	-45.75	132.06	16.05	0

Fig. 1

Fig. 2

A Pβ_{total}

B Pβ_{rich}

C Pβ_{repl}

Fig. 3

inter-assemblages MPD

Fig. 4