HOPS: Automated detection and authentication of pathogen DNA in

archaeological remains

Ron Hübler[1*] (huebler@shh.mpg.de), Felix M. Key[1,2,3*] (key@shh.mpg.de), Christina

Warinner[1] (warinner@shh.mpg.de), Kirsten I. Bos[1] (bos@shh.mpg.de), Johannes

Krause[1] (krause@shh.mpg.de), Alexander Herbig[1] (herbig@shh.mpg.de)

Affiliation

1. Max-Planck Institute for the Science of Human History

2. Institute for Medical Engineering and Sciences, Massachusetts Institute of

Technology, Cambridge, Massachusetts 02139, USA

3. Department of Civil and Environmental Engineering, Massachusetts Institute

of Technology, Cambridge, Massachusetts 02139, USA

* Equal contribution

Correspondence: Felix M. Key (key@shh.mpg.de), Alexander Herbig

(herbig@shh.mpg.de)

Abstract

High-throughput DNA sequencing enables large-scale metagenomic analyses of

complex biological systems. Such analyses are not restricted to present day

environmental or clinical samples, but can also be fruitfully applied to molecular data

from archaeological remains (ancient DNA), and a focus on ancient bacteria can

provide valuable information on the long-term evolutionary relationship between

hosts and their pathogens. Here we present HOPS (**H**euristic **O**perations for **P**athogen

Screening), an automated bacterial screening pipeline for ancient DNA sequence data

that provides straightforward and reproducible information on species identification

and authenticity. HOPS provides a versatile and fast pipeline for high-throughput

screening of bacterial DNA from archaeological material to identify candidates for

subsequent genomic-level analyses.

Keywords

ancient DNA, archaeogenetics, pathogen detection, metagenomics

Background

High-throughput DNA sequencing enables large-scale metagenomic analyses of environments and host tissues, providing an unprecedented understanding of life's microbial diversity. Examples of coordinated efforts to quantify this diversity include the Human Microbiome Project (1), the Tara Ocean Project (2) and the Earth Microbiome Project (3). Metagenomic data from human archaeological remains (*e.g.* bones, teeth or dental calculus), which provide a window into the individuals' metagenomic past, is a welcome addition to the wide landscape of microbial diversity now being revealed. While many ancient DNA (aDNA) studies focus on the analysis of human endogenous DNA isolated from ancient specimens (4-8), the co-recovered metagenomic aDNA can be queried to provide information related to endogenous microbial content at death, with applications ranging from characterizing the natural constituents of the microbiota to identifying systemic infectious diseases (9, 10).

Genomic-level investigations of ancient pathogens have provided valuable information about the evolution of *Yersinia pestis* (11-18), *Mycobacterium leprae* (19, 20), *Mycobacterium tuberculosis* (21, 22), pathogenic *Brucella* species (23, 24), *Salmonella enterica* (25, 26) and *Helicobacter pylori* (27), with others surely on the horizon. Notably, most studies to date have leveraged paleopathological evidence or historical context to pinpoint *a priori* involvement of a specific bacterial pathogen. However, the vast majority of infectious diseases do not lead to the formation of distinct and characteristic bone lesions, and most remains are found in contexts that lack clear associations with a particular disease. Consequently, studies of ancient pathogens must consider a long list of candidate microbes. Therefore, an automated computational screening tool that both detects and evaluates pathogen genetic signals in ancient metagenomic data is needed. Importantly, this tool should also be able to distinguish potential pathogens from the large and diverse microbial background typical of archaeological and other decomposed material, a consideration not typically required for tools developed for clinical applications.

To save computational time and effort, most available metagenomic profiling tools

focus only on individual genes, such as the 16S rRNA gene used by QIIME (28), or

panels of marker genes, such as those used by MetaPhlAn2 (29) and MIDAS (30),

that are information-rich and highly species-specific. However, these genes make up

only a small proportion of a bacterial genome (the 16S rRNA gene, for example,

accounts for only ~0.2% of a bacterial genome), and if a pathogen is present at low

abundance compared to host and environmental DNA, these genes are likely to be

missed in routine metagenomic sequencing screens. As such, although these tools can

be specific, they lack the sensitivity required for ancient pathogen screening from

shallow metagenomic datasets. Screening techniques that accommodate queries of

whole genomes are of clear benefit for archaeological studies (25). However, while

some algorithms, such as Kraken (31), have been developed to query databases that

contain thousands of complete reference genomes using k-mer matching, this

approach does not produce the alignment information necessary to further evaluate

species identification accuracy or authenticity.

In addition to taxonomic classification (32), it is also critical to distinguish ancient

bacteria from modern contaminants (9, 10). Genuine aDNA, especially pathogen

bacterial DNA, is usually only present in small amounts and can be distinguished

from modern DNA contamination by applying an established set of authenticity

criteria (9, 10), the most important of which is the assessment of DNA damage. In

ancient DNA, cytosine deamination accumulates over time at DNA fragment termini

(33, 34), thus leading to a specific pattern of nucleotide misincorporation. The

evaluation of additional authenticity criteria such as edit distances and the distribution

of mapped reads across the reference are also recommended to mitigate against

database bias artifacts and to further validate taxonomic assignments (9, 10). While

manual evaluation of species identification and aDNA authenticity using standalone

tools might be feasible for a small sample set, it is impractical and too labour

intensive to apply to the large sample sizes typical of recent ancient DNA

investigations. The increased throughput of the ancient DNA field warrants an

automated high-throughput solution for pathogen detection in metagenomic datasets.

Successful ancient pathogen detection is reliant upon three criteria: (i) specificity of

species-level detection against a diverse metagenomic background, (ii) high

sensitivity that allows detection even with a weak signal when only trace amounts of

species-specific DNA are present, and (iii) authentication of its ancient origin.

However, no software currently exists that fulfills all requirements essential for

reliable screening of metagenomic aDNA. Here we introduce HOPS (Heuristic

Operations for Pathogen Screening), an automated computational pipeline that

screens metagenomic aDNA data for the presence of bacterial pathogens and assesses

their authenticity using established criteria. We test HOPS on experimental and

simulated data and compare it to common metagenomic profiling tools designed for

modern DNA analysis. We show that HOPS outperforms available tools, is highly

specific and sensitive, and can perform reliable and reproducible taxonomic

identification and authentication with as few as 50 reads.

RESULTS

HOPS Workflow

Figure 1. Schematic depiction of HOPS workflow. First, MALT aligns the metagenomic
data against its reference database and has an optional mode for processing aDNA reads. MaltExtract
then processes the MALT output with various filters and produces various statistics. Finally, post
processing procedures provide a comprehensive visualization of the output which can be evaluated to
identify putatively positive hits.

117
HOPS consists of three parts (Figure 1): i) a modified version of MALT (25, 35),

which includes optional PCR duplicate removal and optional deamination pattern

tolerance at the ends of reads; ii) The newly developed program MaltExtract, which

provides statistics for the evaluation of species identification as well as aDNA

authenticity criteria for a user-specified set of bacterial pathogens, with additional

functionality to filter the aligned reads by various measures such as read length,

sequence complexity or percent identity; and iii) a post-processing script that provides

a summary overview for all samples and potential bacterial pathogens that have been

identified.

MALT

MALT (Megan Alignment Tool) (25, 35) is a fast alignment and taxonomic binning

tool for metagenomic data that aligns DNA sequencing reads to a user-specified

database of reference sequences. Reads are assigned to taxonomic nodes by the naïve

Lowest Common Ancestor (LCA) algorithm (36, 37) and are thus assigned to

different taxonomic ranks based on their specificity. The default version of MALT is

intended for the analysis of metagenomic datasets deriving from modern DNA, and

thus it was not designed to accommodate the specific requirements of aDNA analyses.

In particular, aDNA damage that manifests as miscoding lesions in sequenced

products can lead to an increased number of mismatches, and extensive damage has

the potential to prevent alignment or alter taxonomic assignment. Loss of target reads

due to DNA damage can hamper species detection as aDNA studies usually begin

with shallow sequence data during initial evaluations of sample quality. In addition,

archaeological remains often show low DNA yields, and library amplification can

result in a high number of PCR duplicates, which can falsely inflate quantitative

estimates of taxa.

To account for such shortcomings, we introduce a modified version of MALT that is

specifically tailored to the analysis of aDNA data. In this modified version, PCR

duplicates are removed by eliminating reads identical to those already aligned. In

addition, reads are optionally filtered for a minimum Wootton & Federhen complexity

(38) in order to remove low complexity reads. Furthermore, to accommodate aDNA

damage during alignment, C>T substitutions are ignored in the first five positions

from the 5'-end and G>A substitutions are ignored in first five positions from the 3'-

end.

MaltExtract and post-processing

The core of HOPS is formed by the newly developed MaltExtract module. Without

MaltExtract the result files produced by MALT (RMA6 format) can only be evaluated

manually with the metagenomic analysis tool MEGAN (39). Such analysis becomes

infeasible when working with large data sets, in which each sample must be

separately searched for a long list of candidate organisms, a process that is both

laborious and prone to user error. MaltExtract provides an automated approach for the

assessment of the alignment information stored in RMA files generated by MALT. It

automatically retrieves and assesses information on various evaluation criteria for all

taxonomic nodes that match a given list of target species.

MaltExtract obtains information on edit distance, read length distribution, coverage

distribution and alignment mismatch patterns in order to identify and authenticate the

presence of species-specific ancient DNA. Furthermore, MaltExtract allows data

filtering for maximum read length, minimum percent identity, minimum complexity,

and aDNA damage patterns.

Figure 2. Post-processing steps in HOPS. Three hierarchical post-processing steps are used

in HOPS. (A) First, the edit distance distribution is required to show a decline (black lines). (B)

Second, the alignments are assessed for C>T and G>A mismatches typical for aDNA; by default, any

such damage is considered sufficient. (C) Third, the edit distance distribution of reads showing damage

is evaluated.

Accuracy in taxonomic read assignment is evaluated in a three-step procedure that

includes ancient authentication criteria (Figure 2). The first step evaluates the read

assignment to a taxonomic node. Incorrect read assignments can occur when

databases are incomplete: many species in a metagenomic sample may have no

representative reference genome in the database, and hence become erroneously

assigned to the closest genetic match, which could belong to a different species, or

even genus. Mapping to an incorrect species generally results in an increased number

of mismatches across the read that is evident in the edit distance distribution (Figure

2A). By contrast, if the sequenced reads are assigned to the correct reference species,

the edit distance distribution should continuously decline, with most of the reads

showing no or only a few mismatches, mostly resulting from aDNA damage or

evolutionary divergence of the modern reference from the ancient genome. We

summarize the shape of the edit distance distribution by a score we term *negative*

difference proportion (-Δ%), which leverages the difference in sequencing read

counts between neighboring mismatch categories (Figure S1). The -Δ% takes values

between 0 and 1, where 1 indicates a strictly declining edit distance distribution.

While true positives have a -Δ% of 1 when enough endogenous species-specific

sequencing reads are present, we use a threshold of -Δ% > 0.9 to account for possible

perturbations due to stochasticity in the edit distance distribution when few reads

(~10-20) are present. As such, this permits the detection of even low abundant taxa.

In a second step, the ancient origin of the DNA is evaluated through analysis of DNA

miscoding lesion patterns (Figure 2B). The most prominent modification observed is

deamination of cytosine into uracil, which is read as a thymine by the polymerase.

This leads to an overrepresentation of C>T substitutions at the 5' end and

correspondingly G>A substitutions at the 3' end (34, 40). Evaluation of damage

patterns is mandatory in any ancient DNA study. MaltExtract reports the rates of

substitutions for the leading and trailing 10 positions of the read alignment. The

default post-processing settings require only a single miscoding lesion to be present in

at least one read to qualify as exhibiting damage. This maximizes sensitivity and

allows authentication to function largely independently of read depth.

As a third and final criterion, we evaluate the accuracy of taxonomic assignment for

all aligned reads exhibiting aDNA damage. For this we assess again the edit distance

distribution using the -Δ% score, but now this is only performed for damaged reads

(Figure 2C). In this step, a greater number of assigned reads (>100) is required for

reliable edit distance evaluation due to the fact that not all ancient reads are expected

to exhibit damage.

The MaltExtract output is saved in a structured output folder with a summary file of

the processed input and subfolders for each evaluation criterion. The post-processing

tool generates a summary highlighting which of the target species passed one or more

evaluation criteria for each sample, as well as detailed diagnostic plots displaying the

evaluation criteria for each supported target species (Figure S2).

Assessment of taxonomic assignment on simulated data

Figure 3. Assignment of simulated reads to taxonomic levels for 33 bacterial pathogens. The fraction of simulated reads (red gradient) per reference (y axis) assigned to a specific node across different levels of the taxonomy (x axis). The levels of taxonomy not defined for a species are shown in grey.

The naïve LCA algorithm (36), which is part of HOPS, assigns reads to different taxonomic levels depending on the specificity of sequence matches. Taxonomic assignment thus depends on the structure of the underlying reference database, and it is critical to understand the expected taxonomic placement of sequenced reads from each microbial pathogen in order to successfully identify them. To analyze the taxonomic placement of a test set of 33 bacterial pathogens and to assess the performance of HOPS we simulated sequencing reads that included artificial DNA damage and spiked them into dentine, dental calculus, bone and soil metagenomic backgrounds (see Table 1).

Applying the HOPS pipeline, we recovered 98% of the simulated reads for 32 of the 33 bacterial taxa of interest (see Figure 3). The one exception was *Mycobacterium avium subspecies paratuberculosis K10* for which 23% of simulated reads were assigned to an incorrect *Mycobacterium avium subspecies paratuberculosis* strain. Our analysis shows that in most cases the taxonomic levels "species" and "complex" (e.g. *Mycobacterium tuberculosis* complex and *Yersinia pseudotuberculosis* complex) correctly accumulate the vast majority of the simulated pathogen reads. However,

noteworthy exceptions were *Brucella abortus*, *Brucella melitenis* and *Bordetella*

pertussis. Upon further investigation, we found that many species within the genera

Brucella and *Bordetella* show a high degree of sequence similarity, thus causing the

majority of the reads deriving from these pathogens to be assigned at the genus level.

By contrast, read assignment was found to be very specific for five taxa (*Treponema*

denticola ATCC 35405, *Clostridium tetani* E89, *Clostridium botulinum* E3 str. Alaska

E43*, Streptococcus gordonii* str. Challis substr. CH1 and *Clostridium botulinum*

BKT015925), resulting in the majority of reads deriving from these taxa to be

correctly assigned at the strain level. For *Salmonella enterica* subsp. *enterica* most

reads were assigned at the subspecies level. The results of this test provide a guide for

the levels of taxonomic identification that should be considered when searching for

any of the 33 queried bacterial species in experimental ancient datasets.

Optimization of MALT for aDNA

Figure 4 Comparison of the number of successfully recovered *Y. pestis* reads using standard (SD) and

damage tolerant (DT) MALT with minimum percent identities of (A) 85%, (B) 95% and (C) 99%.

Shown are the recovered reads from the "default" (all reads) and "ancient" (reads with damage) modes

in MALT, with the same 500 reads being spiked into the metagenomic backgrounds. Error bars show

the standard error of five independent technical replicates for each analysis.

Because MALT was designed for taxonomic binning of modern genetic data,

adapting it to be used on aDNA required altering the original MALT implementation

to tolerate terminal substitutions consistent with aDNA damage so that they would not

interfere with the percent identity filter. To evaluate the efficacy of this modification,

we compared the performance of the modified, damage tolerant version of MALT to

the default version using simulated *Y. pestis* data with high damage (~40%) and three

different percent identity filters: 85%, 95% and 99% (Figure 4).

As expected, the greatest difference was observed when applying the stringent 99%

identity filter, for which the damage tolerant MALT version recovered ~20% more

reads than the standard MALT version. Additionally, only the modified version was

able to recover reads with simulated damage under these parameters. At 95% identity,

only a small difference could be observed between the two MALT versions, while

results were almost identical at an 85% identity level. Taken together, the damage

tolerant MALT version provides an advantage when searching for a given pathogen

using stringent filtering criteria.

Performance comparison of HOPS, Kraken and MIDAS on simulated

data

We next evaluated the performance of HOPS by comparing it to two commonly used

metagenomics profiling tools: MIDAS (30), a marker gene-based taxonomic

classifier, and Kraken (31), which performs taxonomic classification based on k-mer

matching to a database of complete genomes. The marker gene database of MIDAS

lacked representation for *Yersinia pseudotuberculosis, Bordetella pertussis* and

Brucella melitensis. Therefore, MIDAS could only be evaluated for 30 of the 33

bacterial pathogens in the simulated data sets. For Kraken, we downloaded the

bacterial database, which lacked a reference genome to *Clostridium sporogenses*.

HOPS consistently detected all 33 pathogens in all backgrounds and among replicates

with as few as 50 reads (see Figure 5A). Kraken failed to identify *Brucella abortus*

and *Mycobacterium tuberculosis* in some replicates with only 50 simulated pathogen

reads, but otherwise had a sensitivity of 100%; however, it was prone to a high false

positive rate (see below). The sensitivity of MIDAS was far lower than for Kraken

and HOPS. Even with 5000 simulated pathogen reads for each species, MIDAS

detected only 11 of the 30 possible bacterial pathogens. This can be explained by the

limited sensitivity of marker gene based approaches, which require relatively high

sequencing coverage in order to ensure adequate representation of the marker genes

needed for identification. This is further evident as MIDAS' sensitivity is more

heavily influenced by the number of simulated reads than Kraken and HOPS.

Figure 5. Performance comparison of HOPS, Kraken and MIDAS. (A) HOPS outperforms other tools, successfully and consistently identifying all 33 target bacteria, even when represented by as few as 50 reads. (B) Number of target species identified in the metagenomic background files (negative controls) for HOPS, Kraken and MIDAS.

Negative controls

To assess false positive assignments, we queried all five metagenomic datasets for detectable signatures of the 33 test bacterial pathogens using HOPS, Kraken and MIDAS in the five metagenomic backgrounds prior to the addition of simulated pathogen reads. Kraken showed the highest susceptibility to false positives (see Figure 5B; Table S1). In this analysis, Kraken detected 24 (73%) pathogens in calculus, 19 (58%) in dentine, 13 (39%) in bone and 18 (55%) in soil. Most problematically, *Mycobacterium tuberculosis* and *Bordetella pertussis* were detected by Kraken in every metagenomic background.

Unexpectedly, MIDAS detected oral streptococci, *Tannerella forsythia, Treponema denticola* and *Porphyromonas gingivalis* in the dentine samples but not in calculus, where they are normally found. Overall, MIDAS produced fewer identifications than

Kraken, but such a result is expected given its reliance on marker gene-based

detection, which limits identification to only abundant taxa.

HOPS detected four test pathogens in the metagenomic background datasets:

Clostridium tetani (soil), *Streptococcus mutans* (calculus, dentine), *Treponema*

denticola (calculus, dentine), and *Porphyromonas gingivalis* (calculus only). Because

C. tetani is ubiquitous in soil, and all other detected bacteria are commensals of the

human oral cavity, their identification via both MIDAS and HOPS likely reflects true

positives. Taken together, HOPS and MIDAS have a lower tendency toward false

positive assignments. Kraken's increased vulnerability for aberrant assignments likely

relates to the absence of an alignment step, which is necessary for reliable species

evaluation in both modern and ancient contexts.

Positive Controls

In addition to performing tests using simulated data, we also tested HOPS, Kraken

and MIDAS on 27 ancient metagenomic datasets known to be positive for bacterial

pathogens (Table 2). They consisted of both shotgun and capture data and they varied

in sequencing depth in accordance with experimental conditions and method of data

generation.

HOPS and Kraken share 100% sensitivity for the detection of target bacterial

pathogens in every sample. By contrast, MIDAS only detected the correct bacterial

pathogen in 22 out of 27 samples. Again, MIDAS sensitivity was likely reduced due

to the marker gene-based approach. These results highlight the advantage of whole-

genome based approaches like MALT and Kraken that take advantage of every

sequenced read.

Runtimes

To calculate the runtime for each program, we used simulated metagenomic files that

each contained five million sequencing reads (see Methods). For each file, HOPS

required an average of 3307±820 seconds for the MALT step, 16±1 seconds for the

MaltExtract step and 1±0 seconds for post processing, for a total of approximately 55

minutes of analysis time per file. Kraken took on average 72±16 seconds to run

Kraken_alignment and 22±3 for *Kraken_translate*, a total of 1.5 minutes, and the

MIDAS pipeline processed each file in an average of 4±1 seconds. HOPS by far

required the highest runtimes of the three tools, but most of this time was required for

sequence alignment, a step that, although time consuming, increases detection

sensitivity, reduces false positives, and enables the authentication of aDNA reads.

Discussion

The field of archaeogenetics faces several challenges, such as the low amount of endogenous target DNA, the highly degraded nature of the DNA, and an unknown and diverse metagenomic background signal that accumulates during decomposition and centuries spent in a depositional environment. This makes reliable identification and authentication of genuine ancient DNA challenging, particularly when targeting bacterial DNA that is usually only present in small amounts. Furthermore, many bacterial pathogens have close relatives in soil, which necessitates meticulous care when making pathogen identifications.

HOPS provides an automated pipeline for high-throughput ancient bacterial species detection and authentication from metagenomic sequencing data. We compare HOPS to Kraken and MIDAS, two widely used methods for estimating both the presence and abundance of bacterial taxa in metagenomic data. These tools, however, have limited application to the specific challenges of aDNA in terms of degradation and chemical modifications manifest as miscoding lesions. Our analyses highlight the need for a pathogen identification pipeline that accommodates qualities of aDNA data and includes an essential and robust authentication for all ancient read assignments. HOPS provides a fast, reliable, and user-friendly solution to these established limitations.

HOPS was tested on simulated ancient pathogen DNA reads, and it successfully detected all targeted species spiked into metagenomic backgrounds with as few as 50 pathogen reads, representing less than 0.00001 % of the total dataset. In this context, our modified version of MALT, which tolerates mismatches resulting from DNA degradation, prevents a decrease in sensitivity even in cases of heavily damaged aDNA. We demonstrate that the marker gene-based metagenomic profiling tool MIDAS had a much lower sensitivity for pathogen detection compared to HOPS, especially for low coverage data, which is typical of ancient DNA screening datasets. Although the sensitivity of Kraken was similar to HOPS, and while Kraken's k-mer matching is considerably faster than the precise alignments used in HOPS, Kraken is incapable of validating species assignment and aDNA authenticity, and thus has a

lower specificity. This is most clearly demonstrated by our analysis of a metagenomic

soil sample in which Kraken detected numerous false positives, including

Mycobacterium tuberculosis and *Bordetella pertussis* (whooping cough). This is

likely due to many soil dwelling bacteria that harbor genetic similarities to these

pathogens, such as diverse mycobacerial species and *B. petrii,* a close relative to *B.*

pertussis that is a common constituent of environmental datasets. These effects are

further compounded by the fact that many environmental microbes have not been

genomically characterised and are not part of any reference database, which only

increases the potential of false assignments to well-sequenced pathogens. The

alignment-based validation procedure implemented in HOPS minimises such false

positive assignments, and thus offers greater accuracy in pathogen identification

during screening when environmental backgrounds comprise the dominant molecular

signal.

A previously published pipeline for the assessment of metagenomic data in

archaeogenetics is metaBIT (41). It implements a variety of methods for the detailed

assessment of metagenomic composition, which also includes validation of aDNA

damage patterns. As metaBIT is based on MetaPhlAn (42), which employs a marker

gene based approach in the initial detection step similar to MIDAS, pathogens in low

abundance could be missed in its initial steps when applied to shallow sequencing

data. An integrated approach combining HOPS and metaBIT might be a promising

future strategy for a detailed characterization of microbiomes while at the same time

proving a high level of sensitivity for the detection of pathogens. In particular, the

analysis of ancient samples that preserve their original microbiome signature, such as

dental calculus (43) or coprolites (44) would benefit from a combined application of

both methodologies, by using metaBIT to assess the microbial make up and using

HOPS for more in depth species authentication.

For all taxonomic classifiers, correct assignment of metagenomic reads is strongly

dependent on the quality of the underlying reference sequences. Currently we use a

curated database for MALT that contains completed reference sequences and

assemblies for bacteria from RefSeq (December 2016). Database sizes are constantly

increasing, but much of this growth derives from the addition of redundant sequence

data from model organisms, which also creates biases. In this context, methodologies

such as SPARSE (45) aim to mitigate against database redundancy by hierarchically

structuring reference sequences, which could be employed to further improve HOPS'

specificity and runtime.

In addition, analysis of our simulated dataset allowed for insights into the taxonomic

structure of each of the bacterial pathogens in our target list. It became apparent that

for some targets the taxonomic species level is not sufficient for identification. This

applies to historically important pathogens such as *Y. pestis* or *M. tuberculosis*. Here,

evaluation of a higher taxonomic level such as complex is more reliable, while in the

case of *Salmonella typhi* (typhoid fever) a lower level (subspecies) is favorable.

Therefore, our simulations provide a valuable resource for the optimization of

pathogen screening approaches in general.

Here, HOPS was evaluated for its success in screening for bacterial pathogens.

Because the reference database is user defined and can amended to include, for

example, the NCBI full nucleotide collection (46) or hand-curated sets of reference

genomes, tremendous flexibility exists in molecular detection, which could extend to

viruses, fungi, and eukaryotes.

Conclusions

We present a fast, reliable, and user-friendly computational pathogen screening

pipeline for ancient DNA that has the flexibility of handling large datasets. HOPS

successfully identifies both simulated and actual ancient pathogen DNA within

complex metagenomic datasets, exhibiting a higher sensitivity than MIDAS and with

fewer false positives than Kraken. HOPS provides a high level of automatization that

allows for the screening of thousands of datasets with very little hands-on time, and it

offers detailed visualizations and statistics at each evaluation step, enabling a high

level of quality control and analytical transparency. HOPS is a powerful tool for high-

throughput pathogen screening in large-scale archaeogenetic studies, producing

reliable and reproducible results even from remains with exceptionally low levels of

pathogen DNA. Such qualities make HOPS a valuable tool for pathogen detection in

the rapidly growing field of archaeogenetics.

Methods

Implementation of MaltExtract

MaltExtract is implemented in Java. It integrates parts of MEGAN's (39) source code for accessing the RMA file structure and functions from *forester* (https://github.com/cmzmasek/forester) for traversing the taxonomic tree.

Simulating data to analyse read assignment using the MALT LCA algorithm

Depending on the database structure and sequence similarity between reference sequences, the naïve LCA (36) algorithm will assign reads to different taxonomic units. To inquire how reads are assigned to the taxonomic tree for 33 bacterial pathogens (Table S2), we simulated ancient pathogen DNA reads using gargammel (47) and spiked them into five ancient metagenomic background datasets obtained from bone, dentine, dental calculus and soil (Table 1). The simulated reads carry a unique identifier in their header in order to differentiate them from metagenomic background sequences, which exhibit either full damage patterns or attenuated damage patterns following UDG-half treatment (48). To simulate aDNA damage in the pathogen sequences, we applied damage profiles obtained from previously published ancient *Yersinia pestis* genomes with (13) and without UDG-half (18) treatment. Simulated reads were processed with EAGER (49) and spiked into the metagenomic backgrounds in different amounts (50, 500 or 5000 reads). For each metagenomic background, a typical screening sequencing depth of five million reads were used.

Table 1. Metagenomic backgrounds used for simulated data sets

ID	Source	Age (Period)	Treatment	Reference
KT31calc	Calculus	Medieval	No UDG	(50)
LP39.10	Dentine	2920-2340 BCE	No UDG	(51)
MK5.001	Dentine	3348-3035 BCE 3619-3366 BCE	UDG half	(52)
TÖSM_1a	Bone	6000-5500 BCE	UDG half	(53)
Soil	Soil	-	No UDG	(25)

Evaluation of the damage tolerant version of MALT

To preserve damage patterns when mapping reads with MALT, we modified the

source code and compared the performance of the modified and default versions.

We therefore created with gargammel (47) test samples that show twice the amount of

damage (~40%) usually found in ancient samples (13). Here, we compare both

MALT versions for the bacterial pathogen *Yersinia pestis* (CO92 reference). Both

versions of MALT were tested with 85%, 95% and 99% minimum percent identity

filtering, to investigate the effects of percent identity filtering on the read alignment of

aDNA reads.

Comparison of HOPS to Kraken and MIDAS

HOPS was compared to two metagenomic taxonomic classification tools: Kraken (31)

and MIDAS (30). We only executed the first step of MIDAS that matches reads to the

marker gene database to determine species abundance. This step was executed on 32

cores with default parameters. The first step is sufficient, as any species undetected in

this step would not be detected in the remaining ones. Kraken was set to use 32 cores

to align the sample data against its reference database with the preload parameter to

load the entire database into memory before starting k-mer alignment. In a second

step kraken-translate was executed to transform taxonomy ids into proper species

names. For Kraken and MIDAS, we judged a pathogen as correctly identified if at

least one read matches to a strain of the correct species to account for the differences

in the database contents, methodologies and output formats.

Databases

In our study, HOPS uses a database containing all complete prokaryotic reference

genomes obtained from NCBI (December 1st 2016) with entries containing 'multi'

and 'uncultured' removed (13 entries). In total, 6,249 reference genomes are included

in the database. For Kraken we downloaded the bacterial database with Kraken's

kraken-build script (June 1 2017). The Kraken database contains no strain references

for *Clostridium sporogenses.* Otherwise it contains at least one reference for all of the

simulated bacterial pathogens (Table S2). For MIDAS we used the default reference

database (May 24 2016), which contained no representation of *Yersinia*

pseudotuberculosis, Bordetella pertussis and *Brucella melitensis.*

Positive controls

We compare the sensitivity and specificity of HOPS, MIDAS and Kraken using 27

metagenomic datasets previously shown to be positive for one of four microbial

pathogens: *Yersinia pestis, Mycobacterium tuberculosis*, *Salmonella enterica* and

Heliobacter pylori (Table 2). These positive control samples represent real

metagenomic data and therefore contain an unknown number of modern species in

addition to the actual recovered bacterial pathogen. Read counts across all samples

ranged from 70,897 to 7,000,000 reads. While most datasets were generated by

shotgun library screening, four datasets were enriched for pathogen DNA prior to

sequencing using DNA capture methods. For all captured datasets and a subset of

shotgun datasets, DNA was treated with UDG prior to library construction to remove

DNA damage. Both types of datasets were included to evaluate the performance of

HOPS on samples with different levels of DNA damage and pathogen abundance.

Table 2 Metagenomic samples used as positive controls

ID	Reconstructed Bacteria	Sequencing reads	Data type	Reference
10C	Salmonella enterica	1,017,400	Shotgun	(25)
35C	Salmonella enterica	986,908	Shotgun	(25)
AD388nUSG	Salmonella enterica	4,027,556	Shotgun	(25)
AD340nUSG	Salmonella enterica	3,929,877	Shotgun	(25)
RK1001.C0101	Yersinia pestis	7,023,370	Shotgun	(17)
GEN_72	Yersinia pestis	7,663,408	Shotgun	(17)
549_O	Yersinia pestis	1,520,471	Shotgun	(16)
JK3031UDG	Yersinia pestis	4,059,016	Shotgun (UDG)	(16)
JK2370UDG	Yersinia pestis	52,858,027	Shotgun (UDG)	(16)
RT6	Yersinia pestis	6,706,316	Shotgun (UDG)	(18)
1343UnTal85	Yersinia pestis	3,462,216	Shotgun	(17)
6Post	Yersinia pestis	2,546,695	Shotgun	(17)
KunilaII	Yersinia pestis	1,007,417	Shotgun	(17)
RISE00	Yersinia pestis	6,000,000	Shotgun	(13)
RISE139	Yersinia pestis	6,000,000	Shotgun	(13)
RISE386	Yersinia pestis	6,000,000	Shotgun	(13)

RISE397	Yersinia pestis	6,000,000	Shotgun	(13)
RISE505	Yersinia pestis	6,000,000	Shotgun	(13)
RISE509	Yersinia pestis	6,000,000	Shotgun	(13)
RISE511	Yersinia pestis	6,000,000	Shotgun	(13)
54	Mycobacterium tuberculosis	70,897	Shotgun	(21)
58	Mycobacterium tuberculosis	114,555	Shotgun	(21)
64	Mycobacterium tuberculosis	160,310	Shotgun	(21)
54	Mycobacterium tuberculosis	5,000,000	Capture (UDG)	(21)
58	Mycobacterium tuberculosis	5,000,000	Capture (UDG)	(21)
64	Mycobacterium tuberculosis	5,000,000	Capture (UDG)	(21)
P1P2	Helicobacter pylori	5,000,000	Capture (UDG)	(27)

Runtimes

To calculate the runtimes for HOPS, Kraken and MIDAS, we used a subset of the simulated files. The subset consisted of all metagenomic background datasets spiked with 5000 reads without technical replicates resulting in a total of 330 metagenomic files. A total of 64 cores and 700 GB of RAM were allocated to each program.

Declarations

Availability of data and material

The complete source code of HOPS is available from GitHub
(https://github.com/rhuebler/HOPS).

Competing interests

The authors declare that they have no competing interests

Funding

This research was funded by the Max Planck Society. The funding body had no
involvement in the design of the study, collection, analysis, and interpretation of data
or in writing the manuscript.

Authors' contributions

RH, FMK and AH conceived the study. RH, FMK, CW, KIB, JK and AH designed
experiments. RH and FMK implemented software. RH, FMK and AH performed
analyses. RH, FMK and AH wrote the manuscript with contributions from all
coauthors.

Acknowledgements

We thank the Department of Archaeogenetics of the Max Planck Institute for the
Science of Human History and Julian Susat for beta testing and helpful discussions.

References

References

1.   Turnbaugh PJ, Ley RE, Hamady M, Fraser-Liggett C, Knight R, Gordon JI.
The human microbiome project: exploring the microbial part of ourselves in a
changing world. Nature. 2007;449(7164):804.
2.   Karsenti E, Acinas SG, Bork P, Bowler C, De Vargas C, Raes J, et al. A
holistic approach to marine eco-systems biology. PLoS Biol. 2011;9(10):e1001177.
3.   Gilbert JA, Jansson JK, Knight R. The Earth Microbiome project: successes
and aspirations. BMC Biol. 2014;12(1):69.
4.   Haak W, Lazaridis I, Patterson N, Rohland N, Mallick S, Llamas B, et al.
Massive migration from the steppe was a source for Indo-European languages in
Europe. Nature. 2015;522(7555):207-11.
5.   Mallick S, Li H, Lipson M, Mathieson I, Gymrek M, Racimo F, et al. The
Simons Genome Diversity Project: 300 genomes from 142 diverse populations.
Nature. 2016;538(7624):201-6.
6.   Orlando L, Ginolhac A, Zhang G, Froese D, Albrechtsen A, Stiller M, et al.
Recalibrating Equus evolution using the genome sequence of an early Middle
Pleistocene horse. Nature. 2013;499(7456):74-8.
7.   Schlebusch CM, Skoglund P, Sjodin P, Gattepaille LM, Hernandez D, Jay F,
et al. Genomic variation in seven Khoe-San groups reveals adaptation and complex
African history. Science. 2012;338(6105):374-9.
8.   Skoglund P, Thompson JC, Prendergast ME, Mittnik A, Sirak K, Hajdinjak
M, et al. Reconstructing Prehistoric African Population Structure. Cell.
2017;171(1):59-71 e21.
9.   Warinner C, Herbig A, Mann A, Fellows Yates JA, Weiss CL, Burbano HA,
et al. A Robust Framework for Microbial Archaeology. Annu Rev Genomics Hum
Genet. 2017;18(0):321-56.
10.   Key FM, Posth C, Krause J, Herbig A, Bos KI. Mining Metagenomic Data
Sets for Ancient DNA: Recommended Protocols for Authentication. Trends Genet.
2017;33(8):508-20.
11.   Bos KI, Schuenemann VJ, Golding GB, Burbano HA, Waglechner N,
Coombes BK, et al. A draft genome of Yersinia pestis from victims of the Black
Death. Nature. 2011;478(7370):506-10.
12.   Wagner DM, Klunk J, Harbeck M, Devault A, Waglechner N, Sahl JW, et al.
Yersinia pestis and the plague of Justinian 541-543 AD: a genomic analysis. Lancet
Infect Dis. 2014;14(4):319-26.
13.   Rasmussen S, Allentoft ME, Nielsen K, Orlando L, Sikora M, Sjogren KG, et
al. Early divergent strains of Yersinia pestis in Eurasia 5,000 years ago. Cell.
2015;163(3):571-82.
14.   Feldman M, Harbeck M, Keller M, Spyrou MA, Rott A, Trautmann B, et al. A
High-Coverage Yersinia pestis Genome from a Sixth-Century Justinianic Plague
Victim. Mol Biol Evol. 2016;33(11):2911-23.
15.   Bos KI, Herbig A, Sahl J, Waglechner N, Fourment M, Forrest SA, et al.
Eighteenth century Yersinia pestis genomes reveal the long-term persistence of an
historical plague focus. Elife. 2016;5:e12994.
16.   Spyrou MA, Tukhbatova RI, Feldman M, Drath J, Kacki S, Beltran de Heredia
J, et al. Historical Y. pestis Genomes Reveal the European Black Death as the Source
of Ancient and Modern Plague Pandemics. Cell Host Microbe. 2016;19(6):874-81.

17.   Valtueña AA, Mittnik A, Key FM, Haak W, Allmńe R, Belinskij A, et al. The
Stone Age plague and its persistence in Eurasia. Current biology. 2017;27(23):3683-
91. e8.
18.   Spyrou MA, Tukhbatova RI, Wang CC, Valtuena AA, Lankapalli AK,
Kondrashin VV, et al. Analysis of 3800-year-old Yersinia pestis genomes suggests
Bronze Age origin for bubonic plague. Nat Commun. 2018;9(1):2234.
19.   Schuenemann VJ, Singh P, Mendum TA, Krause-Kyora B, Jager G, Bos KI, et
al. Genome-wide comparison of medieval and modern Mycobacterium leprae.
Science. 2013;341(6142):179-83.
20.   Schuenemann VJ, Avanzi C, Krause-Kyora B, Seitz A, Herbig A, Inskip S, et
al. Ancient genomes reveal a high diversity of Mycobacterium leprae in medieval
Europe. PLoS Pathog. 2018;14(5):e1006997.
21.   Bos KI, Harkins KM, Herbig A, Coscolla M, Weber N, Comas I, et al. Pre-
Columbian mycobacterial genomes reveal seals as a source of New World human
tuberculosis. Nature. 2014;514(7523):494-7.
22.   Kay GL, Sergeant MJ, Zhou Z, Chan JZ, Millard A, Quick J, et al. Eighteenth-
century genomes show that mixed infections were common at time of peak
tuberculosis in Europe. Nat Commun. 2015;6:6717.
23.   D'Anastasio R, Staniscia T, Milia ML, Manzoli L, Capasso L. Origin,
evolution and paleoepidemiology of brucellosis. Epidemiol Infect. 2011;139(1):149-
56.
24.   Kay GL, Sergeant MJ, Giuffra V, Bandiera P, Milanese M, Bramanti B, et al.
Recovery of a medieval Brucella melitensis genome using shotgun metagenomics.
MBio. 2014;5(4):e01337-14.
25.   Vagene AJ, Herbig A, Campana MG, Robles Garcia NM, Warinner C, Sabin
S, et al. Salmonella enterica genomes from victims of a major sixteenth-century
epidemic in Mexico. Nat Ecol Evol. 2018;2(3):520-8.
26.   Zhou Z, Lundstrom I, Tran-Dien A, Duchene S, Alikhan NF, Sergeant MJ, et
al. Pan-genome Analysis of Ancient and Modern Salmonella enterica Demonstrates
Genomic Stability of the Invasive Para C Lineage for Millennia. Curr Biol.
2018;28(15):2420-8 e10.
27.   Maixner F, Krause-Kyora B, Turaev D, Herbig A, Hoopmann MR, Hallows
JL, et al. The 5300-year-old Helicobacter pylori genome of the Iceman. Science.
2016;351(6269):162-5.
28.   Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello
EK, et al. QIIME allows analysis of high-throughput community sequencing data. Nat
Methods. 2010;7(5):335-6.
29.   Truong DT, Franzosa EA, Tickle TL, Scholz M, Weingart G, Pasolli E, et al.
MetaPhlAn2 for enhanced metagenomic taxonomic profiling. Nat Methods.
2015;12(10):902-3.
30.   Nayfach S, Rodriguez-Mueller B, Garud N, Pollard KS. An integrated
metagenomics pipeline for strain profiling reveals novel patterns of bacterial
transmission and biogeography. Genome Res. 2016;26(11):1612-25.
31.   Wood DE, Salzberg SL. Kraken: ultrafast metagenomic sequence
classification using exact alignments. Genome Biol. 2014;15(3):R46.
32.   Velsko IM, Frantz LAF, Herbig A, Larson G, Warinner C. Selection of
Appropriate Metagenome Taxonomic Classifiers for Ancient Microbiome Research.
mSystems. 2018;3(4).

33.   Jonsson H, Ginolhac A, Schubert M, Johnson PL, Orlando L. mapDamage2.0:
fast approximate Bayesian estimates of ancient DNA damage parameters.
Bioinformatics. 2013;29(13):1682-4.
34.   Briggs AW, Stenzel U, Johnson PL, Green RE, Kelso J, Prufer K, et al.
Patterns of damage in genomic DNA sequences from a Neandertal. Proc Natl Acad
Sci U S A. 2007;104(37):14616-21.
35.   Herbig A, Maixner F, Bos KI, Zink A, Krause J, Huson DH. MALT: Fast
alignment and analysis of metagenomic DNA sequence data applied to the Tyrolean
Iceman. BioRxiv. 2016:050559.
36.   Huson DH, Auch AF, Qi J, Schuster SC. MEGAN analysis of metagenomic
data. Genome Res. 2007;17(3):377-86.
37.   Droge J, McHardy AC. Taxonomic binning of metagenome samples generated
by next-generation sequencing technologies. Brief Bioinform. 2012;13(6):646-55.
38.   Wootton JC, Federhen S. Statistics of Local Complexity in Amino-Acid-
Sequences and Sequence Databases. Computers & Chemistry. 1993;17(2):149-63.
39.   Huson DH, Beier S, Flade I, Gorska A, El-Hadidi M, Mitra S, et al. MEGAN
Community Edition - Interactive Exploration and Analysis of Large-Scale
Microbiome Sequencing Data. Plos Comput Biol. 2016;12(6):e1004957.
40.   Kircher M. Analysis of high-throughput ancient DNA sequencing data.
Methods Mol Biol. 2012;840:197-228.
41.   Louvel G, Der Sarkissian C, Hanghoj K, Orlando L. metaBIT, an integrative
and automated metagenomic pipeline for analysing microbial profiles from high-
throughput sequencing shotgun data. Mol Ecol Resour. 2016;16(6):1415-27.
42.   Haft DH, Tovchigrechko A. High-speed microbial community profiling. Nat
Methods. 2012;9(8):793-4.
43.   Warinner C, Speller C, Collins MJ. A new era in palaeomicrobiology:
prospects for ancient dental calculus as a long-term record of the human oral
microbiome. Philos Trans R Soc Lond B Biol Sci. 2015;370(1660):20130376.
44.   Warinner C, Speller C, Collins MJ, Lewis CM, Jr. Ancient human
microbiomes. J Hum Evol. 2015;79:125-36.
45.   Zhou Z, Luhmann N, Alikhan N-F, Quince C, Achtman M, editors. Accurate
Reconstruction of Microbial Strains from Metagenomic Sequencing Using
Representative Reference Genomes. International Conference on Research in
Computational Molecular Biology; 2018: Springer.
46.   Pruitt KD, Tatusova T, Maglott DR. NCBI reference sequences (RefSeq): a
curated non-redundant sequence database of genomes, transcripts and proteins.
Nucleic Acids Res. 2007;35(Database issue):D61-5.
47.   Renaud G, Hanghøj K, Willerslev E, Orlando L. gargammel: a sequence
simulator for ancient DNA. Bioinformatics. 2016;33(4):577-9.
48.   Rohland N, Harney E, Mallick S, Nordenfelt S, Reich D. Partial uracil-DNA-
glycosylase treatment for screening of ancient DNA. Philos Trans R Soc Lond B Biol
Sci. 2015;370(1660):20130624.
49.   Peltzer A, Jager G, Herbig A, Seitz A, Kniep C, Krause J, et al. EAGER:
efficient ancient genome reconstruction. Genome Biol. 2016;17(1):60.
50.   Mann AE, Sabin S, Ziesemer K, Vagene AJ, Schroeder H, Ozga AT, et al.
Differential preservation of endogenous human and microbial DNA in dental calculus
and dentin. Sci Rep. 2018;8(1):9822.
51.   Olalde I, Brace S, Allentoft ME, Armit I, Kristiansen K, Booth T, et al. The
Beaker phenomenon and the genomic transformation of northwest Europe. Nature.
2018;555(7695):190.

52.   Wang C-C, Reinhold SR, Kalmykov A, Wissgott A, Brandt G, Jeong C, et al.
The genetic prehistory of the Greater Caucasus. bioRxiv. 2018:322347.
53.   Lipson M, Szecsenyi-Nagy A, Mallick S, Posa A, Stegmar B, Keerl V, et al.
Parallel palaeogenomic transects reveal complex genetic history of early European
farmers. Nature. 2017;551(7680):368-72.

Supplementary Information

HOPS: Automated detection and authentication of pathogen DNA in

archaeological remains

Figure S1 The first and third steps in the HOPS postprocessing protocol require a decline in the Edit

Distance distribution.

Damage plot for Mycobacterium_tuberculosis_complex

Node	Mycobacterium_tuberculosis_complex
Top Reference	Mycobacterium_africanum
all reads	203
nonDup	203
readDis	1
nonStacked	203
destacking?	On
C>T_1	0.1818
G>A_−1	0.2069
mean length (sd)	42 (10.451)

Mycobacterium_africanum;_CP010334.1 100%
Mycobacterium_africanum_GM041182;_FR878060.1 099%
Mycobacterium_microti;_CP010333.1 098%
Mycobacterium_tuberculosis;_CP010335.1 098%
Mycobacterium_tuberculosis_CCDC5180;_CP002885.1 098%
Mycobacterium_tuberculosis_CCDC5180;_CP001642.1 098%
Mycobacterium_tuberculosis_CCDC5079;_CP002884.1 098%
Mycobacterium_bovis_BCG_str._Mexico;_CP002095.1 098%
Mycobacterium_tuberculosis_BT2;_CP002882.1 098%
Mycobacterium_tuberculosis_K;_CP007803.1 098%

Figure S2 HOPS summary output for a tuberculosis positive sample. Upper left: Edit distance

distribution for all reads assigned to M. tuberculosis. Upper right: Edit distance distribution for

assigned reads that show a possible DNA damage signal. Middle left: DNA damage plot for assigned

reads. Lower left: Top ten references with percentage of aligned reads. Middle right: Summary

statistics for assigned reads.

Table S1 Results for negative controls. For HOPS the step in the post processing that was reached for

each species is indicated (0: No detection; 1: detected with good edit distance distribution; 2:

additionally indication for damage; 3: additionally good edit distance distribution for damaged reads).

For Kraken the number of k-mers assigned to the species and for MIDAS the number of assigned reads

is listed.

Species	bone_ TOSM1a	calculus_ KT31calc	dentine_M K5.001	dentine_LP39 .10	soil	Software
Bacillus anthracis	0	0	0	0	0	HOPS
Bordetella pertussis	0	0	0	0	0	HOPS
Borrelia	0	0	0	0	0	HOPS

burgdorferi B31						
Brucella abortus	0	0	0	0	0	HOPS
Brucella melitensis	0	0	0	0	0	HOPS
Clostridium botulinum	0	0	0	0	1	HOPS
Clostridium sporogenes	0	0	0	0	0	HOPS
Clostridium tetani	0	0	0	0	1	HOPS
Corynebacterium diphtheriae	0	0	0	0	0	HOPS
Haemophilus influenzae	0	0	0	0	0	HOPS
Helicobacter pylori	0	0	0	0	0	HOPS
Mycobacterium avium subsp. paratuberculosis	0	0	0	0	0	HOPS
Mycobacterium leprae	0	0	0	0	0	HOPS
Mycobacterium tuberculosis	0	0	0	0	0	HOPS
Neisseria gonorrhoeae	0	0	0	0	0	HOPS
Neisseria meningitidis	0	0	0	0	0	HOPS
Porphyromonas gingivalis	0	3	0	0	0	HOPS
Salmonella enterica subsp. enterica	0	0	0	0	0	HOPS
Staphylococcus aureus subsp. aureus	0	0	0	0	0	HOPS
Streptococcus gordonii str.	0	0	0	1	0	HOPS
Streptococcus mutans	0	3	0	2	0	HOPS
Streptococcus pneumoniae	0	2	0	0	0	HOPS
Tannerella forsythia	0	0	0	0	0	HOPS
Treponema denticola	0	0	0	1	0	HOPS
Treponema pallidum subsp. pallidum	0	0	0	0	0	HOPS
Vibrio cholerae	0	0	0	0	0	HOPS
Yersinia pestis	0	0	0	0	0	HOPS
Yersinia pseudotuberculosis	0	0	0	0	0	HOPS
	bone_TOSM1a	calculus_KT31calc	dentine_MK5.001	dentine_LP39.10	soil	Kraken
Bacillus anthracis	0	5	1	0	0	Kraken
Bordetella pertussis	1	12	20	4	10	Kraken
Borrelia burgdorferi B31	0	0	0	0	0	Kraken

Brucella abortus	0	0	0	0	1	Kraken
Brucella melitensis	0	4	0	0	1	Kraken
Clostridium botulinum	40	182	0	21	40	Kraken
Clostridium sporogenes	0	0	0	0	0	Kraken
Clostridium tetani	13	50	1	2	240	Kraken
Corynebacterium diphtheriae	4	79	9	13	11	Kraken
Haemophilus influenzae	2	1617	0	2	3	Kraken
Helicobacter pylori	8	12	0	0	4	Kraken
Mycobacterium avium subsp. paratuberculosis	13	8	39	6	26	Kraken
Mycobacterium leprae	16	19	22	13	21	Kraken
Mycobacterium tuberculosis	53	46	208	55	117	Kraken
Neisseria gonorrhoeae	4	2407	2	3	8	Kraken
Neisseria meningitidis	4	4448	0	9	12	Kraken
Porphyromonas gingivalis	2	13925	0	6	0	Kraken
Salmonella enterica subsp. enterica	8	54	16	24	16	Kraken
Staphylococcus aureus subsp. aureus	1	19	2	4	4	Kraken
Streptococcus gordonii	0	71751	0	15	0	Kraken
Streptococcus mutans	0	401	0	17	0	Kraken
Streptococcus pneumoniae	0	3224	0	0	0	Kraken
Tannerella forsythia	8	31294	14	18	14	Kraken
Treponema denticola	0	25264	0	4	0	Kraken
Treponema pallidum subsp. pallidum	0	0	0	0	0	Kraken
Vibrio cholerae	6	60	1	6	5	Kraken
Yersinia pestis	0	4	0	0	0	Kraken
Yersinia pseudotuberculosis	15	21	7	39	7	Kraken
	bone_ TOSM1a	calculus_ KT31calc	dentine_M K5.001	dentine_LP39 .10	soil	
Bacillus anthracis	0	0	0	0	0	Midas
Bordetella pertussis	0	0	0	0	0	Midas
Borrelia burgdorferi B31	0	0	0	0	0	Midas
Brucella abortus	0	0	0	0	0	Midas

Brucella melitensis	0	0	0	0	0	Midas
Clostridium botulinum	0	0	0	0	0	Midas
Clostridium sporogenes	0	0	0	0	0	Midas
Clostridium tetani	0	0	0	0	0	Midas
Corynebacterium diphtheriae	0	0	0	0	0	Midas
Haemophilus influenzae	0	0	0	0	0	Midas
Helicobacter pylori	0	0	0	0	0	Midas
Mycobacterium avium subsp. paratuberculosis	0	0	0	0	0	Midas
Mycobacterium leprae	0	0	0	0	0	Midas
Mycobacterium tuberculosis	0	0	0	0	0	Midas
Neisseria gonorrhoeae	0	0	0	0	0	Midas
Neisseria meningitidis	0	0	0	0	0	Midas
Porphyromonas gingivalis	0	0	0	29	0	Midas
Salmonella enterica subsp. enterica	0	0	0	0	0	Midas
Staphylococcus aureus subsp. aureus	0	0	0	0	0	Midas
Streptococcus gordonii str.	0	0	0	0	0	Midas
Streptococcus mutans	0	0	0	0	0	Midas
Streptococcus pneumoniae	0	0	0	1	0	Midas
Tannerella forsythia	1	0	1	26	0	Midas
Treponema denticola	0	0	0	42	0	Midas
Treponema pallidum subsp. pallidum	0	0	0	0	0	Midas
Vibrio cholerae	0	0	0	0	0	Midas
Yersinia pestis	0	0	0	0	0	Midas
Yersinia pseudotuberculosis	0	0	0	0	0	Midas

743   **Table S2** Genomes used to generate simulated ancient pathogen DNA data sets

Bacillus anthracis str Ames	Neisseria meningitidis MC58
Bordetella pertussis Tohama I	Porphyromonas gingivalis W83
Borrelia burgdorferi B31	Salmonella enterica subsp enterica serovar Enteritidis str P125109
Brucella abortus 2308	Salmonella enterica subsp enterica serovar Typhi str CT18
Brucella melitensis bv 1 str 16M	Salmonella enterica subsp enterica serovar Typhimurium str LT2
Clostridium botulinum A str ATCC 3502	Staphylococcus aureus subsp aureus NCTC 8325
Clostridium botulinum BKT015925	Streptococcus gordonii str Challis substr CH1
Clostridium botulinum E3 str Alaska E43	Streptococcus mutans UA159
Clostridium sporogenes NCIMB 10696	Streptococcus pneumoniae R6
Clostridium tetani E88	Tannerella forsythia 92A2
Corynebacterium diphtheriae NCTC 13129	Treponema denticola ATCC 35405
Haemophilus influenzae Rd KW20	Treponema pallidum subsp pallidum str nichols
Helicobacter pylori 26695	Vibrio cholerae M66-2
Mycobacterium avium subsp paratuberculosis K10	Vibrio cholerae O1 biovar El Tor str N16961
Mycobacterium leprae TN	Yersinia pestis CO92
Mycobacterium tuberculosis anc	Yersinia pseudotuberculosis IP31758
Neisseria gonorrhoeae FA 1090	

744