1

Title:

Development of new food-sharing relationships among nonkin vampire bats

Authors and affiliation:

Gerald G. Carter[1,2], Damien R. Farine[3,4,5,6], Rachel J. Crisp[2,7], Julia K. Vrtilek[2,8], Simon P. Ripperger[1,2,9], Rachel A. Page[2]

[1]Department of Ecology, Evolution, and Organismal Biology; The Ohio State University, Columbus, OH, USA

[2]Smithsonian Tropical Research Institute, Balboa, Ancón, Panamá

[3]Edward Grey Institute of Field Ornithology, Department of Zoology, University of Oxford, Oxford, UK

[4]Department of Collective Behaviour, Max Planck Institute of Animal Behavior, Konstanz, Germany

[5]Chair of Biodiversity and Collective Behaviour, Department of Biology, University of Konstanz, Konstanz, Germany

[6]Center for the Advanced Study of Collective Behaviour, University of Konstanz, Konstanz, Germany

[7]Centre for Research in Animal Behaviour, University of Exeter, Exeter, UK

[8]Department of Biology, Eidgenössische Technische Hochschule Zürich, Zürich, Switzerland

[9]Museum für Naturkunde, Leibniz-Institute for Evolution and Biodiversity Science, Berlin, Germany

Corresponding author:

Gerald Carter, 318 W. 12th Avenue, 300 Aronoff Laboratory, Columbus, OH 43210,

carter.1640@osu.edu

Abstract

Social relationships that involve costly helping occur most often among kin, but in many complex and individualized animal societies, nonkin also demonstrate stable cooperative relationships that share similarities with human friendship. How do such cooperative bonds form between complete strangers? One theory suggests that strangers should 'test the waters' of a new relationship by initially making low-cost cooperative investments and gradually escalating them with good partners. This 'raising-the-stakes' strategy is evident in humans playing short-term economic games, but it remains unclear whether it applies to the natural helping behaviors that underlie a long-term cooperative relationship. Here, we show evidence that unfamiliar nonkin vampire bats (*Desmodus rotundus*) selectively escalate low-cost investments in allogrooming before developing higher-cost food-sharing relationships. By introducing females from geographically distant sites in pairs or groups and fasting them repeatedly over 15 months, we observed that bats first established new reciprocal grooming relationships, and that increasing grooming rates predicted the occurrence of first food donations, at which point grooming rates no longer increased. New food-sharing relationships emerged reciprocally in 14.5% of 608 female pairs and formed faster when strangers lacked alternative familiar partners. Our results suggest that 'raising-the-stakes' might be easier to detect when tracking multiple types of behavior during relationship formation, rather than measuring a single behavior within an established relationship. The general principle of 'testing the waters' might play an underappreciated role across many other social decisions with long-term consequences, such as joining a new social group or choosing a long-term mate.

Keywords: cooperation, social relationships, vampire bats

Significance statement

Vampire bats form long-term cooperative social bonds that involve reciprocal regurgitated blood sharing. But how do two individuals go from complete strangers to reciprocal food donors? By introducing unfamiliar bats, we found evidence that low-cost grooming paves the way for higher-cost food donations. Food sharing emerged in a reciprocal fashion and it emerged faster when two strangers could not access their original groupmates. Bats that did form new food-sharing relationships had a history of escalating reciprocal grooming up until the food sharing began. The finding that unfamiliar nonkin vampire bats appeared to gradually and selectively transition from low-cost to high-cost cooperative behaviors is the first evidence that nonhuman animals 'raise the stakes' when forming new cooperative relationships.

\body

Animal societies are fundamentally shaped by repeated interactions among individuals over time. Repeated interactions allow individuals to choose to cooperate based on their past experience across different partners (1-3). Organisms as diverse as animals, plants, and fungi have demonstrated partner choice: individuals prevent exploitation by shifting their cooperative investments towards partners that provide better reciprocal returns (3-7). Across several nonhuman species, repeated cooperative interactions can lead to adaptive and enduring social bonds that share similarities with human friendship (8-12), though it remains unclear how these initially form. A significant challenge has been testing how individuals prevent exploitation while forming these stable bonds. How do complete strangers develop a long-term cooperative relationship?

A key idea is that individuals should reduce the risk of exploitation by initially spreading

out smaller cooperative investments across time ('parceling' (13)) or across different partners

('social bet-hedging' (14)), and then gradually escalating investments in the most cooperative

partnerships ('raising the stakes' (15)). For example, one might first assess a potential partner's

tolerance by clustering for warmth, then gain feedback by grooming the partner, and then use the

partner's response to decide whether to provide higher-cost food donations or coalitionary

support (16). Despite its intuitive appeal for explaining how new cooperative relationships

develop, evidence supporting the 20-year-old 'raising-the-stakes' model (15) is surprisingly

scarce. An early test using the cleaner and client fish mutualism suggested that the model does

not apply well to situations with severe asymmetries in partner payoffs or options (17). Studies

with nonhuman primates (18-21) have tested only snapshots of established relationships rather

than the formation of new ones. Human strangers 'raise the stakes' when making monetary bids

in cooperation games (e.g. 22, 23), but the strategy has never been properly tested in a more

ecologically-relevant context of relationship formation. Doing so requires measuring the

emergence of natural helping behaviors between randomly introduced strangers.

We tracked the development of cooperative relationships between previously unfamiliar

wild-caught vampire bats (*Desmodus rotundus*) over 15 months. Cooperative relationships in this

species involve low-cost social grooming (hereafter *grooming*) and higher-cost regurgitations of

ingested blood (*food sharing*), which are provided to close kin and frequent associates (14, 24-

28). Here, we found evidence that vampire bats use reciprocal grooming to gradually establish

new bonds that entail food donations. Our data supported four key predictions: (i) new food-

sharing relationships should form gradually but faster when bats have fewer alternative partners,

(ii) grooming rates should predict the probability that the grooming recipient later donates food

to the groomer, (iii) low initial grooming rates should increase over time but only up until the

first reciprocal food donation, and (iv) the emergence of new food sharing should be reciprocal.

Female vampire bats demonstrate kin-biased fission-fusion social dynamics (24, 27, 28).

New nonkin social bonds can form when an unrelated female joins a social network about once

every two years (24, 28), with individual bats living for up to 16 years in the wild (29). To

observe how new food-sharing relationships form between adults, we captured adult females

from two distant sites in Panamá, Tolé (n=19) and Las Pavas (n=8), and we then ran 638 fasting

trials in which an overnight-fasted subject could be fed by a previously unfamiliar bat from

another site. To test the prediction that new sharing relationships would form faster when

strangers have fewer options of alternative partners, we compared the occurrence of new sharing

when wild-caught strangers were introduced in isolated pairs (one Las Pavas and one Tolé bat),

in small groups (one Las Pavas and three Tolé bats), or in one large mixed group (all bats

together; see Methods, Supplementary Information (SI) Appendix, Fig. S1). New bonds can also

form when individuals are born into a group, and these relationships might form differently. We

therefore also measured the development of non-maternal cooperative relationships between 26

female adults and 13 younger captive-born bats (6 males and 7 females, 11 to 21 months old) in

the large mixed group.

To test our hypotheses, we compared the observed coefficients from general and

generalized linear models (slopes β, and odds ratios OR, respectively) to expected distributions

of coefficient values expected under the null hypotheses using permutations of the network or the

event data (see Methods). We use the term 'potential relationship' for a pair of bats that could

have groomed or shared food, 'relationship' for an *observed* network edge (directed), and 'bond'

to discuss the underlying construct that we *inferred* from the observed relationship (see glossary,

SI Appendix, Table S1).

Over 424 days and 12,012 opportunities for new food donations, new food sharing

developed in 10.8% of the 996 potential relationships among all bats, 14.5% of 608 potential

relationships among females, and 15.6% of 243 potential relationships among wild-caught adult

females (SI Appendix). All bats had at least one donor (range=1-16, mean=6.6). The average

number of new food donors per adult female bat was 2.7 (range=0-7) and the average per

captive-born bat was 2.6 (range=0-6). New grooming relationships developed much more

frequently (all bats=51.9% of 1008; females=58.9% of 618; wild-caught adult females=78.2% of

248; SI Appendix). The average number of new groomers was 7.2 (range=0-16) for adult

females and 14.4 (range=1-23) for captive-born bats. New food sharing emerged after grooming

observations more than expected based on the relative frequency of grooming and sharing (see

supplement).

If social experience and partner availability influence the development of new food-

sharing bonds, then first food donations should occur gradually but faster when bats have fewer

alternative partners. As expected, when strangers from Las Pavas and Tolé were introduced and

housed as isolated pairs, we observed higher rates of new food sharing (β=1.14, p=0.004; effect

on new grooming: β=1.09, p=0.04), compared to when one Las Pavas bat was introduced to three

Tolé bats, despite there being much fewer potential new bonds available to form. When we

aggregated bats from the controlled introduction trials into a large mixed group, bats

preferentially fed and groomed their original familiar groupmates, and new sharing emerged

even more gradually than in the isolated pairs or in small groups (SI Appendix, Fig. S2-S3).

According to the 'raising-the-stakes' hypothesis, strangers should make low-cost investments and use the return rates to inform future investments. We therefore predicted that if grooming helps to build sharing bonds, then the grooming rate should predict the probability of the first food donation in the opposite direction, initial grooming rates should start low and then increase over time but only up until the first reciprocal food donation, and the emergence of new food-sharing should be reciprocal. As expected, the grooming rate given by actor A to recipient B predicted the later occurrence of new food sharing from B back to A (OR=2.15, p<0.0002, n=897). The trajectory of grooming rates over time clearly differed between pairs that developed new food-sharing relationships versus pairs that did not (interaction: OR=1.60, p<0.0002, Fig. 1). The slope of this increase in grooming was also greater *before* the new food-sharing relationship formed than it was after the relationship formed (Fig. 2). New food sharing was more reciprocal than expected if the new donations were random, even when controlling for kinship (estimated via maternal pedigree and microsatellite genotyping, see Methods; MRQAP-DSP; reciprocal sharing: β=0.40, p<0.0002, kinship: β=0.07, p=0.10; SI Appendix). Grooming rates were highest between bats that formed two-way food-sharing relationships, intermediate in relationships where we observed sharing in only one direction, and lowest in pairs where we never saw food sharing (SI Appendix, Fig. S4).

The rarity of new food-sharing relationships corroborates past evidence that food regurgitations are energetically costly and that food-sharing bonds require investments of time and energy (14, 24, 30, 31). Our finding that the emergence of new food sharing between wild-caught adults is influenced by partner behavior and availability provides the clearest evidence to date that nonkin food sharing in vampire bats is not a mere byproduct of kin selection (26). Before this study, one hypothesis was that food sharing decisions could be based entirely on

phenotypic similarity, which could result in a spurious pattern mimicking reciprocal sharing

among nonkin (32, 33). For instance, nonkin pairs with scents or calls that are, by chance, similar

would feed each other in both directions (33). However, this hypothesis incorrectly predicts that

food-sharing relationships should form immediately, and not vary dramatically with partner

availability and past social experience. Under the phenotypic similarity hypothesis, we would

expect to have observed more food-sharing relationships in larger groups simply because there is

a substantially greater opportunity to have pairs of individuals with matching phenotypes.

Previous evidence for 'raising-the-stakes' in nonhuman social relationships came from

observations of grooming among familiar male chimpanzees after the death of an alpha male

(18). The authors suggested that, during this period of social instability, these groupmates may

have needed to re-establish their relationships, and that a diminishing threat of violence led to the

increasing rates of grooming (18). Although the increase in grooming rates is consistent with

each male 'raising the stakes' to assess the risk of aggression from their grooming partner, it

might have also resulted from a general decline in vigilance against possible aggression from any

other groupmate.

Past studies on 'raising the stakes' during relationship development have also focused on

single types of cooperative investments (18-23). However, individuals can raise the stakes either

by increasing the magnitude of one kind of cooperative investment or by adding new higher-cost

forms of cooperative investment. In our study, we show that female vampire bats appear to do

both, first by increasing grooming rates and then by transitioning from low-cost grooming to

high-cost food-sharing.

Past evidence for the 'raise-the-stakes' model (15) has been scarce in part because it is a

variation on the classic 'tit-for-tat' strategy in the iterated prisoner's dilemma (1), a model which

is difficult to test unambiguously using natural forms of cooperation (4, 32, 34). 'Tit-for-tat'

forms of reciprocity have been demonstrated using experiments with trained instrumental tasks

(e.g. pulling a lever to deliver food) and payoffs accruing in distinct rounds (4-6). However, the

'tit-for-tat' model of reciprocity should not be operationalized too literally, because it excludes

many factors crucial to real-world cooperation, such as partner choice, social bonding, different

service types, and the cost-benefit asymmetries resulting from demography, market effects, or

social dominance (4, 7, 34). Also, long-term cooperative social bonds do not produce strict 'tit-

for-tat' patterns of helping. Instead, primates with stronger bonds show *less* evidence for short-

term contingencies in grooming (9), which makes sense if social bonding involves integrating

many different kinds of social interactions into a single positive association.

The relevance of our findings extends beyond high-cost cooperative behaviors. For

example, in some species, courtship behaviors could be seen as a short-term investment in the

formation of longer-term pair bonds with substantial fitness consequences (35). Similarly, the

role of mere physical contact as a low-cost method for building tolerance and trust might be

more general than currently recognized. The key role of grooming for relationship maintenance

in primates is well established, but growing evidence suggests that similar tactile behaviors can

reduce fear and encourage tolerance and cooperation in many other species of mammals, birds,

and fish (e.g. 5, 36-42). Recently developed methods for tracking formation of social bonds at

fine temporal scales (43, 44) could provide new opportunities to test whether gradual escalation

of proximity and body contact is a widespread mechanism for 'testing the water' of a potential

social relationship.

Methods

Animals

We used 41 common vampire bats (*Desmodus rotundus*) as subjects, including 19 female

bats captured exiting a roost in Tolé, Panamá; 8 female bats captured foraging at a cattle pasture

in Las Pavas, Panamá about ~215 km from Tolé; and 14 captive-born bats (8 females, 6 males).

We studied adult females and their young, because these individuals form the basis of food-

sharing networks in the wild, whereas adult males compete for access to territories and females

and do not form stable bonds as often (24-28). To ensure familiarity within groups and

unfamiliarity between groups, we housed the groups separately (Tolé bats for 6 months and Las

Pavas bats for 2 weeks) before the study began. Bats were marked with subcutaneous passive

integrated transponders (Trovan Ltd. USA) and a visually unique combination of forearm bands

(Porzana, National Tag, and birdbands.com). To feed bats, we provided refrigerated or thawed

cattle or pig blood defibrinated with sodium citrate and citric acid.

We used a 3-4 mm biopsy punch to collect tissue samples in 80% or 95% ethanol, then used

a salt–chloroform procedure for DNA isolation, and a LI–COR Biosciences® DNA Analyser

4300 and the SAGA GT allele scoring software to genotype individuals at 17 polymorphic

microsatellite loci. Allele frequencies were based on 100 bats from Tolé and 9 bats from Las

Pavas, respectively. Genotypes were 99.9% complete. To estimate genetic relatedness, we used

the Wang estimator in the R package 'related'. To estimate kinship, we assigned a zero kinship

to known unrelated individuals from different sites and to individuals with negative pairwise

relatedness, and we assigned a kinship of 0.5 for known mother-offspring pairs or pairs with

genetic relatedness estimates greater than 0.5. For all other pairs, we used genetic relatedness as

the estimate for kinship.

Experimental design

We induced allogrooming and regurgitated food sharing using a fasting trial, in which a

focal subject was isolated from the group without food for a night and a day, then released back

to the group of fed bats for 1 hour the following night. During the hour, all grooming or food-

sharing interactions with the subject were recorded using an infrared (IR) light and an IR-

sensitive video camera. Each food sharing bout was estimated by the number of seconds that the

unfed subject spent licking the mouth of a particular groupmate. Grooming was defined as

chewing or licking the fur or wings of another bat. The dyadic sharing or grooming for a trial

was estimated as the sum of all bouts that were at least 5 seconds long. Bats were weighed before

and after trials, and total observed mouth-licking durations predicted weight gain (SI Appendix).

We conducted fasting trials in each group during three experimental phases (SI Appendix,

Fig. S1). First, we conducted 57 'baseline' trials to assess preliminary sharing rates between the

19 Tolé bats housed in a 1.7 x 2.1 x 2.3 m outdoor flight cage in Gamboa, Panama (1 group,

3420 possible sharing interactions). Second, we conducted 106 'controlled introduction' trials to

assess possible formation of new food-sharing bonds between bats introduced as either an

isolated pair (one Las Pavas bat and one Tolé bat) or a quartet (one Las Pavas bat and three Tolé

bats), housed in a 28 x 28 x 40 cm clear plastic observation cage (10 pairs and 8 quartets). These

controlled introductions provided for 162 opportunities for new food sharing between previous

strangers (SI Appendix, Table S2). Finally, we conducted 532 'mixed-group' trials to assess the

formation of new sharing relationships when all bats were housed together in the flight cage

described above (19 Tolé, 7 Las Pavas, and 14 captive-born bats). The introductions in this

combined group provided 11,823 more opportunities for new sharing.

Statistical analyses

During the baseline and mixed-group trials, we estimated food donation size as the number

of seconds that a fasted subject spent mouth-licking a fed groupmate. During the controlled

introduction trials, however, when bats were forced in close proximity, we saw a greater

frequency of begging, defined as mouth-licking that is clearly not food-sharing because the

partner is turning away from the mouth-licking bat and the mouth-licking bat does not gain the

weight that would be expected from food-sharing. To be conservative when measuring sharing,

we therefore did not count mouth-licking as food sharing during the controlled introduction trials

unless the subject weighed more than expected based on the average weight change for bats that

did not perform any mouth-licking.

Durations of sharing and grooming were lognormal. To create a standard index of grooming

rates, we therefore transformed the total duration of directed dyadic interactions in each trial

using natural log (x+1). We call these measures of the log duration per hour 'rates'. When

interaction bout duration and probability had different meanings, we decomposed rates into two

separate response variables: amounts (the magnitude of nonzero rates in a trial) and probabilities

(the presence or absence of a nonzero rate in a trial). We used permutation tests with 5,000

permutations for all hypothesis testing (p-values) and bootstrapping for all 95% confidence

intervals. To approximate two-sided p-values, we doubled any one-sided permutation p-values

that were below 0.05. Null distributions were not always centered on zero due to structure in the

data, so caution must be taken when considering the observed coefficients.

Grooming could occur before sharing simply because it is more frequent. To test whether

grooming preceded new sharing more than expected by chance, we therefore compared the

observed probability and latency of grooming before sharing to the values expected based on

randomizing the days on which grooming and sharing interactions occurred. To test for ingroup-

outgroup biases in sharing for each site, we calculated observed coefficients for the effect of

being from the same capture site on actor grooming rates, then we calculated expected

coefficients by permuting the grooming rates within each actor. To test the effects of kinship and

reciprocal grooming on the formation of new food-sharing relationships in the mixed-group

trials, we used multiple regression quadratic assignment procedure with double semi-partialing

(MRQAP-DSP) via the netlogit function in the sna R package. We also used this method to test

the effect of grooming on occurrence of new sharing only within the controlled introduction

trials. This procedure uses generalized linear models via the glm function in lme4 package to

calculate the observed coefficients and uses network-level permutations to get expected

coefficients. Since MRQAP-DSP cannot test interaction effects, we compared observed and

expected interaction coefficients using permutations in which we shuffled trial rates given by the

actor among different possible receivers and then shuffled the trial rates received by the receiver

among different possible actors. If the interaction coefficients were significant ($p<0.05$), we

conducted separate MRQAP-DSP tests within each group.

To test whether interaction rates changed over time, we generated expected coefficients for

general or generalized linear models by permuting the order of interactions within each potential

relationship. One captive-born bat died for unknown reasons during the mixed-group trials, so

we removed it from all temporal analyses. To test for evidence of reciprocal sharing, we used

MRQAP-DSP to test if the matrix of new sharing in the mixed-group trials was predicted by

reciprocal sharing when controlling for kinship. As an additional test, we also counted the

occurrence of both novel sharing and reciprocal sharing for all new potential relationships, then

counted the same number after randomizing the presence of sharing across potential

relationships.

Data availability

Behavioral data, genotypes, and R code are available as supplementary information.

Acknowledgments

We thank the Smithsonian Tropical Research Institute for logistical support. Isabelle Waurick

conducted the molecular lab work. Frieder Mayer enabled the molecular lab work. Work was

supported by a Smithsonian Postdoctoral Fellowship, a Humboldt Research Fellowship, a

Smithsonian Institution Scholarly Studies Grant and a grant from the National Geographic

Society Committee for Research and Exploration (WW-057R- 17).

Author contributions

Conceptualization, GC; Methodology, GC; Investigation, GC, RC, JV; Genotyping, SR;

Statistical analysis, GC; Original draft, GC; Review & Editing, GC, RC, JV, SR, DF, RP;

Funding acquisition, GC, SR, RP; Resources, RP; Supervision, GC, DF, RP;

References

1   Axelrod R & Hamilton WD (1981) The evolution of cooperation. Science 211:1390-1396.

2   Noë R & Hammerstein P (1994) Biological markets: supply and demand determine the effect of

partner choice in cooperation, mutualism and mating. Behav. Ecol. Sociobiol. 35:1-11.

3 Trivers RL (1971) The evolution of reciprocal altruism. Q. Rev. Biol. 46:35-57.

4 Carter GG (2014) The reciprocity controversy. Anim. Behav. Cogn. 1(3):368-386.

5 Schweinfurth MK & Taborsky M (2018) Reciprocal trading of different commodities in Norway rats.

Curr. Biol. 28:1-6.

6 Rutte C & Taborsky M (2008) The influence of social experience on cooperative behaviour of rats

(*Rattus norvegicus*): direct vs generalised reciprocity. Behav. Ecol. Sociobiol. 62(4):499-505.

7 Fruteau C, Voelkl B, van Damme E, & Noë R (2009) Supply and demand determine the market value

of food providers in wild vervet monkeys. Proc. Natl. Acad. Sci. USA 106:12007-12012.

8 Samuni L, et al. (2018) Social bonds facilitate cooperative resource sharing in wild chimpanzees.

Proc. R. Soc. B 285(1888).

9 Seyfarth RM & Cheney DL (2012) The evolutionary origins of friendship. Ann. Rev. Psych. 63:153-

177.

10 Silk JB, et al. (2010) Strong and consistent social bonds enhance the longevity of female baboons.

Curr. Biol. 20(15):1359-1361.

11 Silk JB, et al. (2009) The benefits of social capital: close social bonds among female baboons

enhance offspring survival. Proc. R. Soc. B 276(1670):3099-3104.

12 Silk JB, Alberts SC, & Altmann J (2003) Social bonds of female baboons enhance infant survival.

Science 302(5648):1231-1234.

13 Connor RC (2010) Cooperation beyond the dyad: on simple models and a complex society. Philos.

Trans. R. Soc. Lond., Ser. B: Biol. Sci. 365(1553):2687-2697.

14 Carter GG, Farine DR, & Wilkinson GS (2017) Social bet-hedging in vampire bats. Biol. Lett.

13(5):20170112.

15 Roberts G & Sherratt TN (1998) Development of cooperative relationships through increasing

investment. Nature 394(6689):175-179.

16 Borgeaud C & Bshary R (2015) Wild vervet monkeys trade tolerance and specific coalitionary

support for grooming in experimentally induced conflicts. Curr. Biol. 25(22):3011-3016.

17 Bshary R (2002) Building up relationships in asymmetric co-operation games between the cleaner

wrasse Labroides dimidiatus and client reef fish. Behav. Ecol. Sociobiol. 52(5):365-371.

18 Kaburu SS & Newton-Fisher NE (2013) Social instability raises the stakes during social grooming

among wild male chimpanzees. Anim. Behav. 86(3):519-527.

19 Barrett L, Henzi SP, Weingrill T, Lycett JE, & Hill RA (2000) Female baboons do not raise the stakes

but they give as good as they get. Anim. Behav. 59(4):763-770.

20 Manson JH, Navarrete CD, Silk JB, & Perry S (2004) Time-matched grooming in female primates?

New analyses from two species. Anim. Behav. 67(3):493-500.

21 Fruteau C, Lemoine S, Hellard E, van Damme E, & Noë R (2011) When females trade grooming for

grooming: Testing partner control and partner choice models of cooperation in two species of

primates. Anim. Behav. 81:1223-1230.

22 Roberts G & Renwick JS (2003) The development of cooperative relationships: an experiment. Proc.

R. Soc. B 270(1530):2279-2283.

23 Van den Bergh B & Dewitte S (2006) The robustness of the 'Raise-The-Stakes' strategy: Coping with

exploitation in noisy Prisoner's Dilemma Games. Evol. Hum. Behav. 27(1):19-28.

24 Wilkinson GS (1984) Reciprocal food sharing in the vampire bat. Nature 308:181-184.

25 Wilkinson GS (1986) Social grooming in the common vampire bat, *Desmodus rotundus*. Anim.

Behav. 34:1880-1889.

26 Carter GG & Wilkinson G (2013) Does food sharing in vampire bats demonstrate reciprocity?

Commun. Integr. Biol. 6(6):e25783.

27 Wilkinson GS (1985) The social organization of the common vampire bat. I. Pattern and cause of

association. Behav. Ecol. Sociobiol. 17:111-121.

28 Wilkinson GS (1985) The social organization of the common vampire bat: II. Mating system, genetic

structure, and relatedness. Behav. Ecol. Sociobiol. 17:123-134.

29 Delpietro H, Russo R, Carter G, Lord R, & Delpietro G (2017) Reproductive seasonality, sex ratio

and philopatry in Argentina's common vampire bats. R. Soc. Open Sci. 4(4):160959.

30 Carter GG & Wilkinson GS (2015) Social benefits of non-kin food sharing by female vampire bats.

Proc. R. Soc. B 282(1819):20152524-20152524.

31 Carter GG, Wilkinson GS, & Page RA (2017) Food-sharing vampire bats are more nepotistic under

conditions of perceived risk. Behav. Ecol. 28(2):565-569.

32 Clutton-Brock T (2009) Cooperation between non-kin in animal societies. Nature 462:51-57.

33 Hammerstein P (2003) Why is reciprocity so rare in social animals? A protestant appeal. Genetic and

cultural evolution of cooperation (ed. & Hammerstein P), pp. 83–93. Cambridge, MA: MIT Press.

34 Noë R (2006) Cooperation experiments: coordination through communication versus acting apart

together. Anim. Behav. 71:1-18.

35 Ihle M, Kempenaers B, & Forstmeier W (2015) Fitness benefits of mate choice for compatibility in a

socially monogamous species. PLoS Biol. 13(9):e1002248.

36 Kern JM & Radford AN (2018) Experimental evidence for delayed contingent cooperation among

wild dwarf mongooses. Proc Natl Acad Sci U S A 115(24):6255-6260.

37 Kenny E, Birkhead TR, & Green JP (2017) Allopreening in birds is associated with parental

cooperation over offspring care and stable pair bonds across years. Behav. Ecol. 28(4):1142-1148.

38 Grutter AS (2004) Cleaner fish use tactile dancing behavior as a preconflict management strategy.

Curr. Biol. 14(12):1080-1083.

39 Soares MC, Oliveira RF, Ros AF, Grutter AS, & Bshary R (2011) Tactile stimulation lowers stress in

fish. Nat. Comm. 2:534.

40 Schirmer A, Jesuthasan SJ, & Mathuru A (2013) Tactile stimulation reduces fear in fish. Front.

Behav. Neuro. 7:167.

41 Radford AN (2012) Post-allogrooming reductions in self-directed behaviour are affected by role and

status in the green woodhoopoe. Biol. Lett. 8(1):24-27.

42 Radford AN (2011) Preparing for battle? Potential intergroup conflict promotes current intragroup

affiliation. Biol. Lett. 7(1):26-29.

43 Alarcón-Nieto G, et al. (2018) An automated barcode tracking system for behavioural studies in birds.

Meth. Ecol. Evol. 9(6):1536-1547.

44 Ripperger S, Günther L, Wieser H, Duda N, Hierold M, Cassens B, Kapitza R, Koelpin A, & Mayer F

(2019) Proximity sensors on common noctule bats reveal evidence that mothers guide juveniles to

roosts but not food. Biol. Lett. 15(2):20180884.

Figure Legends

Fig. 1. Increasing A-to-B grooming led to new B-to-A food-sharing relationships. In cases

where a new food-sharing relationship formed (solid line), the grooming rate towards the future

donor increased over time before the first donation occurred (OR=1.40, n=33, p<0.0002), but the

grooming rate towards a potential donor remained low in cases where no food-sharing

relationship formed (dashed line; OR=1.00, n=420, p=0.37). This divergence existed for all

potential new relationships (panel a), for previously unfamiliar adults (panel b), and for

relationships with captive-born bats (panel c; no interaction effect: OR=0.152, p=0.34).

Relationships with captive-born bats had more divergent grooming trajectories (SI Appendix).

Shading shows the 95% CI for the fitted model's predictions.

412
Fig. 2. Grooming rates increased before, but not after, new food-sharing occurred. The

probability of a focal bat grooming the new donor in a 1-h trial (y-axis) increased before the first

day that the donor fed the focal bat (i.e. 'day zero'; OR=1.4, p=0.0004), but not after this day

zero (OR=0.946, p=0.61; interaction: OR=1.69, p<0.0002). This effect was seen in new food-

sharing relationships with or without captive-born bats (SI Appendix). Shading shows the 95%

CI for the fitted model's predictions.

Supplementary Information Appendix
For the paper, Carter et al. *Development of new food-sharing relationships among nonkin vampire bats.*

Evidence of food sharing
      Fasted subjects gained an average of 51 mg of mass per minute of mouth-licking (R^2 = 0.75, p < 0.001, 95% CI: 45 to 57 mg/min, n = 619 trials without missing data), which is comparable to previous estimates from another captive colony (38 mg of mass per minute of mouth-licking, R^2 = 0.67, 95% CI: 33 to 46 mg/min, n = 121 trials; colony described in [1-3]).

Rates of food sharing
      Across all trials, the probability that a given bat received food from any groupmate was 61% (95% CI = 57 to 64%, 41 bats, 693 trials), which is much lower than the roughly 95% success rate observed in the previous long-term captive colony (95% CI = 92 to 98%, 29 bats, 183 trials; described in [1-3]). Assuming that mouth-licking events over 5 seconds were food donations, 64% of the 340 mixed-group trials with food sharing involved one donor, 24% had two donors, 9% had three donors, 2% had four donors, and only two trials had five donors.

Development of new food-sharing relationships
      We induced 12 of the 38 new food-sharing relationships between wild-caught adult females during the 106 'controlled introduction' fasting trials (Table S2), in which a single stranger Las Pavas bat was introduced to either one unfamiliar Tolé bat (forming an isolated pair) or to three Tolé bats (forming a quartet). The 26 other new food-sharing relationships between wild-caught adult females developed gradually during the 532 'mixed-group' fasting trials in the subsequent period when all the bats from both groups could freely interact (see Figures S1, S2, S3).
      During the mixed-group period, there was a 10% chance that a new food-sharing relationship would develop between an adult and a captive-born bat (7 females, 6 males, 3-19 months old; 68 of 748 potential relationships) and 3.9% chance of a new food-sharing relationship between two captive-born bats. Captive-born bats groomed each other less than they groomed adult females, even when excluding the mother (β = -0.10, n = 460, p = 0.012).
      During the mixed-group trials, the Las Pavas bats were biased towards feeding and grooming other Las Pavas bats (sharing: β = 0.28, n = 158, p < 0.0002, grooming: β = 0.53, n = 161, p < 0.0002). The same 'within-group bias' was seen in the Tolé bats (sharing: β = 0.11, n = 342, p = 0.0008; grooming: β = 0.14, n = 460, p = 0.028). New sharing relationships with captive-born bats were also more likely among bats from the same source population (OR = 1.73, p = 0.035). After controlling for this within-group bias, we found no evidence for a kinship bias (MRQAP-DSP, β = 0.21, p = 0.13). We lacked the statistical power to test for increases in food donation sizes over time within new dyads, but when pooling donations across all dyads, new donations that occurred later in time were not significantly larger (β = 5.6, n = 37, p = 0.95).

Relationships appeared to develop faster during controlled introductions
      During the mixed-group trials, the appearance of first donations became more probable over time (OR = 1.56, n = 3072, p = 0.0188), so new sharing relationships appeared to form gradually (Figures S2, S3). During the mixed-group trials, first donations in new dyads were observed on average 247 days after their introduction (95% CI = 227 to 267, range = 66 to 556 days, n = 83 dyads, Figure S2), and the first evidence of new grooming was seen on average 198 days after

their introduction (95% CI = 186 to 209, range = 7 to 546, n = 351 dyads, Figure S2). During the controlled introductions trials, first donations were observed on average 33 days post-introduction (95% CI = 1 to 56, range = 1 to 193, n = 12 dyads) and first grooming was observed on average 24 days post-introduction (95% CI = 0 to 43, range = 1 to 205, n = 23 dyads).

Food sharing relationships emerged faster in isolated pairs than in quartets
    The seven food donations in new relationships in isolated pairs also tended to occur sooner on average (mean latency = 3.6 days, 95% CI = 1.9 to 5, range = 1-8 days) than the three donations that occurred in quartets during the same time period (latency = 6, 32, and 34 days). During controlled introduction trials, food sharing occurred in 6 of 11 possible cases between familiar bats in the quartets but only in 2 of 20 possible cases between unfamiliar bats in those same quartets (OR = 0.09, df = 1, p = 0.012).

New grooming preceded new food sharing more than expected by chance
    It is important to note that our tests of whether new grooming occurs before new food sharing are highly conservative (i.e. biased away from detecting new grooming before new food sharing), because the actual first grooming events in a new pair almost certainly occurred before our first observations of it, whereas the first food donations we observed were very likely to be the actual first donations. Bats were only focal sampled when in need of food, and they were only in need during the fasting trial, because we isolated and fed them immediately after. Food donations were therefore only necessary and likely during the 1-hour trial when we observed them. In contrast, grooming in the same pair could occur in any hour during the several days between the trials where the same dyad was sampled again (median = 8 days, inter-quartile range = 5 to 14 days). We sampled close to 100% of the time when food sharing was actually necessary, but less than 2% of the time when grooming could have occurred. Moreover, fasting trials increase the probability the subject will receive food, they also decrease the probability it will groom others (see 'Grooming Symmetry' below). In summary, our approach was very conservative: even if we observed the first grooming and sharing events during the same fasting trial, it is very likely that the first grooming actually occurred in the days before this trial, making any evidence for raising-the-stakes hard to detect.
    Despite this conservative bias, we still observed new grooming events before new sharing more than expected based on their relative frequencies. We observed new food sharing in 87 new pairs (30 adult past stranger pairs and 57 pairs with a captive-born bat). In most of these new sharing pairs (50 of 87), we observed new grooming on previous days before the fasting trial with the first food donation. In 28 of 87 pairs, we observed the first grooming during the same trial as the first sharing event. In the remaining 9 pairs (2 adult female pairs and 7 pairs with captive-born bats), we observed the first new donation in a trial without grooming. Note that this does not mean that absolutely no grooming occurred in these trials: food-sharing interactions almost always begin with sniffing and social grooming immediately prior to mouth-licking, but we only counted grooming bouts that were 5 seconds or longer, and the initial interactions were missed by the observer. Regardless, we observed grooming before or during the first trial with food sharing in 90% of the new sharing pairs, which is more than three times higher than expected when the dates of trial observations were shuffled within each new pair and type of behavior (p < 0.0002, expected frequency = 27%, 95% CI = 18% to 37%). To test whether the duration of the time lag between new grooming and sharing observations was larger than expected, we also applied the same permutation procedure to test the significance of the mean

latency between new grooming and sharing observed on different days. We found that the mean latency was also greater than expected based on the relative frequency of new grooming and new sharing (observed = 85 days, expected = 58 days, n = 59, p = 0.013).

Grooming trajectories over time predicted new sharing

The age composition of new potential relationships affected the pre-donation grooming rate trajectories. For adult past strangers, the grooming probabilities increased for all recipients, including those that never donated (OR = 1.12, p = 0.004), and they increased significantly faster for grooming recipients that later donated (OR = 1.49, p < 0.0002; interaction: OR = 1.45, p = 0.032, Figure 1b). For new potential relationships with captive-born bats, however, the grooming probabilities actually decreased for grooming recipients that never donated (OR = 0.9, p = 0.01), but tended to increase for recipients that did later donate (OR = 1.32, p = 0.051; interaction: OR = 1.70, p < 0.0002), Figure 1c).

Grooming before versus after new sharing

Grooming increased before but not after first donations in new relationships. The same pattern (Fig. 2) was found in new relationships between adults (interaction: OR = 1.60, p = 0.033; before: OR = 1.49, p = 0.024; after: OR = 1.01, p = 0.88) and in new relationships with captive-born bats (interaction: OR = 1.7, p = 0.0004; before: OR = 1.32, p = 0.010; after: OR = 0.92, p = 0.6).

Reciprocal development of food sharing

New food sharing rates were more symmetrical across dyads than expected by chance (Mantel test: r = 0.54, p = 0.0002). Among adult past strangers, the proportion of previous trials in which bat A fed B predicted the occurrence of the first new reciprocal donation from bat B to A (OR = 6.00, n = 235, p = 0.017). The number of previously unfamiliar pairs that donated food in both directions during the study period was greater than expected if new donations were random (p < 0.0002, observed bidirectional pairs = 13, expected = 4.6, expected 95% CI = 1 to 9). The probability of reciprocation is low because new food sharing rates were overall low, all bats had access to multiple donors, and most sharing occurred among familiar bats with established relationships.

Grooming symmetry

Previous studies of raising-the-stakes have focused on grooming symmetry within short time periods [4-9], but our experimental design did not allow for this kind of analysis. Grooming rates in new relationships increased over time, predicted reciprocal food-sharing rates, and were symmetrical across dyads over the study period (mantel test: r = 0.77, p < 0.0002), but we could not reliably test whether grooming symmetry increased over time within each dyad, for three main reasons. First, fasting trials reduced grooming symmetry, because the fasted bat was often initially greeted by many groupmates and it also spent less time grooming and more time begging (trying to lick the mouth of a potential donor). Consequently, a fasted subject was twice as likely to be groomed by a groupmate (13% probability) than to groom a groupmate (6% probability). Second, observed grooming rates that are undersampled are expected to become more balanced over time merely as a byproduct of grooming rates increasing over time, and we know that grooming rates increased over time. Third, increases in grooming symmetry over time

could be driven by age effects, because mutual grooming (and hence grooming symmetry) is much lower when one bat is not yet an adult [10].

Evidence that new grooming and sharing are not both caused by proximity

One null hypothesis is that bats initiate new grooming and sharing based entirely on proximity, and the correlation between new grooming and new sharing is spurious. In this case, grooming rates should correlate with new sharing among strangers that are able to freely associate during the mixed-group period, but when strangers are forced into close proximity as in the controlled introduction trials, then the apparent grooming effect should disappear or be much smaller. In other words, if proximity is actually driving the correlation between grooming and sharing, then removing variation in proximity (forced close contact) should reveal the lack of an association between grooming and sharing. In sharp contrast to this prediction, the effect of grooming given on new food received for the strangers held in constant close proximity during the controlled introduction periods was even *greater* than the effect during the mixed-group trials where proximity was allowed to vary greatly across dyads and over time (forced close proximity: OR = 5.44, p = 0.036; variable proximity: OR = 1.63, p = 0.037; network logistic regression in the sna R package).

References
1. Carter GG & Wilkinson GS (2013) Carter GG & Wilkinson GS (2015) Social benefits of non-kin food sharing by female vampire bats. Proc. R. Soc. B 280(1753): 20122573.
2. Carter GG & Wilkinson G (2013) Does food sharing in vampire bats demonstrate reciprocity? Commun. Integr. Biol. 6(6):e25783.
3. Carter GG & Wilkinson GS (2015) Social benefits of non-kin food sharing by female vampire bats. Proc. R. Soc. B 282(1819):20152524.
4. Bshary R (2002) Building up relationships in asymmetric co-operation games between the cleaner wrasse *Labroides dimidiatus* and client reef fish. Behav. Ecol. Sociobiol. 52(5):365-371.
5. Kaburu SS & Newton-Fisher NE (2013) Social instability raises the stakes during social grooming among wild male chimpanzees. Anim. Behav. 86(3):519-527.
6. Barrett L, Henzi SP, Weingrill T, Lycett JE, & Hill RA (2000) Female baboons do not raise the stakes but they give as good as they get. Anim. Behav. 59(4):763-770.
7. Manson JH, Navarrete CD, Silk JB, & Perry S (2004) Time-matched grooming in female primates? New analyses from two species. Anim. Behav. 67(3):493-500.
8. Fruteau C, Lemoine S, Hellard E, van Damme E, & Noë R (2011) When females trade grooming for grooming: Testing partner control and partner choice models of cooperation in two species of primates. Anim. Behav. 81:1223-1230.
9. Roberts G & Renwick JS (2003) The development of cooperative relationships: an experiment. Proc. R. Soc. B 270(1530):2279-2283.
10. Wilkinson GS (1986) Social grooming in the common vampire bat, *Desmodus rotundus*. Anim. Behav. 34:1880-1889.

SI Figures and Tables

Figure S1 | Experiment overview
To see how vampire bats form new social bonds, we created groups of bats from two different sites (colors), then we induced and sampled food sharing and grooming events between bats that are either previously familiar or unfamiliar. Red arrows depict food sharing events during repeated fasting trials. For details of controlled introductions, see SI Table 2. Icons from icons8.com used under a Linkware license.

Figure S2 | First food donations over time
New food-sharing relationships accumulated gradually over time. Black rectangles above X-axis
show the occurrence of fasting trials.

Figure S3 | Gradual development of new food-sharing relationships
Panel A shows food donations (points) over time (x-axis) within new actor-receiver relationships
(y-axis) between two adult females (grey points) or with a captive-born bat (black points).
Repeated dyadic donations are connected by horizontal lines. The end of the controlled
introduction period, after which all bats could interact freely (months 1-4), is shown by the
vertical dotted line. Black rectangles above the x-axis show the fasting trials, when new
donations could be observed. Panel B shows the monthly formation of the food-sharing network
between Las Pavas bats (orange), Tolé bats (purple), and captive-born bats (white). Grey edges
show one-way sharing and red edges show two-way sharing. Two-way sharing occurred more
often than expected by chance (see results).

Figure S4 | Dyadic grooming rates predict new food-sharing relationships.
Mean within-dyad grooming rates, with bootstrapped 95% confidence intervals, are shown for
three possible outcomes (y-axis) and for all potential relationships (black), potential relationships
with captive-born bats (grey), and potential relationships between adult strangers (light grey).

Table S1 | Glossary.

Definition of terms used in the text.

Term	Definition
Dyad	An undirected pair of bats (e.g. AB, BC, AC)
Potential relationship	A directed pair of actor and receiver bats (e.g. AB, BA, AC)
Relationship	A directed actor-receiver pair that is observed to groom or share food during fasting trials.
New relationship	Relationship between bats that first met during the experiment, excluding mother-offspring dyads.
Social bond	The unobserved underlying social relationship (as experienced by the animal) that we infer from observations.

Table S2 | Controlled introductions
The same bats were used in multiple introductions. Bats were moved to and from groups to make new combinations or because of health issues (pregnancy, weight loss). Bats not in a small cage group during controlled introduction trials were kept with familiar individuals in a flight cage.

No.	Group type	No. trials (range of days together)	Adult female bats (*Las Pavas stranger)	Opportunities for new sharing	Introduction date
1	quartet	1 (1 day)	scs, hilga, rc, eve*	3	2016.07.06
2	quartet	1 (1 day)	ccs, sss, sc, una*	3	2016.07.06
3	quartet	1 (1 day)	scc, sd, c, dos*	3	2016.07.06
4	quartet	1 (1 day)	csc, ss (w/pup), s, tes*	3	2016.07.06
5	pair	1 (1 day)	ccc, cat*	1	2016.07.06
6	pair	1 (1 day)	dcd, ivy*	1	2016.07.06
7	pair	1 (1 day)	dd, six*	1	2016.07.06
8	pair	1 (4 days)	d (w/pup), ola* (w/pup)	1	2016.07.02
9	quartet	17 (1–44 days)	sd, scs, d (w/pup), una*	32	2016.08.24
10a	quartet	5 (1–9 days)	s, rc, hilga, dos*	9	2016.08.24
10b	quartet	12 (1–44 days)	s, rc, ccc (w/pup), dos*	21	2016.09.21
11	quartet	17 (1–44 days)	ccs, sc, sss, tes*	27	2016.08.24
12	pair	10 (1–44 days)	dd, cat*	10	2016.08.24
13	pair	10 (1–44 days)	c, ivy*	10	2016.08.24
14	pair	5 (1–10 days)	csc, six*	5	2016.08.24
15	pair	9 (1–44 days)	dcd, eve	9	2016.08.24
16	pair	9 (1–97 days)	ss (w/pup), ola* (w/pup)	19	2016.08.24
17	pair	4 (1–7 days)	cd, six*	4	2016.09.21

<u>Summary of supplied data</u>
Dataset S1. genotypes.csv
Microsatellite genotypes used to assess relatedness.
Dataset S2. vampire_maternal_kinship.csv
Maternal pedigree data
Dataset S3. new_bonds_data.Rdata
Food sharing and allogrooming data
Dataset S4. new_bonds_analysis20.txt
R script for analyzing data