1
2 **Title:**
3 Development of new food-sharing relationships among nonkin vampire bats
4

5 **Authors and affiliation:**
6 Gerald G. Carter[1,2], Damien R. Farine[3,4,5,6], Rachel J. Crisp[2,7], Julia K. Vrtilek[2,8], Simon P. Ripperger[1,2,9],
7 Rachel A. Page[2]
8
[1]Department of Ecology, Evolution, and Organismal Biology; The Ohio State University, Columbus, OH,
USA
[2]Smithsonian Tropical Research Institute, Balboa, Ancón, Panamá
[3]Edward Grey Institute of Field Ornithology, Department of Zoology, University of Oxford, Oxford, UK
[4]Department of Collective Behaviour, Max Planck Institute of Animal Behavior, Konstanz, Germany
[5]Chair of Biodiversity and Collective Behaviour, Department of Biology, University of Konstanz,
Konstanz, Germany
[6]Center for the Advanced Study of Collective Behaviour, University of Konstanz, Konstanz, Germany
[7]Centre for Research in Animal Behaviour, University of Exeter, Exeter, UK
[8]Department of Biology, Eidgenössische Technische Hochschule Zürich, Zürich, Switzerland
[9]Museum für Naturkunde, Leibniz-Institute for Evolution and Biodiversity Science, Berlin, Germany
20
21 **Corresponding author:**
22 Gerald Carter, 318 W. 12th Avenue, 300 Aronoff Laboratory, Columbus, OH 43210, carter.1640@osu.edu

Abstract

23
Social relationships that involve costly helping occur most often among kin, but in many
complex and individualized animal societies, nonkin also demonstrate stable cooperative relationships
that share similarities with human friendship. How do such cooperative bonds form between complete
strangers? Here, we show evidence that unfamiliar nonkin vampire bats (*Desmodus rotundus*) selectively
escalate low-cost investments in allogrooming before developing higher-cost food-sharing relationships.
By introducing female bats from geographically distant sites in captive pairs or groups and fasting them
repeatedly over 15 months, we observed that increasing rates of grooming a partner predicted the
occurrence of that partner's first food donation back to the groomer, after which grooming rates no
longer increased. New food-sharing relationships formed in 14.5% of 608 possible female pairs,
emerged in a reciprocal pattern, and developed more often when strangers lacked alternative familiar
partners. Our results are consistent with predictions from the 'raising-the-stakes' hypothesis that
strangers 'test the waters' of new relationships by initially making low-cost investments in grooming
before making higher-cost food donations. This form of 'raising the stakes' (e.g., transitions from
clustering to grooming to food sharing) might play an underappreciated role in many other social
decisions with long-term consequences, such as joining a new social group or choosing a long-term
mate.
Keywords: cooperation, social relationships, vampire bats
41

Significance statement
Vampire bats form long-term cooperative social bonds that involve reciprocal regurgitated blood
sharing. But how do two individuals go from complete strangers to reciprocal food donors? By
introducing unfamiliar bats, we found evidence that low-cost grooming paves the way for higher-cost
food donations. Bats that formed new food-sharing relationships had a history of escalating reciprocal
grooming up until the food sharing began. Food sharing emerged in a reciprocal fashion and it emerged
more often when two strangers could not access their original groupmates. The finding that unfamiliar
nonkin vampire bats appeared to gradually and selectively transition from low-cost to high-cost
cooperative behaviors is the first evidence that nonhuman animals 'raise the stakes' when forming new
cooperative relationships.
52
53

Text

Animal societies are fundamentally shaped by repeated interactions among individuals over time. Repeated interactions allow individuals to choose to cooperate based on their past experience across different partners (1-3). Organisms as diverse as animals, plants, and fungi have demonstrated partner choice: individuals prevent exploitation by shifting their cooperative investments towards partners that provide better reciprocal returns (3-7). Across several nonhuman mammals, repeated cooperative interactions lead to adaptive and enduring social bonds that share similarities with human friendship (8-12), but it remains unclear how these initially form. A significant challenge has been understanding how individuals prevent exploitation while forming these stable bonds. How do complete strangers develop a long-term cooperative relationship?

A key idea is that individuals should reduce the risk of exploitation by initially spreading out smaller cooperative investments across time ('parceling' (13)) or across different partners ('social bet-hedging' (14)), and then gradually escalating investments in the most cooperative partnerships ('raising the stakes' (15)). For example, one might first assess a potential partner's tolerance by clustering for warmth, then gain feedback by grooming the partner, and then use the partner's response to decide whether to provide higher-cost food donations or coalitionary support (16). Despite its intuitive appeal for explaining how new cooperative relationships develop, evidence supporting the 20-year-old 'raising-the-stakes' model (15) is surprisingly scarce. An early test using the cleaner and client fish mutualism suggested that the model does not apply well to situations with severe asymmetries in partner payoffs or options (17). Studies with nonhuman primates (18-21) have tested only snapshots of established relationships rather than the formation of new ones. Human strangers 'raise the stakes' when making monetary bids in cooperation games (e.g. 22, 23), but we currently lack supporting evidence for this strategy in the more ecologically-relevant context of relationship formation. Gathering this evidence requires measuring the emergence of natural helping behaviors between randomly introduced strangers.

We tracked the development of new cooperative relationships between previously unfamiliar wild-caught vampire bats (*Desmodus rotundus*) over 15 months. Cooperative relationships in this species involve low-cost allogrooming (hereafter *grooming*) and higher-cost regurgitations of ingested blood, or *food sharing* (14, 24-28). We found support for several lines of evidence that vampire bats use reciprocal grooming to gradually establish new bonds that include food donations. If bats choose partners, then new food-sharing relationships should form more often when bats have fewer alternative partners. If grooming gradually leads to sharing, then grooming rates should predict the probability that the grooming recipient later donates food back to the groomer. Grooming rates should also start low and increase over time but only up until the first reciprocal food donation. Finally, new food sharing should be rarer, occur after mutual grooming, and emerge in a reciprocal fashion.

Female vampire bats demonstrate kin-biased fission-fusion social dynamics (24, 27, 28). New nonkin social bonds can form when an unrelated female joins a social network about once every two years (24, 28), with individual bats living for up to 16 years in the wild (29). To observe how new food-sharing relationships form between adults, we captured adult females from two distant sites in Panamá, Tolé (n=19) and Las Pavas (n=8), and we then ran 638 fasting trials in which an overnight-fasted subject could be fed by a previously unfamiliar bat from another site. To test the prediction that new sharing relationships would form more often when strangers have fewer partner options, we compared the occurrence of new sharing when wild-caught strangers were introduced in isolated pairs (one Las Pavas and one Tolé bat), in small groups (one Las Pavas and three Tolé bats), or in one large mixed group (all bats together; see Methods, Supplementary Information (SI) Appendix, Fig. S1). New bonds can also form when individuals are born into a group, and these relationships might form differently. We therefore also measured the development of non-maternal cooperative relationships between 26 female adults and 13 younger captive-born bats (6 males and 7 females, 11 to 21 months old) in the large mixed group.

To test our hypotheses, we compared the observed coefficients from general and generalized linear models (slopes β, and odds ratios OR, respectively) to expected distributions of coefficient values expected under the null hypotheses using permutations of the network or the event data (see Methods). We use the term 'potential relationship' for a pair of bats that could have groomed or shared food, the word 'relationship' for an *observed* network edge (directed), and the word 'bond' to discuss the underlying construct that we *inferred* from the observed relationship (see SI, Table S1).

Fasted bats were fed by at least one donor in 61% of trials (SI Appendix 2). Over 424 days and
12,012 opportunities for new food donations, new food sharing developed in 10.8% of the 996 potential
relationships among all bats, 14.5% of 608 potential relationships among females, and 15.6% of 243
potential relationships among wild-caught adult females (SI Appendix 3). All bats had at least one donor
(range=1-16, mean=6.6). The average number of new food donors per adult female bat was 2.7
(range=0-7) and the average per captive-born bat was 2.6 (range=0-6). New grooming relationships
developed far more frequently (all bats=51.9% of 1008; females=58.9% of 618; wild-caught adult
females=78.2% of 248). The average number of new groomers was 7.2 (range=0-16) for adult females
and 14.4 (range=1-23) for captive-born bats.

If bats choose new partners based on their phenotype alone, then relationships should form
more often when bats have more alternative partners. On the other hand, if bats 'test the waters' of each
new relationships, they should choose partners based on both the availability of different partners and
their past experiences with each, and food-sharing relationships should therefore form more often when
bats have *fewer* alternative partners. As expected, when strangers from Las Pavas and Tolé were
introduced and housed as isolated pairs, we observed higher rates of new food sharing (β=1.14,
p=0.002) and new grooming (β=1.09, p=0.02) compared to when one Las Pavas bat was introduced to
three Tolé bats, despite there being fewer potential new bonds available to form (SI Appendix 4). When
we aggregated bats from the controlled introduction trials into a large mixed group, bats preferentially
fed and groomed their original familiar groupmates, and new sharing emerged even more gradually than
in the isolated pairs or in small groups (SI Appendix 5, Fig. S2 and S3).

If the bats use low-cost grooming to build higher-cost sharing bonds, then the grooming rate
should predict the probability of the first food donation in the opposite direction. As expected, new food
sharing emerged on days after we observed mutual grooming more than expected by chance (SI
Appendix 6), and the grooming rate given by actor A to recipient B predicted the later occurrence of new
food sharing from B back to A (OR=2.15, p=0.0002, n=897). The trajectory of grooming rates over time
clearly differed between pairs that developed new food-sharing relationships versus pairs that did not
(interaction: OR=1.60, p<0.0001, Fig. 1). The slope of this increase in grooming was also greater *before*
the first reciprocal food donation than after. Initial grooming rates started low, then increased over time
up until the new food-sharing relationship formed (Fig. 2).

137
138

139
Fig. 1. Increasing A-to-B grooming led to new B-to-A food-sharing relationships. In cases where a
new food-sharing relationship formed (solid line), the grooming rate towards the future donor increased
over time before the first donation occurred (OR=1.40, n=33, p<0.0001), but the grooming rate towards a
potential donor remained low in cases where no food-sharing relationship formed (dashed line; OR=0.99,
n=420, p=0.58). This divergence in all potential new relationships (panel a) was also detected within
previously unfamiliar adults (panel b), and within relationships with captive-born bats (panel c), which
had more divergent grooming trajectories (SI Appendix 7). Shading shows the 95% CI for the fitted
model's predictions.
148

Fig. 2. Grooming rates increased before, but not after, new food-sharing occurred. The probability of a focal bat grooming the new donor in a 1-h trial (y-axis) increased before the first day that the donor fed the focal bat (i.e. 'day zero'; OR=1.4, p=0.0005), but not after this day zero (OR=1.01, p=0.47; interaction: OR=1.57, p=0.0003). This effect was seen in new food-sharing relationships with or without captive-born bats (SI Appendix 8). Shading shows the 95% CI for the fitted model's predictions.

Emergence of new food sharing was more reciprocal than expected by chance, even when controlling for kinship (MRQAP-DSP; reciprocal sharing: β=0.33, p<0.0002, kinship: β=0.02, p=0.65; SI Appendix 9). Grooming rates in new relationships were also symmetrical across dyads but we lacked the power to determine whether grooming symmetry increased over time within dyads (see SI Appendix 10). Grooming rates were highest between bats that formed two-way food-sharing relationships, intermediate in relationships where we observed sharing in only one direction, and lowest in pairs where we never saw food sharing (SI, Fig. S4).

The rarity of new food-sharing relationships corroborates past evidence that food regurgitations are energetically costly and that food-sharing bonds require investments of time and energy (14, 24, 30, 31). The relationship between new grooming and new food sharing was unlikely to be caused by mere proximity because the effect of new grooming on new food sharing remained evident regardless of whether or not bats were forced into close proximity (SI Appendix 11).

These findings provide the clearest evidence to date that nonkin food sharing in vampire bats is not a byproduct of kin selection (26). Before this study, one hypothesis was that food sharing decisions among nonkin could depend entirely on heuristics based on phenotypic similarity, resulting in a spurious pattern of symmetrical helping that looks like reciprocity (32, 33). However, this hypothesis incorrectly predicts that food-sharing relationships should form immediately and occur most frequently in larger groups simply because there are more opportunities for similar matching phenotypes.

Our results were consistent with the hypothesis that relationships formation occurs through some form of 'raising the stakes' (15). This model has yet to be tested during the transition from 'strangers' to 'friends' because these changes are difficult to document in nature (see SI Appendix 12). Past evidence for the 'raising-the-stakes' strategy (15) has also been scarce in part because it is a variation on the classic 'tit-for-tat' strategy in the iterated prisoner's dilemma (1), a model which is difficult to test using natural forms of cooperation (4, 32, 34). 'Tit-for-tat' forms of reciprocity are demonstrated by experiments with trained instrumental tasks and payoffs that accrue in distinct rounds, such as rats taking turns to pull a lever to deliver food (4-6). However, the 'tit-for-tat' model excludes many factors crucial in the real world, including partner choice, partner fidelity, exchange of multiple service types, and the many cost-benefit asymmetries resulting from demography, market effects, and social rank (4, 7, 34). If social bonding involves integrating many different kinds of social interactions into a single positive association, one should not expect clearly alternating exchanges of help. In primates,

cooperation within long-term social bonds does not produce strict 'tit-for-tat' exchanges of help;
strongly bonded partners show *less* evidence for short-term contingencies in grooming (9).
To clearly demonstrate that an actor's cooperative investments are contingent on a partner's
previous behavior, one must prevent reciprocation and then detect a subsequent decrease in the actor's
cooperative investment. This evidence of reciprocity has yet to be experimentally demonstrated in food-
sharing vampire bats or in any other long-term social relationship (SI Appendix 13). Our findings show
that such an experiment would be most powerful if researchers targeted newly developing relationships
rather than established ones, and if they tracked multiple cooperative behaviors rather than just one.
Past studies on 'raising the stakes' during relationship development have focused on increasing rates of
a single cooperative behavior (18-23), but individuals can also raise the stakes by adding new higher-
cost behaviors. Our findings suggest that female vampire bats do both, first increasing grooming rates
and then transitioning from low-cost grooming to high-cost food-sharing.
The relevance of our findings extends beyond high-cost cooperative behaviors. For example, in
some species, courtship behaviors could be seen as a short-term investment in the formation of longer-
term pair bonds with substantial fitness consequences (35). Similarly, the role of mere physical contact
as a low-cost method for building tolerance and trust might be more general than currently recognized.
The key role of grooming for relationship maintenance in primates is well established, but growing
evidence suggests that similar tactile behaviors can reduce fear and encourage tolerance and
cooperation in many other species of mammals, birds, and fish (e.g. 5, 36-42). Recently developed
methods for tracking formation of social bonds at fine temporal scales (43, 44) could provide new
opportunities to test whether gradual escalation of proximity and body contact is a widespread
mechanism for socially 'testing the water'.
208
Methods
Animals
We conducted experiments at the Smithsonian Tropical Research Institute in Gamboa, Panama.
We used 41 common vampire bats (*Desmodus rotundus*) as subjects, including 19 female bats captured
exiting a roost in Tolé, Panamá; 8 female bats captured foraging at a cattle pasture in Las Pavas,
Panamá about ~215 km from Tolé; and 14 captive-born bats (8 females, 6 males). We studied adult
females and their young, because these individuals form the basis of food-sharing networks in the wild,
whereas adult males compete for access to territories and females and do not form stable bonds as
often (24-28). To ensure familiarity within groups and unfamiliarity between groups, we housed the
groups separately (Tolé bats for 6 months and Las Pavas bats for 2 weeks) before the study began. Bats
were marked with subcutaneous passive integrated transponders (Trovan Ltd. USA) and a visually
unique combination of forearm bands (Porzana, National Tag, and birdbands.com). To feed bats, we
provided refrigerated or thawed cattle or pig blood defibrinated with sodium citrate and citric acid.
We used a 3-4 mm biopsy punch to collect tissue samples in 80% or 95% ethanol, then used a
salt–chloroform procedure for DNA isolation, and a LI–COR Biosciences® DNA Analyser 4300 and the
SAGA GT allele scoring software to genotype individuals at 17 polymorphic microsatellite loci. Allele
frequencies were based on 100 bats from Tolé and 9 bats from Las Pavas, respectively. Genotypes were
99.9% complete. To estimate genetic relatedness, we used the Wang estimator in the R package
'related'. To estimate kinship, we assigned a zero kinship to known unrelated individuals from different
sites and to individuals with negative pairwise relatedness, and we assigned a kinship of 0.5 for known
mother-offspring pairs or pairs with genetic relatedness estimates greater than 0.5. For all other pairs,
we used genetic relatedness as the estimate for kinship.
231
Experimental design
We induced allogrooming and regurgitated food sharing using a fasting trial, in which a focal
subject was isolated from the group without food for a night and a day, then released back to the group
of fed bats for 1 hour the following night. During the hour, all grooming or food-sharing interactions with
the subject were recorded using an infrared (IR) light and an IR-sensitive video camera. Each food
sharing bout was estimated by the number of seconds that the unfed subject spent licking the mouth of
a particular groupmate. Grooming was defined as chewing or licking the fur or wings of another bat. The
dyadic sharing or grooming for a trial was estimated as the sum of all bouts that were at least 5 seconds

long. We weighed bats before and after trials. Observed mouth-licking durations predicted weight gain
(SI Appendix 1).
We conducted fasting trials in each group during three experimental phases (SI, Fig. S1). First, we
conducted 57 'baseline' trials to assess preliminary sharing rates between the 19 Tolé bats housed in a
1.7 x 2.1 x 2.3 m outdoor flight cage (3,420 possible sharing interactions in one group). Second, we
conducted 106 'controlled introduction' trials to assess possible formation of new food-sharing bonds
between bats introduced as either an isolated pair (one Las Pavas bat and one Tolé bat) or a quartet
(one Las Pavas bat and three Tolé bats), housed in a 28 x 28 x 40 cm clear plastic observation cage (10
pairs and 8 quartets). These controlled introductions provided for 162 opportunities for new food sharing
between previous strangers (SI, Table S2). Finally, we conducted 532 'mixed-group' trials to assess the
formation of new sharing relationships when all bats were housed together in the flight cage described
above (19 Tolé, 7 Las Pavas, and 14 captive-born bats). The introductions in this combined group
provided 11,823 more opportunities for new sharing.
253
Statistical analyses
During the baseline and mixed-group trials, we estimated food donation size as the number of
seconds that a fasted subject spent mouth-licking a fed groupmate. During the controlled introduction
trials, however, when bats were forced in close proximity, we saw a greater frequency of begging,
defined as mouth-licking that is clearly not food-sharing because the partner is turning away from the
mouth-licking bat and the mouth-licking bat does not gain the weight that would be expected from food-
sharing. To be conservative when measuring sharing, we therefore did not count mouth-licking as food
sharing during the controlled introduction trials unless the subject weighed more than expected based
on the average weight change for bats that did not perform any mouth-licking.
Durations of sharing and grooming were lognormal. To create a standard index of grooming rates,
we therefore transformed the total duration of directed dyadic interactions in each trial using natural log
(x+1). We call these measures of the log duration per hour 'rates'. When interaction bout duration and
probability had different meanings, we decomposed rates into two separate response variables:
amounts (the magnitude of nonzero rates in a trial) and probabilities (the presence or absence of a
nonzero rate in a trial). We used permutation tests with 10,000 permutations for p-values and
bootstrapping for all 95% confidence intervals. Null distributions were not always centered on zero due
to structure in the data, so caution must be taken when considering the observed coefficients.
Grooming could occur before sharing simply because it is more frequent. To test whether mutual
grooming preceded new sharing more than expected by chance, we compared the observed probability
of observing mutual grooming before new sharing to the values expected from a null model based on
randomly swapping the label of interactions (grooming versus sharing) within each dyad. This
permutation test controls for the relative frequency and timing of grooming and sharing events in new
dyads. To test for ingroup-outgroup biases in sharing for each site, we calculated observed coefficients
for the effect of the actor and receiver being from the same capture site on actor grooming rates, then
we calculated expected coefficients by permuting the grooming rates within each actor to different
possible recipients.
To test the effects of kinship and reciprocal grooming on the formation of new food-sharing
relationships in the mixed-group trials, we used multiple regression quadratic assignment procedure
with double semi-partialing (MRQAP-DSP) via the netlogit function in the sna R package. We also used
this method to test the effect of grooming on occurrence of new sharing only within the controlled
introduction trials. This procedure uses generalized linear models via the glm function in lme4 package
to calculate the observed coefficients and uses network-level permutations to get expected coefficients.
Since MRQAP-DSP cannot test interaction effects, we compared observed and expected interaction
coefficients using permutations in which we shuffled trial rates given by the actor among different
possible receivers and then shuffled the trial rates received by the receiver among different possible
actors. If the interaction coefficients were significant (p<0.05), we conducted separate MRQAP-DSP
tests within each group.
To test whether interaction rates changed over time, we generated expected coefficients for
general or generalized linear models by permuting the order of interactions within each potential
relationship. One captive-born bat died for unknown reasons during the mixed-group trials, so we

removed it from all temporal analyses. To test for evidence of reciprocal sharing, we used MRQAP-DSP
to test if the matrix of new sharing in the mixed-group trials was predicted by reciprocal sharing when
controlling for kinship. As an additional test, we also counted the occurrence of both novel sharing and
reciprocal sharing for all new potential relationships, then counted the same number after randomizing
the presence of sharing across potential relationships.

Data availability
  Behavioral data, genotypes, and R code are available as supplementary information.

Acknowledgments
We thank the Smithsonian Tropical Research Institute for logistical support. Isabelle Waurick conducted
the molecular lab work. Frieder Mayer enabled the molecular lab work. Work was supported by a
Smithsonian Postdoctoral Fellowship, a Humboldt Research Fellowship, a Smithsonian Institution
Scholarly Studies Grant and a grant from the National Geographic Society Committee for Research and
Exploration (WW-057R- 17).

Author contributions
Conceptualization, GC; Methodology, GC; Investigation, GC, RC, JV; Genotyping, SR; Statistical
analysis, GC; Original draft, GC; Review & Editing, GC, RC, JV, SR, DF, RP; Funding acquisition, GC,
SR, RP; Resources, RP; Supervision, GC, DF, RP;

References
1   Axelrod R & Hamilton WD (1981) The evolution of cooperation. Science 211:1390-1396.
2   Noë R & Hammerstein P (1994) Biological markets: supply and demand determine the effect of
    partner choice in cooperation, mutualism and mating. Behav. Ecol. Sociobiol. 35:1-11.
3   Trivers RL (1971) The evolution of reciprocal altruism. Q. Rev. Biol. 46:35-57.
4   Carter GG (2014) The reciprocity controversy. Anim. Behav. Cogn. 1(3):368-386.
5   Schweinfurth MK & Taborsky M (2018) Reciprocal trading of different commodities in Norway rats.
    Curr. Biol. 28:1-6.
6   Rutte C & Taborsky M (2008) The influence of social experience on cooperative behaviour of rats
    (*Rattus norvegicus*): direct vs generalised reciprocity. Behav. Ecol. Sociobiol. 62(4):499-505.
7   Fruteau C, Voelkl B, van Damme E, & Noë R (2009) Supply and demand determine the market value
    of food providers in wild vervet monkeys. Proc. Natl. Acad. Sci. USA 106:12007-12012.
8   Samuni L, et al. (2018) Social bonds facilitate cooperative resource sharing in wild chimpanzees.
    Proc. R. Soc. B 285(1888).
9   Seyfarth RM & Cheney DL (2012) The evolutionary origins of friendship. Ann. Rev. Psych. 63:153-
    177.
10 Silk JB, et al. (2010) Strong and consistent social bonds enhance the longevity of female baboons.
    Curr. Biol. 20(15):1359-1361.
11 Silk JB, et al. (2009) The benefits of social capital: close social bonds among female baboons
    enhance offspring survival. Proc. R. Soc. B 276(1670):3099-3104.
12 Silk JB, Alberts SC, & Altmann J (2003) Social bonds of female baboons enhance infant survival.
    Science 302(5648):1231-1234.
13 Connor RC (2010) Cooperation beyond the dyad: on simple models and a complex society. Philos.
    Trans. R. Soc. Lond., Ser. B: Biol. Sci. 365(1553):2687-2697.
14 Carter GG, Farine DR, & Wilkinson GS (2017) Social bet-hedging in vampire bats. Biol. Lett.
    13(5):20170112.
15 Roberts G & Sherratt TN (1998) Development of cooperative relationships through increasing
    investment. Nature 394(6689):175-179.
16 Borgeaud C & Bshary R (2015) Wild vervet monkeys trade tolerance and specific coalitionary
    support for grooming in experimentally induced conflicts. Curr. Biol. 25(22):3011-3016.
17 Bshary R (2002) Building up relationships in asymmetric co-operation games between the cleaner
    wrasse *Labroides dimidiatus* and client reef fish. Behav. Ecol. Sociobiol. 52(5):365-371.

18 Kaburu SS & Newton-Fisher NE (2013) Social instability raises the stakes during social grooming
among wild male chimpanzees. Anim. Behav. 86(3):519-527.
19 Barrett L, Henzi SP, Weingrill T, Lycett JE, & Hill RA (2000) Female baboons do not raise the stakes
but they give as good as they get. Anim. Behav. 59(4):763-770.
20 Manson JH, Navarrete CD, Silk JB, & Perry S (2004) Time-matched grooming in female primates?
New analyses from two species. Anim. Behav. 67(3):493-500.
21 Fruteau C, Lemoine S, Hellard E, van Damme E, & Noë R (2011) When females trade grooming for
grooming: Testing partner control and partner choice models of cooperation in two species of
primates. Anim. Behav. 81:1223-1230.
22 Roberts G & Renwick JS (2003) The development of cooperative relationships: an experiment. Proc.
R. Soc. B 270(1530):2279-2283.
23 Van den Bergh B & Dewitte S (2006) The robustness of the 'Raise-The-Stakes' strategy: Coping with
exploitation in noisy Prisoner's Dilemma Games. Evol. Hum. Behav. 27(1):19-28.
24 Wilkinson GS (1984) Reciprocal food sharing in the vampire bat. Nature 308:181-184.
25 Wilkinson GS (1986) Social grooming in the common vampire bat, *Desmodus rotundus*. Anim.
Behav. 34:1880-1889.
26 Carter GG & Wilkinson G (2013) Does food sharing in vampire bats demonstrate reciprocity?
Commun. Integr. Biol. 6(6):e25783.
27 Wilkinson GS (1985) The social organization of the common vampire bat. I. Pattern and cause of
association. Behav. Ecol. Sociobiol. 17:111-121.
28 Wilkinson GS (1985) The social organization of the common vampire bat: II. Mating system, genetic
structure, and relatedness. Behav. Ecol. Sociobiol. 17:123-134.
29 Delpietro H, Russo R, Carter G, Lord R, & Delpietro G (2017) Reproductive seasonality, sex ratio and
philopatry in Argentina's common vampire bats. R. Soc. Open Sci. 4(4):160959.
30 Carter GG & Wilkinson GS (2015) Social benefits of non-kin food sharing by female vampire bats.
Proc. R. Soc. B 282(1819):20152524-20152524.
31 Carter GG, Wilkinson GS, & Page RA (2017) Food-sharing vampire bats are more nepotistic under
conditions of perceived risk. Behav. Ecol. 28(2):565-569.
32 Clutton-Brock T (2009) Cooperation between non-kin in animal societies. Nature 462:51-57.
33 Hammerstein P (2003) Why is reciprocity so rare in social animals? A protestant appeal. Genetic and
cultural evolution of cooperation (ed. & Hammerstein P), pp. 83–93. Cambridge, MA: MIT Press.
34 Noë R (2006) Cooperation experiments: coordination through communication versus acting apart
together. Anim. Behav. 71:1-18.
35 Ihle M, Kempenaers B, & Forstmeier W (2015) Fitness benefits of mate choice for compatibility in a
socially monogamous species. PLoS Biol. 13(9):e1002248.
36 Kern JM & Radford AN (2018) Experimental evidence for delayed contingent cooperation among
wild dwarf mongooses. Proc Natl Acad Sci U S A 115(24):6255-6260.
37 Kenny E, Birkhead TR, & Green JP (2017) Allopreening in birds is associated with parental
cooperation over offspring care and stable pair bonds across years. Behav. Ecol. 28(4):1142-1148.
38 Grutter AS (2004) Cleaner fish use tactile dancing behavior as a preconflict management strategy.
Curr. Biol. 14(12):1080-1083.
39 Soares MC, Oliveira RF, Ros AF, Grutter AS, & Bshary R (2011) Tactile stimulation lowers stress in
fish. Nat. Comm. 2:534.
40 Schirmer A, Jesuthasan SJ, & Mathuru A (2013) Tactile stimulation reduces fear in fish. Front. Behav.
Neuro. 7:167.
41 Radford AN (2012) Post-allogrooming reductions in self-directed behaviour are affected by role and
status in the green woodhoopoe. Biol. Lett. 8(1):24-27.
42 Radford AN (2011) Preparing for battle? Potential intergroup conflict promotes current intragroup
affiliation. Biol. Lett. 7(1):26-29.
43 Alarcón-Nieto G, et al. (2018) An automated barcode tracking system for behavioural studies in
birds. Meth. Ecol. Evol. 9(6):1536-1547.
44 Ripperger S, Günther L, Wieser H, Duda N, Hierold M, Cassens B, Kapitza R, Koelpin A, & Mayer F
(2019) Proximity sensors on common noctule bats reveal evidence that mothers guide juveniles to
roosts but not food. Biol. Lett. 15(2):20180884.