Distinct RopGEFs successively drive polarization and outgrowth of root hairs

AUTHORS:

Philipp Denninger[1,6], Anna Reichelt[1,6], Vanessa A. F. Schmidt[1], Dietmar G. Mehlhorn[3], Lisa Y.
Asseck[3], Claire E. Stanley[4,5], Nana F. Keinath[1], Jan-Felix Evers[1], Christopher Grefen[3] and
Guido Grossmann[1,2,*]

AFFILIATIONS:

[1]Centre for Organismal Studies (COS), Heidelberg University, Im Neuenheimer Feld 230,
69120 Heidelberg, Germany
[2]Excellence Cluster CellNetworks, Heidelberg University, Im Neuenheimer Feld 267, 69120
Heidelberg, Germany
[3]Center for Plant Molecular Biology, Developmental Genetics, University of Tübingen, Auf
der Morgenstelle 32, 72076 Tübingen, Germany
[4]Institute for Chemical and Bioengineering, ETH Zürich, Vladimir-Prelog-Weg 1, 8093 Zürich,
Switzerland
[5]Agroecology and Environment Research Division, Agroscope, Reckenholzstrasse
191, 8046 Zürich, Switzerland

[6]These authors contributed equally.

*Correspondence:
Guido Grossmann
Centre for Organismal Studies
Im Neuenheimer Feld 230
69120 Heidelberg
Germany
email: guido.grossmann@cos.uni-heidelberg.de
ORCID ID: 0000-0001-7529-9244

SUMMARY

Root hairs are tubular protrusions of the root epidermis that significantly enlarge the exploitable soil volume in the rhizosphere. Trichoblasts, the cell type responsible for root hair formation, switch from cell elongation to tip growth through polarization of the growth machinery to a pre-defined root hair initiation domain (RHID) at the plasma membrane. The emergence of this polar domain resembles the establishment of cell polarity in other eukaryotic systems [1-3]. Rho-type GTPases of plants (ROPs) are among the first molecular determinants of the RHID [4, 5] and later play a central role in polar growth [6]. Numerous studies have elucidated mechanisms that position the RHID in the cell [7-9] or regulate ROP activity [10-18]. The molecular players that target ROPs to the RHID and initiate outgrowth, however, have not been identified. We dissected the timing of the growth machinery assembly in polarizing hair cells and found that positioning of molecular players and outgrowth are temporally separate processes that are each controlled by specific ROP guanine nucleotide exchange factor (GEFs). A functional analysis of trichoblast-specific GEFs revealed GEF3 to be required for normal ROP polarization and thus efficient root hair emergence, while GEF4 predominantly regulates subsequent tip growth. Ectopic expression of GEF3 induced the formation of spatially confined, ROP-recruiting domains in other cell types, demonstrating the role of GEF3 to serve as a membrane landmark during cell polarization.

RESULTS AND DISCUSSION

Temporal analysis of hair cell polarization reveals phased deployment of the tip-growth machinery

To dissect the process of breaking cellular symmetry and initiating polar growth in plants, we analyzed the gradual assembly of the tip growth machinery in trichoblasts using specific markers for cytoskeletal rearrangement, plasma membrane specialization, cell wall modification, vesicle trafficking, and Rho-GTPase signaling. Using stable transgenic *Arabidopsis* lines expressing fluorescently labeled versions of these protein markers, we determined the timing of their polarization at the RHID during trichoblast differentiation. As the *Arabidopsis* root represents a time-axis of development from stem cells to mature cells, we numbered the developmental stages (-7 to +3) within a cell file, with the last cell before bulging labeled -1 and the first cell after the onset of bulging labeled +1 (**Figure 1A**). We used the integral plasma membrane marker GFP-LTI6B [19] as a reference and calculated the polarity index of fluorescently labeled marker proteins to quantify protein accumulation at the RHID (**Figure 1B, C**). Our survey unveiled a two-phase assembly of the tip growth machinery: an initiation phase, during which the RHID is positioned and predefined, followed by the tip growth phase. Consistent with previous reports [4, 5], mCitrine-labeled (mCit) GTPase ROP2 associated with the RHID during the early initiation phase (stage -4), well before any detectable cell bulging (**Figure 1B-D, Figure S1**). The timing of mCit-ROP2 polarization appeared to be largely independent from promoter activity, as analysis of an estradiol-inducible reporter construct for mCit-ROP2 resulted in a similar timing of RHID association, albeit with higher cytosolic signal (**Figure 1D, Figure S1**). Closely related GTPases ROP4 and ROP6 exhibited a similar timing as ROP2, when inducibly expressed (**Figure 1D, Figure S1**), which indicates that ROP GTPases are prepositioned well before growth is activated. The only tested ROP-effectors that were found to polarize after ROPs but still during the initiation phase, were the PHOSPHATIDYLINOSITOL-4-PHOSPHATE 5-KINASE 3 (PIP5K3) [20], and the NADPH oxidase ROOT HAIR DEFECTIVE 2 (RHD2) [21, 22]. PIP5K3 and RHD2 have both been shown to directly interact with ROPs during tip growth regulation [18, 23] and modulate vesicle traffic to the plasma membrane and cell wall rigidity, respectively [24].

The second phase of root hair emergence was characterized by the polarization of known ROP effectors such as (ARPC2), a subunit of the actin polymerizing ARP2/3 complex [25], and the markers for vesicle transport and secretion SYP123 and SEC3A [26-28] (**Figure 1D, Figure S1**). The onset of bulge formation was further characterized by local elevation of cytosolic Ca^{2+} as reported by the sensor RGECO-1 [29, 30], as well as the formation of $PI(4,5)P_2$ at the emerging apex as indicated by the reporter P15Y [31]. The receptor-like kinase FERONIA, which activates ROPs during root hair tip growth [32], accumulated after bulging in stage +2 (**Figure 1D, Figure S1**). Even though our visualization of total ROP distribution does not provide information on the ROP activity state, the temporal separation of ROP positioning and growth activation led us to

90 hypothesize that distinct upstream, phase-specific regulators drive ROP GTPase recruitment and
91 activation.
92

RopGEFs exhibit specific spatiotemporal patterns of expression and subcellular
localization

ROPs have been extensively studied during cell elongation [33], stomata regulation [34],
pavement cell development [35, 36], and in regulation of the two tip-growing plant cell types, pollen
tubes [37] and root hairs [4, 5]. As molecular "switches", the activity-state of GTPases depends on
regulators that "toggle" the switch between active or inactive [38, 39]. GTPase-activating proteins
(GAPs) accelerate hydrolysis of a bound GTP to GDP, rendering the GTPase inactive. Examples
for GAPs with described roles in pollen tube growth are RopGAP1 [40] and REN1 [14]. To switch
the GTPase into an active state, guanyl-nucleotide exchange factors (GEFs) facilitate GDP release
from the GTPase and the binding of another GTP molecule, and thereby promote interactions with
downstream effectors [39, 41]. RopGEFs of the PRONE family [42, 43] and SPK1, the only DOCK
family GEF [44], play known roles in tip growth regulation in pollen tubes [17, 45, 46] and leaf
trichomes[44, 47]. However, temporal relationships between ROPs and RopGEFs have not yet
been established in these cell types and much less is known about RopGEF functions in root hairs.
To identify candidate ROP regulators that play potential roles in root hair initiation and growth
regulation we mined cell-type-specific gene expression data [48-50] and compared expression
profiles of RopGEFs in root trichoblasts and atrichoblasts. Among these RopGEFs (hereafter
abbreviated as GEFs), the expression of GEF3 and GEF4 showed the strongest root hair-
specificity, followed by GEF12, GEF11, GEF14, and GEF10 in descending order (**Figure S2A**).
Based on their trichoblast-specific expression, these GEFs were selected for subsequent analyses.
To determine in which stage of hair cell maturation the candidate GEFs play their roles, we
analyzed their expression patterns in growing roots expressing mCit fusions under the control of
the respective native promoters. We observed a diverse distribution of expression patterns among
the remaining GEFs, with a surprisingly unique pattern for each gene (**Figure 2A**). We further
analyzed the timing of polarization at the RHID for each mCit-labeled GEF and compared it to the
timing of mCit-ROP2. We found that mCit-GEF3 and mCit-GEF14 reached significant polarization
consistently one cell stage earlier than mCit-ROP2 (**Figure 2B, Figure S1**). A major difference
between both expression patterns was that GEF14 expression lasted from the meristematic zone
(MZ) to the elongation zone (EZ), whereas GEF3 expression was first observed in the root transition
zone (TZ, in-between MZ and EZ) and lasted beyond hair emergence (**Figure 2A**). The beginning
of gene expression of GEF4 and GEF12 was detected first in EZ and the polarization of both
proteins at the RHID coincided with the onset of bulge formation (**Figure 2A, B, Figure S1**). This
specific expression pattern and the timing of polarization of both, GEF4 and GEF12, indicate similar
functions during a defined time window in differentiating trichoblasts. This is distinctly different
compared to GEF3 and GEF14, as these GEFs both polarize at a significantly earlier timepoint and
show a more defined and restricted localization to the RHID.

In addition to the specific expression pattern of GEF3, its localization stood out in comparison to all other GEFs, as mCit-GEF3 appeared highly polarized at the cell periphery of the RHID, soon after the first fluorescence signal became detectable (**Figure 2A-C**). Polarization of mCit-GEF14 occurred less rapidly but reached a similar degree as mCit-GEF3 at stage +1 (**Figure 2B,D, Figure S1),** while mCit-GEF4 and mCit-GEF12, exhibited a lower degree of polarization and association with the plasma membrane (**Figure 2B, E, F, Figure S1**). mCit-GEF3 appeared highly concentrated in a circular polar domain, being barely detectable outside this domain (polarity index at cell "+1": 14.4 ± 2.2) (**Figure 2C, Figure S1**). We tested whether the timing of GEF polarization was dependent on the timing of gene expression and analyzed the polarity indices of mCit-GEF3 and mCit-GEF4 under the control of estradiol-inducible promoters. In both cases, the earlier gene expression did not result in an earlier RHID polarization of the protein (**Figure S1**), but rather delayed the time point of significant polarization of mCit-GEF4, likely due to higher cytosolic levels that affected the polarity index calculation. Its independence from gene expression indicates that the timing of polarization is an intrinsic feature that may be controlled by further partners of unknown identity.

To determine the functional role of the investigated GEFs we analyzed T-DNA insertion lines for phenotypes in root hair initiation and development. Root hair density was used as a measure for the general ability to develop root hairs, distance of the first bulge from the root tip was used to infer delays in the timing of root hair initiation and root hair length was used as measure for regulation of tip growth. Only in *gef3* mutant alleles, root hair density was reduced and root hair initiation was delayed, (**Figure 2G-I**), while hair length seemed only slightly affected (**Figure S2E**). In *gef3-1*, which lacks almost all full length transcript **(Figure S2C)**, the observed phenotypes could be rescued with a GEF3::mCit-GEF3 construct (**Figure 2H, I**), further verifying that loss of GEF3 function causes the phenotypes and confirming the functionality of the fusion construct. Furthermore, other mutant alleles of *gef3* showed a similar delay in root hair initiation (**Figure S2B,D**). None of the *gef3* T-DNA insertion lines lacked root hairs entirely, pointing towards functional redundancy among GEFs or compensatory mechanisms. A mutant line carrying both *gef3-1* and *rop2-1* alleles did not exhibit a significant increase in phenotype strength (**Figure 2G-I**), which indicates that such compensation involves other ROP-GEF combinations present in trichoblasts. Except for *gef3-1* no other *gef* single mutants, including *gef4-*2, exhibited any significant negative effects on bulge distance or root hair density (**Figure 2G-I**), but *gef4-2* lines did show reduced hair length (**Figure S2E**). A combination of *gef3-1* and *gef4-2* exhibited similar root hair density in *gef3-1*, but showed a small increase in distance of the first bulge from the root tip, i.e. stronger delay in hair initiation (**Figure 2G-I**). This phenotypic analysis suggests different, yet overlapping roles for GEF3 and GEF4, where GEF3 is mainly, but not exclusively, responsible for root hair initiation and bulging, while GEF4 is regulating root hair tip growth.

GEF3 is required for efficient root hair emergence and polar recruitment of ROP2

We have shown that GEF3 accumulated very specifically at the RHID, preceded ROP2 accumulation, and was required for proper root hair development. Therefore, GEF3 was selected as a promising potential regulator of ROP dependent root hair initiation.

To determine the temporal relationship of GEF3 and ROP2 recruitment to the RHID in detail, we observed the localization of both proteins at higher resolution over time. We employed a new microfluidic device, termed RootChip-8S, for long-term imaging of growing *Arabidopsis* roots. The RootChip-8S, is an adapted version of previously published RootChip designs [51-53] and consists of eight separate observation chambers that can be individually perfused to provide stable growth conditions and restrict root movement in Z to prevent focal drift (**Figure S3A, B**). The first detectable signal of GEF3::mCit-GEF3 was observed in the transition zone of the root, shortly after the last cell division (**Figures 2 A and 3 A**). Time-lapse imaging showed early targeting of mCit-GEF3 to the rootward cell pole (**Figure 3 A**). As the protein gradually enriched at the RHID over approximately 30 min (**Figure 3 B**), its localization at the rootward cell pole vanished (**Figure 3 A**). The GEF3 domain eventually stabilized at a consistent diameter of 9.6 μm (± 2.5 μm; *n*=17; FWHM), with its center 9.0 μm (± 2.2 μm) from the rootward cell pole (**Figure 3 B, Figure S3C**). To validate the earlier RHID accumulation of GEF3 compared to ROP2 through simultaneous visualization in the same cells, we imaged their localization in growing roots co-expressing fluorescently labeled versions of both proteins and found that mCit-GEF3 accumulated at the RHID one stage earlier than mTurquoise2-ROP2 (mTq2-ROP2) (**Figure 3 C**).

To test if GEF3 directly influences ROP2 recruitment to the RHID, we expressed mCit-ROP2 in the *gef3-1* T-DNA insertion mutant. Strikingly, we did not observe any significant polarization of mCit-ROP2 in the *gef3-1* mutant (**Figure 3 D, E**). Some cells in *gef3-1* were still able to produce bulges, but these bulges lacked polar ROP2 accumulation (**Figure 3 D, E**). Interestingly, the general membrane association of mCit-ROP2 was unaffected, as the localization appeared similar to that in other cell types. This shows that GEF3 function is required for ROP2 recruitment specifically to the RHID. We thus hypothesized that the observed root hair phenotypes in *gef3* mutants could be explained by failed polar recruitment and activation of downstream ROPs. Consequently, ROP loss-of-function mutants would exhibit similar phenotypes as *gef3*. We combined two previously described T-DNA lines for ROP2 and ROP4 [35, 54], and analyzed single and double mutants for phenotypes in root hair initiation. To account for the different genetic backgrounds of both single mutants (*rop2-1*, Col-0; *rop4-1*, WS), we transformed the double mutant *rop2-1/rop4-1* with a ROP2p::mTq2-ROP2 construct, and used the generated line as control for phenotype analysis. For both traits, root hair density and distance of the first bulge to the root tip, the double mutant *rop2-1/rop4-1* showed a phenotype that exceeded the strength of phenotypes in *gef3-1* (**Figure 3 F - H, Figure S2F,G**). The single mutant lines *rop2-1 and rop4-1* showed, however, no significant effects and our ROP2p::mTq2-ROP2 construct was able to fully revert phenotypes in *rop2-1/rop4-1*, demonstrating functional redundancy among both ROPs. In contrast to the loss of ROP2 polarity in *gef3-1* mutants, polar mCit-GEF3 domains were still observed in *rop2-1/rop4-1* mutants (**Figure 3 I, J**). mCit-GEF3 polarity was unaffected in the early elongation

zone (**Figure 3 I, J; stage -4**), but was slightly reduced in later stages (**Figure 3 I, J; stages -1,**
+1) compared to mCit-GEF3 polarity in Col-0 background (**Figure S1**). This indicates that initial
GEF3 polarization is largely ROP independent, while ROPs have a positive effect on GEF3 polarity
in later stages of root hair development. The question remained how GEF3 fulfills its function as a
factor directing polarization of ROP2. Previous studies have demonstrated that GEF-catalyzed
nucleotide exchange of ROPs involves the formation of a stable G protein-GEF complex [55-58].
To test if GEF3 was capable of recruiting ROP2 through physical interaction, we used a ratiometric
bimolecular fluorescence complementation (rBiFC) assay with GEF3 fused to a C-terminal
fragment of YFP (cYFP) and either an N-terminal fragment of YFP (nYFP)-carrying ROP2 (wild-
type), a constitutively active CA-ROP2 (G14V), a dominant-negative DN-ROP2 (D120A), or a
truncated *rop2ΔN* (Δ1-79) that lacks its interaction domain (**Figure S3D**) [5, 59], co-expressed
transiently in *Nicotiana benthamiana* leaf epidermis cells [60]. All GEF3-cYFP/nYFP-ROP2
combinations, except for *rop2ΔN*, resulted in detectable YFP fluorescence (**Figure 3 K, L, Figure**
S3H-J). This finding was corroborated by a split-ubiquitin assay in yeast [61], testing the same
putative interactor combinations. Except for *rop2ΔN*, all forms of ROP2 yielded yeast growth,
indicating physical interaction with GEF3 (**Figure S3E-G**). Consistent with the hypothesis that
physical interaction with GEF3 is required for polar recruitment of ROP2 to the RHID, mCit-*rop2ΔN*
expressed in *Arabidopsis* trichoblasts did not assume polar localization at the RHID (stage +1,
Figure 3 M, N). Our interaction assays show further that GEF3 physically interacts with the active
as well as the inactive form of ROP2 via its interaction domain. While physical interaction between
GEF3 and CA-ROP2 was unexpected at first, specific binding of certain RopGEFs to both, GDP-
and GTP-bound ROPs has been demonstrated before [43].

Taken together, these results show that GEF3 is necessary for ROP polarization by recruiting
ROP2 to the RHID via direct physical interaction. Furthermore, the presence of ROP2 and ROP4
is not needed for GEF3 polarization, but enhances GEF3 polarity.

GEF3 defines the RHID and guides ROP polarization

To investigate whether GEF3 is not only necessary but also sufficient for ROP2 polarization, we
studied transgenic lines, inducibly overexpressing mCit-GEF3 (mCit-GEF3ox) in Col-0 or in
ROP2::mTq2-ROP2 background. Upon induction of mCit-GEF3 expression we observed that
additional patch-like GEF3 domains formed in trichoblasts and, surprisingly, also in atrichoblasts
(**Figure 4 A**). We frequently found mCit-GEF3 accumulation at the shootward sides of cells in the
meristem or the early elongation zone (**Figure 4 A**). Importantly, all observed normal and ectopic
mCit-GEF3 accumulations resulted in the simultaneous recruitment of mTq2-ROP2 (**Figure 4 A**).

We took advantage of the ubiquitous overexpression of mCit-GEF3 and determined its
localization in cells outside the root. In epidermal cells of the hypocotyl, we occasionally observed
occurrences of ectopic GEF3 accumulation into domains that strikingly resembled those of RHIDs
in trichoblasts (**Figure 4 B, Figure S4A**). Consistently, these ectopic mCit-GEF3 domains also lead
to recruitment of mTq2-ROP2, which never occurred in hypocotyl cells under control conditions

(**Figure 4 B**). Except for trichoblasts, mCit-GEF3ox did not lead to bulge formation in other cells, neither did induced expression of mCit-GEF3ox influence the timing of bulging (**Figure S4B**), indicating that ROP polarization alone was not sufficient to trigger outgrowth and further trichoblast-specific factors are required.

Since the analysis of gene expression, protein localization and loss-of-function phenotypes all suggested a differential timing of GEF3 and GEF4 function, we hypothesized that gain-of-function phenotypes should also reflect their specific roles in root hair initiation and growth. We therefore compared the phenotypic effects of mCit-GEF3ox to mCit-GEF4ox, which is known to induce root hair branching in trichoblasts [62]. In mCit-GEF4ox lines, bulges with branched tips, which failed to form long tube-like hairs, occurred in 30.5% of the observed cells (**Figure 4 C, D**). mCit-GEF3ox roots also displayed root hairs with branched tips in 20.6% of all trichoblasts, resembling mCit-GEF4 overexpression (**Figure 4 C, D**). However, mCit-GEF3ox further led to the formation of additional bulges, with 5.5% of trichoblasts showing multiple unbranched bulges and 21.4% of the cells showing multiple branched bulges within one cell (**Figure 4 C, D**). This phenotype bears striking similarities to *scn1* mutants that lack a ROP-inhibiting RhoGDI [11], which further points towards a gain of ROP activity in mCit-GEF3ox roots at an early stage during RHID positioning. We observed mCit-GEF3 accumulations in tips of branched root hairs and mTq2-ROP2 was generally present in the same locations (**Figure 4 E**), thus confirming that mCit-GEF3ox was also responsible for the formation of additional polar domains in branching hairs. These differential effects of GEF3 and GEF4 overexpression resemble the differential timing of their polar localization and further support the hypothesis that both proteins serve specific functions during different phases of root hair development, with GEF3 being decisive for the early formation of the RHID and ROP2 recruitment, and GEF4 being involved in downstream activation of ROP2 to trigger growth.

GEFs serve as positional cues during cell polarization

To our knowledge, GEF3 is not only the first RopGEF member reported to temporally precede ROP polarization during tip growth initiation, but it is also the first example for a plant protein that, when overexpressed, is capable of inducing the ectopic formation of polar, ROP-recruiting domains. Our finding, that GEF3 acts upstream of ROP2, challenges the idea of ROP self-organization during the initiation of cell polarization in trichoblasts [63]. A self-organizing reaction-diffusion mechanism, involving the interplay of ROP-activating RopGEFs and ROP-inactivating RopGAPs has been found to underlie the regular cell wall patterning in differentiating metaxylem vessels [64, 65]. In contrast to the multiple, regularly spaced cell wall pits of the metaxylem, the dependable positioning and establishment of a single polar RHID in trichoblast is likely to require a mechanism that eliminates randomness during its formation. Our results suggest that such non-random polarization can be achieved through a GEF3-mediated recruitment and temporally controlled ROP activation by differentially timed GEFs at a predefined location.

Evidence for a GTPase-positioning function of GEFs can also be found in other organisms. Recent work shows that during cellularization in Drosophila embryos, the GEF Dizzy and the

heteromeric GEF complex ELMO-Sponge are required for polarization of the GTPase Rap1 [41, 66]. It remains, however, challenging to establish the molecular sequence of GEF and GTPase functions in the context of a developing embryo. In budding yeast, Cdc42 is activated by its GEF Cdc24 during both budding and shmoo formation [67-69]. To initiate budding in yeast, the GEF Bud5 is recruited to the edge of the previous cell-division site by the landmark Axl2 [70]. Stabilized by scaffold proteins, Bud5 and the GEF Cdc24 then locally activate the GTPases Rsr4 and Cdc42 [71-76].

Using a GEF-scaffold complex as landmark for the targeted deployment of the growth machinery may therefore represent a common design principle among eukaryotic cells to prevent spontaneous and random polarization and, instead, drive growth in a directed manner. Yet, the fact that RopGEFs share no conserved features with animal or fungal GEFs [42], indicates that the problem of robust cell morphogenesis has been solved several times during evolution to account for specific differences in cellular growth mechanisms, organ development and lifestyle.

ACKNOWLEDGMENTS

We are grateful to Dominique Bergmann (Stanford) and Sheila McCormick (UC Berkeley) for detailed comments on our manuscript. We thank Karin Schumacher (U. Heidelberg) and Thomas Dresselhaus (U. Regensburg) for helpful advice, all members of the Grossmann lab and our colleagues at COS Heidelberg for inspiring discussions and their active support. We thank Jan Lohmann (Heidelberg), Markus Grebe (Potsdam), Masa Sato (Kyoto), Tijs Ketelaar (Wageningen), Kateřina Schwarzerová (Prague), Yvon Jaillais (Lyon), and Ueli Grossniklaus (Zurich) for sharing materials and Karin Schumacher and Thomas Holstein (both Heidelberg) for access to microscopes. This work was supported by research group funds from the excellence cluster CellNetworks to G.G., grants from the Deutsche Forschungsgemeinschaft (GR 4559/3-1 to G.G. and EV 198/1-1 to J.F.E.), seed funding through the SFB1101, an Emmy Noether fellowship (GR 4251/1-1) to C.G., a PhD-student grant from the Carl-Zeiss-Stiftung to D.G.M., and financial support by ETH Zürich to C.E.S.

AUTHOR CONTRIBUTIONS

The study was conceived by P.D. and G.G.; P.D., A.R., V.A.F.S., D.G.M., L.Y.A., and N.F.K. performed experiments; P.D., A.R., V.A.F.S., D.G.M., L.Y.A., C.G. and G.G. analyzed data. C.E.S. and G.G. designed the microfluidic device. J.F.E. designed the spinning disk microscope and gave advice about the imaging procedure. P.D., A.R. and G.G. prepared figures and wrote the manuscript with input from all coauthors. All authors read and approved the final version of the manuscript.

DECLARATION OF INTERESTS

327 The authors declare no competing interests.

FIGURES AND FIGURE LEGENDS

Figure 1. Timeline analysis of protein accumulation at the RHID.

(**A**) Model of root hair development in an epidermal cell file. During the initiation phase the root hair initiation domain (RHID) emerges in elongating trichoblasts and the tip growth machinery is assembled using an unknown mechanism. Stages -1/+1 mark the stages before and after the onset of bulging followed by the transition into polar growth. (**B**) Comparison of mCit-ROP2 and GFP-

LTI6B protein localization at the RHID in different stages of root hair development. For mCit-ROP2, cross sections (c. s.) and surface views (s. v.) of the same cells are shown. Intensities are represented in false-color, covering the full range of measured values within each data set. Scale bar, 10 μm. (**C**) Polarity index values (signal intensity in/out of RHID; $\log_2$ scale) for mCit-ROP2 and GFP-LTI6B along the different root hair stages. Error bars indicate SEM (n(stage - 1)=11(ROP2); 14(LTI6B)). The black triangle indicates the first stage with a significant difference between mCit-ROP2 and the evenly distributed GFP-LTI6B ($p < 0.01$). (D) Protein accumulation at the RHID throughout the early development of root hairs in 13 different marker lines. Genes are sorted by the timing of accumulation within their group of protein function. Color coding represents the polarity index for each developmental step. (n.d.: no data). For each stage average polarity index values from multiple cells are shown (see F**igure S1** for details). Stages -7 to -1 represent cells in the initiation phase before bulging, +1 to +3 cells after bulging. Solid vertical lines indicate the first stage with a significant difference between the respective marker and GFP-LTI6B. Listed genes were fused to either eGFP or mCitrine (mCit), under control of their own promoters, except for UBQ10::P15Y (PI(4,5)P_2 reporter), 35S::LTI6B and genes under control of an estradiol inducible promoter (indicated by an asterisk). See also **Figure S1**.

Figure 2. Functional analysis of RopGEFs in the root epidermis identifies GEF3 to be required for efficient root hair emergence.

(**A**) Roots expressing mCit fusion proteins (yellow) under their respective promoters, stained with propidium iodide (magenta) to visualize root and cell outlines. Top panel: Maximum intensity projection of the root tip. Scale bar, 200 μm. lower Panel: Optical cross section through the root in the elongation zone to visualize the tissue-specific expression. Scale bar, 20 μm. (**B**) Timeline analysis of protein accumulation at the RHID for selected GEFs in comparison to ROP2. GEFs are sorted by timing of accumulation at the RHID. Polarity index measurement and calculation as in **Figure 1**. (**C-F**) Subcellular localization (intensity-coded representation) of mCit-GEF3, mCit-GEF4, mCit-GEF14 and mCit-GEF12, in one cell before (-1) and after bulging (+1), respectively. Confocal cross sections (top) and surface views (bottom) of the same cells are shown. Scale bars, 20μm. (**G**) Representative root tips of WT (Col-0), and *gef* single and double mutant lines. Arrowheads indicate the first bulge. (**H**, **I**) Quantification of root hair density (**H**) and distance (**I**) of first bulge to the root tip for selected single and double mutants. Values were normalized to corresponding Col-0 values for each independent experiment. Box-plots: number of measurements (n); center lines show medians; notches of boxes represent the 95% confidence interval of the medians; box limits indicate 25th and 75th percentiles as determined by R software; whiskers extend 1.5 times the interquartile range from the 25th and 75th percentiles, dots represent outliers; crosses represent sample means; grey bars indicate 95% confidence intervals of the means. Letters above box-plots show results of an ANOVA-Test (significance value *p*=0.01), with same letters indicating no significant differences. See also **Figure S2**.

Figure 3. GEF3 is necessary for polar ROP2 recruitment during root hair initiation.

(**A**) Time-lapse imaging of GEF3::mCit-GEF3 in the RootChip-8S (see **Figure S2**). A root hair cell file in the early elongation zone is shown. Multiple stages of protein accumulation can be seen in each still image. The time series was registered to compensate for cell movement due to root growth. Numbers indicate time (mm:ss). Scale bar, 20µm. (**B**) Kymograph of intensity distribution (horizontal axis) along the outer cell flank (left-right orientation) between arrow heads in (**A**) over time (vertical axis). Note the gradual expansion of the emerging mCit-GEF3 domain until approximately 30 min and a stable domain size thereafter. (**C**) Colocalization and intensity profiles of relative signal distribution for mCit-GEF3 and mTq2-ROP2 along root hair cell files. Top: example root hair cell file expressing ROP2::mTq2-ROP2 (cyan) and GEF3::mCit-GEF3 (yellow). Cells are labeled from youngest cells (-6) to oldest cell (-1) before bulging. Bottom: average profiles of relative mTq2-ROP2 (cyan) and mCit-GEF3 (yellow) intensities along the outer cell periphery (line width 3 px) of 6 root hair cell files (arrowhead indicates direction of line measurement). To compensate for intensity differences between measurements, the signal intensities of each construct are normalized to the value at position 0 µm of each cell and afterwards averaged. (**D**) Subcellular localization and (**E**) quantification of mCit-ROP2 polarity at the RHID in Col-0 and the *gef3-1* mutant in root hair cells one cell before (-1) and after bulging (+1). Scale bar, 20µm. (**F**) Representative root tips of *rop2-1*, *rop4-1* and *rop2-1/rop4-1* mutant lines. Arrowheads indicate the first bulge. (**G**, **H**) Quantification of root hair density and distance of first bulge to the root tip in ROP single and double mutants, as well as in *rop2-1/rop4-1* carrying a ROP2::mCit-ROP2 construct. Values are normalized to corresponding Col-0 values for each independent experiment (for comparison, data for Col-0 shown in Figure 2H,I are re-plotted). (**I**) GEF3::mCit-GEF3 expression and localization in *rop2-1/rop4-1* mutant background. Upper panel: overview of the elongation zone. Lower panel: subcellular localization of mCit-GEF3 in cells that correspond to stages -4, -1, and +1. Due to the lack of bulges in this mutant, the reference point used in other lines is missing here and stage designations are therefore estimates. (**J**) Quantification of the polarity index of GEF3::mCit-GEF3 in *rop2-1/rop4-1* mutant background. Stages are chosen as indicated in (**I**). (**K**) Representative images of an rBiFC assay in tobacco epidermal cells for each combination tested. Positive YFP signal indicates protein-protein interaction, RFP signal in the same cells serves as expression control (see Materials and Methods and **Figure S2** for details). All images were recorded using identical conditions and settings, correction for brightness and contrast during image processing was kept equal except for the exemplary YFP image of *rop2ΔN*, which was set brighter to visualize the very low background signal. (**L**) Quantification of ratiometric BiFC experiments. YFP signals were normalized to the corresponding RFP signal to allow comparison of interaction strength. (**M**) Confocal image of root hair cells expressing estradiol inducible mCit-*ROP2* or mCit-*rop2ΔN*. Scale bar, 10µm. (**N**) Quantification of polarity index of the constructs shown in (**N**) and compared to the non-polar membrane marker GFP-LTI6B and free mCitrine. For a detailed explanation of all shown box-plots, see the description given in the legend of Figure 2. See also **Figure S3**.

Figure 4. GEF3 initiates the formation of polar growth domains

(**A**) Localization of *ROP2*::mTq2-ROP2 (cyan) in roots overexpressing mCit-GEF3 (yellow). Top: representative cell files of trichoblasts and atrichoblasts, both show ectopic accumulations of both markers. Bottom: intensity profiles along cell periphery of cells shown on the left. Dashed lines indicate cell borders. (**B**) Subcellular localization of *ROP2*::mTq2-ROP2 (cyan) in hypocotyl cells ectopically expressing mCit-GEF3 (yellow). To visualize the background signal in the mCit channel of the mock-treated seedling, intensities were 10-fold enhanced during image processing as compared to the image of estradiol-treated specimen. Other channels are scaled identically. Scale bar, 20 µm. (**C**) Top panels: Representative images of Col-0 (left) and roots overexpressing mCit-*GEF4* (middle) and mCit-*GEF3* (right) (24h after estradiol induction). Inserts show an idealized sketch of the predominant root hair shape. Bottom panels: Higher magnification views of the root

zones indicated as C' and C'' in the top panels of GEF4ox and GEF3ox, respectively. Scale bars,

100 µm. (**D**) Frequency of phenotypes (induction of multiple hairs, branched bulges, and combined

phenotypes) induced by overexpression of lines shown in (**C**) in % of total observed hairs. Bottom

numbers indicate *n*. (**E**) Colocalization of mCit-GEF3 (yellow) and ROP2::mTq2-ROP2 (cyan) in

branched root hairs, 24 hours after estradiol induction of mCit-GEF3 overexpression. Scale bar,

20 µm. See also **Figure S4**.

MATERIALS AND METHODS

Plant material

Arabidopsis thaliana ecotype Col-0 was used as wildtype in this study. The mutant lines *rop2-1* (SALK_055328C), *gef3-1* (SALK_079879C), *gef3-2* (SALK_079885), *gef3-3* (SAIL_294_A09), *gef3-4* (SALK_021751), *gef3-5* (SALK_046978C), *gef4-2* (SALK_107520) *gef10-1* (SALK_009456), *gef11-1* (SALK_126725C), *gef14-2* (SALK_046067) were obtained from NASC (Nottingham Arabidopsis Stock Centre http://arabidopsis.info/). The *rop4-1* mutant [36] was kindly provided by Markus Grebe (Potsdam). Double mutants were made by crossing and selection for double homozygous plants by PCR (Primers used for genotyping, see Table S3). The GFP-SYP123 and GFP-SYP132 lines [26] were kindly provided by Masa H. Sato (Kyoto), SEC3A-GFP lines [27] were kindly provided by Tijs Ketelaar (Wageningen), the GFP-ARPC2A line [25] was kindly provided by Kateřina Schwarzerová (Prague), and the PI(4,5)P$_2$-reporter line (P15Y) [31] was kindly provided by Yvon Jaillais (Lyon). To generate transgenic Col-0 lines expressing GFP-tagged FER, Col-0 was transformed with pFER::FER-GFP containing pMDC111 [77], kindly provided by Ueli Grossniklaus (Zurich).

Information on all genes referenced in this work, including mutant alleles and sources is provided in **Supplementary Data S1**, **Table S1**.

RNA extraction and expression analysis

Seedlings were grown on mesh on a ¼ MS plate. Root tissue was harvested 10 dag, immediately frozen in liquid nitrogen and ground in a tissue lyser (Qiagen) with metal beads. Total RNA was extracted using the RNeasy Plant Mini Kit (Qiagen). After DNAse digest (Thermo Fischer Scientific), 0.5 µg total RNA was used for cDNA synthesis with oligo(dT) primers (RevertAid First Strand cDNA Synthesis Kit, Thermo Fischer Schientific). To test for the presence of Gef3 mRNA, 2µl of this cDNA was used for PCR amplification (40 cycles). The primers used for the expression analysis are listed in Supplementary Information, **Table S2**.

Data mining of published cell type-specific gene expression was performed using Genevestigator [78].

Growth conditions and inducible expression

Plants were grown under long day conditions (16 h light) at 21°C. Seedlings used for experiments were grown for seven days on ¼ MS medium (pH 5.7) without sucrose, supplied with 0.8% plant agar (Duchefa). Transgenic plants transformed with an estradiol inducible construct were induced by covering with cellulose tissues soaked with ¼ MS medium containing 20 µM Estradiol (Stock 20 mM in Ethanol) and 0.01% Silwet L77 for 15 minutes. The time after induction differed between the individual expression constructs and ranged from 3 – 12 h to get sufficient signal for imaging. In

471 general, for imaging of protein localization using inducible expression constructs, time after
472 induction was as low as possible to avoid overexpression artefacts.
473

Plant transformation and selection

The agrobacterium strain GV3101::pMP90RK pre-transformed with the helper plasmid pSoup was
used for all transformations of *Arabidopsis thaliana*. Plants were transformed using floral dipping
and transgenic T1 plants were recovered after the appropriate selection. Selection of constructs
containing mTurquoise2 (mTq2) was done on plates containing 50 µg/ml Kanamycin. T1 Selection
of constructs containing mCitrine (mCit) was done on soil spraying 3x within 2 weeks (7 – 21 day
old seedlings) with Basta®-Solution containing 200 µg/ml Glufosinate-ammonium and 0.05%
Tween20. Further generations were selected on plates containing Basta®-Solution to a
concentration of 7.5 µg/ml Glufosinate-ammonium. At least 10 independent transgenic lines per
expression construct were recovered and checked for fluorescence. Lines that had clear
fluorescence and limited phenotypic anomalies were confirmed by PCR and used for further study.
485

Molecular cloning

All cloning for expression vectors used in Arabidopsis was performed using the GreenGate system
[79] and base vector pGGZ003 was used for all expression constructs. A List of all expression
constructs including module composition and tagging site is provided in Supplementary
Information, **Table S3**. Where internal BsaI (Eco31I) sites were present, they were mutated fusing
two PCR fragments, with one mutating the internal BsaI site by an altered primer sequence at this
site. After cloning the individual modules into the entry vectors, the complete fragment was checked
by sequencing. To confirm the correct assembly of modules into pGGZ003, the presence and
orientation was checked by sequencing. As fluorophores, mCitrine (mCit) [80] and mTurqoise2
(mTq2) [81] were fused to the gene of interest separated by a short GAGAGA-Linker. mCit was
cloned from the Ca^{2+} indicator CerTN-L15, containing the common folding mutations V163A and
S175G [82], and made monomeric by introducing the A206K mutation. mTq2 was cloned from the
Addgene vector pLifeAct-mTurquoise2 (ID #36201). A List of all primers used for molecular cloning
is provided in Supplementary Information, **Table S2**.
For estradiol inducible expression constructs [83], the XVE gene was fused to the promoter (635
bp) and terminator of *UBI10* (AT4G05320). This expression cassette was cloned in reverse
orientation into the entry vector pGGA000 together with the oLexA minimal 35S promoter element.
Both elements were separated by a 560 bp spacer. This resulted in a single module that could be
used as a promoter module for GreenGate cloning.
For expression of genes under control of their native promoter, the following sequences upstream
of the start codon were cloned into pGGA000: *GEF3* -1743 bp; *GEF4* -1500 bp; *GEF10* -1534 bp;
GEF11 -1458 bp; *GEF12* -701 bp; *ROP2* -1503 bp; *PIP5K3* -1193 bp.

For the mbSUS and rBiFC analysis, constructs (listed in Supplementary Information, **Table S3**)
were generated using Gateway technology or the 2in1 cloning system (Grefen and Blatt 2012).
Coding sequences were PCR-amplified and inserted via 'BP' recombination into either pDONR207,
pDONR221-P1P4, or pDONR221-P3P2, respectively, and confirmed by sequencing. Resulting
'Entry constructs' were cloned via 'LR' recombination into Destination vectors for either mbSUS or
rBiFC [84].

Phenotyping of T-DNA insertion lines and inducible overexpression mutant lines

Primary roots (7 dag) were used for all phenotyping experiments. For phenotyping of T-DNA
insertion lines the distance of the first bulge to the root tip was defined. The first visible swelling of
the cell outline was defined as first bulge, even when this morphological change was very little. The
root hair density was analyzed in the next 2 mm. The root hair length was analyzed measuring root
hairs growing perpendicular to the root in a region 3-6 mm away from the root tip. For phenotyping
the overexpression lines, roots were induced by spraying with medium containing 20 µM Estradiol.
The position of the root tip was marked just after induction and normal growing roots were imaged
and analyzed 24-26 hours after induction (h. a. i.). A region approximately 2-4 mm from the root tip
was used for the analysis of the overexpression phenotype.

Propidium iodide (PI) staining

Primary roots (7 dag) were stained for 5 min in 10 µg/ml propidium iodide (PI; diluted in ¼ MS, from
1 mg/ml stock), washed in medium and imaged directly. As PI staining varies substantially between
different developmental zones of the root, we employed gamma correction (0.75) exclusively in the
PI channel of the images to equalize stronger and weaker stained regions in the final figures (see
image processing).

Ratiometric Bimolecular Fluorescence analyses (rBiFC)

Agrobacterium-mediated transient transformation of 4-6 week old *N. benthamiana* leaves with 2in1
rBiFC constructs was performed as described previously [85]. Fluorescence intensities were
recorded 36h post infiltration for approximately 30 images per construct as described in [60]. Ratios
between complemented YFP fluorescence and RFP were calculated and plotted using BoxplotR.
Immunoblotting verified protein expression utilizing anti-MYC conjugated to peroxidase antibodies
and anti-HA peroxidase conjugated antibodies, respectively.

Mating-based Split Ubiquitin Assay ('Cyto-SUS')

Bait/Cub fusion and preys/Nub fusions were generated by cloning the coding region for *GEF3* and
different *ROP2* versions into the Gateway-compatible vectors pMetOYC-Dest and pNX35-Dest,
respectively. These were transformed into haploid yeast strains THY.AP4 (Cub fusion) and

THY.AP5 (Nub fusions) as described previously [86]. After mating, diploid yeasts were dropped in

10 times OD dilutions on vector- (CSM-Leu-, Trp-, Ura) and interaction-selective media (CSM-Leu-

, Trp-, Ura-, Ade-, His-) supplemented with 50 or 500 µM methionine. The non-mutated N-terminal

ubiquitin moiety (NubI) was used as positive control, and NubI13G (empty vector) as negative

control [84]. Protein expression was verified in haploid yeast using antibodies against the VP16

domain for the Cub fusion or the HA epitope tag for the Nub fusions.

Microfluidic device (RootChip-8S) fabrication and on-chip plant cultivation

553 Microfluidic device fabrication and on-chip plant cultivation were performed as described in

554 [87, 88]. Briefly, Arabidopsis seedlings were grown in pipette tips filled with solidified medium and

555 interfaced with poly(dimethylsiloxane) (PDMS) RootChip-8S devices 4-5 days after germination

556 under sterile conditions. Media perfusion through the microfluidic channels was controlled using a

557 peristaltic pump (DNE GmbH; volumetric flow rate in each channel, 5 µl/min).

Bright-field and fluorescence imaging

Live-cell fluorescence imaging was performed on a custom-built spinning-disk confocal

microscope, consisting of a Nikon Ti-E stand, equipped with 20x multi-immersion (N.A. 0.75) and

60x water immersion (N.A. 1.2) objectives (Nikon), a motorized stage (Applied Scientific

Instrumentation, USA), a spinning disk with customized hole pattern (CREST Optics, Italy), a

motorized filter wheel (Cairn Research, UK), a laser launch (Omicron, Germany), and an EMCCD

camera (Photometrics, USA). Image acquisition was operated through Nikon NIS Elements

software or Micro-Manager [89]. mCitrine-labeled proteins were excited using a laser emitting at

the wavelength of 515 nm; mTq2-labeled proteins were excited at 440 nm. Bandpass filter

specifications were CWL (central wavelength) = 542 nm, BW (band width) = 20 nm for mCit

emission and CWL = 480 nm, BW = 40 for mTq2. Chlorophyll autofluorescence was recorded using

561 nm excitation and emission filter specifications of CWL = 630 nm, BW = 92. Root hair

phenotyping was performed by transmitted light microscopy on a Nikon SMZ18 stereo microscope,

equipped with SHR Plan Apo 0.5x (N.A. 0.075) and 2x (N.A. 0.3) objectives (Nikon) and an Orca

Flash 4.0 sCMOS camera (Hamamatsu, Japan). Imaging of the propidium iodide stained root tips

expressing different *GEF*::mCit-*GEF* fusions was performed on a Leica SP5 point scanning

confocal microscope with a 20x water immersion objective (N.A 0.7), using 1x Zoom, Pinhole 1AU,

1024x1024 pixel scanning field, 400 Hz scanning speed, 2x line average as settings. mCit was

excited with the 514 nm line of an Argon-Laser and emission light was collected between 520-550

nm. Propidium iodide was exited with a 561 nm laser and emission light was collected between

570-650 nm. Both channels were imaged in sequential scans to avoid crosstalk using Hybrid-

Detectors (Gain 200). Z-Stacks were acquired with 2 µm step size for 20 X objectives and 0.6 µm

for 60/63 X objectives. The top-view of the RHID was reconstructed from Z-Stacks using the built-

in function of FIJI (3D-Project with interpolation).

Image processing, data analysis and software

All image processing, image analysis and measurements was done using FIJI (ImageJ) [90], further data analysis was done in Microsoft Excel, if not mentioned otherwise. A gaussian filter (radius 0.8 px) and a rolling ball (radius 100 px) background subtraction was applied to all shown images. The "Royal" lookup table was used for images shown in intensity pseudo color. Kymographs were generated using the multi-kymograph plugin in FIJI. For images of propidium iodide (PI) staining for visualization of the root outlines, the gamma settings were adjusted to 0.75 to reduce the contrast between weakly and strongly stained regions of the root. The adjustment was only done in the PI channel. The signal of the mCitrine-fusions was not altered.

To obtain vertical root cross sections, Z-stacks (voxel size: 0.7568x0.7568x2.0142 μm^3) were processed using the Reslice function in FIJI. Maximum intensity projections of approximately 100 µm along the root axis, in a distance of approximately 700 to 1000 µm away from the root tip were generated.

Polarity indices were analyzed on the original unprocessed images by measuring in 3 by 15 pixel regions at the RHID, outside the RHID and outside the root, as a background value. Before bulge formation a region approximately 10 µm from the rootward cell edge was used as the RHID. The background value was subtracted from both regions and the value outside the RHID was divided by the value inside the RHID $[(Int_{in} - Int_{bkgd}) / Int_{out} - Int_{Bkgd})]$. This polarity index was calculated for each cell within a root hair file, all files were aligned to the first cell with a visible bulge and the average polarity index for each developmental step was calculated. A signal was defined as polar, if its value differed (p-value <0.05, according to two-tailed t-test with unequal variances) to the polarity index of the membrane marker GFP-LTI6B.

To measure GEF3 domain size, intensity profiles along the outer plasma membrane were generated from confocal cross sections of trichoblasts. Since gradually decreasing fluorescence signals impede an unambiguous determination of the outer boundary of polar accumulations, domain size was quantified by measuring the full signal width at half-maximal signal intensity (FWHM; **Figure S3**).

Box plots were created using BoxPlotR (http://shiny.chemgrid.org/boxplotr/), an R-based web-tool provided by the Tyers (Montreal) and Rappsilber (Edinburgh) labs. For all plots number of measurements (n) is given. Center lines show the medians; notches of boxes represent the 95% confidence interval of the medians; box limits indicate the 25th and 75th percentiles as determined by R software; whiskers extend 1.5 times the interquartile range from the 25th and 75th percentiles, outliers are represented by dots; crosses represent sample means; bars indicate 95% confidence intervals of the means. As statistical test, t-tests (two-tailed, unequal variances) were performed in Microsoft Excel, ANOVA tests were performed in R and the used script is provided in **Data S1**.

SUPPLEMENTAL INFORMATION

Table S1. List of identifiers for genes and single mutant alleles used in this study.

Table S2. List of PCR primers used for GreenGate and Gateway cloning.

Table S3. List of used expression vectors.

Data S1. R script for ANOVA/Tukey tests performed on mutant phenotype data.

REFERENCES

1.   Riquelme, M. (2013). Tip growth in filamentous fungi: a road trip to the apex. Annu Rev
631   Microbiol *67*, 587–609.

632   2.   Rounds, C. M., and Bezanilla, M. (2013). Growth Mechanisms in Tip-Growing Plant Cells.
633   Annu Rev Plant Biol.

634   3.   Russell, S. A., and Bashaw, G. J. (2017). Axon guidance pathways and the control of gene
635   expression.

636   4.   Molendijk, A. J., Bischoff, F., Rajendrakumar, C. S., Friml, J., Braun, M., Gilroy, S., and
637   Palme, K. (2001). Arabidopsis thaliana Rop GTPases are localized to tips of root hairs and
638   control polar growth. EMBO J *20*, 2779–2788.

639   5.   Jones, M. A., Shen, J.-J., Fu, Y., Li, H., Yang, Z., and Grierson, C. S. (2002). The
640   Arabidopsis Rop2 GTPase is a positive regulator of both root hair initiation and tip growth.
641   Plant Cell *14*, 763–776.

642   6.   Bi, E., and Park, H.-O. (2012). Cell polarization and cytokinesis in budding yeast. Genetics
643   *191*, 347–387.

644   7.   Ikeda, Y., Men, S., Fischer, U., Stepanova, A. N., Alonso, J. M., Ljung, K., and Grebe, M.
645   (2009). Local auxin biosynthesis modulates gradient-directed planar polarity in
646   Arabidopsis. Nat Cell Biol *11*, 731–738.

647   8.   Fischer, U., Ikeda, Y., Ljung, K., Serralbo, O., Singh, M., Heidstra, R., Palme, K., Scheres,
648   B., and Grebe, M. (2006). Vectorial information for Arabidopsis planar polarity is mediated
649   by combined AUX1, EIN2, and GNOM activity. Curr Biol *16*, 2143–2149.

650   9.   Kiefer, C. S., Claes, A. R., Nzayisenga, J.-C., Pietra, S., Stanislas, T., Hüser, A., Ikeda, Y.,
651   and Grebe, M. (2015). Arabidopsis AIP1-2 restricted by WER-mediated patterning
652   modulates planar polarity. Development *142*, 151–161.

653   10.   Fu, Y., Li, H., and Yang, Z. (2002). The ROP2 GTPase controls the formation of cortical
654   fine F-actin and the early phase of directional cell expansion during Arabidopsis
655   organogenesis. Plant Cell *14*, 777–794.

656   11.   Carol, R. J., Takeda, S., Linstead, P., Durrant, M. C., Kakesova, H., Derbyshire, P., Drea,
657   S., Zárský, V., and Dolan, L. (2005). A RhoGDP dissociation inhibitor spatially regulates
658   growth in root hair cells. Nature *438*, 1013–1016.

659   12.   Chang, F., Gu, Y., Ma, H., and Yang, Z. (2013). AtPRK2 Promotes ROP1 Activation via
660   RopGEFs in the Control of Polarized Pollen Tube Growth. Mol Plant *6*, 1187–1201.

661   13.   Hwang, J.-U., Gu, Y., Lee, Y.-J., and Yang, Z. (2005). Oscillatory ROP GTPase activation
662   leads the oscillatory polarized growth of pollen tubes. Mol Biol Cell *16*, 5385–5399.

663   14.   Hwang, J.-U., Vernoud, V., Szumlanski, A., Nielsen, E., and Yang, Z. (2008). A tip-
664   localized RhoGAP controls cell polarity by globally inhibiting Rho GTPase at the cell apex.
665   Curr Biol *18*, 1907–1916.

666   15.   Lin, Y., Wang, Y., Zhu, J. K., and Yang, Z. (1996). Localization of a Rho GTPase Implies a
667   Role in Tip Growth and Movement of the Generative Cell in Pollen Tubes. Plant Cell *8*,
668   293–303.

669   16.   Yalovsky, S., Bloch, D., Sorek, N., and Kost, B. (2008). Regulation of Membrane

670   Trafficking, Cytoskeleton Dynamics, and Cell Polarity by ROP/RAC GTPases. Plant
671   Physiol *147*, 1527–1543.

672 17.   Zhang, Y., and McCormick, S. (2007). A distinct mechanism regulating a pollen-specific
673   guanine nucleotide exchange factor for the small GTPase Rop in Arabidopsis thaliana.
674   Proc Natl Acad Sci USA *104*, 18830–18835.

675 18.   Kost, B., Lemichez, E., Spielhofer, P., Hong, Y., Tolias, K., Carpenter, C., and Chua, N. H.
676   (1999). Rac homologues and compartmentalized phosphatidylinositol 4, 5-bisphosphate
677   act in a common pathway to regulate polar pollen tube growth. J Cell Biol *145*, 317–330.

678 19.   Cutler, S. R., Ehrhardt, D. W., Griffitts, J. S., and Somerville, C. R. (2000). Random
679   GFP::cDNA fusions enable visualization of subcellular structures in cells of Arabidopsis at
680   a high frequency. Proc Natl Acad Sci USA *97*, 3718–3723.

681 20.   Kusano, H., Testerink, C., Vermeer, J. E. M., Tsuge, T., Shimada, H., Oka, A., Munnik, T.,
682   and Aoyama, T. (2008). The Arabidopsis Phosphatidylinositol Phosphate 5-Kinase PIP5K3
683   is a key regulator of root hair tip growth. Plant Cell *20*, 367–380.

684 21.   Foreman, J., Demidchik, V., Bothwell, J. H., Mylona, P., Miedema, H., Torres, M. A.,
685   Linstead, P., Costa, S., Brownlee, C., Jones, J. D., et al. (2003). Reactive oxygen species
686   produced by NADPH oxidase regulate plant cell growth. Nature *422*, 442–446.

687 22.   Takeda, S., Gapper, C., Kaya, H., Bell, E., Kuchitsu, K., and Dolan, L. (2008). Local
688   positive feedback regulation determines cell shape in root hair cells. Science *319*, 1241–
689   1244.

690 23.   Wong, H. L., Pinontoan, R., Hayashi, K., Tabata, R., Yaeno, T., Hasegawa, K., Kojima, C.,
691   Yoshioka, H., Iba, K., Kawasaki, T., et al. (2007). Regulation of Rice NADPH Oxidase by
692   Binding of Rac GTPase to Its N-Terminal Extension. Plant Cell *19*, 4022–4034.

693 24.   Mendrinna, A., and Persson, S. (2015). Root hair growth: it's a one way street.
694   F1000Prime Rep *7*, 23.

695 25.   Havelková, L., Nanda, G., Martinek, J., Bellinvia, E., Sikorová, L., Šlajcherová, K.,
696   Seifertová, D., Fischer, L., Fišerová, J., Petrásek, J., et al. (2015). Arp2/3 complex subunit
697   ARPC2 binds to microtubules. Plant Science *241*, 96–108.

698 26.   Ichikawa, M., Hirano, T., Enami, K., Fuselier, T., Kato, N., Kwon, C., Voigt, B., Schulze-
699   Lefert, P., Baluska, F., and Sato, M. H. (2014). Syntaxin of Plant Proteins SYP123 and
700   SYP132 Mediate Root Hair Tip Growth in Arabidopsis thaliana. Plant Cell Physiol *55*, 790–
701   800.

702 27.   Zhang, Y., Immink, R., Liu, C.-M., Emons, A.-M., and Ketelaar, T. (2013). The Arabidopsis
703   exocyst subunit SEC3A is essential for embryo development and accumulates in transient
704   puncta at the plasma membrane. New Phytol *199*, 74–88.

705 28.   Xing, S., Mehlhorn, D. G., Wallmeroth, N., Asseck, L. Y., Kar, R., Voss, A., Denninger, P.,
706   Schmidt, V. A. F., Schwarzländer, M., Stierhof, Y.-D., et al. (2017). Loss of GET pathway
707   orthologs in Arabidopsis thaliana causes root hair growth defects and affects SNARE
708   abundance. Proc Natl Acad Sci USA *114*, E1544–E1553.

709 29.   Zhao, Y., Araki, S., Wu, J., Teramoto, T., Chang, Y.-F., Nakano, M., Abdelfattah, A. S.,
710   Fujiwara, M., Ishihara, T., Nagai, T., et al. (2011). An expanded palette of genetically
711   encoded Ca^{2+} indicators. Science *333*, 1888–1891.

712 30.   Keinath, N. F., Waadt, R., Brugman, R., Schroeder, J. I., Grossmann, G., Schumacher, K.,
713   and Krebs, M. (2015). Live Cell Imaging with R-GECO1 Sheds Light on flg22- and Chitin-
714   Induced Transient $[Ca^{2+}]_{cyt}$ Patterns in Arabidopsis. Mol Plant *8*, 1188–1200.

715   31.   Simon, M. L. A., Platre, M. P., Assil, S., van Wijk, R., Chen, W. Y., Chory, J., Dreux, M.,
716   Munnik, T., and Jaillais, Y. (2014). A multi-colour/multi-affinity marker set to visualize
717   phosphoinositide dynamics in Arabidopsis. Plant J *77*, 322–337.

718   32.   Duan, Q., Kita, D., Li, C., Cheung, A. Y., and Wu, H.-M. (2010). FERONIA receptor-like
719   kinase regulates RHO GTPase signaling of root hair development. Proc Natl Acad Sci
720   USA *107*, 17821–17826.

721   33.   Lin, D., Nagawa, S., Chen, J., Cao, L., Chen, X., Xu, T., Li, H., Dhonukshe, P., Yamamuro,
722   C., Friml, J., et al. (2012). A ROP GTPase-dependent auxin signaling pathway regulates
723   the subcellular distribution of PIN2 in Arabidopsis roots. Curr Biol *22*, 1319–1325.

724   34.   Jeon, B. W., Hwang, J. U., Hwang, Y., Song, W. Y., Fu, Y., Gu, Y., Bao, F., Cho, D.,
725   Kwak, J. M., Yang, Z., et al. (2008). The Arabidopsis Small G Protein ROP2 Is Activated
726   by Light in Guard Cells and Inhibits Light-Induced Stomatal Opening. Plant Cell *20*, 75–87.

727   35.   Xu, T., Wen, M., Nagawa, S., Fu, Y., Chen, J.-G., Wu, M.-J., Perrot-Rechenmann, C.,
728   Friml, J., Jones, A. M., and Yang, Z. (2010). Cell surface- and rho GTPase-based auxin
729   signaling controls cellular interdigitation in Arabidopsis. Cell *143*, 99–110.

730   36.   Fu, Y., Gu, Y., Zheng, Z., Wasteneys, G., and Yang, Z. (2005). Arabidopsis interdigitating
731   cell growth requires two antagonistic pathways with opposing action on cell
732   morphogenesis. Cell *120*, 687–700.

733   37.   Li, H., Lin, Y., Heath, R. M., Zhu, M. X., and Yang, Z. (1999). Control of pollen tube tip
734   growth by a Rop GTPase-dependent pathway that leads to tip-localized calcium influx.
735   Plant Cell *11*, 1731–1742.

736   38.   Bourne, H. R., Sanders, D. A., and McCormick, F. (1990). The GTPase superfamily: a
737   conserved switch for diverse cell functions. Nature *348*, 125–132.

738   39.   Hodge, R. G., and Ridley, A. J. (2016). Regulating Rho GTPases and their regulators. Nat
739   Rev Mol Cell Biol *17*, 496–510.

740   40.   Hwang, J.-U., Wu, G., Yan, A., Lee, Y.-J., Grierson, C. S., and Yang, Z. (2010). Pollen-
741   tube tip growth requires a balance of lateral propagation and global inhibition of Rho-family
742   GTPase activity. J Cell Sci *123*, 340–350.

743   41.   Schmidt, A., Lv, Z., and Großhans, J. (2018). ELMO and Sponge specify subapical
744   restriction of Canoe and formation of the subapical domain in early Drosophila embryos.
745   Development *145*, dev157909.

746   42.   Berken, A., Thomas, C., and Wittinghofer, A. (2005). A new family of RhoGEFs activates
747   the Rop molecular switch in plants. Nature *436*, 1176–1180.

748   43.   Gu, Y., Li, S., Lord, E. M., and Yang, Z. (2006). Members of a novel class of Arabidopsis
749   Rho guanine nucleotide exchange factors control Rho GTPase-dependent polar growth.
750   Plant Cell *18*, 366–381.

751   44.   Qiu, J.-L., Jilk, R., Marks, M. D., and Szymanski, D. B. (2002). The Arabidopsis SPIKE1
752   Gene Is Required for Normal Cell Shape Control and Tissue Development. Plant Cell *14*,
753   101–118.

754   45.   Gu, Y., Li, S., Lord, E. M., and Yang, Z. (2006). Members of a Novel Class of Arabidopsis
755   Rho Guanine Nucleotide Exchange Factors Control Rho GTPase-Dependent Polar
756   Growth. Plant Cell *18*, 366–381.

757   46.   Guan, Y., Guo, J., Li, H., and Yang, Z. (2013). Signaling in Pollen Tube Growth: Crosstalk,
758   Feedback, and Missing Links. Mol Plant *6*, 1053–1064.

47. Yanagisawa, M., Alonso, J. M., and Szymanski, D. B. (2018). Microtubule-Dependent
Confinement of a Cell Signaling and Actin Polymerization Control Module Regulates
Polarized Cell Growth. Curr Biol *28*, 2459–2466.e4.

48. Birnbaum, K., Shasha, D. E., Wang, J. Y., Jung, J. W., Lambert, G. M., Galbraith, D. W.,
and Benfey, P. N. (2003). A gene expression map of the Arabidopsis root. Science *302*,
1956–1960.

49. Brady, S. M., Orlando, D. A., Lee, J.-Y., Wang, J. Y., Koch, J., Dinneny, J. R., Mace, D.,
Ohler, U., and Benfey, P. N. (2007). A high-resolution root spatiotemporal map reveals
dominant expression patterns. Science *318*, 801–806.

50. Borges, F., Gomes, G., Gardner, R., Moreno, N., McCormick, S., Feijó, J. A., and Becker,
J. D. (2008). Comparative Transcriptomics of Arabidopsis Sperm Cells. Plant Physiol *148*,
1168–1181.

51. Grossmann, G., Guo, W.-J., Ehrhardt, D. W., Frommer, W. B., Sit, R. V., Quake, S. R.,
and Meier, M. (2011). The RootChip: an integrated microfluidic chip for plant science.
Plant Cell *23*, 4234–4240.

52. Jones, A. M., Danielson, J. Å., Manojkumar, S. N., Lanquar, V., Grossmann, G., and
Frommer, W. B. (2014). Abscisic acid dynamics in roots detected with genetically encoded
FRET sensors. eLife *3*, e01741.

53. Stanley, C. E., Shrivastava, J., Brugman, R., Heinzelmann, E., van Swaay, D., and
Grossmann, G. (2018). Dual-flow-RootChip reveals local adaptations of roots towards
environmental asymmetry at the physiological and genetic levels. New Phytol *217*, 1357–
1369.

54. Kang, E., Zheng, M., Zhang, Y., Yuan, M., Yalovsky, S., Zhu, L., and Fu, Y. (2017). The
Microtubule-Associated Protein MAP18 Affects ROP2 GTPase Activity during Root Hair
Growth. Plant Physiol *174*, 202–222.

55. Thomas, C., Fricke, I., Weyand, M., and Berken, A. (2009). 3D structure of a binary ROP-
PRONE complex: the final intermediate for a complete set of molecular snapshots of the
RopGEF reaction. Biol Chem *390*, 427–435.

56. Thomas, C., Fricke, I., Scrima, A., Berken, A., and Wittinghofer, A. (2007). Structural
evidence for a common intermediate in small G protein-GEF reactions. Mol Cell *25*, 141–
149.

57. Cherfils, J., and Chardin, P. (1999). GEFs: structural basis for their activation of small
GTP-binding proteins. Trends Biochem Sci *24*, 306–311.

58. Wittinghofer, F. (1998). Ras signalling - Caught in the act of the switch-on. Nature News
394, 317–.

59. Li, H., Shen, J. J., Zheng, Z. L., Lin, Y., and Yang, Z. (2001). The Rop GTPase switch
controls multiple developmental processes in Arabidopsis. Plant Physiol *126*, 670–684.

60. Grefen, C., and Blatt, M. (2012). A 2in1 cloning system enables ratiometric bimolecular
fluorescence complementation (rBiFC). Biotechniques *53*.

61. Grefen, C., Obrdlik, P., and Harter, K. (2008). The Determination of Protein-protein
Interactions by the Mating-based Split-ubiquitin system (mbSUS). Methods Mol Biol *479*,
1–17.

62. Huang, G. Q., Li, E., Ge, F.-R., Li, S., Wang, Q., Zhang, C. Q., and Zhang, Y. (2013).
Arabidopsis RopGEF4 and RopGEF10 are important for FERONIA-mediated

developmental but not environmental regulation of root hair growth. New Phytol *200*,
1089–1101.

63. Payne, R. J. H., and Grierson, C. S. (2009). A theoretical model for ROP localisation by
auxin in Arabidopsis root hair cells. PLoS ONE *4*, e8337.

64. Nagashima, Y., Tsugawa, S., Mochizuki, A., Sasaki, T., Fukuda, H., and Oda, Y. (2018). A
Rho-based reaction-diffusion system governs cell wall patterning in metaxylem vessels.
Sci. Rep. *8*, 11542.

65. Oda, Y., Iida, Y., Kondo, Y., and Fukuda, H. (2010). Wood Cell-Wall Structure Requires
Local 2D-Microtubule Disassembly by a Novel Plasma Membrane-Anchored Protein. Curr
Opin Plant Biol *20*, 1197–1202.

66. Bonello, T. T., Perez-Vale, K. Z., Sumigray, K. D., and Peifer, M. (2018). Rap1 acts via
multiple mechanisms to position Canoe and adherens junctions and mediate apical-basal
polarity establishment. Development *145*, dev157941.

67. Peterson, J., Zheng, Y., Bender, L., Myers, A., Cerione, R., and Bender, A. (1994).
Interactions between the bud emergence proteins Bem1p and Bem2p and Rho-type
GTPases in yeast. J Cell Biol *127*, 1395–1406.

68. Nern, A., and Arkowitz, R. A. (1998). A GTP-exchange factor required for cell orientation.
Nature *391*, 195–198.

69. Zheng, Y., Bender, A., and Cerione, R. A. (1995). Interactions among proteins involved in
bud-site selection and bud-site assembly in Saccharomyces cerevisiae. J Biol Chem *270*,
626–630.

70. Roemer, T., Madden, K., Chang, J., and Snyder, M. (1996). Selection of axial growth sites
in yeast requires Axl2p, a novel plasma membrane glycoprotein. Genes & Development
10, 777–793.

71. Chant, J., Corrado, K., Pringle, J. R., and Herskowitz, I. (1991). Yeast BUD5, encoding a
putative GDP-GTP exchange factor, is necessary for bud site selection and interacts with
bud formation gene BEM1. Cell *65*, 1213–1224.

72. Bender, A., and Pringle, J. R. (1989). Multicopy suppression of the cdc24 budding defect
in yeast by CDC42 and three newly identified genes including the ras-related gene RSR1.
Proc Natl Acad Sci USA *86*, 9976–9980.

73. Chiou, J.-G., Balasubramanian, M. K., and Lew, D. J. (2017). Cell Polarity in Yeast. Annu
Rev Cell Dev Biol *33*, annurev–cellbio–100616–060856.

74. Wedlich-Soldner, R., Wai, S. C., Schmidt, T., and Li, R. (2004). Robust cell polarity is a
dynamic state established by coupling transport and GTPase signaling. J Cell Biol *166*,
889–900.

75. Yamaguchi, Y., Ota, K., and Ito, T. (2007). A Novel Cdc42-interacting Domain of the Yeast
Polarity Establishment Protein Bem1 IMPLICATIONS FOR MODULATION OF MATING
PHEROMONE SIGNALING. J Biol Chem *282*, 29–38.

76. Butty, A. C., Pryciak, P. M., Huang, L. S., Herskowitz, I., and Peter, M. (1998). The role of
Far1p in linking the heterotrimeric G protein to polarity establishment proteins during yeast
mating. Science *282*, 1511–1516.

77. Escobar-Restrepo, J.-M., Huck, N., Kessler, S., Gagliardini, V., Gheyselinck, J., Yang, W.-
C., and Grossniklaus, U. (2007). The FERONIA receptor-like kinase mediates male-female
interactions during pollen tube reception. Science *317*, 656–660.

78. Hruz, T., Laule, O., Szabó, G., Wessendorp, F., Bleuler, S., Oertle, L., Widmayer, P.,
Gruissem, W., and Zimmermann, P. (2008). Genevestigator v3: a reference expression
database for the meta-analysis of transcriptomes. Adv Bioinformatics *2008*, 420747.

79. Lampropoulos, A., Sutikovic, Z., Wenzl, C., Maegele, I., Lohmann, J. U., and Forner, J.
(2013). GreenGate---a novel, versatile, and efficient cloning system for plant transgenesis.
PLoS ONE *8*, e83043.

80. Griesbeck, O., Griesbeck, O., Baird, G. S., Baird, G. S., Campbell, R. E., Campbell, R. E.,
Zacharias, D. A., Zacharias, D. A., Tsien, R. Y., and Tsien, R. Y. (2001). Reducing the
Environmental Sensitivity of Yellow Fluorescent Protein. J Biol Chem *276*, 29188–29194.

81. Goedhart, J., Stetten, von, D., Noirclerc-Savoye, M., Lelimousin, M., Joosen, L., Hink, M.
A., van Weeren, L., Gadella, T. W. J., and Royant, A. (2012). Structure-guided evolution of
cyan fluorescent proteins towards a quantum yield of 93%. Nat Commun *3*, 509.

82. Heim, N., Garaschuk, O., Friedrich, M. W., Mank, M., Milos, R. I., Kovalchuk, Y., Konnerth,
A., and Griesbeck, O. (2007). Improved calcium imaging in transgenic mice expressing a
troponin C–based biosensor. Nat Meth *4*, 127–129.

83. Zuo, J., Niu, Q.-W., and Chua, N.-H. (2000). An estrogen receptor-based transactivator
XVE mediates highly inducible gene expression in transgenic plants. Plant J *24*, 265–273.

84. Xing, S., Wallmeroth, N., Berendzen, K. W., and Grefen, C. (2016). Techniques for the
Analysis of Protein-Protein Interactions in Vivo. Plant Physiol *171*, 727–758.

85. Mehlhorn, D. G., Wallmeroth, N., Berendzen, K. W., and Grefen, C. (2018). 2in1 Vectors
Improve In Planta BiFC and FRET Analyses. In The Plant Endoplasmic Reticulum
Methods in Molecular Biology. (New York, NY: Humana Press, New York, NY), pp. 139–
158.

86. Asseck, L. Y., Wallmeroth, N., and Grefen, C. (2018). ER Membrane Protein Interactions
Using the Split-Ubiquitin System (SUS). In The Plant Endoplasmic Reticulum Methods in
Molecular Biology. (New York, NY: Humana Press, New York, NY), pp. 191–203.

87. Grossmann, G., Meier, M., Cartwright, H. N., Sosso, D., Quake, S. R., Ehrhardt, D. W.,
and Frommer, W. B. (2012). Time-lapse fluorescence imaging of Arabidopsis root growth
with rapid manipulation of the root environment using the RootChip. J Vis Exp *65*, e4290.

88. Stanley, C. E., Shrivastava, J., Brugman, R., Heinzelmann, E., Frajs, V., Bühler, A., van
Swaay, D., and Grossmann, G. (2018). Fabrication and Use of the Dual-Flow-RootChip for
the Imaging of Arabidopsis Roots in Asymmetric Microenvironments. Bio-protocol *8*, 1–19.

89. Edelstein, A., Amodaj, N., Hoover, K., Vale, R., and Stuurman, N. (2010). Computer
control of microscopes using µManager. Curr Protoc Mol Biol *Chapter 14*, Unit14.20.

90. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
platform for biological-image analysis. Nat Meth *9*, 676–682.