Multi-Neuromodulator Measurements across Fronto-Striatal Network Areas of the Behaving Macaque using Solid-Phase Microextraction

Seyed-Alireza Hassani[1,2,†], Sofia Lendor[3,†], Ezel Boyaci[3], Janusz Pawliszyn[3,‡,*] and Thilo Womelsdorf[1,2,‡,*]

1 Department of Psychology, Vanderbilt University, PMB 407817, 2301 Vanderbilt Place, Nashville TN 37240, USA
2 Department of Biology, Centre for Vision Research, York University, Toronto, Ontario M6J 1P3, Canada
3 Department of Chemistry, University of Waterloo, 200 University Avenue West, Waterloo, Ontario N2L 3G1, Canada

* Corresponding author 1: Tel. 519-888-4641, Fax. 519-746-0435, E-mail: janusz@uwaterloo.ca
* Corresponding author 2: Tel. 615-322-0070, E-mail: thilo.womelsdorf@vanderbilt.edu
† ‡ These authors contributed equally

Short title: Reliable in-vivo sensing of multiple neurotransmitters using SPME

Keywords: Glutamate; Dopamine; Acetylcholine; Excitation-inhibition balance; Basal Ganglia; Prefrontal Cortex

Number of figures: 2
Number of supplementary material: 4
    supplementary methods, supplementary discussion, figures (2), tables (1)

Abstract

Different neuromodulators rarely act independent from each other to modify neural processes.

To understand this interdependence in local circuits during different brain states, it is necessary

to reliably extract local concentrations of multiple neuromodulators in vivo. Here we describe

results using solid phase microextraction (SPME), a method providing sensitive, multi-

neuromodulator measurements. Reliable measurements of glutamate, dopamine, acetylcholine

and choline were made simultaneously within frontal cortex and striatum of two macaque

monkeys during goal-directed behavior. We find glutamate concentrations several orders of

magnitude higher than acetylcholine and dopamine in all brain regions. Dopamine was reliably

detected in the striatum at tenfold higher concentrations than acetylcholine. Acetylcholine and

choline concentrations were detected with high consistency across brain areas, within and

between monkeys. These findings illustrate that SPME microprobes provide a versatile novel

tool to characterize multiple neuromodulators across different brain areas in vivo to understand

the interdependence and co-variation of neuromodulators during goal directed behavior.

Introduction

Extracellular concentrations of neuromodulators influence firing regimes, input-output

relationships and neural interactions in local circuits and long-range brain networks [1,2], and are

dysregulated in virtually all psychiatric disorders [3,4]. Accumulating evidence suggests that

these fundamental roles of neuromodulators for circuit functioning are unlikely realized by single

neuromodulators operating in isolation. Rather, neuromodulatory systems are heavily intertwined

[4–6], and operate simultaneously on individual cells and circuits [2,7–9]. In each circuit, local

mechanisms exert control over the release of neuromodulators from terminals of brainstem-

originating projection neurons. This local control proceeds through activation of pre-synaptic

glutamatergic receptors such as AMPAR and NMDAR [10–12]. These insights suggest that an

understanding of the contribution of neuromodulators to circuit functioning requires measuring,

simultaneously, multiple neuromodulators in conjunction with ongoing glutamatergic

neurotransmitter concentrations and action. Consistent with this conclusion single

neuromodulator theories often fail to account for all observable symptoms in psychiatric diseases

[13–15].

Despite the accumulating evidence for the interdependence of neuromodulator actions,

few methods exist for their simultaneous measurement in vivo and across multiple brain areas

(**Table S1**). Most of these existing neurochemical sensing methods have a high barrier to entry

by requiring specialized equipment and trained experts preventing data collection by scientists

who are otherwise interested in the role of endogenous and exogenous neuroactive chemicals in

cognition and psychiatric disorders. Electrochemical methods such as fast scan cyclic

voltammetry (FSCV) and amperometry have sub-second temporal resolution, but are limited to

the measurement of a few compounds [16,17] and are not robust for in vivo application in

nonhuman primates [18]. Imaging techniques such as positron emission tomography (PET) are

also limited to the measurement of one or a few compounds simultaneously [19]. Microdialysis

(MD) paired with mass spectrometry is the most commonly used method for measuring multiple

neuromodulators in awake behaving animals. MD provides a data collection method which is

then analyzed post hoc to identify and quantify collected analytes. It operates with a semi-

permeable membrane which allows for the continuous collection of the available extracellular

neuromodulators through passive diffusion, and can even be used to locally release

pharmacological agents [20–22]. However, MD does have several disadvantages including

disrupting the tissue during its initial placement of the probe or a guiding cannula resulting in

damage-induced release of neuromodulators that can last several hours before stable

measurements become possible. Moreover, MD has low affinity for hydrophobic compounds and

comparatively broad spatial and temporal resolution that is in the range of 200-400 μm in

diameter and 10-20 minutes, respectively. These values are dependent on the surface area of the

permeable membrane, the exact method of MD, flow rate, resolution of detection methods for

analytes of interest, tissue tortuosity and more [20–22].

In light of these limitations of available measurement techniques, we set out to

characterize a novel protocol for measuring multiple neuromodulators in vivo using probes

optimized for solid phase microextraction (SPME) [23]. SPME probes are thin (200 μm) wires of

arbitrary length coated with an inert porous polymeric matrix using biocompatible binder on one

end where molecules with appropriate size and affinity migrate via passive diffusion and are

retained by chemical interactions (*see* Methods). SPME provides an alternate method for data

collection which can then be analyzed by tools such as mass spectrometers. This method has

been shown to extract in neural tissue dynamic changes in dopamine (DA) and serotonin (5-HT)

levels with comparable precision to MD [24,25]. Additionally, due to the similarity of SPME

probes to commonly used micoelectrodes in electrophysiological recordings, relatively minor

adjustments will allow for the adaptation of conventional microelectrode driving systems for

SPME use. This, combined with post collection analysis through standard chemistry facilities

makes SPME an attractive and easy-to-use tool for electrophysiology labs.

The ability of SPME to report major neuromodulators as well as glutamate and GABA

were tested in two behaving rhesus macaques. Probes were repeatedly and simultaneously

inserted into two cortical regions and the striatum to observe inter-areal differences between

extracellular neuromodulator concentrations. We found that extracellular concentrations of

glutamate, dopamine, acetylcholine and choline could be reliably distinguished and differed

systematically between brain regions.

Results

During all measurements animals were engaged in a cognitively demanding task with stable

behavioral performance to minimize state dependent fluctuations of basal extracellular

concentrations of neuromodulators. Three brain regions were selected to provide a sample of

extracellular neuromodulatory concentrations. Two cortical regions: prefrontal cortex (PFC) area

8, premotor cortex (PreMC) area 6 and the head of the caudate nucleus (CD) were selected (**Fig**

1F). In 3 daily sessions, we sampled 3 times per session simultaneously from all three areas.

Monkey As had an additional 3 days of recording with one sample collected on each day.

Overall we collected and analyzed 12 probes in monkey As and 9 probes in monkey Ke. We

were specifically interested in major neuromodulators and neurotransmitters but successfully

measured other compounds such as amino acids (e.g. glutamine, taurine, phenylalanine etc) that

we do not discuss here.

To allow comparison of the SPME extraction results to those typically reported in

microdialysis studies, we calculated the relative change in measured concentrations across the

three successive sampling events per session pooled across monkeys to enhance the statistical

power of the analysis (**Fig 2**). Relative to the end of the first sampling event, the second and third

sampling events started after 40 minutes and 100 minutes respectively. We expected that the

variability of measurements (indexed as standard error of the median) is comparable to

repeatedly measured microdialysis of an identical, active brain state. We found that measured

concentrations did not change significantly across sampling events for any compound area

combination (Wilcoxon rank sum test; **Fig 2**).

We found that four target neurotransmitters and neuromodulators were reliably detected

in each animal: glutamate, dopamine, acetylcholine, and choline. Serotonin was also detected on

several probes but always near the limit of detection (LOD) and therefore was excluded from

analyses here (a report of serotonin is provided in the **Supplementary Results**) [26]. Glutamate

concentrations were several orders of magnitude higher than all other observed compounds of

interest in all areas and both animals. Relative to glutamate, choline concentrations were >15

times lower, DA concentrations were >700 lower and ACh concentrations were >8300 times

lower (**Fig 3**).

Glutamate concentrations (across areas) ranged from 159,147 ($\pm$19,395) to 233,659

ng/mL ($\pm$33,917) in monkey As, and from 135,523 ($\pm$22,945) to 184,333 ng/mL ($\pm$33,516) in

monkey Ke (median, SEMedian). Neither animal showed a significant difference in glutamate

concentrations between areas (As: kruskal-wallis test, $p = 0.063$; Ke: kruskal-wallis test, $p =$

0.48). However, in both monkeys, glutamate concentrations were highest in the head of the

caudate, which showed a trend to be higher in the caudate than in premotor cortex in monkey As

(Wilcoxon rank sum test, $p = 0.0226$, not significant at the Bonferroni corrected alpha level of $p=$

0.0167). Comparisons between monkeys showed no significant differences between the

glutamate concentrations in any of the measured areas (**Fig 3**).

In addition to glutamate, dopamine could be reliably detected in both monkeys in the

caudate at concentrations of 232 ng/mL ($\pm$47) in monkey As, and 226 ng/mL ($\pm$31) in monkey

As. These concentrations were similar between monkeys (Wilcoxon rank sum test p = 1, n.s.).

Dopamine was not found above detection limits in frontal cortex.

Acetylcholine measurements in the three brain areas ranged from (across areas) 25 ($\pm$2)

to 36 ng/mL ($\pm$5) in monkey As, and from 29 ($\pm$2) to 36 ng/mL ($\pm$4) in monkey Ke. Extracellular

ACh concentrations were significantly different between areas in monkey As (kruskal-wallis test,

p = 0.008). The caudate showed significantly higher concentrations than premotor cortex and a

trend to higher concentrations than PFC (PFC8a: Wilcoxon rank sum test, p = 0.04; PreM6:

Wilcoxon rank sum test, p = 0.005); only the difference between PreM6 and caudate survived

multiple comparisons correction (Bonferroni alpha = 0.0167). Cortical areas were not

significantly different from one another (Wilcoxon rank sum test, p = 0.17). In monkey Ke, only

one cortical region had concentrations of ACh above detection limits, which approached

significant difference to the concentrations measured in the caudate (Wilcoxon rank sum test, p =

0.08). Measurements of extracted ACh in the pre-motor cortex of monkey Ke were consistently

below detection limits (**Fig 3**).

We next measured choline which is a main product of the enzymatic breakdown of the

highly regulated ACh and which is a main indicator of attentional modulation of cholinergic

activity in frontal cortex [27]. Choline concentrations ranged across areas from 2,914 ($\pm$590) to

5,078 ng/mL ($\pm$675) in monkey As, and 3,408 ($\pm$3,705) to 5,533 ng/mL ($\pm$860) in monkey Ke.

Choline concentrations between areas were significantly different in monkey As (kruskal-wallis

test, p = 0.0262), but not monkey Ke (kruskal-wallis test, p = 0.25). Pairwise comparisons in

monkey As showed that concentrations after multiple comparison correction were significantly

different between PreM6 and the caudate (Wilcoxon rank sum test, p = 0.01; Bonferroni alpha =

0.0167), and approached significance for PFC8 *versus* caudate (Wilcoxon rank sum test, p =

0.06; Bonferroni alpha = 0.0167). Comparisons of choline concentrations between monkeys were

not significant in any of the measured areas.

Discussion

We demonstrated the reliable measurements of glutamate, dopamine, acetylcholine and choline

simultaneously within cortical and subcortical regions of awake and behaving macaques using

coated microprobes optimized for solid phase microextraction. Glutamate concentrations were

several orders of magnitude higher than dopamine, acetylcholine and choline across brain

regions. Extracellular concentrations of choline, dopamine and acetylcholine were detected at

>15, >700, and >8,300 times lower than glutamate respectively. Dopamine was readily detected

in the caudate but not observed at detectable concentrations in the cortical regions measured.

Acetylcholine concentrations showed a statistical trend of being different between the caudate

and measured cortical areas and with high consistency between monkeys. Choline

concentrations, a product of acetylcholine degradation as well as a precursor for its synthesis,

were negatively correlated with acetylcholine (R = -0.355; p = 0.01). Together, these findings

provide new and rare insights about the neurochemical circuit profiles during an active brain

state in three areas of the primate fronto-striatal network.

The high consistency of measured concentrations within animals, between brain areas

and between monkeys suggests that SPME microprobes could provide a versatile neuro-

technique for understanding how variations of the neurochemical milieu relate to local and long-

range circuit operations and ultimately cognitive functioning.

Extracellular Concentrations of Glutamate, Dopamine, Acetylcholine and Choline. Our

results revealed particularly high levels of glutamate within all measured brain regions (PFC,

PreMC, and CD) relative to other neuromodulators. This finding is consistent with its ubiquitous

role as major excitatory neurotransmitter. This holds true even in the caudate which consists

exclusively of GABAergic neurons, but receives glutamatergic inputs from the cortex [28].

Tonic concentrations of glutamate within the striatum have been previously reported to be

comparable to that of the hippocampus, prefrontal cortex and other cortical areas due to

glutamatergic inputs, extra-synaptic release and glial release [29–31]. In contrast, classical

neuromodulators such as DA and ACh, are released in extracellular space through volume

transmission where they are highly regulated and may bind to receptors on multiple neurons [32–

35]. This regulation may help explain the relatively low observed concentrations of DA and ACh

and may be a large source of extracellular choline through the enzymatic breakdown of ACh.

Extracellular DA was measured well above the detection threshold in the striatum. In

combination with glutamatergic and cholinergic concentrations, our results show promise in the

application of SPME to further our understanding of the relationship between glutamatergic

inputs to the striatum and the neuromodulatory ACh and DA signals impinging on striatal

circuits. Such circuits are strongly involved in hypo- and hyperkinetic diseases [3,15].

Dopaminergic signals did not exceed the detection threshold in either cortical regions. The

presence of DA in the PFC is well documented [36] and thus has failed to exceed our detection

limit. A possible explanation for the lack of detectable cortical DA is the susceptibility of

catecholamines to oxidation. This may be partially addressed by faster coupling of sample

collection to quantitation by MS (see **Supplementary Discussion**).

Few previous studies have documented ACh concentrations in vivo due to its rapid

breakdown by acetylcholinesterase [35,37,38]. ACh plays a major role in organizing local

circuits, deployment of attention, locomotion and reward through different receptors present in

cortical and subcortical regions [6,39,40]. It has been difficult to discern which ACh

concentration corresponds to efficient endogenous circuit operations. Our finding of

measurements in all sampling locations except for premotor cortex in monkey Ke, is therefore a

promising starting point for future studies comparing ACh concentrations, in conjunction with

other neuromodulators, during different cognitive states. The fast turn-around of ACh due to

AChE however, likely leads to an underestimation of ACh concentrations by SPME. Although

AChE cannot traverse the porous SPME coating due to its size (~7.6nm diameter), preventing

any enzymatic degradation of trapped ACh (**Supplementary Material**), the dynamic

equilibration process continuously leads to a transfer of ACh out of the SPME coating following

the concentration gradient of the rapidly depleted ACh into the extracellular space as discussed

in the case of calibration in the coating approach [41,42].

Amongst other sources, choline is created as a byproduct of ACh degradation by AChE.

Choline is also a precursor for the synthesis of ACh. Given this relationship, a negative

correlation between choline concentrations and ACh is expected which we indeed observed in

our dataset. Despite the difference in the measured concentrations of choline and acetylcholine,

other studies with similar differences have demonstrated the relationship of ACh and choline to

behavior within the cortex [27]. Future studies may focus on quantifying this relationship across

a wider range of brain states.

Reliable Measurement of Individual Differences of State Specific Neuromodulatory Tone.

Our results suggest that SPME probes provide reliable and sensitive measurements in consecutive sampling events within an experimental session and between sessions on consecutive days. These results were obtained while we controlled brain states across measurement events by engaging animals in a cognitive task. This experimental control could have contributed to the comparable extracellular levels of neuromodulators from sample to sample within *and* between days from similar brain locations (**Fig 2**) [43–46]. The reproducibility of multiple measurements at the same anatomical location within and between days suggests that the SPME penetrations did not significantly disturb tissue [24] (**Supplementary Information**). This conclusion contrasts with reported experience from microdialysis measurements, where the initial placement of the probe or guiding cannula causes transient measurement instabilities that can require several hours of settling time before reliable, steady state neuromodulator concentrations are measurable [29,47–50]. These waiting periods must be taken into account when designing microdialysis experiments for in vivo tracking of neuromodulators during behavioral tasks in awake and restrained animals. Furthermore, the reliability of observations for the reported neurochemicals with SPME is extremely high and outperforms many reported microdialysis studies. We present 100% reliability in reporting glutamate and choline concentrations in all brain regions and subjects. Acetylcholine concentrations were reported with 100% reliability in one animal and 92% (11/12 probes) in the other within the prefrontal cortex, 0% reliability (below detection threshold) in one animal and 83% (10/12 probes) in the other within the premotor cortex and 100% reliability in one animal and 92% (11/12 probes) in the other within the caudate. Dopamine concentrations were reported

with 100% reliability in both animals in the caudate and were not observed above detection

thresholds in the cortex.

Qualities and Advantages of SPME. Our results illustrate several inherent advantages of using

SPME to measure neurochemical profiles in brain circuits [26]. As illustrated in **Table S1**,

SPME complements or outperforms alternative techniques in various domains. Practically, the

arbitrary length of the wire where the SPME coating is placed allows for robust placement within

the brain. The narrow diameter of the SPME probe that is within the range common to electrodes

for electrophysiological recordings, allows for repeated, simultaneous sampling at multiple sites

without observable changes to detected extracellular neuromodulator levels using 20-minute

sampling times.

Spatial Resolution. In principle, the spatial resolution of the SPME measurements are

capable of reflecting laminar concentration gradients across areas as small as several tens of

micrometers. In our protocol we coated 3mm in length with 50 μm of thickness and desorbed the

entire area into an organic solvent mixture for subsequent LC-MS/MS analysis. However, with

the use of other techniques, such as desorption electrospray ionization (DESI), which represents

a direct-to-MS approach [51], much smaller areas of the SPME matrix may be analyzed at a time

allowing for laminar or near-laminar resolution. Direct coupling of SPME fibres to MS is the

preferred mode, as it provides the most rapid analysis thus preventing diffusion of compounds

within the coating and smearing the imprinted layer information. If diffusion of molecules within

the fibre is high, it can be prevented by, for example, the inclusion of a divider between two

halves of the coating. This would allow the simultaneous measurement of separated

compartments of a cortical column. The absolute spatial resolution of SPME is difficult to

estimate as SPME operates through diffusion and dynamic equilibration similar to MD. As a

consequence, the sampled volume will depend on sampling time, the compound's diffusion

coefficient, as well as the target tissue and its properties affecting the rate of diffusion such as

tortuosity [32].

Temporal Resolution. Several factors determine the appropriate extraction time using

SPME. Depending on the goals of an experimenter, resolutions below 5 minutes may be

achieved as the detection limit is reached within the first few minutes for most compounds tested

(**Fig 1D-E**). As sampling continues, the SPME coating will eventually reach an equilibrium point

with the external environment (**Fig 1D-E**). In principle, sampling times beyond a stable

equilibrium point will result in the same extracted values of compounds. However, in the brain

there is no single stable equilibrium point and thus, even with long sampling times, there will be

some variation in extracted concentrations. Within the dynamic regime, we can still calculate

environmental concentrations from which sampling occurred given knowledge of the extraction

time and the dynamic range. The lower limit of this dynamic range that can be informative is

determined by the coating's physicochemical characteristics and sample's properties and in

practice also volume of desorption and sensitivity of the MS. The analytical technique's

sensitivity also practically determines the thickness of the SPME coating as thinner coatings

could in principle increase temporal resolution but would extract less compounds overall

requiring higher sensitivity to detect. With higher sensitivity, less time is required for the thin

coating to extract sufficient amounts to exceed limits of detection and quantitation. This then

means that the compound with regards to which the sensitivity is the lowest, limits the temporal

resolution because it will have the slowest-to-reach detectable concentration. However, making

measurements in the dynamic range requires consistency of extraction times as differences in

time will lead to variations of extraction yields in a range that can be estimated from the SPME

extraction time profiles (**Fig 1D-E**). Given the dynamic nature of brain networks and the lack of

a stable equilibrium point, we expect consistency in brain states to be one of the biggest factors

in reducing replicated variability. Moreover, SPME measurements are inherently an average of

the temporal dynamics of the target tissue milieu. This means that due to the bi-directional

exchange of neuromodulators from the extracellular space and the SPME coating, longer periods

of stable concentration gradients will be more strongly reflected in the extracted measurements.

Utility and Ease of Use. SPME based microprobe extraction allows for multiple

simultaneous measurements that can be reliably repeated within the same measurement sessions

at the same locations with no evidence of damage induced disruption of the neurochemical

environment. Another critical advantage of SPME over comparable MS-analysis based methods

such as MD is its ease of use and accessibility. Due to their similar size to recording

microelectrodes used for electrophysiology, little adjustment is required for conducting SPME

measurements using existing acute microelectrode positioning systems. Measurements in deeper

structures should use an accompanying guide tube (**Fig 1C**) as travelling through non-target

tissue will result in unwanted chemical collection. Beyond the collection, a chemistry core could

apply LC-MS/MS to the probes allowing for relatively easy quantitation of extracellular

compounds of interest.

Detection Methods. SPME is ultimately a sampling method, much like MD, that provides

data for analytical tools such as chromatography and mass spectrometry. In fact, once SPME

analytes have been desorbed from the probe into a solvent, data provided by SPME and MD are

treated very much the same. Thus, the advancement in detection limits and reliability of post-hoc

analytical methods utilized by MD also benefit SPME, and vice versa. However, SPME sample

analysis is more reliably and conveniently coupled to LC-MS/MS as it lacks issues such as high

salt content of collected dialysate that MD suffers from [52].

Extension of SPME measurements beyond classical neuromodulators. A unique

advantage of SPME over alternative methods of in vivo detection of compounds within the brain

is its potential affinity for hydrophobic compounds. Although MD is capable of collecting

metabolomics data similarly to SPME [20,53,54], the artificial cerebra-spinal fluid that is

commonly traversing the semi-permeable membrane better facilitates the collection of

hydrophilic molecules. This means that molecules that cross the semi-permeable membrane and

are therefore collected and quantified through LC-MS are much more likely to be hydrophilic.

SPME, in contrast, provides balanced analyte coverage, which means that in principle certain

SPME matrix coatings have similar affinity for both hydrophilic or hydrophobic compounds

[55]. Data collected here includes other neurochemicals of interest such as amino acids (e.g.

glutamine, taurine, phenylalanine, etc) that we reliably detected but do not discuss here. In

practice, the detection of very polar molecules such as monoamines and catecholamines requires

coating chemistry that facilitates hydrophilic compound extraction. Comparatively many lipids

play important roles in intracellular signaling and have been suggested to provide biomarkers for

psychiatric disorders [56,57], therefore making SPME a potentially versatile and comprehensive

tool for brain neurochemistry studies.

Implications for Understanding and Treating Psychiatric Disease States. Most psychiatric

disorders are accompanied by neuromodulatory dysregulation [3]. However, this fact is seldom

studied in a multi-modulatory manner with a single or few nuclei or neurochemicals being

observed at a time. This is an increasingly evident problem because models attributing symptoms

of a disorder to a single neuromodulator often fall short in explaining many symptoms [13,14].

For example, in Parkinson's disease, outside of the well characterized dopaminergic deficits,

there is evidence for deficits in noradrenergic, cholinergic and serotonergic systems as well

[6,15,58]. Many cognitive deficits observed within Parkinson's disease are linked to such non-

dopaminergic deficits. Such findings emphasize the need to simultaneously observe multiple

neuromodulatory systems. Thus the simultaneous measurement of multiple neuromodulators, in

multiple brain regions within healthy and clinical populations will allow for a better

understanding of the underlying causes of symptoms and progression of psychiatric disorders

[3,4,59].

A better understanding of psychiatric disorders through multi-modulator methods may

also lead to more accurate understanding of the action of pharmacological agents. Previously,

many studies aimed at identifying the locus of action of a pharmacological agent have used local

injection methods such as iontophoresis or microinjections [60,61]. Although highly informative

about the role of neuromodulators in modulating the activity of individual cells and circuits, this

approach does not allow physiologically realistic exploration of a systemically administered

pharmacological agent. Pharmacological agents are often administered in some systemic fashion

and even with highly specific receptor affinities, may interact with multiple neuromodulatory

systems through the actions of heteroreceptors [4,5,62]. Multi-modulator measurements, as

described here, will allow for a better understanding of pharmacological agents, as well as

provide novel insights into the development of more effective drugs or combinations of drugs to

better treat the clinical population.

Conclusion. We described a novel SPME protocol capable of simultaneous, multi-modulator

measurements of multiple brain regions. Our results suggest that SPME both supplements

current methods of neuromodulator detection and allows for novel measurements previously not

possible for the investigation and dissection of neuromodulatory systems, their role in

physiological brain processes and their modulation by pharmacological agents.

Methods

Animals

Data was collected from two 8 year-old male rhesus macaques (*Macaca mulatta*) weighing 8-12

kg. All animal care and experimental protocols were approved by the York University Animal

Care Committee and were in accordance with the Canadian Council on Animal Care guidelines.

Details regarding the experimental setup, recording procedures, and reconstruction of recording

sites have been described previously [63]. Briefly, animals were implanted with a 20 mm by 28

mm recording chamber over the frontal region of the right hemisphere guided by stereotaxic

coordinates [64] and MR images. The animals were seated in a custom made primate chair and

head stabilized with their eyes 65cm away from a 21' LCD monitor refreshed at 85 Hz. Eye

traces were collected by a video-based eye-tracking system (Eyelink 1000 Osgoode, Ontario,

Canada, 500 Hz sampling rate). The animals were engaged in an over-trained attention tasks in

which they would use saccadic eye movements to acquire juice reward. The specifics of the task

are described elsewhere [65]. Both animals showed stable performance and acquired similar

reward volumes on all recording days. In both tasks, stimulus presentation and reward delivery

was controlled through MonkeyLogic (http://www.brown.edu/Research/monkeylogic/).

SPME protocol and Fabrication of SPME Probes

We provide a visual overview of the complete SPME protocol used here in **Fig S1** and delineate the chemical materials, LC-MS/MS analysis, and detailed quantitation of neuromodulators in the ***Supplementary Methods*** and in a companion paper [26]. All measurements used miniaturized SPME probes that were manufactured by repeated dip-coating of stainless steel wires in a suspension of extracting phase in a binder. This general procedure reported previously [66], has been modified to fit the purpose of *in vivo* brain sampling [26]. The 3 mm long tip of the wire was acid-etched down to approx. 100 μm to create a recession capable of accommodating thicker layer of coating without significantly increasing the total probe diameter. The extracting phase was an in-house synthesized hydrophilic-lipophilic balance polymer functionalized with benzenesulfonic acid to introduce strong cation exchange properties (HLB-SCX). The monodispersed polymeric particles with diameter of approx. 1 μm were suspended in the binder consisting of 7% polyacrylonitrile dissolved in N,N-dimethylformide (w/v), ensuring the particles-to-binder ratio at 15% (w/v). The extracting phase suspension was prepared one day before the probe coating, homogenized by sonication and stirred at 800 rpm overnight. Several layers of coating were deposited on the modified wires, resulting in total coating thickness of ≈50 μm and total probe diameter of 195 μm. The average pore size of the extracting phase was measured at 1.2nm. Before the use, the probes were cleaned with the mixture of methanol, acetonitrile and 2-propanol (50:25:25, v/v/v), activated in the mixture of methanol and water (50:50. v/v) and sterilized in steam for 15 min at 121°C. During the sampling, SPME probes were inserted into sheathing cannulas to protect the coating and prevent it from extracting compounds on the way to the target brain area. The cannulas underwent the same cleaning and

sterilization procedure as the probes and then the lengths of both components of the SPME

assembly were adjusted to 60-70 and 70-80 mm for the probes and cannulas, respectively.

MRI Guided Electrophysiological Mapping of Target Tissue

The anatomical coordinates of the brain regions of interest were first identified through 3 T MR

images. The MR images were then verified with extracellular electrophysiological recordings of

the target areas, which provided the gray and white matter boundaries for the cortical sites and

the dorsal most aspect of the head of the caudate nucleus. Tungsten microelectrodes were 200

μm thick with an impedance of 1-2 MΩ. All electrodes, SPME probes and their accompanying

guiding cannulas were driven down into the brain and later out using software-controlled

precision microdrives (Neuronitek, ON, Canada). Electrodes were connected to a multichannel

acquisition processor (Neuralynx Digital Lynx system, Inc., Bozeman, Montana, USA) which

was used for data amplification, filtering and acquisition of spiking activity. Spiking activity was

obtained by applying a 600-8000 Hz bandpass filter, with further amplification and digitization

at a 32 KHz sampling rate. For every recording day, electrodes were lowered until the first

detection of spiking activity (indicative of gray matter) at the depth suggested by the MR images.

SPME Sampling and Post-Processing Procedures

All three SPME assemblies (example of sampling using one assembly in **Fig 1A-C**) were

simultaneously driven to 200 μm above the point of first spiking detection. SPME assemblies

were located ~1 mm away from the electrode penetration location. Then, only the SPME probes

were inserted 3 mm into the areas of interest. On average macaque cortical thickness is only 2

mm while the head of the caudate at the point of sampling was well over 3 mm. An extraction

duration of 20 min was selected to reflect stable extraction time profiles for all compounds of

interest. Extraction time profiles reflect the detectable concentration of collected analyte over

time, which is captured in sequence by a linear "kinetic" regime, a "dynamic" regime and an

"equilibrium" regime over extraction time. We illustrate in **Fig 1D-E** these regimes for two main

compounds measured with the SPME probe we used in vivo. After the 20 min extraction event,

all SPME probes were driven back 3 mm into the guiding cannulas and all SPME assemblies

were withdrawn from the brain (**Fig 1A-C**).

The microdrives were removed from the chamber to enable unclamping of the SPME

probes, a brief wash and then storage in glass vials surrounded by dry ice until placing them into

a -80°C freezer. One entire sampling event (one extraction in 3 different brain areas together

with assembling the SPME probes and cannulas, driving into and out of brain, washing and

preparing for storage) was performed within 50 min, except for the first sampling event in each

sampling day. The first sampling event, with the area identification using electrophysiology

recording, was performed within 75 min.

Neuromodulators detection and quantitation

Detection and quantitiation of neuromodulators followed previously established procedures

described in the **Supplementary Material**. In brief, SPME probes were defrosted and desorbed

in an aqueous solution containing water, acetonitrile, methanol, formic acid and deuterated

isotopologues of target neuromodulators as internal standards (IS). Chromatographic separation

of target compounds was conducted as previously reported [25]. Tandem mass spectrometry

(MS/MS) was performed with electrospray ionization with two MS/MS transitions for each

neuromodulator and one for each IS [26].

Stock solutions of all target neuromodulators were prepared and used to generate

standard calibration curves and to calculate the amounts of neuromodulator extracted by each

experimental SPME probe. Conversion of the amount of neuromodulator extracted to in-brain

concentrations was done using matrix-matched external calibrations. LOD is defined as 3 times

the signal obtained from blank samples.

460
461

Acknowledgments

This research was supported by grants from the Canadian Institutes of Health Research (CIHR),

the Natural Sciences and Engineering Research Council of Canada (NSERC) and the Ontario

Ministry of Economic Development and Innovation (MEDI). We thank Dr. Hongying Wang for

invaluable help with animal care. We thank Thermo Scientifc for providing the triple quadrupole

mass spectrometer TSQ Quantiva used in this work as a loan to our laboratory. The authors

thankfully acknowledge Pfizer Canada Inc., Merck Canada Inc., Quebec Consortium for Drug

Discovery (CQDM), Brain Canada, and Ontario Brain Institute for the grant "Solid phase

microextraction-based integrated platform for untargeted and targeted in vivo brain studies".

Conflicts of interest

None of the authors have any conflict to disclose.

References

1. Thiele A, Bellgrove MA. Neuromodulation of Attention. Neuron. Elsevier Inc.; 2018;97: 769–785. doi:10.1016/j.neuron.2018.01.008
2. Marder E. Overview Neuromodulation of Neuronal Circuits: Back to the Future. Neuron. Elsevier Inc.; 2012;76: 1–11. doi:10.1016/j.neuron.2012.09.010
3. Millan MJ, Agid Y, Brüne M, Bullmore ET, Carter CS, Clayton NS, et al. Cognitive dysfunction in psychiatric disorders: characteristics, causes and the quest for improved therapy. Nat Rev Drug Discov. Nature Publishing Group; 2012;11: 141–168. doi:10.1038/nrd3628
4. Avery MC, Krichmar JL. Neuromodulatory Systems and Their Interactions : A Review of Models , Theories , and Experiments. Front Neural Circuits. 2017;11: 1–18. doi:10.3389/fncir.2017.00108
5. Gobert A, Audinot V, Cistarelli L, Millan MJ. Simultaneous quantification of serotonin, dopamine and noradrenaline levels in single frontal cortex dialysates of freely-moving rats reaveals a complex pattern of reciprocal auto- and heteroreceptor-mediated control of release. Neuroscience. 1998;84: 413–429.
6. Moehle MS, Pancani T, Byun N, Yohn SE, Wilson GH, Dickerson JW, et al. Cholinergic Projections to the Substantia Nigra Pars Reticulata Inhibit Dopamine Modulation of Basal Ganglia through the M4Muscarinic Receptor. Neuron. 2017;96: 1358–1372. doi:10.1016/j.neuron.2017.12.008
7. Santana N, Mengod G, Artigas F. Expression of a1-adrenergic receptors in rat prefrontal cortex: cellular co-localization with 5-HT2A receptors. Int J Neuropsychopharmacol. 2018;16: 1139–1151. doi:10.1017/S1461145712001083
8. Arnsten AFT, Paspalas CD, Gamo NJ, Yang Y, Wang M. Dynamic network connectivity: A new form of neuroplasticity. Trends Cogn Sci. Elsevier Ltd; 2010;14: 365–375. doi:10.1016/j.tics.2010.05.003
9. Hassan SF, Zumut S, Burke PG, Mcmullan S, Cornish JL, Goodchild AK. Comparison of noradrenlaine, dopamine and serotonin in mediating the tachycardic and thermogenic effects of methamphetamine in the ventral medial prefrontal cortex. Neuroscience. IBRO; 2015;295: 209–220. doi:10.1016/j.neuroscience.2015.03.028
10. Wang JKT, Andrews H, Thukral V. Presynaptic Glutamate Receptors Regulate Noradrenaline Release from Isolated Nerve Terminals. J Neurochem. 1992;58: 204–211.
11. Ghersi C, Bonfanti A, Manzari B, Feligioni M, Raiteri M, Pittaluga A. Pharmacological heterogeneity of release-regulating presynaptic AMPA / kainate receptors in the rat brain : study with receptor antagonists. Neurochem Int. 2003;42: 283–292.
12. Pittaluga A, Feligioni M, Longordo F, Luccini E, Raiteri M. Trafficking of presynaptic AMPA receptors mediating neurotransmitter release : Neuronal selectivity and relationships with sensitivity to cyclothiazide. Neuropharmacology. 2006;50: 286–296. doi:10.1016/j.neuropharm.2005.09.004
13. Bohnen NB, Albin R, Muller MLTM, Petrou M, Kotagal V, Koeppe RA, et al. Frequency of Cholinergic and Caudate Nucleus Dopaminergic Deficits Across the Predemented Cognitive Spectrum of Parkinson Disease and Evidence of Interaction Effects. JAMA Neurol. 2015;72: 194–200. doi:10.1001/jamaneurol.2014.2757
14. Remy P, Doder M, Lees A, Turjanski N, Brooks D. Depression in Parkinson ' s disease: loss of dopamine and noradrenaline innervation in the limbic system. Brain. 2005;128:

1314–1322. doi:10.1093/brain/awh445

15. Halliday GM, Leverenz JB, Schneider JS, Adler CH. The Neurobiological Basis of
Cognitive Impairment in Parkinson's Disease. Mov Disord. 2014;29: 634–650.
doi:10.1002/mds.25857

16. Jacobs CB, Peairs MJ, Venton BJ. Review: Carbon nanotube based electrochemical
sensors for biomolecules. Anal Chim Acta. 2010;662: 105–127.
doi:10.1016/j.aca.2010.01.009

17. Dale N, Hatz S, Tian F, Llaudet E. Listening to the brain: Microelectrode biosensors for
neurochemicals. Trends Biotechnol. 2005;23: 420–428. doi:10.1016/j.tibtech.2005.05.010

18. Vartak D, Togt C Van Der, Vugt B Van, Roelfsema PR. Electrochemical measurement of
acetylcholine in the dorsolateral prefrontal cortex : A technical report. BioRXiv. 2017; 1–
25.

19. Fisher RE, Morris ED, Alpert NM, Fischman AJ. In Vivo Imaging of Neuromodulatory
Synaptic Transmission Using PET : A Review of Relevant Neurophysiology. Hum Brain
Mapp. 1995;3: 24–34. doi:https://doi.org/10.1002/hbm.460030104

20. Anderzhanova E, Wotjak CT. Brain microdialysis and its applications in experimental
neurochemistry. Cell Tissue Res. 2013;354: 27–39. doi:10.1007/s00441-013-1709-4

21. Watson CJ, Venton BJ, Kennedy RT. In Vivo Measurements of Neurotransmitters by
Microdialysis Sampling. Anal Chem. 2006;78: 1391–1399. doi:10.1021/ac0693722

22. Kennedy RT. Emerging trends in in vivo neurochemical monitoring by microdialysis.
Curr Opin Chem Biol. Elsevier Ltd; 2013;17: 860–867. doi:10.1016/j.cbpa.2013.06.012

23. Pawliszyn J. Theory of Solid-Phase Microextraction. In: Pawliszyn J, editor. Handbook of
Solid Phase Microextraction. Elsevier; 2012. pp. 13–59.

24. Cudjoe E, Bojko B, de Lannoy I, Saldivia V, Pawliszyn J. Solid-Phase Microextraction: A
Complementary In Vivo Sampling Method to Microdialysis. Angew Chemie Int Ed.
2013;52: 12124–12126. doi:10.1002/anie.201304538

25. Cudjoe E, Pawliszyn J. Optimization of solid phase microextraction coatings for liquid
chromatography mass spectrometry determination of neurotransmitters. J Chromatogr A.
Elsevier B.V.; 2014;1341: 1–7. doi:10.1016/j.chroma.2014.03.035

26. Lendor S, Hassani S-A, Boyaci E, Singh V, Womelsdorf T, Pawliszyn J. Development of
solid phase microextraction-based miniaturized probe and protocol for extraction of
neurotransmitters from brain in vivo.

27. Parikh V, Kozak R, Martinez V, Sarter M. Prefrontal Acetylcholine Release Controls Cue
Detection on Multiple Timescales. Neuron. 2007;56: 141–154.
doi:10.1016/j.neuron.2007.08.025

28. Haber SN, Knutson B. The reward circuit: Linking primate anatomy and human imaging.
Neuropsychopharmacology. Nature Publishing Group; 2010;35: 4–26.
doi:10.1038/npp.2009.129

29. Moussawi K, Riegel A, Nair S, Kalivas PW, Bargas J, Nacional U. Extracellular
glutamate : functional compartments operate in different concentration ranges. Front Syst
Neurosci. 2011;5: 1–9. doi:10.3389/fnsys.2011.00094

30. Moghaddam B. Stress Preferentially Increases Extraneurona Levels of Excitatory Amino
Acids in the Prefrontal Cortex : Comparison to Hippocampus and Basal Ganglia. J
Neurochem. 1993;60: 1650–1657.

31. Ascenzo MD, Fellin T, Terunuma M, Revilla-sanchez R, Meaney DF, Auberson YP, et al.
mGluR5 stimulates gliotransmission in the nucleus accumbens. Proc Natl Acad Sci.

2006;104: 1995–2000.

32.   Syková E. Extrasynaptic volume transmission and diffusion parameters of the
extracellular space. Neuroscience. 2004;129: 861–876.
doi:10.1016/j.neuroscience.2004.06.077

33.   Vizi ES. Role of high-affinity receptors and membrane transporters in nonsynaptic
communication and drug action in the central nervous system. Pharmacol Rev. 2000;52:
63–89.

34.   Santello M, Calì C, Bezzi P. Gliotransmission and the Tripartite Synapse. In: Kreutz M,
Sala C, editors. Synaptic Plasticity Advances in Experimental Medicine and Biology.
2012. pp. 307–331.

35.   Schwartz J, Javitch J. Neurotransmitters. In: Kandel E, Schwartz J, Jessell T, Siegelbaum
S, Hudspeth A, editors. Principles of Neural Science. 2013. pp. 289–306.

36.   Goldman-Rakic PS, Lidow MS, Gallager DW. Overlap of dopaminergic, adrenergic, and
serotoninergic receptors and complementarity of their subtypes in primate prefrontal
cortex. J Neurosci. 1990;10: 2125–2138.

37.   Kreitzer AC. Physiology and pharmacology of striatal neurons. Annu Rev Neurosci.
2009;32: 127–47. doi:10.1146/annurev.neuro.051508.135422

38.   Zapata A, Chefer VI, Parrot S, Denoroy L. Detection and Quantification of
Neurotransmitters in Dialysates. Curr Protoc Neurosci. 2013;63: 1–35.
doi:10.1002/0471142301.ns0704s63

39.   Granger AJ, Mulder N, Saunders A, Sabatini BL. Cotransmission of acetylcholine and
GABA. Neuropharmacology. Elsevier Ltd; 2016;100: 40–46.
doi:10.1016/j.neuropharm.2015.07.031

40.   Howe WM, Gritton HJ, Lusk NA, Roberts EA, Hetrick VL, Berke JD, et al. Acetylcholine
Release in Prefrontal Cortex Promotes Gamma Oscillations and Theta–Gamma Coupling
during Cue Detection. J Neurosci. 2017;37: 3215–3230. doi:10.1523/JNEUROSCI.2737-
16.2017

41.   Zhang X, Es-haghi A, Musteata FM, Ouyang G, Pawliszyn J. Quantitative in Vivo
Microsampling for Pharmacokinetic Studies Based on an Integrated Solid-Phase
Microextraction System. Anal Chem. 2007;79: 4507–4513. doi:10.1021/ac070177c

42.   Alam N, Pawliszyn J. Numerical Simulation and Experimental Validation of Calibrant-
Loaded Extraction Phase Standardization Approach. Anal Chem. 2016;88: 8632–8639.
doi:10.1021/acs.analchem.6b01802

43.   Marder E, O'Leary T, Shruti S. Neuromodulation of Circuits with Variable Parameters:
Single Neurons and Small Circuits Reveal Principles of State-Dependent and Robust
Neuromodulation. Annu Rev Neurosci. 2014;37: 329–346. doi:10.1146/annurev-neuro-
071013-013958

44.   McGinley MJ, Vinck M, Reimer J, Batista-Brito R, Zagha E, Cadwell CR, et al. Waking
State: Rapid Variations Modulate Neural and Behavioral Responses. Neuron. Elsevier
Inc.; 2015;87: 1143–1161. doi:10.1016/j.neuron.2015.09.012

45.   Reimer J, Froudarakis E, Cadwell CR, Yatsenko D, Denfield GH, Tolias AS. Pupil
Fluctuations Track Fast Switching of Cortical States during Quiet Wakefulness. Neuron.
Elsevier Inc.; 2014;84: 355–362. doi:10.1016/j.neuron.2014.09.033

46.   Reimer J, McGinley MJ, Liu Y, Rodenkirch C, Wang Q, McCormick DA, et al. Pupil
fluctuations track rapid changes in adrenergic and cholinergic activity in cortex. Nat
Commun. Nature Publishing Group; 2016;7: 1–7. doi:10.1038/ncomms13289

47.   Watanabe M, Kodama T, Hikosaka K. Increase of Extracellular Dopamine in Primate
Prefrontal Cortex During a Working Memory Task. J Neurophysiol. 1997;78: 2795–2798.
doi:https://doi.org/10.1152/jn.1997.78.5.2795
48.   Kodama T, Hikosaka K, Honda Y, Kojima T, Tsutsui K, Watanabe M. Dopamine and
glutamate release in the anterior default system during rest : A monkey microdialysis
study. Behav Brain Res. Elsevier B.V.; 2015;294: 194–197.
doi:10.1016/j.bbr.2015.08.013
49.   Kodama T, Kojima T, Honda Y, Hosokawa T, Tsutsui K, Watanabe M. Oral
Administration of Methylphenidate (Ritalin) Affects Dopamine Release Differentially
Between the Prefrontal Cortex and Striatum: A Microdialysis Study in the Monkey. J
Neurosci. 2017;37: 2387–2394. doi:10.1523/JNEUROSCI.2155-16.2017
50.   Kodama T, Hikosaka K, Watanabe M. Differential changes in glutamate concentration in
the primate prefrontal cortex during spatial delayed alternation and sensory-guided tasks.
Exp Brain Res. 2002;145: 133–141. doi:10.1007/s00221-002-1084-y
51.   Deng J, Yang Y, Wang X, Luan T. Strategies for coupling solid-phase microextraction
with mass spectrometry. TrAC - Trends Anal Chem. Elsevier Ltd; 2014;55: 55–67.
doi:10.1016/j.trac.2013.12.004
52.   Guihen E, Connor WTO. Current separation and detection methods in microdialysis the
drive towards sensitivity and speed. Electrophoresis. 2009;30: 2062–2075.
doi:10.1002/elps.200900039
53.   Li J, von Pföstl V, Zaldivar D, Zhang X, Logothetis NK, Rauch A. Measuring multiple
neurochemicals and related metabolites in blood and brain of the rhesus monkey by using
dual microdialysis sampling and capillary hydrophilic interaction chromatography-mass
spectrometry. Anal Bioanal Chem. 2012;402: 2545–2554. doi:10.1007/s00216-011-5427-
z
54.   Zhang X, Rauch A, Lee H, Xiao H, Rainer G, Logothetis N. Capillary hydrophilic
interaction chromatography/mass spectrometry for simultaneous determination of multiple
neurotransmitters in primate cerebral cortex. Rapid Commun Mass Spectrom. 2007;21:
3621–3628.
55.   Reyes-garcés N, Gionfriddo E, Gómez-Ríos GA, Alam MN, Boyaci E, Bojko B, et al.
Advances in Solid Phase Microextraction and Perspective on Future Directions. Anal
Chem. 2018;90: 302–360. doi:10.1021/acs.analchem.7b04502
56.   Tamiji J, Crawford A. The Neurobiology of Lipid Metabolism in. Eur PMC. 2011;18: 98–
112. doi:10.1159/000323189
57.   Yehuda S, Rabinovitz S, Mostofsky DI. Mini-Review Essential Fatty Acids Are Mediators
of Brain Biochemistry and Cognitive Functions. J Neurosci Res. 1999;56: 565–570.
58.   Halliday GM, Li YW, Blumbergs PC, Joh TH, Cotton PRGH, Howe PRC, et al.
Neuropathology of Immunohstochemically Idendied Brainstem Neurons in Parkinson's
Disease. Am Neurol Assoc. 1990;27: 373–385.
59.   Anderzhanova E, Wotjak CT. NextGen Brain Microdialysis : Applying Modern
Metabolomics Technology to the Analysis of Extracellular Fluid in the Central Nervous
System. Mol Neuropsychiatry. 2015;1: 60–67. doi:10.1159/000381855
60.   Wang M, Ramos BP, Paspalas CD, Shu Y, Simen A, Duque A, et al. alpha2A-
Adrenoceptors Strengthen Working Memory Networks by Inhibiting cAMP-HCN
Channel Signaling in Prefrontal Cortex. Cell. 2007;129: 397–410.
doi:10.1016/j.cell.2007.03.015

61.   Lapiz MDS, Morilak DA. Noradrenergic modulation of cognitive function in rat medial
prefrontal cortex as measured by attentional set shifting capability. Neuroscience.
2006;137: 1039–1049. doi:10.1016/j.neuroscience.2005.09.031

62.   Millan MJ, Goodwin GM, Meyer-lindenberg A, Ove S. Learning from the past and
looking to the future : Emerging perspectives for improving the treatment of psychiatric
disorders. Eur Neuropsychopharmacol. 2015;25: 599–656.
doi:10.1016/j.euroneuro.2015.01.016

63.   Oemisch M, Westendorff S, Oemisch M, Westendorff S, Everling S, Womelsdorf XT.
Interareal Spike-Train Correlations of Anterior Cingulate and Dorsal Prefrontal Cortex
during Attention Shifts Interareal Spike-Train Correlations of Anterior Cingulate and
Dorsal Prefrontal Cortex during Attention Shifts. J Neurosci. 2015;35: 13076–13089.
doi:10.1523/JNEUROSCI.1262-15.2015

64.   Paxinos G, Huang X, Toga A. The Rhesus Monkey Brain in Stereotaxic Coordinates.
2000.

65.   Hassani SA, Oemisch M, Balcarras M, Westendorff S, Ardid S, van der Meer MA, et al.
A computational psychiatry approach identifies how alpha-2A noradrenergic agonist
Guanfacine affects feature-based reinforcement learning in the macaque. Sci Rep. Nature
Publishing Group; 2017;7: 40606. doi:10.1038/srep40606

66.   Gómez-Ríos GA, Tascon M, Reyes-garcés N, Boyacı E. Quantitative analysis of biofluid
spots by coated blade spray mass spectrometry , a new approach to rapid screening. Sci
Rep. 2017;7: 1–7. doi:10.1038/s41598-017-16494-z

678
679

Fig 1. SPME sampling procedure, extraction time profiles and measurement locations.
(A) Dimensions of the SPME probe and the accompanying cannula. There was a 3000 μm buffer from the opening of the cannula and the start of the SPME coating. The SPME coating was 195 μm at the tip. The SPME coating made up the terminal 3000 μm of the entire probe and was placed in a cannula with an internal diameter of 200 μm and an external diameter of 270 μm. **(B)** Tungsten microelectrodes were lowered into the brain guided by 3 T magnetic resonance (MR) images in order to map the depth at which detectable spiking was observed matching expectations from the MR images. (C) SPME probes and their accompanying cannulas in all brain areas were then simultaneously lowered to 200 μm above the point of first observable firing in the target brain region and the SPME probes were lowered to expose only the 3000 μm SPME coating. After 20 minutes of extraction, the SPME probes were then retracted 3000 μm back into the cannula and the probes plus their accompanying cannulas were removed from the brain. The table inset into the figure provides the velocities and times of each probe and cannula as they transitioned between the extraction steps. (D) Extraction quantities of Glutamate in ng and (E) Dopamine in ng over time. The linear "kinetic" regime and "equilibrium" regime are labelled with the dynamic range being in between. (F**)** Anatomical locations of SPME sampling events in the right hemisphere of two Rhesus Macaques. Target brain regions are highlighted in red and include two cortical regions, prefrontal cortex (PFC) area 8 and premotor cortex (PreM) area 6 as well as a subcortical region: the head of the caudate.

Fig 2. Changes in the normalized concentrations of the observed neuromodulators across daily sampling sessions in each sampled brain region. The median concentration of (A) Glutamate (B) acetylcholine (C) choline and dopamine (D) with standard error of the median plotted for each sampling time and location. Data was normalization for each subject separately and for the respective neurochemical and area combinations. Sampling locations are represented by different lines: solid line for the prefrontal cortex, dashed line for the premotor cortex and dotted line for the caudate nucleus. Sampling events were 20 minutes in length and the time between the end of one sampling event to the start of the next sampling event was 40 minutes (STD 2 minutes) making the second sampling event 40 minutes and the third 100 minutes from the initial measurement. No significant change was observed for any neuromodulator in any brain region (Wisconsin rank sum test). Note that a single prefrontal choline data point was excluded in this analysis for being >4 standard deviations from the median (data point is present in **Fig 3a**).

Fig 3. SPME sampling of the macaque brain.
SPME probes extract from the cortex (PFC and PreM6) as well as the caudate of two adult macaques after 20 minutes of extraction within the respective brain region. Average of n= 12 probes for animal As and n= 9 probes for animal Ke. Four neuromodulators were reliably measured: glutamate, dopamine, acetylcholine, and choline (orange, green, purple, red respectively). All brain regions had high concentrations of glutamate and relatively much lower concentrations of acetylcholine and choline extracted (with the exception of animal Ke PreM6 lacking detectable acetylcholine). The only tested region to yield measurable dopamine was the caudate. The dashed blue line depicts the limit of detection (LOD).

A

3000 μm

3000 μm

195 μm

ID: 200 μm

OD: 270 μm

B

Gray matter

200 μm

C

i ii iii iv

	fibre	cannula	time
i	30 μm/s	30 μm/s	variable
ii	30 μm/s	-	200s
iii	30 μm/s	-	1200s* + 200s
iv	30 μm/s	30 μm/s	variable

*stationary: extraction period of 20 min

D

Glutamate

Equilibrium regime

Kinetic regime

Amount Extracted [ng]

Extraction time [min]

E

Dopamine

Equilibrium regime

Amount Extracted [ng]

Extraction time [min]

F

Macaque Brain Surface

PFC 8

PreM 6

Caudate head

Supplementary Methods

Chemicals, Reagents and Materials

The LC-MS-grade solvents methanol (MeOH), acetonitrile (ACN), isopropanol (IPA) and water, as well as the acetylcholinesterase inhibitor phenylmethylsulfonyl fluoride (PMSF) were purchased from Fisher Scientifc. Formic acid, polyacrylonitrile (PAN), N,N-dimethylformamide (DMF) and the standards of neurotransmitters: γ-aminobutyric acid (GABA), glutamic acid (Glu), acetylcholine (ACh), histamine (Hist), serotonin (5-HT), dopamine (DA) and choline (Cho) as well as their deuterated analogues were purchased from Millipore Sigma (Oakville, ON, Canada). Epinephrine (Epi), norepinephrine (NE) and their deuterated analogues were obtained from Cerilliant Corporation (Round Rock, TX, USA). Choline-D9, was purchased from Cambridge Isotope Laboratories (Tewksbury, MA, USA). The reagents used for synthesis of hydrophilic-lipophilic balance polymer particles functionalized with strong cation exchange groups, as well as compounds for preparation of PBS were purchased from Millipore-Sigma. The stainless steel wire (stainless steel grade AISI 304, 150 μm diameter) used for manufacturing of SPME probes was purchased from Unimed S.A. (Lausanne, Switzerland). The stainless steel tubing used as guiding cannulas (270 μm O.D.; 200±5 μm I.D.) was obtained from Vita Needle (Needham, MA, USA).

LC-MS/MS Analysis

On the day of analysis, the SPME probes were defrosted and desorbed into 40 μL of water/ACN/MeOH 40:30:30 (v/v/v) mixture containing 1 % of formic acid and a mixture of deuterated isotopologues of targeted neuromodulators (used as internal standards, IS) at 20 ng/mL. The desorption was carried out for 1 h with agitation at 1500 rpm. The extracts were injected into the LC-MS system for targeted neurotransmitter analysis within a few hours after desorption. All experiments were carried out using an Ultimate 3000RS HPLC system coupled to TSQ Quantiva triple quadrupole mass spectrometer (Thermo Fisher Scientific, San Jose, California, USA). Data processing was performed using Thermo software Xcalibur 4.0 and Trace Finder 4.1. For chromatographic separation of the target compounds, a modified method of what is previously reported (Cudjoe and Pawliszyn 2014) was used, adjusted to include more

targeted neuromodulators and their corresponding IS. A Kinetex® PFP LC column (100 x 2.1 mm, 1.7 µm; Phenomenex, Torrance, CA, USA) was held at 30°C with the mobile phase flow rate at 400 µL/min. The mobile phase A consisted of water/MeOH/ACN 90:5:5 (v/v/v) with 0.1 % formic acid, and mobile phase B was ACN/water 90:10 (v/v) with 0.1% of formic acid. The chromatographic gradient was applied starting from 100% B for 1 min and increasing the aqueous mobile phase A to 100% for 3 min with convex gradient function, held for 0.5 min and subsequent linear return to initial conditions and re-equilibration for 1 min, yielding total time of 6.5 min. The injection volume was 10 µL. MS/MS analysis was performed with electrospray ionization in positive mode under selected reaction monitoring conditions, with two MS/MS transitions for each neuromodulator (quantifier and qualifier) and one for each IS.

Quantitation of neuromodulators

Individual stock standard solutions of all targeted neuromodulators were prepared in methanol or water with 0.1% formic acid at a concentration of 1 mg/mL and stored at −80 °C for maximum of one month. In order to calculate the amounts of neuromodulators extracted by each probe, calibrator standards prepared in the same desorption solvent as real samples were analyzed in the same batch. This instrumental calibration curve was prepared in the range of 0.1-200 ng/mL by a serial dilution of the stock standard mixture of all compounds at 1µg/mL. The IS concentration was kept constant at 20 ng/mL in all calibrators, identically as in the real samples. The amounts extracted were calculated based on linear regression equation obtained from the analytical signal (the ratio of chromatographic peak areas of analytes and their corresponding IS) plotted against the concentration.

In order to calculate the concentrations of neuromodulators in brain, matrix-matched external calibration approach was used. The surrogate matrix consisted of 2% agar gel mixed with brain homogenate in the ratio 1:1 (*v/w*). The homogenized brain tissue was earlier incubated with 1 mM PMSF for 1 h at 37°C to prevent enzymatic digestion of acetylcholine in the calibrator samples. Due to several target compounds being present in brain homogenate at high concentrations (e.g. for glutamate and choline the "blank" brain homogenate matrix doesn't exist), their quantitation was based on signals of their deuterated isotopologues. The calibrator samples were prepared in the surrogate matrix with concentrations of neuromodulators ranging

from 5 to 3000 µg/mL for the isotopically labelled compounds or from 10 to 2000 ng/mL for the remaining compounds.

The extractions were carried out with SPME probes manufactured and pre-treated identically to the probes used for *in vivo* sampling and using the same 20 min extraction time and desorption conditions as for the real samples. The amounts of neuromodulators extracted from the calibrator samples were determined in the same way as described above and plotted against concentrations of calibrators. The resulting weighted linear regression equations were applied to the amounts of neuromodulators extracted from the *in vivo* samples, yielding values of concentrations of the compounds of interest in brain.

The limits of detection (LOD) were estimated as the levels corresponding to the signal to noise ratio of 3 and were calculated based on the signal of blank calibrator sample (considered as the noise).

Supplementary Results

Observed serotonin concentrations. Overall, serotonin concentrations were not observed as reliably as the other reported neurochemicals and therefore were excluded from the main analyses. However, in one of the animals, serotonin was observed near the LOD in the prefrontal cortex as well as the caudate in a subset of probes. Within the 9 probes placed in both areas, serotonin was detected in 22% of samples (2/9 probes) in the prefrontal cortex and 33% of samples (3/9 probes) in the caudate nucleus. Serotonin observations ranged from 149 ng/mL to 232 ng/mL with a median of 171 ng/mL (±18.5 ng/mL; standard error of the median) with a LOD of 100 ng/mL. The proximity of the measurements to the LOD suggest that the other probes likely collected concentrations of serotonin below the detection threshold. No serotonin observations were made in the second animal.

Supplementary Discussion

Future adjustments to the SPME neurochemical sensing. Several improvements to the protocol can be made in order to make SPME more informative. Using a more sensitive MS would reduce detection thresholds and increase sensitivity in detecting compounds not

successfully measured here such as GABA, serotonin or norepinephrine. This would be very impactful, especially in the case of Serotonin for reasons discussed above. Additionally, efforts could be made to improve the extracting phase synthesis and functionalization protocols and increase the extracting capabilities of SPME probes. Moreover, in order to decrease the MS background and interferences in the range of small molecules, derivatization strategy could be considered.

For GABA, the most promising strategy is post-desorption derivatization by reagent increasing the compound's hydrophobicity resulting in better MS signal such as benzoyl chloride (Wong 2016). Catecholamine and acetylcholine detection may be improved by optimizing the post-sampling analysis pipeline. Improvements can be made at several steps in order to better preserve and quantify catecholamines: (1) faster coupling of sampling to desorption (2) trying various antioxidant solutions and other preservatives for maintaining catecholamine integrity and (3) faster analysis of the desorbed analytes. Faster coupling of sampling to desorption could be achieved through a more stream-lined process of fibre placement and retraction involving a static cannula maintained through several sampling events. This mechanism would also allow for a faster replacement of SPME probes resulting in a shorter delay between consecutive measurements.

A preliminary experiment using spiked artificial cerebra-spinal fluid (aCSF) with very high, known concentrations of all target compounds yielded detectable concentrations of all target compounds indicating that the extracting phase of the SPME probe is capable of capturing all compounds of interest. Furthermore, simultaneously collected replicates displayed very little loss of collected compounds over several days when placed in a -80°C freezer (data not shown). Such findings neither support nor antagonize the suggested strategies for improving SPME yield and sensitivity in primates as detection properties within the aCSF do not seem to accurately reflect those observed in vivo in NHPs. This is likely contributed to by the dynamic and highly regulated nature of target compounds within the extracellular space of the brain and the dynamic equilibrium between the extracting phase of the SPME probes and the extracellular environment. Further tests of the possible differences between these two mediums and improvement strategies to SPME probes are subject for future studies.

Supplementary Figures

Fig S1. Experimental procedure from SPME probe fabrication to quantitation. Fabrication described the in-house procedure to prepare SPME probes. Preparation describes experimental set-up. Sampling describes the actual data collection process. LC/MS analysis describes the chemical quantitation of collected data samples.

Supplementary Tables

Table 1. Overview of in vivo neurochemical measurement methods

	PET imaging	Electro-chemistry	Micro-dialysis	Solid phase micro-extraction
Temporal Resolution	Minutes	Highest (millisecond range)	1-30 minutes (usually 20 minutes); dependent on MS sensitivity, target etc.	<5-30 minutes; dependent on MS sensitivity, coating thickness, target etc.
Spatial Resolution	Voxel	High; surface area may vary (relevant for enzyme based methods)	Diffusion based; surface area may vary	Diffusion based; surface area may vary
Sensitivity	Indirect measurement via competitive radiolabeled species	High	Depending on post-hoc methods (i.e. MS)	Depending on post-hoc methods (i.e. MS)
Neuro-active targets	A few at most	A few at most	Many	Many
Non-neuro-active targets	No	No	Yes, greater efficacy for hydrophilic compounds	Yes, greater potential efficacy for hydrophobic compounds
In vivo feasibility	Difficult in awake, behaving animal models; movement highly restricted	Good (low reliability in NHPs)	Good	Best
Cost	High	Requires special equipment	Requires special equipment	Easy to port to an acute micro-electrode setup; requires a chemistry core

Table S1. A comparison of methods capable of measuring single or multiple neurochemicals in vivo. Temporal resolution, spatial resolution, sensitivity, capability to measure neuro-active and non-neuro-active compounds, in vivo feasibility and cost. PET: positron emission tomography; NHP: non-human primate; MS: mass spectrometer.