Genomics evolutionary history and diagnostics of the *Alternaria alternata* species group including apple and Asian pear pathotypes

Andrew D. Armitage[1]*, Helen M. Cockerton[1], Surapareddy Sreenivasaprasad[2], James
Woodhall[3], Charles R. Lane[4], Richard J. Harrison[1] and John P. Clarkson[5]

[1] NIAB EMR, New Road, East Malling, Kent, ME19 6BJ, UK

[2] University of Bedfordshire, University Square, Luton, Bedfordshire, LU1 3JU, UK

[3] Parma Research and Extension Center, University of Idaho, Parma 83660, USA

[4] FERA Science Ltd, Sand Hutton, York, YO41 1LZ, UK

[5] University of Warwick, Wellesbourne, Warwick, CV35 9EF, UK

* Correspondence:
Andrew Armitage
Andrew.Armitage@EMR.ac.uk

Keywords: CDC, host-specific toxin, mating type, Dothideomycete, *Alternaria mali*, *Alternaria*
***gaisen*, nanopore.**

Abstract - 249

Introduction – 826

Methods – 2037

Results – 2291

Discussion - 1315

Manuscript Length – 6469

Abstract

The *Alternaria* section *alternaria* (*A. alternata* species group) represents a diverse group of saprotroph, human allergens and plant pathogens. *Alternaria* taxonomy has benefited from recent phylogenetic revision but the basis of differentiation between major phylogenetic clades within the group is not yet understood. Furthermore, genomic resources have been limited for the study of host-specific pathotypes. We report near complete genomes of the apple and Asian pear pathotypes as well as draft assemblies for a further 10 isolates representing *Alternaria tenuissima* and *Alternaria arborescens* lineages. These assemblies provide the first insights into differentiation of these taxa as well as allowing the description of effector and non-effector profiles of apple and pear conditionally dispensable chromosomes (CDCs). We define the phylogenetic relationship between the isolates sequenced in this study and a further 23 *Alternaria* spp. based on available genomes. We determine which of these genomes represent MAT1-1-1 or MAT1-2-1 idiomorphs and designate host-specific pathotypes. We show for the first time that the apple pathotype is polyphyletic, present in both the *A. arborescens* and *A. tenuissima* lineages. Furthermore, we profile a wider set of 89 isolates for both mating type idiomorphs and toxin gene markers. Mating-type distribution indicated that gene flow has occurred since the formation of *A. tenuissima* and *A. arborescens* lineages. We also developed primers designed to *AMT14,* a gene from the apple pathotype toxin gene cluster with homologs in all tested pathotypes. These primers allow identification and differentiation of apple, pear and strawberry pathotypes, providing new tools for pathogen diagnostics.

1   Introduction

Species within the genus *Alternaria* encompass a range of lifestyles, acting as saprotroph,

opportunistic pathogens and host-adapted plant pathogens (Thomma 2003). Large spored species

include *A. solani*, a major pathogen of potato, whereas small spored taxa include the *Alternaria*

alternata species group (*Alternaria* sect. *alternaria*), which are found ubiquitously in the

environment acting as saprotroph and opportunistic necrotrophs. This species group is responsible for

opportunistic human infections and a range of host adapted plant diseases.

Taxonomy within this presumed asexual genus has been subject to recent revision (Lawrence et al.

2013). Large spored species can be clearly resolved by standard phylogenetic markers such as ITS

and are supported by morphological characters. However, small spored species within the *A.*

alternata species group overlap in morphological characters, possess the same ITS haplotype

(Kusaba and Tsuge, 1995a) and show low variation in other commonly used barcoding markers

(Woudenberg et al. 2015; Lawrence et al. 2013; Armitage et al. 2015). Highly variable phylogenetic

markers have provided resolution between groups of isolates that possess morphological patterns

typical of descriptions for *A. gaisen*, *A. tenuissima* and *A. arborescens* (Armitage et al. 2015).

However, taxonomy of the species group is further complicated by designation of isolates as

pathotypes, each able to produce polyketide host-selective toxins (HST) adapted to apple, Asian pear,

tangerine, citrus, rough lemon or tomato (Tsuge et al. 2013). Genes involved in the production of

these HSTs are located on conditionally dispensable chromosomes (CDCs) (Hatta et al. 2002). CDCs

have been estimated to be 1.05 Mb in the strawberry pathotype (Hatta et al. 2002), 1.1 - 1.7 Mb in the

apple pathotype (Johnson et al. 2001), 1.1 - 1.9 Mb in the tangerine pathotype (Masunaka et al. 2000;

Masunaka et al. 2005) and 4.1 Mb in the pear pathotype (Tanaka et al. 1999; Tanaka and Tsuge

2000). These CDCs are understood to have been acquired through horizontal gene transfer and as

such, the evolutionary history of CDCs may be distinct from the core genome.

The polyketide synthase genes responsible for the production of the six HSTs are present in clusters.

Some genes within these clusters are conserved between pathotypes (Miyamoto et al. 2009; Hatta et

al. 2002), while genes are also present within these clusters that are unique to particular pathotypes

(Ajiro et al. 2010; Miyamoto et al. 2010). This is reflected in structural similarities between the pear

(AKT) and strawberry (AFT) and tangerine (ACTT) toxins with each containing a 9,10-epoxy-8-

hydroxy-9-methyl-decatrienoic acid moiety. In contrast, the toxin produced by the apple pathotype

(AMT) does not contain this moiety and is primarily cyclic in structure (Tsuge et al. 2013).

Studies making use of bacterial artificial chromosomes (BAC) have led to the sequencing of toxin

gene cluster regions from three apple pathotype isolates (Genbank accessions: AB525198,

AB525199, AB525200; unpublished). These sequences are 100-130 Kb in size and contain 17 genes

that are considered to be involved in synthesis of the AMT apple toxin (Harimoto et al. 2007). *AMT1*,

AMT2, *AMT3* and *AMT4* have been demonstrated to be involved in AMT synthesis, as gene

disruption experiments have led to loss of toxin production and pathogenicity (Johnson et al. 2000;

Harimoto et al. 2007; Harimoto et al. 2008). However, experimental evidence has not been provided

to show that the remaining 13 *AMT* genes have a role in toxin production. Four genes present in the

CDC for the pear pathotype have been identified and have been named *AKT1*, *AKT2*, *AKT3*, *AKTR-1*

(Tanaka et al. 1999; Tanaka & Tsuge 2000; Tsuge et al. 2013) and a further two genes (*AKT4*,

AKTS1) have been reported (Tsuge et al. 2013).

The toxicity of a HST is not restricted to the designated host for that pathotype. All or some of the

derivatives of a toxin may induce necrosis on "non-target" host leaves. For example, AMT from the

apple pathotype can induce necrosis on the leaves of Asian pear (Kohmoto et al. 1976). Therefore

non-host resistance may be triggered by recognition of non-HST avirulence genes.

Alternaria spp. are of phytosanitary importance, with apple and pear pathotypes subject to quarantine

regulations in Europe under Annex IIAI of Directive 2000/29/EC as *Alternaria alternata* (non-

European pathogenic isolates). As such, rapid and accurate diagnostics are required for identification.

Where genes on essential chromosomes can be identified that phylogenetically resolve taxa, then

these can be used for identification of quarantine pathogens (Bonants et al. 2010; Quaedvlieg et al.

2012). Regulation and management strategies also need to consider the potential for genetic

exchange between species. The *Alternaria* sect. *alternaria* are presumed asexual but evidence has

been presented for either the presence of sexuality or a recent sexual past. Sexuality or parasexuality

provides a mechanism for reshuffling the core genome associated with CDCs of a pathotype. It is

currently unknown whether pathotype identification can be based on sequencing of phylogenetic loci,

or whether the use of CDC-specific primers is more appropriate. This is of particular importance for

the apple and Asian pear pathotypes due the phytosanitary risk posed by their potential establishment

and spread in Europe.

2   Methods

Twelve genomes were sequenced, selected from a collection of isolates whose phylogenetic identity

was determined in previous work (Armitage et al. 2015). These twelve isolates were three *A.*

arborescens clade isolates (*FERA 675*, *RGR 97.0013* and *RGR 97.0016*), four *A. tenuissima* clade

isolates (*FERA 648*, *FERA 1082*, *FERA 1164* and *FERA 24350*), three *A. tenuissima* clade apple

pathotype isolates (*FERA 635*, *FERA 743*, *FERA 1166* and *FERA 1177*) and one *A. gaisen* clade pear

pathotype isolate (*FERA 650*).

2.1   DNA and RNA extraction and sequencing

Apple pathotype isolate *FERA 1166* and Asian pear pathotype isolate *FERA 650* were sequenced

using both Illumina and nanopore MinION sequencing technologies and the remaining ten isolates

were sequenced using Illumina sequencing technology. For both illumina and MinION sequencing,

DNA extraction was performed on freeze dried mycelium grown in PDB for 14 days.

High molecular weight DNA was extracted for MinION sequencing using the protocol of

Schwessinger & McDonald (2017), scaled down to a starting volume of 2 ml. This was followed by

phenol-chloroform purification and size selection to a minimum of 30 Kb using a Blue Pippin. The

resulting product was concentrated using ampure beads before library preparation was performed

using a Rapid Barcoding Sequencing Kit (SQK-RBK001) modified through exclusion of LLB beads.

Sequencing was performed on a Oxford Nanopore GridION generating an additional 39 and 44 times

coverage of sequence data for isolates *FERA 1166* and *FERA 650* respectively.

gDNA for illumina sequencing of isolate *FERA 1166* was extracted using a modified CTAB protocol

(Li *et al.*, 1994). gDNA for illumina sequencing of the eleven other isolates was extracted using a

Genelute Plant DNA Miniprep Kit (Sigma) using the manufacturers protocol with the following

modifications: the volume of lysis solutions (PartA and PartB) were doubled; an RNase digestion

step was performed as suggested in the manufacturer's protocol; twice the volume of precipitation

solution was added; elution was performed using elution buffer EB (Qiagen). A 200 bp genomic

library was prepared for isolate *FERA 1166* using a TrueSeq protocol (TrueSeq Kit, Illumina) and

sequenced using 76 bp paired-end reads on an Illumina GA2 Genome Analyser. Genomic libraries

were prepared for the other eleven isolates using a Nextera Sample Preparation Kit (Illumina) and

libraries sequenced using a MiSeq Benchtop Analyser (Illumina) using 250 bp, paired end reads.

RNAseq was performed to aid training of gene models. mRNA was extracted from isolates *FERA*

1166 and *FERA 650* grown in full strength PDB, 1% PDB, Potato Carrot Broth (PCB) and V8 juice

broth (V8B). The protocol for making PCB and V8B was as described in (Simmons, 2007)for

making Potato Carrot Agar and V8 juice agar, with the exception that agar was not added to the

recipe. Cultures were grown in conical flasks containing 250 mls of each liquid medium for 14 days.

mRNA extraction was performed on freeze dried mycelium using the RNeasy Plant RNA extraction

Kit (Qiagen). Concentration and quality of mRNA samples were assessed using a Bioanalyzer

(Agilent Technologies). mRNA from the sample grown in 1% PDB for isolate *FERA 650* showed

evidence of degradation and was not used further. Samples were pooled from growth mediums for

each isolate and 200 bp cDNA libraries prepared using a TrueSeq Kit (Illumina). These libraries were

sequenced in multiplex on a MiSeq (Illumina) using 200 bp paired end reads.

2.2   Genome assembly and annotation

De-novo genome assembly was performed for all 12 isolates. Assembly for isolate *FERA 650* was

generated using SMARTdenovo (https://github.com/ruanjue/smartdenovo, February 26, 2017 github

commit), whereas assembly for isolate *FERA 1166* was generated by merging a SMARTdenovo

assembly with a MinION-Illumina hybrid SPAdes v3.9.0 assembly using quickmerge v0,2 (Antipov

et al. 2016, Chakraborty et al. 2016). Prior to assembly, adapters were removed from MinION reads

using Porechop v0.1.0 and reads were further trimmed and corrected using Canu v1.6 (Ruan 2016;

Koren et al. 2017). Following initial assembly, contigs were corrected using MinION reads through

ten rounds of Racon (May 29, 2017 github commit) correction (Vaser et al. 2017)and one round of

correction using MinION signal information with nanopolish (v0.9.0) (Loman et al. 2015). Final

correction was performed through ten rounds of Pilon v1.17 (Walker et al. 2014) using Illumina

sequence data. Assemblies for the ten isolates with Illumina-only data were generated using SPAdes

v.3.9.0 (Bankevich et al. 2012). Assembly quality statistics were summarized using Quast v.4.5

(Gurevich et al. 2013). Single copy core Ascomycete genes were identified within the assembly using

BUSCO v3 and used to assess assembly completeness (Simão et al. 2015). RepeatModeler,

RepeatMasker and TransposonPSI were used to identify repetitive and low complexity regions

(http://www.repeatmasker.org, http://transposonpsi.sourceforge.net). Visualisation of whole genome

alignments between *FERA 1166* and *FERA 650* was performed using circos v0.6 (Krzywinski et al.

2009), following whole genome alignment using the nucmer tool as part of the MUMmer package

v4.0 (Marçais *et al.* 2018).

Gene prediction was performed on softmasked genomes using Braker1 v.2 (Hoff et al. 2016), a

pipeline for automated training and gene prediction of AUGUSTUS 3.1 (Stanke & Morgenstern

2005). Additional gene models were called in intergenic regions using CodingQuarry v.2 (Testa et al.

2015). Braker1 was run using the "fungal" flag and CodingQuarry was run using the "pathogen" flag.

RNAseq data generated from *FERA 1166* and *FERA 650* were aligned to each genome using STAR

v2.5.3a (Dobin et al. 2013), and used in the training of Braker1 and CodingQuarry gene models.

Orthology was identified between the 12 predicted proteomes using OrthoMCL v.2.0.9 (Li et al.

2003) with an inflation value of 5. Orthology was identified between the 12 predicted proteomes

using OrthoMCL v.2.0.9 (Li et al. 2003) with an inflation value of 5.

Draft functional genome annotations were determined for gene models using InterProScan-5.18-57.0

(Jones et al. 2014) and through identifying homology (BLASTP, E-value > $1x10^{-100}$) between

predicted proteins and those contained in the March 2018 release of the SwissProt database (Bairoch

and Apweiler 2000). Putative secreted proteins were identified through prediction of signal peptides

using SignalP v.4.1 and removing those predicted to contain transmembrane domains using

TMHMM v.2.0 (Käll et al. 2004; Krogh et al. 2001). Additional programs were used to provide

evidence of effectors and pathogenicity factors. EffectorP v1.0 was used to screen secreted proteins

for characteristics of length, net charge and amino acid content typical of fungal effectors

(Sperschneider et al. 2016). Secreted proteins were also screened for carbohydrate active enzymes

using HMMER3 (Mistry et al. 2013) and HMM models from the dbCAN database (Huang et al.

2018). DNA binding domains associated with transcription factors (Shelest 2017) were identified

along with two additional fungal-specific transcription factors domains (IPR007219 and IPR021858).

Annotated assemblies were submitted as Whole Genome Shotgun projects to DDBJ/ENA/Genbank

(Table 1).

2.3   Phylogenetics

BUSCO hits of single copy core ascomycete genes to assemblies were extracted and retained if a

single hit was found in all of the 12 sequenced genomes and 23 publically available *Alternaria* spp.

genomes from the Alternaria genomes database (Dang et al. 2015). Sequences from the resulting hits

of 500 loci were aligned using MAFFT v6.864b (Katoh and Standley 2013), before alignments were

trimmed using trimAl v1.4.1 (Capella-Gutiérrez et al. 2009), and trees calculated for each locus using

RAxML v8.1.17 (Liu et al. 2011). The most parsimonious tree from each RAxML run was used to

determine a single consensus phylogeny of the 500 loci using ASTRAL v5.6.1 (Zhang et al. 2018).

The resulting tree was visualised using the R package GGtree v1.12.4 (Yu et al. 2016).

2.4   CDC identification

Contigs unique to apple and pear pathotypes were identified through read alignment to assembled

genomes. Read alignment was performed using Bowtie2 (Langmead and Salzberg 2012), returning a

single best alignment for each paired read. Read coverage was quantified from these alignments

using Samtools (Li et al. 2009).

2.5   Toxin-synthesis genes in *Alternaria* genomes

Sequence data for 40 genes located in *A. alternata* HST gene clusters were downloaded from

Genbank. BLASTn searches were performed for all 40 gene sequences against one another to

identify homology between these sequences. Genes were considered homologous where they had >

70 % identical sequences over the entire query length, and an e-value of 1×10^{-30}. tBLASTx was used

to search for the presence of these genes in assemblies.

2.6   Signatures of genetic exchange

Mating type idiomorphs present in publicly available genomes were identified using BLASTn

searches. A wider assessment within 89 characterised *Alternaria* isolates (Armitage et al. 2015) was

undertaken using specific PCR primers (Arie et al. 2000). PCR primers (AAM1-3: 5'-

TCCCAAACTCGCAGTGGCAAG-3'; AAM1-3: 5'-GATTACTCTTCTCCGCAGTG-3; M2F: 5'-

AAGGCTCCTCGACCGATGAA-3;   M2R: 5'-CTGGGAGTATACTTGTAGTC-3) were run in

multiplex with PCR reaction mixtures consisting of 10 µl redtaq (REDTaq ReadyMix PCR Reaction

Mix, Sigma-Aldrich), 2 µl DNA, 1 µl of each primer (20 µM), and 4 µl purified water (Sigma-

Aldrich). PCR reaction conditions comprised of an initial 60 second denaturing step at 94°C followed

by 30 cycles of a melting step of 94°C for 30 seconds, an annealing step at 57°C for 30 seconds, and

an extension step at 72°C for 60 seconds, these cycles were followed by a final extension step at

72°C for 420 seconds. MAT1-1-1 or MAT1-2-1 idiomorphs were determined through presence of a

271 or 576 bp product following gel electrophoresis, respectively.

2.7   PCR screens for apple and pear toxin-synthesis genes

A set of 90 previously characterised isolates was used to further investigate the distribution of

pathotypes throughout the *A. alternata* species group. PCR primers were designed for the

amplification of three genes (*AMT4*, *AMT14*, *AKT3*) located within CDC gene clusters involved in

toxin synthesis. Primers for *AMT4* were designed to amplify apple pathotype isolates, *AKT3* to

amplify pear pathotype isolates and *AMT14* to identify both apple and pear pathotype isolates. These

primers were then used to screen isolates for the presence of these genes in 30 cycles of PCR using

0.25 ul Dream taq, 1 ul of 10x PCR buffer, 1 ul of dNTPs, 1 ul of gDNA, 1 ul of each primer (5

µM), and 4.75 ul purified water (Sigma-Aldrich). PCR products were visualised using gel

electrophoresis and amplicon identity confirmed through Sanger sequencing. Primers AMT4-EMR-F

(5'-CTCGACGACGGTTTGGAGAA-3) and AMT4-EMR-R (5'-TTCCTTCGCATCAATGCCCT-

3) were used for amplification of AMT4. Primers AKT3-EMR-F (5'-

GCAATGGACGCAGACGATTC-3) and AKT3-EMR-R (5'-CTTGGAAGCCAGGCCAACTA-3)

were used for amplification of *AKT3*. Primers AMT14-EMR-F (5'-TTTCTGCAACGGCGKCGCTT-

3) and AMT14-EMR-R (5'-TGAGGAGTYAGACCRGRCGC-3) were used for amplification of

AMT14. PCR reaction conditions were the same as described above for mating type loci, but with

annealing performed at 66°C for all primer pairs.

2.8   Virulence assay

Pathogenicity assays were performed on apple *cv.* Spartan and *cv.* Bramley's seedling to determine

differences in isolate virulence between *A. tenuissima* isolates possessing the apple pathotype CDC

(*FERA 635*, *FERA 743* or *FERA 1166*) and non-pathotype isolates lacking the CDC (*FERA 648*,

FERA 1082 or *FERA 1164*). Detailed pathogenicity assay methods are provided in supplementary

information 5. Briefly, leaves were inoculated with 10 ul of 1×10^{5} spore suspensions at six points

and the number of leaf spots counted at 14 days post inoculation. One isolate was infected per leaf,

with 10 replicates per cultivar. Binomial regression using a generalised linear model (GLM) was

used to analyse the number of resulting lesions per leaf.

Unfolded adult apple leaves, less than 10 cm in length were cut from young (less than 12 months old)

apple *cv.* Spartan trees or *cv.* Bramley's seedling trees. These were quality-checked to ensure that

they were healthy and free from disease. Leaves were grouped by similar size and age and organised

into ten experimental replicates of nine leaves. Leaves placed in clear plastic containers, with the

abaxial leaf surface facing upwards. The base of these boxes was lined with two sheets of paper

towel, and wetted with 50 mls of sterile distilled water (SDW). The cultivars were assessed in two

independent experiments.

Spore suspensions were made by growing *A. alternata* isolates on 1% PDA plates for four weeks at

23 °C before flooding the plate with 2 mls of SDW, scraping the plate with a disposable L-shaped

spreader. Each leaf was inoculated with 10 ml of 1×10^{5} spores.ml^{-1} *A. alternata* spore suspension or

10 ml of sterile-distilled water at six points on the abaxial leaf surface. Of the nine leaves in each

box, three leaves were inoculated with a spore suspensions from isolates carrying apple pathotype

CDC, three leaves were inoculated with non-pathotype isolates lacking the CDC, and three leaves

were inoculated with SDW. Following inoculation, each container was sealed and placed in plastic

bags to prevent moisture loss. Boxes were then kept at 23 °C with a 12 hr. light / 12 hr. dark cycle.

3   Results

3.1   Generation of near-complete genomes for the apple and pear pathotype using MinION
**   sequencing**

Assemblies using nanopore long-read sequence data for the apple pathotype isolate *FERA 1166*, and

pear pathotype isolate *FERA 650* were highly contiguous, with the former totaling 35.7 Mb in 22

contigs and the latter totaling 34.3 Mb in 27 contigs (Table 2). Whole genome alignments of these

assemblies to the 10 chromosomes of *A. solani* showed an overall macrosynteny between genomes

(Fig. 1), but with structural rearrangement of apple pathotype chromosomes in comparison to *A.*

solani chromosomes 1 and 10. The Asian pear pathotype had distinct structural rearrangements in

comparison to *A. solani*, chromosomes 1 and 2 (Fig. 1). Scaffolded contigs of *FERA 1166* spanned

the entire length of *A. solani* chromosomes 2, 3, 6, 8 and 9 and chromosomes 4, 5, 6 and 10 for *FERA*

650 (Fig. 1). Interestingly, sites of major structural rearrangements within *A. solani* chromosome 1

were flanked by telomere-like TTAGGG sequences.

Genome assembly of 10 Illumina sequenced isolates yielded assemblies of a similar total size to

MinION assemblies (33.9 - 36.1 Mb) but fragmented into 167-912 contigs. Assembled genomes were

repeat sparse, with 1.41 - 2.83 % of genomes repeat masked (Table 2). Genome assemblies of *A.*

arborescens isolates (33.8 - 33.9 Mb), were of similar total size to non-pathotype *A. tenuissima*

isolates and had similar repetitive content (2.51 – 2.83 and 1.41 – 2.83 % respectively). Despite this,

identification of transposon families in both genomes showed expansion of DDE ($T_{5df} = 5.36$, P >

0.01) and gypsy ($T_{5df} = 6.35$, P > 0.01,) families in *A. arborescens* genomes (Fig. 2).

3.2   Phylogeny of sequenced isolates

The relationship between the 12 sequenced isolates and 23 *Alternaria* spp. with publically available

genomes was investigated through phylogenetic analysis of 500 shared core ascomycete genes. *A.*

pori and *A. destruens* genomes were excluded from the analysis due to low numbers of complete

single copy ascomycete genes being found in their assemblies (Supp. information 1). The 12

sequenced isolates were distributed throughout *A. gaisen*, *A. tenuissima* and *A. arborescens* clades

(Fig. 3). The resulting phylogeny (Fig. 3), formed the basis for later assessment of CDC presence and

mating type distribution among newly sequenced and publicly available genomes, as discussed

below.

3.3   Gene and effector identification

Gene prediction resulted in 12757 – 13733 genes from the our assemblies (Table 2), with

significantly more genes observed in the apple pathotype isolates than in *A. tenuissima* clade non-

pathotype isolates ($P > 0.01$, $F_{2,8df} = 51.19$). BUSCO analysis identified that gene models included

over 97% of the single copy conserved ascomycete genes, indicating well trained gene models. Apple

pathotype isolates possessed greater numbers of secondary metabolite clusters ($P > 0.01$, $F_{2,8df} =$

8.96) and secreted genes ($P > 0.01$, $F_{2,8df} = 44.21$) than non-pathotype *A. tenuissima* isolates,

indicating that CDCs contain additional secreted effectors. Non-pathotype *A. tenuissima* clade

isolates were found to possess greater numbers of genes encoding secreted proteins than *A.*

arborescens isolates ($P > 0.01$, $F_{2,8df} = 44.21$), including secreted CAZYmes ($P > 0.01$, $F_{2,8df} = 9.83$).

The basis of differentiation between these taxa was investigated further.

3.4   Genomic differences between *A. tenuissima* and *A. arborescens* clades

Orthology analysis was performed upon the combined set of 158,280 total proteins from the 12

sequenced isolates. In total, 99.2 % of proteins clustered into 14,187 orthogroups. Of these, 10,669

orthogroups were shared between all isolates, with 10,016 consisting of a single gene from each

isolate. This analysis allowed the identification of 239 orthogroups that were either unique to *A.*

arborescens isolates or expanded in comparison to non-pathotype *A. tenuissima* isolates.

Expanded and unique genes to *A. arborescens* isolates was further investigated using *FERA 675*

(Supp. information 2). Genes involved in reproductive isolation were in this set, including 21 of the

148 heterokayon incompatibility (HET) loci from *FERA 675*. CAZymes were also identified within

this set, three of which showed presence of chitin binding activity and the other three having roles in

xylan or pectin degradation. In total, 25 genes encoding secreted proteins were within this set,

secreted proteins with pathogenicity-associated functional annotations included a lipase, a

chloroperoxidase, an aerolysin-like toxin, a serine protease and an aspartic peptidase. A further six

secreted genes had an effector-like structure by EffectorP but no further functional annotations.

Furthermore, one gene from this set was predicted to encode a fungal-specific transcription factor
unique to *A. arborescens* isolates.

Further to the identification of genes unique or expanded in *A. arborescens*, 220 orthogroups were
identified as unique or expanded in the *A. tenuissima*. These orthogroups were further investigated
using isolate *FERA 648* (Supp. information 2). This set also contained genes involved in reproductive
isolation, including nine of the 153 from *FERA 648*. CAZymes within the set included two chitin
binding proteins, indicating a divergence of LysM effectors between *A. tenuissima* and *A.*
arborescens lineages. The five additional CAZymes in this set represented distinct families from
those expanded/unique in *A. arborescens*, including carboxylesterases, chitooligosaccharide oxidase
and sialidase. In total, 18 proteins from this set were predicted as secreted, including proteins with
cupin protein domains, leucine rich-repeats, astacin family peptidase domains and with four predicted
to have effector-like structures but no further annotations. *A. tenuissima* isolates had their own
complement of transcription factors, represented by four genes within this set.

3.5   Identification of CDC contigs and assessment of copy number

Alignment of Illumina reads to the apple and Asian pear pathotype MinION reference assemblies
identified variable presence of some contigs, identifying these as contigs representing CDCs (CDC
contigs). Six contigs totaling 1.87 Mb were designated as CDCs in the apple pathotype reference
(Table 3) and four contigs totaling 1.47 Mb designated as CDCs in the pear pathotype reference
(Table 4). Two *A. tenuissima* clade non-pathotype isolates (*FERA 1082*, *FERA 1164*) were noted to
possess apple pathotype CDC contigs 14 and 19 totaling 0.78 Mb (Table 3).

Read alignments showed that CDC contigs were present in multiple copies within *A. alternata*
pathotype isolates. *FERA 1166* Illumina reads aligned to its own assembly showed two-fold coverage
over contigs 14, 15, 20 and 21 in comparison to core contigs (Table 3). This was more pronounced in
isolate *FERA 1177* that had between two- and eight-fold coverage of these contigs. The same was
observed in pear pathotype CDC regions, with contigs 14 and 24 in isolate *FERA 650* showing two-
fold coverage from Illumina reads in comparison to core contigs (Table 4).

3.6   Toxin gene clusters are present on multiple CDC contigs

Homologs to 15 of the 17 AMT cluster genes were located on contigs 20 and 21 in the apple
pathotype reference genome (e-value $< 1\times10^{-30}$, $> 70\%$ query alignment), confirming them as CDC-
regions (Table 5). Of the remaining two genes, *AMT11* had low-confidence BLAST homologs on

contigs 18 and 21 (e-value $< 1\text{x}10^{-30}$) whereas the best BLAST hit of *AMT15* was located on contig

18 (e-value $< 1\text{x}10^{-30}$). Duplication of toxin gene regions was observed between CDC contigs, with

contig 20 carrying homologs to 16 toxin genes, but with contig 21 also carrying the *AMT1* to *AMT12*

section of the cluster (Table 5). The three other apple pathotype isolates (*FERA 635*, *FERA 743* and

FERA 1177) also showed presence of 15 of the 17 AMT genes (e-value $< 1\text{x}10^{-30}$, $> 70\%$ query

alignment), and with some AMT genes present in multiple copies within the genome indicating that

the AMT toxin region has also been duplicated in these isolates.

The Asian pear pathotype was also found to carry toxin gene clusters in multiple copies, with

homologs to the four AKT cluster genes present on contig 14 of the *FERA 650* assembly (e-value $<$

$1\text{x}10^{-30}$, $> 70\%$ query alignment), with three of these also present on contig 24 (e-value $< 1\text{x}10^{-30}$,

two with $> 70\%$ query alignment). BLAST hit results from AKT genes were supported by their

homologs from strawberry and tangerine pathotypes also found in these regions (Table 5). The pear

pathotype genome was also found to contain additional homologs from apple (*AMT14*), strawberry

(*AFT9-1*, *AFT10-1*, *AFT11-1* and *AFT12-1*) and citrus (*ACTT5* and *ACTT6*) located on CDC contigs

14 and 24 (Table 5).

3.7   CDCs carry effectors alongside secondary metabolites

A total of 624 proteins were encoded on the six contigs designated as CDCs in the reference apple

pathotype genome, with 502 proteins encoded on the four Asian pear pathotype CDC contigs (Supp.

information 3). We further investigated the gene complements of these regions.

Approximately a quarter of gene models on apple pathotype CDC contigs were involved in

secondary metabolism, with 153 genes present in six secondary metabolite gene clusters. This

included AMT toxin gene homologs on contigs 20 and 21, which were located within NRPS

secondary metabolite gene clusters. Three other secondary metabolite clusters were located on CDC

contigs with two of these involved in the production of T1PKS secondary metabolites and the third

with unknown function. A further two secondary metabolite clusters were located on contig 14

shared with two non-pathotype isolates, one of which is involved in the production of a T1PKS. The

pear pathotype also carried 153 genes in secondary metabolite gene clusters. These 30 % of CDC

genes were located in four clusters, with the AKT toxin genes in T1PKS clusters of contigs 14 and

24. A second cluster was present on contig 14 with unknown function and a T1PKS cluster was

present on contig 16.

Approximately 5 % of the genes on apple CDC contigs encoded secreted proteins, with 32 in isolate

*FERA 116*6 many of which had potential effector functions with six designated as CAZymes and 12

testing positive by EffectorP. Similarly, a total of 41 secreted proteins were predicted on the CDC

regions of the Asian pear pathotype, with eight of these designated as secreted CAZymes and 13

testing positive by EffectorP. Further investigation into the 32 secreted proteins from the apple

pathotype identified three CAZYmes from the chitin-active AA11 family, two from the cellulose-

active GH61 family and one cellulose-active GH3 family protein. Six of the 13 EffectorP proteins

also had domains identifiable by interproscan: four carried NTF2-like domains, which are envelope

proteins facilitating protein transport into the nucleus; one was a fungal hydrophobin protein; one

was a member of an panther superfamily PTHR40845 that shares structural similarity with proteins

from the plant pathogens *Phaeospharia nodorum*, *Sclerotinia sclerotiorum* and *Ustilago maydis*. Of

the 38 secreted proteins identified from the pear pathotype, two CAZYmes were also identified from

the chitin-active AA11 family, two from the AA3 family with single proteins from GH5, CBM67 and

AA7 families. Ten of the twelve secreted EffectorP proteins had no functional information as

predicted by interproscan, with the other two identified as carrying WSC domains IPR002889, which

are cysteine-rich domains involved carbohydrate binding. CDCs may also play important roles in

transcriptional regulation with 29 putative transcription factors identified in the apple pathotype CDC

contigs(4.6 % CDC genes) and 35 identified in pear pathotype CDC contigs (7.0 % CDC genes).

3.8   Polyphyletic distribution of apple and tangerine pathotypes

The evolutionary relationship between *A. alternata* pathotypes sequenced in this study and publicly

available genomes was analyzed by the core gene phylogeny (Fig. 2). We identified four isolates as

tangerine pathotypes (Z7, BMP2343, BMP2327, BMP3436) two as tomato pathotypes (BMP0308,

EGS39-128), one Asian pear pathotype (MBP2338), one rough-lemon (BMP2335) and two apple

pathotypes (BMP3063, BMP3064) through searches for genes from HST-gene clusters (Supp.

information 4). When plotted on the genome phylogeny, we found the apple and tangerine pathotypes

to be polyphyletic (Fig. 2). Five of the six sequenced apple pathotype isolates were located in the *A.*

tenuissima clade and one in the *A. arborescens* clade, whereas the tangerine pathotype was present in

both the *A. tenuissima* clade and in the *A. tangelonis / A. longipes* clade.

3.9   Molecular tools for identification of apple, pear and strawberry pathotypes

PCR primers for three loci *(AMT4, AKT3* and *AMT14)* were designed to identify the distribution of

pathotypic isolates through the *A. alternata* species group and were screened against a set of 89

previously characterised isolates (Fig. 4). Five isolates tested positive for the presence of *AMT4*, each

of which was from the *A. tenuissima* clade *(FERA 635, FERA 743, FERA 1166, FERA 1177)*. Five

isolates tested positive for the presence of *AKT3*, including the three isolates from Asian pear in the

A. gaisen clade and a further two isolates from the *A. tenuissima* clade that were from strawberry.

Sequencing of the *AKT3* amplicons from the two isolates *ex.* strawberry identified them as the *AFT3-*

2 ortholog of *AKT3*, showing that these isolates were strawberry pathotypes rather than pear

pathotypes. Sequencing of PCR products from the other isolates confirmed them to be apple or pear

pathotypes as expected. All of the isolates testing positive for *AMT4* or *AKT3* also tested positive for

AMT14, indicating its suitability as a target gene for identification of a range of pathotypes.

Presence of apple pathotype CDCs was confirmed to be associated with pathogenicity through

detached apple leaf assays. Apple pathotype isolates showed significantly greater numbers of necrotic

lesions when inoculated onto *cv.* Spartan ($F_{72df} = 100.64$) and *cv.* Bramley's Seedling ($F_{72df} = 69.64$)

leaves than non-pathotype *A. tenuissima* isolates (Fig. 5).

3.10   Signatures of genetic exchange

Of the 12 sequenced isolates, BLAST searches identified five as carrying the MAT1-1-1 idiomorph

and seven as carrying MAT1-2-1. Both idiomorphs showed distribution throughout the *A. alternata*

genome phylogeny (Fig. 2). These results were supported by PCR assays identifying the mating type

of 89 previously characterised isolates (Fig. 4). Idiomorphs did not deviate from a 1:1 ratio within *A.*

tenuissima (21 MAT1-1-1 : 23 MAT1-2-1; χ2 = 0.09, 1df, P>0.05) or *A. arborescens* clades (18

MAT1-1-1 : 24 MAT1-2-1; χ2 = 0.86; 1df; P>0.05), as expected under a random mating population.

All three of the *A. gaisen* clade isolates carried the MAT1-2-1 idiomorph.

4   Discussion

This work builds upon the current genomic resources available for *Alternaria*, including the *A.*

brassicicola and *A. solani* genomes (Belmas et al. 2018; Wolters et al. 2018), *A. alternata* from onion

(Bihon et al. 2016) the additional 25 *Alternaria spp.* genomes available on the *Alternaria* Genomes

Database (Dang et al. 2015) as well as recent genomes for other pathotype and non-pathotype *A.*

alternata (Hou et al. 2016; Wang et al. 2016; Nguyen et al. 2016). Of the previously sequenced

genomes, *A. solani*, the citrus pathotype and a non-pathotype *A. alternata* isolate have benefitted

from long read sequencing technology with each comprising less than 30 contigs (Wolters et al.

2018; Wang et al. 2016; Nguyen et al. 2016). Total genome sizes in this study (33-36 Mb) were in

line with previous estimates for *A. alternata*, with the tomato pathotype also previously assembled

into 34 Mb (Hu et al. 2012). Synteny analysis of our two reference genomes against the

chromosome-level *A. solani* genome revealed structural differences for chromosomes 1 and 10 in the

apple pathotype and for chromosomes 1 and 2 in the pear pathotype. These structural differences may

represent distinct traits between clades of the *A. alternata* species group, and may represent a barrier

to genetic exchange involved in the divergence of *A. gaisen* and *A. tenuissima* lineages. The number

of essential chromosomes in our reference genomes is in line with previous findings in *A. alternata*

(Kodama et al. 1998), with 9-11 core.

Species designations within the species group have been subject to recent revision (Woudenberg et

al. 2015; Lawrence et al. 2013; Armitage et al. 2015) leading to potential confusion when selecting

isolates for study. For example, the available *A. fragariae* genome (Dang et al. 2015), did not

represent a strawberry pathotype isolate and was located in the *A. gaisen* clade. As such, the

phylogenetic context for sequenced *Alternaria* genomes described in this study, along with pathotype

identification provides a useful framework for isolate selection in future work.

4.1   Evidence of genetic exchange

A 1:1 ratio of MAT loci was observed within *A. arborescens* and *A. tenuissima* clades. This supports

previous identification of both idiomorphs within *A. alternata, A. brassicae* and *A. brassicicola*

(Berbee et al. 2003). Furthermore, presence of both MAT idiomorphs within apple pathotype isolates

indicates that genetic exchange (sexuality or parasexuality) has occurred since the evolution of

CDCs, providing a mechanism of transfer of CDCs. Furthermore, we show that some recent or

historic genetic exchange has occurred between *A. tenuissima* and *A. arborescens* clades, with both

apple and tangerine pathotypes exhibiting a polyphyletic distribution throughout the phylogeny.

4.2   Duplication of toxin-gene contigs

Toxin genes have been proposed to be present in multiple copies within *A.* sect. *alternaria* pathotype

genomes with *AMT2* proposed to be present in at least three copies in the apple pathotype CDC

(Harimoto et al. 2008), and multiple copies of *AKTR* and *AKT3* in the pear pathotype (Tanaka et al.

1999; Tanaka and Tsuge 2000). Through read mapping we demonstrated that this is the case.

Furthermore, we show that toxin gene clusters are present on multiple contigs, with differences in the

gene complements between these clusters. At this stage, it is unclear whether these different clusters

are responsible for the production of the variant R-groups previously characterised in AMT or AKT

toxins (Nakashima et al. 1985; Harimoto et al. 2007). Differences were also noted between non-

pathotype isolates from the *A. tenuissima* clade in the presence/absence of contigs 14 and 19,

representing a total of 775 Kb. Chromosomal loss has been reported in the apple pathotype (Johnson

et al. 2001), and it is not clear if this represents chromosomal instability in culture or additional

dispensable chromosomes within *A. tenuissima* clade isolates.

4.3   PCR primers for diagnostics

It is now clear that genes on essential chromosomes do not provide reliable targets for identification

of different pathotypes and hence loci located directly on CDCs should be used. We found *AMT14*

homologs to be present in all pathotype genomes and designed primers to this region. These

demonstrated specificity to apple, pear and strawberry pathotypes within a set of 86 *Alternaria*

isolates. Furthermore, Sanger sequencing of these amplicons confirmed this to be a single locus that

can both identify and discriminate a range of pathotypes. Wider validation of this primer set is now

required to test its suitability across other pathotypes.

4.4   Divergence of *A. arborescens* and *A. tenuissima*

The divergence of *A. tenuissima* and *A. arborescens* lineages was investigated through identification

of expanded and unique gene compliments. We identified HET loci unique to *A. arborescens* or *A.*

tenuissima lineages. HET loci may act as incompatibility barriers to common genetic exchange

between these taxa (Glass and Kaneko 2003). Taxa also showed divergence in effector profiles,

including chitin binding effectors, with *A. arborescens* isolates possessing unique xylan/pectin

degradation CAZymes, while *A. tenuissima* isolates possessed unique carboxylesterase,

chitooligosaccharide and sialidase CAZymes. Chitin binding proteins are important in preventing

MAMP triggered host recognition by plants and animals during infection, and may also aid

persistence of resting bodies outside of the host (Kombrink and Thomma 2013). Putative

transcription factors were also amongst the proteins specific to *A. arborescens* or *A. tenuissima*,

indicating that these taxa not only possess distinct gene complements but also differ in how they

respond to stimuli. Dispersed repeat sequences such as transposable elements have been shown to

serve as sites of recombination within and between fungal chromosomes (Zolan 1995) and we also

show distinct transposon profiles between *A. arborescens* and *A. tenuissima*. Transposons are known

to aid host adaptation in plant pathogens (Faino et al. 2016; Gijzen 2009; Schmidt et al. 2013) and

have been a mechanism for differentiation of these taxa.

4.5   Effectors on CDC regions

Alternaria HSTs are capable of inducing necrosis on non-host leaves (Kohmoto et al. 1976), meaning

that non-host resistance must be associated with recognition of other avirulence genes. We

investigated the complements of other putative pathogenicity genes and effectors produced by the

apple and Asian pear pathotypes and identified additional CAZymes and secondary metabolite

profiles on CDC regions, distinct between pathotypes, suggesting additional host-adapted tools for

pathogenicity. Additional secondary metabolites clusters were present on both apple and pear

pathotype CDCs as well as unique complements of secreted CAZymes. CAZyme families AA3, AA7

and AA9 have previously been reported to be in greater numbers in the citrus pathotype in

comparison to non-pathotypes (Wang et al. 2016). Furthermore, putative transcription factor genes

were identified in CDCs indicating that these regions may have some level of transcriptional

autonomy from the core genome. This has been shown in *Fusarium*, where effector proteins are

regulated by the SGE transcription factor on the core genome but also by FTF and other transcription

factor families (TF1-9) located on lineage specific chromosomes (van der Does et al. 2016).

4.6   Conclusions

We report near-complete reference genomes for the apple and Asian pear pathotypes of *A*. sect.

alternaria and provide genomic resources for a further ten diverse isolates from this clade. For the

first time we show sequenced *Alternaria* genomes in a phylogenetic context allowing the

identification of both mating type idiomorphs present in *A. arborescens* and *A. tenuissima*, with a

distribution throughout subclades that was indicative of recent genetic exchange. The presence of the

apple CDC in isolates of both mating types supports gene flow between isolates. Furthermore, the

distribution of isolates from different pathotypes throughout the phylogeny indicated that apple and

tangerine pathotypes are polyphyletic. This means that gene flow is not limited to within, but has also

occurred between *A. tenuissima* and *A. arborescens* lineages. We also developed PCR primers to aid

identification of pathotypes, with those targeting the AMT14 locus identifying a range of pathotypes

due to its conservation between CDCs. Despite evidence of genetic exchange between *A. arborescens*

and *A. tenuissima* clades, we show that these taxa are sufficiently isolated to have diverged, with

significant differences in core effector profiles and transposon content.

5   Figure Legends and Tables

Figure 1

Genome alignment between the reference *A. solani* genome and long-read assemblies of *A. alternata*
apple (A) and Asian pear (B) pathotypes. Links are shown between aligned regions. Locations of
telomere repeat sequences are marked within assembled contigs. Contig order in reference to *A.*
solani chromosomes is summarized, with those contigs displaying evidence of structural
rearrangement marked with an asterisk.

Figure 2

Distinct DDE and gypsy family transposon families between genomes of *A. arborescens* (arb) and *A.*
tenuissima (ten) clade isolates. Numbers of identified transposons are also shown for hAT, TY1
copia, mariner, cacta, LINE, MuDR / Mu transposases, helitrons and the Ant1-like mariner elements.

Figure 3

Phylogeny of sequenced and publicly available *Alternaria* spp. genomes. Maximum parsimony
consensus phylogeny of 500 conserved single copy loci. Dotted lines show branches with support
from < 80% of trees. Mating type idiomorphs MAT1-1-1 and MAT1-2-1 show distribution
throughout the phylogeny. Isolates pathotype is labelled following identification of genes involved in
synthesis of apple, pear, strawberry, tangerine, rough lemon and tomato toxins.

Figure 4

Presence/absence of toxin and mating type genes for 89 *Alternaria* isolates. Results are plotted onto
the 5-gene phylogeny of Armitage (2015). MAT1-1-1 and MAT1-2-1 mating type idiomorphs are
designated.

Figure 5

Number of *Alternaria* lesions per apple leaf at 14 dpi for treatments in virulence assays on *cv.*
Spartan or *cv.* Bramley's Seedling leaves. Apple pathotype isolates (*FERA 635*, *FERA 743* and *FERA*
1166) cause significant disease symptoms in comparison to control leaves, in contrast to non-
pathotype isolates (*FERA 648*, *FERA 1082* and *FERA 1164*). Number of lesions (±SE) are shown
with significance at $P < 0.05$, as determined from a GLM at the isolate and pathotype level.

Table 1

NCBI biosample and genome accession numbers of data generated in this study.

Isolate	BioSample	Accession
FERA 675	SAMN06205217	PDUP01000000
RGR 97.0013	SAMN06205218	PDWY01000000
RGR 97.0016	SAMN06205219	PEJP01000000
FERA 650	SAMN06205220	PDWZ02000000
FERA 1082	SAMN06205221	PDXA01000000
FERA 1164	SAMN06205222	PDXB01000000

FERA 1166	SAMN06205223	PDXC01000000
FERA 1177	SAMN06205224	PDXD01000000
FERA 24350	SAMN06205225	PDXE01000000
FERA 635	SAMN06205226	PDXF01000000
FERA 648	SAMN06205227	PDXG01000000
FERA 743	SAMN06205228	PDXH01000000

Organism	A. gaisen	A. tenuissima								A. arborescens		
Pathotype	pear	apple				non-pathotype				non-pathotype		
Isolate	FERA 650	FERA 1166	FERA 635	FERA 743	FERA 1177	FERA 648	FERA 1082	FERA 1164	FERA 24350	FERA 675	RGR 97.0013	RGR 97.0016
Assembly size (Mb)	34.3	35.7	36.1	35.9	35.6	33.5	33.9	34.7	33.0	33.9	33.8	33.8
Contigs	27	22	912	788	735	124	345	250	167	325	287	339
Largest contig (Mb)	6.26	3.90	1.43	3.88	4.61	3.50	1.72	2.04	1.90	2.09	1.61	1.06
N50 (Kb)	2,110	1,584	319	622	537	1,341	547	709	608	862	685	528
Repeatmasked (Kb)	749	1,268	953	869	865	658	628	983	466	853	955	784
Repeatmasked (%)	2.18	3.55	2.64	2.42	2.43	1.96	1.85	2.83	1.41	2.51	2.83	2.32
Conserved Ascomycete genes in genome (%)	98.7	99.0	98.8	98.9	98.8	99.0	98.6	98.8	99.0	98.6	98.6	98.7
Total genes	13169	13576	13733	13707	13580	12757	13028	13114	12806	12896	12766	12820
Total proteins	13220	13633	13812	13776	13647	12798	13091	13169	12856	12936	12813	12863
Conserved Ascomycete genes in gene models (%)	98.7	99.0	98.1	98.2	97.7	98.3	98.2	98.5	98.4	97.7	98.1	98.1
Secreted proteins	1208	1251	1261	1247	1270	1228	1246	1235	1225	1199	1189	1185
Secreted EffectorP proteins	246	248	264	263	268	233	252	252	241	236	229	229
Secreted CAZYme proteins	383	389	397	390	397	392	401	385	391	375	372	382
Secondary metabolite clusters	30	34	36	36	30	28	30	29	26	28	28	30

Table 3

Identification of CDC regions in the *A. alternata* apple pathotype reference genome. Read depth is shown from illumina reads against each contig, by isolate. Contigs showing reductions in coverage from non-pathotype isolates were identified as regions of conditionally dispensable chromosomes (CDCs).

FERA 1166 contig	Length (bp)	*A. gaisen* pear FERA 650	*A. tenuissima* apple FERA 1166	FERA 635	FERA 743	FERA 1177	*A. tenuissima* - FERA 648	FERA 1082	FERA 1164	FERA 24350	*A. arborescens* - FERA 675	RGR 97.0013	RGR 97.0016
1	3902980	32	49	26	40	60	34	21	28	36	53	35	26
2	3781932	31	49	26	40	58	33	20	29	35	51	34	26
3	3000390	33	50	26	40	59	34	21	28	36	53	35	26
4	2851745	32	49	27	41	59	34	21	28	36	53	36	27
5	2693844	33	49	27	40	60	34	21	29	36	53	36	27
6	2583941	33	50	27	40	60	34	21	29	36	53	36	27
7	2502671	33	49	26	40	59	34	21	28	36	52	35	26
8	2455819	32	49	27	40	60	33	21	29	35	52	35	26
9	2451092	32	49	26	40	59	33	20	28	35	52	35	26
10	2402550	31	49	26	40	58	32	21	28	35	51	34	25
11	2194253	32	49	26	40	59	34	21	28	35	52	35	26
12	1303685	33	49	27	40	59	34	21	28	36	53	36	27
13	767820	23	48	24	36	47	25	17	25	29	44	26	20
14 (CDC)	549494	0	81	29	79	115	0	23	33	0	0	0	0
15 (CDC)	435297	62	90	51	79	407	0	0	0	0	0	0	0
16	433285	23	48	26	40	53	26	18	28	33	51	27	20
17	391795	26	48	20	33	56	30	19	27	29	51	30	23
18 (CDC)	368761	0	49	22	40	32	0	0	0	0	0	0	0
19 (CDC)	225742	0	48	22	61	72	0	18	31	0	0	0	0
20 (CDC)	154051	0	69	20	45	214	0	0	0	0	0	0	0
21 (CDC)	143777	0	49	20	36	219	0	0	0	0	0	0	0
22	109256	25	50	20	30	50	26	14	22	30	35	27	22

570 **Table 4**

571 Identification of CDC regions in the *A. alternata* Asian pear pathotype reference genome. Read depth
572 is shown from illumina reads against each contig, by isolate Contigs showing reductions in coverage
573 from non-pathotype isolates were identified as regions of conditionally dispensable chromosomes
574 (CDCs).

FERA 650 contig	Length (bp)	*A. gaisen* pear FERA 650	*A. tenuissima* apple FERA 1166	FERA 635	FERA 743	FERA 1177	*A. tenuissima* - FERA 648	FERA 1082	FERA 1164	FERA 24350	*A. arborescens* - FERA 675	RGR 97.0013	RGR 97.0016
1	6257968	35	43	25	38	58	33	20	28	35	50	34	26
2	2925786	35	45	25	39	58	34	20	28	35	51	35	26
3	2776589	35	44	26	39	58	34	20	28	35	51	35	26
4	2321443	35	45	26	39	58	34	20	28	35	51	35	27
5	2116911	35	46	26	40	59	35	21	29	36	52	36	27
6	2110033	35	44	25	39	58	33	20	28	35	51	35	26
7	1975041	35	42	25	38	56	33	19	27	34	50	34	26
8	1822907	35	43	25	39	58	33	20	28	35	51	35	26
9	1811374	34	42	25	38	57	32	19	27	34	49	34	25
10	1696734	34	42	25	38	56	32	19	27	34	50	34	26
11	1617057	35	43	25	38	57	33	19	27	34	50	34	26
12	1416541	35	47	26	40	60	35	21	29	36	53	36	27
13	1110254	35	39	24	36	55	31	19	26	33	48	32	24
14 (CDC)	629968	68	49	21	41	202	0	0	0	0	0	0	0
15	554254	34	40	24	37	56	32	19	27	33	48	33	25
16 (CDC)	547262	34	8	13	18	22	8	5	14	11	20	8	6
17	522351	35	37	23	35	55	31	18	26	33	47	33	24
18	463030	34	30	22	33	49	25	17	24	29	44	28	21
19	353531	34	30	21	34	49	26	16	23	30	43	28	21
20	350965	34	20	18	27	45	22	15	19	26	40	20	18
21	313768	35	24	19	29	46	25	15	22	27	38	28	19
22	288922	33	35	23	35	51	28	21	24	31	44	30	22
23 (CDC)	206183	34	1	3	5	43	0	0	10	1	0	0	0
24 (CDC)	89891	70	0	0	0	0	0	0	0	0	0	0	0
25	25201	16	0	0	0	0	0	0	0	0	0	0	0
26	23394	5	0	0	0	0	0	0	0	0	0	0	0
27	19592	17	0	0	0	0	0	0	0	0	0	0	0

Table 5

Genomic location (contig number) of homologs to genes from apple, pear, strawberry, tangerine,
rough lemon, and tomato toxin gene clusters. Results from reference genome isolates *FERA 1166* and
FERA 650 indicate toxin clusters are present in multiple copies within the genome. This is supported
by identification of multiple AMT1 homologs in *FERA 635* and *FERA 743*. Homology between
query sequences is shown (homolog groups), as determined from reciprocal BLAST searches
between queries. Homologs are identified by e-value $< 1x10^{-30}$, $> 70\%$ query alignment. * Marks
lower-confidence hits with $< 70\%$ query alignment.

Pathotype	Gene	NCBI Accession	Group	FERA 635	FERA 743	FERA 1166	FERA 1177	FERA 650
Apple	AMT1	AB525198		201,224	146,160	20, 21	81	
	AMT2	AB525198	a	224	160	20, 21	81	
	AMT3	AB525198		224	160	21	81	
	AMT4	AB525198		178	140	20, 21	81	
	AMT5	AB525198		178	140	20, 21	81	
	AMT6	AB525198		178	140	20, 21	81	
	AMT7	AB525198		178	140	20, 21	81	
	AMT8	AB525198		178	140	20, 21	81	
	AMT9	AB525198		178	140	20, 21	81	
	AMT10	AB525198		178	143	20	81	
	AMT11	AB525198		*199**	*144**	*21**	*81**	
	AMT12	AB525198		199	144	20	129	
	AMT13	AB525198		200	129	20	128	
	AMT14	AB525198		200	129	20	128	14
	AMT15	AB525198				*18***		*14**
	AMT16	AB525198		200, 12	129, 13	20, 7	128, 21	12
	AMTR1	AB525198		230	143	20	81	14
Pear	AKT1	AB015351	b					14, 24
	AKT2	AB015352	c					14
	AKT3	AB035492	d					14, 24
	AKTR	AB035491	e					14, 24
Strawberry	AFT1-1	AB070711	b					14, 24
	AFT3-1	AB070713	d					14, 24
	AFT3-2	AB179766	d					14, 24
	AFT9-1	AB179766						14, 24
	AFT10-1	AB179766						14, 24
	AFT11-1	AB179766						14, 24
	AFT12-1	AB179766						24
	AFTS1	AB119280	a	224	160	20, 21	81	
	AFTR-1	AB070712	e					14, 24
	AFTR-2	AB179766	e					14, 24
Tangerine	ACTT1	AB034586	b					14, 24

	ACTT2	AB432914	c	14
	ACTT3	AB176941	d	14, 24
	ACTTR	AB176941	e	14, 24
	ACTT5	AB444613		24
	ACTT6	AB444614		14
Rough Lemon	ACRTS1	AB688098		
	ACRTS2	AB725683		
Tomato	ALT1	AB465676		

Supp. information 1

586   Identification of single copy Ascomycete genes in reference *Alternaria* spp. genomes

Supp. information 2

588   Gene IDs, location and functional annotations of genes in expanded orthogroups between *A.*
589   *tenuissima* and *A. arborescens*.

Supp. information 3

591   Gene IDs, location and functional annotations of genes located on CDC contigs from apple and Asian
592   pear pathotype isolates.

Supp. information 4

Toxin gene BLAST hits in publicly available *Alternaria* spp. genomes, allowing identification of
pathotype isolates.

6   Data Availability

Accession numbers for genomic data are provided in Table 1. Sanger sequence data is deposited on
NCBI under accession numbers MK255031-MK255052.

7   Conflict of Interest

CL was employed by FERA Science Ltd. All other authors declare no competing interests.

8   Author Contributions

AA, SS, CL and JC contributed conception and design of the study; AA, HC and RH performed lab
work including library preparation and sequencing; AA performed bioinformatic analyses and wrote
the manuscript. All authors contributed to manuscript revision, read and approved the submitted
version.

9   Funding

610   AA was supported Defra Plant Health Taxonomic fellowship 2010 – 2014.

10   Acknowledgments

613   Thanks are given to the FERA Science Ltd, Dr. P. Gannibal, Dr. R. Roberts and Dr. E. Simmons for
614   access to *Alternaria* isolates. This manuscript has been released as a Pre-Print on bioRxiv (Armitage
615   et al. Preprint).
616

11   References

Ajiro, N., Miyamoto, Y., Masunaka, A., Tsuge, T., Yamamoto, M., Ohtani, K., Fukumoto, T., Gomi,
K., Peever, T.L., Izumi, Y., Tada, Y. and Akimitsu, K. 2010. Role of the host-selective ACT-toxin
synthesis gene ACTTS2 encoding an enoyl-reductase in pathogenicity of the tangerine pathotype of
Alternaria alternata. Phytopathology 100(2), pp. 120–126.

Antipov, D., Korobeynikov, A., McLean, J.S. and Pevzner, P.A., 2015. hybridSPAdes: an algorithm
for hybrid assembly of short and long reads. *Bioinformatics 32*(7), pp.1009-1015.

Arie, T., Kaneko, I., Yoshida, T., Noguchi, M., Nomura, Y. and Yamaguchi, I. 2000. Mating-type
genes from asexual phytopathogenic ascomycetes *Fusarium oxysporum* and *Alternaria alternata.*
Molecular Plant-Microbe Interactions 13(12), pp. 1330–1339.

Armitage, A.D., Barbara, D.J., Harrison, R.J., Lane, C.R., Sreenivasaprasad, S., Woodhall, J.W. and
Clarkson, J.P. 2015. Discrete lineages within *Alternaria alternata* species group: Identification using
new highly variable loci and support from morphological characters. *Fungal biology* 119(11), pp.
994–1006.

Armitage, A.D., Cockerton, H.C., Sreenivasaprasad, S., Woodhall, J.W., Lane, C.R., Harrison, R.J.,
and Clarkson, J.P. Preprint. Genomics, evolutionary history and diagnostics of the *Alternaria
alternata* species group including apple and Asian pear pathotypes. bioRxiv 534685; doi:
https://doi.org/10.1101/534685.

Bairoch, A. and Apweiler, R. 2000. The SWISS-PROT protein sequence database and its supplement
TrEMBL in 2000. *Nucleic Acids Research* 28(1), pp. 45–48.

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S., Lesin, V.M.,
Nikolenko, S.I., Pham, S., Prjibelski, A.D., Pyshkin, A.V., Sirotkin, A.V., Vyahhi, N., Tesler, G.,
Alekseyev, M.A. and Pevzner, P.A. 2012. SPAdes: a new genome assembly algorithm and its
applications to single-cell sequencing. *Journal of Computational Biology* 19(5), pp. 455–477.

Belmas, E., Briand, M., Kwasiborski, A., Colou, J., N'Guyen, G., Iacomi, B., Grappin, P., Campion,
C., Simoneau, P., Barret, M. and Guillemette, T. 2018. Genome sequence of the necrotrophic plant
pathogen *Alternaria brassicicola* Abra43. *Genome announcements* 6(6).

Berbee, M.L., Payne, B.P., Zhang, G., Roberts, R.G. and Turgeon, B.G. 2003. Shared ITS DNA

substitutions in isolates of opposite mating type reveal a recombining history for three presumed
asexual species in the filamentous ascomycete genus *Alternaria*. *Mycological Research* 107(Pt 2),
pp. 169–182.

Bihon, W., Cloete, M., Gerrano, A.S., Oelofse, D. and Adebola, P. 2016. Draft Genome Sequence of
Alternaria alternata Isolated from Onion Leaves in South Africa. *Genome announcements* 4(5).

Bonants, P., Groenewald, E., Rasplus, J.Y., Maes, M., de Vos, P., Frey, J., Boonham, N., Nicolaisen,
M., Bertacini, A., Robert, V., Barker, I., Kox, L., Ravnikar, M., Tomankova, K., Caffier, D., Li, M.,
Armstrong, K., Freitas-Astúa, J., Stefani, E., Cubero, J. and Mostert, L. 2010. QBOL: a new EU
project focusing on DNA barcoding of Quarantine organisms. *EPPO Bulletin* 40(1), pp. 30–33.

Capella-Gutiérrez, S., Silla-Martínez, J.M. and Gabaldón, T. 2009. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 25(15), pp. 1972–1973.

Chakraborty, M., Baldwin-Brown, J.G., Long, A.D. and Emerson, J.J., 2016. Contiguous and
accurate de novo assembly of metazoan genomes with modest long read coverage. *Nucleic acids*
research 44(19), pp.e147-e147.

Cingolani, P., Platts, A., Wang, L.L., Coon, M., Nguyen, T., Wang, L., Land, S.J., Lu, X. and Ruden,
D.M. 2012. A program for annotating and predicting the effects of single nucleotide polymorphisms,
SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3. *Fly* 6(2), pp.
80–92.

Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker, R.E.,
Lunter, G., Marth, G.T., Sherry, S.T., McVean, G., Durbin, R. and 1000 Genomes Project Analysis
Group 2011. The variant call format and VCFtools. *Bioinformatics* 27(15), pp. 2156–2158.

Dang, H.X., Pryor, B., Peever, T. and Lawrence, C.B. 2015. The *Alternaria* genomes database: a
comprehensive resource for a fungal genus comprised of saprophytes, plant pathogens, and allergenic
species. *BMC Genomics* 16, p. 239.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M. and
Gingeras, T.R. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29(1), pp. 15–21.

van der Does, H.C., Fokkens, L., Yang, A., Schmidt, S.M., Langereis, L., Lukasiewicz, J.M.,
Hughes, T.R. and Rep, M. 2016. Transcription factors encoded on core and accessory chromosomes
of *Fusarium oxysporum* induce expression of effector Genes. *PLoS Genetics* 12(11), p. e1006401.

Faino, L., Seidl, M.F., Shi-Kunne, X., Pauper, M., van den Berg, G.C.M., Wittenberg, A.H.J. and
Thomma, B.P.H.J. 2016. Transposons passively and actively contribute to evolution of the two-speed
genome of a fungal pathogen. *Genome Research* 26(8), pp. 1091–1100.

Gijzen, M. 2009. Runaway repeats force expansion of the *Phytophthora infestans* genome. *Genome*
Biology 10(10), p. 241.

Glass, N.L. and Kaneko, I. 2003. Fatal attraction: nonself recognition and heterokaryon
incompatibility in filamentous fungi. *Eukaryotic Cell* 2(1), pp. 1–8.

Gurevich, A., Saveliev, V., Vyahhi, N. and Tesler, G. 2013. QUAST: quality assessment tool for
genome assemblies. *Bioinformatics* 29(8), pp. 1072–1075.

Harimoto, Y., Hatta, R., Kodama, M., Yamamoto, M., Otani, H. and Tsuge, T. 2007. Expression
profiles of genes encoded by the supernumerary chromosome controlling AM-toxin biosynthesis and
pathogenicity in the apple pathotype of *Alternaria alternata*. *Molecular Plant-Microbe Interactions*
20(12), pp. 1463–1476.

Harimoto, Y., Tanaka, T., Kodama, M., Yamamoto, M., Otani, H. and Tsuge, T. 2008. Multiple
copies of AMT2 are prerequisite for the apple pathotype of Alternaria alternata to produce enough
AM-toxin for expressing pathogenicity. *Journal of General Plant Pathology* 74(3), pp. 222–229.

Hatta, R., Ito, K., Hosaki, Y., Tanaka, T., Tanaka, A., Yamamoto, M., Akimitsu, K. and Tsuge, T.
2002. A conditionally dispensable chromosome controls host-specific pathogenicity in the fungal
plant pathogen *Alternaria alternata. Genetics* 161(1), pp. 59–70.

Hoff, K.J., Lange, S., Lomsadze, A., Borodovsky, M. and Stanke, M. 2016. BRAKER1:
Unsupervised RNA-Seq-Based Genome Annotation with GeneMark-ET and AUGUSTUS.
Bioinformatics 32(5), pp. 767–769.

Hou, Y., Ma, X., Wan, W., Long, N., Zhang, J., Tan, Y., Duan, S., Zeng, Y. and Dong, Y. 2016.
Comparative genomics of pathogens causing brown spot disease of tobacco: *Alternaria longipes* and
Alternaria alternata. Plos One 11(5), p. e0155258.

Hu, J., Chen, C., Peever, T., Dang, H., Lawrence, C. and Mitchell, T. 2012. Genomic characterization
of the conditionally dispensable chromosome in *Alternaria arborescens* provides evidence for
horizontal gene transfer. *BMC Genomics* 13, p. 171.

Huang, L., Zhang, H., Wu, P., Entwistle, S., Li, X., Yohe, T., Yi, H., Yang, Z. and Yin, Y. 2018.
dbCAN-seq: a database of carbohydrate-active enzyme (CAZyme) sequence and annotation. *Nucleic*
Acids Research 46(D1), pp. D516–D521.

Johnson, L.J., Johnson, R.D., Akamatsu, H., Salamiah, A., Otani, H., Kohmoto, K. and Kodama, M.
2001. Spontaneous loss of a conditionally dispensable chromosome from the *Alternaria alternata*
apple pathotype leads to loss of toxin production and pathogenicity. *Current Genetics* 40(1), pp. 65–
72.

Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J.,
Mitchell, A., Nuka, G., Pesseat, S., Quinn, A.F., Sangrador-Vegas, A., Scheremetjew, M., Yong, S.-
Y., Lopez, R. and Hunter, S. 2014. InterProScan 5: genome-scale protein function classification.
Bioinformatics 30(9), pp. 1236–1240.

Käll, L., Krogh, A. and Sonnhammer, E.L.L. 2004. A combined transmembrane topology and signal
peptide prediction method. *Journal of Molecular Biology* 338(5), pp. 1027–1036.

Katoh, K. and Standley, D.M. 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Molecular Biology and Evolution* 30(4), pp. 772–780.

Kodama, M., Akamatsu, H., Itoh, Y., Narusaka, Y., Sanekata, T., Otani, H. and Kohmoto, K. 1998.
Host-specific toxin deficient mutants of the tomato pathotype of *Alternaria alternata* obtained by
restriction enzyme-mediated integration. In: Kohmoto, K. and Yoder, O. C. eds. *Molecular Genetics*
of Host-Specific Toxins in Plant Disease. Developments in plant pathology. Dordrecht: Springer
Netherlands, pp. 35–42.

Kohmoto, K., Khan, I.D., Renbutsu, Y., Taniguchi, T. and Nishimura, S. 1976. Multiple host-specific
toxins of *Alternaria mali* and their effect on the permeability of host cells. *Physiological Plant*
Pathology 8(2), pp. 141–153.

Kombrink, A. and Thomma, B.P.H.J. 2013. LysM effectors: secreted proteins supporting fungal life.
PLoS Pathogens 9(12), p. e1003769.

Koren, S., Walenz, B.P., Berlin, K., Miller, J.R., Bergman, N.H. and Phillippy, A.M. 2017. Canu:
scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation.
Genome Research 27(5), pp. 722–736.

Krzywinski, M., Schein, J., Birol, I., Connors, J., Gascoyne, R., Horsman, D., Jones, S.J. and Marra,
M.A., 2009. Circos: an information aesthetic for comparative genomics. *Genome research* 19(9),
pp.1639-1645.

Krogh, A., Larsson, B., von Heijne, G. and Sonnhammer, E.L.L. 2001. Predicting transmembrane
protein topology with a hidden Markov model: application to complete genomes. *Journal of
Molecular Biology* 305(3), pp. 567–580.

Langmead, B. and Salzberg, S.L. 2012. Fast gapped-read alignment with Bowtie 2. *Nature Methods*
9(4), pp. 357–359.

Lawrence, D.P., Gannibal, P.B., Peever, T.L. and Pryor, B.M. 2013. The sections of *Alternaria*:
formalizing species-group concepts. *Mycologia* 105(3), pp. 530–546.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
Durbin, R. and 1000 Genome Project Data Processing Subgroup 2009. The Sequence
Alignment/Map format and SAMtools. *Bioinformatics* 25(16), pp. 2078–2079.

Li, L., Stoeckert, C.J. and Roos, D.S. 2003. OrthoMCL: identification of ortholog groups for
eukaryotic genomes. *Genome Research* 13(9), pp. 2178–2189.

Liu, K., Linder, C.R. and Warnow, T. 2011. RAxML and FastTree: comparing two methods for
large-scale maximum likelihood phylogeny estimation. *Plos One* 6(11), p. e27731.

Loman, N.J., Quick, J. and Simpson, J.T. 2015. A complete bacterial genome assembled de novo
using only nanopore sequencing data. *Nature Methods* 12(8), pp. 733–735.

Masunaka, A., Ohtani, K., Peever, T.L., Timmer, L.W., Tsuge, T., Yamamoto, M., Yamamoto, H.
and Akimitsu, K. 2005. An Isolate of *Alternaria alternata* That Is Pathogenic to Both Tangerines and
Rough Lemon and Produces Two Host-Selective Toxins, ACT- and ACR-Toxins. *Phytopathology*
95(3), pp. 241–247.

Marçais, G., Delcher, A.L., Phillippy, A.M., Coston, R., Salzberg, S.L. and Zimin, A., 2018.
MUMmer4: a fast and versatile genome alignment system. *PLoS computational biology* 14(1),
p.e1005944.

Masunaka, A., Tanaka, A., Tsuge, T., Peever, T.L., Timmer, L.W., Yamamoto, M., Yamamoto, H.
and Akimitsu, K. 2000. Distribution and Characterization of AKT Homologs in the Tangerine
Pathotype of *Alternaria alternata*. *Phytopathology* 90(7), pp. 762–768.

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella, K.,
Altshuler, D., Gabriel, S., Daly, M. and DePristo, M.A. 2010. The Genome Analysis Toolkit: a
MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Research*
20(9), pp. 1297–1303.

Mistry, J., Finn, R.D., Eddy, S.R., Bateman, A. and Punta, M. 2013. Challenges in homology search:
HMMER3 and convergent evolution of coiled-coil regions. *Nucleic Acids Research* 41(12), p. e121.

Miyamoto, Y., Ishii, Y., Honda, A., Masunaka, A., Tsuge, T., Yamamoto, M., Ohtani, K., Fukumoto,
T., Gomi, K., Peever, T.L. and Akimitsu, K. 2009. Function of genes encoding acyl-CoA synthetase
and enoyl-CoA hydratase for host-selective act-toxin biosynthesis in the tangerine pathotype of
Alternaria alternata. *Phytopathology* 99(4), pp. 369–377.

Miyamoto, Y., Masunaka, A., Tsuge, T., Yamamoto, M., Ohtani, K., Fukumoto, T., Gomi, K.,
Peever, T.L., Tada, Y., Ichimura, K. and Akimitsu, K. 2010. ACTTS3 encoding a polyketide
synthase is essential for the biosynthesis of ACT-toxin and pathogenicity in the tangerine pathotype
of *Alternaria alternata*. *Molecular Plant-Microbe Interactions* 23(4), pp. 406–414.

Nakashima, T., Ueno, T., Fukami, H., Taga, T., Masuda, H., Osaki, K., Otani, H., Kohmoto, K. and
Nishimura, S. 1985. Isolation and structures of AK-toxin I and II. Host-specific phytotoxic
metabolites produced by *Alternaria alternata* Japanese pear pathotype. *Agricultural and biological*
chemistry 49(3), pp. 807–815.

Nguyen, H.D.T., Lewis, C.T., Lévesque, C.A. and Gräfenhan, T. 2016. Draft Genome Sequence of
Alternaria alternata ATCC 34957. *Genome announcements* 4(1).

Nieuwenhuis, B.P.S. and James, T.Y. 2016. The frequency of sex in fungi. *Philosophical*
Transactions of the Royal Society of London. Series B, Biological Sciences 371(1706).

Pfeifer, B., Wittelsbürger, U., Ramos-Onsins, S.E. and Lercher, M.J. 2014. PopGenome: an efficient
Swiss army knife for population genomic analyses in R. *Molecular Biology and Evolution* 31(7), pp.
1929–1936.

Quaedvlieg, W., Groenewald, J.Z., de Jesús Yáñez-Morales, M. and Crous, P.W. 2012. DNA
barcoding of *Mycosphaerella* species of quarantine importance to Europe. *Persoonia* 29, pp. 101–
115.

Ruan, J. 2016. Ultra-fast de novo assembler using long noisy reads [Online]. Available at:
https://github.com/ruanjue/smartdenovo [Accessed: 26 June 2018].

Schmidt, S.M., Houterman, P.M., Schreiver, I., Ma, L., Amyotte, S., Chellappan, B., Boeren, S.,
Takken, F.L.W. and Rep, M. 2013. MITEs in the promoters of effector genes allow prediction of
novel virulence genes in *Fusarium oxysporum. BMC Genomics* 14, p. 119.

Shelest, E. 2017. Transcription Factors in Fungi: TFome Dynamics, Three Major Families, and Dual-
Specificity TFs. *Frontiers in genetics* 8, p. 53.

Simão, F.A., Waterhouse, R.M., Ioannidis, P., Kriventseva, E.V. and Zdobnov, E.M. 2015. BUSCO:
assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*
31(19), pp. 3210–3212.

Sperschneider, J., Gardiner, D.M., Dodds, P.N., Tini, F., Covarelli, L., Singh, K.B., Manners, J.M.
and Taylor, J.M. 2016. EffectorP: predicting fungal effector proteins from secretomes using machine
learning. *The New Phytologist* 210(2), pp. 743–761.

Stanke, M. and Morgenstern, B. 2005. AUGUSTUS: a web server for gene prediction in eukaryotes
that allows user-defined constraints. *Nucleic Acids Research* 33(Web Server issue), pp. W465-7.

Stewart, J.E., Thomas, K.A., Lawrence, C.B., Dang, H., Pryor, B.M., Timmer, L.M.P. and Peever,
T.L. 2013. Signatures of recombination in clonal lineages of the citrus brown spot pathogen,
Alternaria alternata sensu lato. *Phytopathology* 103(7), pp. 741–749.

Stumpf, M.P.H. and McVean, G.A.T. 2003. Estimating recombination rates from population-genetic
data. *Nature Reviews. Genetics* 4(12), pp. 959–968.

Tanaka, A., Shiotani, H., Yamamoto, M. and Tsuge, T. 1999. Insertional mutagenesis and cloning of
the genes required for biosynthesis of the host-specific AK-toxin in the Japanese pear pathotype of
Alternaria alternata. Molecular Plant-Microbe Interactions 12(8), pp. 691–702.

Tanaka, A. and Tsuge, T. 2000. Structural and functional complexity of the genomic region
controlling AK-toxin biosynthesis and pathogenicity in the Japanese pear pathotype of Alternaria
alternata. *Molecular Plant-Microbe Interactions* 13(9), pp. 975–986.

Testa, A.C., Hane, J.K., Ellwood, S.R. and Oliver, R.P. 2015. CodingQuarry: highly accurate hidden
Markov model gene prediction in fungal genomes using RNA-seq transcripts. *BMC Genomics* 16, p.

170.

Thomma, B.P.H.J. 2003. Alternaria spp.: from general saprophyte to specific parasite. *Molecular*
Plant Pathology 4(4), pp. 225–236.

Tsuge, T., Harimoto, Y., Akimitsu, K., Ohtani, K., Kodama, M., Akagi, Y., Egusa, M., Yamamoto,
M. and Otani, H. 2013. Host-selective toxins produced by the plant pathogenic fungus *Alternaria*
alternata. *FEMS Microbiology Reviews* 37(1), pp. 44–66.

Vaser, R., Sović, I., Nagarajan, N. and Šikić, M. 2017. Fast and accurate de novo genome assembly
from long uncorrected reads. *Genome Research* 27(5), pp. 737–746.

Walker, B.J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S., Cuomo, C.A., Zeng, Q.,
Wortman, J., Young, S.K. and Earl, A.M. 2014. Pilon: an integrated tool for comprehensive
microbial variant detection and genome assembly improvement. *Plos One* 9(11), p. e112963.

Wang, M., Sun, X., Yu, D., Xu, J., Chung, K. and Li, H. 2016. Genomic and transcriptomic analyses
of the tangerine pathotype of *Alternaria alternata* in response to oxidative stress. *Scientific reports* 6,
p. 32437.

Wolters, P.J., Faino, L., van den Bosch, T.B.M., Evenhuis, B., Visser, R.G.F., Seidl, M.F. and
Vleeshouwers, V.G.A.A. 2018. Gapless Genome Assembly of the Potato and Tomato Early Blight
Pathogen *Alternaria solani*. *Molecular Plant-Microbe Interactions* 31(7), pp. 692–694.

Woudenberg, J.H.C., Seidl, M.F., Groenewald, J.Z., de Vries, M., Stielow, J.B., Thomma, B.P.H.J.
and Crous, P.W. 2015. *Alternaria* section *Alternaria*: Species, formae speciales or pathotypes?
Studies in mycology 82, pp. 1–21.

Yu, G., Smith, D.K., Zhu, H., Guan, Y. and Lam, T.T.-Y. 2016. *ggtree*□: an *R* package for
visualization and annotation of phylogenetic trees with their covariates and other associated data.
Methods in ecology and evolution / British Ecological Society.

Zhang, C., Rabiee, M., Sayyari, E. and Mirarab, S. 2018. ASTRAL-III: polynomial time species tree
reconstruction from partially resolved gene trees. *BMC Bioinformatics* 19(Suppl 6), p. 153.

Zolan, M.E. 1995. Chromosome-length polymorphism in fungi. *Microbiological reviews* 59(4), pp.
686–698.

A

A. solani
NL03003

Apple pathotype
FERA 1166

NL03003 chromosome	FERA 1166 contig
1	8, 6*, 2
2	17, 12, 1
3	3, 16
4	22, 5
5	4
6	11, 13
7	7
8	10
9	9
10	6*
-	14, 15, 18, 19, 20, 21

B

A. solani
NL03003

Asian pear pathotype
FERA 650

NL03003 chromosome	FERA 650 contig
1	9, 5, 1*
2	10, 1*
3	2, 19
4	20, 3
5	17, 4
6	6, 18
7	13, 12
8	22, 7, 16
9	15, 8
10	21, 11
-	14, 23, 24, 25, 26, 27

RGR 97.0021	*Malus domestica*	MAT1–2–1	
RGR 97.0022	*Malus domestica*	MAT1–2–1	
RGR 97.0010	*Malus domestica*	MAT1–1–1	
RGR 97.0015	*Malus domestica*	MAT1–2–1	
RGR 97.0008	*Malus domestica*	MAT1–1–1	
RGR 97.0014	*Malus domestica*	MAT1–2–1	
RGR 97.0007	*Malus domestica*	MAT1–2–1	
FERA 681	*Pyrus sp.*	MAT1–2–1	
RGR 97.0012	*Malus domestica*	MAT1–1–1	
FERA 631	*Pyrus communis*	MAT1–1–1	
FERA 488	*Prunus persica*	MAT1–2–1	
RGR 97.0025	*Malus domestica*	MAT1–1–1	
FERA 1539	*Juglans regia*	MAT1–2–1	
FERA 348	*Malus domestica*	MAT1–2–1	
RGR 97.0018	*Malus domestica*	MAT1–1–1	
FERA 2040	*Malus domestica*	MAT1–2–1	
RGR 97.0009	*Malus domestica*	MAT1–2–1	
RGR 97.0017	*Malus domestica*	MAT1–2–1	
RGR 97.0019	*Malus domestica*	MAT1–1–1	
RGR 97.0011	*Malus domestica*	MAT1–1–1	
RGR 97.0016	*Malus domestica*	MAT1–2–1	
RGR 97.0026	*Malus domestica*	MAT1–2–1	
FERA 1521	*Prunus armeniaca*	MAT1–2–1	
RGR 97.0027	*Malus domestica*	MAT1–1–1	
RGR 97.0067	*Malus domestica*	MAT1–1–1	
RGR 97.0013	*Malus domestica*	MAT1–1–1	
FERA 675	*Pyrus pyrifolia*	MAT1–2–1	
RGR 97.0024	*Malus domestica*	MAT1–2–1	
RGR 97.0020	*Malus domestica*	MAT1–1–1	
FERA 484		MAT1–2–1	
FERA 1518	*Solanum lycopersicum*	MAT1–2–1	
FERA 1171	*Malus domestica*	MAT1–2–1	
FERA 2277	*Impatiens walleriana*	MAT1–1–1	
FERA 2280	*Impatiens walleriana*	MAT1–1–1	
FERA 9285	*Impatiens walleriana*	MAT1–1–1	
FERA 1169	*Malus domestica*	MAT1–1–1	
FERA 1170	*Malus domestica*	MAT1–2–1	
FERA 1172	*Malus domestica*	MAT1–1–1	
FERA 1512	*Dendrathema sp.*	MAT1–1–1	
FERA 479	*Fragaria sp.*	MAT1–1–1	
FERA 478	*Fragaria sp.*	MAT1–1–1	
FERA 2038	*Solanum lycopersicum*	MAT1–2–1	
FERA 1522	*Actinidia sp.*	MAT1–2–1	
EGS 39.128	*Solanum lycopersicum*	MAT1–2–1	
FERA 647	*Pyrus sp.*	MAT1–1–1	
FERA 21677	*Pyrus sp.*	MAT1–2–1	
FERA 538	*Pyrus pyrifolia*	MAT1–2–1	
FERA 326	*Pyrus pyrifolia*	MAT1–1–1	
FERA 540	*Pyrus sp.*	MAT1–2–1	
FERA 704	*Pyrus pyrifolia*	MAT1–2–1	
FERA 2044	*Malus domestica*	MAT1–2–1	
FERA 679	*Malus domestica*	MAT1–2–1	
FERA 678	*Malus domestica*	MAT1–2–1	
FERA 2043	*Malus domestica*	MAT1–1–1	
FERA 2041	*Malus domestica*	MAT1–2–1	
FERA 1551	*Malus domestica*	MAT1–2–1	AMT4 + AMT14
FERA 743	*Malus domestica*	MAT1–2–1	AMT4 + AMT14
FERA 24350	*Pyrus pyrifolia*	MAT1–1–1	
FERA 635	*Malus domestica*	MAT1–2–1	AMT4 + AMT14
FERA 833	*Malus domestica*	MAT1–1–1	
FERA 1166	*Malus domestica*	MAT1–2–1	AMT4 + AMT14
FERA 1490	*Fragaria sp.*	MAT1–1–1	AKT3 + AMT14*
FERA 482	*Fragaria sp.*	MAT1–1–1	
EGS 34.015	*Dianthus sp.*	MAT1–2–1	
FERA 648	*Pyrus sp.*	MAT1–1–1	
O–159	*Malus domestica*	MAT1–1–1	AMT4 + AMT14
EGS 38.029	*Malus domestica*	MAT1–1–1	
FERA 469	*Citrus reticulata*	MAT1–1–1	
FERA 2712	*Citrus sp.*	MAT1–2–1	
EGS 34.016	*Arachis sp.*	MAT1–1–1	
FERA 1105		MAT1–1–1	
FERA 1410		MAT1–2–1	
FERA 2103	*Citrus sp.*	MAT1–2–1	
HA–1	*Malus domestica*	MAT1–1–1	
FERA 23766	*Pyrus pyrifolia*	MAT1–2–1	
FERA 1310	*Malus domestica*	MAT1–2–1	
FERA 1491	*Fragaria sp.*	MAT1–1–1	AKT3 + AMT14*
FERA 1523	*Citrullus lanatus*	MAT1–2–1	
FERA 1520	*Lactuca sp.*	MAT1–1–1	
FERA 3250		MAT1–2–1	
FERA 1082	*Malus domestica*	MAT1–2–1	
FERA 1519	*Gossypium sp.*	MAT1–2–1	
FERA 634	*Malus domestica*	MAT1–1–1	
FERA 15968	*Solanum tuberosum*	MAT1–1–1	
FERA 1177	*Malus domestica*	MAT1–1–1	AMT4 + AMT14
FERA 1164	*Malus domestica*	MAT1–2–1	
FERA 1165	*Malus domestica*	MAT1–2–1	
FERA 650	*Pyrus sp.*	MAT1–1–1	AKT3 + AMT14
FERA 632	*Pyrus sp.*	MAT1–1–1	AKT3 + AMT14
EGS 90.0512	*Pyrus pyrifolia*	MAT1–1–1	AKT3 + AMT14

arborescens clade

tenuissima clade

gaisen clade

0.2