Title: Increased Replication Rates of Dissimilatory Nitrate-Reducing Bacteria Lead to Decreased

Anammox Bioreactor Performance

Authors: Ray Keren[1]*, Jennifer E. Lawrence[1,2]*, Weiqin Zhuang[1,3], David Jenkins[1], Jillian F. Banfield[4],

Lisa Alvarez-Cohen[1], Lijie Zhou[1,5]+, Ke Yu[1,6]+

*These authors contributed equally to the work

+Corresponding authors

Affiliations

1- Department of Civil and Environmental Engineering, University of California Berkeley,

Berkeley, CA, USA

2- Tighe & Bond, Westwood, MA, USA

3- Department of Civil and Environmental Engineering, The University of Auckland,

Auckland, New Zealand

4- Earth and Planetary Sciences, University of California Berkeley, Berkeley, CA, USA

5- College of Chemistry and Environmental Engineering, Shenzhen University, Shenzhen,

China

6- Shenzhen Graduate School, Peking University, Shenzhen, China

Abstract

Anaerobic ammonium oxidation (anammox) is a biological process employed to remove reactive nitrogen

from wastewater. While a substantial body of literature describes the performance of anammox

bioreactors under various operational conditions and perturbations, few studies have resolved the

metabolic roles of its community members. Here, we use metagenomics to study the microbial

community within a laboratory-scale anammox bioreactor from inoculation, through performance

destabilizations, to stable steady-state. Metabolic analyses reveal that dissimilatory nitrate reduction to

ammonium (DNRA) is the primary nitrogen removal pathway that competes with anammox in the

bioreactor. Increased replication rates of bacteria capable of DNRA lead to the out-competition of

annamox bacteria, which is the key source of fixed carbon, and the loss of bioreactor performance.

Ultimately, our findings underline the importance of metabolic interdependencies related to nitrogen and

carbon-cycling within anammox bioreactors and highlight the potentially detrimental effects of bacteria

that are otherwise considered core community members.

Main Text

Anammox bacteria obtain energy from the conversion of ammonium and nitrite into molecular

nitrogen gas (N_2)[1]. The only currently known bacteria that catalyze this process are Planctomycetes[2,3],

none of which have yet been isolated[3,4]. In practice, anammox bacteria are employed in an eponymous

process in combination with the partial nitritation (PN) process to remove ammonium from wastewaters

or anaerobic digestors though side-streams. First, in PN, approximately half of the ammonium in solution

is aerobically oxidized to nitrite. Second, in anammox, both ammonium and nitrite are anaerobically

converted to N_2[5,6]. PN/anammox is beneficial because it consumes 60% less energy, produces 90% less

biomass, and emits a significantly smaller volume of greenhouse gases than conventional nitrogen

removal by nitrification and denitrification processes[7].

Even though over 100 full-scale PN/anammox processes have been installed across the globe at

municipal and industrial wastewater treatment plants,[8] anammox bacteria have very low growth rates

within engineered environments and are easily inhibited by a variety of factors including fluctuating

substrate and metabolite concentrations[9,10]. Furthermore, recovery from an inhibition event can take up to

six months, which is unacceptably long for municipalities who must meet strict nitrogen discharge

limits[11]. These problems are compounded by what is currently only a cursory understanding of the

microbial communities responsible for stable, robust anammox performance. The broad application of

PN/anammox in wastewater treatment processes requires a more comprehensive understanding of the

complex interactions among the numerous bacterial species within the bioreactors.

Previous research suggests that a core microbial community exists within anammox bioreactors[12–

16]. In the majority of anammox bioreactors, uncultured members of the phyla Bacteroidetes, Chloroflexi,

Ignavibacteria, and Proteobacteria have been identified alongside Planctomycetes. Since these phyla have

primarily been identified through 16S rRNA studies, their interplay with anammox performance has yet

to be elucidated[12–16]. From their taxonomic identity, it is assumed that the additional groups of bacteria

cooperate to transform and remove nitrate, a product of anammox metabolism[17–19].

Here, we illuminate the metabolic relationships between the anammox bacterium and its

supporting community members during the start-up and operation of a laboratory-scale anammox

bioreactor. We used genome-centric metagenomics to recover 337 draft genomes from six time-points

spanning 440 days of continuous bioreactor operation. We combined our reconstruction of the microbial

community's dynamic metabolic potential with bioreactor performance data and relative abundance

profiles based on both metagenomic and 16S rRNA sequencing. As a result, we were able to identify core

metabolic activities and potential interdependencies that inform the performance and stability of the

anammox bioreactor. We found that certain metabolic interactions between the anammox bacterium and

associated community members may be responsible for the destabilization of anammox bioreactors. To

our knowledge, this is the first time-series-based study to link anammox metagenomic insights and

community composition to bioreactor functionality[20]. Our findings bolster the fundamental, community-

level understanding of the anammox process. Ultimately, these results will enable better understanding of

this important microbial process and a more comprehensive control of this promising technology to help

facilitate its widespread adoption at wastewater treatment plants.

Bioreactor performance. The performance of a laboratory-scale anaerobic membrane bioreactor

(described in methods) was tracked for 440 days from initial inoculation, through several performance

crashes, to stable and robust anammox activity (Figure 1). Performance was quantified in a variety of

ways, including by its nitrogen removal rate (NRR, g-N L^{-1} d^{-1}) and by its effluent quality (g-N L^{-1} d^{-1}).

Bioreactor performance generally improved over the first 103 days of operation. At this point, the

hydraulic residence time was reduced from 48 to 12 hours and influent concentrations were reduced to

maintain a stable loading rate. Additional biomass from a nearby pilot-scale PN/anammox process was

added on Day 145 and bioreactor performance improved, enabling influent ammonium and nitrite

concentrations to be steadily increased until the NRR approached 2 g-N L^{-1} d^{-1}. On Day 189 the

bioreactor experienced a technical malfunction and subsequent performance crash, identified by a rapid

decrease in the NRR and the effluent quality. On Day 203, the bioreactor was again amended with a

concentrated stock of biomass and the NRR and the effluent quality quickly recovered. Influent

ammonium and nitrite concentrations were again increased until the NRR reached 2 g-N L^{-1} d^{-1}.

The bioreactor subsequently maintained steady performance for approximately 75 days, until Day

288, when effluent concentrations of ammonium and nitrite unexpectedly began to increase and nitrate

concentrations disproportionately decreased. Seven days later, the NRR rapidly plummeted. No technical

malfunctions had occurred, indicating that a destabilized microbial community may have been

responsible for the performance crash. At that time, the cause of the performance decline was not

understood, so the bioreactor was not re-seeded with biomass. After 50 days of limited performance,

concentrations of copper, iron, molybdenum, and zinc in the bioreactor influent were increased[21–24] and

the NRR rapidly recovered. Stable and robust bioreactor performance was subsequently maintained.

Metagenomic sequencing and binning. Whole community DNA was extracted and sequenced at six

time-points throughout the study: Day 0 (D0), for inoculant composition; Day 82 (D82), during nascent,

positive anammox activity; Day 166 (D166), three weeks after an additional biomass amendment; Day

284 (D284), after a long period of stable and robust anammox activity and just before destabilization; Day

328 (D328), in the midst of the destabilization period; and Day 437 (D437), during mature, stable, and

robust anammox activity.

From all samples, 337 genomes were binned, 244 of which were estimated to be >70% complete.

The genomes were further dereplicated across the six time-points into clusters at 95% average nucleotide

identity (ANI). This resulted in 127 representative and unique genomes (Table 1), which were used for all

downstream analyses. Mapping showed an average read recruitment of 76% to representative genomes

(Table 2). The number of genomes present at each time-point (using threshold values of coverage > 1 and

breadth > 0.5) ranged from 60 (D437) to 103 (D166). In addition, nine strains were detected that differed

from the representative genome by 2% ANI (Supplemental Information, Supplemental Table 1). With the

exception of the anammox bacterium, referred to at the genus level, all genomes are referred to at their

phylum level.

Core anammox community. Resulting genomes from our study, in combination with genomes from two

previous anammox metagenomic studies, Speth et al.[18] (22 genomes) and Lawson et al.[25] (15 genomes),

provide strong evidence to support a core anammox community (Figure 2). The relative abundances of

bacteria from the dominant phyla across these three bioreactors are fairly similar: in each bioreactor the

anammox, along with Chloroflexi, Ignavibacteria, and Proteobacteria bacteria, compose >70% of the

community (Figure 2B).

Due to the significantly larger genome yield and time-series analysis in this study, our bioreactor

shared more genomes with each of the other bioreactors than the other bioreactors shared between

themselves. Nevertheless, three genomes from closely related bacteria were identified across all three

bioreactors: *Brocadia* (responsible for anammox), an unclassified Chloroflexi, and an unclassified

Ignavibacteria. All three of these genomes were present in our bioreactor during stable operation on

D437, and two of them (*Brocadia* and an Ignavibacterium) were among the ten most abundant genomes

at that time. In total, 21 genomes from our bioreactor were closely related to those from at least one of the

two other bioreactors, 17 of which were present at D437 (Supplemental Table 2). The related bacteria

accounted for 50% and 93% of the Speth et al. and Lawson et al. genomes, respectively. The bioreactor

studied by Speth et al. was different from the other two bioreactors because it was amended with oxygen

to perform partial nitritation and anammox within the same bioreactor, while the others performed

anammox only.

A more focused phylogenetic tree of Planctomycetes showed that the *Brocadia* in our bioreactor

and in the Lawson et al. bioreactor are the same species (*Brocadia sapporensis*[26]), while the *Brocadia*

species from the Speth et al. bioreactor is different (*Brocadia sinica*) (Supplemental Figure 1).

Community dynamics. The relative abundances of organisms represented by genomes were calculated

by multiplying genome coverage and breadth. Additional 16S rRNA gene sequencing, executed at 56

timepoints across the lifespan of the bioreactor, allowed us to expand our view of the relative abundances

of bacterial genomes over time. 38 of the 127 genomes contained 16S rRNA gene sequences that matched

16S rRNA gene sequencing efforts, and these 38 genomes accounted for the majority of the bioreactor

microbial community (Figure 3).

The *Brocadia* genus accounted for a small fraction of the bacteria in the inoculating biomass.

Consistent with previous research of a combination PN/anammox bioreactor, members of the phyla

Acidobacteria, Bacteroidetes, Ignavibacteriae, and Proteobacteria were also present[12–16]. During the first

100 days of bioreactor operation, *Brocadia* increased in relative abundance. Its replication rate at D82

was high (Supplemental Table 3), which corroborates its overall enrichment in the community. Following

the bioreactor malperformance and biomass amendment on Day 147, the bioreactor became dominated by

a bacterium represented by a single genome of the phylum Bacteroidetes (order Sphingobacteriales). The

bacterium's calculated replication rate was low on D166, and over the next 100 days its relative

abundance steadily declined. In contrast, the *Brocadia* replication rate was extremely high on D166,

allowing it to once again dominate the microbial community. *Brocadia* remained dominant until Day 290,

when the relative abundances of several Chloroflexi (most notably, one from the class Anaerolineae) and

an Ignavibacteria dramatically increased. Shortly after this shift, the bioreactor experienced an

unexplained period of performance decline and subsequent performance crash. During this period the

Brocadia replication rated dramatically declined, while the Chloroflexi replication rates increased

(Supplemental Table 3). These shifts in replication rates six days before a response in relative abundance

profiles and 12 days before a response in NRR are consistent with an instability in population dynamics

directly impacting the bioreactor performance.

The relative abundances of *Brocadia* and the Chloroflexi, as well as their replication rates,

remained fairly constant over the next 44 days. After the influent media trace metal concentrations were

increased, the relative abundance of Chloroflexi decreased and that of *Brocadia* increased. By D437,

Brocadia again dominated the bioreactor, and the replication rates of *Brocadia* and Chloroflexi bacteria

became similar.

Community grouping. As both internal and external factors can work in combination to affect the

structure of a bioreactor community, we hypothesized that there would be groups of bacteria (or sub-

communities) associated with different phases of the bioreactor lifespan. To test for grouping, all of the

genomes were cross-correlated (Figure 4A). The resulting heatmap revealed four distinct clusters (A-D)

that are highly correlated. Cluster A was the largest, with 52 genomes, while Clusters B-D had 25, 24, and

26 genomes respectively.

To better examine the clustering of the genomes in relation to the different time-points, we ran

nonmetric multidimensional scaling (nMDS) analyses on the abundance data (Figure 4B). The nMDS

projection showed that genome groups are strongly associated with specific time-points: Group A is

associated with the inoculant source biomass at D0 and D166, while Group C is associated with the

nascent anammox community at D82. Group B is associated with the times of destabilized anammox

performance (Days 284-328), and Group D is associated with the mature, stable anammox community at

D437. *Brocadia* is part of Group D, although its location on the nMDS projection is skewed to the left

because of its high abundance throughout most of the experiment. Group A dominated at Days 0 and 166,

but was highly reduced at other times (Figure 4C). Group B dominated at D328, and maintained a similar

abundance at all other time points. Group C was mostly unique to D82, although a few of its members

remained in the bioreactor after the crash at low abundance. Group D bacteria showed little change up to

D284 (except a spike at D82), after which they increased in abundance.

It is interesting to note that the nascent anammox community is different from that of the

destabilized and the mature anammox communities. Because the nascent anammox community was

supplemented by a source inoculant biomass amendment, we cannot resolve a linear trajectory for the

microbial community between the initial and final states. Groups B and D, while distinct, share many

similarities, and the majority of the genomes associated with Group B were still present in the bioreactor

on D437.

For all subsequent analyses, we split the genomes into two groups: those that are associated with

the mature anammox community at D437 (Anammox Associated, AA, nMDS groups B and D), and those

that are not (Source Associated, SA, nMDS groups A and C). The AA community includes all of the

genomes that are present at D437 while the SA community includes the rest of the genomes that are not

present at D437. Some of these genomes are associated with the sludge amendments, and some are

associated with the nascent anammox community; at no point is there a community exclusively comprised

of SA genomes. The relative abundance of each group by time aligned with the previous analysis (Figure

4C). Since *Brocadia* dominated the community from D82 and onwards, it was removed from group D for

the purpose of this comparison.

Metabolic profiles. For the purpose of analyzing the metabolic potential of the microbial community we

evaluated only genomes with > 70% completeness (n = 88). Using Hidden Markov Model (HMM)

searches of the KEGG database, we checked for the presence of genes (with KO number) and calculated

KEGG module completeness[27,28]. The genomes were clustered by KO presence/absence (Supplemental

Figure 2) and their module completeness (Figure 5). The clustering by the two methods resulted in similar

groupings.

The module clustering resolved five groups (α, β, γ, δ, ε) (Figure 5A). Groups α and β contain

more anammox-associated genomes (90% and 60% respectively) while groups γ, δ, and ε contain 65%,

70% and 60% of source-associated genomes. The taxonomy of the bacteria also strongly influenced the

clustering (Figure 5B). Group α is composed solely of Gram (+) bacteria, while Group β is composed of

Microgenomates (Candidate Phyla Radiation (CPR) bacteria). Group γ is composed entirely of Gram (-)

bacteria (including *Brocadia*), Group δ is composed of Ignavibacteria and Bacteroidetes (other members

of these phyla were clustered in Group γ). Only Ignavibacteria of Group δ are associated with the AA

group, so further analysis of the group refers to them and not the Bacteroidetes. Group ε is composed of

Proteobacteria.

Based on the KEGG module clustering, we reconstructed the representative metabolisms of the

groups (Figure 6). We used a module completeness threshold of 67% per genome, and considered it

representative if it was complete in >50% of its members. Group δ is not represented since it diverged

from group γ by auxotrophies in several modules (Figure 5A, red rectangle). The *Brocadia* metabolism is

shown in Supplemental Figure 3.

While module completeness was used for most of the analyses, in several cases it was not

sufficient (e.g., overlap between modules, no module for path). In the cases of oxidative phosphorylation,

fermentation, carbon fixation, several amino acid synthesis pathways, and nitrogen metabolism we

analyzed gene presence manually.

Nitrogen cycling. We evaluated the quality genomes for the presence of all genes related to nitrogen

metabolism appearing in KEGG (Figure 7). Four additional HMMs were added for anammox genes

(hydrazine synthase subunit A (hzsA), hydrazine oxidoreductase subunit A (hzoA)), and nitrification

(nitrite oxidoreductase subunits nrxA and B)[29]. For the latter, the similarity of the gene to the nitrate

reductase narGH was taken into consideration.

With the exception of two CPR, all of the genomes in the bioreactor contained genes encoding

assimilation of ammonia into glutamate (Figure 7A). More than half (49) of the bacteria could reduce

nitrate, and the same number could further reduce nitrite to nitrogen monoxide (NO), however only 26

bacteria could do both steps. The most commonly-found gene encoding for nitrate reduction was narGH;

niK was more common than nirS (36 and 19 occurrences, respectively). The remaining steps of

denitrification were encoded in a smaller number of genomes. The nrxAB gene was only identified in two

genomes, one of which was *Brocadia.*

One-step DNRA was identified in 22 genomes, predominantly with nrfAH. While ammonia

assimilation and nitrate reduction were fairly similar in the AA and SA bacteria, DNRA was more

common in AA and denitrification beyond nitrite in the SA genomes (Figure 7C).

Supporting bacteria could improve bioreactor performance if they remove nitrate (nitrate

reducers) and excess nitrite, but they could be detrimental if they compete with anammox for nitrite

(DNRA and denitrification from nitrite). To check for changes in the abundances of these groups, we

classified bacteria by the presence of genes encoding for DNRA, denitrification, or nitrate reduction

(Figure 7B). Some bacteria classified as denitrifiers or DNRA also encoded nitrate reduction. A few

genomes in the D0 sample encoded both denitrification and DNRA, but their abundances were negligible.

Brocadia had genes required for DNRA but, given the overall bioreactor performance, was expected to be

primarily performing anammox for energy generation. DNRA could potentially be used by *Brocadia* for

detoxification by cycling potentially toxic excess nitrite back to ammonium where it could then

participate in the anammox reactions[18,25].

In the inoculant source community, the nitrate reducers were the most dominant group (38%),

with similar amounts of denitrifiers and DNRA (26% and 25% respectively). The abundance of anammox

was consistent with the bioreactor performance (Figure 1). The denitrifying group of bacteria decreased in

relative abundance to 8% around D284. On the other hand, bacteria capable of DNRA were relatively

abundant throughout the bioreactor start up. Most notably, these bacteria dominated the bioreactor during

its destabilization, reaching 48% at D328, compared to 23% for the anammox bacteria. An increase of

bacteria capable of DNRA was consistent with the bioreactor performance data which showed a decline

in the amount of ammonium consumed relative to overall bioreactor performance. At this time period the

following four DNRA bacteria were highly abundant: (LAC_IGN05, LAC_CHLX01, LAC_CHLX10,

and LAC_CHLX09). Three of the four are Group B bacteria, and one is Group D. All four bacteria

showed an increase in relative abundance between D284 and D328. Three of the four also showed

increased replication rates just before the onset of the crash, as mentioned above. The two other abundant

bacteria (apart from the anammox bacterium) were LAC_BAC23 (nitrate reducer) and LAC_PROT27

(denitrifier). The former was among the most abundant when the community is not SA dominated, while

the latter was always one of three most abundant bacteria. These 7 bacteria constituted >75% of the

community at D328.

Carbon fixation. Several bacteria (n =12) in the community are potentially capable of carbon fixation via

the Wood-Ljungdahl pathway or the Calvin cycle. *Brocadia* was confirmed as a primary producer, fixing

carbon via the Wood-Ljungdahl pathway using energy from the anammox pathway. The other ten

bacteria had genes for reduction of nitrogen compounds. To confirm that these bacteria are likely

autotrophs, we checked for genes conferring the ability to use inorganic electron donors. Three of these

bacteria had no potential electron donor and therefore were classified as heterotrophs. The remainder had

genes for oxidizing sulfide or hydrogen and were classified as potential autotrophs. Of these nitrate

reducing bacteria (n = 8), only one was relatively abundant after D166, and increased in abundance

between D284 and D328.

LAC_PROT27 can fix carbon by the Calvin cycle, is a denitrifier, and can possibly oxidize

sulfide to sulfite (dsrAB are present in the genome). This bacterium was among the most abundant at all

time-points; it increased significantly in abundance between D284 and D328 and the increase continued

to D437. However, the replication rate of the bacterium decreased from D166 onwards (Supplemental

Table 3), so it is not likely competing with, or destabilizing *Brocadia*.

Electron transfer. Apart from nitrogen reduction, another common anaerobic respiration pathway within

the community is acetate fermentation (genes detected in 60% of the genomes). This process was much

more common in AA (69%) bacteria than in SA (51%) bacteria. Ni-Fe Hydrogenase was present in 31%

of the genomes, but was most common among the Chloroflexi of group α (87% and 48% of all

occurrences of hydrogenases found) (Figure 6B).

290   The majority of bacteria in the bioreactor are potentially facultative aerobes (58%). All had high

affinity complex IV, which differed between AA and SA bacteria. In the AA bacteria, the bd type was

found in all aerobic members of group α (one also has a cbb3 type) and the Ignavibacteria, and the cbb3

type occurred mostly in Proteobacteria. For the SA bacteria, the cbb3 type was found in 24/25 aerobes

and the bd-type was only found in 6/25 (only in one bacterium it is the sole variant). Complex III, which

is also essential to aerobic respiration, was only found in 14 Proteobacteria, one Actinobacterium, and one

Chloroflexi. It is possible that other bacteria have an alternative Complex III[30] that cannot be found by

current KEGG annotations. Complexes I/II were found in nearly all of the bacteria, except CPR. Only

five bacteria lacked the F-type ATPase; two had the V-type ATPase instead.

Central carbon metabolism. It is likely that nearly all bacteria (98%) can oxidize sugar by glycolysis

(Figure 6A and E), while fewer bacteria (69%) had the pentose phosphate pathway (PPP). Acetyl-CoA

could be synthesized from pyruvate (90% general, 98% AA, and 81% SA), or by beta-oxidation (49%

general, 57% AA, and 43% SA). The majority of bacteria had the full TCA cycle (84%, or 88% after

excluding CPR). A possible major carbon source for the bacteria in the bioreactor are amino acids (aa.),

with 95% being able to incorporate aa. into their central carbon metabolism. The most common aa.

(aspartate) could be converted into oxaloacetate and fed into the TCA cycle. Three aa. (serine, alanine,

and cysteine) could be converted into pyruvate. Of these, only cysteine conversion is unidirectional, so

aa., as a carbon source, cannot be ascertained. Group α had additional genes that support a reliance on

proteins for their metabolism (Figure 6B). They also had a set of peptidases, as well as multiple

transporters covering all forms of aa., peptides, and polyamines.

Some metabolic groups could use aa. as precursors for synthesis of other metabolites. Glutamate

and histidine could be converted to PRPP, and glutamine to pyrimidines (Figure 6A). Groups ɣ and ε

could use aspartate to synthesize NAD^+, and glutamine to synthesis IMP (Figure 6C). NAD^+ and IMP

could not be synthesized by all of the bacteria, indicating that there are potential metabolic

interdependencies in the community. Members of group ε (Figure 6D) could use leucine as a precursor to

acetyl-CoA, lysine for acetoacetyl-CoA, glutamate for glutathione, and chorismate for ubiquinone. The

last two could only be synthesized by group ε, indicating additional potential metabolic interdependencies

in the community.

Comparing AA and SA. To examine why certain bacteria were enriched in the bioreactor while others

were removed, we compared the synthesis of metabolites to the utilization of nutrients in the bioreactor.

For synthesis we checked 24 KEGG modules for aa., 16 modules for vitamins or cofactors, and 11

modules for lipids and fatty acids. For nutrient utilization we looked at 52 modules of transporters. A

difference larger than 10% in the ratio of bacteria with or without a complete module was considered

relevant.

In all synthesis categories, SA bacteria had higher completeness ratios in the majority of the

modules investigated (14 of 24 aa. modules, 13 of 16 vitamins and cofactor modules, and 8 of 11 lipid

and fatty acid modules). The transportation modules showed an opposite trend, with 38 of 50 modules

having higher completeness ratios in AA bacteria.

This comparison shows that the selective driver in the anammox community is the ability of the

bacteria to acquire nutrients from the environment, rather than the ability to synthesize them. The larger

ratio of bacteria with auxotrophies in the AA bacteria hints of a greater reliance on other members of the

community.

Metabolic interdependencies. The bacteria in the AA community maintain a complex metabolic system. In the mature functioning bioreactor, *Brocadia* is (almost) the only primary producer present. It is also the only bacterium capable of synthesizing vitamin B12. For most other metabolites (vitamins, and cofactors) the possible metabolic interdependencies[31] are less straightforward (Supplemental table 4). Seven of 20 aa. could be synthesized by the majority of all metabolic groups (Figures 5-6). Members of group δ had the largest set of auxotrophies, lacking the genes conferring the ability to synthesize eight aa. The four other aa. could be synthesized by most group ε members and a few of the group ɣ members. Only a single cofactor (CoA) could be commonly synthesized by all groups. Group α had auxotrophies for most other cofactors, with the exception of pimeloyl-ACP (in 45% of members). Most other vitamins and cofactors were commonly synthesized by only a single group, usually group ε. With lipids and fatty acids, many modules were irrelevant to compare since group α differed from all other groups as a solely Gram (+) bacteria, while the rest are all Gram (-). However, even after the Gram (-) specific modules are excluded, group α still had multiple auxotrophies. *Brocadia* and group ɣ also had few commonly complete modules.

When combining all of the above data, we found that groups ɣ and ε both had mutualistic associations with *Brocadia* (Figure 8). Group ε potentially provided more metabolites to *Brocadia* than it received whereas groups α and δ seemed to gain more from *Brocadia* than they provided. Interestingly, four members of group α and one member of group δ were identified as the possible cause of the performance destabilization.

By the end of the experiment (D437) when bioreactor performance had stabilized, members of group α had become the second most abundant group after *Brocadia*. The ten most abundant bacteria at this point included four members of group δ and three members of group ε. Comparing these relative abundances to bacterial abundances during lowest bioreactor performance (D328), we found that *Brocadia* and group ε were reduced in abundance by about 50%, while groups α and δ were increased by 70% and 100%, respectively.

Discussion

In this study we present an in-depth analysis of the development of an anammox community from

seed to stable state (through several perturbations) in an anaerobic membrane bioreactor. By combining

several methodologies, we were able to gain important insights into the dynamics and interactions of

more than 100 species in the bioreactor community.

Previous studies have discussed a potential core anammox community[12–16]. With the exception of

very few studies, all such work has been conducted with single gene markers. Our metagenomic analysis

of an anammox community is the largest to-date and thus expands the ability to test this hypothesis. Our

results support the existence of a core community, while identifying factors that differentiate

communities. The high similarity between bacteria originating from three distinct anammox

bioreactors[18,24] strongly suggests a global core anammox microbial community. In the construction of the

phylogenetic tree we used >3000 reference genomes originating from diverse environments. Even with

the sheer number and diversity of sources, we found that the anammox community forms distinct clades

at the species level. More than half of the bacteria did not have species level relatives, and an additional

26% only had a relative found in our anammox bioreactor or in a previous anammox study[18,24]. Together,

nearly 80% of the bacteria are unique to anammox bioreactors, so it is clear that the anammox bioreactor

selects for a unique set of bacteria. Parameters that increased the differences between communities are the

species of the anammox bacterium and the bioreactor configuration. Since both parameters relate to the

same bioreactor[18], we cannot conclude which has a stronger effect.

We identified several potential bacterial destabilizers of the anammox process. Analysis of

replication rates days prior to the destabilization event revealed that these destabilizing bacteria increased

their replication rates, while *Brocadia* nearly ceased replication. These results imply a causative nature to

the change. Genes conferring DNRA capability were detected in these bacteria, which would allow them

to compete with *Brocadia* for nitrite. This supposition is consistent with the bioreactor performance

which exhibited decreased nitrogen removal and increased ammonium in the effluent during this period.

The dominating bacteria during bioreactor malperformance were heterotrophs. In full-scale anammox

bioreactors where influent organic carbon is essentially ubiquitous, heterotrophic dominance could

continue indefinitely without some sort of active countermeasure. Therefore, future research should target

the inhibition of potential destabilizing heterotrophs.

A broader investigation of metabolic interdependencies within the community shed light on the

stability of the anammox community. *Brocadia* is the source of organic material in the community, but

obtains essential metabolites from community members, especially Proteobacteria. This forms a basis for

a mutual symbiotic relationship. On the other hand, Chloroflexi, comprising the largest group of bacteria

besides *Brocadia*, receive numerous metabolites while apparently providing few in return. They are

characterized by an array of extracellular proteases and amylases, likely used to breakdown the

extracellular matrices formed by *Brocadia*. Chloroflexi, as a group, are most associated with anammox

bacteria and form a large fraction of the core community. They also account for the majority of the

destabilizing bacteria. Together, the results point to a parasitic symbiosis. Further investigation into these

relations is warranted.

While anammox bacteria generate sufficient organic carbon to support the growth of its co-

occurring heterotrophic microorganisms, the tipping point between stable and unstable operation and the

factors that control it have not been fully identified. Input changes may be able to restore anammox

activity, but this is just an empirical solution. Our findings improve the understanding of nitrogen-cycling

within an anammox bioreactor and advance the comprehensive control of this promising technology.

However, further work is needed to elucidate the precise mechanisms that control community

interactions.

Methods

Bioreactor operation. A laboratory-scale, anaerobic membrane bioreactor with a working volume of 1L

was constructed and operated for over 440 days (Supplemental Figure 4). The bioreactor was originally

inoculated with approximately 2 g VSS L^{-1} of biomass from a pilot-scale deammonification process

treating sidestream effluent at San Francisco Public Utilities Commission (SFPUC) in San Francisco, CA.

The bioreactor was re-inoculated with similar concentrations of biomass from the same source on Days

147 and 203. Synthetic media containing ammonium, nitrite, bicarbonate, and trace nutrients (meant to

mimic sidestream effluent at a municipal wastewater treatment plant) was fed to the bioreactor

(Supplemental Table 4). For the first 154 days of operation, the bioreactor was kept under nitrite-limiting

conditions to prevent inhibitory conditions, and influent ammonium and nitrite concentrations ranged

from 200-300 mg N L^{-1} and 100-300 mg N L^{-1}, respectively. On Day 154, ammonium and nitrite

concentrations were adjusted to the anammox stoichiometric ratio, 1:1.32. Afterwards, influent

ammonium and nitrite concentrations were maintained at this ratio and ranged from 200-500 mg N L^{-1}

and 265-660 mg N L^{-1}, respectively. On Day 353, influent concentrations of copper, iron, molybdenum,

and zinc were increased based on literature suggestions[21–24].

424   The bioreactor was operated in a continuous flow mode. For the first 145 days, the hydraulic

retention time (HRT) was maintained at 48 hours; afterwards it was reduced to 12 hours. No solids were

removed from the bioreactor for the first 100 days of operation; afterwards, the solids retention time

(SRT) was reduced to 50 days. A polyvinylidene fluoride hollow fiber membrane module with a 0.4 µm

pore size and total surface area of 260 cm^2 (Litree Company, China) was mounted in the bioreactor.

Temperature was maintained at 37° C with an electric heating blanket (Eppendorf, Hauppauge, NY).

Mixing was provided by an impeller at a rate of 200 rpm. Mixed gas was supplied continuously to the

bioreactor (Ar:CO_2 = 95:5; 50 mL min^{-1}) to eliminate dissolved oxygen and maintain pH at 7.2. Influent

and effluent concentrations of ammonium, nitrite, and nitrate were measured approximately every other

day using HACH test kits (HACH, Loveland, CO), as described in the manufacturer's methods 10031,

10019, and 10020, respectively.

Biomass collection and DNA extraction. Biomass samples were extracted via syringe from the

bioreactor every 2-10 days, flash frozen in liquid nitrogen, and stored frozen at -80 °C until use. Genomic

DNA was extracted from the samples using the DNeasy PowerSoil Kit (Qiagen, Carlsbad, CA), as

described in the manufacturer's protocol. The concentration and purity of extracted DNA was measured

with a NanoDrop Spectrophotometer (Thermo Scientific, Waltham, MA). The concentration of genomic

DNA in all samples was normalized to 10 ng/µL with nuclease-free water (Thermo Scientific, Waltham,

MA). All genomic DNA samples were stored at -20 °C until use.

Metagenomic sequencing, assembly, and binning. Genomic DNA samples from six time-points were

sent to the Joint Genome Institute (JGI) in Walnut Creek, CA for sequencing on the Illumina HiSeq 2500

1T sequencer (Illumina, San Diego, CA). Resulting sequences from each time-point were processed

separately, following the ggKbase SOP (https://ggkbase-help.berkeley.edu/overview/data-preparation-

metagenome/). In summary, Illumina adapters and trace contaminants were removed (BBTools, GJI) and

raw sequences were quality-trimmed with Sickle[32]. Paired-end reads were assembled using IDBA_UD

with the pre-correction option and default settings[33]. For coverage calculations, reads were mapped with

bowtie2[34]. Genes were predicted by Prodigal[35] and predicted protein sequences were annotated using

usearch[36] against KEGG, UniRef100, and UniProt databases. The 16S rRNA gene and tRNA prediction

was done with an in-house script and tRNAscanSE[37] respectively. At this point, the processed data was

uploaded to ggKbase for binning.

Manual binning was performed using the ggKbase tool. The binning parameters for binning were

GC% and coverage (CV) distribution, and phylogeny of the scaffolds. Quality of the manual bins was

assessed by the number of Bacterial Single Copy Genes (BSCG) and Ribosomal Proteins (RP) found in

each bin (aiming at finding the full set of genes, while minimizing the multiple copies). In addition to

manual binning, automated binning was done using four binners: ABAWACA1[38], ABAWACA2,

CONCOCT[39], and Maxbin2[40]. For all, the default parameters were chosen.

All bins from both automatic and manual binning tools were input into DASTool[41] to iterate

through bins from all binning tools and choose the optimal set of bins. checkM was run to analyze

genome completeness[42]. The scaffold-to-bin file created by DASTool was uploaded back to ggKbase and

all scaffolds were rebinned to match the DASTool output. Each of the new bins were manually inspected

and scaffolds that were suspected to be falsely binned were removed.

After we inspected the first round of binning, we decided to improve the high coverage bins, by

subsampling the read file, followed by the same SOP as above[43]. In addition, refinement of the *Brocadia*

Genome bins was done with ESOMs[44] (Supplemental methods).

Post binning analysis. Unique representative genomes were determined by the dereplication tool, dRep[45],

using a 95% threshold for species level clustering. Within each cluster, the representative genome was

chosen based on their completeness, length, N50, contamination, and strain heterogeneity. In several

clusters with higher heterogeneity, a second strain was chosen (Supplemental Table 1). The strain

threshold was set at 2% difference (but lower than 5%).

All the representative and strain genomes were curated by correcting scaffolding errors

introduced by idba_ud, using the ra2.py program[38]. Following curation, the genomes were processed

again for gene calling and annotation (see above for details). Analysis of replication rates at different

time-points was performed with the iRep program[46] using the default parameters.

Both raw reads and genomes were submitted to the National Center for Biotechnology

Information (NCBI) Genbank, under project accession number PRJNA511011. In addition, the

Representative and strains genomes were uploaded to ggkbase as two separate projects

(https://ggkbase.berkeley.edu/LAC_reactor_startup/organisms and

https://ggkbase.berkeley.edu/LAC_reactor_strains/organisms).

Phylogenetic analysis and core anammox analysis. The taxonomic affiliation of each genome was

initially assigned in ggKbase. This was based on the taxonomic annotation of genes in the scaffolds. For

each hierarchical taxonomic level, the taxonomy was decided if at least 50% of genes had a known

taxonomic identification.

Phylogenetic analysis of the genomes (current study, Speth et al.[18], and Lawson et al.[25]) was

based on a set of 15 ribosomal proteins[47]. Each gene was aligned separately to a set of 3225 reference

genomes, followed by concatenation while keeping the aligned length of each gene intact. A preliminary

tree was created by adding the queried genomes to the reference tree using pplacer v1.1.alpha19[48] and a

set of in-house scripts. The tree was uploaded to iTOL[49] for visualization and editing. After initial

inspection we decided to reduce the tree in preparation of creating a maximum likelihood tree. Large

phyla with no representatives in an anammox sample were removed (approximately 1000 sequences). The

remaining sequences were aligned by MUSCLE[50] and a RAxML tree built in The CIPRES Science

Gateway V. 3.3[50,51].

For the analysis of phylogenetic distance between different anammox community members, we

used the APE package[52] in R[53,54] to extract the distance matrix. Species level distance was set at 5% of the

longest measured distance on the tree.

16S rRNA gene sequencing, processing, and analysis. DNA samples, taken at 56 timepoints across the

lifespan of the bioreactor, were sent to the Institute for Environmental Genomics at the University of

Oklahoma (Norman, OK) for amplification of the variable 4 (V4) region of the 16S rRNA gene, library

preparation, and amplicon sequencing. The full protocol was previously described in Wu et al. (Wu

2015). In summary, the V4 region of the bacterial 16S rRNA gene was amplified from DNA samples

using primers 515F (5'-GTGCCAGCMGCCGCGG-3') and 806R (3'-TAATCTWTGGVHCATCAG-5'),

with barcodes attached to the reverse primer. Amplicons were pooled at equal molality and purified with

the QIAquick Gel Extraction Kit (QIAGEN Sciences, Germantown, MD). Paired-end sequencing was

then performed on the barcoded, purified amplicons with the Illumina MiSeq sequencer (Illumina, San

Diego, CA).

Subsequent sequence processing and data analysis were performed in-house using MOTHUR

v.1.39.5, following the MiSeq SOP[55,56]. In summary, sequences were demultiplexed, merged, trimmed,

and quality filtered. Unique sequences were aligned against the SILVA 16S rRNA gene reference

alignment database[57]. Sequences that did not align to the position of the forward primer were discarded.

Chimeras were detected and removed. Remaining sequences were clustered into operational taxonomic

units (OTUs) within a 97% similarity threshold using the Phylip-formatted distance matrix.

Representative sequences from each OTU were assigned taxonomic identities from the SILVA gene

reference alignment database[57]. Sequences that were not classified as bacteria were removed. Remaining

OTUs were counted, and the 137 most abundant OTUs (accounting for up to 99% of sequence reads

within individual samples) were transferred to Microsoft Excel (Microsoft Office Professional Plus 2016)

for downstream interpretation and visualization. The 137 most abundant OTUs were up loaded to figshare

(https://figshare.com/account/projects/59324/articles/7640396).

In order to correlate genome-based OTUs to 16S rRNA gene-based OTUs, 16S rRNA sequences

were extracted from the representative genomes and combined with the representative sequences from the

137 most abundant 16S rRNA gene-based OTUs. If a representative genome did not contain the V4

region of the 16S rRNA gene, the region was pulled from another genome in the same cluster. The

combined 16S rRNA sequences were aligned following the protocol described above, and those sharing at

least 99% average nucleotide identity were assumed to represent the same microorganism (Evans 2006,

Blast).

Community dynamics analysis. The paired sequence reads from all time-points were mapped to the set

of reference genomes using bowtie2[34], followed by a calculation of coverage (average number of reads

mapped per nucleotide) and breadth (% of genome that was covered by at least one read in the mapping)

for each genome per time-point[58]. The multiplication of the two values was then used to calculate the

estimated abundance. This was done to negate biases that can be created by repetitive sequences that

occur more often in very partial genome bins (i.e. only the repetitive sequences associated with the

genome bin are found in a given time-point).

Association between genomes was tested by calculating pairwise correlation for all genomes by

abundance. The Rho values (ranging from -1 to 1) were used to create a distance table (Euclidean

distance), followed by clustering with the ward.D method. The resulting clusters were marked A-D. To

test the association of genomes and clusters to time-points, we ran a nMDS analysis (non-parametric

MultiDimensional Scaling) with the genomes and the time-point. Each genome was colored by its

abundance cluster on the 2D projection of the nMDS.

For relative abundance changes, the estimated abundances of genomes were divided by the sum

of all estimated abundance values per time-point. For a clearer resolution of changes in the four

abundance groups, the *Brocadia* (part of group D) was presented separately.

Metabolic analysis. The functional profiles of the genomes were evaluated using KEGG KAAS[59], with

Hidden Markov Models for shared KEGG orthologies (KOs)[27,28,60]. From this, we received the KEGG

annotation (KO number) for all open reading frames and a completeness value for each KEGG module.

KO annotations that were questionable were removed from analysis.

From the KO list, we created a presence absence matrix (Jaccard index), and clustered using the

Complete method. From module completeness, we created a Euclidean distance matrix, followed by

clustering with the ward.D method. Based on module completeness clustering, we assigned genomes to

metabolic groups α-ε.

For each metabolic group a representative metabolic map was created. A module completeness

greater than 67% in at least half of the group members was considered as representative of the group.

Once the modules were selected, they were drawn and connected based on metabolic KEGG maps.

Additional reaction, complexes, and transporters were added according to KO presence (e.g., aa.

synthesis, oxidative phosphorylation complexes, flagellar motor, etc.).

For nitrogen metabolism, all relevant KOs were examined. For the purpose of this study, nitrate

reduction was considered as a separate path from denitrification/DNRA, since it could be the first step in

both, using the same enzymes. Denitrifying bacteria were considered as bacteria capable of full

conversion of nitrite to N_2. DNRA bacteria were considered as bacteria capable of conversion of nitrite to

ammonium using the nrfAH enzymes. No partial nitrogen process was considered for this paper, although

it is present, according to per step analysis.

Acknowledgements

This research was supported by the National Science Foundation through the Engineering Research

Center for ReInventing the Nation's Water Infrastructure (ReNUWIt) ECC-1028962. This material is also

based upon work supported by the National Science Foundation Graduate Research Fellowship under

Grant No. DGE 1106400. Any opinion, findings, and conclusions or recommendations expressed in this

material are those of the authors and do not necessarily reflect the views of the National Science

Foundation.

Author contribution

K.Y. and L.Z. supervised the study. L.Z., L.A-C., D.J., and K.Y designed the study. L.Z. built the

bioreactor. R.K. analyzed metagenomics data and wrote the manuscript. J.L. analyzed 16S rRNA data,

analyzed bioreactor performance, and wrote the manuscript. J.F.B supervised the metagenomics analysis.

W.Z. and K.Y. contributed to bioreactor maintenance and analysis, sampling and 16S rRNA data analysis.

All authors read the manuscript and contributed with inputs.

Note: The authors declare no competing financial interest.

References

1.   Mulder, A., Graaf, A. A., Robertson, L. A. & Kuenen, J. G. Anaerobic ammonium oxidation
discovered in a denitrifying fluidized bed reactor. *FEMS Microbiol. Ecol.* **16**, 177–184 (1995).

2.   Kuenen, J. G. Anammox bacteria: from discovery to application. *Nat. Rev. Microbiol.* **6**, 320–326
(2008).

3.   Sonthiphand, P., Hall, M. W. & Neufeld, J. D. Biogeography of anaerobic ammonia-oxidizing
(anammox) bacteria. *Front. Microbiol.* **5**, (2014).

4.   Connan, R. *et al.* Batch enrichment of anammox bacteria and study of the underlying microbial
community dynamics. *Chem. Eng. J.* **297**, 217–228 (2016).

5.   Strous, M. *et al.* Deciphering the evolution and metabolism of an anammox bacterium from a
community genome. *Nature* **440**, 790–794 (2006).

6.   Kartal, B. *et al.* Molecular mechanism of anaerobic ammonium oxidation. *Nature* **479**, 127–130
(2011).

7.   Paques. Anammox sustainable nitrogen removal. Available at:
https://en.paques.nl/mediadepot/1818a31cd232/WEBbrochureAnammox.pdf. (Accessed: 26th
September 2018)

8.   Lackner, S. *et al.* Full-scale partial nitritation/anammox experiences – An application survey. *Water*
Res. **55**, 292–303 (2014).

9.   Ali, M. & Okabe, S. Anammox-based technologies for nitrogen removal: Advances in process start-
up and remaining issues. *Chemosphere* **141**, 144–153 (2015).

10.   Jin, R.-C., Yang, G.-F., Yu, J.-J. & Zheng, P. The inhibition of the Anammox process: A review.
Chem. Eng. J. **197**, 67–79 (2012).

11.   Li, X., Klaus, S., Bott, C. & He, Z. Status, Challenges, and Perspectives of Mainstream Nitritation-

Anammox for Wastewater Treatment. *Water Environ. Res.* **90**, 634–649 (2018).

12.   Li, X.-R. *et al.* The bacterial diversity in an anaerobic ammonium-oxidizing (anammox) reactor

community. *Syst. Appl. Microbiol.* **32**, 278–289 (2009).

13.   Gonzalez-Martinez, A. *et al.* Bacterial community structure of a lab-scale anammox membrane

bioreactor. *Biotechnol. Prog.* **31**, 186–193 (2015).

14.   Gonzalez-Martinez, A. *et al.* Comparison of bacterial diversity in full scale anammox bioreactors

operated under different conditions. *Biotechnol. Prog.* **31**, 1464–1472 (2015).

15.   Gonzalez-Martinez, A. *et al.* Microbial community analysis of a full-scale DEMON bioreactor.

Bioprocess Biosyst. Eng. **38**, 499–508 (2015).

16.   Pereira, A. D., Cabezas, A., Etchebehere, C., de Lemos Chernicharo, C. A. & de Araújo, J. C.

Microbial communities in anammox reactors: a review. *Environmental Technology Reviews* **6**, 74–

93 (2017).

17.   Bagchi, S., Lamendella, R., Strutt, S., Van Loosdrecht, M. C. M. & Saikaly, P. E.

Metatranscriptomics reveals the molecular mechanism of large granule formation in granular

anammox reactor. *Sci. Rep.* **6**, 28327 (2016).

18.   Speth, D. R., in 't, M., Guerrero-Cruz, S., Dutilh, B. E. & Jetten, M. S. M. Genome-based microbial

ecology of anammox granules in a full-scale wastewater treatment system. *Nat. Commun.* **7**, 11172

(2016).

19.   Castro-Barros, C. M., Jia, M., van Loosdrecht, M. C. M., Volcke, E. I. P. & Winkler, M. K. H.

Evaluating the potential for dissimilatory nitrate reduction by anammox bacteria for municipal

wastewater treatment. *Bioresour. Technol.* **233**, 363–372 (2017).

20.   Tang, X., Guo, Y., Jiang, B. & Liu, S. Metagenomic approaches to understanding bacterial

communication during the anammox reactor start-up. *Water Res.* **136**, 95–103 (2018).

21.   van de Graaf, A. A. *et al.* Anaerobic oxidation of ammonium is a biologically mediated process.

Appl. Environ. Microbiol. **61**, 1246–1251 (1995).

22.   Chen, H., Yu, J.-J., Jia, X.-Y. & Jin, R.-C. Enhancement of anammox performance by Cu(II), Ni(II)

and Fe(III) supplementation. *Chemosphere* **117**, 610–616 (2014).

23.   Bi, Z., Qiao, S., Zhou, J., Tang, X. & Zhang, J. Fast start-up of Anammox process with appropriate

ferrous iron concentration. *Bioresour. Technol.* **170**, 506–512 (2014).

24.   Zhang, X. *et al.* Impacts of the heavy metals Cu (II), Zn (II) and Fe (II) on an Anammox system

treating synthetic wastewater in low ammonia nitrogen and low temperature: Fe (II) makes a

difference. *Sci. Total Environ.* **648**, 798–804 (2018).

25.   Lawson, C. E. *et al.* Metabolic network analysis reveals microbial community interactions in

anammox granules. *Nat. Commun.* **8**, 15416 (2017).

26.   Narita, Y. *et al.* Enrichment and physiological characterization of an anaerobic ammonium-oxidizing

bacterium ' Candidatus Brocadia sapporoensis'. *Syst. Appl. Microbiol.* **40**, 448–457 (2017).

27.   Burstein, D. *et al.* Major bacterial lineages are essentially devoid of CRISPR-Cas viral defence

systems. *Nat. Commun.* **7**, 10613 (2016).

28.   Anantharaman, K. *et al.* Analysis of five complete genome sequences for members of the class

Peribacteria in the recently recognized Peregrinibacteria bacterial phylum. *PeerJ* **4**, e1607 (2016).

29.   Anantharaman, K. *et al.* Thousands of microbial genomes shed light on interconnected

biogeochemical processes in an aquifer system. *Nat. Commun.* **7**, 13219 (2016).

30.   Sun, C. *et al.* Structure of the alternative complex III in a supercomplex with cytochrome oxidase.

Nature **557**, 123–126 (2018).

31.   Zomorrodi, A. R. & Segrè, D. Genome-driven evolutionary game theory helps understand the rise of

metabolic interdependencies in microbial communities. *Nat. Commun.* **8**, 1563 (2017).

32.   Joshi NA, F. J. N. Sickle: A sliding-window, adaptive, quality-based trimming tool for FastQ files

(Version 1.33) [Software]. Available at https://github.com/najoshi/sickle. (2011).

33.   Peng, Y., Leung, H. C. M., Yiu, S. M. & Chin, F. Y. L. IDBA-UD: a de novo assembler for single-

cell and metagenomic sequencing data with highly uneven depth. *Bioinformatics* **28**, 1420–1428

(2012).

34.   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**, 357–

359 (2012).

35. Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site identification.

BMC Bioinformatics **11**, 119 (2010).

36. Edgar, R. C. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**,

2460–2461 (2010).

37. Lowe, T. M. & Eddy, S. R. tRNAscan-SE: a program for improved detection of transfer RNA genes

in genomic sequence. *Nucleic Acids Res.* **25**, 955–964 (1997).

38. Brown, C. T. *et al.* Unusual biology across a group comprising more than 15% of domain Bacteria.

Nature **523**, 208–211 (2015).

39. Alneberg, J. *et al.* Binning metagenomic contigs by coverage and composition. *Nat. Methods* **11**,

1144–1146 (2014).

40. Wu, Y.-W., Simmons, B. A. & Singer, S. W. MaxBin 2.0: an automated binning algorithm to

recover genomes from multiple metagenomic datasets. *Bioinformatics* **32**, 605–607 (2015).

41. Sieber, C. M. K. *et al.* Recovery of genomes from metagenomes via a dereplication, aggregation, and

scoring strategy. (2017). doi:10.1101/107789

42. Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM: assessing the

quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res.*

25, 1043–1055 (2015).

43. Hug, L. A. *et al.* Critical biogeochemical functions in the subsurface are associated with bacteria

from new phyla and little studied lineages. *Environ. Microbiol.* **18**, 159–173 (2016).

44. Ultsch, A. & Mörchen, F. *ESOM-maps: Tools for Clustering, Visualization, and Classification with*

Emergent SOM. (2005).

45. Olm, M. R., Brown, C. T., Brooks, B. & Banfield, J. F. dRep: a tool for fast and accurate genomic

comparisons that enables improved genome recovery from metagenomes through de-replication.

ISME J. **11**, 2864–2868 (2017).

46. Brown, C. T., Olm, M. R., Thomas, B. C. & Banfield, J. F. Measurement of bacterial replication

rates in microbial communities. *Nat. Biotechnol.* **34**, 1256–1263 (2016).

47. Hug, L. A. *et al.* A new view of the tree of life. *Nat Microbiol* **1**, 16048 (2016).

48. Matsen, F. A., Kodner, R. B. & Armbrust, E. V. pplacer: linear time maximum-likelihood and

Bayesian phylogenetic placement of sequences onto a fixed reference tree. *BMC Bioinformatics* **11**,

538 (2010).

49. Letunic, I. & Bork, P. Interactive tree of life (iTOL) v3: an online tool for the display and annotation

of phylogenetic and other trees. *Nucleic Acids Res.* **44**, W242–5 (2016).

50. Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high throughput.

Nucleic Acids Res. **32**, 1792–1797 (2004).

51. Miller, M. A., Pfeiffer, W. & Schwartz, T. Creating the CIPRES Science Gateway for inference of

large phylogenetic trees. in *2010 Gateway Computing Environments Workshop (GCE)* (2010).

doi:10.1109/gce.2010.5676129

52. Paradis, E., Claude, J. & Strimmer, K. APE: Analyses of Phylogenetics and Evolution in R language.

Bioinformatics **20**, 289–290 (2004).

53. *RStudio Team (2015). RStudio: Integrated Development for R. RStudio, Inc., Boston, MA.*

54. R: The R Project for Statistical Computing. Available at: http://www.R-project.org. (Accessed: 26th

September 2018)

55. Schloss, P. D. *et al.* Introducing mothur: open-source, platform-independent, community-supported

software for describing and comparing microbial communities. *Appl. Environ. Microbiol.* **75**, 7537–

7541 (2009).

56. Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K. & Schloss, P. D. Development of a

dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the

MiSeq Illumina sequencing platform. *Appl. Environ. Microbiol.* **79**, 5112–5120 (2013).

57. Pruesse, E. *et al.* SILVA: a comprehensive online resource for quality checked and aligned

ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Res.* **35**, 7188–7196 (2007).

58. Olm, M. R. *et al.* Identical bacterial populations colonize premature infant gut, skin, and oral

microbiomes and exhibit different in situ growth rates. *Genome Res.* **27**, 601–612 (2017).

59. Moriya, Y., Itoh, M., Okuda, S., Yoshizawa, A. C. & Kanehisa, M. KAAS: an automatic genome

annotation and pathway reconstruction server. *Nucleic Acids Res.* **35**, W182–5 (2007).

60. Finn, R. D. *et al.* HMMER web server: 2015 update. *Nucleic Acids Res.* **43**, W30–8 (2015).

Figures

719
720

Figure 1 | Performance of the anaerobic membrane bioreactor. Influent and effluent concentrations of
ammonium, nitrite, and nitrate (all as N) (primary y-axis) within the anaerobic membrane bioreactor
performing anammox monitored over a period of 440 days. The influent did not contain nitrate, so
influent nitrate is not plotted. The nitrogen removal rate (NRR), is plotted against the secondary y-axis.
Sampling time points for metagenomes are indicated with purple stars below the x-axis.

A

B

	WS6		Deltaproteobacteria
	Acidobacteria		Deinococcus Thermus
	Actinobacteria		Bacteroidetes
	Armatimonadetes		Proteobacteria
	Parcubacteria		Chloroflexi
	Omnitophica		Ignavibacteria
	Nitrospirae		Planctomycetes
	Microgenomates		

Current (n= 127)
Speth et al (n= 22)
Lawson et al (n= 14)

Tree scale: 0.1

Figure 2 | Phylogenetic analysis of three anammox microbial communities. (A) A maximum
likelihood tree based on the alignment of 15 concatenated ribosomal proteins. In the construction of the
tree 3225 reference sequences were used, with genomes from current and previous genome-centric studies
on anammox communities. Genomes from the current anammox community are marked with a red dash,
genomes from two previously studied communities; Speth et al. and Lawson et al., are marked with green
and blue dashes, respectively. (B) relative abundance of major phyla in the three microbial communities.
Current community reference data was calculated from day 437 only. The relative abundance *Brocadia*
sp. comprises nearly all of the abundance attributed to phylum Planctomycetes (with small contributions
from other members of the phylum). The most abundant phyla (Chloroflexi, Ignavibacteria, and
Proteobacteria) consistently account for >70% of the communities. The phyla colors follow the ggkbase
color scheme and the major phyla are shown in the legend.

Phylum	Class	Order	Genus
Unmatched			
Acidobacteria	-	-	-
Bacteroidetes	-	-	-
Bacteroidetes	Sphingobacteria	Sphingobacteriales	-
Chloroflexi	-	-	-
Chloroflexi	Anaerolineae	-	-
Deinococcus-Thermus	Deinococci	Deinococcales	Truepera
Ignavibacteriae	Ignavibacteria	-	-
Nitrospirae	Nitrospira	Nitrospirales	Nitrospira
Planctomycetes	-	-	-
Planctomycetes	Planctomycetia	Brocadiales	Brocadia
Proteobacteria	-	-	-
Proteobacteria	Alphaproteobacteria	-	-
Proteobacteria	Betaproteobacteria	Burkholderiales	-
Proteobacteria	Betaproteobacteria	Rhodocyclales	-
Proteobacteria	Gammaproteobacteria	-	-
Proteobacteria	Gammaproteobacteria	Xanthomonadales	-
Verrucomicrobia	-	-	-

764
765
Figure 3 | Relative abundances of bacterial taxa over the lifespan of the bioreactor. Relative
abundances of bacterial taxa, as identified by metagenomic and 16S rRNA gene sequencing, are plotted
against the primary y-axis. Results derived from metagenomic sequencing are indicated with a purple star
below the x-axis; all remaining results are derived from 16S rRNA gene sequencing. For visual clarity,
sequencing results falling within three days of each other have been merged. "Unmatched" includes the
OTUs and genomes that were not able to be matched across the two sequencing platforms. The similar
relative abundance profiles at shared time points across metagenomic and 16S rRNA gene sequencing
platforms (highlighted in the black boxes) provided us with the confidence to extrapolate high-resolution
relative abundance profiles of our representative genomes from our 16S rRNA gene sequencing efforts.
775
776
777
778

779
780
Figure 4| Analysis of the bioreactor community dynamics by the estimated abundance of the
bacteria. (A) clustering heatmap of bacteria based on pairwise cross correlations in the six time points
(matrix values are Rho values). Color scales mark high positive correlation in green and high negative
correlation in brown. The row and column dendrograms are identical. The row dendrogram shows the
calculated distance between the clusters with a dashed red line marking the threshold distance for

clustering. Three columns of annotations between the row dendrogram and heatmap; Core- labels bacteria
related to previously studied bacteria (see relevant section); AA/SA- association of bacteria with either
mature anammox or source inoculum; Group- assigned group based on correlation and clustering. (B)
Two dimensional nMDS projection of bacteria and time points, showing the association of the bacteria
(and abundance groups to certain time points). Each colored dot represents the centroid of a bacterium,
with colors matching the abundance group. The location of *Brocadia* is marked with a red star. (C)
Relative abundance of Groups A-D by time points. *Brocadia* (of group D) is presented apart from the
group to more easily show changes in the other group members.

Figure 5| Metabolic profiling of bacterial community based on KEGG module completeness. (A)
heatmap showing the reciprocal clustering of genomes (rows) and KEGG modules (columns). The
heatmap is based on a Euclidean distance matrix and clustering with the ward.D method. Genome
clustering resulted in 5 clusters (groups α-ε). Module clustering resulted in 11 clusters (blocks 1-11).
Black rectangles on heatmap show module blocks that have increased completeness in a group of bacteria
(compared to the others), and red rectangles show decreased completeness. The three columns on the left
of the heatmap denote core association, AA/SA division, and abundance grouping respectively. B)
Relative abundance by phyla of members in the metabolic clusters.
809
810

Figure 6| Representative metabolic maps of bacterial groups in the bioreactor. To prevent
redundancy, the metabolism is presented in a nested approach with each panel showing only paths unique
to the relevant metabolic group. Two exceptions are group β (all detected paths are shown), and group δ.
The latter is not presented here since it shares all paths with group ɣ and only differs by auxotrophies. (A)
Metabolic map of paths that are common to all bacteria in the bioreactor (except Microgenomates and
Brocadia sp.). The vast majority of bacteria in the bioreactor are heterotrophs, capable of carbohydrate-
based metabolism (glycolysis, pentose phosphate pathway) and amino acid-based metabolism. Some
bacteria can respire oxygen, but can also ferment (acetate/alanine). (B) Paths unique to group α. The
bacteria have genes for hydrogen oxidation, supporting anaerobic growth, as well as genes for oxidative
phosphorylation with cytochrome BD complex. These bacteria have a cassette of extracellular proteases
and decarboxylases, paired with a wide array of transporters. The bacteria are also potentially capable of
synthesizing long chain isoprenoids. (C) Paths found in Gram (-) bacteria (groups ɣ, δ, and ε). Most paths
are related to fatty acid and lipid synthesis. Several important precursors (chorismate and IMP) can
potentially be synthesized by these bacteria. Motility is also a common feature in these bacteria (via a
flagellar motor) (D) Unique paths of group ε (Proteobacteria). This group has the potential to synthesize
multiple vitamins and cofactors (biotin, pyridoxal, glutathione, etc.), as well as several aa. (tyrosine,
phenylalanine, proline). Another unique feature is the multiple secretion systems present in the bacteria.
(E) Metabolic profile of CPR bacteria (Microgenomates). The bacteria are obligate anaerobes that ferment
pyruvate. They can only utilize carbohydrates as their carbon source. Some of the bacteria in this group
might also be able to synthesize long chain isoprenoids, in the same path as group α.

832
833
834

835
836 **Figure 7| Nitrogen cycle in the anammox bioreactor. (**A) The steps in the nitrogen cycle are color
837 coded by their association to different types of metabolism. The number of bacteria with genes encoding a

given step is listed and the bar chart depicts the ratio of bacteria within the AA/SA groups associated with the step. (B) Changes in relative abundance of bacterial groups by their nitrogen metabolism. Only a single path was assigned to each genome for the purpose of this analysis. Since nitrate reduction is also considered a first step in denitrification and DNRA, is was assigned only when other paths were not present. Bacteria with no complete metabolic path are depicted in light grey. Anammox is the dominant nitrogen metabolic path at Days 82, 284, and 437. This matches the bioreactor performance monitoring (Figure 1). At times when the source community is predominant (Days 0 and 166), nitrogen reduction is the most common metabolic path, followed by DNRA. During the period of bioreactor destabilization (Day 328) the DNRA bacteria dominate the community. (C) The number of bacteria in which a given metabolic path was detected. The bars are divided by AA/SA association. Unlike panel B, overlap of functions was allowed for the genome counts. The largest group are the nitrate reducers, followed by denitrifiers and DNRA. Denitrifiers are more common among the SA group, while DNRA are more common among AA group.

Amino acids

Peptides

Vitamins/Co-factors

Lipid/Fatty acids

Figure 8| Potential metabolic hand-offs between the MO groups in the anammox bioreactor. Arrows were assigned according to absence of ability to synthesis a metabolite and connect to all groups that do have the ability (meaning there is redundancy in arrows). The arrowhead points at the group that receives the metabolite. The width of the arrow is proportional to the ratio of metabolites of a given type that are provided; amino acids - 20 metabolites; Peptides - deduced from proteases and transporters (Figure 6B); Vitamins/Co-factors - 10 metabolites; Lipids/Fatty acids - 7 metabolites. The size of each group is proportional to their relative abundance at Day 437. Group β is not shown since the assumption is that its members obtain all of their nutrients and metabolites from their host. Overall, groups α and δ receive the most metabolites and group ε the least. Group δ has the highest number of aa. synthesis auxotrophies and can potentially acquire these from many other community members. Group ε has only a single auxotrophy in vitamin/Co-factor synthesis while most other groups have multiple auxotrophies (group α capable of only a single metabolite). *Brocadia* sp. is the only bacterium capable of vitamin B12 synthesis.

    Tables

868
869   **Table 1 | Genome parameters for the representative bacteria from the anammox bioreactor**
870

Genome	Genome code (ggkbase)	Genome length (bp)	GC%	Contigs	ORF	Longest contig	Completeness (%)
LAC_ACD01	anamox1_Acidimicrobiia_62_4_curated	298722	58.08	173	496	5546	65.54
LAC_ACD02	anamox1_Acidobacteria_62_5_curated	3111074	61.59	565	3251	20877	65.54
LAC_ACD03	anamox1_Pyrinomonas_methylaliphatogenes_54_8_curated	2071852	54.23	87	1989	105081	95.69
LAC_ACD04	anamox3_Acidobacteria_71_4_curated	1334157	69.62	772	1759	7429	39.71
LAC_ACD05	LAC_NA06_Candidatus_Solibacter_usitatus_62_12_curated	4513182	61.85	167	4021	141967	91.38
LAC_ACD06	LAC_NA07_Acidobacteria_70_38_curated	3262818	69.91	40	2917	270970	94.83
LAC_ACD07	LAC_NA07_Candidatus_Solibacter_usitatus_59_6_curated	1779371	57.94	742	2242	9528	56.02
LAC_ACT01	anamox2_Actinobacteria_65_5_curated	2799571	63.61	662	3147	20411	67.46
LAC_ACT02	LAC_NA07_Actinobacteria_71_14_curated	3047960	70.65	64	3117	193413	93.97
LAC_ACT03	LAC_NA07_Actinobacteria_74_18_curated	1235923	73.53	141	1261	48493	64.86
LAC_ACT04	LAC_NA07_Actinotalea_fermentans_75_19_curated	2905150	75.39	31	2745	349240	100
LAC_ARCH01	anamox2_Methanosarcina_thermophila_41_9_curated	2959285	41.19	82	2703	143092	35.55
LAC_ARM01	LAC_NA06_Fimbriimonas_ginsengisoli_61_14_curated	2673573	61.04	54	2542	245447	88.45
LAC_BAC01	anamox1_Bacteria_33_9_curated	775494	32.38	45	793	69755	70.85
LAC_BAC02	anamox1_Bacteria_45_8_curated	2579669	45	166	2221	61316	87.77
LAC_BAC03	anamox1_Bacteria_50_18_curated	820566	49.55	22	874	204925	77.9
LAC_BAC04	anamox1_Bacteria_53_17_curated	3465172	53.24	12	2895	903458	100
LAC_BAC05	anamox1_Bacteria_55_18_curated	4100779	54.56	216	3365	90642	98.28
LAC_BAC06	anamox1_Bacteria_57_32_curated	2939326	56.91	76	2571	186230	95.69
LAC_BAC07	anamox1_Bacteria_57_5_curated	3332363	56.3	813	3846	19658	68.62
LAC_BAC08	anamox1_Bacteria_57_9_curated	3469651	57.28	74	3012	281471	93.65
LAC_BAC09	anamox1_Bacteria_64_7_curated	3988790	63.42	306	3679	75294	93.03
LAC_BAC10	anamox1_Bacteria_65_5_curated	2089721	64.54	434	2237	15423	51.28
LAC_BAC11	anamox1_Bacteria_72_15_curated	2937143	72.45	24	2507	371794	87.93
LAC_BAC12	anamox2_Bacteria_34_5_curated	434538	33.34	88	508	19334	50.78
LAC_BAC13	anamox2_Bacteria_61_6_curated	2415824	60.87	385	2646	23948	65.91
LAC_BAC14	anamox2_Bacteria_68_6_curated	2291336	67.96	460	2621	29805	64.03
LAC_BAC15	anamox3_Bacteria_50_5_curated	1042006	49.54	203	1076	20276	42.87
LAC_BAC16	anamox3_Bacteria_66_7_curated	2266342	65.35	286	2150	45201	81.9
LAC_BAC17	anamox3_Bacteria_67_13_curated	1642722	66.5	47	1442	89691	67.24
LAC_BAC18	anamox3_Bacteria_67_15_curated	2765727	66.48	144	2980	91127	95.69
LAC_BAC19	anamox4_Bacteria_63_7_curated	3451341	62.64	351	3276	43084	93.1
LAC_BAC20	anamox4_Bacteria_69_43_curated	3827900	69.37	15	3388	1008601	86.21
LAC_BAC21	LAC_NA06_Bacteria_60_18_curated	3689918	59.98	50	3221	327084	94.83

LAC_BAC22	LAC_NA07_Bacteria_38_171_curated	2383463	37.56	22	2157	344212	98.28
LAC_BAC23	LAC_NA07_Bacteria_70_305_curated	2792777	70.13	54	2465	247763	98.28
LAC_BAC24	LAC_NA07_Bacteria_71_12_curated	2770642	70.73	206	2877	57439	81.41
LAC_BACT01	anamox1_Bacteroidetes_38_5_curated	1414251	37.34	293	1501	13351	62.96
LAC_BACT02	anamox1_Bacteroidetes_39_16_curated	2896628	38.84	40	2472	542771	100
LAC_BACT03	anamox1_Bacteroidetes_63_11_curated	3356307	63.21	73	2915	220821	100
LAC_BACT04	anamox1_Sphingobacteriales_42_27_curated	3490519	42	95	3068	154972	100
LAC_BACT05	anamox1_Sphingobacteriales_43_8_curated	3156182	42.95	130	2826	130033	97.41
LAC_BACT06	anamox2_BJP_IG2103_Bacteroidetes_37_22_46_7_curated	2490972	45.52	190	2226	53299	93.1
LAC_BACT07	anamox2_Sphingobacteriales_41_11_curated	2768577	41.16	129	2490	79962	87.07
LAC_BACT08	anamox3_Bacteroidetes_39_15_curated	2533542	39.26	23	2188	636516	100
LAC_BACT09	anamox3_Burkholderiales_71_6_curated	1367674	69.86	648	1904	18351	32.85
LAC_BACT10	anamox3_Sphingobacteriales_44_6_curated	3033899	42.77	512	2768	29727	73.9
LAC_BACT11	anamox3_Sphingobacteriales_50_9_curated	4305233	49.73	216	2999	129109	99.14
LAC_BACT12	anamox4_Bacteroidetes_40_74_curated	2627547	39.91	15	2219	539103	100
LAC_BACT13	LAC_NA06_Bacteroidetes_30_9_curated	2233050	29.92	225	2055	41036	85.06
LAC_CHLX01	anamox1_Bacteria_56_37_curated	4970250	55.88	43	4404	437022	94.83
LAC_CHLX02	anamox1_Chloroflexi_52_59_curated	2864537	52.47	96	2759	102167	100
LAC_CHLX03	anamox2_Chloroflexi_60_8_curated	2023353	60.06	111	1885	73961	47.49
LAC_CHLX04	anamox3_Chloroflexi_59_6_curated	1407767	53.68	351	1507	12267	40.22
LAC_CHLX05	anamox3_Chloroflexi_68_6_curated	2294191	67.35	479	2330	19055	53.92
LAC_CHLX06	anamox4_Chloroflexi_66_15_curated	4383590	65.96	133	3602	139118	91.22
LAC_CHLX07	LAC_NA06_Anaerolineales_42_27_curated	2409642	41.78	210	2380	75774	93.1
LAC_CHLX08	LAC_NA06_Chloroflexi_57_14_curated	2719486	57.25	159	2567	157158	86.91
LAC_CHLX09	LAC_NA06_sub_Chloroflexi_59_14_curated	2594752	59.45	164	2359	101841	91.22
LAC_CHLX10	LAC_NA06_sub_Chloroflexi_61_22_curated	2755473	61.22	170	2707	68305	96.55
LAC_CHLX11	LAC_NA07_Caldilinea_aerophila_61_12_curated	3885866	60.59	83	3168	234454	80.88
LAC_CHLX12	LAC_NA07_Chloroflexi_57_23_curated	6358497	57.22	93	5201	299767	89.66
LAC_CHLX13	LAC_NA07_Chloroflexi_57_9_curated	2081669	55.57	325	2239	31406	71.76
LAC_CHLX14	LAC_NA07_Chloroflexi_58_12_curated	2711149	57.43	154	2459	125878	60.42
LAC_CHLX15	LAC_NA07_Chloroflexi_60_59_curated	3858643	60.1	13	3478	869846	94.83
LAC_CHLX16	LAC_NA07_Chloroflexi_65_58_curated	3316034	65.08	327	3017	48064	87.93
LAC_CHLX17	LAC_NA07_Chloroflexi_67_63_curated	3446202	66.61	65	2852	450062	96.55
LAC_CHLX18	LAC_NA07_RBG_16_RIF_CHLX_72_14_curated_75_20_curated	2506614	74.69	63	2330	167021	91.38
LAC_CLO01	anamox1_Candidatus_Cloacimonas_acidaminovorans_38_6_curated	1192038	35.28	246	1132	15612	56.47
LAC_D-T01	anamox1_Truepera_radiovictrix_71_5_curated	435656	69.12	317	694	3402	24.49
LAC_D-T02	LAC_NA07_Truepera_radiovictrix_72_29_curated	1353035	72.21	165	1373	45517	82.6
LAC_DADA01	LAC_NA06_RIFCSPHIGHO2_12_FULL_Dadabacteria_53_21_curated_58_6_curated	869333	55.01	539	1358	4902	33.68
LAC_GMT01	anamox1_Gemmatimonas_aurantiaca_57_6_curated	2631895	56.94	255	2549	56382	91.22

LAC_IGN01	anamox1_RBG_16_Ignavibacteria_36_9_curated_35_5_curated	1163158	34.03	347	1374	12559	70.14
LAC_IGN02	anamox2_Ignavibacteriales_33_72_curated	3331053	33.19	161	2995	201964	98.28
LAC_IGN03	anamox2_Ignavibacteriales_33_9_curated	2936140	33.09	166	2753	82489	86.05
LAC_IGN04	anamox2_Ignavibacteriales_41_12_curated	3156563	41.32	42	2805	297034	96.55
LAC_IGN05	anamox2_sub_Ignavibacterium_album_33_16_curated	3391206	33.43	195	3029	116818	97.81
LAC_IGN06	anamox3_BJP_IG2069_Ignavibacteriae_38_11_30_7_curated	1803433	30.38	132	1600	73197	73.51
LAC_IGN07	anamox3_sub_Ignavibacteriales_42_14_curated	3163923	42.21	22	2498	452672	96.55
LAC_MIC01	anamox2_Microgenomates_45_6_curated	740215	44.28	99	904	30865	73.35
LAC_MIC02	anamox2_Microgenomates_49_6_curated	950204	49.23	109	1082	36723	68.42
LAC_MIC03	anamox2_Roizmanbacteria_38_11_curated	663231	37.62	14	693	114830	58.31
LAC_MIC04	anamox4_Microgenomates_48_8_curated	999881	47.64	49	1055	99690	80.02
LAC_MIC05	LAC_NA06_Microgenomates_41_17_curated	1110978	41.17	7	1216	569682	80.88
LAC_MIC06	LAC_NA07_Roizmannbacteria_52_60_curated	957307	52.33	6	1013	885168	73.98
LAC_NIT01	anamox4_Candidatus_Nitrospira_defluvii_60_9_curated	3085099	60.3	94	3084	176639	95.69
LAC_OMN01	anamox3_Omnitrophica_63_14_curated	365510	62.81	4	402	274544	25.86
LAC_PLT01	anamox1_Planctomycetia_64_8_curated	4138393	63.58	783	3420	35628	87.62
LAC_PLT02	anamox4_sub_Candidatus_Brocadia_sinica_42_75_curated	3107335	42.29	64	2859	160056	98.28
LAC_PROT01	anamox1_Betaproteobacteria_69_7_curated	2106569	66.39	531	2636	12349	66.55
LAC_PROT02	anamox1_Burkholderiales_70_40_curated	3601855	70.09	60	3356	242838	100
LAC_PROT03	anamox1_Burkholderiales_71_17_curated	3639459	70.81	114	3448	149014	99.22
LAC_PROT04	anamox1_Gammaproteobacteria_64_6_curated	2447777	64.02	391	2594	28213	84.25
LAC_PROT05	anamox1_Lysobacter_67_10_curated	783854	66.76	125	840	26187	34.64
LAC_PROT06	anamox1_Nitrosomonas_europaea_50_14_curated	2128781	50.43	37	2006	177381	98.28
LAC_PROT07	anamox1_Proteobacteria_65_15_curated	2240508	65.45	348	2476	63177	58.59
LAC_PROT08	anamox1_Proteobacteria_67_8_curated	2912885	66.93	203	2910	122132	81.27
LAC_PROT09	anamox1_Rhodocyclales_69_13_curated	2263347	68.53	185	2364	100335	96.55
LAC_PROT10	anamox2_Burkholderiales_67_5_curated	1246117	66.28	298	1434	12340	38.05
LAC_PROT11	anamox2_Burkholderiales_68_9_curated	3527974	68.25	181	3452	116403	91.22
LAC_PROT12	anamox2_Burkholderiales_70_13_curated	3074974	70.33	39	2841	297272	84.48
LAC_PROT13	anamox2_Gammaproteobacteria_67_23_curated	2663638	67.06	116	2584	140445	79.31
LAC_PROT14	anamox2_Hydrogenophilales_66_19_curated	2257110	65.94	122	2331	92964	91.95
LAC_PROT15	anamox2_Myxococcales_71_5_curated	1434218	70.21	495	1687	8213	30.21
LAC_PROT16	anamox2_Rhizobiales_67_45_curated	4837226	66.56	24	4668	717639	100
LAC_PROT17	anamox2_Xanthomonadales_68_9_curated	1242229	67.63	21	1168	141855	50
LAC_PROT18	anamox2_Xanthomonadales_70_8_curated	2605543	68.54	346	2555	32162	82.76
LAC_PROT19	anamox3_Nitrosomonas_europaea_51_28_curated	2372894	50.54	93	2302	91833	98.12
LAC_PROT20	anamox3_Proteobacteria_68_11_curated	3218375	68.07	66	3078	349840	100
LAC_PROT21	anamox4_Alphaproteobacteria_58_14_curated	2593178	57.86	5	2558	1525488	98.28
LAC_PROT22	anamox4_Gammaproteobacteria_67_14_curated	3265689	66.91	73	3049	145604	98.28

LAC_PROT23	anamox4_Nitrosomonas_eutropha_48_11_curated	1880399	48.46	103	1862	88908	94.83
LAC_PROT24	LAC_NA06_Betaproteobacteria_71_7_curated	4060980	69.45	1841	5695	17332	69.75
LAC_PROT25	LAC_NA06_Burkholderiales_73_13_curated	3708868	72.67	335	3469	53457	94.51
LAC_PROT26	LAC_NA06_Rhizobiales_66_24_curated	3228463	66.32	148	3270	125494	98.28
LAC_PROT27	LAC_NA07_Burkholderiales_70_312_curated	2688511	69.75	290	2827	57610	93.32
LAC_PROT28	LAC_NA07_Proteobacteria_68_32_curated	2747459	68.4	14	2541	557607	98.28
LAC_PROT29	LAC_NA07_Rhodobacterales_68_7_curated	1308110	67.47	473	1651	13528	52.93
LAC_PROT30	LAC_NA07_Rhodocyclales_67_14_curated	2723825	66.65	212	2909	51017	92.95
LAC_SCH01	LAC_NA06_Candidatus_Saccharibacteria_40_6_curated	463567	39.93	148	580	16266	58.46
LAC_SPR01	anamox1_Turneriella_parva_44_7_curated	2667359	44.12	240	2748	87610	92.24
LAC_VER01	anamox2_Verrucomicrobia_58_8_curated	3592680	57.68	250	3293	90598	87.93
LAC_VER02	anamox2_Verrucomicrobia_62_8_curated	3685167	61.78	161	3134	100565	98.28
LAC_VER03	anamox3_Opitutus_terrae_67_4_curated	780321	65.72	503	1046	5713	24.42
LAC_VER04	anamox3_Pedosphaera_parvula_66_5_curated	1233438	65.98	543	1384	10642	56
LAC_VER05	anamox3_Verrucomicrobia_59_12_curated	2712092	59.27	50	2592	173980	96.55

Table 2 | Read counts to representative genomes across time points

Time piont (days)	# Total reads	# Total maped reads	% Mapped to rep	# Rep number
0	55398280	40291503	72.73	92
82	62544544	43877427	70.15	82
166	60931806	46030350	75.54	103
284	56282006	48644523	86.43	68
328	127048582	95132145	74.88	87
437	119945232	93737087	78.15	60

* number of representatives based on threshold of >1 coverage and > 0.5 breadth