Title: Increased Replication of Dissimilatory Nitrate-Reducing Bacteria Leads to Decreased

Anammox Bioreactor Performance

Authors: Ray Keren[1]*, Jennifer E. Lawrence[2]*, Weiqin Zhuang[3], David Jenkins[1], Jillian F.

Banfield[4], Lisa Alvarez-Cohen[1,5], Lijie Zhou[6]+, Ke Yu[7]+

*These authors contributed equally to the work

+Corresponding authors

Affiliations

1- Department of Civil and Environmental Engineering, University of California Berkeley,

Berkeley, CA, USA

2- Tighe & Bond, Westwood, MA, USA

3- Department of Civil and Environmental Engineering, The University of Auckland,

Auckland, New Zealand

4- Department of Earth and Planetary Sciences, University of California Berkeley,

Berkeley, CA, USA

5- Earth and Environmental Sciences Division, Lawrence Berkeley National Laboratory,

Berkeley, CA, USA

6- College of Chemistry and Environmental Engineering, Shenzhen University, Shenzhen,

China

7- Shenzhen Graduate School, Peking University, Shenzhen, China

Email of Authors:

Ray Keren: ray_keren@berkeley.edu; Jennifer E. Lawrence: jlawrence@tighebond.com; Weiqin

Zhuang: wq.zhuang@auckland.ac.nz ; David Jenkins: dijenkins@berkeley.edu ; Jillian F.

Banfield: jbanfield@berkeley.edu; Lisa Alvarez-Cohen: alvarez@ce.berkeley.edu : Lijie Zhou:

pakerzhou@szu.edu.cn ; Ke Yu: yuke.sz@pku.edu.cn

Abstract

Background Anaerobic ammonium oxidation (anammox) is a biological process employed to

remove reactive nitrogen from wastewater. While a substantial body of literature describes the

performance of anammox bioreactors under various operational conditions and perturbations,

few studies have resolved the metabolic roles of their core microbial community members.

Results Here, we used metagenomics to study the microbial community of a laboratory-scale

anammox bioreactor from inoculation, through a performance destabilization event, to robust

steady-state performance. Metabolic analyses revealed that nutrient acquisition from the

environment is selected for in the anammox community. Dissimilatory nitrate reduction to

ammonium (DNRA) was the primary nitrogen removal pathway that competed with anammox.

Increased replication of bacteria capable of DNRA led to the out-competition of annamox

bacteria, and the loss of the bioreactor's nitrogen removal capacity. These bacteria were highly

associated with the anammox bacterium and considered part of the core microbial community.

Conclusions Our findings highlight the importance of metabolic interdependencies related to

nitrogen- and carbon-cycling within anammox bioreactors and the potentially detrimental effects

of bacteria that are otherwise considered core microbial community members.

Background

Anaerobic ammonium oxidizing (anammox) bacteria obtain energy from the conversion of ammonium and nitrite to molecular nitrogen gas (N_2) [1]. Currently, the only bacteria known to catalyze this process are members of the phylum *Planctomycetes* [2, 3], none of which have been isolated [3, 4]. In practice, anammox bacteria are employed in an eponymous process in combination with the partial nitritation (PN) process to remove ammonium from nitrogen-rich wastewaters. First, in PN, approximately half of the ammonium in solution is aerobically oxidized to nitrite. Second, in anammox, both ammonium and nitrite are anaerobically converted to N_2 [5, 6]. The PN/anammox (i.e., deammonification) process is beneficial because it consumes 60% less energy, produces 90% less biomass, and emits a significantly smaller volume of greenhouse gas than conventional nitrogen removal by nitrification and denitrification processes [7]. To date, over 100 full-scale deammonification process bioreactors have been installed at municipal and industrial wastewater treatment plants across the globe [8].

Within an engineered environment, anammox bacteria have very low growth rates and can easily become inhibited by fluctuating substrate and metabolite concentrations [9, 10]. When these two limitations are coupled, recovery from an inhibition event can take up to six months (which is unacceptably long for municipalities that must meet strict nitrogen discharge limits) [11]. Furthermore, these problems are compounded by a cursory understanding of the microbial communities that exist alongside anammox bacteria. A deeper understanding of the complex interactions occurring among bacterial species in an anammox bioreactor is required for the broad application of the deammonification process for wastewater treatment.

Previous research has suggested that a core microbial community exists within anammox bioreactors [12–16]. In the majority of studied bioreactors, uncultured members of the phyla

Bacteroidetes, *Chloroflexi*, *Ignavibacteria*, and *Proteobacteria* have been identified alongside

Planctomycetes, the phylum that contains anammox bacteria. These phyla have primarily been

identified through 16S rRNA gene studies, so their interplay with anammox performance has not

yet been fully elucidated [12–16]. From their taxonomic identity and performance studies, it is

assumed that the additional phyla compete for nitrite and cooperate to transform (i.e.

dissimilatory nitrate reduction to ammonium - DNRA) and remove (i.e. denitrifiers) nitrate, a

product of anammox metabolism [17–19].

Here, we illuminate deeper metabolic relationships between an anammox bacterium,

Brocadia, and its supporting community members during the start-up and operation of a

laboratory-scale anammox bioreactor. We begin by analyzing the formation of the "anammox

community" through a combination of genome-centric metagenomics and 16S rRNA gene

sequencing. Metabolic features of positively enriched bacteria are compared to negatively

enriched bacteria during the start-up process. Next, we focus our investigation on an anammox

performance destabilization event that was driven by microbial interactions. Last, we conduct a

comparative analysis of our anammox community to similarly-studied anammox communities

[18, 20] to highlight the broader relevance of our results. To our knowledge, this is the first time-

series-based study to link anammox metagenomic insights and community composition to

anammox bioreactor functionality [21]. Our findings bolster the fundamental, community-level

understanding of the anammox process. Ultimately, these results will enable more

comprehensive control of this promising technology and facilitate its widespread adoption at

wastewater treatment plants.

Results

Bioreactor performance. The performance of a laboratory-scale anammox anaerobic membrane bioreactor (MBR) (described in methods) was tracked for 440 days from initial inoculation, through several performance crashes, to stable and robust anammox activity (Figure 1). Performance was quantified by nitrogen removal rate (NRR, g-N L^{-1} d^{-1}) and effluent quality (g-N L^{-1} d^{-1}). Bioreactor performance generally improved over the first 103 days of operation. At this point, the hydraulic residence time (HRT) was reduced from 48 to 12 hours and influent nitrogen concentrations were reduced to maintain a stable loading rate. Additional seed biomass from a nearby pilot-scale deammonification process was added on Day 145 following a performance crash and bioreactor performance improved, enabling influent ammonium and nitrite concentrations to be steadily increased until the NRR approached 2 g-N L^{-1} d^{-1}. On Day 189 the bioreactor experienced a technical malfunction and subsequent performance crash, identified by a rapid decrease in the NRR and the effluent quality. On Day 203, the bioreactor was again amended with a concentrated stock of seed biomass and the NRR and the effluent quality quickly recovered. Influent ammonium and nitrite concentrations were again increased until the NRR reached 2 g-N L^{-1} d^{-1}.

The bioreactor subsequently maintained steady performance for approximately 75 days, until Day 288, when effluent concentrations of ammonium and nitrite unexpectedly began to increase and nitrate concentrations disproportionately decreased. Seven days later, the NRR rapidly plummeted. No technical malfunctions had occurred, indicating that the destabilization of the anammox process may have been caused by interactions among its microbial community members. At the time, the cause of the performance decline was not understood, so the bioreactor was not re-seeded with biomass. After 50 days of limited performance, concentrations

of copper, iron, molybdenum, and zinc in the bioreactor influent were increased based on

literature recommendations [22–25] and the NRR rapidly recovered. Stable and robust bioreactor

performance was subsequently maintained.

Metagenomic sequencing and binning. Whole community DNA was extracted and sequenced

at six time points throughout the study: Day 0 (D0), for inoculant composition; Day 82 (D82),

during nascent, positive anammox activity; Day 166 (D166), three weeks after an additional

biomass amendment; Day 284 (D284), after a long period of stable and robust anammox activity

and just before the bioreactor performance was destabilized; Day 328 (D328), in the midst of the

performance destabilization period; and Day 437 (D437), during mature, stable, and robust

anammox activity.

From all samples, 337 genomes were binned, 244 of which were estimated to be >70%

complete by checkM [26]. The genomes were further dereplicated across the six time points into

clusters at 95% average nucleotide identity (ANI). This resulted in 127 representative and unique

genomes (Table S1) that were used for all downstream analyses. Mapping showed an average

read recruitment of 76% to the representative genomes (Table 1). The number of genomes

present at each time point (using threshold values of coverage > 1 and breadth > 0.5) ranged

from 60 (D437) to 103 (D166). In addition, nine strains were detected that differed from the

representative genome by 2% ANI (Additional file 1, Table S2). Except for the anammox

bacterium, which is referenced at the genus level (*Brocadia*), all representative genomes are

referenced at the phylum level.

Community structure and temporal dynamics. As both internal and external factors can work

in combination to affect the structure of a bioreactor community, we hypothesized that different

groups of bacteria (i.e., sub-communities) would be associated with different phases of the

bioreactor's lifespan. To test for grouping, all genomes were pairwise-correlated (Figure 2A).

The resulting heatmap revealed four distinct clusters (Groups A-D). Group A was the largest,

with 52 genomes, while Groups B-D had 25, 24, and 26 genomes respectively (Table S3).

To better examine the clustering of genomes in relation to the bioreactor's lifespan, we

ran nonmetric multidimensional scaling (nMDS) analyses on the genomes' relative abundance

data (Figure 2B). The nMDS projection revealed that genome groups were strongly associated

with specific time points: Group A was associated with the inoculant biomass at D0 and D166,

while Group C was associated with the nascent anammox community at D82. Group B was

associated with the times of destabilized anammox performance (Days 284-328), and Group D

was associated with the mature, stable anammox community at D437. *Brocadia* is a part of

Group D, although its location on the nMDS projection is skewed to the left because of its high

relative abundance throughout the majority of the bioreactor's lifespan. Because the nascent

anammox community was amended with additional biomass, we could not resolve a linear

trajectory for the microbial community between the initial and final states. Nevertheless, Groups

B and D shared many similarities, and the majority of the genomes associated with Group B

were still present in the bioreactor on D437.

To further resolve the relative abundances of Groups A-D over the bioreactor's lifespan,

16S rRNA genes from genomes were combined with direct 16S rRNA sequencing data, then

clustered into operational taxonomic units (OTUs). Of the 127 representative genomes, 34

contained a 16S rRNA sequence that was successfully clustered to an OTU. The matching OTUs

represented 55% of the total 16S rRNA reads at day 9, but quickly increased to an average

representation of 86%. The OTUs that match assembled genomes were grouped (Figure 2) and

their relative abundance summed (Figure 3A). The matching genomes comprised 18/52, 10/25,

3/24, and 7/26 of Groups A-D respectively. The matching OTUs represented 55% of the total

16S rRNA reads at day 9, but quickly increased to an average representation of 86%.

Group A was dominant on Day 0, but rapidly decreased in abundance by the first 16S

rRNA gene sequencing time point on Day 9 (Figure 3B). Group A was again dominant after the

addition of new inoculum on Day 145 (Figure 3A). Group B (and to a certain extent Group A)

became dominant just before Day 300 when the anammox performance was destabilized.

To check the accuracy of the 16S rRNA matching to the metagenomic data, we compared

the relative abundances of Groups A-D across three data sub-sets (Figure 3B): all metagenomes

(marked 'met' on the x-axis), only metagenomes with matching 16S rRNA OTUs (marked 'sub-

met' on the x-axis), and the 16S rRNA OTUs. Overall, the three datasets were compatible, with

slight variations in over/under-estimations of particular Groups. In comparison to the

metagenomically-derived data, the 16S rRNA data tended to overestimate the relative abundance

of Group A and underestimate the relative abundance of Group D. A large fraction of the

Chloroflexi in Group D were not matched to 16S rRNA OTUs, so the underestimation was

consistent with expectations.

For all subsequent analyses, we split the representative genomes into two groups (Table

S3): those that are associated with the mature anammox community at D437 (Anammox

Associated, AA), and those that are not (Source Associated, SA). The AA community includes

all of the genomes that are present at D437, while the SA community includes the rest of the

genomes that are not present at D437. Some of these genomes are associated with the sludge

amendments, and some are associated with the nascent anammox community; at no point is there

a community exclusively comprised of SA genomes.

Metabolic profiles. For the purpose of analyzing the metabolic potential of the microbial

community, we evaluated only genomes with > 70% completeness (n = 88) [26]. Using Hidden

Markov Model (HMM) searches of the Kyoto Encyclopedia of Genes and Genomes (KEGG)

database, we checked for the presence of genes based on their KEGG Orthology (KO) number,

and calculated KEGG module completeness [28, 29]. The genomes were clustered by KO

presence/absence (Figure S1) and their module completeness (Figure 4). Clustering by the two

methods resulted in similar groupings.

Module clustering resolved five groups ($\square$, β, $\square$, δ, ε) (Figure 4A and Table S3). Groups

$\square$ and β contained more anammox-associated genomes (90% and 60%, respectively), while

groups $\square$, δ, and ε contained 65%, 70% and 60% of source-associated genomes, respectively.

The clustering was also strongly influenced by bacterial taxonomy (Figure 4B). Group $\square$ was

composed solely of Gram (+) bacteria, primarily *Chloroflexi*. Group β was composed of

Candidate Phyla Radiation (CPR) bacteria, *Microgenomates*. This group of bacteria has reduced

genomes and metabolism [30], and thus has an unknown effect on the community metabolism.

Group $\square$ was composed entirely of Gram (-) bacteria from a wide range of phyla, and includes

the anammox bacterium *Brocadia*. Group δ was composed of *Ignavibacteria* and *Bacteroidetes*,

but only the *Ignavibacteria* from Group δ were associated with the AA group. Accordingly,

further analysis of Group δ refers only to the *Ignavibacteria*. Group ε was composed entirely of

Proteobacteria.

Based on the KEGG module clustering, we reconstructed the representative metabolisms

of the five groups (Figure 5). We used a module completeness threshold of 67% per genome and

considered it representative if it was complete in >50% of the group's members. Group δ was not

represented since it diverged from group ☐ by auxotrophies in several modules (Figure 4A, red

rectangle). The *Brocadia* metabolism is shown in Figure S2.

While module completeness was used for most of the analyses, in several cases it was

not sufficient (e.g., overlap between modules, no module for path). For oxidative

phosphorylation, fermentation, carbon fixation, several amino acid synthesis pathways, and

nitrogen metabolism, we analyzed gene presence manually. For anammox, four additional

HMMs were added: hydrazine synthase subunit A (*hzsA*), hydrazine oxidoreductase subunit A

(*hzoA*), and nitrite oxidoreductase subunits *nrxA* and *nrxB* [31]. For the latter, the similarity of

the gene to the nitrate reductase *narGH* was taken into consideration.

With the exception of two CPR bacteria, all of the genomes in the bioreactor contained

genes encoding assimilation of ammonia into glutamate (Figure 6). More than half (49) of the

bacteria could potentially reduce nitrate, and the same number had the genes needed to further

reduce nitrite to nitrogen monoxide (NO); however, only 26 bacteria had the genes to do both

steps. The remaining steps of denitrification were encoded in an even smaller number of

genomes. The *nrxAB* gene was only identified in two genomes, one of which was *Brocadia.*

One-step DNRA was identified in 22 genomes. While the number of genes encoding ammonia

assimilation and nitrate reduction to nitrite were fairly similar in the AA and SA groups'

genomes, DNRA was more common in the AA genomes and denitrification beyond nitrite in the

SA genomes.

Carbon fixation is a necessary step in the anammox bioreactor since the influent media

did not contain organic carbon. Only two bacteria in the community could be considered

autotrophic primary producers. *Brocadia* was confirmed as a primary producer, fixing carbon via

the Wood-Ljungdahl pathway and obtaining energy from the anammox pathway. The second

bacterium, LAC_PROT27 (*Proteobacteria*, group AA), might be able to fix carbon by the Calvin

cycle and obtain energy from denitrification, and could possibly oxidize sulfide to sulfite (*dsrAB*

is present in the genome). While LAC_PROT27 was consistently highly abundant in the

bioreactor, it was always at least 3-folds less abundant than *Brocadia* (except at time 0). Several

other bacteria were also potential autotrophs (or mixotrophs), but were low in relative abundance

over the bioreactor's lifespan. Additional information about carbon metabolism and electron

transfer can be found in supplementary information file 1.

Analysis of metabolic selection in the anammox bioreactor. During the maturation of the

anammox bioreactor, the number (Table 1) and diversity (Table S4) of genomes were both

reduced. To examine why certain bacteria were enriched (group AA) while others were removed

(group SA), we compared the genomes' ability to synthesize metabolites against their ability to

acquire nutrients from the environment. For synthesis we checked 24 KEGG modules for amino

acids (aa.), 18 modules for vitamins and cofactors, and 28 modules for lipids and fatty acids. For

nutrient acquisition we checked 54 KEGG modules for transporters. The mean module

completeness across these categories was compared. A complete module implies a bacterium has

the functional capability (be it synthesis or transport). Thus, the higher the module completeness

of a group, the more likely it is that its members have the relevant functional capability. For

statistical analysis, when both data sets (AA and SA) fit a normal distribution, a two sample T-

test was conducted. When the values did not fit a normal distribution, the ratio between groups

AA and SA was calculated, and used to set a confidence interval (CI: mean $\pm 1.64*(SD/n^{0.5})$,

alpha = 0.05). Values outside of the CI were considered significantly different than the mean.

Synthesis modules for aa. (p-value = 0.68) and vitamins/cofactors (p-value = 0.51) both

fit a normal distribution, and no statistically significant differences were found for these

categories between the AA and SA bacterial groups. Synthesis modules for lipids and fatty acids

did not have a normal distribution, so their ratio was inspected (CI upper = 1.22. CI lower =

0.80). Six modules were significantly higher and 14 were significantly lower in the AA group vs.

the SA group. Group AA had a higher proportion of Gram (+) bacteria, which also added to the

difference in module completeness. Transport modules also did not fit a normal distribution.

Inspection of the module ratio showed that both the upper CI (2.69) and lower CI (1.74) were

higher than a ratio of 1. Of the 26 transport modules with a ratio >1.74, 18 were transport

systems for organic carbon molecules (sugars, lipids, a.a., and cofactors).

These comparisons suggest that a bacterium's ability to acquire nutrients from its

environment may be a selective driver in the anammox bioreactor's community. This was

particularly highlighted when observing Metabolic group α, the dominant metabolic group

within the AA bacteria. Members of group α have a cassette of extracellular proteases and

decarboxylases paired with a wide array of transporters (Figure 5B) that enable acquisition of

nutrients from the environment. In addition, the larger ratio of bacteria with auxotrophies in the

AA bacteria (Figure 4A, red rectangles for groups α and δ) hints of a greater reliance on external

metabolites from other members of the community.

Metabolic interdependencies between community members. The bacteria in the AA

community have a complex metabolic system, with many bacteria relying on other members to

provide them with necessary metabolites. In the mature functioning bioreactor, *Brocadia* was the

only primary producer present. It was also the only bacterium capable of synthesizing vitamin

B12. For most other metabolites (e.g., vitamins and cofactors) the possible metabolic

interdependencies [33] are less straightforward (Figure 7 and Table S5). In the schematic figure,

the size of each group reflects its relative abundance in the bioreactor at D437. The arrows point

towards the group that potentially receives metabolites it cannot synthesize, and the arrows' sizes

reflect the proportion of the examined metabolites that the group needs. Members of group α (the

most dominant group aside from *Brocadia*) had multiple auxotrophies in the synthesis of

vitamins, cofactors, fatty acids, and lipids. Group α could produce a few metabolites needed by

other groups. Members of group ε were the most metabolically diverse and could produce many

metabolites needed by other groups. This group could account for most of the auxotrophies of

Brocadia. Members of group ε could support 75% of *Brocadia*'s auxotrophies in aa, vitamins

and cofactors, as well as 60% of its auxotrophies in fatty acids and lipid synthesis (Table S5).

Group γ was the smallest group in the bioreactor at D437 and had a mix of auxotrophies and

metabolic support potential.

When combining all of the above data, we found that groups ⬚ and ε both had

mutualistic associations with *Brocadia* (Figure 7). Group ε potentially provided more

metabolites to *Brocadia* than it received, whereas groups ⬚ and δ seemed to gain more from

Brocadia than they provided.

Investigating a microbial driven anammox performance destabilization event Just before

day 300 of the bioreactor's lifespan, an unexpected anammox performance destabilization event

occurred. We hypothesized that some interaction between the anammox bacterium and the co-

existing community members led to this event. To evaluate which bacteria might have influenced

the destabilization event and which might have been influenced, we relied on two parameters:

replication and relative abundance (using coverage). First, we checked for changes in a genome's

replication rate (as calculated by iRep [34], see methods for detailed explanation) between D166

and D284. Second, we investigated the log-ratio (LR) of coverage [35] between D284 and D328.

To eliminate bias in microbial loading or sequencing depth, we used four different reference

frame genomes (RFg) for the LR calculation [35]. Difference in sequencing depth or microbial

load can highly bias comparisons between samples. Higher sequencing depth means higher

coverage across the entire sample so it might seem that the abundance of bacteria has increased.

To remove that bias a reference frame genome(s), with little change in relative abundance over

time, is chosen. The abundance values of all other genomes within a given time point are divided

by the abundance of the RFg before calculating the log-ratio between samples. The internal ratio

removes the aforementioned bias between samples. By combining these two parameters, we

were able see which bacteria were actively replicating before the performance destabilization

event and eliminate biases created by relative abundance results.

Genomes with iRep values for three out of the four time points between D166 and D437

(27, accounting for > 80% of community) were chosen for this analysis. For each RFg in the LR

calculations, significant change was considered for values outside of the confidence interval (CI),

calculated for all 127 genomes (Table S6). A bacterium was considered to influence the

destabilization if it both increased in replication prior to the event and had a positive and

significantly high LR relative to each RFg. A bacterium was considered to be influenced by the

destabilization if it decreased its replication rate prior to the event and its LR was significantly

low.

Two *Chloroflexi* (LAC_CHLX01 and LAC_CHLX10) showed consistent, significant

growth across all RFgs (Figure 8A and Table 2), as well as increased replication rates prior to the

destabilization event (Figure 8B). These bacteria likely influenced the destabilization event.

Three additional bacteria (the *Chloroflexi* LAC_CHLX06, LAC_CHLX09, and *Ignavibacteria*

LAC_IGN05) were also potential influencers, having significant growth based on some of the

RFgs. On the other hand, *Brocadia* (LAC_PLT02), showed significantly decreased replication

rates prior to the destabilization event. *Brocadia*'s replication rate on D284 (1.07) indicates that

only 7% of the bacterium's population was actively replicating. At other time points, *Brocadia's*

replication rate climbed as high as 2.13, indicating that 100% of the bacterium's population was

actively replicating. One additional bacterium (LAC_PROT22, a *Proteobacteria* from group

AA) showed decreased replication and growth (under two RFgs).

Next, we next investigated the nitrogen metabolism of the 27 bacteria, specifically the

three metabolic paths that compete for nitrite i.e., anammox, denitrification, and DNRA (Figure

8C). Denitrification was subdivided into three steps (NO_2 reduction, NO reduction, and N_2O

reduction); DNRA is a single step process. Anammox could only be performed by *Brocadia*.

Only a single bacterium (LAC_BAC20) could perform full denitrification. All bacteria capable

of DNRA were also capable of partial denitrification. When allowing for a bacterium to have

multiple pathways (Figure 8C), NO_2 reduction via DNRA (*nrfAH*) and denitrification (*nirS/nirK*)

appeared to be equally dominant during the anammox performance destabilization event (D328).

While N_2O reduction was also dominant, a bottleneck appeared to exist at the second step, NO

reduction. When we assume that the bacteria capable of both DNRA and partial denitrification

choose the path that yields the most energy [36], we can remove the partial denitrification

pathway (Figure 8C). Under this assumption, DNRA was clearly revealed as the dominant

process occurring in the bioreactor during the anammox performance destabilization event. All

bacteria that were shown to affect the destabilization event were DNRA bacteria. In addition, the

destabilizing bacteria belong to metabolic groups α (LAC_CHLX01, LAC_CHLX06,

LAC_CHLX09, and LAC_CHLX10) and δ (LAC_IGN05). Both of these groups were reliant on

other members for organic carbon (Figure 5B and Figure 7).

Core anammox community. Resulting genomes from our study, in combination with genomes

from two previous anammox metagenomic studies, Speth et al. [18] (22 genomes) and Lawson et

al. [20] (15 genomes), provide strong evidence to support a core anammox community (Figure

9). The relative abundances of bacteria from the dominant phyla across these three bioreactors

were fairly similar: in each bioreactor the anammox, along with *Chloroflexi*, *Ignavibacteria*, and

Proteobacteria bacteria, composed >70% of the community (Figure 9B).

Due to the significantly larger genome yield and time-series analysis in this study, our

bioreactor shared more genomes with each of the other bioreactors than the other bioreactors

shared between themselves. In total, 21 genomes from our bioreactor were closely related to

those from at least one of the two other bioreactors, 17 of which were present at the last time

point, D437 (Table S7). The related bacteria accounted for 50% and 93% of the Speth et al. and

Lawson et al. genomes, respectively. The bioreactor studied by Speth et al. was different from

the other two bioreactors because it was amended with oxygen to perform partial nitrition and

anammox within the same bioreactor, while the others performed anammox only.

A more focused phylogenetic tree of *Planctomycetes* shows that the *Brocadia* in our

bioreactor and in the Lawson et al. bioreactor are the same species (*Brocadia sapporensis [27]*),

while the *Brocadia* species from the Speth et al. bioreactor is different (*Brocadia sinica*) (Figure

S3).

Discussion

In this study we present an in-depth analysis of the development of an anammox

community from seed to stable state (through several perturbations) in an anaerobic membrane

bioreactor. By combining several methodologies, we are able to gain important insights into the

dynamics and interactions of more than 100 species in the bioreactor community [37, 38].

The first perturbation of the bioreactor, a mechanical malfunction combined with

inoculum amendments, changed the trajectory of the community succession. This can be seen

from the relative abundance-based grouping (Figures 2-3) and the detected strain shifts. The first

inoculum amendment had a much stronger effect on community assembly than the subsequent

bioreactor malfunction and second amendment [39, 40]. The large shift in the community

occurred between days 96 and 152, after which the community trajectory became fairly

consistent until day 290. The oscillations after days 350 are likely due to differences in

sequencing depths. Large changes in community structure due to inoculations are a real concern

for large-scale bioreactors, where the influent contains constantly-changing communities of

bacteria [41, 42]. It is unclear if members of group C, which were dominant in the nascent

anammox community, would have better supported the bioreactor's anammox performance [39].

Looking at their metabolism may offer a few clues: of the 14 genomes clustered into the nascent

anammox community, six were of group ε and four were of group γ, both of which were

considered to maintain a mutualistic or commensal relationship with the anammox bacterium. In

addition, denitrification was more common than DNRA in group C. Two genomes

(LAC_ACD03 and LAC_PROT30) from the nascent anammox community were included in the

focused investigation of the anammox performance destabilization. Both had no significant

effect on the event nor were they significantly affected by it.

The metabolic analysis of the matured community showed that transport systems for

nutrients (mainly organic carbon) were the most enriched in the community. The ability of

bacteria to utilize available nutrients in the environment [43] has been shown before and has

been proposed to explain dominance in nutrient-rich environments [44]. However, when such

bacteria favor acquisition over synthesis, it can further stress a community that is reliant on a

slow growing primary producer for nutrients [45–48]. Members of groups α and δ have these

characteristics, and some are implicated in the anammox performance destabilization event.

The aforementioned destabilization event took place after nearly 100 days of high performance,

and no external factors could explain the sudden destabilization and performance crash. By

combining information about replication rates and changes in relative abundance of community

members, we were able to identify several bacteria that likely affected the performance crash.

Analysis of composite data (such as relative abundance) for true changes in community structure

has many pitfalls [35]. In addition, the response of a bacterium to an event, in this case an

increase in relative abundance following the performance destabilization, cannot be deduced to

mean it had any effect on the event. We were fortunate to have taken a metagenomic sample the

week prior to the destabilization event. By measuring replication rate changes as well as relative

abundance changes, we could better deduce a possible causative effect. A bacterium that

increases its replication rates prior to the event and increases in relative abundance due to the

event is more likely to have a causative effect. While most bacteria had higher replication rates at

D284 compared to D166 (17 of 22 bacteria with values in both days), only five bacteria

significantly increased in relative abundance following the event. Genes conferring DNRA and

partial denitrification capabilities were detected in these bacteria. These types of bacteria could

improve bioreactor performance if they remove nitrate and excess nitrite, but they could be

detrimental if they compete with anammox for nitrite or allow for the build-up of nitrite. Here,

the equilibrium between support of the anammox process and disruption of the anammox process

was tilted towards the latter.

Two possible scenarios for nitrogen metabolism are consistent with the bioreactor's

performance that exhibited decreased nitrogen removal and increased ammonium in the effluent

leading up to the destabilization event. One scenario is described by DNRA dominance and the

second by nitrite reduction to nitric oxide and its subsequent leakage from the system. The first

scenario provides more energy to the bacteria [36], so we speculate it is more likely. *Brocadia*

has genes required for DNRA, but given the high nitrogen removal rate by the anammox process

leading up the disturbance, it can be assumed that *Brocadia* would not be primed to carry out the

DNRA reaction on such a short timescale. However, DNRA could potentially be used by

Brocadia for detoxification by cycling potentially toxic excess nitrite back to ammonium, where

it could then participate in the anammox reactions [18, 20].

The replication rate of *Brocadia* prior to the performance destabilization was down to

1.07, from 2.13 on D166. A replication rate of 1.07 is equivalent to only 7% of the population

actively replicating, which can explain the large decrease in relative abundance in the following

time point. This also hints that the process leading to the destabilization event occurred prior to

D284. Whether the process could be assigned to a single specific occurrence, or if it is an

additive process culminating in a break from equilibrium among members of the community, is a

gap that future research should address.

A broader investigation of metabolic interdependencies within the community shed light

on the stability of the anammox community. *Brocadia* is the source of organic material in the

community, but obtains essential metabolites from community members, especially

Proteobacteria. This forms a basis for a mutual symbiotic relationship. On the other hand,

Chloroflexi, comprising the largest group of bacteria besides *Brocadia*, receive numerous

metabolites while apparently providing few in return. They are characterized by an array of

extracellular proteases and amylases, likely used to breakdown the extracellular matrices formed

by *Brocadia*. *Chloroflexi*, as a group, are most associated with anammox bacteria and form a

large fraction of the core community. They also account for the majority of the destabilizing

bacteria. Together, the results point to a parasitic symbiosis. While anammox bacteria generate

sufficient organic carbon to support the growth of its co-occurring heterotrophic microorganisms,

the tipping point between stable and unstable operation and the factors that control it have not

been fully identified. Input changes may be able to restore anammox activity, but this is only an

empirical solution. In full-scale anammox bioreactors where influent organic carbon is

essentially ubiquitous, heterotrophic dominance could persist without some sort of active

countermeasure. Therefore, future research should target the inhibition of potential destabilizing

heterotrophs.

Previous studies have discussed a potential core anammox community [12–16]. With the

exception of very few studies, all such work has been conducted with single gene markers. Our

analysis of an anammox community is the largest to-date and thus expands the ability to test this

hypothesis. Our results support the existence of a core community, while identifying factors that

differentiate communities. The high similarity among bacterial communities originating from

three distinct anammox bioreactors [18, 20] strongly suggests a global core anammox microbial

community. In the construction of the phylogenetic tree, we used >3000 reference genomes

originating from diverse environments. Through this analysis, we found that the anammox

community forms distinct clades at the species level, despite the sheer number and diversity of

sources. More than half of the bacteria did not have species level relatives, and an additional

26% only had a relative found in our anammox bioreactor or in a previous anammox study [18,

20]. Together, nearly 80% of the bacteria are unique to anammox bioreactors, so it is clear that

the anammox bioreactor selects for a unique set of bacteria. Parameters that increased the

differences between communities are the species of the anammox bacterium and the bioreactor

configuration. Since both parameters relate to the same bioreactor [18], we cannot conclude

which has a stronger effect.

Conclusions

Here we present the largest-to-date metagenomic analysis of a microbial community in an

anammox bioreactor. Our results support the growing body of literature that suggests that

anammox communities are unique and may share a core microbial community. We identified a

distinct phylogenetic profile across reported metagenomic analyses of anammox bioreactors. In

subsequent analyses of our metagenomes, we identified metabolic traits associated with the core

microbial community that are distinguishable from other bacteria present in the source sludge

inoculum. In addition, our time-series analysis included a biologically-driven period of anammox

performance destabilization. We identified an increase in replication rates for several bacteria

just prior to the event. Further analysis revealed that these bacteria contain genes conferring

DNRA, which puts them in direct competition with the *Brocadia* sp. for nitrogen resources.

Combined, our results provide a possible mechanistic explanation for the performance shift of

the anammox bioreactor and advance the comprehensive control of this promising technology.

However, further work is needed to elucidate the precise mechanisms that govern anammox

community interactions and to predict performance destabilization events.

Methods

Bioreactor operation. A laboratory-scale, anammox anaerobic membrane bioreactor (MBR)

with a working volume of 1L was constructed and operated for over 440 days (Figure S4). The

bioreactor was originally inoculated with approximately 2 g volatile suspended solids (VSS) L^{-1}

of biomass from a pilot-scale deammonification process treating sidestream effluent at San

Francisco Public Utilities Commission (SFPUC) in San Francisco, CA. The bioreactor was re-

inoculated with similar concentrations of biomass from the same source on Days 147 and 203.

Synthetic media containing ammonium, nitrite, bicarbonate, and trace nutrients (meant to mimic

sidestream effluent at a municipal wastewater treatment plant) was fed to the bioreactor (Table

S8). For the first 154 days of operation, the bioreactor was kept under nitrite-limiting conditions

to prevent inhibitory conditions due to the buildup of nitrite, and influent ammonium and nitrite

concentrations ranged from 200-300 mg N L^{-1} and 100-300 mg N L^{-1}, respectively. On Day 154,

ammonium and nitrite concentrations were adjusted to the theoretical anammox stoichiometric

ratio, 1:1.32. Afterwards, influent ammonium and nitrite concentrations were maintained at this

ratio. Ammonium ranged from 200-500 mg N L^{-1} and nitrite 265-660 mg N L^{-1}. On Day 353,

influent concentrations of copper, iron, molybdenum, and zinc were increased based on literature

suggestions [22–25].

The bioreactor was operated in a continuous flow mode. For the first 145 days, the

hydraulic retention time (HRT) was maintained at 48 hours; afterwards it was reduced to 12

hours. No solids were removed from the bioreactor for the first 100 days of operation;

afterwards, the solids retention time (SRT) was reduced to 50 days. A polyvinylidene fluoride

hollow fiber membrane module with a 0.4 µm pore size and total surface area of 260 cm^2 (Litree

Company, China) was mounted in the bioreactor. Temperature was maintained at 37° C with an

electric heating blanket (Eppendorf, Hauppauge, NY). Mixing was provided by an impeller at a

rate of 200 rpm. Mixed gas was supplied continuously to the bioreactor (Ar:CO$_2$ = 95:5; 50 mL

min^{-1}) to eliminate dissolved oxygen and maintain a circumneutral pH range of 6.9-7.2. Influent

and effluent concentrations of ammonium, nitrite, and nitrate were measured approximately

every other day using HACH test kits (HACH, Loveland, CO), as described in the

manufacturer's methods 10031, 10019, and 10020, respectively.

Biomass collection and DNA extraction. Biomass samples were extracted via syringe from the

bioreactor every 2-10 days, flash frozen in liquid nitrogen, and stored frozen at -80 ℃ until use.

Genomic DNA was extracted from the samples using the DNeasy PowerSoil Kit (Qiagen,

Carlsbad, CA), as described in the manufacturer's protocol. Extracted DNA was quantified with

a NanoDrop Spectrophotometer (Thermo Scientific, Waltham, MA), and normalized to

approximately 10 ng/µL with nuclease-free water (Thermo Scientific, Waltham, MA). All

genomic DNA samples were stored at -20 ℃ until use. For shotgun metagenomic sequencing,

samples were sent to the Joint Genome Institute (JGI) in Walnut Creek, CA. There, DNA quality

was assessed prior to library preparation and sequencing (150bp pair-end) on the Illumina HiSeq

2500 1T sequencer (Illumina, San Diego, CA). For 16S rRNA sequencing, samples were sent to

the Institute for Environmental Genomics at the University of Oklahoma. There, DNA quality

was assessed prior to library preparation and amplicon sequencing on the Illumina MiSeq

sequencer (Illumina, San Diego, CA).

Metagenomic sequencing, assembly, and binning. Resulting sequences from each time point

were processed separately, following the ggKbase SOP (https://ggkbase-

help.berkeley.edu/overview/data-preparation-metagenome/). Briefly, Illumina adapters and trace

contaminants were removed (BBTools, GJI) and raw sequences were quality-trimmed with

Sickle [49]. Paired-end reads were assembled using IDBA_UD with the pre-correction option

and default settings [50]. For coverage calculations, reads were mapped with bowtie2 [51].

Genes were predicted by Prodigal [52] and predicted protein sequences were annotated using

usearch [53] against KEGG, UniRef100, and UniProt databases. The 16S rRNA gene and tRNA

prediction was done with an in-house script and tRNAscanSE [54] respectively. At this point, the

processed data was uploaded to ggKbase for binning.

Manual binning was performed using the ggKbase tool. The binning parameters were

GC% and coverage (CV) distribution, and phylogeny of the scaffolds. Quality of the manual bins

was assessed by the number of Bacterial Single Copy Genes (BSCG) and Ribosomal Proteins

(RP) found in each bin (aiming at finding the full set of genes, while minimizing multiple

copies). In addition to manual binning, automated binning was done using four binners:

ABAWACA1 [30], ABAWACA2, CONCOCT [55], and Maxbin2 [56]. For all, the default

parameters were chosen.

All bins from both automatic and manual binning tools were input into DASTool [57] to

iterate through bins from all binning tools and choose the optimal set of bins. checkM was run to

analyze genome completeness [26]. The scaffold-to-bin file created by DASTool was uploaded

back to ggKbase and all scaffolds were rebinned to match the DASTool output. Each of the new

bins were manually inspected and scaffolds suspected of being falsely binned were removed.

After inspecting the first round bins, we improved the high coverage bins by subsampling

the read files, and repeating the SOP above [58]. In addition, refinement of the *Brocadia*

Genome bins was done with ESOMs [59] (Supplemental methods).

Post binning analysis. Unique representative genomes were determined by the dereplication

tool, dRep [60], using a 95% threshold for species level clustering. Within each cluster, the

representative genome was chosen based on their completeness, length, N50, contamination, and

strain heterogeneity. In several clusters with higher heterogeneity, a second strain was chosen

(Table S2). The strain threshold was set at 2% difference.

All representative and strain genomes were curated by correcting scaffolding errors

introduced by idba_ud, using the ra2.py program [30]. Following curation, the genomes were

processed again for gene calling and annotation (see above for details). The replication rates of

bacteria can be inferred from examining the coverage ratio between the origin of replication and

the terminus of replication. In a population that is not replicating there will be no difference in

the coverage so the ratio would be one. If the population is replicating, we expect the ratio to be

>1 since there would be replication forks that have not finished replicating and thus the coverage

towards the origin of replication would be higher than that of the terminus. Calculating

replication rate is more complicated in metagenomic samples but it is still possible to look at

overall trends in coverage across the genome. Analysis of replication rates at different time

points was performed with the iRep program [34] using the default parameters (Table S9).

Briefly, iRep calculates replication rate by measuring sequencing coverage trend that results

from bi-directional genome replication from a single origin of replication. The program uses only

high-quality draft genomes (>=75% complete, <=175 fragments/Mbp sequence, and <=2%

contamination). Since iRep is a measure of a trend it does not have any units.

Raw reads were submitted to the National Center for Biotechnology Information (NCBI)

Genbank, under project accession number PRJNA511011. In addition, the Representative and

Strain genomes were uploaded to ggkbase as two separate projects

(https://ggkbase.berkeley.edu/LAC_reactor_startup/organisms and

https://ggkbase.berkeley.edu/LAC_reactor_strains/organisms).

Phylogenetic analysis and core anammox analysis. The taxonomic affiliation of each genome

was initially assigned in ggKbase, based on the taxonomic annotation of genes in the scaffolds.

For each hierarchical taxonomic level, the taxonomy was decided if at least 50% of genes had a

known taxonomic identification.

Phylogenetic analysis of the genomes (current study, Speth et al. [18], and Lawson et al.

[20]) was based on a set of 15 ribosomal proteins [61]. Each gene was aligned separately to a set

of 3225 reference genomes, followed by concatenation while keeping the aligned length of each

gene intact. A preliminary tree was created by adding the queried genomes to the reference tree

using pplacer v1.1.alpha19 [62] and a set of in-house scripts. The tree was uploaded to iTOL

[63] for visualization and editing. After initial inspection we decided to reduce the tree in

preparation for creating a maximum likelihood tree. Large phyla with no representatives in an

anammox sample were removed (approximately 1000 sequences). The remaining sequences

were aligned by MUSCLE [64] and a RAxML tree built in The CIPRES Science Gateway V. 3.3

[64, 65].

For the analysis of phylogenetic distance between different anammox community

members, we used the APE package [66] in R [67, 68] to extract the distance matrix. Species

level distance was set at 5% of the longest measured distance on the tree. The R script and RData

files for the analysis of related species, community dynamics, and metabolic capacities were

uploaded to figshare (https://figshare.com/projects/Anammox_Bioreactor/59324).

16S rRNA gene sequencing, processing, and analysis. DNA samples, taken at 55 time points

across the lifespan of the bioreactor, were sent to the Institute for Environmental Genomics at the

University of Oklahoma (Norman, OK) for amplification of the variable 4 (V4) region of the 16S

rRNA gene, library preparation, and amplicon sequencing. The full protocol was previously

described in Wu et al. [69]. In summary, the V4 region of the bacterial 16S rRNA gene was

amplified from DNA samples using primers 515F (5'-GTGCCAGCMGCCGCGG-3') and 806R

(3'-TAATCTWTGGVHCATCAG-5'), with barcodes attached to the reverse primer. Amplicons

were pooled at equal molarity and purified with the QIAquick Gel Extraction Kit (QIAGEN

Sciences, Germantown, MD). Paired-end sequencing was then performed on the barcoded,

purified amplicons with the Illumina MiSeq sequencer (Illumina, San Diego, CA).

Subsequent sequence processing and data analysis were performed in-house using

MOTHUR v.1.39.5, following the MiSeq SOP [70, 71]. In summary, sequences were

demultiplexed, merged, trimmed, and quality filtered. Unique sequences were aligned against the

SILVA 16S rRNA gene reference alignment database [72]. Sequences that did not align to the

position of the forward primer were discarded. Chimeras were detected and removed. The

remaining sequences were clustered into operational taxonomic units (OTUs) within a 97%

similarity threshold using the Phylip-formatted distance matrix. Representative sequences from

each OTU were assigned taxonomic identities from the SILVA gene reference alignment

database [72]. Sequences that were not classified as bacteria were removed. Remaining OTUs

were counted, and the 137 most abundant OTUs (accounting for up to 99% of sequence reads

within individual samples) were transferred to Microsoft Excel (Microsoft Office Professional

Plus 2016) for downstream interpretation and visualization. The read files from all time points,

as well as the 137 most abundant OTUs were uploaded to figshare

(https://figshare.com/projects/Anammox_Bioreactor/59324).

In order to correlate genome-based OTUs to 16S rRNA gene-based OTUs, 16S rRNA

gene sequences were extracted from the representative genomes and combined with the

representative sequences from the 137 most abundant 16S rRNA gene-based OTUs. If a

representative genome did not contain the V4 region of the 16S rRNA gene, the region was

pulled from another genome in the same cluster. The combined 16S rRNA gene sequences were

aligned following the protocol described above, and those sharing at least 99% average

nucleotide identity were assumed to represent the same microorganism [73] .

Community dynamics analysis. The paired sequence reads from all time points were mapped to

the set of reference genomes using bowtie2 [51], followed by calculations for coverage (average

number of reads mapped per nucleotide) and breadth (percent of genome covered by at least one

read) for each genome per time point [74]. The multiplication of the two values was then used to

calculate the estimated relative abundance. These steps were done to negate the biases created by

repetitive sequences that occur more often in partial genome bins.

Association between genomes was tested by calculating pairwise correlation for all

genomes by relative abundance. Rho values (ranging from -1 to 1) were used to create a distance

table (Euclidean distance), followed by clustering with the ward.D method. The resulting clusters

were marked A-D. To test for the association of genomes and clusters to time points, we ran a

nMDS analysis (non-parametric MultiDimensional Scaling) with genomes and time points. Each

genome was colored by its relative abundance cluster on the 2D projection of the nMDS.

For relative abundance changes, the estimated relative abundances of genomes were

divided by the sum of all estimated relative abundance values per time point. For a clearer

resolution of changes in the four relative abundance groups, the *Brocadia* (part of group D) was

presented separately.

Metabolic analysis. The functional profiles of the genomes were evaluated using KEGG KAAS

[75], with Hidden Markov Models for shared KEGG orthologies (KOs) [28, 29, 76]. From this,

we received the KEGG annotation (KO number) for all open reading frames and a completeness

value for each KEGG module. KO annotations that were questionable were removed from the

analysis.

From the KO list, we created a presence-absence matrix (Jaccard index), and clustered

the genomes using the Complete method. From module completeness, we created a Euclidean

distance matrix, followed by clustering with the ward.D method. Based on module completeness

clustering, we assigned genomes to metabolic groups □-ε.

For each metabolic group, a representative metabolic map was created. Module

completeness greater than 67% in at least half of the group members was considered

representative of the group. Once the modules were selected, they were drawn and connected

based on metabolic KEGG maps. Additional reaction, complexes, and transporters were added

according to KO presence (e.g., aa. synthesis, oxidative phosphorylation complexes, flagellar

motor, etc.).

For nitrogen metabolism, all relevant KOs were examined. For the purpose of this study,

nitrate reduction was considered a separate path from denitrification/DNRA, since it could be the

first step in both pathways, using the same enzymes. Denitrifying bacteria were considered to be

bacteria capable of full conversion of nitrite to N_2. DNRA bacteria were considered to be

bacteria capable of conversion of nitrite to ammonium using the nrfAH enzymes. No partial

nitrogen process was considered for this paper, although it was present, according to per-step

analysis.

Notes

Ray Keren and Jennifer E. Lawrence share senior authorship.

List of abbreviations

Anammox anaerobic ammonium oxidation; **DNRA** dissimilatory nitrate reduction to

ammonium; **PN** partial nitritation; **NRR** nitrogen removal rate; **D0** day 0; **D82** day 82; **D166** day

166; **D284** day 284; **D328** day 328; **D437** day 437; **ANI** average nucleotide identity; **nMDS**

nonmetric multidimensional scaling; **AA** anammox associated; **SA** source Associated; **HMM**

Hidden Markov Model; **KEGG** Kyoto Encyclopedia of Genes and Genomes; **KO** KEGG

Orthology; **CPR** Candidate Phyla Radiation; **CI** confidence interval; **LR** log-ratio; **RFg**

reference frame genome.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

The metagenomic datasets analyzed during the current study are available on the National Center

for Biotechnology Information (NCBI) Genbank, under project accession number

PRJNA511011.

The 16S rRNA gene datasets analyzed during the current study are available on the Figshare

repository: https://figshare.com/projects/Anammox_Bioreactor/59324

The R script used to analyze the metagenomic data is also available on the Figshare repository,

under the same link as above.

All other data generated during this study is either included in this published article (and its

supplementary information files) or can be made available from the corresponding author upon

reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This research was supported by the National Science Foundation through the Engineering

Research Center for ReInventing the Nation's Water Infrastructure (ReNUWIt) ECC-1028962.

This material is also based upon work supported by the National Science Foundation Graduate

Research Fellowship under Grant No. DGE 1106400. Any opinion, findings, and conclusions or

recommendations expressed in this material are those of the authors and do not necessarily

reflect the views of the National Science Foundation.

Author contribution

K.Y. and L.Z. supervised the study. K.Y., L.Z., L.A-C., and D.J. designed the study. L.Z. built

and maintained the bioreactor. R.K. analyzed metagenomics data and wrote the manuscript. J.L.

analyzed 16S rRNA gene data, analyzed bioreactor performance, and wrote the manuscript.

J.F.B. supervised the metagenomics analysis. W.Z. and K.Y. contributed to bioreactor

maintenance and analysis, sampling, and 16S rRNA gene data analysis. All authors read the

manuscript and contributed with their input.

Acknowledgements

The authors would like to thank Edmund Antell for his help with evaluating energy gains via

nitrogen metabolism, as well as his feedback on manuscript revisions.

References

1. Mulder A, Graaf AA, Robertson LA, Kuenen JG. Anaerobic ammonium oxidation discovered in a
denitrifying fluidized bed reactor. FEMS Microbiol Ecol. 1995;16:177–84.

2. Kuenen JG. Anammox bacteria: from discovery to application. Nat Rev Microbiol. 2008;6:320–6.

3. Sonthiphand P, Hall MW, Neufeld JD. Biogeography of anaerobic ammonia-oxidizing (anammox)
bacteria. Front Microbiol. 2014;5. doi:10.3389/fmicb.2014.00399.

4. Connan R, Dabert P, Khalil H, Bridoux G, Béline F, Magrí A. Batch enrichment of anammox bacteria
and study of the underlying microbial community dynamics. Chem Eng J. 2016;297:217–28.

5. Strous M, Pelletier E, Mangenot S, Rattei T, Lehner A, Taylor MW, et al. Deciphering the evolution
and metabolism of an anammox bacterium from a community genome. Nature. 2006;440:790–4.

6. Kartal B, Maalcke WJ, de Almeida NM, Cirpus I, Gloerich J, Geerts W, et al. Molecular mechanism of
anaerobic ammonium oxidation. Nature. 2011;479:127–30.

7. Paques. Anammox sustainable nitrogen removal.
https://en.paques.nl/mediadepot/1818a31cd232/WEBbrochureAnammox.pdf. Accessed 26 Sep 2018.

8. Lackner S, Gilbert EM, Vlaeminck SE, Joss A, Horn H, van Loosdrecht MCM. Full-scale partial
nitritation/anammox experiences – An application survey. Water Res. 2014;55:292–303.

9. Ali M, Okabe S. Anammox-based technologies for nitrogen removal: Advances in process start-up and
remaining issues. Chemosphere. 2015;141:144–53.

10. Jin R-C, Yang G-F, Yu J-J, Zheng P. The inhibition of the Anammox process: A review. Chemical
Engineering Journal. 2012;197:67–79. doi:10.1016/j.cej.2012.05.014.

11. Li X, Klaus S, Bott C, He Z. Status, Challenges, and Perspectives of Mainstream
Nitritation Anammox for Wastewater Treatment. Water Environ Res. 2018;90:634–49.

12. Li X-R, Du B, Fu H-X, Wang R-F, Shi J-H, Wang Y, et al. The bacterial diversity in an anaerobic
ammonium-oxidizing (anammox) reactor community. Syst Appl Microbiol. 2009;32:278–89.

13. Gonzalez-Martinez A, Osorio F, Rodriguez-Sanchez A, Martinez-Toledo MV, Gonzalez-Lopez J,
Lotti T, et al. Bacterial community structure of a lab-scale anammox membrane bioreactor. Biotechnol
Prog. 2015;31:186–93.

14. Gonzalez-Martinez A, Osorio F, Morillo JA, Rodriguez-Sanchez A, Gonzalez-Lopez J, Abbas BA, et
al. Comparison of bacterial diversity in full scale anammox bioreactors operated under different
conditions. Biotechnol Prog. 2015;31:1464–72.

15. Gonzalez-Martinez A, Rodriguez-Sanchez A, Muñoz-Palazon B, Garcia-Ruiz M-J, Osorio F, van
Loosdrecht MCM, et al. Microbial community analysis of a full-scale DEMON bioreactor. Bioprocess
Biosyst Eng. 2015;38:499–508.

16. Pereira AD, Cabezas A, Etchebehere C, de Lemos Chernicharo CA, de Araújo JC. Microbial
communities in anammox reactors: a review. Environmental Technology Reviews. 2017;6:74–93.

17. Bagchi S, Lamendella R, Strutt S, Van Loosdrecht MCM, Saikaly PE. Metatranscriptomics reveals
the molecular mechanism of large granule formation in granular anammox reactor. Sci Rep.
2016;6:28327.

18. Speth DR, in 't M, Guerrero-Cruz S, Dutilh BE, Jetten MSM. Genome-based microbial ecology of
anammox granules in a full-scale wastewater treatment system. Nat Commun. 2016;7:11172.

19. Castro-Barros CM, Jia M, van Loosdrecht MCM, Volcke EIP, Winkler MKH. Evaluating the
potential for dissimilatory nitrate reduction by anammox bacteria for municipal wastewater treatment.
Bioresour Technol. 2017;233:363–72.

20. Lawson CE, Wu S, Bhattacharjee AS, Hamilton JJ, McMahon KD, Goel R, et al. Metabolic network
analysis reveals microbial community interactions in anammox granules. Nat Commun. 2017;8:15416.

21. Tang X, Guo Y, Jiang B, Liu S. Metagenomic approaches to understanding bacterial communication
during the anammox reactor start-up. Water Res. 2018;136:95–103.

22. van de Graaf AA, Mulder A, de Bruijn P, Jetten MS, Robertson LA, Kuenen JG. Anaerobic oxidation
of ammonium is a biologically mediated process. Appl Environ Microbiol. 1995;61:1246–51.

23. Chen H, Yu J-J, Jia X-Y, Jin R-C. Enhancement of anammox performance by Cu(II), Ni(II) and
Fe(III) supplementation. Chemosphere. 2014;117:610–6.

24. Bi Z, Qiao S, Zhou J, Tang X, Zhang J. Fast start-up of Anammox process with appropriate ferrous
iron concentration. Bioresour Technol. 2014;170:506–12.

25. Zhang X, Chen Z, Zhou Y, Ma Y, Ma C, Li Y, et al. Impacts of the heavy metals Cu (II), Zn (II) and
Fe (II) on an Anammox system treating synthetic wastewater in low ammonia nitrogen and low
temperature: Fe (II) makes a difference. Sci Total Environ. 2018;648:798–804.

26. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the quality of
microbial genomes recovered from isolates, single cells, and metagenomes. Genome Res. 2015;25:1043–
55.

27. Narita Y, Zhang L, Kimura Z-I, Ali M, Fujii T, Okabe S. Enrichment and physiological
characterization of an anaerobic ammonium-oxidizing bacterium " Candidatus Brocadia sapporoensis."
Syst Appl Microbiol. 2017;40:448–57.

28. Burstein D, Sun CL, Brown CT, Sharon I, Anantharaman K, Probst AJ, et al. Major bacterial lineages
are essentially devoid of CRISPR-Cas viral defence systems. Nat Commun. 2016;7:10613.

29. Anantharaman K, Brown CT, Burstein D, Castelle CJ, Probst AJ, Thomas BC, et al. Analysis of five
complete genome sequences for members of the class Peribacteria in the recently recognized
Peregrinibacteria bacterial phylum. PeerJ. 2016;4:e1607.

30. Brown CT, Hug LA, Thomas BC, Sharon I, Castelle CJ, Singh A, et al. Unusual biology across a
group comprising more than 15% of domain Bacteria. Nature. 2015;523:208–11.

31. Anantharaman K, Brown CT, Hug LA, Sharon I, Castelle CJ, Probst AJ, et al. Thousands of microbial
genomes shed light on interconnected biogeochemical processes in an aquifer system. Nat Commun.
2016;7:13219.

32. Sun C, Benlekbir S, Venkatakrishnan P, Wang Y, Hong S, Hosler J, et al. Structure of the alternative
complex III in a supercomplex with cytochrome oxidase. Nature. 2018;557:123–6.

33. Zomorrodi AR, Segrè D. Genome-driven evolutionary game theory helps understand the rise of
metabolic interdependencies in microbial communities. Nat Commun. 2017;8:1563.

34. Brown CT, Olm MR, Thomas BC, Banfield JF. Measurement of bacterial replication rates in
microbial communities. Nat Biotechnol. 2016;34:1256–63.

35. Morton JT, Marotz C, Washburne A, Silverman J, Zaramela LS, Edlund A, et al. Establishing
microbial composition measurement standards with reference frames. Nat Commun. 2019;10:1–11.

36. Strohm TO, Griffin B, Zumft WG, Schink B. Growth Yields in Bacterial Denitrification and Nitrate
Ammonification. Appl Environ Microbiol. 2007;73:1420–4.

37. Probst AJ, Castelle CJ, Singh A, Brown CT, Anantharaman K, Sharon I, et al. Genomic resolution of
a cold subsurface aquifer community provides metabolic insights for novel microbes adapted to high CO
concentrations. Environ Microbiol. 2017;19:459–74.

38. High definition for systems biology of microbial communities: metagenomics gets genome-centric
and strain-resolved. Curr Opin Biotechnol. 2016;39:174–81.

39. Knelman JE, Nemergut DR. Changes in community assembly may shift the relationship between
biodiversity and ecosystem function. Front Microbiol. 2014;5. doi:10.3389/fmicb.2014.00424.

40. Fukami T, Dickie IA, Paula Wilkie J, Paulus BC, Park D, Roberts A, et al. Assembly history dictates
ecosystem functioning: evidence from wood decomposer communities. Ecol Lett. 2010;13:675–84.

41. Johnston J, LaPara T, Behrens S. Composition and Dynamics of the Activated Sludge Microbiome
during Seasonal Nitrification Failure. Sci Rep. 2019;9:4565.

42. Ofiteru ID, Lunn M, Curtis TP, Wells GF, Criddle CS, Francis CA, et al. Combined niche and neutral
effects in a microbial wastewater treatment community. Proceedings of the National Academy of
Sciences. 2010;107:15345–50. doi:10.1073/pnas.1000604107.

43. DeLong EF, Preston CM, Mincer T, Rich V, Hallam SJ, Frigaard N-U, et al. Community genomics
among stratified microbial assemblages in the ocean's interior. Science. 2006;311:496–503.

44. Trivedi P, Anderson IC, Singh BK. Microbial modulators of soil carbon storage: integrating genomic
and metabolic knowledge for global prediction. Trends Microbiol. 2013;21:641–51.

45. Lackner S, Terada A, Smets BF. Heterotrophic activity compromises autotrophic nitrogen removal in
membrane-aerated biofilms: results of a modeling study. Water Res. 2008;42:1102–12.

46. Vanniel E, Arts P, Wesselink B, Robertson L, Kuenen J. Competition between heterotrophic and
autotrophic nitrifiers for ammonia in chemostat cultures. FEMS Microbiology Letters. 1993;102:109–18.
doi:10.1016/0378-1097(93)90006-n.

47. Brown EJ, Button DK, Lang DS. Competition between heterotrophic and autotrophic microplankton
for dissolved nutrients. Microbial Ecology. 1981;7:199–206. doi:10.1007/bf02010303.

48. Vet S, de Buyl S, Faust K, Danckaert J, Gonze D, Gelens L. Bistability in a system of two species
interacting through mutualism as well as competition: Chemostat vs. Lotka-Volterra equations. PLOS
ONE. 2018;13:e0197462. doi:10.1371/journal.pone.0197462.

49. Joshi NA FJN. Sickle: A sliding-window, adaptive, quality-based trimming tool for FastQ files
(Version 1.33) [Software]. Available at https://github.com/najoshi/sickle. 2011.

50. Peng Y, Leung HCM, Yiu SM, Chin FYL. IDBA-UD: a de novo assembler for single-cell and
metagenomic sequencing data with highly uneven depth. Bioinformatics. 2012;28:1420–8.

51. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012;9:357–9.

52. Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic gene
recognition and translation initiation site identification. BMC Bioinformatics. 2010;11:119.

53. Edgar RC. Search and clustering orders of magnitude faster than BLAST. Bioinformatics.
2010;26:2460–1.

54. Lowe TM, Eddy SR. tRNAscan-SE: a program for improved detection of transfer RNA genes in
genomic sequence. Nucleic Acids Res. 1997;25:955–64.

55. Alneberg J, Bjarnason BS, de Bruijn I, Schirmer M, Quick J, Ijaz UZ, et al. Binning metagenomic
contigs by coverage and composition. Nat Methods. 2014;11:1144–6.

56. Wu Y-W, Simmons BA, Singer SW. MaxBin 2.0: an automated binning algorithm to recover
genomes from multiple metagenomic datasets. Bioinformatics. 2015;32:605–7.

57. Sieber CMK, Probst AJ, Sharrar A, Thomas BC, Hess M, Tringe SG, et al. Recovery of genomes
from metagenomes via a dereplication, aggregation, and scoring strategy. 2017. doi:10.1101/107789.

58. Hug LA, Thomas BC, Sharon I, Brown CT, Sharma R, Hettich RL, et al. Critical biogeochemical
functions in the subsurface are associated with bacteria from new phyla and little studied lineages.
Environ Microbiol. 2016;18:159–73.

59. Ultsch A, Mörchen F. ESOM-maps: Tools for Clustering, Visualization, and Classification with
Emergent SOM. 2005.

60. Olm MR, Brown CT, Brooks B, Banfield JF. dRep: a tool for fast and accurate genomic comparisons
that enables improved genome recovery from metagenomes through de-replication. ISME J.
2017;11:2864–8.

61. Hug LA, Baker BJ, Anantharaman K, Brown CT, Probst AJ, Castelle CJ, et al. A new view of the tree
of life. Nat Microbiol. 2016;1:16048.

62. Matsen FA, Kodner RB, Armbrust EV. pplacer: linear time maximum-likelihood and Bayesian
phylogenetic placement of sequences onto a fixed reference tree. BMC Bioinformatics. 2010;11:538.

63. Letunic I, Bork P. Interactive tree of life (iTOL) v3: an online tool for the display and annotation of
phylogenetic and other trees. Nucleic Acids Res. 2016;44:W242–5.

64. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic
Acids Res. 2004;32:1792–7.

65. Miller MA, Pfeiffer W, Schwartz T. Creating the CIPRES Science Gateway for inference of large
phylogenetic trees. In: 2010 Gateway Computing Environments Workshop (GCE). 2010.
doi:10.1109/gce.2010.5676129.

66. Paradis E, Claude J, Strimmer K. APE: Analyses of Phylogenetics and Evolution in R language.
Bioinformatics. 2004;20:289–90.

67. RStudio Team (2015). RStudio: Integrated Development for R. RStudio, Inc., Boston, MA.
http://www.rstudio.com/.

68. R: The R Project for Statistical Computing. http://www.R-project.org. Accessed 26 Sep 2018.

69. Wu L, Wen C, Qin Y, Yin H, Tu Q, Van Nostrand JD, et al. Phasing amplicon sequencing on
Illumina Miseq for robust environmental microbial community analysis. BMC Microbiol. 2015;15:125.

70. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, et al. Introducing mothur:
open-source, platform-independent, community-supported software for describing and comparing
microbial communities. Appl Environ Microbiol. 2009;75:7537–41.

71. Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. Development of a dual-index
sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina
sequencing platform. Appl Environ Microbiol. 2013;79:5112–20.

72. Pruesse E, Quast C, Knittel K, Fuchs BM, Ludwig W, Peplies J, et al. SILVA: a comprehensive
online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB.
Nucleic Acids Res. 2007;35:7188–96.

73. Evans J, Sheneman L, Foster J. Relaxed neighbor joining: a fast distance-based phylogenetic tree
construction method. J Mol Evol. 2006;62:785–92.

74. Olm MR, Brown CT, Brooks B, Firek B, Baker R, Burstein D, et al. Identical bacterial populations
colonize premature infant gut, skin, and oral microbiomes and exhibit different in situ growth rates.
Genome Res. 2017;27:601–12.

75. Moriya Y, Itoh M, Okuda S, Yoshizawa AC, Kanehisa M. KAAS: an automatic genome annotation
and pathway reconstruction server. Nucleic Acids Res. 2007;35 Web Server issue:W182–5.

76. Finn RD, Clements J, Arndt W, Miller BL, Wheeler TJ, Schreiber F, et al. HMMER web server: 2015
update. Nucleic Acids Res. 2015;43:W30–8.

Additional Files

Additional file 1: **Table S1** Genome parameters for the representative bacteria from the

anammox bioreactor. **Temporal strain shifts in the bioreactor**. Analysis of shifts between bacterial

strains during the first bioreactor crash and subsequent biomass amendment. **Table S2.** Genome

parameters for the bacterial strains related to representative bacteria from the bioreactor. **Table S3**

assignment of genomes to different clusters. **Figure S1**. Analysis of community metabolism by gene

presence. **Figure S2.** Reduced metabolic map of the Brocadia sp. Bacterium. **Electron transfer**

pathways in the community. Analysis of different electron donors and acceptors utilized by the

anammox community members. **Central carbon metabolism in the community**. Analysis of carbon

utilization in the anammox microbial community. **Table S4** Diversity indices of the anammox

community. **Table S5**. Synthesis and transport of metabolites in the metabolic groups. **Table S6**

Confidence intervals of the Log-Ratio changes for each Reference Frame genome. **Table S7.** Closest

relative based on distance in ML tree for anammox bacterial community. **Figure S3.** Maximum likelihood

tree of the PVC clade (Planctomycetes, Verrucomicrobia, Chlamydiae, and Lentisphaerae). **Figure S4.**

Schematic of the anaerobic membrane bioreactor. **Table S8**. Composition of the synthetic wastewater

feed. **Supplemental methods.** Refinement of the *Candidatus Brocadia* sp. genome. **Table S9**.

Replication rates of genomes for each time point, calculated using the iRep program.

Figure legends

Figure 1 | Performance of the anaerobic membrane bioreactor. Influent and effluent

concentrations of ammonium, nitrite, and nitrate (all as N) (primary y-axis) within the anaerobic

membrane bioreactor performing anammox monitored over a period of 440 days. The influent

did not contain nitrate, so influent nitrate is not plotted. The nitrogen removal rate (NRR), is

plotted against the secondary y-axis. Sampling time points for metagenomes are indicated with

purple stars below the x-axis. Biomass amendments are indicated with green stars below the x-

axis. Bioreactor crashes (either mechanical or biologically driven) are marked with a dashed red

line.

Figure 2| Analysis of the bioreactor community clustering by the estimated relative

abundance of the bacteria. (A) clustering heatmap of bacteria based on pairwise cross

correlations for the six time points (matrix values are Rho values). Color scales mark high

positive correlation in green and negative correlation in brown. The row and column

dendrograms are identical (consisting of the genomes). The row dendrogram shows the

calculated distance between the clusters with a dashed red line marking the split to clusters.

Colored squares as well as bars to the left of the heat map show relative abundance based

grouping: Yellow - Group A, Blue - Group B, Purple - Group C, Red - Group D. A black star to

the right of the heatmap marks the anammox bacterium (*Brocadia*). (B) Two-dimensional nMDS

projection of bacteria and time points, showing the association of the bacteria (and relative

abundance groups to certain time points). Each colored dot represents the centroid of a

bacterium, with colors matching the relative abundance group. Black marks represent the

centroid of the time points, while the shape represents the state of the bioreactor: Circle -

anammox dominated, Triangle - bioreactor crash (either mechanical or biological), Square - time

zero. The location of *Brocadia* is marked with a red star.

Figure 3 | Relative abundances of bacterial groups over the lifespan of the bioreactor. (A)
Relative abundances of bacterial groups A-D, based on 16S rRNA OTUs that have a match to a
draft genome. Group colors are matched by the analysis in Figure 2, with the exception of
Brocadia, which was removed from group D and is depicted in black. Green points above the
graph show timepoints where the community was sampled. (B) Comparison of relative
abundances of the different groups at the same time (or the closest time). Relative abundance
was calculated for groups based on all the draft genomes (marked met on the x-axis), the subset
of genomes with matched to the 16S rRNA OTUs (marked sub-met on the x-axis) or the 16S
rRNA OTUs (only day is marked on x-axis).

Figure 4| Metabolic profiling of bacterial community based on KEGG module
completeness. (A) heatmap showing clustering of genomes (rows) by their KEGG module
completeness (columns). Completeness ranges from 1 (green) to zero (white). The heatmap is
based on a Euclidean distance matrix and clustering with the ward.D method. Genome clustering
resulted in 5 clusters (groups α-ε). Rectangles on the heatmap mark module blocks that
differentiate the genome groups. Black rectangles on heatmap show module blocks that have
increased completeness in a group of bacteria (compared to the others), and red rectangles show
decreased completeness. Marks to the left of the heatmap show the division of AA and SA
bacteria. A black star to the right of the heatmap marks *Brocadia*. B) Relative abundance by
phyla of members in each metabolic cluster.

Figure 5| Representative metabolic maps of bacterial groups in the bioreactor. To prevent

redundancy, the metabolism is presented in a nested approach with each panel showing only

paths unique to the relevant metabolic group. Two exceptions are group β (all detected paths are

shown), and group δ. The latter is not presented here since it shares all paths with group □ and

only differs by auxotrophies. (A) Metabolic map of paths that are common to all bacteria in the

bioreactor (except *Microgenomates* and *Brocadia* sp.). The vast majority of bacteria in the

bioreactor are heterotrophs, capable of carbohydrate-based metabolism (glycolysis, pentose

phosphate pathway) and amino acid-based metabolism. Some bacteria can respire oxygen, and

can also ferment (acetate/alanine). (B) Paths unique to group □. These bacteria have genes for

hydrogen oxidation, supporting anaerobic growth, as well as genes for oxidative phosphorylation

with cytochrome BD complex. These bacteria have a cassette of extracellular proteases and

decarboxylases, paired with a wide array of transporters. Theyare also potentially capable of

synthesizing long chain isoprenoids. (C) Paths found in Gram (-) bacteria (groups □, δ, and ε).

Most paths are related to fatty acid and lipid synthesis. Several important precursors (chorismate

and IMP) can potentially be synthesized by these bacteria. Motility is also a common feature in

these bacteria (via a flagellar motor) (D) Unique paths of group ε (*Proteobacteria*). This group

has the potential to synthesize multiple vitamins and cofactors (biotin, pyridoxal, glutathione,

etc.), as well as several amino acids (tyrosine, phenylalanine, proline). Another unique feature is

the multiple secretion systems present in the bacteria. (E) Metabolic profile of CPR bacteria

(*Microgenomates*). These bacteria are obligate anaerobes that ferment pyruvate. They can only

utilize carbohydrates as their carbon source. Some of the bacteria in this group might also be able

to synthesize long chain isoprenoids, in the same path as group □.

Figure 6| Nitrogen cycle in the anammox bioreactor. The steps in the nitrogen cycle are color-

coded by their association to different types of metabolism. The number of bacteria with genes

encoding a given step is listed, and the pie chart depicts the ratio between AA and SA bacteria

associated with the step.

1000

Figure 7| Potential metabolic interdependencies between the metabolic groups in the

anammox bioreactor. (A) All potential metabolic interdependencies combined. (B) Metabolic

interdependencies for vitamins/cofactors alone. (C) Metabolic interdependencies for lipids/fatty

acids alone. (D) Metabolic interdependencies for amino acids and peptides alone. Arrows were

assigned according to the absence of a group's ability to synthesize a metabolite, and they

connect to all groups that have the ability (there is redundancy in arrows). The arrowhead points

at the group that receives the metabolite. The width of the arrow is proportional to the ratio of

metabolites of a given type that are provided; amino acids - 20 metabolites; Peptides - deduced

from proteases and transporters (Figure 5B); Vitamins/Co-factors - 10 metabolites; Lipids/Fatty

acids - 7 metabolites. The size of each group is proportional to their relative abundance at Day

437. Group β is not shown since it is assumed that all members obtain all of their nutrients and

metabolites from their host. Overall, groups ☐ and δ receive the most metabolites and group ε

receives the least. Group δ has the highest number of aa. synthesis auxotrophies and can

potentially acquire these from many other community members. Group ε has only a single

auxotrophy in vitamin/Co-factor synthesis, while most other groups have multiple auxotrophies.

Brocadia sp. is the only bacterium capable of vitamin B12 synthesis.

1017

Figure 8 | Tracking bacterial growth and nitrogen metabolism in relation to the anammox

performance destabilization. (A) Distribution of Log-Ratio changes for selected bacteria

between D328 and D284, using different genomes as reference frames. Bacteria that are

significantly affected or considered to affect the destabilization event are color-coded. Down-

pointing triangles depict bacteria negatively affected by the destabilization and upwards-pointing

triangles indicate bacteria that likely caused (or contributed) to the destabilization. (B)

Replication rate values on days 166 and 284. Bacteria are color coded the same as in panel A.

(C) Relative abundance of nitrogen metabolism pathways in the selected bacteria. Denitrification

is divided into its reaction steps. Anammox is considered as a single path since only a single

bacterium can perform it (*Brocadia* – LAC_PLT02). DNRA from NO2 is a single step reaction.

Abundance was calculated twice, once allowing multiple pathways to occur within each

bacterium, and once after choosing a single pathway per bacterium (based on potential energy

gain).

1031

Figure 9 | Phylogenetic analysis of three anammox microbial communities. (A) A maximum

likelihood tree based on the alignment of 15 concatenated ribosomal proteins. In the construction

of the tree 3225 reference sequences were used, with genomes from current and previous

genome-centric studies on anammox communities. Genomes from the current anammox

community are marked with a red dashed line, genomes from two previously studied

communities; Speth et al. and Lawson et al., are marked with green and blue dashed lines,

respectively. Nodes containing only reference genomes were collapsed for ease of view.

Collapsed nodes are depicted as triangles and their size is relative to the number of bacteria they

contain. A black star marks *Brocadia*. (B) relative abundance of major phyla in the three

microbial communities. Current community reference data was calculated from day 437 only.

The relative abundance of *Brocadia* sp. comprises nearly all of the relative abundance attributed

to phylum *Planctomycetes* (with small contributions from other members of the phylum). The

most abundant phyla (*Chloroflexi*, *Ignavibacteria*, and *Proteobacteria*) consistently account for

>70% of the communities. The phyla colors follow the ggkbase color scheme and the major

phyla are shown in the legend.

1047

1048
1049   **Table 1 | Read counts to representative genomes across time points**

Time point (days)	Total read number	Mapped read number	% of reads mapped	Number of genomes detected*
0	55398280	40291503	72.73	92
82	62544544	43877427	70.15	82
166	60931806	46030350	75.54	103
284	56282006	48644523	86.43	68
328	127048582	95132145	74.88	87
437	119945232	93737087	78.15	60

1050

1051   * number of representative genomes based on threshold of >1 coverage and > 0.5 breadth

1052   **Table 2 | Log-Ratio and replication rates (iRep) for select bacteria**

Genome	LR$_{(RF1)}$	LR$_{(RF2)}$	LR$_{(RF3)}$	LR$_{(RF4)}$	iRep D166	iRep D284
LAC_CHLX01	1.05	1.17	2.27	1.73	1.16	1.32
LAC_PROT02	-1.54	-1.41	-0.31	-0.85	1.17	1.30
LAC_PROT03	-1.50	-1.37	-0.27	-0.81	1.32	1.40
LAC_CHLX02	-2.91	-2.79	-1.68	-2.23	1.16	1.17
LAC_ACD03	-0.99	-0.86	0.24	-0.30	1.19	1.35
LAC_BACT04	-0.77	-0.64	0.46	-0.08	1.26	1.26
LAC_IGN05	-0.33	-0.20	0.90	0.36	1.38	1.51
LAC_PROT19	-2.09	-1.96	-0.86	-1.40	1.22	1.32
LAC_IGN07	-0.98	-0.85	0.25	-0.29	1.30	1.48
LAC_PROT21	-1.18	-1.06	0.04	-0.50	NA	1.29
LAC_BAC19	-0.13	0.00	1.10	0.56	NA	1.38

LAC_BAC20	-1.17	-1.04	0.06	-0.48	1.13	1.19
LAC_BACT12	-1.72	-1.59	-0.49	-1.03	1.14	1.18
LAC_NIT01	-1.93	-1.80	-0.70	-1.24	1.25	1.31
LAC_CHLX06	-0.18	-0.05	1.05	0.51	1.21	1.28
LAC_PROT22	-1.23	-1.10	0.00	-0.54	1.36	1.24
LAC_PLT02	-2.12	-1.99	-0.89	-1.43	2.13	1.07
LAC_CHLX07	0.00	0.13	1.23	0.69	1.69	NA
LAC_BAC21	-1.31	-1.18	-0.08	-0.62	1.13	1.25
LAC_CHLX09	0.25	0.37	1.48	0.93	1.56	1.63
LAC_CHLX10	0.43	0.55	1.66	1.11	1.26	1.42
LAC_BAC22	0.27	0.40	1.50	0.96	NA	1.39
LAC_BAC23	-0.31	-0.18	0.92	0.38	1.17	1.17
LAC_PROT27	-0.35	-0.23	0.88	0.33	2.06	1.56
LAC_PROT28	-0.69	-0.56	0.54	0.00	1.33	1.27
LAC_PROT30	-0.48	-0.36	0.75	0.20	NA	1.89
LAC_D-T02	-0.43	-0.30	0.80	0.25	NA	1.59

1053

1054

A

B

A

Amino acids
Peptides
Vitamins/Co-factors
Lipid/Fatty acids

B

Vitamins/Co-factors

C

Lipid/Fatty acids

D

Amino acids
Peptides

A

Causing
destabilization ▲

Affected by
destabilization ▼

B

LAC_PLT02 ■ LAC_CHLX01 ■ LAC_CHLX06 ■ LAC_CHLX09 ■ LAC_CHLX10 ■ LAC_IGN05 ■ LAC_PROT22 ■

C

Metabolic reaction

■ Anammox
■ NO$_2$ reduction
■ NO reduction
■ N$_2$O reduction
■ DNRA

A

Current (n= 127)
Speth et al (n= 22)
Lawson et al (n= 14)

Archaea
TM7
Microgenomates
Spirochaetes
Alphaproteobacteria
Gammaproteobacteria
Betaproteobacteria
Chloroflexi
Acidobacteria
Nitrospirae
Dadabacteria
Deltaproteobacteria
Actinobacteria
Omnitrophica
Deinococcus Thermus
Verrucomicrobia
Bacteroidetes
Planctomycetes
Ignavibacteriae

Tree scale: 0.1

B

Acidobacteria
Actinobacteria
Armatimonadetes
Bacteroidetes
Chloroflexi
Deinococcus Thermus
Deltaproteobacteria
Ignavibacteria
Microgenomates
Nitrospirae
Omnitophica
Parcubacteria
Planctomycetes
Proteobacteria

1.0

0.8

0.6

0.4

0.2

0.0

LAC   Speth 1.4m   Speth 3.8m   Lawson