1 TITLE: Invasive Siberian chipmunks *Eutamias sibiricus* in Italy: a socio-ecological analysis
2 indicates that they could, and should, be removed

3

4 AUTHORS: Jacopo Cerri^{1*}, Emiliano Mori², Rudy Zozzoli³, Andrea Gigliotti⁴, Antony Chirco⁵,
 5 Sandro Bertolino⁶

6

7 AUTHORS' INSTITUTIONAL AFFILIATIONS:

- Istituto Superiore per la Protezione e la Ricerca Ambientale, Piazzale dei Marmi 12, 57123,
 Livorno, Italy. Email: jacopo.cerri@isprambiente.it, mobile phone: (+39)3395692346.
 ORCID ID: 0000-0001-5030-0376. Contribution: major contribution in the conception and
 design of the study, altogether with the analysis and interpretation of data and with
 manuscript drafting;
- Dipartimento di Scienze della Vita, Università degli Studi di Siena. Via P.A. Mattioli 4,
 53100, Siena, Italy. Email: <u>moriemiliano@tiscali.it</u>. Italy. ORCID ID: 0000-0001-8108 7950. Contribution: conception and design of the study, data collection and interpretation of
 the data. Important contribution in manuscript editing;
- Dipartimento di Scienze Chimiche, della Vita e della Sostenibilità Ambientale, Università
 degli Studi di Parma, Parco Area delle Scienze, 11a, 43124, Parma, Italy. Email:
 <u>zozzoli.rudy@gmail.com</u>. Contribution: conception and design of the study, data
 acquisition on the field, intellectual role in manuscript revision;
- 4. Dipartimento di Scienze delle Produzioni Animali e dell'Ambiente, Università degli Studi di 21 22 Firenze, Piazzale delle Cascine, 18, 50144 Firenze, Italy. Email: 23 andrea.gigliotti1988@libero.it. Contribution: data acquisition on the field, data intepretation, intellectual role in manuscript revision; 24

25	5. Dipartimento di Biologia, Università degli Studi di Firenze, Via Madonna del Piano, 6,
26	50019 Sesto Fiorentino FI. Email: <u>antony.chirco@stud.unifi.it</u> . Contribution: data
27	acquisition on the field, data intepretation, intellectual role in manuscript revision;
28	6. Department of Life Sciences and Systems Biology, University of Turin, Via Accademia
29	Albertina 13, 10123 Torino, Italy. Email: <u>sandro.bertolino@unito.it</u> . Telephone:
30	(+39)0116704513. ORCID ID: 0000-0002-1063-8281. Contribution: conception and design
31	of the study, analysis and interpretation of data and manuscript drafting;
32	
33	CORRESPONDING AUTHOR: Jacopo Cerri, Istituto Superiore per la Protezione e la Ricerca
34	Ambientale, Piazzale dei Marmi 12, 57123, Livorno, Italy. Email: jacopo.cerri@isprambiente.it,
35	mobile phone: (+39)3395692346. ORCID ID: 0000-0001-5030-0376.
36	
37	
38	
39	
40	
41	
42	
43	
44	
45	
46	
47	
48	
49	
50	

51 ABSTRACT

BACKGROUND: Eradication of invasive alien species is a form of pest control linked to 52 biodiversity conservation that usually involves killing animals. Squirrels are prominent among 53 invasive alien species in Italy with four species introduced. Three of them are included within the 54 list of alien species of European concern and their eradication and control is recommended. 55 However, their local control is not an easy task, being highly appreciated by the general public. We 56 propose a socio-ecological approach to evaluate the feasibility of eradicating Siberian chipmunks 57 (Eutamias sibiricus) populations. We performed a structured questionnaire to assess the social 58 perception of invasive Siberian chipmunks in urban parks where they occur and to identify groups 59 of visitors who might oppose eradication. We also carried out geographic profiling to test for the 60 spatial expansion of chipmunk populations. 61

RESULTS: Overall, park visitors regarded chipmunks positively and appreciated to see them, but human-chipmunk interactions were still rare. We did not identify any group of visitors with a strong attachment to chipmunks, who might oppose future control programs. Geographic profiling showed that chipmunks in Valeggio sul Mincio are starting to expand outside of their introduction site.

67 CONCLUSIONS: Data from questionnaires show that chipmunks eradication, coupled with 68 adequate communication initiatives, might be feasible. Moreover, geographic profiling indicates 69 that time for a rapid removal is running out. Socio-ecological approaches, combining the analysis of 70 structured questionnaires administered to stakeholders and statistical modeling of pest observations, 71 could be a valuable tool to decide the feasibility and the urgency of invasive pest control.

- 72
- 73 Keywords: Alien squirrels, eradication, geographic profiling, pest control, socio-ecological model
- 74
- 75
- 76

77 1 INTRODUCTION

The Siberian chipmunk *Eutamias sibiricus* is a widespread species in Russia and the Far East, 78 which has become invasive in some European countries since the 1960s after its widespread trade 79 as a pet species.¹ Chipmunks established viable populations In Italy¹, chipmunks that escaped from 80 captivity established four viable populations in Northern Italy, and at two urban parks in Rome. 81 Siberian chipmunks are not a mainstream invader, as the Eastern grey squirrel *Sciurus carolinensis*;² 82 however, they are considered an invasive alien species of European concern being listed within the 83 EU Regulation 1143/2014. Particularly, chipmunks can act as a vector of tick-borne diseases and 84 zoonoses.^{3,4} The European regulation requires member states to eradicated listed species from their 85 territories when it is still possible and this appear the situation in Italy where the species is still 86 localized with small populations.^{5,6} 87

Management interventions aimed at containing or removing invasive alien mammals are more feasible when two ideal conditions occur. First, they are more cost-effective and face a higher success rate whenever the target species are still in the early stages of their invasion.⁷⁻⁹ Second, eradication initiatives tend to be more feasible when target species have minor interactions with society.¹⁰ Attempts to remove iconic alien mammals could result into strong opposition from some stakeholders.^{11,12} For instance, a trial eradication of the gray squirrel from Italy attempted in 1990s was stopped by animal right groups who brought the case in front of the court.¹³

95 Preliminary social impact assessments can tell managers whether control interventions will be 96 opposed by relevant stakeholders and how they can be designed accordingly, to be successful. ^{14,15} 97 Socio-ecological assessments go one step further, by combining information from relevant 98 stakeholders, obtained through qualitative or quantitative methods from the social sciences, with 99 information about the ecology, distribution and population dynamics of target species¹⁶. Ecological 100 information can also be spatially explicit, as most ecological processes incorporate a geographical 101 dimension¹⁷.

Eradicating chipmunks from Italy is required under the European and national legal framework, but, to the best of our knowledge, no study was conducted to verify if the preconditions for a successful eradication occur. Social prerequisites are particular sensitive considering the strong opposition faced by managers aiming to control or eradicate the gray squirrel in this country, even in recent vears.^{13, 18, 19}

Our research aims to fill this gap, by conduction a socio-ecological analysis combining spatial data of the species altogether with information from a structured surveys administered to a sample of visitors. Our analysis aims to test whether chipmunks became an iconic species and to inspect patterns in their geographical spread over time, to identify at which stage their invasion might be.

111

112 2 MATERIALS AND METHODS

In this research we focused on the three urban parks where chipmunks established viable populations in Italy. The first one is Sigurtà Garden Park, in Valeggio Sul Mincio, where Korean chipmunks were released in 1978, establishing the largest Italian populations of chipmunks in Italy.^{1, 20-22} The latter two areas where two urban parks in Rome, Villa Ada, where chipmunks were introduced at multiple times since the early 1908s and Villa Doria-Pamphilii, where chipmunks were observed for the first time in 2018.⁶

We surveyed a sample of visitors, administering a structured questionnaire measuring some their 119 interactions with chipmunks and some psychological drivers of human-chipmunk interactions: 120 attitudes, emotional dispositions, core affect, existence beliefs, social norms and behavioral 121 intention about the presence of chipmunks. Attitudes were measured by means of a Likert scale and 122 they were conceptualized as divided in some beliefs, characterized each one by its strength and the 123 evaluation of its outcome.²³ The attitudinal scale was built up by considering all the potential 124 impacts of a species of ground squirrels living in a park, after a pilot study (S1). Emotions were 125 measured as emotional disposition (joy, fear, surprise, disgust, interest) and core affect, or the extent 126 respondents would have felt positive or negative at the idea of encountering a chipmunk.²⁴⁻²⁷ We 127

128 measured existence beliefs by asking respondents to rate the importance of chipmunks in the park, both for future generations and per se.²⁸ We measured social norms about the appropriateness of 129 chipmunk presence in the park, by using three items measuring moral beliefs, empirical and 130 131 normative expectations, and the willingness to enforce them by reporting the presence of chipmunks to local authorities.²⁹ Visitors were also asked whether they had ever heard of chipmunks living in 132 the park and if they had ever seen, fed or touched them. A complete list of the various questions 133 adopted in the questionnaire, altogether with their summary is available in Table 1 and a complete 134 of copy the questionnaire 135 at

136 (https://docs.google.com/forms/d/e/1FAIpQLSelkWacunZsTWDB3Qxy3QCYvfaFKg-

hhd9LFmhcLcEZAbyfuA/viewform?usp=sf_link). The questionnaires were administered in Rome in both parks, but the collected data were grouped together as the two parks share the same pool of visitors. The questionnaire was implemented on GoogleForms. Most respondents (93.95%) were recruited on the field and they completed the questionnaire on a tablet. An online version was also administered on some Facebook groups on these urban parks. Questionnaires were confidential and they took approximately 15 minutes to be filled.

We assessed the reliability of our attitudinal scale through McDonald's Omega,³⁰ and we tested for 143 construct validity through Confirmatory Factor Analysis (CFA), with a Maximum Likelihood 144 estimator with robust standard errors and a Satorra-Bentler scaled test statistic. Both indicator and 145 latent variables were standardized and all the factor loadings were estimated. We adopted two 146 correlated latent variables reflecting the strength of each beliefs ad the evaluation of its outcome 147 and we also specified some residual correlations between each couple of items describing a specific 148 impact. We selected the best subset of items and the best latent variable structure by comparing 149 150 models through likelihood-ration testing and some fitness indexes. Attitudes were aggregated into a final score by summing the product of each couple of items.²³ 151

We segmented respondents on the basis of their attitudes, emotional dispositions, core affect and their moral, empirical and normative expectations about the presence of chipmunks in the park.

Segmentation aimed to identify clusters of respondents who strongly supported chipmunk presence and could oppose their eradication. We tested for the presence of clusters in the data through the Hopkins index and we compared k-means, hierarchical and k-medoid cluster analysis³¹ to assess which one clustered observation the best.

We carried out Generalized Linear Modeling to highlight differences in the two areas, in terms of attitudes scores, core affect, existence beliefs and moral, empirical and normative expectations about the presence of chipmunks in the park. In each model we included a dummy variable assessing whether respondents ecountered chipmunks at least once in their lifetime, and an interaction term, to account for the effect of past behavior and belief saliency.

Finally, we tested for chipmunk expansion outside of their introduction sites by means of 163 geographic profiling (GP). GP is common in criminology, where the spatial locations of crimes are 164 used to calculate the probability of occurrence of the offender's residence for each point over a 165 certain geographical area. GP outperforms classical measures of spatial tendency, and many 166 ecologists found it good for tracing back the origin of individuals that could move across space.³²⁻³⁵ 167 We adopted a Bayesian GP algorithm,³⁶ requiring only the specification of a distribution parameter, 168 indicating a plausible maximum extent to which individuals could move. Based on available 169 evidence indicating that chipmunks usually disperse within a few hundred meters from their 170 birthplace,³⁷⁻³⁸ we opted for a dispersal parameter of 1 km. We used available observations collected 171 in Villa Ada, from 2011 to 2014 (n=26), and in Valeggio sul Mincio, from 1997 to 2018 (n=87), as 172 the input for the GP algorithm. We did not use observations from Villa Doria-Pamphilii as the park 173 is embedded in an urbanized matrix, which prevents chipmunks from dispersing around. It is 174 important to note that we were not interested to identify where chipmunks were released, but to 175 176 reconstruct a probabilistic profile for the origin of the observations: the inspection of its shape told us whether observed chipmunks came from disjoint hotspots, as expected for an expanding invasive 177 alien species, or from a single one, like in the case of a species which is not expanding. Statistical 178 analyses were carried out with the statistical software R (R Core Team 2018) and a detailed 179

information about statistical analysis, altogether with a reproducible software code is available inthe Supplementary Information (S2).

182

183 3 RESULTS AND DISCUSSION

Respondents had generally positive emotions towards chipmunks. Moreover, they generally agreed 184 with the idea that the presence of chipmunks in the park was important for future generations and 185 that it was important to have chipmunks living in the park even if one does not see them. Finally, 186 most respondents deemed right and common for chipmunks to live in urban parks (Table 1). 187 188 However, most respondents were not aware of the presence of chipmunks in the park where they were interviewed and about half of them had never observed these animals before. The proportion 189 of respondents who had fed (14.6%) or touched (6.3%) chipmunks were even lower. Moreover, 190 191 14.11% of respondents reported to have observed chipmunks, despite they were not aware of their presence (Figure 1). 192

193 CFA and McDonald's Omega did not support an overall attitudinal construct, but they identified 194 two separate groups of beliefs. The first one included items about the impact of chipmunks over the 195 quality of recreation at the park: increasing the aesthetic appeal of the park, attracting new visitors 196 and making visitors more prone to visit the park again. The second group included the potential 197 impacts of chipmunks over human health: rummaging garbage from bins, transmitting disease to 198 humans and to visitors' dogs (Table 2; Table 3).

Hierarchical cluster analysis with Euclidean distance and a complete link indicated the presence of a small segment of respondents, characterized by negative attitudes about chipmunk impacts over human health, fear and disgust towards chipmunks (Figure 2). These could be people who are scared or disgusted by rodents and concerned about their impact over hygiene, two aspects that are often related.³⁹⁻⁴¹

The two sites differed only in respondents' score about the positive impact of chipmunks over the recreational experience, with Rome having slightly higher scores. Visitors in Rome also agreed

slightly more with the idea that most people deemed appropriate for chipmunks to live in urbanparks in Italy (Figure 3; Table 4).

Overall, these findings indicate that visitors regard chipmunks positively. Apart from a small cluster 208 209 of respondents, most of them have positive emotions towards chipmunks, deem appropriate the fact that they live in an urban park and value their presence as something having an intrinsic value. 210 However, this positive perception probably stems from a more general, favorable, disposition 211 towards the presence of wildlife at urban areas. Respondents do not have a coherent system of 212 attitudes about the presence of chipmunks, probably because their real interactions were limited: 213 attitudes are shaped and reinforced by our everyday experience with a certain issue, that make it 214 salient for ourselves.^{23,42} On the other hand, visitors had stable beliefs about those impacts of 215 chipmunks that could affect their recreational experience at the park, as well as fears about those 216 impacts that could undermine hygiene. These two sub-dimensions probably indicate that visitors' 217 beliefs are embedded in broader belief networks encompassing different, and more salient, topics.⁴³ 218 For example, our respondents could have stable belief networks diseases, and they could have tied 219 to them some of their beliefs about chipmunks. Framing experiments, where participants are primed 220 to think about some precise topics and where the effect of this priming over beliefs is measured⁴⁴ 221 might be a valuable tool to better investigate how human-wildlife interactions are embedded into 222 broader nomological networks, and influenced by beliefs about relevant social issues. Framing 223 experiments could also be used to test for attitude certainty and strength.^{45,46} 224

The idea that respondents' attitudes were not grounded into experience is reinforced by the limited interactions between visitors and chipmunks: approximately, only half visitors observed chipmunks in the park, 15% of them fed chipmunks and only about 6% of them reported to have touched a chipmunk. Moreover, some visitors who observed chipmunks were not aware of their stable presence in the park: our questionnaire was arguably the first time they were introduced to this aspect. These superficial interactions are also reflected by the low differences between the two areas. Respondents in Rome and in Valeggio sul Mincio had similar scores for almost all the psychological antecedents of their interactions with chipmunks. They showed minor differences only in their beliefs about the impact of chipmunks over the recreational experience, and in their normative beliefs about the presence of chipmunks in an urban green area. Moreover, hierarchical cluster analysis did not divide respondents into meaningful segments, on the basis of their attitudes, emotions, existence values and social norms towards chipmunks and it did not identify any group of strong supporters of chipmunks.

Taken together, these three points are important for the future management of chipmunks in the 238 study area. Attitudes are an antecedent of human behavior and often a good barometer to forecast an 239 eventual opposition to the management of invasive native²³ and introduced wildlife.⁴⁷ As visitors do 240 not have stable attitudes and no segments of highly motivated visitors exist, it is reasonable to say 241 that an eradication campaign would not face any strong opposition from local visitors. Chipmunks 242 243 at the two sites do not seem to be an iconic species vet like the grav squirrel in many urban parks if the UK.⁴⁸ Their interactions with visitors, especially those creating emotional bindings, like feeding, 244 are still limited. However, considered that respondents regard chipmunks favorably and that they 245 value their presence as a legacy for the younger generations, we believe that eradication initiatives 246 should be coupled with an adequate communication strategy, to avoid polarization and the 'backfire 247 effect'. Considered that respondents from the two sites did not show any particular difference, 248 communication actions might be similar for Rome and Valeggio sul Mincio. 249

Geographic profiling confirmed that invasive chipmunks disperse less than 500 m from the place 250 where they are born. However, while chipmunk observations in Rome come from a single source, 251 observations in Valeggio sul Mincio are likely to have involved individuals coming from two 252 distinct spatial cores (Figure 4). One of these two cores was found to be outside of the boundaries of 253 254 the urban park where the species was introduced: although slowly, chipmunks are expanding outside Sigurtà Garden Park, their introduction site in Valeggio sul Mincio. As chipmunks increase 255 their density we expect them to continue their expansion outside of the park. This perspective is not 256 encouraging, because Valeggio sul Mincio is surrounded by a countryside environment and 257

cultivations, that might promote chipmunks dispersal at the landscape scale²⁰ and maybe even their role as a pest species damaging crops, in the near future. Considered that chipmunks are still distributed over a relatively small area, but there are large areas in Italy suitable for the species ^{49,50}, we recommend their quick removal, as it will be easy and cost-effective.

262

263 4 CONCLUSION

In this research, we showed how spatially-explicit data about a biological invasion and survey data 264 about its social perception can inform decision makers about the feasibility, and the urgency, of 265 management actions. Geographic profiling can be used not only to identify introduction, or to locate 266 dens of invasive pests but also to signal the emergence of source-sink systems. These systems 267 indicate the end of an early invasion stage and the spatial expansion of the invasive alien species, 268 269 often due to their numerical increase, which can make eradication or control harder and expensive. Moreover, structured surveys could inform conservationists about the interactions between 270 stakeholders and biological invaders, altogether with their social perception. 271

Our findings indicate that visitors still have limited interactions with invasive chipmunks, at the 272 urban parks in Italy where they have been introduced. They do not have stable attitudes, and there 273 seems not to be any group of visitors who regard chipmunks as an iconic species. Perspectives 274 about chipmunks are positive, but probably weak. At the same time, chipmunks are expanding 275 outside of their introduction site in Northern Italy. We deem that initiatives aimed at removing 276 chipmunks are still feasible, if properly planned, and urgent. Postponing any management 277 intervention could complicate eradications, both for the spatial spread of the species in one of the 278 two areas and for the risk of change in visitors' attitude due to a greater confidence with chipmunks 279 280 that have become more abundant.

281

282 ACKNOWLEDGMENTS

283 The "U.O. Manutenzione e Valorizzazione del Verde Urbano" office of the Municipality of Roma

284	provided us with permits to set the survey within urban parks in Rome. Authors would like to thank
285	Davide Sogliani, who kindly helped in the fieldwork, and Giulia Benassi, who provided them with
286	data from previous surveys about chipmunks presence in Rome.

287

5 REFERENCES 288

- 1. Mori E , Zozzoli R, Menchetti M, Global distribution and status of introduced Siberian 289 chipmunks Eutamias sibiricus. Mammal Rev 48:139-152 (2018). 290
- 2. Bertolino S, Introduction of the American grey squirrel (Sciurus carolinensis) in Europe: a 291 292 case study in biological invasion. Curr Sci India 95:903-906 (2008).
- 3. Pisanu B, Marsot M, Marmet J, Chapuis JL, Reale D, Vourc'h G, Introduced Siberian 293 chipmunks are more heavily infested by ixodid ticks than are native bank voles in a 294 295 suburban forest in France. Int J Parasitol 40:1277-1283 (2010).
- 4. Marsot M, Chapuis JL, Gasqui P, Dozières A, Massèglia S, Pisanu B, Ferquel E, Vourc'h G, 296 Introduced Siberian chipmunks (Tamias sibiricus barberi) contribute more to Lyme 297 borreliosis risk than native reservoir rodents. PLoS One 8: e55377 (2013). 298
- 5. Benassi G, Bertolino S, Distribution and activity of the introduced Siberian chipmunk 299 (Laxmann 1769) in an urban park in Rome, Italy. Mammalia 75:87–90 (2011). 300
- 6. Zozzoli R, Menchetti M, Mori E, Spatial behaviour of an overlooked alien squirrel: the case 301 of Siberian chipmunks Eutamias sibiricus. Behav Process 153:107-111 (2018). 302
- 7. Pluess T, Cannon R, Jarošík V, Pergl J, Pyšek P, Bacher S, When are eradication campaigns 303 successful? A test of common assumptions. Biol Invasions 14:1365-1378 (2012). 304
- 8. Bomford M, O'Brien P, Eradication or control for vertebrate pests? Wildlife Soc B 23:249-305 306 255 (1995).
- 9. Robertson PA, Adriaens T, Lambin X, Mill A, Roy S, Shuttleworth CM, Sutton-Croft M, 307 The large-scale removal of mammalian invasive alien species in Northern Europe. Pest 308 Manag Sci 73:273-279 (2016). 309

310	10. McNeely JA, The great reshuffling: human dimensions of invasive alien species The great
311	reshuffling: human dimensions of invasive alien species. Workshop on the Human
312	Dimensions of Invasive Alien Species, Cape Town, ZA, 15-17 September 2000: 213-237
313	(2001).

- 314 11. Blackburn TM, Pettorelli N, Katzner T, Gompper ME, Mock K, Garner TWJ, Altwegg R,
 315 Redpath S, Gordon IJ , Dying for conservation: eradicating invasive alien species in the face
 316 of opposition. *Anim Conserv* 13:227-228 (2010).
- 317 12. Shine R, Doody JS, Invasive species control: understanding conflicts between researchers
 318 and the general community. *Front Ecol Environ* **9**: 400-406 (2011).
- Bertolino S, Genovesi P, Spread and attempted eradication of the grey squirrel (*Sciurus carolinensis*) in Italy, and consequences for the red squirrel (*Sciurus vulgaris*) in Eurasia.
 Biol Conserv 109: 351-358 (2003).
- 14. Crowley SL, Hinchliffe S, McDonald RA, Conflict in invasive species management. *Front Ecol Environ* 15: 133-141 (2017).
- 15. Crowley SL, Hinchliffe S, McDonald RA, Invasive species management will benefit from
 social impact assessment. *J Appl Ecol* 54: 351-357 (2017).
- 16. Gálvez N, Guillera-Arroita G, St. John FA, Schüttler E, Macdonald DW, Davies ZG, A
 spatially integrated framework for assessing socioecological drivers of carnivore decline. *J Appl Ecol* 55:1393-1405 (2018).
- Struebig MJ, Linkie M, Deere NJ, Martyr DJ, Millyanawati B, Faulkner SC, Le Comber SC,
 Mangunjaya FM, Leader-Williams N, McKay JE, St. John FA, Addressing human-tiger
 conflict using socio-ecological information on tolerance and risk. *Nat Commun* 9:3455
 (2018).
- 18. Genovesi P, Bertolino S, Human dimension aspects in invasive alien species issues: the case
 of the failure of the grey squirrel eradication project in Italy. In: McNeely JA (Eds) The

- Great Reshuffling: human dimensions of Invasive Alien Species. IUCN, Gland, Switzerland
 and Cambridge, UK, Vi + 242 pp: 113-119 (2001).
- 19. La Morgia V, Paoloni D, Genovesi P, Eradicating the grey squirrel *Sciurus carolinensis* from
 urban areas: an innovative decision-making approach based on lessons learnt in Italy. *Pest Manag Sci* 73:354-363 (2017).
- 20. Mori E, Zozzoli R, Mazza G, Coming in like a wrecking-ball: are native Eurasian red
 squirrels displacing invasive Siberian chipmunks? A study from an urban park. *Urban Ecosyst* 21:975–981 (2018).
- Mori E, Milanesi P, Menchetti M, Zozzoli R, Monaco A, Capizzi D, Nerva L, Genetics
 reveals that free-ranging chipmunks introduced to Italy have multiple origins. *Hystrix* DOI:
 10.4404/hystrix-00116-2018 (2018).
- 346 22. Mori E, Menchetti M, Zozzoli R, Milanesi P, The importance of taxonomy in species
 347 distribution models at a global scale: the case of an overlooked alien squirrel facing
 348 taxonomic revision. *J Zool* **307**:43-52 (2019).
- 23. Manfredo MJ, Who cares about wildlife? Springer, New York, NY, USA: 1-2 (2008).
- 24. Jacobs MH, Human emotions toward wildlife. *Hum Dimens Wildl* **17**:1-3 (2012).
- 25. Jacobs MH, Vaske JJ, Roemer JM, Toward a mental systems approach to human relationships with wildlife: the role of emotional dispositions. *Hum Dimens Wildl* **17**:4-15 (2012).
- 26. Jacobs MH, Vaske JJ, Dubois S, Fehres P, More than fear: role of emotions in acceptability
 of lethal control of wolves. *Eur J Wildlife Res* 60:589-598 (2014).
- 27. Larson LR, Cooper CB, Hauber ME, Emotions as drivers of wildlife stewardship behavior:
 Examining citizen science nest monitors' responses to invasive house sparrows. *Hum Dimens Wildl* 21:18-33 (2016).
- 28. Frank B, Glikman JA, Sutherland M, Bath AJ, Predictors of extreme negative feelings
 toward coyote in Newfoundland. *Hum Dimens Wildl* 21:297-310 (2016).

- 361 29. Bicchieri C, Norms in the wild: How to diagnose, measure, and change social norms.
 362 Oxford University Press, Oxford, UK (2016).
- 363 30. Dunn TJ, Baguley T, Brunsden V, From alpha to omega: A practical solution to the pervasive
 364 problem of internal consistency estimation. *Brit J Psychol* 105:399-412 (2014).
- 365 31. Kassambara A, Practical guide to cluster analysis in R. CreateSpace: North Charleston,
 366 South Carolina, USA (2017).
- 367 32. Raine NE, Rossmo DK, Le Comber SC, Geographic profiling applied to testing models of
 368 bumble-bee foraging. *J R Soc Interface* 6:307-319 (2008).
- 369 33. Martin RA, Rossmo DK, Hammerschlag N, Hunting patterns and geographic profiling of
 white shark predation. Journal of Zoology (London) 279: 111-118 (2009).
- 34. Faulkner SC, Stevenson MD, Verity R, Mustari AH, Semple S, Tosh DG, Le Comber SC,
 Using geographic profiling to locate elusive nocturnal animals: a case study with spectral
 tarsiers. *J Zool* 295: 261-268 (2015).
- 374 35. Faulkner SC, Verity R, Roberts D, Roy SS, Robertson PA, Stevenson MD, Le Comber SC,
 375 Using geographic profiling to compare the value of sightings vs trap data in a biological
 376 invasion. *Divers Distrib* 23:104-112 (2017).
- 377 36. Verity R, Stevenson MD, Rossmo DK, Nichols RA, Le Comber SC, Spatial targeting of
 378 infectious disease control: identifying multiple, unknown sources. *Methods Ecol Evol* 5:647379 655 (2014).
- 37. Marmet J, Pisanu B, Chapuis JL, Home range, range overlap, and site fidelity of introduced
 Siberian chipmunks in a suburban French forest. *Eur J Wildlife Res* 55:497-504 (2009).
- 38. Marmet J, Pisanu B, Chapuis JL, Natal dispersal of introduced Siberian chipmunks, *Tamias* sibiricus, in a suburban forest. *J Ethol* 29:23-29 (2012).
- 384 39. Davey GC, Macdonald BA, Brierley L, The effect of disgust on anxiety ratings to fear 385 relevant, disgust-relevant and fear-irrelevant stimuli. J Anxiety Disord 22:1347-1354 (2008).

- 40. Curtis V, Aunger R, Rabie T, Evidence that disgust evolved to protect from risk of disease. *P R Soc B-Biol Sci* 271:S131-S133 (2004).
- 388 41. Prokop P, Usak M, Fančovičová J, Risk of parasite transmission influences perceived
 389 vulnerability to disease and perceived danger of disease-relevant animals. *Behav Process* 390 85:52-57 (2010).
- 42. Heberlein TA, Navigating Environmental Attitudes. Oxford University Press, Oxford, UK
 (2012).
- 43. Nilsson NJ, Understanding beliefs. MIT (Massachusetts Institute of Technology) Press,
 Cambridge, Massachusetts, USA (2014).
- 44. Chong D, Druckman JN, Framing theory. *Annu Rev Polit Sci* **10**:103-126 (2007).
- 45. Howe LC, Krosnick JA, Attitude strength. *Annu Rev Psychol* **68**:327-351 (2017).
- 46. Rucker DD, Petty RE, Briñol P, What's in a frame anyway? A meta-cognitive analysis of the
 impact of one versus two sided message framing on attitude certainty. J Consum Psychol
 18:137-149 (2008).
- 400 47. Sharp RL, Larson LR, Green GT, Factors influencing public preferences for invasive alien
 401 species management. *Biol Conserv* 144:2097-2104 (2011).
- 402 48. Dunn M, Marzano M, Forster J, Gill RM, Public attitudes towards "pest" management:
 403 Perceptions on squirrel management strategies in the UK. *Biol Conserv* 222:52-63 (2018).
- 404 49. Di Febbraro M, Martinoli A, Russo D, Preatoni D, Bertolino S, Modelling the effects of
 405 climate change on the risk of invasion by alien squirrels. *Hystrix* 27:22–29 (2016).
- 50. Di Febbraro M, Menchetti M, Russo D, Ancillotto L, Aloise G, Roscioni F, Preatoni DG,
 Loy A, Martinoli A, Bertolino S, Mori E, Integrating climate and land–use change scenarios
 in modelling the future spread of invasive squirrels in Italy. *Divers Distrib*, DOI:
 10.1111/ddi.12890 (2019).

410 Figure legends

- 411
- 412 Figure 1. Sankey plot about visitors-chipmunks interactions.

413

414 Figure 2. Dendrogram of the hierarchical cluster analysis.

415

- 416 Figure 3. Marginal effects of the GLM, based on a Gamma distribution of the error and an identity
- 417 link: average scores of normative expectations in the two areas, both for respondents who had

418 observed chipmunks (blue) and for respondents who didn't (red).

419

- 420 Figure 4. Location of the three study sites (left), heatmap with the posterior probabilities of the
- 421 origins of observed tamias (center) and posterior probability of the dispersal parameter (right).

422 TABLES

423 Table 1. Questionnaire structure, item summary and completion rate.

Construct	Format	Question	Mean ± SD
Belief strength	Bipolar scale, from "Strongly disagree" (-3) to "Strongly agree" (+3)	Nowadays the presence of chipmunks in an urban park would"	0.55 ± 1.57
		improve its overall environmental quality.	
		make it more aesthetically appealing.	1.14 ± 1.50
		lead people who already visited the park to return there more often.	1.35 ± 1.34
		attract new visitors.	1.43 ± 1.31
		make visiting the park a more satisfying experience.	1.69 ± 1.39
		cause disease transmission to visitors.	-1.18 ± 1.64
		cause disease transmission to the dogs which are walked in the park.	-1.04 ± 1.48
		cause damaging to the vegetation of the park.	-1.04 ± 1.57
		cause nest depredation on those birds that breed there.	-0.61 ± 1.68
		affect garbage disposal, as chipmunks could rummage it around while searching for food.	-0.88 ± 1.58
Evaluation of each belief	Unipolar scale from with 7 degrees, from "Bad" to "Good"	"How would you evaluate the following consequences of chipmunks presence in the park?"	1.73 ± 1.86
		The park improves its environmental quality.	
		The park becomes more aesthetically appealing.	2.02 ± 1.40
		People who already visited the park, now return there	1.65 ± 1.48

		more often.	
		New people come to see the park.	1.93 ± 1.30
		Visiting the park becomes a more satisfying experience.	1.76 ± 1.41
		Chipmunks transmit diseases to visitors.	-2.20 ± 1.22
		Chipmunks transmit diseases to the dogs which are walked in the park.	-2.08 ± 1.18
		The vegetation of the park is damaged by chipmunks.	-1.71 ± 1.43
		Chipmunks prey on the nests of the birds that breed in the park.	-1.67 ± 1.34
		Chipmunks rummage the garbage around garbage bins, while searching for food.	-1.60 ± 1.19
Core affect	Bipolar scale, from "Extremely negative" (-3) to "Extremely positive " (+3)	"Use the following scale to describe how you would feel about living each one of these experience".	0.97 ± 0.18
		Spotting one, or more, chipmunks, while strolling in the park.	
Emotional disposition (Happiness)	Bipolar scale, from "Extremely negative" (-3) to "Extremely positive " (+3)	"Could you please tell us to what extend you agree or disagree with the following statements?"	1.51 ± 1.48
		I would feel happy, if I spotted some chipmunks when I go to the park. (Happiness)	
		I would feel scared, if I spotted some chipmunks when I go to the park. (Fear)	-2.55 ± 1.11
		I would be surprised, if I spotted some chipmunks when I go to the park. (Surprise)	0.24 ± 1.93
		I would feel disgusted, if I spotted some chipmunks when I go to the park. (Disgust)	-2.46 ± 1.18
		I would feel interested, if I spotted some chipmunks	1.48 ± 1.54

		when I go to the park. (Interest)	
Existence beliefs	Bipolar scale, from "Extremely negative" (-3) to "Extremely positive " (+3)	"To what extend do you agree with the following statements?"	1.25 ± 1.83
		It is important to maintain the presence of chipmunks in the park, for future generations.	
		Whether or not I would get to see chipmunks in the park, it is important to me that they live there.	1.12 ± 1.74
Moral beliefs	Bipolar scale, from "Extremely negative" (-3) to "Extremely positive " (+3)	"To what extend do you agree with the following statements?"	1.28 ± 1.60
		Chipmunks should not be allowed to live in an urban green space, like a park.	
Empirical expectations	Bipolar scale, from "Extremely negative" (-3) to "Extremely positive " (+3)	"To what extend do you agree with the following statements?"	0.94 ± 1.50
		I expect that most chipmunks living in Italy inhabit urban parks; (empirical expectations).	
Normative expectations	Bipolar scale, from "Extremely negative" (-3) to "Extremely positive " (+3)	"To what extend do you agree with the following statements?"	1.13 ± 1.37
		I expect most people to believe that chipmunks should be allowed to lived in an urban green space, like the park; (normative expectations).	
Perceived willingness to report chipmunks presence to local authorities (other people)	Dichotomous (Yes/No)	I expect most people to report chipmunks to local authorities, if they found they were living in an urban park.	Yes = 4.03%, No = 95.67%
Personal comittment to report chipmunks presence to local	Dichotomous (Yes/No)	I would report chipmunks to local authorities, if they found they were living in an urban park.	Yes = 9.27%, No = 90.73%

authorities			
Awareness about he presence of chipmunks	Dichotomous (Yes/No)	Prior to this survey, have you ever heard that some chipmunks inhabited the park?	Yes = 41.12%, No = 58.87%
Experience with chipmunks	Ordered scale (Multiple times per week; Multiple times per month; Once; Never)	"Could you please tell us how often did you have the following experiences at the park?" Spotting one, or more, chipmunks at the park.	Never = 50.40% Once = 27.82% Multiple times per month = 16.12% Multiple times per week = 5.64%
		Feeding the chipmunks at the park.	Never = 85.48% Once = 10.09% Multiple times per month = 2.82% Multiple times per week = 1.61%
		Touching, or caressing, a chipmunks at the park.	Never = 94.76% Once = 4.03% Multiple times per month = 0.40% Multiple times per week = 0.80%
Park usage	Dichotomous (Yes/No)	"Could you please tell us which of the following activities you usually practice at the park?" Sport (e.g jogging, cycling, football, volley).	Yes = 0.28, No = 0.72
		Birdwatching.	Yes = 6.85, No = 93.15%
		Strolling.	Yes = 79.43%, No = 20.57%
		Picnicking.	Yes = 22.98%, No = 77.02%
		Walking the dog.	Yes = 24.19%, No = 75.81%
		Painting / photography.	Yes = 25.80%, No = 74.19%
		Playing a musical instrument.	Yes = 4.43%, No = 93.57%
		Feeding animals.	Yes = 6.04%, No = 93.96%
		Working or volunteering.	Yes = 2.82%, No = 97.18%
Age	Ordered scale (15-20 years, 21-30 years, 31-40 years, 41-50 years, 51-65 years, more than 65 years)	Could you please tell us in which year were you born?	15-20 years = 10.5%, 21-30 years = 29.44 % 31-40 years = 22.98 %

			41-50 years = 19.75 % 51-65 years = 10.88 % more than 65 years = 6.5%
Sex	Dichotomous (Male/Female)	We know that sex and gender are not the same concept. Could please just tell us your sex?	Male = 50.81% Female = 49.29%
Education	Ordered scale	Which is your level of education?	Elementary school = 1.61% Secondary shool = 8.47 % High school = 41. 94 % University degree = 47.98%

424 Table 2. Confirmatory Factor Analysis, model comparison.

Beliefs about chipmunks' impact over the recreational quality of the park.						
	Full model	Model without the items about the "Environmental quality"	Model without the items about the "Environmental quality" and the "Visitors' satisfaction"			
SRMR	0.04213852	0.03863779	0.03520523			
RMR	0.04196861	0.03848199	0.03506327			
TLI	0.9477260	0.9502502	0.9739350			
CFI	0.9663123	0.9733483	0.9913117			
IFI	0.9666189	0.9735970	0.9914187			
ECVI	0.5632614	0.3781471	0.1772727			
BIC	5471.809	4381.852	3500.436			
AIC	5380.460	4308.070	3444.221			
BIC2	5389.389	4315.282	3449.716			
GFI	0.9341394	0.9495378	0.9838887			
AGFI	0.8750919	0.8788907	0.9323327			
PGFI	0.4925462	0.3956407	0.2342592			
MFI	0.8884080	0.9285285	0.9860585			
Scaled chi-squared difference test	df = 5, AIC = 3444.2, BIC = 3500.4, chisq = 11.964	df = 15, AIC = 4308.1, BIC = 4381.9, chisq = 51.781, chisq.diff = 28.519, df.diff = 10, p- value = 0.00149	df = 29, AIC = 5380.5, BIC = 5471.8, chisq = 87.689, chisq.diff = 24.250, df.diff = 14, p- value = 0.04274			
Cronbach's alpha	0.93	0.92	0.88			
McDonald's Omega	Omega Hierarchical = 0.74, Omega H asymptotic = 0.78, Omega Total = 0.95	Omega Hierarchical = 0.76, Omega H asymptotic = 0.81, Omega Total = 0.95	Omega Hierarchical = 0.68, Omega H asymptotic = 0.73, Omega Total = 0.93			
Beliefs about chipmun	ks' impact over the hygien	e of the park.				
	Full model	Model without the items about the "Vegetation damaging"	Model without the items about the "Nest depredation"			
SRMR	0.06205949	0.05537003	0.03495205			
RMR	0.06180925	0.05514676	0.03481111			
TLI	0.8835495	0.9502502	0.9742254			
CFI	0.9249541	0.9733483	0.9914085			
IFI	0.9265352	0.9735970	0.9916257			

ECVI	0.5515900	0.3781471	0.1625515
BIC	6467.589	4381.852	3912.691
AIC	6376.239	4308.070	3856.477
BIC2	6385.168	4315.282	3861.971
GFI	0.9386140	0.9495378	0.9888453
AGFI	0.9386140	0.9495378	0.9888453
PGFI	0.4949056	0.3956407	0.2354394
MFI	0.8936076	0.9285285	0.9933433
Scaled chi-squared difference test	df = 5, AIC = 3856.5, BIC = 3912.7, chisq = 8.3128	df = 15, AIC = 5146.0, BIC = 5219.8, chisq = 32.8162 , chisq.diff = 16.867, df.diff = 10 , p- value = 0.077360	df = 29, AIC = 6376.2, BIC = 6467.6, chisq = 84.7943, chisq.diff = 37.521, df.diff = 14, p- value = 0.000615
Cronbach's alpha	0.79	0.74	0.74
McDonald's Omega	Omega Hierarchical = 0.4, Omega H asymptotic = 0.47, Omega Total = 0.85	Omega Hierarchical = 0.42, Omega H asymptotic = 0.51, Omega Total = 0.82	Omega Hierarchical = 0.43, Omega H asymptotic = 0.52, Omega Total = 0.83

425 Table 3. Confirmatory Factor Analysis: factor loadings, covariances, variances and communalities.

Beliefs about chipmunks' impact over the recreational quality of the park.


		Estimate	S.E	z-value	p-value
Strength of the belief	Aesthetic quality	0.697	0.076	9.188	0.000
	Increased visits in the near future	0.809	0.076	10.589	0.000
	Increase in the number of visitors	0.799	0.081	9.850	0.000
Evaluation of the outcomes	Aesthetic quality	0.874	0.084	10.409	0.000
	Increased visits in the near future	0.840	0.076	10.999	0.000
	Increase in the number of visitors	0.840	0.093	9.036	0.000
Covariances betw	een the strength o	of each belief	and the evalua	ation of its outcor	ne
	Latent constructs (Strength x Evaluation)	0.684	0.070	9.807	0.000
	Aesthetic quality	0.017	0.041	0.408	0.683
	Increased visits in the near future	0.093	0.041	2.300	0.021
	Increase in the number of visitors	0.036	0.040	0.893	0.372
Variances					
Strength of the belief	Aesthetic quality	0.510	0.086	5.951	0.000
	Increased visits in the near future	0.340	0.073	4.625	0.000
	Increase in the number of visitors	0.357	0.068	5.250	0.000
Evaluation of the	Aesthetic	0.227	0.074	3.058	0.002


outcomes	quality				
	Increased visits in the near future	0.298	0.059	5.037	0.000
	Increase in the number of visitors	0.296	0.067	4.410	0.000
Communalities					
Strength of the belief: aesthetic quality = 0.4879791	Strength of the belief: increased visits in the near future = 0.6580856	Strength of the belief: increase in the number of visitors = 0.6409658		Evaluation of the outcomes: increased visits in the near future = 0.7031584	Evaluation of the outcomes: increase in the number of visitors = 0.7043100
Beliefs about chip	munks' impact o	ver the hygiene	of the park.		
Factor loadings					
		Estimate	S.E	z-value	p-value
Strength of the belief	Disease transmission to visitors	0.878	0.067	13.016	0.000
	Disease transmission to visitors' dogs	0.771	0.062	12.407	0.000
	Garbage rummaging	0.540	0.070	7.672	0.000
Evaluation of the outcomes	Disease transmission to visitors	0.678	0.089	7.664	0.000
	Disease transmission to visitors' dogs	0.865	0.096	9.025	0.000
	Garbage rummaging	0.438	0.096	4.543	0.000
Covariances between the strength of each belief and the evaluation of its outcome					
	Latent constructs (Strength x Evaluation)	0.335	0.084	3.992	0.000
	Disease	0.039	0.042	0.929	0.353

	transmission to visitors				
	Disease transmission to visitors' dogs	-0.025	0.042	-0.596	0.551
	Garbage rummaging	0.057	0.067	0.849	0.396
Variances					
Strength of the belief	Disease transmission to visitors	0.228	0.089	2.553	0.011
	Disease transmission to visitors' dogs	0.404	0.076	5.308	0.000
	Garbage rummaging	0.709	0.077	9.175	0.000
Evaluation of the outcomes	Disease transmission to visitors	0.532	0.153	3.479	0.001
	Disease transmission to visitors' dogs	0.243	0.130	1.870	0.061
	Garbage rummaging	0.802	0.109	7.385	0.000
Communalities					
Strength of the belief: disease transmission to visitors = 0.7721118	Strength of the belief: disease transmission to visitors' dogs = 0.5953521	Strength of the belief: garbage rummaging = 0.2915509	the outcomes: disease	Evaluation of the outcomes: disease transmission to visitors' dogs = 0.7546028	Evaluation of the outcomes: garbage rummaging = 0.1927527

Table 4. Ouput of the Generalized Linear Model. Normative expectations were measured on a rescaled biploar Likert scales, ranging from 1 to 7.

Response variable: normative expectations about chipmunks presence.							
Predictor	Estimate	S.E.	t-value	p-value			
Intercept	5.57407	0.19862	28.063	<2e-16 ***			
Location	-0.64550	0.25149	-2.567	0.0109 *			
Previous experience with chipmunks	0.04131	0.30804	0.134	0.8934			
Location * Previous experience	-0.16988	0.37050	-0.459	0.6470			
Null deviance = 24.126		Residual devi	Residual deviance = 22.937				






Cluster dendrogram with AU/BP values (%)

Distance: euclidean Cluster method: complete





Predicted values for Normative expectations





latitude

