Predator-prey nutrient competition undermines predator coexistence

Toni Klauschies*,[1] and Ursula Gaedke[1]

January 30, 2019

* Corresponding author, e-mail: toni.klauschies@uni-potsdam.de

1. University of Potsdam, Germany

Abstract

Contemporary theory of predator coexistence through relative non-linearity in their functional responses strongly relies on the Rosenzweig-MacArthur equations (1963) in which the (autotrophic) prey exhibits logistic growth in the absence of the predators. This implies that the prey is limited by a resource which availability is independent of the predators. This assumption does not hold under nutrient limitation where both prey and predators bind resources such as nitrogen or phosphorus in their biomass. Furthermore, the prey's resource uptake-rate is assumed to be linear and the predator-prey system is considered to be closed. All these assumptions are unrealistic for many natural systems. Here, we show that predator coexistence on a single prey is strongly hampered when the prey and predators indirectly compete for the limiting resource in a flow-through system. In contrast, a non-linear resource uptake rate of the prey slightly promotes predator coexistence. Our study highlights that predator coexistence does not only depend on differences in the curvature of their functional responses but also on the type of resource constraining the growth of their prey. This has far-reaching consequences for the relative importance of fluctuation-dependent and -independent mechanisms of species coexistence in natural systems where autotrophs experience light or nutrient limitation.

Key words: Coexistence, Competition, Fluctuations, Consumer-resource oscillations, Predator-prey cycles, Relative non-linearity between functional responses

1. Introduction

Since the seminal work of Lotka (1925), Volterra (1928), Gause (1932) and Hutchinson (1961) ecologists are wondering about the mechanisms that underlie the astonishing diversity observed in natural systems. Contemporary theory distinguishes two broad categories of mechanisms that promote species coexistence. Fluctuation-independent mechanisms such as resource partitioning or specialist predation may allow different species to coexist under constant environmental conditions (Chesson 2000; Chase and Leibold 2003). This contrasts with fluctuation-dependent mechanisms of species coexistence that strongly rely on temporal variation in the environmental conditions (Abrams and Holt 2002; Chesson 2003). For example, species may be able to coexist due to species-specific differences in the demographic responses to exogenously driven fluctuations in the abiotic environment (Chesson and Warner 1981; Ellner et al. 2016) or because of endogenously generated oscillations in species abundances (Armstrong and McGehee 1980; Yamamichi et al. 2011) or traits (Abrams 2006; Klauschies et al. 2016). In this study, we investigate how the type of resource, limiting prey growth influences coexistence of two predators on a single fluctuating prey through relative non-linearity in their functional responses, as it represents an important food web module in more complex and diverse communities.

Previous work demonstrated that two predators can coexist on a single prey for a broad range of parameters when the system exhibits pronounced predator-prey oscillations and when the predators show substantial differences in the curvature of their functional responses (Koch 1974; Hsu et al. 1978; Armstrong and McGehee 1980; Abrams and Holt 2002). The predator with the more linear functional response is the better competitor when the system exhibits strong predator-prey oscillations but promotes less variation in the species' abundances when it reaches high densities. In contrast, the predator with the more non-linear functional response is competitive superior at steady state dynamics while favoring larger fluctuations in the species abundances when it becomes dominant (Abrams 2004; Mittelbach 2012; Xiao and Fussmann 2013). Hence, the dominant predator alters the environment in a way that is beneficial for the rare predator, allowing its regrowth from low densities (Figure 1A). This negative frequency-dependent selection may enable stable coexistence of the two different predators. In line with the gleaner-opportunist trade-off (Grover 1997; Litchman and Klausmeier 2001), one predator (the gleaner) usually exhibits a higher net-growth growth rate at lower prey densities whereas the other predator (the opportunist) benefits from a higher net-growth rate at higher prey densities.

Most theoretical investigations in this field rest on the classical equations of Rosenzweig and MacArthur (1963) where the prey grows logistically in the absence of the two predators (e.g. Hsu et al. 1978; Xiao and Fussmann 2013). This implies three very strong assumptions about the type, the uptake and the replenishment of the resource that is limiting the growth of the prey, which may or may not hold in nature.

First of all, logistic growth entails that the predators will not directly affect the amount of resources freely available for the uptake by the prey. This might be reasonable for resources such as light or space, which are hardly influenced by predators. However, when mineral nutrients such as nitrogen or phosphorus are limiting autotrophic or bacterial growth, the resource is often substantially stored within the biomasses of both the prey and the predators as particulate organic nutrients. As a result, the predators affect the prey not only by consumption but also by emergent competition for nutrients (Figure 1B). Given that herbivores have much higher nutrient to carbon ratios (N/C ratios) than nutrient depleted plants (Sterner and Elser 2002; Andersen 2004; Hessen et al. 2013), nutrient competition between predators and prey may be substantial even when the total biomass of the predators is relatively low.

Second, under logistic growth, the uptake rate of the prey for the resource is assumed to be linear (Kooi et al. 1998). This assumption lacks empirical evidence (Eppley et al. 1969; Sibly and Hone 2002) and is not supported by mechanistic considerations (Dugdale 1967; Droop 2003). Rather, in line with classical enzyme kinetics (Michaelis and Menten 1913) and functional response theory (Holling 1961), it typically depends non-linearly on the external resource concentration. A linear uptake rate is only to be expected at very low resource concentrations, e.g. when the prey is strongly bottom-up controlled. In contrast, at higher resource concentrations, the uptake rate of the prey is likely to be saturating.

Finally, the Rosenzweig-MacArthur predator-prey model basically describes a closed autochthonous system with a given constant amount of resources, the carrying capacity K, for the prey. In this case, nutrients are thought to be internally replenished. However, most natural systems receive allochthonous nutrient inputs and experience losses of dissolved and particulate nutrients, i.e. they are more realistically described as flow-through systems.

Hence, we investigate to what extent previous results on the coexistence of two predators sharing a single prey depend on the above mentioned simplifying assumptions. We show that coexistence most likely occurs in autochthonous systems where the predators do not interfere with the resource availability of the prey, e.g. under light or space limitation, and when the preys' uptake rate for the resource is strongly non-linear. Therefore, predator coexistence does not only depend on differences in the non-linearity of their functional responses but also

on the type of resource constraining the growth of their prey. This is crucial for our understanding of species coexistence in natural systems because we can expect that fluctuation-dependent mechanisms of species coexistence are more relevant in light or space limited systems as compared to nutrient limited systems.

Figure 1: *Predator coexistence through relative non-linearity in their functional responses and associated fluctuations in species abundances. (A) Negative frequency-dependent selection: the dominant predator (in-transparent) alters the environment in a way that is beneficial for the rare predator (transparent). A dominance of the predator with a type I functional response promotes stasis in the predator-prey dynamics although it is the better competitor under fluctuating environmental conditions. In contrast, a dominance of the predator with a type II functional response promotes predator-prey oscillations while being competitive superior under constant environmental conditions. (B) The type of resource limitation influences predator-prey interactions beyond consumptive effects of the predators on the prey. Predators mainly affect the prey by consumption under light or energy limitation whereas predators also compete with the prey for shared resources under nutrient limitation of the prey. (C) The predator-prey model defined by equation [1] in the main text can be derived from a generic predator-prey model that explicitly accounts for the replenishment, uptake and recycling of nutrients (for the derivation see Appendix A; for a detailed model description see also methods in the main text).*

2. Methods

2.1. Description of the predator-prey model

We consider a predator-prey system that comprises two different predators and one prey. This model is derived from a more general predator-prey system that explicitly accounts for the uptake and replenishment of inorganic mineral nutrients in a flow-through system (for details see Appendix A). Hence, the biomasses of the prey R and the two predators C_1 and C_2 are changing over time according to the following set of equations:

$$\frac{dR}{dt} = \left[\frac{a_R\left(N_I - \left(\eta_R R + \eta_{C_1} C_1 + \eta_{C_2} C_2\right)\right)}{1 + h_R a_R\left(N_I - \left(\eta_R R + \eta_{C_1} C_1 + \eta_{C_2} C_2\right)\right)} - e_{C_1} a_{C_1} C_1 - \frac{e_{C_2} a_{C_2} C_2}{1 + h_{C_2} a_{C_2} R} - \delta \right] R \qquad [1a]$$

$$\frac{dC_1}{dt} = \left[e_{C_1} a_{C_1} R - d_{C_1} - \delta \right] C_1 \qquad [1b]$$

$$\frac{dC_2}{dt} = \left[\frac{e_{C_2} a_{C_2} R}{1 + h_{C_2} a_{C_2} R} - d_{C_2} - \delta \right] C_2 \qquad [1c]$$

Nutrient-uptake of the prey is modelled by using a type II functional response with uptake rate a_R and processing time h_R (Michaelis & Menten 1913; Holling, 1961). The concentration of dissolved inorganic nutrients which are freely available to the prey, is given by $N = N_I - \left(\eta_R R + \eta_{C_1} C_1 + \eta_{C_2} C_2\right)$ where N_I denotes the total amount of dissolved inorganic and particulate organic nutrients in the system. The parameters η_R and η_{C_i} represent the carbon to nutrient ratios of the prey and predator species, respectively, and correspond to species-specific competition coefficients for the total amount of resources in Lotka-Volterra competition models (cf. MacArthur and Levins 1967; Schoener 1974). The term $\left(\eta_R R + \eta_{C_1} C_1 + \eta_{C_2} C_2\right)$ accounts for the fact that the predators contain not only carbon but also other nutrients in their biomasses. In this bounded state, the nutrients are not available for prey uptake. Hence, predators may not only feed on the prey but also diminish the nutrients available for their own prey and thus compete with the prey for shared nutrients. The overall relevance of this process for the predator-prey interaction will generally depend on the carbon to nutrient ratios of the different species and on the extent to what mineral nutrients are the primarily limiting factor of the prey growth. As detailed in Appendix A, the terms in eq. [1] reflecting nutrient competition between predators and prey emerge during the derivation of our predator-prey model from a corresponding chemostat model that also explicitly describes the nutrient dynamics (Figure 1C). Hence, they are based on a mechanistic description of the resource uptake by the prey.

Previous studies have shown that two predators are most likely to coexist on a single prey through endogenously generated predator-prey oscillations when they exhibit substantial differences in their functional responses (Hsu et al. 1978; Chesson 2000; Xiao and Fussmann

2013). Hence, without loss of generality, we assumed the grazing rate of C_1 to follow a linear, type I, functional response with attack rate a_{C_1}. In contrast, the grazing rate of C_2 was assumed to follow a type II functional response with attack rate a_{C_2} and handling time h_{C_2} (Armstrong and McGehee 1980; Abrams and Holt 2002). The ingested prey biomass is converted into predator biomass with species-specific transfer efficiencies e_{C_i}. In addition, all species are washed out from the system with the dilution rate δ, representing a shared loss rate. To account also for senility or mortality of the individual predators, we introduced the species-specific death rates d_{C_i}. We did not incorporate natural mortality of the prey through senescence assuming it to be negligible compared to the overall losses through predation and dilution, which facilitates comparison with previous results.

When assuming the absence of nutrient competition among predators and prey $(\eta_{C_i} = 0)$, a linear nutrient uptake rate of the prey ($h_R = 0$), and a closed predator-prey system (batch conditions where $\delta = 0$), our system described by eq. [1] simplifies to the predator-prey system studied extensively in previous work (e.g. Armstrong and McGehee 1980; Abrams and Holt 2002; Xiao and Fussmann 2013). In this case, the intrinsic growth rate (r) and the carrying capacity (K) of the prey are given by $r = a_R N_I$ and $K = N_I$. Hence, the classical predator-prey model based on the Rosenzweig-MacArthur equations (Rosenzweig and MacArthur 1963), is a special case of our general predator-prey model described by eq. [1]. This allows us to directly compare our results to previous findings.

2.2. Rescaled predator-prey model

We simplified the analysis by rescaling the time, state variables and parameters of our predator-prey model as follows: $t' = t/(a_R N_I)$, $R' = \eta_R R/N_I$, $C_i' = \eta_R C_i/(e_{C_i} N_I)$, $h_R' = (h_R a_R N_I)$, $a_{C_i}' = \eta_R e_{C_i} a_{C_i}/a_R$, $d_{C_i}' = d_{C_i}/(a_R N_I)$, $\delta' = \delta/(a_R N_I)$, $h_{C_1}' = h_{C_1} a_R N_I/e_{C_1}$ and $\eta_{C_i}' = \eta_{C_i} e_{C_i}/\eta_R$. Substituting the rescaled state variables and effective parameters into eq. [1] and dropping the primes, we obtain the following non-dimensionalized predator-prey model:

$$\frac{dR}{dt} = \left[\frac{\left(1-(R+\eta_{C_1}C_1+\eta_{C_2}C_2)\right)}{1+h_R\left(1-(R+\eta_{C_1}C_1+\eta_{C_2}C_2)\right)} - a_{C_1}C_1 - \frac{a_{C_2}C_2}{1+h_{C_2}a_{C_2}R} - \delta \right] R \qquad [2a]$$

$$\frac{dC_1}{dt} = \left[a_{C_1}R - d_{C_1} - \delta \right] C_1 \qquad [2b]$$

$$\frac{dC_2}{dt} = \left[\frac{a_{C_2}R}{1+h_{C_2}a_{C_2}R} - d_{C_2} - \delta \right] C_2 \qquad [2c]$$

Hence, the overall population dynamics of the different species depend on the curvature of the functional responses, the death rates and the competition coefficients for nutrients of the two

different predators, the non-linearity of the nutrient uptake rate of the resource and the shared loss rate for all species. In contrast, the total amount of nutrients in the system, the transfer efficiencies of the two predators, the resource uptake rate of the prey and the competition coefficient of the prey for the total amount of nutrients can be omitted for further analysis.

2.3. Model analysis

We combined analytical and numerical approaches to achieve a clear and comprehensive understanding of the impact of nutrient competition among the predator and prey species and of a non-linear nutrient uptake rate of the prey on the likelihood of two predators to coexist on a single prey in a batch or a flow-through (chemostat) system.

The potential for two predators to coexist on a single prey mainly depends on the death rates of the two predators when their functional responses are fixed (Abrams and Holt 2002). Following Abrams and Holt (2002) we thus evaluated predator coexistence based on the region in the two-dimensional $d_{C_1} - d_{C_2}$ parameter space that allows for predator coexistence. In line with Hsu et al. (1978) and Xiao and Fussmann (2013), we first derived the necessary conditions for species coexistence that define a range of parameters that may permit predator coexistence (for details see section 2.3.1 and Appendix B). Within this parameter space, we used numerical simulations to obtain also the sufficient conditions that define the range of parameters enabling predator coexistence (for details see section 2.3.2). In accordance with previous studies we first investigated the influence of the curvature of the type II functional response of C_2 on predator coexistence assuming logistic growth of the prey in the absence of the predators, thereby setting the baseline for further analysis. Afterwards, we explored the impact (1) of resource competition among predators and prey $\left(\eta_{C_i} > 0\right)$, (2) of a non-linear resource uptake of the prey $(h_R > 0)$ and (3) of a shared loss rate for all species $(\delta > 0)$ on the likelihood of predator coexistence on a single prey (for details see section 2.3.3).

2.3.1. Parameter range of potential species coexistence

In the following we derive necessary conditions defining a region in the $d_{C_1} - d_{C_2}$ parameter space that may allow for predator coexistence. A basic requirement for predator coexistence is that each predator is able to persist with the prey alone and thus, in the absence of the other predator. This prerequisite gives rise to upper limits in the death rates of the two predators as they are given by $d_{C_1}^U$ for C_1 and $d_{C_2}^U$ for C_2 (for details see Appendix B.1; cf. Figure 2A – upper limits of the x and y axes).

Furthermore, species are generally viewed to stably coexist when they can mutually invade each other's resident community (Chesson 2000). Hence, there has to be negative frequency-dependent selection on the different predators, which allows all species to regrow from low densities. The linear functional response of C_1 ensures that the $R - C_1$ system approaches a stable equilibrium in which the equilibrium density of the prey $(R_{C_1}^*)$ represents the minimum prey requirement of C_1. Therefore, C_2 will be able to invade the $R - C_1$ system when it exhibits a positive per-capita net growth for $R_{C_1}^*$. This implies that the minimum prey requirement of C_2 $(R_{C_2}^*)$ has to be lower than the prey requirement of C_1: $R_{C_2}^* < R_{C_1}^*$.(Appendix B.2; Figure 2A – solid line).

In contrast to the linear functional response of C_1, the non-linear functional response of C_2 may destabilize the interior equilibrium of the $R - C_2$ system, provoking predator-prey oscillations. However, due to C_1's linear functional response, its invasion growth rate for the $R - C_2$ system equals its per-capita net-growth rate for the temporal average of the prey density $\bar{R}_{C_2}$. Hence, C_1 can invade the $R - C_2$ system when its per-capita net-growth rate for $\bar{R}_{C_2}$ is larger than zero and, thus, when its minimum prey requirement is lower than $\bar{R}_{C_2}$, i.e. $R_{C_1}^* < \bar{R}_{C_2}$. Unfortunately, a general expression for $\bar{R}_{C_2}$ does not exist. We can, however, derive an upper limit for $\bar{R}_{C_2}$ $(\bar{R}_{C_2}^U)$ which generally depends on the death rate and the parameters of the functional response of C_2. Hence, C_1 cannot invade the $R - C_2$ system for $R_{C_1}^* > \bar{R}_{C_2}^U$ (Appendix B.3; Figure 2A – dashed line).

In line with the reasoning above, the two different predators are able to coexist on the single prey when the following inequality is satisfied: $R_{C_2}^* < R_{C_1}^* < \bar{R}_{C_2}$ (cf. Figure 2B; Abrams and Holt 2002). Hence, the minimum prey requirement of C_2 has to be lower than the temporal average of the prey density in the $R - C_2$ system, i.e. $R_{C_2}^* < \bar{R}_{C_2}$. This inequality only holds when the equilibrium value $R_{C_2}^*$ is unstable, i.e., the corresponding predator-prey system exhibits oscillations in the species' abundances. Hence, the death rate of C_2 has to be smaller than the threshold value $d_{C_2}^H$, at which the $R - C_2$ system exhibits a Hopf-bifurcation which gives rise to a limit cycle in the corresponding predator-prey dynamics (Figure 2A – dashed-dotted line; for details see Appendix B.4).

Figure 2: *Necessary conditions for the coexistence of two different predators through relative non-linearity in their functional responses and associated fluctuations in the prey abundance R. (A) Parameter range, i.e. death rates d_{C_1} and d_{C_2}, that potentially enables coexistence of predator one, C_1, and predator two, C_2 (P_a; grey shaded area). The scales of d_{C_1} and d_{C_2} are chosen in accordance with the first necessary condition requiring each predator to persist in*

the absence of the other one. In line with the second necessary condition C_2 can invade the $R -$ C_1 system when d_{C_2} is lower than a threshold value that depends on d_{C_1} (grey solid line). The third necessary condition predicts the exclusion of C_1 when d_{C_2} falls below a critical value that is also depending on d_{C_1} (grey dashed line). Finally, according to the fourth necessary condition, N_4, predator coexistence is only possible when the $R - C_2$ system exhibits predator-prey oscillations. For this, d_{C_2} has to be lower than the Hopf-bifurcation point $d_{C_2}^H$ of the $R -$ C_2 system (grey dashed-dotted line). Dots indicate parameter combinations used for the individual simulations shown in (C-F) (for details see methods). (B) Mutual invasibility of the two predators depends on the predators' per-capita net-growth rates. While C_1 is assumed to exhibit a linear functional response (black solid line), C_2 has a non-linear functional response (black dashed-dotted line). C_2 can invade the $R - C_1$ system when its minimum prey requirement ($R_{C_2}^*$, vertical dashed-dotted line) is lower than the minimum prey requirement of C_1 ($R_{C_1}^*$, vertical solid line). In contrast, C_1 can invade the $R - C_2$ system when the abundances of R and C_2 exhibit oscillations and $R_{C_1}^*$ is lower than the temporal average of the corresponding prey density ($\bar{R}_{C_2}$, vertical dashed line). (C-F) Dynamics of R (black solid line), C_1 (grey solid line) and C_2 (grey dashed-dotted line) in dependence of the predators' death rates. For $d_{C_1} = 0.2$ and $d_{C_2} = 0.2$ (dark grey dot in (A)), both predators are able to coexist on a limit cycle (C). For $d_{C_1} = 0.1$ and $d_{C_2} = 0.3$ (black dot in (A)), C_1 outcompetes C_2 (D). In contrast, C_2 excludes C_1 for $d_{C_1} = 0.8$ and $d_{C_2} = 0.4$ (E, stable equilibrium; white dot in (A)) as well as for $d_{C_1} = 0.9$ and $d_{C_2} = 0.3$ (F, predator-prey oscillations; light grey dot in (A)). All simulations are based on the following parameter values: $a_{C_1} = 1$, $a_{C_2} = 4$, $h_{C_2} = 2$, $h_R = 0$, $\eta_{C_i} = 0$, $\delta = 0$.

As detailed in Appendix C, the differences $R_{C_1}^* - R_{C_2}^*$ and $\bar{R}_{C_2} - R_{C_2}^*$ measure the average fitness and niche differences between the two different predators, respectively. Hence, whenever the average fitness differences between the two different predators are smaller than their niche differences, i.e. $R_{C_1}^* - R_{C_2}^* < \bar{R}_{C_2} - R_{C_2}^* \leftrightarrow R_{C_1}^* < \bar{R}_{C_2}$, the predators will be able to coexist on the single prey species under recurrent predator-prey oscillations. Otherwise, one of the two predators will be excluded from the system.

In line with Xiao and Fussmann (2013), the necessary conditions outlined above specify a region in the parameter space in which predator coexistence is possible (cf. Figure 2A – grey shaded area). The absolute size of this parameter region (P_a) quantifies the overall strength of relative non-linearity in the predators' functional responses and associated fluctuations in the prey's abundance to allow for predator coexistence (for details on the quantification of P_a see Appendix B.5). In addition, we calculated a relative size of this parameter region

$\left(P_r = P_a / \left(d_{C_1}^U \cdot d_{C_2}^U\right)\right)$ that scales P_a with respect to the upper limits of the predators' death rates $d_{C_i}^U$. Hence, P_r quantifies the proportion of the parameter space of predator persistence that enables predator coexistence (Figure 2A). Assuming that each predator is able to persist with the prey alone and that the death rates of the different predators are drawn randomly from their feasible ranges, P_r provides a probability that two predators are able to coexist on a single fluctuating prey.

2.3.2. *Parameter range of realized species coexistence*

Since $\bar{R}_{C_2} \leq \bar{R}_{C_2}^U$, the parameter range of potential species coexistence P_a will generally overestimate the parameter region that really allows for coexistence of the two different predators. The magnitude of $\bar{R}_{C_2}^U - \bar{R}_{C_2}$ determines to what extent the range of potential species coexistence differs from the range of realized species coexistence. Hence, we quantified the latter by conducting numerical simulations of our predator-prey model defined by eq. [2].

Following Abrams (2004), we considered the coexistence of the two different predators to be stable when each predator is able to invade the resident system comprising the other predator and the single prey at its long-term equilibrium, combined with persistence of the two predators after invasion. First, we tested for mutual invasibility of the two predators by conducting numerical simulations where either C_1 or C_2 was set to zero at the beginning. The initial density of the resident predator species was set to 0.2 whereas the initial prey density was chosen to equal 0.1. Based on the simulation results of the two different resident systems, we calculated the invasion growth rates, i.e. the time-averaged per-capita net-growth rates, of the two different predators according to the following equations:

$$I_{C_1} = \int_o^T \frac{1}{C_1} \frac{dC_1}{dt} dt = a_{C_1} \bar{R}_{C_2} - d_{C_1} - \delta$$

$$I_{C_2} = \int_o^T \frac{1}{C_2} \frac{dC_2}{dt} dt = \frac{a_{C_2} R_{C_1}^*}{1 + h_{C_2} a_{C_2} R_{C_1}^*} - d_{C_2} - \delta$$

with I_{C_i} being the invasion growth rate of predator i for the corresponding resident system comprising the other predator and the prey. The overall simulation time T was set to 10^4. The two predators can mutually invade each other when both of them exhibit positive invasion growth rates for the corresponding resident systems, i.e. $I_{C_i} > 0$. In contrast, if any $I_{C_i} < 0$, the relevant predator will not be able to recover from low densities, and thus, cannot invade.

We complemented this approach by also testing for persistence of the two predators after invasion, by conducting additional numerical simulations where all species were initially present. Resolving the $d_{C_1} - d_{C_2}$ parameter space with a grid of 50 equally spaced values of

d_{C_1} and d_{C_2}, respectively, resulted in a total amount of simulations (Z_{total}) of $50 \cdot 50 = 2500$. All simulations were initialized with equally abundant predators and prey, i.e. $C_i = R = 0.1$. We determined species coexistence based on the presence or absence of long-term trends in the population dynamics. First, for each predator, we calculated *Pearson's correlation coefficient* (ρ) between the predators' abundance and time using the last 5000 time units of the corresponding simulation runs (cf. Klauschies et al. 2016). We then considered the two predators to persist when both correlation coefficients were non-significant, i.e. $\rho_i \approx 0$. In contrast, when one of the two predators exhibited a significant negative correlation coefficient, it is assumed to experience competitive exclusion.

Finally, we estimated the absolute (O_a) and the relative (O_r) range of realized species coexistence as follows:

$$O_a = \frac{Z_{coexist}}{Z_{total}} \left(d_{C_1}^U \cdot d_{C_2}^U \right)$$

$$O_r = \frac{Z_{coexist}}{Z_{total}}$$

with $Z_{coexist}$ being the number of simulations where the two predators were able to coexist. Simulations and analyses were performed in MATLAB, version 7.13, using solver ode23 for ODEs (The MathWorks Inc., Natick, MA, 2011). We increased the precision of the solver by reducing the absolute and relative tolerances to 10^{-9} and 10^{-11}, respectively, and the maximum step size to 0.1.

3. Results

In this study we investigated the possibility of two predators coexisting on a single fluctuating prey through differences in the curvature of their functional responses. The likelihood for predator coexistence mainly depends on the death rates of the two predators $\left(d_{C_i} \right)$ when their functional responses are locked (Abrams and Holt 2002). Hence, we evaluated the region in the two-dimensional $d_{C_1} - d_{C_2}$ parameter space that may allow for predator coexistence (Figure 2A; grey shaded area). The absolute size of this parameter region (P_a) measures the overall strength of the corresponding coexistence mechanism. In contrast, the relative size of this parameter region $\left(P_r = \frac{P_a}{d_{C_1}^U \cdot d_{C_2}^U} \right)$ scales P_a with respect to $d_{C_1}^U$ and $d_{C_2}^U$, i.e. the upper limits of the predators' death rates and quantifies the proportion of the parameter space of predator persistence that may allow for predator coexistence. It thus represents a probability for predator coexistence when both predators are able to persist in the absence of their competitor. During

the whole study, we assumed that predator one (C_1) exhibits a linear functional response and predator two (C_2) a type II functional response (Figure 2B).

Depending on the predator's death rates, either both predators coexist on the single prey due to endogenously generated predator-prey oscillations (Figure 2C) or one predator is competitive excluded by the other predator (Figure 2D-F). When C_1 is the only remaining predator, the predator-prey dynamics approach a stable equilibrium over time because of its linear functional response (Figure 2D). In contrast, when the predators' death rates select for C_2 instead, the corresponding predator-prey dynamics either show stasis (Figure 2E) or a limit cycle (Figure 2F).

To clearly relate our results to previous studies we first considered the classical predator-prey model where two predators are feeding on a single prey that exhibits logistic growth in the absence of predation (Hsu et al. 1978; Abrams and Holt 2002; Xiao and Fussmann 2013). Hence, we first evaluated how P_a and P_r depend on the feeding traits, i.e. attack rates and handling times, of the different predators. For this, we set the corresponding parameters h_R, η_{C_i}, and δ of eq. [2] all to zero. After having established a baseline for further analysis, we investigated the impact of nutrient competition among the different predator and prey species $\left(\eta_{C_i} > 0\right)$, and of a non-linear nutrient uptake rate of the prey ($h_R > 0$) on P_a and P_r in the presence or absence of a shared loss rate ($\delta \geq 0$) simulating e.g. batch or chemostat conditions.

3.1 Relative non-linearity in the predators' functional responses fosters coexistence

The possibility of two predators to coexist on a single prey species essentially depends on the curvature of the type-II functional response of C_2. It has to be sufficiently non-linear for the ecologically feasible prey densities in order to allow for predator-prey oscillations. Therefore, a necessary requirement for predator coexistence is that the attack rate (a_{C_2}) and the handling time (h_{C_2}) of C_2 are sufficiently high, i.e. $a_{C_2} h_{C_2} > 1$ (for details see Appendix B). In addition, stable predator coexistence strongly relies on differences in their functional responses and thus, on deviations in their attack rates and handling times.

The absolute size (P_a) and the relative size (P_r) of the parameter region that may allow for predator coexistence increase with a_{C_2} while keeping the attack rate of C_1 (a_{C_1}) constant (Figure 3A-B). This happens because higher values of a_{C_2} strongly increase the competitive ability of C_2 compared to C_1 at lower prey densities R, while hardly affecting its competitive ability at higher densities of R, thereby preventing the competitive exclusion of C_2 by C_1. Regarding the handling time h_{C_2}, P_a shows a humped-shape pattern, independent of the exact

value of a_{C_2} (Figure 4A): when starting from low values, an increase in h_{C_2} leads to an increase in P_a up to a value of h_{C_2} where P_a reaches a maximum. Increasing h_{C_2} further causes P_a to decrease again due to the ambivalent effects of higher values of h_{C_2} on the predator-prey dynamics. An increase in h_{C_2} generally promotes predator-prey oscillations and thereby predator coexistence but it also reduces the maximum grazing rate of C_2 and thus, the maximum death rate it can withstand. Consequently, depending on the net-effect of both processes, an increase of h_{C_2} can lead either to an increase or a decrease of P_a. This strongly contrasts with the impact of h_{C_2} on P_r that always increases with higher values of h_{C_2} because P_r is scaled with respect to the maxima of the predators' feasible death rates (Figure 4B).

Comparing P_a with the size of the parameter region where the predators definitely coexist (O_a), for different values of h_{C_2}, further reveals that P_a strongly overestimates O_a when h_{C_2} is small, but is of similar magnitude as O_a when h_{C_2} is sufficiently high. For example, for $h_{C_2} = 1$ $P_a = 0.146$ is more than twice as large as $O_a = 0.062$ (Figure 5A) but for $h_{C_2} = 4$ $P_a = 0.084$ is nearly as large as $O_a = 0.073$ (Figure 5B). This happens because the amplitude of the predator-prey oscillations of the $R - C_2$ system increases much slower as d_{C_2} decreases from its Hopf-bifurcation point $d_{C_2}^H$ when h_{C_2} is small (Figure 4C, D). As a result, the difference between the temporal average of $\bar{R}_{C_2}$ and its equilibrium value $R_{C_2}^*$ is substantially lower for $h_{C_2} = 1$ than for $h_{C_2} = 4$ (Figure 4C, D). Hence, for $h_{C_2} = 1$, i.e. when the non-linearity of the type II functional response of C_2 is rather weak, C_1 cannot take advantage of very long periods of high resource densities, reducing its potential to invade the $R - C_2$ system. Another consequence of this property is that the value of h_{C_2} that maximizes O_a will be generally higher than the value that maximizes P_a.

Finally, when the two predators are able to coexist, the shape of the predator-prey oscillations could look very different, depending on the parameters of the functional responses and the death rates of the two predators (Figure 4E-F). For most parameter combinations, the oscillations are quite regular, typically exhibiting one dominant time scale at which the two predators are almost completely synchronized, i.e. their phase difference is close to zero (for an example see Figure 4E; further details are given in Appendix D). However, the two predators differ in the temporal variation of their abundances. C_1 strongly increases when the prey density is sufficiently high but also strongly decreases when the prey density is falling below its minimum prey requirement promoting a rather high temporal variation (*Coefficient of variation* (CV) = 0.77). In contrast, C_2 increases less in response to high prey densities due to its lower maximum growth rate but also decreases less when the prey density is rather low because of its

lower death rate and thus, higher starvation resistance at food shortage. As a result, the abundance of C_2 often varies less (CV = 0.60). This type of dynamics nicely reflects the gleaner-opportunist trade-off faced by the two predators, where C_1 exhibits a competitive advantage at times of high prey densities whereas C_2 is doing better at periods of low prey densities.

Figure 3: *The absolute size, P_a, (A, C, E, G) and the relative size, P_r, (B, D, F, H) of the parameter range that may allow for predator coexistence in dependence of the attack rate of C_2, a_{C_2} (A, B), the strength of nutrient competition between the predators and the prey, η_{C_i}, (C, D), the non-linearity of the prey's nutrient uptake rate, h_R, (E, F) or the shared loss rate for all species, δ (G, F) for three different values of the handling time of C_2 ($h_{C_2} = 1$, solid line; $h_{C_2} = 2$, dashed-dotted line; $h_{C_2} = 4$, dashed line). Higher values of P_a indicate a larger region in the parameter space that may enable predator coexistence. The attack rate of C_1, a_{C_1}, was set to 1. Increasing a_{C_2} enhances the relative non-linearity between the functional responses of the two different predators. The higher η_{C_i}, the more nutrients are stored in the biomasses of the predators and, thus, the higher is the competitive effect of the predators on the prey. Increasing δ enhances the overall mortality of all species which is most relevant for the persistence of C_2. All results are based on the analytically derived necessary conditions of species coexistence (for details see Methods and Appendix B).*

This pattern strongly contrasts with the dynamics observed for some other parameter combinations, where the shape of the predator-prey oscillations is much more complex, often displaying two different dominant time scales (Figure 4F). This type of dynamics is only observed when the non-linearity of the type II functional response of C_2 is very strong. As a result of temporal niche differentiation between the two predators, the abundances of C_1 and C_2 vary asynchronously on the longer time scale, displaying a phase difference of 0.74 π (for details see Appendix D). In contrast, the predator-prey oscillations on the shorter time scale are only clearly visible in the dynamics of R and C_2, preventing any meaningful quantification of the phase difference between the predators' population dynamics. In line with the core mechanism that allows species coexistence through relative non-linearity in the predators' functional responses, a temporary dominance of C_1 strongly stabilizes the predator-prey oscillations. Under these conditions, C_2 exhibits positive net-growth rates due to its lower prey requirements, allowing its regrowth from low densities. The enhanced grazing pressure by C_2 then causes a decrease in R below the minimum prey requirement of C_1 leading to a strong decrease in its abundance. The associated release from competition, in turn, allows C_2 to establish high abundances. As a consequence of the curved functional response of C_2 and its temporary dominance, the predator-prey dynamics start to exhibit fast predator-prey oscillations. As a result, the corresponding high mean values in R favor the regrowth of C_1 compared to C_2. This induces low abundances of C_2 again, resetting the initial conditions of the competitive cycle.

Figure 4: *Panels A and B show the absolute size, P_a, and the relative size, P_r, of the parameter region that may allow for predator coexistence in dependence of the handling time of C_2, h_{C_2}, for three different values of the attack rate of C_2 ($a_{C_2} = 2$, solid line; $a_{C_2} = 4$, dashed-dotted line; $a_{C_2} = 8$, dashed line). Higher values of P_a indicate a larger region in the parameter space that may enable predator coexistence. Panels C and D show the maximum (solid black line), minimum (dashed-dotted black line), temporal average (dashed grey line) and equilibrium value (solid grey line) of the prey's abundance (R) in dependence of the death rate of C_2 (d_{C_2}) for the $R - C_2$ system ($a_{C_2} = 4$), in case of $h_{C_2} = 1$ (C) and $h_{C_2} = 4$ (D). The solid vertical grey line indicates the Hopf-bifurcation point $(d_{C_2}^H)$ of the corresponding $R - C_2$ system.*

Panels E and F display the population dynamics of R (black solid line), C_1 (grey dashed-dotted line) and C_2 (grey solid line). The shape of the predator-prey oscillations depends on the attack rate and handling time of C_2 and the death rates of the two different predators, ranging from simple predator-prey oscillations (E, $d_{C_1} = 0.2$, $d_{C_2} = 0.2$, $a_{C_1} = 1$, $a_{C_2} = 4$, $h_{C_2} = 2$) to complex population dynamics (F, $d_{C_1} = 0.1$, $d_{C_2} = 0.1$, $a_{C_1} = 1$, $a_{C_2} = 16$, $h_{C_2} = 8$).

3.2 Nutrient competition among predators and prey hampers coexistence

The possibility of the two predators to coexist on the single prey does not only depend on the parameters of the functional responses of the predators but also on the type of resource that is limiting the growth of the prey. Compared to light or space limitation (Figures 3A-B, 5A-B), nutrient limitation of the prey and the emergent nutrient competition between the predator and prey species strongly reduces the parameter region for which the two predators coexisted (Figures 3C-D, 5C-D). This happens because nutrient competition stabilizes the predator-prey dynamics (Appendix B.4). The ecological mechanism behind this stabilizing effect is that predators absorb mineral nutrients in their biomasses which are not further available for the uptake by the prey. Hence, high predator densities not only negatively affect the prey directly by predation, but also indirectly through nutrient competition. Therefore, the carrying capacity of the prey is effectively reduced in the presence of the predators, which is known to stabilize predator-prey dynamics.

The negative impact of nutrient competition on the absolute (P_a) and relative (P_r) sizes of the parameter range that may enable predator coexistence is more severe for lower values of h_{C_2} (Figures 5C-D vs. A-B). For example, P_a decreases by roughly 26% for $h_{C_2} = 1$ (Figure 5A&C) whereas the decrease is only about 13% for $h_{C_2} = 4$ (Figure 5B&D). The negative impact of nutrient competition on predator coexistence is even more pronounced when considering the parameter range of realized predator coexistence (O_a), especially for higher values of h_{C_2}. In fact, nutrient competition between the predators and the prey reduces the corresponding absolute size of the parameter range of realized species coexistence (O_a) by nearly 55% for $h_{C_2} = 1$ (Figure 5A&C) and by 44% for $h_{C_2} = 4$ (Figure 5B&D). Hence, the impact of nutrient competition on the likelihood of predator coexistence is very pronounced for a broad range of parameters.

Figure 5: *The outcome of predator competition in dependence of the death rates d_{C_i} of two different predators C_1 and C_2 assuming logistic growth for the prey (A, B; $h_R = 0$, $\eta_{C_i} = 0$,*

$\delta = 0$), *nutrient competition between predators and prey (C, D; $h_R = 0$, $\eta_{C_i} = 1$, $\delta = 0$), a non-linear nutrient uptake of the prey (E, F; $h_R = 2$, $\eta_{C_i} = 0$, $\delta = 0$) or an overall loss rate for all species (G, H; $h_R = 0$, $\eta_{C_i} = 0$, $\delta = 0.1$). Results are shown for a lower (A, C, E, G; $h_{C_2} = 1$) and a higher (B, D, F, H; $h_{C_2} = 4$) handling time of C_2. The attack rates of the two different predators were set to $a_{C_1} = 1$ and $a_{C_2} = 4$. The parameter region that may potentially enable predator coexistence, P_a, is indicated by the grey shaded area whereas the parameter region for which the two predators do actually coexist, O_a, is marked by the white area. The black area indicates parameter combinations for which one of the two predators goes extinct (for further details see the caption of Figure 1).*

3.3 Non-linear resource uptake rates of the prey promote predator coexistence

Increasing the non-linearity of the resource uptake rate of the prey enhances slightly the likelihood for predator coexistence based on relative non-linearity in their functional responses (Figures 3E, F; 5E, F) because it promotes unstable predator-prey dynamics, especially for lower values of h_{C_2}. This can be seen by the higher values of the Hopf-bifurcation point $d_{C_2}^H$ (dashed dotted lines in Figure 5E, F) at which the $R - C_2$ system gives rise to predator-prey oscillations. In our example, a non-linear nutrient uptake rate of the prey increases the absolute size of the parameter range of potential species coexistence (P_a) by 18% and the absolute size of the parameter range of realized species coexistence (O_a) by 48% for $h_2 = 1$ (Figure 5A&E). In contrast, the positive effect of a non-linear nutrient uptake rate is less pronounced for $h_2 = 4$, in which case the increase was limited to 2% for P_a and 8% for O_a (Figure 5B&F). This observation can be explained by the fact that P_a and O_a were already close to their maxima for $h_2 = 4$, i.e. the Hopf-bifurcation point $d_{C_2}^H$ of the $R - C_2$ system almost coincides with the upper limit of d_{C_2} ($d_{C_2}^U$) (cf. Figure 5B&F). In contrast, for $h_{C_2} = 1$, the positive effect of a non-linear nutrient uptake rate of the prey is quite large given that $d_{C_2}^H$ was substantially below $d_{C_2}^U$ (cf. Figure 5A&E).

3.4 Coexistence is more likely in closed systems than in flow-through systems

Adding a shared loss rate to all species by considering a chemostat (i.e. flow-through) system with a dilution rate δ substantially reduces the potential of the two predators to coexist on the single prey. An increase in δ always strongly decreases the size of the parameter ranges of potential (P_a) and realized (O_a) species coexistence, independent of h_{C_2}, i.e. the exact curvature of the type-II functional response of C_2 (Figures 3G, 5G-H vs. 5A-B). For example, P_a and O_a are reduced by 28% and 42%, respectively, when $h_{C_2} = 1$ and by 36% and 40% when $h_2 = 4$.

This happens because δ reduces the preys' carrying capacity, which stabilizes the predator-prey dynamics (Appendix B), and decreases the maximum death rates of the two predators they can withstand (cf. scale of y axes in Figure 5).

In contrast to P_a, the relative size of the parameter region that potentially allows for predator coexistence (P_r) increases with δ, independent of h_{C_2} (Figure 3H). For instance, P_r increases by 6% when $h_{C_2} = 1$ and by 29% when $h_2 = 4$ (Figure 5G-H vs. 5A-B). This is explicable by the larger negative impact of δ on C_2 than C_1 due to its lower maximum growth and death rates, which renders C_2 more sensitive to additional mortality (cf. Figure 2A). Consequently, C_2 cannot exclude C_1 anymore when the death rate of C_1 is sufficiently low, e.g. below ≈ 0.3 in Figure 5H (dashed line). Furthermore, P_r may exceed 0.5 when the dilution rate is sufficiently high (Figure 3H), which strongly contrasts with closed systems where $\delta = 0$.

In contrast to P_a, O_a and P_r, the response of the relative size of the parameter region that actually allows for predator coexistence (O_r) to alterations of δ depends on h_{C_2}. For example, O_r decreases by 14% when $h_{C_2} = 1$ (Figure 5G vs. 5A) because the stabilizing effect of δ on the predator-prey dynamics reduces predator coexistence more than any beneficial effect arising through a reduced risk of C_1 to get excluded by C_2 at lower death rates of both predators. In comparison, O_r increases by 25% when $h_2 = 4$ (Figure 5H vs. 5B) because the positive effect of δ on predator coexistence at lower death rates outweighs the weakly stabilizing effect of δ on the predator-prey dynamics and thus negative impact on predator coexistence.

4. Discussion

For almost 50 years, it has fascinated ecologists that two different predators may be able to coexist on a single fluctuating prey when they exhibit substantial differences in the curvature of their functional responses (Koch 1974; Chesson 2000; Abrams and Holt 2002). Previous studies explored this mechanism of species coexistence in predator-prey models that are based either on the classical Rosenzweig-MacArthur (RM) equations (1963) with logistically growing prey (Hsu et al. 1978; Armstrong and McGehee 1980; Abrams 2004; Xiao and Fussmann 2013), or on chemostat equations with explicit nutrient dynamics (Butler and Waltman 1981; Butler et al. 1983; Keener 1985; Smith 1995). The main motivation for using the latter was to derive predictions, which can be experimentally tested (e.g. Butler and Waltman 1981). The RM equations are often believed to mainly differ from corresponding chemostat models by assuming a linear uptake rate of the prey for the limiting resource (e.g. Grover and Holt 1998). This impression arises because logistic growth, as it is modelled by RM equations, can be

derived from an explicit consumer-resource model that assumes a linear uptake rate of the consumer for the limiting resource (Armstrong and McGehee 1980). However, this derivation is based on the important assumption that no other trophic levels, such as predators, are present. By adding another trophic level to the original system, we show that predators may influence their prey not only by predation but also by emergent nutrient competition (Figure 1; Appendix B). Hence, the RM model reflects conditions where the prey is limited by a resource not affected by the predators such as light or space, rather than nutrients. The two different types of predator-prey models thus substantially differ in their assumption about the type, uptake and replenishment of the resource that is limiting the growth of the prey.

Therefore, we analyzed a generic predator-prey model that accounts explicitly for the uptake and replenishment of nutrients and compared it to the classical RM predator-prey model. We evaluated the impact of nutrient competition between the two different trophic levels on the potential of two predators to coexist on a single fluctuating prey through relative non-linearity in their functional responses. Our results demonstrate that nutrient competition between the predators and the prey strongly stabilizes the predator-prey dynamics and thereby substantially reduces the potential for predator coexistence. The stabilizing effect of nutrient competition originates from the negative impact of the predators on the effective carrying capacity of their own prey. Our study thus demonstrates that the type of resource limitation may substantially influence the stability of predator-prey dynamics and the likelihood for predator coexistence through fluctuation-dependent mechanisms, with potentially far-reaching consequences for the overall stability of entire food-webs.

Since nutrient limitation of the (autotrophic) prey is common in aquatic (Sommer and Lampert 2007) and terrestrial ecosystems (Santiago and Goldstein 2016), we can expect that predators are very likely to influence the amount of the limiting resource for their prey in many natural systems. This holds in particular for (aquatic) ecosystems that frequently exhibit top-heavy biomass distributions in which the heterotrophic biomass is at least of similar magnitude as the autotrophic biomass (del Giorgio and Gasol 1995; Burkepile 2013; McCauley et al. 2018; Shurin et al. 2006). Given the high nutrient to carbon ratios of animals compared to plants (Sterner and Elser 2002; Hall 2009; Hessen et al. 2013), the majority of nutrients in aquatic ecosystems is stored in the consumer biomass (Hassett et al. 1997; Gaedke et al. 2002). Hence, our predator-prey model described by eq. [2] captures an essential property of many natural systems, which is not entailed in the Rosenzweig-MacArthur (RM) equations (1963). This is particularly relevant, because important ecological features such as the paradox of enrichment (Rosenzweig 1971), the hydra effect (Abrams 2009; Sieber and Hilker 2012), biological chaos

in three species food chains (Hastings and Powell 1991; McCann and Yodiz 1994), the impact of trait adaptation on species coexistence (Abrams 2006; Klauschies et al. 2016) and the stability of complex food webs (Brose et al. 2006; Williams and Martinez 2008) were initially investigated in models based on the RM equations.

Nutrient competition between predators and prey may even be relevant for (terrestrial) ecosystems with bottom-heavy biomass distributions (Polis 1999; Shurin et al. 2006; Hatton et al. 2015). On a global scale, the biomass ratio of heterotrophs to autotrophs is about 0.05 in terrestrial ecosystems (Bar-On et al. 2018). Assuming a five times higher nutrient to carbon ratio for heterotrophs than for autotrophs (e.g. Lemoine et al. 2014; Sterner and Elser 2002), the heterotrophic biomass contains 20% of the total amount of nutrients. Hence, nutrient competition between herbivores and plants is likely to be important also for the overall population dynamics in terrestrial ecosystems. Only if predators mainly affect their prey by consumption and less by nutrient competition, predator-prey models based on the Rosenzweig-MacArthur equations (1963) may well capture the observed predator-prey dynamics, e.g. in hyper-eutrophic systems where prey are rather light- than nutrient-limited (Hautier et al. 2009).

Our study further shows that the potential for predator coexistence on a single prey also depends on the non-linearity of the prey's uptake rate for the resource. A linear uptake rate is to be expected under strong bottom-up control of the prey where the limiting resource is scarce, e.g. severe light limitation in winter or ultra-oligotrophic conditions in the open ocean (Lampert and Sommer 2007; Sommer 2005). Otherwise non-linear relationships are more realistic (Michaelis and Menten 1913; Murdoch 1973). This is very important given that our results show that a non-linear nutrient-uptake rate of the prey may strongly promote predator-prey oscillations, and thereby facilitate predator coexistence when the relative non-linearity between the predators' functional responses is relatively weak. In contrast, a non-linear resource uptake rate of the prey hardly influenced the likelihood for predator coexistence when the relative non-linearity between the predators' functional responses was rather strong. This is in line with previous findings from a modified RM predator-prey model that includes theta-logistic growth to reflect a non-linear nutrient uptake rate of the prey (Abrams and Holt 2002). Depending on whether the stabilizing effect of resource competition between predators and prey on the populations dynamics prevails over the destabilizing effect of a non-linear resource uptake rate of the prey, predator coexistence in our model is more or less likely compared to the corresponding Rosenzweig-MacArthur model with logistically growing prey.

Finally, our results demonstrate that the likelihood for predator coexistence on a single prey species decreases if all species experience a shared loss rate because it reduces the effective

carrying capacity of the prey, and the maximum death rates the predators can withstand. Furthermore, in line with previous results for standard chemostat models (Smith 1995), a higher dilution rate stabilizes predator-prey dynamics and thereby also reduces the likelihood for predator coexistence through relative non-linearity in their functional responses. While adding a shared loss rate to our predator-prey system does not alter the total amount of nutrients in the system, it does affect the distribution of nutrients, i.e. the ratio between particulate organic nutrients and dissolved nutrients. A shared loss rate may naturally arise through water exchange or an omnivorous top predator feeding on predators and prey.

There are two fundamental requirements for predator coexistence through relative non-linearity in their functional responses. One are fluctuations in the abundance of the single prey that can be externally driven (Sommer 1985; Grover 1988; Abrams 2004) or endogenously generated by predator-prey interactions (Armstrong and McGehee 1980). The latter essentially depends on the potential of the predators' functional response to destabilize the corresponding predator-prey dynamics, for which the functional response has to be sufficiently non-linear for the ecologically feasible prey densities. This condition is likely to be met in many natural systems, given ample experimental evidence for invertebrate predators to exhibit substantial curvature in their functional responses (Hassell et al. 1977; Rothaupt 1988; Jeschke et al. 2004; Sarnelle and Wilson 2008; Seifert et al. 2014).

The second requirement for predator coexistence are differences in the predators' abilities to exploit efficiently periods of low or high prey abundances in line with the gleaner-opportunist trade-off (Fredrickson and Stephanopoulos 1981; Xiao and Fussmann 2013). Experimental results and field observations suggest that the gleaner-opportunist trade-off may result from size differences among predators or from differences in their feeding or life-history strategies. For example, smaller rotifers (the gleaners) had lower prey requirements than larger rotifers allowing the smaller species to outcompete the larger species at lower prey densities (Stemberger and Gilbert 1985; Sarma et al. 1996). In contrast, larger rotifers (the opportunists) are better competitors at higher prey densities, enabling them to exploit favorable conditions of high prey densities (Boraas et al. 1990; Sarma et al. 1999). Another example are filter-feeding zooplankton such as cladocerans with often linear functional responses (Jeschke et al. 2004) and high maximum growth rates (Sommer and Stibor 2002). In contrast, interception feeders such as copepods show non-linear functional responses (Jeschke et al. 2004) and lower maximum growth rates (Sommer and Stibor 2002).

However, a direct experimental demonstration of predator coexistence through endogenously generated fluctuations in the prey abundance and differences in the functional

responses of different predators is still lacking. An important reason might be that different coexistence mechanisms frequently operate in parallel and that it is rather difficult to experimentally test for the individual effect of a particular coexistence mechanism (Ellner et al. 2018). In fact, predator coexistence on a single prey is also possible in the absence of predator-prey oscillations, for example, when predators show interference between their conspecifics (Vance 1984; Hsu et al. 2013), utilize different life stages of the prey (Haigh and Maynard Smith 1972), or are limited by different currencies of the prey such as food density and food quality (Loladze et al. 2000, 2004; Moe et al. 2005; Elser et al. 2012).

Conclusions

We show that the potential of two predators to coexist on a single fluctuating prey species through differences in the curvature of their functional responses depends on the type of resource that is limiting the growth of the prey. In contrast to light or space limitation of the prey, predators and prey compete for the limiting resource when the prey exhibits nutrient limitation. This feature strongly stabilizes the predator-prey dynamics and thereby reduces the likelihood for predator coexistence. In contrast, a non-linear resource uptake rate of the prey promotes predator-prey oscillations, and thus, predator coexistence. Predator coexistence through relative non-linearity in their functional responses is also less likely to occur in flow-through systems where all species suffer from an additional shared loss rate. Hence, fluctuation-dependent mechanisms of species coexistence are most likely to promote biodiversity in closed systems where the basal prey exhibits light or space limitation. In contrast, under nutrient competition between predators and prey, the maintenance of biodiversity may strongly rely on fluctuation-independent mechanisms.

Author contributions

Toni Klauschies conceived the idea of this study, developed the mathematics included, coded and simulated the model, analyzed the data and wrote the manuscript. Ursula Gaedke commented on the manuscript and improved the writing.

Acknowledgments

We thank Michael Sieber, Ruben Ceulemans, and Alice Boit for helpful comments and suggestions. Toni Klauschies was funded by the German Research Foundation (DFG, GA 401/26-1).

References

Abrams, P. A. and R. D. Holt (2002) The impact of consumer-resource cycles on the coexistence of competing consumers. *Theoretical Population Biology* 62:281-295.

Abrams, P. A. (2004) When does periodic variation in resource growth allow robust coexistence of competing consumer species. *Ecology* 85: 372-382.

Abrams, P. A. (2006) The Prerequisites for and Likelihood of Generalist-Specialist Coexistence. *The American Naturalist* 167: 329-342.

Abrams, P. A. (2009) When does greater mortality increase population size? The long history and diverse mechanisms underlying the hydra effect. *Ecology Letters* 12: 462-474.

Armstrong, R. A. and R. Mcgehee (1980) Competitive-Exclusion. *The American Naturalist* 115: 151-170.

Barabás, G., R. D'Andrea and S. M. Stump (2018) Chesson's coexistence theory. *Ecological Monographs* 88: 277-303.

Bar-On, Y. M., R. Phillips and R. Milo (2018) The biomass distribution on Earth. *Proceedings of the National Academy of Sciences of the United States of America* 115: 6506-6511.

Boraas, M. E., D. B. Seale and J. B. Horton (1990) Resource competition between two rotifer species (Brachionus rubens and B. calyciflorus): an experimenal test of mechanistic model. Journal of Plankton Research 12: 77–8.

Brose, U., R. J. Williams and N. D. Martinez (2006) Allometric scaling enhances stability in complex food webs. *Ecology Letters* 9: 1228-1236.

Burkepile, D. E. (2013) Comparing aquatic and terrestrial grazing ecosystems: is the grass really greener? *Oikos* 122: 306-312.

Butler, G. J. and P. Waltman (1981) Bifurcation from a limit cycle in a two predator-one prey ecosystem modeled on a chemostat. *Journal of Mathematical Biology* 12: 295-310.

Butler, G. J., S. B. Hsu and P. Waltman (1983) Coexistence of Competing Predators in a Chemostat. *Journal of Mathematical Biology* 17: 133-151.

Chase, J. M., and M. A. Leibold (2003) Ecological niches: linking classical and contemporary approaches. The University of Chicago Press, Chicago, Illinois, USA.

Chesson, P. 2000. Mechanisms of maintenance of species diversity. *Annual Review of Ecology and Systematics* 31:343-366.

Chesson, P. (2003) Quantifying and testing coexistence mechanisms arising from recruitment fluctuations. *Theoretical Population Biology* 64: 345–357.

Chesson, P. (2018) Updates on mechanisms of maintenance of species diversity. *Journal of Ecology* 106: 1773–1794. DOI: 10.1111/1365-2745.13035

Chesson, P. and R. R. Warner (1981) Environmental Variability Promotes Coexistence in Lottery Competitive Systems. *The American Naturalist* 117: 923-943.

Droop, M. R. (2003) In defence of the Cell Quota model of micro-algal growth. *Journal of Plankton Research* 25: 103-107.

Dugdale, R. C. (1967) Nutrient limitation in the sea: dynamics, identification, and significance. *Limnology and Oceanography* 12: 685-695

Ellner, S. P., R. E. Snyder and P. B. Adler (2016) How to quantify the temporal storage effect using simulations instead of math. *Ecology Letters* 19: 1333-1342.

Ellner, S. P., R. E. Snyder, P. B. Adler and G. Hooker (2018) An expanded modern coexistence theory for empirical applications. *Ecology Letters* 22: 3-18.

Eppley, R. W., J. N. Rogers and J. J. McCarthy (1969) Half-saturation constants for uptake of nitrate and ammonium by marine phytoplankton. *Limnology and Oceanography* 14: 912-920.

Fredrickson, A. G. and G. Stephanopoulos (1981) Microbial Competition. Science 213: 972-979.

Gaedke, U., S. Hochstädter and D. Straile (2002) Interplay between energy limitation and nutrional deficiency: empircial data and food web models. *Ecological Monographs* 72: 251-270.

Gause, G. F. (1932) Experimental Studies on the struggle for existence. *Journal of Experimental Biology* 9: 389-402.

del Giorgio, P. A. and J. M. Gasol (1995) Biomass Distribution in Freshwater Plankton Communities. *The American Naturalist* 146: 135-152.

Grover, J. P. (1988) Dynamics of competition in a variable environment: experiments with two diatom species. *Ecology* 69: 408–417.

Grover, J. P. (1997) Resource Competition. Chapman & Hall, London, UK.

Grover, J. P. and R. D. Holt (1998) Disentangling Resource and Apparent Competition: realistic Models for Plant-herbivore Communities. *Journal of Theoretical Biology* 191: 353-376.

Haigh, J. and J. Maynard Smith (1972) Can There be More Predators Than Prey? *Theoretical Population Biology* 3: 290-299.

Hall, S. R. (2009) Stoichiometrically Explicit Food Webs: Feedbacks between Resource Supply, Elemental Constraints, and Species Diversity. *Annual Review of Ecology, Evolution, and Systematics* 40: 503–528.

Hardin, G. (1960) The competitive exclusion principle. *Science* 131:1292-1298.

Hassel, M. P., J. H. Lawton and J. R. Beddington (1977) Sigmoid Functional Responses by Invertebrate Predators and Parasitoids. *Journal of Animal Ecology* 46: 249-262.

Hasset, R. P., B. Cardinale, L. B. Stabler and J. J. Elser (1997). Ecological stichiometry of N and P in pelagic ecosystems: Comparison of lakes and oceans with emphasis on the zooplankton-phytoplankton interaction. *Limnology and Oceanography* 42. 648-662.

Hastings, A. and T. Powell (1991) Chaos in a three-species food chain. *Ecology* 72: 896-903.

Hatton, I. A., K. S. McCann, J. M. Fryxell, T. J. Davies, M. Smerlak, A. R. E. Sinclair and M. Loreau (2015) The predator-prey power law: Biomass scaling across terestrial and aquatic biomes. *Science* 349: 1070-1083.

Hautier, Y., P. A. Niklaus and A. Hector (2009) Competition for Light Causes Plant Biodiversity Loss After Eutrophication. *Science* 324: 636-638.

Holling, C. S. (1961) Principles of Insect Predation. *Annual Review of Entomology* 6: 163-182.

Hsu, S. B., S. P. Hubbell and P. Waltman (1978) A Contribution to the Theory of Competing Predators. *Ecological Monographs* 48: 337-349.

Hsu, S. B., S. Ruan and T.-H. Yang (2013) On the dynamics of two-consumers-one-resource competing systems with Beddington-Deangelis functional response. *Discrete and Continuous Dynamical Systems Series B* 18: 2331-2353.

Hutchinson, G. E. (1961) The Paradox of the Plankton. *The American Naturalist* 95:137-145

Jeschke, J. M., M. Kopp and R. Tollrian (2004) Consumer-food systems: why type I functional responses are exclusive to filter feeders. *Biological Reviews* 79: 337-349.

Lampert, W. and U. Sommer (2007) Limnoecology. The ecology of lakes and streams. Oxford University Press, New York.

Lemoine, N. P., S. T. Giery and D. E. Burkepile (2014) Differing nutritional constraints of consumers across ecosystems. *Oecologia* 174:1367–137.

Letten, A. D., P.-J. Ke and T. Fukami (2017) Linking modern coexistence theory and contemporary niche theory. *Ecological Monographs* 87: 161–177.

Litchman, E. and C. A. Klausmeier (2001) Competition of Phytoplankton under Fluctuating Light. *The American Naturalist* 157: 170-187.

Loladze, I., Y. Kuang, J. J. Elser and W. F. Fagan (2004). Competition and stoichiometry: coexistence of two predators on one prey. *Theoretical Population Biology* 65: 1-15.

Lotka, A. J. (1925) Elements of Physical Biology. Baltimore, Williams and Wilkins. (Reprinted as Elements of Mathematical Biology (1956) Dover, NY).

McCann, K. and P. Yodiz (1994) Biological Conditions for Chaos in a Three-Species Food Chain. *Ecology* 75: 561-564.

Keener, J. P. (1985) Oscillatory coexistence in a food chain model with competing predators. *Journal of Mathematical Biology* 22: 123-135.

Klauschies, T., D. A. Vasseur and U. Gaedke (2016) Trait adaptation promotes species coexistence in diverse predator and prey communities. *Ecology and Evolution* 6: 4141–4159.

Koch, A. L. (1974) Competitive Coexistence of Two Predators Utilizing the Same Prey Under Constant Environmental Conditions. *Journal of Theoretical Biology* 44: 387-395.

Kooi, B. W., M. P. Boer and S. A. L. M. Kooijman (1998) On the Use of the Logistic Equation in Models of Food Chains. *Bulletin of Mathematical Biology* 60: 231-246.

MacArthur, R. and R. Levins. 1967. Limiting Similarity Convergence and Divergence of Coexisting Species. *The American Naturalist* 101: 377-385.

McCauley, D. J., G. Gellner, N. D. Martinez, R. J. Williams, S. A. Sandin, F. Micheli, P. J. Mumby and K. S. McCann (2018) On the prevalence and dynamics of inverted trophic pyramids and otherwise top-heavy communities. *Ecology Letters* 21: 439-454.

Michalis, L. and M. L. Menten (1913) Die Kinetik der Invertinwerkung. Biochemische Zeitschrift 49: 333-369.

Mittelbach, G. G. (2012) Community Ecology. Oxford Univeristy Press, New York.

Moe, S. J., R. S. Stelzer, M. R. Forman, W. S. Harpole, T. Daufresne and T. Yoshida (2005) Recent advances in ecological stoichiometry: insights for population and community ecology. *Oikos* 109: 29-39.

Murdoch, W. W. (1973) Functional response of predators. *Journal of Applied Ecology* 10: 335–342.

Polis, G. A. (1999) Why Are Parts of the World Green? Multiple Factors Control Productivity and the Distribution of Biomass. *Oikos* 86: 3-15.

Rosenzweig, M. L. and R. H. MacArthur (1963) Graphical Representation and Stability Conditions of Predator-Prey Interactions. *The American Naturalist* 97:209-&.

Rosenzweig, M. L. (1971) Paradox of Enrichment: Destabilization of Exploitation Ecosystems in Ecological Time. *Science* 171: 385-387.

Rothhaupt, K. O. (1988) Mechanistic resource competition theory applied to laboratory experiments with zooplankton. *Nature* 333: 660–662.

Santiago, L. S. and G. Goldstein (2016) Is Photosynthesis Nutrient Limited in Tropical Trees? In: Goldstein G., Santiago L. (editors) Tropical Tree Physiology, pp 299-315. Tree Physiology, volume 6. Springer, Cham.

Sarma, S. S. S., N. Iyer and H. J. Dumont (1996) Competitive interactions between herbivorous rotifers: importance of food concentration and initial population density. Hydrobiologia 331: 1–7.

Sarma, S. S. S., M. Alfredo Fernández Araiza and S. Nandini (1999) Competition between Brachionus calyciflorus Pallas and Brachionus patulus (Müller) (Rotifera) in relation to algal food concentration and initial population density. *Aquatic Ecology* 33: 339–345.

Sarnelle, O. and A. E. Wilson (2008) Type III Functional Response in Daphnia. Ecology 89: 1723-1732.

Schoener, T. W. (1974) Some methods of calculating competition coefficients from resource-utilization spectra. The American Naturalist 108: 332-340.

Seifert, L. I., F. de Castro, A. Marquart, U. Gaedke, G. Weithoff and M. Vos (2014) Heated Relations: Temperature-Mediated Shifts in Consumption across Trophic Levels. PLoS ONE 9: e95046.

Shurin, J. B., D. S. Gruner and H. Hillebrandt (2006) All wet or dried up? Real differences between aquatic and terrestrial food webs. *Proceedings of the Royal Society B* 273: 1-9.

Sibly, R. M. and J. Hone (2002) Population growth rate and its determinants: an overview. *Philosophical Transactions of the Royal Society B* 357: 1153–1170.

Sieber, M. and F. M. Hilker (2012) The hydra effect in predator–prey models. *Journal of Mathematical Biology* 64: 341-360.

Smith, H. L. and P. Waltman (1995) The Theory of the Chemostat. Dynamics of Microbial Competition. Cambridge University Press, New York.

Sommer, U. (1985) Comparison between steady-state and nonsteady state competition-experiments with natural phytoplankton. Limnology and Oceanography 30:335–346.

Sommer, U. (2005) Biologische Meereskunde. Springer-Verlag Berlin Heidelberg, New York.

Stemberger, R. S. and J. J. Gilbert (1985) Body size, food concentration and population growth in planktonic rotifers. Ecology 66: 1151–1159.

Sterner, R. W. and J. J. Elser (2002) Ecological Stoichiometry. The Biology of Elements from Modelcules to the Biosphere. Princeton University Press, New Jersey.

Vance, R. R. (1984) Interference Competition and the Coexistence of Two Competitors on a Single Limiting Resource. *Ecology* 65: 1349-1357.

Volterra, V. (1928) Variations and fluctuations of the number of individuals in animal species living together. *ICES Journal of Marine Science* 3: 3-51.

Williams R. J. and N. D. Martinez (2008) Success and its limits among structural models of complex food webs. *Journal of Animal Ecology* 77: 512-519.

Xiao, X. and G. F. Fussmann (2013) Armstrong-McGehee mechanism revisited: Competitive exclusion and coexistence of non-linear consumers. *Journal of Theoretical Biology* 339: 26-35.

Yamamichi, M., T. Yoshida, and A. Sasaki (2011) Comparing the Effects of Rapid Evolution and Phenotypic Plasticity on Predator-Prey Dynamics. *The American Naturalist* 178:287-304.

Appendix A: Generic predator-prey model

1. Description of a general predator-prey model

In this section, we derive a generic predator-prey model that integrates and extends previous studies studying the potential for predator coexistence on a single prey species, through differences in their functional responses and endogenously generated fluctuations in species abundances under batch (Koch 1974; Armstrong and McGehee 1980; Hsu et al. 1978; Abrams and Holt 2002; Xiao and Fussmann 2013), or chemostat conditions (Butler and Waltman 1981; Butler et al. 1983; Keener 1985). Our model comprises two different predators and one prey species and explicitly accounts for the replenishment, uptake, excretion and remineralization of mineral nutrients such as nitrogen or phosphorus in a flow-through system:

$$\frac{dN}{dt} = \underbrace{\delta(N_I - N)}_{replenishment} - \underbrace{\frac{\eta_R a_R N R}{1 + h_R a_R N}}_{uptake} + \underbrace{\sum_{i=1}^{2}\left(\eta_R - \eta_{C_i} e_{C_i}\right)\frac{a_{C_i} R C_i}{1 + h_{C_i} a_{C_i} R}}_{excretion} + \underbrace{\eta_R d_R R + \sum_{i=1}^{2}\eta_{C_i} d_{C_i} C_i}_{remineralization} \quad [A1a]$$

$$\frac{dR}{dt} = \left[\frac{a_R N}{1 + h_R a_R N} - \sum_{i=1}^{2}\frac{a_{C_i} C_i}{1 + h_{C_i} a_{C_i} R} - d_R - \delta\right]R \quad [A1b]$$

$$\frac{dC_1}{dt} = \left[\frac{e_{C_1} a_{C_1} R}{1 + h_{C_1} a_{C_1} R} - d_{C_1} - \delta\right]C_1 \quad [A1c]$$

$$\frac{dC_2}{dt} = \left[\frac{e_{C_2} a_{C_2} R}{1 + h_{C_2} a_{C_2} R} - d_{C_2} - \delta\right]C_2 \quad [A1d]$$

Hence, our model tracks the concentration of dissolved inorganic nutrients N and the different pools of particulate organic matter, i.e. the carbon biomasses of the prey R and the two predators C_1 and C_2 in a chemostat system. The replenishment of N depends on the dilution rate δ, the concentrations of dissolved inorganic nutrients in the inflowing medium N_I and in the chemostat N, the nutrient-uptake of the prey, the excretion of nutrients by the predators and the remineralization of particulate organic nutrients due to the decomposition of dead material that originates from all trophic levels. Excretion and remineralization depend on the nutrient to carbon ratios $\eta = \left[\frac{N}{C}\right]$ of the different species, which we assumed to be constant.

Nutrient-uptake of the autotrophic prey follows a type II functional response, i.e. Michaelis-Menten-Kinetik (Michaelis & Menten 1913), with uptake rate α_R and handling time h_R (Holling, 1961). Inorganic nutrients are converted into prey biomass according to its specific nutrient to carbon ratio η_R. The two predators also feed on the prey with a type II functional response with species-specific attack rates α_{C_i} and handling times h_{C_i}. The ingested biomass of the prey is converted into biomass of the predators with species-specific transfer efficiencies e_{C_i}. To reflect most natural conditions, predators are assumed to be energy- and thus, carbon-limited. Hence, the transfer efficiencies of the predators ensure that the stoichiometry of the

prey meets the nutritional demands of the predators, i.e. $e_{C_i} < \frac{\eta_R}{\eta_{C_i}} \leftrightarrow \eta_R - \eta_{C_i} e_{C_i} > 0$. Hence,

the food quality of the prey, i.e. $\frac{\eta_R}{\eta_{C_i}} = \left[\frac{N}{C}\right]_R \Big/ \left[\frac{N}{C}\right]_{C_i}$, is sufficiently high to support the

energetically feasible biomass production of the predators in units of carbon. Therefore, the loss of energy through respiration or low assimilation efficiencies of carbon prevents further growth of the predator biomasses so that excretion may substantially contribute to the allocation of dissolved inorganic nutrients to the system. In contrast, we neglected the case where the prey is of low (stoichiometric) food quality, i.e. $e_{C_i} > \frac{\eta_R}{\eta_{C_i}}$, and thus where the predators are nutrient-

limited. In such a case, a lack of essential nutrients in the prey will prevent the production of more predator biomass, which implies that re-mineralization will be more important for filling up nutrients than nutrient excretion.

Furthermore, all nutrients and organisms are washed out from the system with the dilution rate δ, constituting a shared loss rate for all species. Finally, we also accounted for senility or basal mortality of the different species by incorporating the species-specific death rates d_R and d_{C_i} for the prey and the two predators, respectively. Accounting for remineralization and recycling of nutrients ensures that the predator-prey model described by eq. [A1] is mass-balanced.

2. *Model simplification*

As detailed below, the predator-prey model defined by eq. [A1] contains a conserved quantity. This allows us to reduce its dimension and thereby to simplify our analysis. Following the methodology used by Armstrong and McGehee (1980), Jones and Ellner (2007), Scranton and Vasseur (2016) and O'Dwyer (2018), we first define the total amount of dissolved inorganic nutrients and particulate organic nutrients bound in the predator and prey biomasses as follows: $N_T = N + \eta_R R + \eta_{C_1} C_1 + \eta_{C_2} C_2$. Consequently, the rate of change of N_T is given by:

$$\frac{dN_T}{dt} = \frac{dN}{dt} + \eta_R \frac{dR}{dt} + \eta_{C_1} \frac{dC_1}{dt} + \eta_{C_2} \frac{dC_2}{dt} = \delta(N_I - N_T) \tag{A2}$$

We can now solve this simple linear ordinary differential equation analytically by substitution

with $N_T{}' = N_T - N_I$. Since $\frac{dN_T{}'}{dt} = \frac{dN_T}{dt}$ we get:

$$\frac{dN_T{}'}{dt} = \delta(N_I - N_T) = \delta\big(N_I - (N_T{}' + N_I)\big) = -\delta N_T{}' \tag{A3}$$

This equation represents exponential growth, which has the solution:

$$N_T{}'(t) = N_T{}'(0)e^{-\delta t} \tag{A4}$$

Using the backwards substitution terms $N_T(t) = N_T{'}(t) + N_I$ and $N_T{'}(0) = N_T(0) - N_I$ we can now express the solution in terms of $N_T(t)$ again:

$$N_T(t) = N_I + (N_T(0) - N_I)e^{-\delta t} \tag{A5}$$

In the limit as $t \to \infty$ the system reaches its asymptotically stable equilibrium: $N_T(t) \to N_I$. Therefore, we can write a steady-state approximation for N_T:

$$N = N_I - \left(\eta_R R + \eta_{C_1} C_1 + \eta_{C_2} C_2\right) \tag{A6}$$

Substituting eq. [A6] into eq. [A1] results into the dimensionally reduced predator-prey model:

$$\frac{dR}{dt} = \left[\frac{a_R\left(N_I - (\eta_R R + \eta_{C_1} C_1 + \eta_{C_2} C_2)\right)}{1 + h_R a_R\left(N_I - (\eta_R R + \eta_{C_1} C_1 + \eta_{C_2} C_2)\right)} - \sum_{i=1}^{2}\frac{a_{C_i} C_i}{1 + h_{C_i} a_{C_i} R} - d_R - \delta\right] R \tag{A7a}$$

$$\frac{dC_1}{dt} = \left[\frac{e_{C_1} a_{C_1} R}{1 + h_{C_1} a_{C_1} R} - d_{C_1} - \delta\right] C_1 \tag{A7b}$$

$$\frac{dC_2}{dt} = \left[\frac{e_{C_2} a_{C_2} R}{1 + h_{C_2} a_{C_2} R} - d_{C_2} - \delta\right] C_2 \tag{A7c}$$

The nutrient to carbon ratios η_R, η_{C_1} and η_{C_2} determine the strength of nutrient competition among all prey and predator species. Hence, in line with Lotka-Volterra competition models, we can interpret them as species-specific competition coefficients for nutrients. More generally, this model allows us to capture various scenarios of resource limitation. When mineral nutrients such as phosphorus or nitrogen are limiting the growth of the prey, predators influence the prey negatively through consumption and competition. Depending on the actual values of the attack rates and competition coefficients of the two predators, the negative effects of the two predators on the prey by nutrient competition may surpass those by predation. Nutrient competition among the prey and predators arise from the fact that all species contain essential nutrients in their biomasses as particulate organic nutrients. In this bounded state, the limiting nutrients are not available for the growth of the autotrophic prey, which effectively reduces its carrying capacity. In contrast, when light or space is limiting prey growth, the two predators do not compete with the prey for the resource at all because they do not contribute to the self-shading or self-limitation of the prey.

Furthermore, in predator-prey systems that only comprise two predators and one single prey, predator coexistence strongly depends on the relative non-linearity in their functional responses (Hsu et al. 1978; Chesson 2000; Abrams and Holt 2002). Hence, coexistence is most likely to occur if one predator exhibits a linear functional response whereas the functional response of the other predator shows substantial curvature (Abrams and Holt 2002; Xiao and Fussmann 2013). Therefore, we restricted our analysis to the case where C_1 is not limited by handling the prey, i.e. $h_{C_1} = 0$.

In addition, the parameter region for which the two different predators may coexist also increases with a higher carrying capacity of the prey, i.e. the equilibrium biomass of the prey in the absence of the two predators (Hsu et al. 1978). This results from the fact that a higher carrying capacity promotes unstable population dynamics, which are necessary for the two predators to coexist due to differences in their functional responses (for details see Appendix B). Given that an increase in d_R effectively reduces the carrying capacity, i.e. maximum achievable biomass, of the prey and thereby the likelihood of predator coexistence, we focused our analysis on the case, where the basal prey mortality was negligible, i.e. $d_R = 0$. These modifications lead to the simplified predator-prey model used in the main text (cf. eq. [1])

Additional references

Jones, L. E. and S. P. Ellner (2007) Effects of rapid prey evolution on predator–prey cycles. *Journal of Mathematical Biology* 55: 541-573.

Scranton, K. and D. A. Vasseur (2016) Coexistence and emergent neutrality generate synchrony among competitors in fluctuating environments. *Theoretical Ecology* 9: 353-363.

O'Dwyer, J. P. (2018) Whence Lotka-Volterra? Conservation laws and integrable systems in ecology. *Theoretical Ecology* 11: 441-452.

Appendix B: Necessary conditions and parameter range of potential species coexistence

Following the general methodology of Hsu et al. (1978) and Xiao and Fussmann (2013) we here derive ecological conditions for species coexistence that determine a range of parameters for which species coexistence might be possible. Although these conditions are not sufficient, they are necessary, implying that outside this parameter range species coexistence is impossible. For simplicity, we will assume predator one (C_1) to exhibit a linear functional response and predator two (C_2) to have a (non-linear) functional type II response.

1) Persistence of predator-prey systems in the absence of exploitative competition

Predator coexistence necessarily requires persistence of the two different predator-prey systems comprising either C_1 or C_2 next to the prey. Hence, each predator has to be able to invade the system that only includes the prey. In the absence of C_1 and C_2, the prey is approaching its single-species equilibrium $R_S^* = 1 - \frac{\delta}{(1-h_R\delta)}$. The two predators thus exhibit positive invasion growth rates when their grazing rates for R_S^* are larger than the sum of their different loss rates through mortality (d_{C_i}) and dilution (δ), respectively, i.e. $a_{C_1} R_S^* > d_{C_1} + \delta$ for C_1 and $\frac{a_{C_2} R_S^*}{1 + h_{C_2} a_{C_2} R_S^*} > d_{C_2} + \delta$ for C_2. Consequently, the death rates that allow persistence of the two different predator-prey systems are limited to $d_{C_1} < a_{C_1} \left(1 - \frac{\delta}{(1-h_R\delta)}\right) - \delta$ for C_1 and to $d_{C_2} < \frac{a_{C_2}(1-(h_R+1)\delta)}{(1-h_R\delta) + h_{C_2} a_{C_2}(1-(h_R+1)\delta)} - \delta$ for C_2. These conditions put natural limits on the two-dimensional parameter space in the $d_{C_1} - d_{C_2}$ plane for which predator coexistence might be possible. The first necessary condition for predator coexistence is thus given by:

$$N_1: \qquad \begin{matrix} d_{C_1} < d_{C_1}^U \\ d_{C_2} < d_{C_2}^U \end{matrix}$$

with $d_{C_1}^U = a_{C_1}\left(1 - \frac{\delta}{(1-h_R\delta)}\right) - \delta$ and $d_{C_2}^U = \frac{a_{C_2}(1-(h_R+1)\delta)}{(1-h_R\delta) + h_{C_2} a_{C_2}(1-(h_R+1)\delta)} - \delta$ denoting the upper limits of the death rates of two different predators C_1 and C_2, respectively.

2) Invasion of C_2

The linear functional response of C_1 ensures that the $C_1 - R$ system will approach a stable equilibrium with prey density $R_{C_1}^* = \frac{d_{C_1}+\delta}{a_{C_1}}$. A necessary condition for predator coexistence is that C_2 is able to invade the $C_1 - R$ system. Hence, the invasion growth rate of C_2 for $R_{C_1}^*$ has

to be larger than zero, i.e. $\frac{1}{C_2}\frac{dC_2}{dt} = \frac{a_{C_2}R^*_{C_1}}{1+h_{C_2}a_{C_2}R^*_{C_1}} - d_{C_2} - \delta > 0$, which corresponds to

$\frac{a_{C_2}(d_{C_1}+\delta)}{a_{C_1}+h_{C_2}a_{C_2}(d_{C_1}+\delta)} - d_{C_2} - \delta > 0$. Therefore, the second necessary condition for predator

coexistence is given by:

N_2: $\qquad\qquad\qquad d_{C_2} < \frac{a_{C_2}(d_{C_1}+\delta)}{a_{C_1}+h_{C_2}a_{C_2}(d_{C_1}+\delta)} - \delta.$

This implies that the equilibrium density of the prey in the $C_2 - R$ system has to be smaller than

$R^*_{C_1}$, i.e. $R^*_{C_2} = \frac{d_{C_2}+\delta}{(a_{C_2}-(d_{C_2}+\delta)h_{C_2}a_{C_2})} < R^*_{C_1}$. In other words, the minimum prey requirement of

C_2 has to be lower than the minimum prey requirement of C_1. In contrast, when $R^*_{C_2} > R^*_{C_1}$, C_2

cannot invade the $C_1 - R$ system.

3) Competitive exclusion of C_1

Since the curvature of the type II functional response is concave, the time-averaged per-capita

net-growth rate of C_2 will always be larger than or equal to the time-averaged per-capita net-

growth rate of a corresponding predator that has a linear functional response with the same per-

capita net-growth rates at the minimum ($R_{min} = 0$), and the maximum $\left(R_{max} = R^*_S = 1 - \right.$

$\left.\frac{\delta}{(1-h_R\delta)}\right)$ feasible prey densities as C_2. Hence, the following inequality holds:

$\int \frac{1}{C_2}\frac{dC_2}{dt}dt = \int \frac{a_{C_2}R(t)}{1+h_{C_2}a_{C_2}R(t)}dt - d_{C_2} - \delta \geq \frac{a_{C_2}\int R(t)dt}{1+h_{C_2}a_{C_2}R^*_S} - d_{C_2} - \delta = \frac{a_{C_2}\bar{R}_{C_2}}{1+h_{C_2}a_{C_2}R^*_S} - d_{C_2} - \delta$

with $\bar{R}_{C_2} = \int R(t)\,dt$ being the time-averaged prey density of the $C_2 - R$ system. Since

$\int \frac{1}{C_2}\frac{dC_2}{dt}dt = 0$ at equilibrium, the following inequality holds: $\frac{a_{C_2}\bar{R}_{C_2}}{1+h_{C_2}a_{C_2}R^*_S} - d_{C_2} - \delta \leq 0.$

Setting the term $\frac{a_{C_2}}{1+h_{C_2}a_{C_2}R^*}\bar{R}_{C_2} - d_{C_2} - \delta$ to zero and solving for $\bar{R}_{C_2}$, thus gives us an upper

limit for the temporal average of the prey density in the corresponding predator-prey system

that is denoted by $\bar{R}^U_{C_2} = \frac{(d_{C_2}+\delta)}{a_{C_2}}\left(1 + h_{C_2}a_{C_2}R^*_S\right)$. Therefore, whenever C_1 has a minimum prey

requirement $R^*_{C_1}$ higher than $\bar{R}^U_{C_2}$, it will inevitably be outcompeted by C_2. Consequently, the

following inequality has to be met for the two predators to be able to coexist:

$\frac{1}{C_1}\frac{dC_1}{dt} = a_{C_1}\bar{R}^U_{C_2} - d_{C_1} - \delta > 0 \leftrightarrow a_{C_1}\frac{(d_{C_2}+\delta)}{a_{C_2}}\left(1 + h_{C_2}a_{C_2}R^*_S\right) - d_{C_1} - \delta > 0$

This leads to the following third necessary condition of predator coexistence:

N_3: $\qquad\qquad\qquad d_{C_2} > \frac{(d_{C_1}+\delta)a_{C_2}}{a_{C_1}(1+h_{C_2}a_{C_2}R^*_S)} - \delta.$

Nevertheless, $R^*_{C_1} < \bar{R}^U_{C_2}$ does not ensure that C_1 can invade the $C_2 - R$ system. For this to happen, the minimum prey requirement of C_1 has to be lower than the actual temporal average of the prey density in the corresponding predator-prey system, i.e. $R^*_{C_1} < \bar{R}_{C_2}$.

4) *Hopf-bifurcation in predator-prey models with type II functional response*

Predator coexistence through relative non-linearity in their functional responses requires oscillations in the abundances of the different species. It is well known from the theory of non-linear dynamics that limit cycles only emerge in systems of ordinary differential equations with more than 2 dimensions when there is some degree of non-linearity, i.e. non-linear density-dependence, in the per-capita rates of the corresponding state variables. Hence, in the predator-prey system defined by eq. [2], oscillations can only occur in the presence of C_2. Therefore, we performed a linear stability analysis of the two-dimensional $R - C_2$ system, in order to evaluate the influence of nutrient competition among the prey and predators, the effect of a non-linear resource uptake rate of the prey, and the impact of a shared loss rate for all species on the stability of the predator-prey dynamics. For simplicity, we will neglect the index of predator two for its state variable and parameters and thus consider the following predator-prey model:

$$\frac{dR}{dt} = \left[\frac{(1-(R+\eta C))}{1+h_R(1-(R+\eta C))} - \frac{aC}{1+haR} - \delta\right] R \qquad [B1a]$$

$$\frac{dC}{dt} = \left[\frac{aR}{1+haR} - d - \delta\right] C \qquad [B1b]$$

Throughout the analysis, we will focus on parameter values where the predator is able to persist, i.e. $d < d^U$. To start, we calculate the interior equilibrium of the prey (R^*) and predator (C^*) species for the different model parametrizations used in the main text by setting eq. [B1] to zero, i.e. $\frac{dR}{dt} = 0$ and $\frac{dC}{dt} = 0$, and solving for the corresponding values of the state variables satisfying these two conditions (for the calculated equilibria see Table B1). Subsequently, we will calculate the entries of the Jacobian matrix of the linearized predator-prey system and evaluate them at the interior equilibrium in order to describe the local behavior of the predator-prey system closely around its equilibrium state, and to determine its local stability. The stability of the interior equilibrium depends on the properties of the Jacobian matrix:

$$J = \begin{pmatrix} \frac{\partial\left(\frac{dR}{dt}\right)}{\partial R} & \frac{\partial\left(\frac{dR}{dt}\right)}{\partial C} \\ \frac{\partial\left(\frac{dC}{dt}\right)}{\partial R} & \frac{\partial\left(\frac{dC}{dt}\right)}{\partial C} \end{pmatrix}\Bigg|_{R=R^*, C=C^*}$$

The predator-prey model defined by eq. [2] (main text) does not account for interference or cooperation among individual predators or any other kind of direct density-dependence in C.

Consequently, the lower right entry of J is zero, i.e. $\partial \dot{C}/\partial C\big|_{R=R^*,C=C^*} = 0$. Furthermore, an increase in the density of the prey has always a positive effect on the predator density due to its type II functional response, which is a monotonically increasing function in R. Hence, the lower left entry of J is strictly positive, i.e. $\partial \dot{C}/\partial R\big|_{R=R^*,C=C^*} > 0$. In contrast, an increase in the predator density will always have a negative effect on the prey density through grazing or emergent nutrient competition. Therefore, the upper right entry of J is strictly negative, i.e. $\partial \dot{R}/\partial C\big|_{R=R^*,C=C^*} < 0$. Finally, the sign of the upper left entry of J, i.e. $\partial \dot{R}/\partial R\big|_{R=R^*,C=C^*}$ will depend on the actual parameter values, e.g. the death rate of the predator. Hence, the entries of J possess the following signs:

$$J = \begin{pmatrix} \mp & - \\ + & 0 \end{pmatrix}$$

According to the Ruth-Hurwitz criterium for a two-dimensional system of ordinary differential equations, the interior equilibrium is stable when the determinant of the Jacobian matrix evaluated at the interior equilibrium (J) is positive and the corresponding trace is negative (Murray 2002). The determinant of the Jacobian matrix is given by:

$$Det(J) = \left(\frac{\partial \left(\frac{dR}{dt} \right)}{\partial R} \cdot \frac{\partial \left(\frac{dC}{dt} \right)}{\partial C} \right) - \left(\frac{\partial \left(\frac{dC}{dt} \right)}{\partial R} \cdot \frac{\partial \left(\frac{dR}{dt} \right)}{\partial C} \right)$$

In line with our reasoning above, the sign of $Det(J)$ is as follows:

$$sign(Det(J)) = (\mp \cdot 0) - (+ \cdot -) = +$$

Therefore, the determinant of the Jacobian will always be positive. Consequently, the stability of the interior equilibrium entirely depends on the trace of the Jacobian matrix, which is given by:

$$Tr(J) = \frac{\partial \left(\frac{dR}{dt} \right)}{\partial R} + \frac{\partial \left(\frac{dC}{dt} \right)}{\partial C} = \mp + 0 = \mp$$

Since $\partial \dot{C}/\partial C\big|_{R=R^*,C=C^*} = 0$, the trace of the Jacobian is given by the derivative of the rate of change of the prey with respect to its own density. Hence, we calculated $\partial \dot{R}/\partial R\big|_{R=R^*,C=C^*}$ for the different model parametrizations used in the main text, and generally solved for the death rate at which the trace of the Jacobian is zero. At this particular point (d^H), a Hopf-bifurcation occurs, separating two regions for which the system exhibits either a stable equilibrium ($d > d^H$), or a limit cycle ($d < d^H$) (cf. Figure 4C, D). In general, the Hopf-bifurcation point d^H has to be calculated numerically. However, in some cases explicit expressions for $d^H_{C_2}$ can be obtained (for details see Table B1).

For instance, in the absence of nutrient competition among the predator and the prey ($\eta = 0$) and in case of a linear resource uptake rate of the prey ($h_R = 0$) and batch culture conditions ($\delta = 0$) the Hopf-bifurcation point d^H of the classical predator-prey system (Rosenzweig and MacArthur 1963) with logistically growing prey in the vacancy of the predator is given by the following (see also Abrams and Holt 2002):

$$d^H = (ah - 1)/h(ah + 1)$$

Since $d > 0$, a Hopf-bifurcation only occurs when $ah > 1$, so that d^H is also positive.

Furthermore, nutrient competition among the predators and the prey generally reduces the Hopf-bifurcation point of the corresponding $C - R$ system and thus stabilizes its population dynamics. This can be seen when choosing a particular value for the competition coefficient η between the predator and the prey, e.g. $\eta = 1$. In this case, the Hopf-bifurcation point d^H can be written as:

$$d^H = \frac{ah-1}{h(ah+1)} + \underbrace{\frac{1}{h(ah+1)}\left(1 - \sqrt{1 + \frac{(1+ah)}{a}}\right)}_{<0}.$$

The stabilizing effect of nutrient competition between predators and prey on the predator-prey dynamics is stronger, when the non-linearity of the type II functional response is weaker. In contrast to the stabilizing effect of nutrient competition among predators and prey, a non-linear resource uptake rate of the prey has a destabilizing effect on the predator-prey dynamics. When accounting for non-linearity in the preys' resource uptake rate ($h_R > 0$), the Hopf-bifurcation point of the corresponding predator-prey system is given as follows:

$$d^H = \frac{ah-1}{h(ah+1)} + \underbrace{\frac{1-\left(1+h_R\left(1+\frac{1}{ah}\right)\right)^{-\frac{1}{2}}}{h(ah+1)}}_{>0}.$$

Hence, d^H increases with an increase in h_R. The impact of h_R on the overall stability of the predator-prey dynamics depends on the curvature of the type-II functional response of the predator. The larger the product ah is, the larger the factor $\left(1 + h_R\left(1 + \frac{1}{ah}\right)\right)^{-\frac{1}{2}}$ becomes and thus, the smaller the term $1 - \left(1 + h_R\left(1 + \frac{1}{ah}\right)\right)^{-\frac{1}{2}}$ will be. Hence, the destabilizing effect of h_R on the predator-prey dynamics is larger when the non-linearity of the type-II functional response of the predator is weaker.

Finally, a common death rate of all species through the dilution rate δ is stabilizing the predator-prey dynamics. The Hopf-bifurcation point of the corresponding $C - R$ system, i.e. when $\delta > 0$, is given by the following:

$$d^H = \frac{(ah-1)}{h(ah+1)} \underbrace{-\delta\left(1 + \frac{2a}{(ah+1)((1-\delta)ah+1)}\right)}_{<0}.$$

Hence, d^H decreases when δ is increasing. The stabilizing effect of δ on the predator-prey dynamics is larger when the non-linearity of the type-II functional response of the predator is stronger, and thus, when the handling time h is higher. This happens, because the dilution rate does not only affect the stability of the predator-prey dynamics but also the persistence of the corresponding $C - R$ system.

5) Quantifying the size of the parameter range of potential species coexistence

The necessary conditions N_2, N_3 and N_4 of predator coexistence derived above determine a range of parameters that may permit the coexistence of the two different predators (Figure B1A). In order to quantify the size of this parameter region we first calculated three different threshold values of the death rate of C_1, $d_{C_1}^{T_i}$, that allows piecewise integration of the relevant functions. The first threshold value $d_{C_1}^{T_1}$ is given by the death rate d_{C_1} for which the function

$N_2(d_{C_1}) = \frac{a_{C_2}(d_{C_1}+\delta)}{a_{C_1}+h_{C_2}a_{C_2}(d_{C_1}+\delta)} - \delta$ (reflecting the second necessary condition) intersects with

the Hopf-bifurcation point $d_{C_2}^H$ (denoting the fourth necessary condition), i.e. $N_2(d_{C_1}^{T_1}) = d_{C_2}^H$ (cf. Figure B1B). Solving this equation with respect to $d_{C_1}^{T_1}$ leads to the following expression of the first threshold value:

$$d_{C_1}^{T_1} = \frac{(d_{C_2}^H+\delta)a_{C_1}+\delta(d_{C_2}^H+\delta)h_{C_2}a_{C_2}-a_{C_2}\delta}{(a_{C_2}-(d_{C_2}^H+\delta)h_{C_2}a_{C_2})}.$$

Accordingly, the second threshold value $d_{C_1}^{T_2}$ is given by the intersection point of the function

$N_3(d_{C_1}) = \frac{(d_{C_1}+\delta)a_{C_2}}{a_{C_1}(1+h_{C_2}a_{C_2}R_S^*)} - \delta$ (representing the third necessary condition) with the Hopf-

bifurcation point $d_{C_2}^H$, i.e. $N_3(d_{C_1}^{T_2}) = d_{C_2}^H$ (cf. Figure B1C). Evaluating this equation with respect to $d_{C_1}^{T_2}$ results into the following expression of the second threshold value:

$$d_{C_1}^{T_2} = \frac{a_{C_1}}{a_{C_2}}(d_{C_2}^H + \delta)(1 + h_{C_2}a_{C_2}R_S^*) - \delta.$$

Figure B1: *Parameter range, i.e. death rates d_{C_1} and d_{C_2}, that potentially enables coexistence of predator one, C_1, and predator two, C_2 (grey shaded area) (for details see legend of Figure 1). The parameter range of potential species coexistence marked in panel (A) is given by the sum of parameter ranges shown in panels (B) and (C) minus the parameter range shown in panel (D). Striped areas in panels (C)-(D) indicate parameter regions where predator coexistence is impossible.*

Finally, the third threshold value $d_{C_1}^{T_3}$ is given by the the value of d_{C_1} at which the function $N_3(d_{C_1})$ equals zero, i.e. $N_2(d_{C_1}^{T_3}) = 0$ (cf. Figure B1D). Solving the latter for $d_{C_1}^{T_3}$ results into the following expression of the third threshold value:

$$d_{C_1}^{T_3} = \delta\left(\frac{a_{C_1}}{a_{C_2}}\left(1 + h_{C_2}a_{C_2}R_S^*\right) - 1\right).$$

Using these three different threshold values, the absolute size of the parameter region of potential predator coexistence is given by the following sum of integrals (cf. Figure B1):

$$P_a = \int_0^{d_{C_1}^{T_1}} \frac{a_{C_2}(d_{C_1}+\delta)}{a_{C_1}+h_{C_2}a_{C_2}(d_{C_1}+\delta)} - \delta \; \boldsymbol{d}d_{C_1} + d_{C_2}^H \left(d_{C_1}^{T_2} - d_{C_1}^{T_1}\right) - \int_{d_{C_1}^{T_3}}^{d_{C_1}^{T_2}} \frac{(d_{C_1}+\delta)a_{C_2}}{a_{C_1}(1+h_{C_2}a_{C_2}R_S^*)} - \delta \; \boldsymbol{d}d_{C_1}$$

After having solved the first integral with *Wolfram Alpha*, we obtain the following expression of P_a:

$$P_a = \frac{a_{C_2}h_{C_2}d_{C_1}^{T_1} - a_{C_1} \, ln\left(\frac{a_{C_2}h_{C_2}d_{C_1}^{T_1}}{a_{C_1}+a_{C_2}h_{C_2}\delta}+1\right)}{a_{C_2}h_{C_2}^2} - \delta d_{C_1}^{T_1} + d_{C_2}^H \left(d_{C_1}^{T_2} - d_{C_1}^{T_1}\right) - \left(\frac{a_{C_2}\left(\frac{1}{2}d_{C_1}^{T_2}+\delta\right)d_{C_1}^{T_2}}{a_{C_1}(1+h_{C_2}a_{C_2}R_S^*)} - \right.$$

$$\left. \frac{a_{C_2}\left(\frac{1}{2}d_{C_1}^{T_3}+\delta\right)d_{C_1}^{T_3}}{a_{C_1}(1+h_{C_2}a_{C_2}R_S^*)}\right) + \delta\left(d_{C_1}^{T_2} - d_{C_1}^{T_3}\right).$$

Note, although it is not explicitly stated, P_a also depends on η_{C_2} through the dependency of the Hopf-bifurcation point $d_{C_2}^H$ on η_{C_2}. However, P_a does not depend on η_{C_1}. This strongly contrasts with the fact that the parameter range of realized species coexistence will likely also depend on η_{C_1} and potential differences between η_{C_1} and η_{C_2}.

Additional references

Murray, J. D. (2002) Mathematical Biology. I: An Introduction. Springer-Verlag, New York.

Table B1: *Summary of results of a linear stability analysis of the predator-prey model defined by eq. [B1] for the interior equilibrium under different model parameterizations, showing the influence of different ecological processes on the stability of the corresponding predator-prey dynamics. The colors highlight instances where the calculated terms of the particular expressions, i.e. equilibrium values, entrees of the Jacobian Matrix or Hopf-bifurcation point, differ from the ones of the reference scenario, i.e. the terms of the Rosenzweig-MacArthur predator-prey model (Rosenzweig-MacArthur 1963).*

Equilibria	R^*		C^*	
$\delta \geq 0; h_R > 0; \eta > 0$	$(d+\delta)/(1-(d+\delta)h)a$		$\frac{((1-\delta h_R)(1-R)-\delta)(1+haR)}{h_R \eta a} - \frac{(a(1+h_R(1-R))+(1-\delta h_R)(1+haR)\eta)}{h_R \eta a}C^* + C^{*2} = 0$	
$\delta = 0; h_R = 0; \eta = 0$	$d/a(1-dh)$		$(1-R^*)(1+ahR^*)/a$	
$\delta = 0; h_R = 0; \eta > 1$	$d/a(1-dh)$		$(1-R^*)(1+ahR^*)/((1+ahR^*)\eta + a)$	
$\delta = 0; h_R > 0; \eta = 0$	$d/a(1-dh)$		$(1-R^*)(1+ahR^*)/[a(1+h_R(1-R^*))]$	
$\delta > 0; h_R = 0; \eta = 0$	$(d+\delta)/a(1-(d+\delta)h)$		$(1-\delta-R^*)(1+ahR^*)/a$	
Entrees of the Jacobian	$\partial \dot{R}/\partial R$	$\partial \dot{R}/\partial C$	$\partial \dot{C}/\partial R$	$\partial \dot{C}/\partial C$
$\delta > 0; h_R > 0; \eta > 0$	$(1-2R-\eta C)/\left(1+h_R\left(1-(R+\eta C)\right)\right) - \frac{R}{\left(1+h_R(1-(R+\eta C))\right)^2} - \frac{aC}{(1+haR)^2} - \delta$	$-\frac{\eta R}{\left(1+h_R(1-(R+\eta C))\right)^2} - \frac{aR}{(1+haR)}$	$aC/(1+ahR)^2$	$\frac{aR}{(1+ahR)} - d - \delta$
$\delta = 0; h_R = 0; \eta = 0$	$1-2R-\frac{aC}{(1+ahR)^2}$	$-\frac{aR}{(1+ahR)}$	$aC/(1+ahR)^2$	$\frac{aR}{(1+ahR)} - d$
$\delta = 0; h_R = 0; \eta > 1$	$1-2R-\eta C-\frac{aC}{(1+ahR)^2}$	$-\eta R - \frac{aR}{(1+ahR)}$	$aC/(1+ahR)^2$	$\frac{aR}{(1+ahR)} - d$
$\delta = 0; h_R > 0; \eta = 0$	$\frac{(1-R)}{(1+h_R(1-R))} - \frac{R}{(1+h_R(1-R))^2} - \frac{aC}{(1+ahR)^2}$	$-\frac{aR}{(1+ahR)}$	$aC/(1+ahR)^2$	$\frac{aR}{(1+ahR)} - d$
$\delta > 0; h_R = 0; \eta = 0$	$1-\delta-2R-\frac{aC}{(1+ahR)^2}$	$-\frac{aR}{(1+ahR)}$	$aC/(1+ahR)^2$	$\frac{aR}{(1+ahR)} - d - \delta$

Hopf-bifurcation point	d^H
$\delta > 0; h_R > 0; \eta > 0$	$\frac{(1-R^*-\eta C^*)}{\left(1+h_R\left(1-(R^*+\eta C^*)\right)\right)} - \frac{R^*}{\left(1+h_R\left(1-(R^*+\eta C^*)\right)\right)^2} - \frac{aC^*}{(1+haR^*)^2} - \delta = 0$ (implicit expression)
$\delta = 0; h_R = 0; \eta = 0$	$\frac{(ah-1)}{h(ah+1)}$ (explicit expression)
$\delta = 0; h_R = 0; \eta = 1$	$\frac{(ah-1)}{h(ah+1)} + \frac{1}{h(ah+1)}\left(1 - \sqrt{1 + \frac{(ah+1)}{a}}\right)$ (explicit expression)
$\delta = 0; h_R > 0; \eta = 0$	$\frac{ah-1}{h(ah+1)} + \frac{1}{h(ah+1)}\left(1 - \left(1 + h_R\left(1 + \frac{1}{ah}\right)\right)^{-\frac{1}{2}}\right)$ (explicit expression)
$\delta > 0; h_R = 0; \eta = 0$	$\frac{(ah-1)}{h(ah+1)} - \delta\left(1 + \frac{2a}{(ah+1)((1-\delta)ah+1)}\right)$ (explicit expression)

Appendix C: Fluctuations in the prey abundance act as a stabilizing mechanism of predator coexistence

Contemporary theory suggests that species coexistence depends on the balance between stabilizing niche differences and destabilizing fitness differences among species (Chesson 2000; Barabás et al. 2018). Niche differences enable species coexistence by increasing intra- relative to interspecific competition, thereby facilitating the regrowth of species from low densities (Chesson 2018). In contrast, fitness differences promote competitive exclusion of inferior species in the absence of niche differentiation (Letten et al. 2017). Hence, two or more predators cannot coexist on a single prey species in the absence of niche differences, because the presence of inevitable fitness differences among the different predators will drive all but one of them extinct (Chesson 2000). Stabilizing niche differences between two different predators may arise through relative non-linearity in their functional responses and associated fluctuations in the prey abundance. We therefore here derive corresponding terms reflecting average fitness differences and stabilizing niche differences in our predator-prey model defined by eq. [2]. Following Chesson (2018), average fitness differences between the different predators can be inferred from their invasion growth rates in the absence of stabilizing niche differences. Hence, we calculated the invasion growth rates, i.e. temporal averages of the per-capita growth rates, of both predators for the corresponding resident system that comprises the other predator and the prey at its long-term equilibrium, which is assumed to be stable. Hence, the invasion growth rate of predator one for the equilibrium density of the prey in the corresponding resident predator-prey system comprising predator two is given by:

$$I_{C_1} = \frac{dC_1}{C_1 dt}\bigg|_{R^*_{C_2}} = a_{C_1} R^*_{C_2} - d_{C_1} - \delta = a_{C_1}\left(R^*_{C_2} - R^*_{C_1}\right) + \underbrace{a_{C_1} R^*_{C_1} - d_{C_1} - \delta}_{=0} = a_{C_1}\left(R^*_{C_2} - R^*_{C_1}\right)$$

Similarly, the invasion growth rate of predator two for the equilibrium density of the prey in the corresponding resident predator-prey system comprising predator one is given by:

$$I_{C_2} = \frac{dC_2}{C_2 dt}\bigg|_{R^*_{C_1}} = \frac{a_{C_2} R^*_{C_1}}{1 + h_{C_2} a_{C_2} R^*_{C_1}} - d_{C_2} - \delta = \frac{a_{C_2} R^*_{C_1}}{1 + h_{C_2} a_{C_2} R^*_{C_1}} - \frac{a_{C_2} R^*_{C_2}}{1 + h_{C_2} a_{C_2} R^*_{C_2}} +$$

$$\underbrace{\frac{a_{C_2} R^*_{C_2}}{1 + h_{C_2} a_{C_2} R^*_{C_2}} - d_{C_2} - \delta}_{=0} = \frac{a_{C_2} R^*_{C_1}\left(1 + h_{C_2} a_{C_2} R^*_{C_2}\right) - a_{C_2} R^*_{C_2}\left(1 + h_{C_2} a_{C_2} R^*_{C_1}\right)}{\left(1 + h_{C_2} a_{C_2} R^*_{C_1}\right)\left(1 + h_{C_2} a_{C_2} R^*_{C_2}\right)} =$$

$$\frac{a_{C_2}\left(R^*_{C_1} - R^*_{C_2}\right)}{\left(1 + h_{C_2} a_{C_2} R^*_{C_1}\right)\left(1 + h_{C_2} a_{C_2} R^*_{C_2}\right)}$$

Comparing the invasion growth rates of both predators reveals that the competitive outcome entirely depends on the difference between $R^*_{C_2}$ and $R^*_{C_1}$. Hence, the average fitness differences

(FD) between the two predators are given by the difference between the prey requirements of predator two $\left(R^*_{C_2}\right)$ and predator one $\left(R^*_{C_1}\right)$, i.e. $R^*_{C_2} - R^*_{C_1}$.

According to the coexistence mechanism of relative non-linearity in the predator's functional responses, stabilizing niche differences between the two different predators only arise through predator-prey oscillations. Given that the linear functional response of predator one ensures that the equilibrium of the corresponding predator-prey system is always stable, additional terms cannot arise in the invasion growth rate of predator two. However, the type-II functional response of predator two may destabilize the equilibrium of the corresponding predator-prey system. Under such conditions, the invasion growth rate of predator one is given as follows:

$$I_{C_1} = \frac{dC_1}{C_1 dt}\bigg|_{\bar{R}_{C_2}} = a_{C_1}\left(\bar{R}_{C_2} - R^*_{C_1}\right) = a_{C_1}\left(\bar{R}_{C_2} - R^*_{C_1} - R^*_{C_2} + R^*_{C_1} + R^*_{C_2} - R^*_{C_1}\right) = a_{C_1}\left(\bar{R}_{C_2} - \right.$$

$$\left. R^*_{C_2} + R^*_{C_2} - R^*_{C_1}\right)$$

Comparing the invasion growth rate of predator one for the resident system of predator two in the absence of predator-prey oscillations, which prevents coexistence, with the one in the presence of predator-prey oscillations, which potentially allows coexistence, shows that the stabilizing niche differences (ND) between the two predators are given by the difference between $\bar{R}_{C_2}$ and $R^*_{C_2}$, i.e. $\bar{R}_{C_2} - R^*_{C_2}$. Hence, the net-stabilizing effect of relative non-linearity in the predators' functional response on predator coexistence is thus given by the sum of stabilizing niche differences (+) and destabilizing fitness differences (-), i.e. $ND + FD = \bar{R}_{C_2} - R^*_{C_2} + R^*_{C_2} - R^*_{C_1} = \bar{R}_{C_2} - R^*_{C_1}$.

Appendix D: Fourier analysis of predator-prey oscillations

The population dynamics of our predator-prey model described by eq. [2] could look very different when the two predators C_1 and C_2 were able to coexist. For most parameter combinations, the population dynamics showed simple predator-prey oscillations in which the increase of the prey was followed by an increase in the two predators with some time-lag (e.g. Figure 5E). In contrast, the different predator-prey interactions generated much more complex population dynamics for some other parameter combinations (e.g. Figure 5F). To evaluate the differences in the population dynamics in more detail, we performed a discrete Fourier transformation by using the *fft* function implemented in MATLAB, version 7.13 (The MathWorks Inc., Natick, MA, 2011).

Our results show that for the simple predator-prey oscillations, most of the variation in the species' abundances can be attributed to a single dominate time scale (Figure D1A, C and E). Here, the two predators are strongly synchronized, exhibiting a negligible phase-difference of 0.03 π (cf. Figure D1B, E). In contrast, for the more complex population dynamics, our analysis reveals that there are two dominant time scales (Figure D2A, C and E), which can be most clearly seen from the amplitude spectrum of C_2 (Figure D2C). At the longer time scale, the population dynamics of the two predators exhibit a phase difference of 0.74 π (cf. Figure D2B, E). Hence, their abundances cycle almost anti-synchronously, meaning that an increase in the abundance of C_2 coincides with a decrease in the abundance of C_2 and *vice versa* (Figure D2E). In contrast to the prey (R) and C_2, C_1 showed hardly any variation in its abundance at the shorter time scale (Figure D2F) which can be explained by the fact that the shorter time scale in the population dynamics is generated by the predator-prey interaction between R and C_2. In fact, the high frequency component reflects the predator-prey oscillations that potentially emerge in the $R - C_2$ system in the absence of C_1.

Figure D1: *Frequency Domain and Time Domain of the population dynamics showing simple predator-prey oscillations (cf. Figure 4C). Amplitude spectrum (grey) of the time series of the prey (R), predator one (C_1) and predator two (C_2) are shown in panel A, C and E, respectively. The black vertical line indicates the dominate time scale of the population dynamics. Panel B shows the full time series of R (black solid line), C_1 (grey solid line) and C_2 (grey dashed-dotted line). The population dynamics at the dominant time scale are plotted in panel D based on the calculated discrete Fourier transform.*

Figure D2: *Frequency Domain and Time Domain of the population dynamics showing complex predator-prey oscillations (cf. Figure 4D). Amplitude spectrum (grey) of the time series of the prey (R), predator one (C_1) and predator two (C_2) are shown in panel A, C and E, respectively. The black vertical lines indicate the two dominate time scales of the population dynamics. Panel B shows the full time series of R (black solid line), C_1 (grey solid line) and C_2 (grey dashed-dotted line). The population dynamics at the longer and shorter time scales are plotted in panel D and F, respectively, based on the calculated discrete Fourier transform*