Genome-Scale reconstruction of *Paenarthrobacter aurescens* TC1 metabolic model

towards the study of atrazine bioremediation

Shany Ofaim[1,2,¥], Raphy Zarecki[1,¥], Seema Porob[3], Daniella Gat[3], Tamar Lahav[1],

Yechezkel Kashi[2], Radi Aly[1], Hanan Eizenberg[1], Zeev Ronen[3,†] & Shiri Freilich*[†]

[1]Newe Ya'ar Research Center, Agricultural Research Organization, Ramat Yishay,

Israel, [2]Faculty of Biotechnology and Food Engineering, Technion-Israel Institute of

Technology, Haifa, Israel, [3]Department of Environmental Hydrology & Microbiology,

Zuckerberg Institute for Water Research, Jacob Blaustein Institutes for Desert

Research, Ben-Gurion University of the Negev, Midreshet Ben-Gurion, 8499000,

Israel

[¥]equal contribution

[†] equal contribution

* Corresponding author

email   address   (in   the   order   of   author   list):   shashany@gmail.com,

raphy.zarecki@gmail.com,   porob@post.bgu.ac.il,   gat.daniella@gmail.com,

lahav008@gmail.com,   kashi@technion.ac.il,   radi@volcani.agri.gov.il,

eizenber@volcani.agri.gov.il, zeevrone@bgu.ac.il, shiri.freilich@gmail.com.

ABSTRACT

Atrazine is an herbicide and pollutant of great environmental concern that is naturally

biodegraded by microbial communities. *Paenarthrobacter aurescens* TC1 is one of the

most studied degraders of this herbicide. Here, we developed a genome scale

metabolic model for *P. aurescens* TC1, *i*RZ1176, in order to study the atrazine

degradation process and mechanism at organism level. Constraint based flux balance

analysis and time dependent simulations were used to explore the bacteria's

phenotypic landscape. Simulations led to model based predictions for growth and

atrazine degradation efficiency in myriad of media supplemented with different

combinations of carbon and nitrogen sources. Simulations aimed at designing media

optimized for supporting growth and enhancing degradation, bypassing the need in

strain design *via* genetic modifications. Predictions were tested *in vitro* and

experimental growth and degradation measurements support and validate the

reliability of the model. Model simulations have prioritized degradation and growth

enhancement effect of various amino acid and co-factors. The model presented here

can potentially serve as a predictive tool towards achieving optimal biodegradation

efficiencies and for the development of ecologically friendly solutions for pollutant

degradation.

Introduction

Atrazine (2-chloro-4-ethylamino-6-isopropylamino-1,3,5-triazine) is an herbicide and

pollutant of great environmental concern. Its residues are found in soil samples

decades after it was last applied and were shown to chronically leach into local

aquifers[1]. Areas contaminated with hazardous materials, including atrazine, are

rapidly increasing worldwide introducing a need in remediation approaches.

Bioremediation - an environmental bioprocess in which naturally occurring organisms

are used for breaking down hazardous substances into less toxic or non-toxic

substances - is increasingly acknowledged as a cost-effective strategy for

environmental cleaning[2-4]. Critical environmental bioprocesses are naturally carried

out by bacteria and are related to removal of pollutants from water, soil or air[5]. Thus,

bacterial bioremediation is widely applied for the degradation of various organic

pollutants [6-9] including herbicides[9-13]. Bioremediation may occur naturally or through

the stimulating addition of fertilizers, electron acceptors, etc. These amendments

encourage the growth of the degrading microbes within the medium or environment

in a process termed biostimulation [3,14].

One of the well-known atrazine degraders is the gram positive *Paenarthrobacter*

aurescens TC1[15-19]. *P. aurescense* TC1 is described as nutritionally versatile bacterium

that is capable of utilizing a wide array of complex metabolites, including agriculturally

used herbicides[4,20,21]. The full hydrolytic pathway for atrazine degradation was

reported and described in *P. aurescens* TC1[21]. Moreover, This bacterium was

previously reported to biodegrade atrazine much more efficiently than most atrazine

degrading bacteria including *Pseudomonas* sp. ADP [22]. Intermediates, created in the

degradation process are further metabolized by the organism and can provide sole

sources for carbon and nitrogen[18]. While *P.*ADP can only use the hydrolytic products

of the ring cleavage as nitrogen but not as carbon; TC1 can use the side groups of

atrazine as source for carbon, nitrogen and energy consuming up to 3,000 mg of

atrazine per liter [18]. As such, *P. aurescens* TC1 is an ideal candidate for the study of

atrazine bioremediation using a genome scale metabolic model reconstruction.

Modern tools of genomics, transcriptomics, metabolomics, proteomics, signaling systems and synthetic biology have opened new avenues for biotechnological advances and are increasingly applied for promoting better understanding of complex biological systems. In recent years, constraint based metabolic modelling approaches have become widely used as an *in silico* tool for organism-level phenotyping and the subsequent development of metabolic engineering strategies [23,24]. Generally, such approaches follow four key steps: (1) data acquisition – mainly genome sequencing information, basic cell-physiological and biochemical knowledge and some experimental data on cell growth; (2) model reconstruction - translating data into structured mathematical representation; (3) constraint-based optimization simulations - the prediction of growth rate, substrate uptake rates, and byproduct rates under different growth conditions or following knockout mutations, in the absence of kinetic information[25-31]; and (4) experimental validation. The potential of such models for the investigation of optimal processing is now acknowledged and practiced[5,32-35]. Examples for applicative use include the optimal production or utilization of industrial compounds such as xylose[36], biofuels [37],vitamins [38] and drug development[39]. In bioremediation, genome scale metabolic modeling approaches were applied for the design of *Geobacter sulfurreducens* strain capable of increased electron transfer and a higher Fe(III) respiratory that was beneficial for environmental bioremediation[40]. Application of genome scale metabolic modeling for strain design is often based on the Optknock algorithm [41] and its derivatives which search for the optimal gene knockouts for a desired metabolic production. However, unlike industrial use of engineered strain for enhanced production in monoculture, introduction of exogenous species into natural habitat is far from trivial and hampers

the application of modeling approaches towards bioremediation solutions. An

alternative strategy is the use of modeling for the design of optimal conditions that

will enhance processes (degradation or synthesis) carried by the endogenous

community[42]. For example, interactions between *Geobacter* and *Rhodoferax* – a

competitor Fe(III)-reducers in anoxic subsurface environments - were captured and

described using metabolic modelling and allowed identification of optimal

degradation conditions[43]. Models can also predict the benefits of cooperative and

commensal interactions to degradation efficiency. Xu *et al* have demonstrate the use

of metabolic models for predicting interactions between *P.aurescens* TC1 and

endogenous soil species. Exchange metabolites secreted by non-degraders species

contribute for enhancing *P.aurescens* TC1 growth and expediting degradation

efficiency[44,45]. Here, we directly focused on modeling-based identification of media

supplements that can enhance degradation. We describe the process of manual

curation of *P.aurescens* TC1 draft metabolic model, its application for predicting

growth and atrazine degradation in media containing different carbon and nitrogen

sources based on a dynamic (capable of reporting changes over time) and non-

dynamic flux based algorithms, and the experimental testing of the predictions.

Results

P.aurescens TC1 genome scale metabolic network reconstruction

The genome of *P.aurescens* TC1 is composed of a circular chromosome of 4.6 Mb

coding for 4,222 open reading frames (ORFs) as well as two plasmids, 0.3 Mb each,

coding for additional ~600 ORFs[20]. To capture the most comprehensive metabolic

capacities encoded in genome and plasmids, a genome-scale metabolic model was

reconstructed. To this end, the genome and plasmids were annotated and used to

create a draft reconstruction using the Model SEED pipeline[46]. Additional genome

annotations of *P.aurescens* TC1 from four established and comprehensive public

resources and databases (RAST, KEGG, JGI and UniProt) were then added to the draft

network. Model Reactions involved in atrazine degradation pathway and its further

conversion into the cellular building blocks isopropylamine and ethylamine were

added manually (Fig. 1) based on a detailed reports of the pathway in the specific

strain[18,47]. *P.aurescens* TC1 degrades atrazine by hydrolytic dechlorination. The

process is catalyzed by triazine hydrolase (TrzN) followed by two hydrolytic

deamination reactions catalyzed by hydroxyatrazine hydrolase (AtzB) and N-

isopropylammelide isopropylaminohydrolase (AtzC) (reactions rxn06933, rxn03845

and rxn03846 in Fig. 1, respectively). These enzymes convert atrazine to cyanuric acid

and release the alkylamines ethylamine and isopropylamine. Cyanuric acid

accumulates stoichiometrically, whereas the alkylamines are further catabolized into

cellular building blocks[21]. Ethylamine and isopropylamine are converted into

acetaldehyde and L-alanine (Fig. 1), respectively, and can serve as sole sources of

carbon and nitrogen required for growth [18]. Relevant reactions that were missing in

the automatic reconstruction were manually added to the model and are shown in

Fig. 1. Simulations were carried iteratively to ensure the production of all biomass

components under the minimal feasible conditions described in Strong, et al. [18]. The

reconstructed metabolic network presented here covers 25% of the ORFs present in

the genome and includes 1176 genes, 2541 reactions, out of which 125 are exchange,

and 2848 metabolites. The coverage of the genome is similar to those of previously

published reconstructed metabolic networks of gram positive bacteria such as *C.*

difficile or *B. subtilis* with 20% [48,49] and the overall number of reactions is similar to the

number of reactions covered by gold standard, updated models of species with similar

genome sizes such as *i*JO1366 [50]. A detailed description of the network including the

reactions, metabolites, genes, and compartments that comprise the network is

provided in Supplemental data 1. The model is also available here as Systems Biology

Markup Language (SBML) file [51] in Supplemental data 2. The SBML file can be used

with tools such as MATLAB or other SBML compliant software. The minimal media

used for simulations are available in Supplemental data 3.

<u>Validations of *i*RZ1176's *in silico* predictions</u>

As previously reported, *P.aurescens* TC1 degrades atrazine and metabolizes its

intermediates, isopropylamine and ethylamine. Both can be used as sole carbon and

nitrogen sources. Growth simulations confirm that the model captures these reported

physiological capacities.   Experimental growth measurements verify model

simulations and show that atrazine and its degradation products isopropylamine and

ethylamine can be used as sole carbon and nitrogen sources for *P. aurscens* TC1 (Fig.

2A). Supplementing the medium with glucose enhanced growth both *in silico* and *in*

vitro, while the addition of ammonia had not shown an effect on growth. This is in

accordance with previous work stating that *P.aurescens* TC1 growth is not inhibited

by nitrogen sources such as ammonia [22], unlike the inhibition of atrazine degradation

in *Pseudomonas sp. strain* ADP in nitrogen rich environments [52].

Finally, growth simulations under a gradient of nitrogen *vs*. carbon   sources further

illustrate that atrazine serves as a good nitrogen source and as a poor carbon source

(Fig. 2B). The growth promoting potential of combinations of atrazine with ammonium

(NH_4^+) and glucose as nitrogen and/or carbon sources respectively was tested. The

effect of nutrient ratio on growth rate was predicted by optimizing the production of

biomass across uptake gradients while constraining the fluxes through exchange

reactions for carbon and nitrogen sources. As expected, increasing glucose uptake

increased the growth rate (Fig. 2B) and the maximal achievable growth rate was much

higher with glucose as a carbon source in comparison to atrazine as a sole carbon

source. However, atrazine is as equally good nitrogen source as ammonia. This may

be endorsed by ammonium as a byproduct of the atrazine degradation process (Fig.

1).

<u>Media modifications effect on *P.* aurescens TC1 growth</u>

In addition to glucose, several carbon sources were previously reported to enhance

growth of several strains of *Arthrobacter* species in atrazine containing media [21,22,53].

To identify additional media modifications that can potentially increase atrazine

degradation, we simulated growth in 123 different media combinations, each

supplementing the atrazine containing minimal mineral media with a single exchange

metabolite. Several sugars, amino acids and dipeptides were predicted to induce

enhanced growth (Supplemental data 4). This is in accordance with the previous

characterization of *P.aurescens* TC1 as a metabolically versatile species [20,21]. Out of a

variety of potential nutritional sources, *Paenarthrobacter* species have been shown to

utilize different amino acids in different capacities [54,55]. The three amino acids

isoleucine, histidine and methionine are predicted to support growth in high, mediate,

and low efficiency, respectively (Supplemental data 4). Dynamic simulations of growth supplementing media with these three amino acids indicated that their effect is media dependent (Fig. 3). Isoleucine has the strongest predicted effect on growth in atrazine containing medium whereas histidine has stronger effect in the same medium without atrazine; methionine has the weakest effect in all media (Fig. 3). *P.aurescens* TC1 growth in *in vitro* media corresponding to the media used for the *in silico* simulations were generally in agreement with model's predictions, supporting the superiority of isoleucine over histidine in atrazine supplemented media but not in the atrazine deficient medium (Fig. 3). Supplementing media with methionine showed a weaker effect on growth in comparison to the other two amino acids tested, agreeing with its weak effect in simulations. This is also in agreement with a recent work by Deutch [55] showing that whereas isoleucine and histidine serve as potential nitrogen and/or carbon sources of *P.aurescens* TC1, methionine does not serve as a growth supporter. Like atrazine, amino acids can potentially serve as both carbon and nitrogen source. Considering C:N ratio as an indicator of growth support potential, isoleucine and histidine have a ratio of 6:1 and 6:3, respectively, whereas methionine, having no growth promoting effect has a ratio of 5:1 (Supplemental data 4). Other amino acids with 5:1 or 5:3 C:N ratio like proline and serine, respectively, are also predicted to be an efficient supporters of growth, similar to isoleucine (Supplemental data 4). Thus, the growth effect of atrazine cannot be predicted merely based on C:N ratio.

Media modifications effect on atrazine degradation

Following the validation of the model predictions for growth capacities, we directly evaluated atrazine degradation in different media conditions. To this end, we applied

Flux variability analysis (FVA), allowing the identification of the ranges of flux

variability that are possible within a given media[56]. Once the biomass flux was

maximized, we estimated the rate of atrazine degradation according to the relevant

flux of interest (for example, phosphate). The simulation of atrazine degradation over

time was in a medium that contains atrazine as a sole carbon and nitrogen source *vs.*

modified media sets (see Methods). Atrazine degradation was estimated according to

the amount of atrazine left in the media following each time cycle. Simulations

showed that supplementing the media with glucose not only enhances growth (Fig. 2)

but also induced an enhancement of atrazine consumption. Predictions were

confirmed by experimental validation (Fig. 4).

Finally, the addition of phosphate, a main element in fertilizers, to soils has been

reported to have a significant effect on the degradation of herbicides. For instance, its

addition was shown to effect soil bacteria's degradation activity including a reported

effect on *Arthrobacter* sp. HB-5, a close relative of *P.aurescens* TC1[57]. Phosphate has

been previously shown to accelerate atrazine degradation[3]. To test the effect of

phosphate, we used the model to simulate growth and atrazine degradation in high

or limited phosphate conditions. In accordance with these reports, simulations

showed similar trends, which were confirmed by experimental results. These indicate

an effect of phosphate concentrations *on P. aurescens* TC1 growth (Supplemental data

5) and atrazine degradation (Fig. 4).

Discussion

The accumulation and persistence of herbicides and their intermediates in soil and

drinking water has long been suggested to be disruptive, imposing a need for effective

strategies for their degradation and removal [58]. Bioremediation is a widely applied

approach for the degradation of herbicides including atrazine [9-13]. Here, we

reconstructed a genome scale metabolic model, *i*RZ1176, of *P. aurescnes* TC1, a well-

studied atrazine degrader, in order to use the *in silico* simulations for predicting

optimal media to support its growth and atrazine degradation activity. The *in silico*

simulations were carried on an array of minimal media using atrazine and its interim

degradation products as alternative sole carbon and nitrogen sources supplemented

by potential enhancers. The simulations suggested a group of metabolites promoting

the increase in biomass yield including sugars, amino acids and dipeptides. We

hypothesize that the introduction of such an efficient carbon source leads to an

increase in biomass and a subsequent increase in the degradation activity. Simulations

were confirmed by *in vitro* experimental validation. This genome scale reconstruction

may be used for further experimenting and the gain of new biological knowledge of

this metabolically versatile bacteria.

While the metabolic modeling approach presented here was demonstrated useful for

generating predictions, its limitations should be acknowledged. First, model

construction relies on automatic protocols and publically free genomic information.

Biomass composition and energy maintenance requirements are hence

approximations [46,48]. Second, data used to construct the model is eclectic and

annotations were gathered from numerous databases. The lack of data

standardization has led to many overlaps and misnomers resulting in draft models

containing massive gaps. The lion share of the gap filling and network improvement

processes are still done manually due to the differences in data description and

cataloging in the different databases. As technology and standardization advance, the

process of constructing a metabolic model able to produce testable predictions will be faster and have greater accuracy. Despite relying on genomic data and a growth approximation, resulting predictions can successfully capture the relative effect of different media complements on growth and atrazine degradation. Accuracy of predictions is likely to be significantly improved by the addition of other layers of information including metabolomics, transcriptomics and reaction kinetics. These additional layers of information will allow imposing additional constrains on the internal and external fluxes, as well as an accurate description of the biomass composition specific to this strain. Continuation of the manual curation process, also based on added transcription information, is likely to improve the performances of future versions.

Overall, we describe the construction of a mathematical stoichiometric metabolic model of *Paenarthrobacter aurescens* TC1 (*i*RZ1176), allowing simulating its degradation activity. Simulations were used to predict growth and degradation following media modifications such as high/low phosphate content and the addition of variety of amino acids and carbon source to the standard minimal R medium. Effects of such media were observed experimentally to effect atrazine degradation. The simulations go beyond the C:N ratio of media components and take into account the complement of reactions and stoichiometric complexity of cellular processes and may be used to further study this versatile organism for the design of improved degradation strategies in soil. Additionally, we demonstrate that through using metabolic time dependent simulations, an optimal range of nutrient augmentation may also be planned. Though soil communities are complex in terms of their composition and microbial consortia dynamics, in a recent study we demonstrated the

usefulness of modeling approaches for deciphering metabolic complexities[44], pointing

at the future potential of modeling approaches. Despite the complexity of soil

communities, supplementing soil with carbon sources is long known to induce a

change in microbial structure and function. A predictive tool such as the presented

model is a first step towards designing optimal degradation efficiencies that can be

further tested in the field. The integration of species-specific models together with

community approaches is likely to contribute to our understanding of biodegradation

processes in natural environments and promote the development of bioaugmentation

and biostimulation strategies.

Methods

<u>Reconstruction and curation of the *P. aurescens* TC1 metabolic model</u>

An initial draft model of *P. aurescens* TC1 was reconstructed as described in Henry, et

al. [46]. Briefly, the complete genome sequence of *P. aurescens* TC1 was retrieved from

NCBI (GenBank accession ids. NC_008711 including NC_008711.1 NC_008712.1, and

NC_008713.1 – chromosome and two plasmids, respectively), uploaded into RAST [59],

spliced into genes and annotated. RAST annotations were put through the Model SEED

pipeline [46] and an initial metabolic network reconstruction was obtained. This initial

reconstruction contained a list of reactions classified as exchange, transport, and

cytosolic as well as a list of all relevant metabolites and a biomass reaction. Based on

taxonomy, the gram positive biomass reaction was specified in the process of draft

reconstructions using Model SEED and KBase pipelines. The composition of the

biomass reaction summarizes the fractional contribution of a generalized microbial

biomass precursors (e.g., amino acids and lipids) to the synthesis of a new cell and is

similar to the previously published genome scale reconstruction of *B.Subtilis*[48].

Unspecified non-metabolic consuming processes in the cell were defined as ATP

maintenance and set similarly to [48]. Missing reactions were filled through an automatic

gap-filling procedure, using default (optimal) media options [46]. Manual curation

procedures were carried to adapt model performances to reported physiological

capacities. Simulation conditions were modified to reflect experimentally validated

minimal media alternatives that support growth of *P.* aurescnes TC1 [18].

Curation procedures were carried iteratively versus growth simulations to ensure the

production of all biomass components in Minimal Mineral Media (MMM: K^+, Mn^{2+},

CO2, Zn^{2+}, SO_4^-, Cu^{2+}, Ca^{2+}, HPO_4^{2-}, Mg^{2+}, Fe^{2+}, Cl^-), supplemented by alternative carbon

and nitrogen sources including atrazine, ethylamine, isopropylamine, ammonia,

glucose and selected amino acids. As part of the curation process, the hydrolytic

pathway of atrazine degradation was *a priori* added to the model based on previously

published biochemical evidence [18]. Reaction directionality was assigned based on the

KEGG database. Errors in reaction reversibility and elemental balance were manually

identified and corrected according to literature and/or available databases. The model

is available in SBML format and as excel sheet (Supplemental data 1 & 2). The minimal

media used for simulations are available in Supplemental data 3.

<u>Flux balance analyses simulations and dynamic, time dependent simulations</u>

Flux distributions in the metabolic model were determined using Flux Balance Analysis

(FBA) [60-62]. Briefly, a metabolic model includes a network that is described as a

stoichiometric matrix $\mathbf{S}_{m \times n}$, where *m* represents the number of metabolites and *n* the

number of reactions in the model. In the assumed pseudo-steady-state, internal

metabolites concentrations and fluxes are assumed to be constant. The model can be
represented as: $\mathbf{S} \times v = 0$, where vector v signifies the fluxes through the internal
reactions. As the number of reactions (*n*) typically exceeds the number of metabolites
(*m*), the system is under-determined. In order to get a feasible solution space, linear
programming (LP), subject to mass balance preservation and flux constraints, was
used by introducing an optimization problem. Here, model growth was maximized by
maximizing its biomass reaction, which was used as objective function. Specific fluxes
for metabolites consumption and secretion were determined following fixation of the
respective biomass reaction to its maximal value and then using Flux Variability
Analysis (FVA)[56], a linear programming method to find the minimal amount of
metabolites needed to produce a preset amount of biomass. Dynamic modeling was
used for the prediction of the profile of consuming metabolites typical to the biomass
increase of *P. aurescens* TC1 and atrazine degradation across time. To this end, we
simulated the behavior of our metabolic model across time. This was done as
described in [44] and illustrated in Supplemental data 5, based on concepts defined in
[63]. Briefly, the model works under the following assumptions: (1) A finite start amount
of media components is available; (2) A maximal amount of uptake a single cell can
acquire from the media in a given time point is defined (the lower bound of the
exchange reaction value); (3) New substrate concentrations in time t, are determined
by the predicted substrate concentration for the previous step augmented with any
additional substrates provided or consumed in the current iteration. The maximum
uptake was set to a ratio of up to 1 unit of each metabolite available in the media; (4)
After each time tick, the biomass amount was updated according to the flux amount
of the biomass reaction in the model at this time tick. As biomass production rate

serves as a proxy for the size of the population in the simulated environment and substrate uptake/secretion is mainly affected by to population size, the model was used to evaluate the actual substrate uptake/secretion and growth rate given the supplied media across time. Simulations were carried until reaching a state where time cycles did not lead to an increase in biomass. Initial concentration values for all metabolites were set to a fixed amount of 50 units (represented as the initial lower bounds, LB, of the exchange reactions). Simulation parameters and defaults are available in Supplemental data 3. Starting with one bacterium, the flux balance model was used to predict the uptake of phosphate, sulfate, carbon and nitrogen sources (*i.e.* atrazine and glucose) by *P. aurescens* TC1 across time.

All model simulations were done on an Intel i7 quad-core server with 32GB of memory, running Linux. The development programming language of our simulators was JAVA, and our linear programming software was IBM CPLEX. Surface plots were created using the MATLAB 2019a (MathWorks, Inc., Natick, Massachusetts, United States). Pathway maps were created using Escher maps[64].

In vitro bacterial growth conditions and chemical analysis

P.aurescens TC1 (ATCC BAA-1386™) was grown on R medium (based on ATCC 2662 R) plates with glucose (0.2%) and atrazine (30 mg/l, Agan Adama -Ashdod, Israel Israel) as sole nitrogen. Stock cultures were maintained in glycerol at -80°C. Cultures were resuscitated on R medium agar plates (1.5% Difco bacto™) and were used as starter colonies for liquid medium growth experiments. The growth of *P.aurescens* TC1 was tested in R media supplemented with different combinations of carbon and nitrogen

sources, including Glucose (0.2%), Atrazine (30 mg/l, Agan Adama -Ashdod, Israel),

ammonium chloride (37 mg/l), Ethylamine and Isopropylamine (31.1 and 40.8 mg/l,

Sigma-Aldrich -Rehovot, Israel) and the amino acids Histidine, Isoleucine and

Methionine (36, 92 and 103 mg/l respectively). Concentrations of amino acids,

ethylamine and Isopropylamine were calculated to provide the same nitrogen content

as the atrazine. In addition to the carbon and nitrogen sources, growth in media

supplemented by atrazine and atrazine together with glucose was tested under two

phosphate concentrations (K_2HPO_4 in 0.45 and 0.045 g/l). Each substrate was added

to sterile R medium and bacteria was grown in triplicate 250ml flasks incubated in

30°C with shaking in the dark. One flask contained only R medium and was used as

control. A 200 ul sample was taken from each flask every 8 hours for measurement of

OD in microplate reader (520nm, Infinite® 200 PRO Tecan, Männedorf Switzerland)

and 1 ml was taken to study atrazine degradation rate using HPLC. The HPLC analysis

was performed with Agilent 1100 HPLC (Waldbronn Germany) equipped with a DAD

detector. Samples were separated on Kinetex C18 column (Phenomenex Torrance, CA)

and the mobile phase consisted of 70% methanol flowing at 1 ml/min. Detection and

quantitation of atrazine was done at 240 nm with detection limit of 0.1 mg/l. Total

duration of each growth experiment was between 152 to 168 hours.

References

1   Abubucker, S. *et al.* Metabolic reconstruction for metagenomic data and its
application to the human microbiome. *PLoS Comput Biol* **8**, e1002358,
doi:10.1371/journal.pcbi.1002358 (2012).
2   Kastner, M. & Miltner, A. Application of compost for effective bioremediation of
organic contaminants and pollutants in soil. *Appl Microbiol Biotechnol* **100**, 3433-
3449, doi:10.1007/s00253-016-7378-y (2016).

Kanissery, R. G. & Sims, G. K. Biostimulation for the Enhanced Degradation of Herbicides in Soil. *Applied and Environmental Soil Science* **2011**, 10, doi:10.1155/2011/843450 (2011).

Viegas, C. A. *et al.* Evaluating formulation and storage of Arthrobacter aurescens strain TC1 as a bioremediation tool for terbuthylazine contaminated soils: Efficacy on abatement of aquatic ecotoxicity. *Sci Total Environ* **668**, 714-722, doi:10.1016/j.scitotenv.2019.02.355 (2019).

Perez-Garcia, O., Lear, G. & Singhal, N. Metabolic Network Modeling of Microbial Interactions in Natural and Engineered Environmental Systems. *Front Microbiol* **7**, 673, doi:10.3389/fmicb.2016.00673 (2016).

Nilsson, T., Rova, M. & Smedja Backlund, A. Microbial metabolism of oxochlorates: a bioenergetic perspective. *Biochim Biophys Acta* **1827**, 189-197, doi:10.1016/j.bbabio.2012.06.010 (2013).

Mani, D. & Kumar, C. Biotechnological advances in bioremediation of heavy metals contaminated ecosystems: an overview with special reference to phytoremediation. *International Journal of Environmental Science and Technology* **11**, 843-872, doi:10.1007/s13762-013-0299-8 (2013).

Shah, J. & Dahanukar, N. Bioremediation of organometallic compounds by bacterial degradation. *Indian J Microbiol* **52**, 300-304, doi:10.1007/s12088-011-0223-1 (2012).

Megharaj, M., Ramakrishnan, B., Venkateswarlu, K., Sethunathan, N. & Naidu, R. Bioremediation approaches for organic pollutants: a critical perspective. *Environ Int* **37**, 1362-1375, doi:10.1016/j.envint.2011.06.003 (2011).

Nousiainen, A. O. *et al.* Bioremediation strategies for removal of residual atrazine in the boreal groundwater zone. *Appl Microbiol Biotechnol* **99**, 10249-10259, doi:10.1007/s00253-015-6828-2 (2015).

Sagarkar, S. *et al.* Soil mesocosm studies on atrazine bioremediation. *J Environ Manage* **139**, 208-216, doi:10.1016/j.jenvman.2014.02.016 (2014).

Fan, X. & Song, F. Bioremediation of atrazine: recent advances and promises. *Journal of Soils and Sediments* **14**, 1727-1737, doi:10.1007/s11368-014-0921-5 (2014).

Singh, B. & Singh, K. Microbial degradation of herbicides. *Crit Rev Microbiol* **42**, 245-261, doi:10.3109/1040841X.2014.929564 (2016).

Adams, G. O., Fufeyin, P. T., Okoro, S. E. & Ehinomen, I. Bioremediation, biostimulation and bioaugmention: a review. *International Journal of Environmental Bioremediation & Biodegradation* **3**, 28-39 (2015).

Wang, Q. & Xie, S. Isolation and characterization of a high-efficiency soil atrazine-degrading Arthrobacter sp. strain. *International Biodeterioration & Biodegradation* **71**, 61-66, doi:10.1016/j.ibiod.2012.04.005 (2012).

Fang, H., Lian, J., Wang, H., Cai, L. & Yu, Y. Exploring bacterial community structure and function associated with atrazine biodegradation in repeatedly treated soils. *Journal of hazardous materials* **286**, 457-465, doi:10.1016/j.jhazmat.2015.01.006 (2015).

Zhang, Y. *et al.* Metabolic ability and individual characteristics of an atrazine-degrading consortium DNC5. *J Hazard Mater* **237-238**, 376-381, doi:10.1016/j.jhazmat.2012.08.047 (2012).

Strong, L. C., Rosendahl, C., Johnson, G., Sadowsky, M. J. & Wackett, L. P. Arthrobacter aurescens TC1 metabolizes diverse s-triazine ring compounds. *Appl Environ Microbiol* **68**, 5973-5980 (2002).

Sagarkar, S. *et al.* Monitoring bioremediation of atrazine in soil microcosms using molecular tools. *Environ Pollut* **172**, 108-115, doi:10.1016/j.envpol.2012.07.048 (2013).

20   Mongodin, E. F. *et al.* Secrets of soil survival revealed by the genome sequence of
Arthrobacter aurescens TC1. *PLoS genetics* **2**, e214,
doi:10.1371/journal.pgen.0020214 (2006).
21   Deutch, C. E., Bui, A. P. & Ho, T. Growth of Paenarthrobacter aurescens strain TC1 on
atrazine and isopropylamine during osmotic stress. *Annals of Microbiology* **68**, 569-
577, doi:10.1007/s13213-018-1364-9 (2018).
22   Silva, V. P. *et al.* Evaluation of Arthrobacter aurescens Strain TC1 as Bioaugmentation
Bacterium in Soils Contaminated with the Herbicidal Substance Terbuthylazine. *PloS*
one **10**, e0144978 (2015).
23   Xu, C. *et al.* Genome-scale metabolic model in guiding metabolic engineering of
microbial improvement. *Appl Microbiol Biotechnol* **97**, 519-539, doi:10.1007/s00253-
012-4543-9 (2013).
24   Contador, C. A., Rodriguez, V., Andrews, B. A. & Asenjo, J. A. Genome-scale
reconstruction of Salinispora tropica CNB-440 metabolism to study strain-specific
adaptation. *Antonie Van Leeuwenhoek* **108**, 1075-1090, doi:10.1007/s10482-015-
0561-9 (2015).
25   Mahadevan, R. & Lovley, D. R. The degree of redundancy in metabolic genes is linked
to mode of metabolism. *Biophys J* **94**, 1216-1220, doi:10.1529/biophysj.107.118414
(2008).
26   Mahadevan, R. *et al.* Characterization of metabolism in the Fe(III)-reducing organism
Geobacter sulfurreducens by constraint-based modeling. *Appl Environ Microbiol* **72**,
1558-1568, doi:10.1128/AEM.72.2.1558-1568.2006 (2006).
27   Oberhardt, M. A. *et al.* Harnessing the landscape of microbial culture media to predict
new organism-media pairings. *Nat Commun* **6**, 8493, doi:10.1038/ncomms9493
(2015).
28   Lee, J., Yun, H., Feist, A. M., Palsson, B. Ø. & Lee, S. Y. Genome-scale reconstruction
and in silico analysis of the Clostridium acetobutylicum ATCC 824 metabolic network.
Appl Microbiol Biotechnol **80**, doi:10.1007/s00253-008-1654-4 (2008).
29   Thiele, I., Vo, T. D., Price, N. D. & Palsson, B. O. Expanded metabolic reconstruction of
Helicobacter pylori (iIT341 GSM/GPR): an in silico genome-scale characterization of
single- and double-deletion mutants. *J Bacteriol* **187**, 5818-5830,
doi:10.1128/JB.187.16.5818-5830.2005 (2005).
30   Price, N. D., Papin, J. A., Schilling, C. H. & Palsson, B. O. Genome-scale microbial in
silico models: the constraints-based approach. *Trends Biotechnol* **21**, 162-169,
doi:10.1016/S0167-7799(03)00030-1 (2003).
31   Reed, J. L., Vo, T. D., Schilling, C. H. & Palsson, B. O. An expanded genome-scale model
of Escherichia coli K-12 (iJR904 GSM/GPR). *Genome Biol* **4**, R54, doi:10.1186/gb-2003-
4-9-r54 (2003).
32   Fang, Y. *et al.* Direct coupling of a genome-scale microbial in silico model and a
groundwater reactive transport model. *J Contam Hydrol* **122**, 96-103,
doi:10.1016/j.jconhyd.2010.11.007 (2011).
33   Zomorrodi, A. R. & Segre, D. Synthetic Ecology of Microbes: Mathematical Models and
Applications. *J Mol Biol* **428**, 837-861, doi:10.1016/j.jmb.2015.10.019 (2016).
34   Biswas, K., Paul, D. & Sinha, S. N. Biological Agents of Bioremediation: A Concise
Review. *Microbiology* **1**, 39-43 (2015).
35   Henson, M. A. Genome-scale modelling of microbial metabolism with temporal and
spatial resolution. *Biochem Soc Trans* **43**, 1164-1171, doi:10.1042/BST20150146
(2015).
36   Cordova, L. T., Long, C. P., Venkataramanan, K. P. & Antoniewicz, M. R. Complete
genome sequence, metabolic model construction and phenotypic characterization of

Geobacillus LC300, an extremely thermophilic, fast growing, xylose-utilizing
bacterium. *Metab Eng* **32**, 74-81, doi:10.1016/j.ymben.2015.09.009 (2015).

37 de la Torre, A. *et al.* Genome-scale metabolic reconstructions and theoretical
investigation of methane conversion in Methylomicrobium buryatense strain 5G(B1).
Microb Cell Fact **14**, 188, doi:10.1186/s12934-015-0377-3 (2015).

38 Zou, W. *et al.* Reconstruction and analysis of a genome-scale metabolic model of the
vitamin C producing industrial strain Ketogulonicigenium vulgare WSH-001. *J*
Biotechnol **161**, 42-48, doi:10.1016/j.jbiotec.2012.05.015 (2012).

39 Pacheco, M. P., Bintener, T. & Sauter, T. Towards the network-based prediction of
repurposed drugs using patient-specific metabolic models. *EBioMedicine*,
doi:10.1016/j.ebiom.2019.04.017 (2019).

40 Izallalen, M. *et al.* Geobacter sulfurreducens strain engineered for increased rates of
respiration. *Metab Eng* **10**, 267-275, doi:10.1016/j.ymben.2008.06.005 (2008).

41 Burgard, A. P., Pharkya, P. & Maranas, C. D. Optknock: a bilevel programming
framework for identifying gene knockout strategies for microbial strain optimization.
Biotechnol Bioeng **84**, 647-657, doi:10.1002/bit.10803 (2003).

42 Ang, K. S., Lakshmanan, M., Lee, N. R. & Lee, D. Y. Metabolic Modeling of Microbial
Community Interactions for Health, Environmental and Biotechnological Applications.
Curr Genomics **19**, 712-722, doi:10.2174/1389202919666180911144055 (2018).

43 Zhuang, K. *et al.* Genome-scale dynamic modeling of the competition between
Rhodoferax and Geobacter in anoxic subsurface environments. *The ISME journal* **5**,
305-316, doi:10.1038/ismej.2010.117 (2011).

44 Xu, X. *et al.* Modeling microbial communities from atrazine contaminated soils
promotes the development of biostimulation solutions. *The ISME journal* **13**, 494-508,
doi:10.1038/s41396-018-0288-5 (2019).

45 Faust, K. Microbial Consortium Design Benefits from Metabolic Modeling. *Trends*
Biotechnol **37**, 123-125, doi:10.1016/j.tibtech.2018.11.004 (2019).

46 Henry, C. S. *et al.* High-throughput generation, optimization and analysis of genome-
scale metabolic models. *Nature biotechnology* **28**, 977-982, doi:10.1038/nbt.1672
(2010).

47 Kundu, K. *et al.* Defining lower limits of biodegradation: atrazine degradation
regulated by mass transfer and maintenance demand in Arthrobacter aurescens TC1.
The ISME journal, doi:10.1038/s41396-019-0430-z (2019).

48 Oh, Y. K., Palsson, B. O., Park, S. M., Schilling, C. H. & Mahadevan, R. Genome-scale
reconstruction of metabolic network in Bacillus subtilis based on high-throughput
phenotyping and gene essentiality data. *The Journal of biological chemistry* **282**,
28791-28799, doi:10.1074/jbc.M703759200 (2007).

49 Larocque, M., Chenard, T. & Najmanovich, R. A curated C. difficile strain 630 metabolic
network: prediction of essential targets and inhibitors. *BMC systems biology* **8**, 117,
doi:10.1186/s12918-014-0117-z (2014).

50 Orth, J. D. *et al.* A comprehensive genome-scale reconstruction of Escherichia coli
metabolism--2011. *Molecular systems biology* **7**, 535, doi:10.1038/msb.2011.65
(2011).

51 Hucka, M. *et al.* The systems biology markup language (SBML): a medium for
representation and exchange of biochemical network models. *Bioinformatics* **19**, 524-
531, doi:10.1093/bioinformatics/btg015 (2003).

52 Govantes, F. *et al.* Regulation of the atrazine-degradative genes in Pseudomonas sp.
strain ADP. *FEMS microbiology letters* **310**, 1-8, doi:10.1111/j.1574-
6968.2010.01991.x (2010).

53 Guo, Q., Zhang, J., Wan, R. & Xie, S. Impacts of carbon sources on simazine
biodegradation by Arthrobacter strain SD3-25 in liquid culture and soil microcosm.
International Biodeterioration & Biodegradation **89**, 1-6 (2014).
54 Dsouza, M., Taylor, M. W., Turner, S. J. & Aislabie, J. Genomic and phenotypic insights
into the ecology of Arthrobacter from Antarctic soils. *BMC Genomics* **16**, 36,
doi:10.1186/s12864-015-1220-2 (2015).
55 Deutch, C. E. L-Proline catabolism by the high G + C Gram-positive bacterium
Paenarthrobacter aurescens strain TC1. *Antonie Van Leeuwenhoek* **112**, 237-251,
doi:10.1007/s10482-018-1148-z (2019).
56 Mahadevan, R. & Schilling, C. H. The effects of alternate optimal solutions in
constraint-based genome-scale metabolic models. *Metab Eng* **5**, 264-276 (2003).
57 Zhang, Q. *et al.* Impacts of nitrogen and phosphorus on atrazine-contaminated soil
remediation and detoxification by Arthrobacter sp. strain HB-5. *Environmental earth*
sciences **71**, 1465-1471 (2014).
58 Zhang, C., Li, Y., Wang, C., Niu, L. & Cai, W. Occurrence of endocrine disrupting
compounds in aqueous environment and their bacterial degradation: A review.
Critical Reviews in Environmental Science and Technology **46**, 1-59,
doi:10.1080/10643389.2015.1061881 (2015).
59 Meyer, F. *et al.* The metagenomics RAST server – a public resource for the automatic
phylogenetic and functional analysis of metagenomes. *BMC Bioinformatics* **9**, 1-8,
doi:10.1186/1471-2105-9-386 (2008).
60 Orth, J. D., Thiele, I. & Palsson, B. O. What is flux balance analysis? *Nat Biotech* **28**,
245-248,
doi:http://www.nature.com/nbt/journal/v28/n3/abs/nbt.1614.html#supplementary
-information (2010).
61 Varma, A. & Palsson, B. O. Stoichiometric flux balance models quantitatively predict
growth and metabolic by-product secretion in wild-type Escherichia coli W3110.
Applied and environmental microbiology **60**, 3724-3731 (1994).
62 Thiele, I. & Palsson, B. O. A protocol for generating a high-quality genome-scale
metabolic reconstruction. *Nat Protoc* **5**, 93-121, doi:10.1038/nprot.2009.203 (2010).
63 Harcombe, W. R. *et al.* Metabolic resource allocation in individual microbes
determines ecosystem interactions and spatial dynamics. *Cell reports* **7**, 1104-1115,
doi:10.1016/j.celrep.2014.03.070 (2014).
64 King, Z. A. *et al.* Escher: A Web Application for Building, Sharing, and Embedding Data-
Rich Visualizations of Biological Pathways. *PLoS computational biology* **11**, e1004321,
doi:10.1371/journal.pcbi.1004321 (2015).

Acknowledgments

This work was funded by grants from the NSFC-ISF joint program (31461143009),

Israel Science Foundation Grant no. 1416/14. Atrazine was a gift of Agan Adama

(Ashdod, Israel). We thank Omer Kapiluto for statistical support.

Author Contributions

S.O, R.Z, T.L and S.F designed the study, constructed the model and performed *in*

silico analysis. Z.R, S.P, R.A, and D.G designed and performed the *in vitro*

experiments. Y.K and H.E analyzed and reviewed the results. All authors and

approved the final version of the manuscript.

Additional Information

Supplementary information:

Supplemental data 1 – Model reactions and metabolites

Supplemental data 2 – SBML model file

Supplemental data 3 – Simulation parameters and defaults.

Supplemental data 4 – Alternative media compositions

Supplemental data 5 – Supplemental figures

Competing Interests:

The authors declare that they have no conflict of interest.

FIGURE LEGENDS

Figure 1. Atrazine metabolism as implemented in the cytoplasm compartment of *P. aurescens* TC1 genome scale metabolic model reconstruction (*i*RZ1176). External metabolites are notated with an e0;

Figure 2. Growth performances of *P.aurescens* TC1 in dependence on the availability of different carbon and nitrogen sources. (A) Predicted and observed *A.aurescens* TC1 growth in mineral minimal media supplemented with Atrazine or its degradation intermediates. ATZ-Atrazine, NH4, GLU-Glucose, ETH - Ethylamine, ISO – Isopropylamine, CTRL - control. Values are normalized by dividing in maximal value. (B) Surface plot rendering *P.aurescens* TC1 growth (biomass) in dependence on atrazine, glucose and NH4 availability. Color scale is indicative of biomass (Z-axis).

628   Figure 3. Predicted and observed growth performances of *P. aurescens* TC1 growth in mineral minimal media with glucose complemented by different amino acids and atrazine. ATZ-Atrazine, HIS- Histdine, ILE-Isoluecine and MET- Metionine. Observed values represents mean in triplicates; bars represent SD.

633   Figure 4. Predicted and observed atrazine degradation in modified media. High and low phosphate concentrations: 50 and 15 mmol/gdw h for predicted and 0.081 and 0.081 g/l for observed, respectively. ATZ- atrazine, GLU- glucose. Observed values represents mean in triplicates; bars represent SD.