Genome-Scale reconstruction of *Paenarthrobacter aurescens* TC1 metabolic model

towards the study of atrazine bioremediation

Shany Ofaim[1,2]* [ŧ], Raphy Zarecki[1]*, Seema Porob[3], Daniella Gat[3], Tamar Lahav[1], Xihui

Xu[1, 4], Yechezkel Kashi[2], Radi Aly[1], Jiandong Jiang[4], Hanan Eizenberg[1], Zeev Ronen[3 †] &

Shiri Freilich[ŧ][1][†]

[1]Newe Ya'ar Research Center, Agricultural Research Organization, Ramat Yishay,

Israel, [2]Faculty of Biotechnology and Food Engineering, Technion-Israel Institute of

Technology, Haifa, Israel, [3]Department of Environmental Hydrology & Microbiology,

Zuckerberg Institute for Water Research, Jacob Blaustein Institutes for Desert

Research, Ben-Gurion University of the Negev, Midreshet Ben-Gurion, 8499000,

Israel, [4]Department of Microbiology, Agricultural University of Nanjing, No. 6

Tongwei RD, 210095, Nanjing, Jiangsu Province, China

*equal contribution

[†] equal contribution

ŧ Corresponding author

email address (in the order of author list): shashany@gmail.com,

raphy.zarecki@gmail.com, porob@post.bgu.ac.il, gat.daniella@gmail.com,

lahav008@gmail.com, xuxihui01@gmail.com, kashi@technion.ac.il,

radi@volcani.agri.gov.il, jiang_jjd@njau.edu.cn, eizenber@volcani.agri.gov.il,

zeevrone@bgu.ac.il, shiri.freilich@gmail.com.

ABSTRACT

Atrazine is an herbicide and pollutant of great environmental concern that is naturally biodegraded by microbial communities. The efficiency of biodegradation can be improved through the stimulating addition of fertilizers, electron acceptors, etc. In recent years, metabolic modelling approaches have become widely used as an *in silico* tool for organism-level phenotyping and the subsequent development of metabolic engineering strategies including biodegradation improvement. Here, we constructed a genome scale metabolic model, *i*RZ960, for *Paenarthrobacter aurescens* TC1 – a widely studied atrazine degrader - aiming at simulating its degradation activity. A mathematical stoichiometric metabolic model was constructed based on a published genome sequence of *P. aurescens* TC1*.* An Initial draft model was automatically constructed using the RAST and KBase servers. The draft was developed into a predictive model through semi-automatic gap-filling procedures including manual curation. In addition to growth predictions under different conditions, model simulations were used to identify optimized media for enhancing the natural degradation of atrazine without a need in strain design via genetic modifications. Model predictions for growth and atrazine degradation efficiency were tested in myriad of media supplemented with different combinations of carbon and nitrogen sources that were verified *in vitro*. Experimental validations support the reliability of the model's predictions for both bacterial growth (biomass accumulation) and atrazine degradation. Predictive tools, such as the presented model, can be applied

for achieving optimal biodegradation efficiencies and for the development of

ecologically friendly solutions for pollutant degradation in changing environments.

KEYWORDS

Genome scale metabolic models; Flux Balance Analysis; *Paenarthrobacter aurescens*

TC1; atrazine; bioremediation

Background

Areas contaminated with hazardous materials are rapidly increasing worldwide

introducing a need in remediation approaches. Bioremediation - an environmental

bioprocess in which naturally occurring organisms are used for breaking down

hazardous substances into less toxic or non-toxic substances, is increasingly

acknowledged as a cost-effective strategy for environmental cleaning (Kanissery and

Sims 2011; Kastner and Miltner 2016). As critical environmental bioprocesses carried

by bacteria are either related to removal of pollutants from water, soil or air or

involved in the production of commercially valuable compounds (Perez-Garcia et al.

2016), bacterial bioremediation is widely applied for the degradation of various

organic pollutants (Mani and Kumar 2013; Megharaj et al. 2011; Nilsson et al. 2013;

Shah and Dahanukar 2012) such as herbicides (Fan and Song 2014; Megharaj et al.

2011; Nousiainen et al. 2015; Sagarkar et al. 2014; Singh and Singh 2016).

Bioremediation may occur naturally or through the stimulating addition of fertilizers,

electron acceptors, etc. These amendments encourage the growth of the degrading

microbes within the medium or environment in a process termed biostimulation

(Adams et al. 2015; Kanissery and Sims 2011).

Atrazine is an herbicide and pollutant of great environmental concern. Its residues are

still found in soil samples decades after it was last applied. These residues were shown

to chronically leach into the local aquifer (Abubucker et al. 2012). Atrazine has been

widely reported to be biodegraded by bacteria (Chen et al. 2014; Fan and Song 2014;

Monard et al. 2011; Nousiainen et al. 2015; Sagarkar et al. 2013; Sagarkar et al. 2014;

Udikovic-Kolic et al. 2012; Wang and Xie 2012; Xie et al. 2013; Zhang et al. 2012). One

of the well-known atrazine degraders is *Paenarthrobacter aurescens* TC1 (Fang et al.

2015; Sagarkar et al. 2013; Strong et al. 2002; Wang and Xie 2012; Zhang et al. 2012).

P. aurescense TC1 is described as nutritionally versatile bacterium that is capable of

utilizing a wide array of complex metabolites, including agriculturally used herbicides

(Deutch et al. 2018; Mongodin et al. 2006). The full hydrolytic pathway for atrazine

degradation was reported and described in *P. aurescens* TC1. Intermediates, created

in the degradation process are further metabolized by the organism and can provide

sole sources for carbon and nitrogen (Strong et al. 2002). This bacterium was

previously reported to biodegrade atrazine much more efficiently than most atrazine

degrading bacteria including *Pseudomonas* sp. ADP (Silva et al. 2015). While PADP can

only use the hydrolytic products of the ring cleavage as nitrogen but not as carbon;

TC1 can use the side groups of atrazine as source for carbon, nitrogen and energy

consuming up to 3,000 mg of atrazine per liter (Strong et al. 2002). As such, *P.*

aurescens TC1 can be a model for the study of atrazine bioremediation.

Modern tools of genomics, transcriptomics, metabolomics, proteomics, signaling

systems and synthetic biology have opened new avenues for biotechnological

advances and are increasingly applied for promoting better understanding of complex

biological systems. In recent years, constraint based metabolic modelling approaches

have become widely used as an *in silico* tool for organism-level phenotyping and the

subsequent development of metabolic engineering strategies (Contador et al. 2015;

Xu et al. 2013). Generally, such approaches follow four key steps: (1) data acquisition

– mainly genome sequencing information, basic cell-physiological and biochemical

knowledge and some experimental data on cell growth; (2) model reconstruction -

translating data into structured mathematical representation; (3) constraint-based

optimization simulations - the prediction of growth rate, substrate uptake rates, and

byproduct rates under different growth conditions or following knockout mutations,

in the absence of kinetic information (Lee et al. 2008; Mahadevan et al. 2006;

Mahadevan and Lovley 2008; Oberhardt et al. 2015; Price et al. 2003; Reed et al. 2003;

Thiele et al. 2005); and (4) experimental validation. The potential of such models for

the investigation of optimal processing is now acknowledged and practiced (Biswas et

al. 2015; Fang et al. 2011; Henson 2015; Perez-Garcia et al. 2016; Zomorrodi and Segre

2016). Examples for applicative use include the optimal production or utilization of

industrial compounds such as xylose (Cordova et al. 2015), biofuels (de la Torre et al.

2015) and vitamins (Zou et al. 2012). In bioremediation, genome scale metabolic

modeling approaches were applied for the design of *Geobacter sulfurreducens* strain

capable of increased electron transfer and a higher Fe(III) respiratory that was

beneficial for environmental bioremediation (Izallalen et al. 2008). Strain design is

often based on the Optknock algorithm (Burgard et al. 2003) and its derivatives which

search for the optimal gene knockouts for a desired metabolic production. However,

unlike industrial use of engineered strain for enhanced production in monoculture,

introduction of exogenous species into natural habitat is far from trivial and hampers

the application of modeling approaches towards bioremediation solutions. An

alternative strategy is the use of modeling for the design of optimal conditions that

will enhance processes (degradation or synthesis) carried by the endogenous

community. In a recent project we demonstrate the use of metabolic models for

predicting interactions between *P.aurescens* TC1 and endogenous soil species.

Exchange metabolites secreted by non-degraders species contribute for enhancing

P.aurescens TC1 growth and expediting degradation efficiency (Xu et al. 2019). Here,

we describe in detail the process of manual curation of *P.aurescens* TC1 draft

metabolic model, its application for predicting growth and atrazine degradation in

media containing different carbon and nitrogen sources based on a dynamic (capable

of reporting changes over time) and non-dynamic flux based algorithms, and the

experimental testing of the predictions.

Materials and Methods

<u>Reconstruction of the initial draft for *P. aurescens* TC1 metabolic model</u>

An initial draft model of *P. aurescens* TC1 was reconstructed as described in (Xu et al.

2019). Briefly, the genome sequence of *P. aurescens* TC1 was retrieved from NCBI

(GenBank accession id. NC_008711), uploaded into RAST (Meyer et al. 2008), spliced

into genes and annotated. RAST annotations were put through the Model SEED

pipeline (Henry et al. 2010) and an initial metabolic network reconstruction was

obtained. This initial reconstruction contained a list of reactions classified as exchange,

transport, and cytosolic as well as a list of all relevant metabolites and a biomass

reaction. The composition of the biomass reaction summarizes the fractional

contribution of a generalized microbial biomass precursors (e.g., amino acids and

lipids) to the synthesis of a new cell, as provided in the draft reconstructions from

Model SEED and KBase. Unspecified non-metabolic consuming processes in the cell

were defined as ATP maintenance and set similarly to (Oh et al. 2007). Missing

reactions were filled through an automatic gap-filling procedure, using default

(optimal) media options (Henry et al. 2010). The model was exported from KBase in

SBML format (Additional File 4).

<u>Model curation and evaluation</u>

Manual curation procedures were carried to adapt model performances to reported

physiological capacities. Simulation conditions were modified to reflect

experimentally validated minimal media alternatives that support growth of *P.*

aurescnes TC1 (Strong et al. 2002). Missing reactions (network gaps) were filled based

on integrating information extracted from public databases such as KEGG (Kanehisa

and Goto 2000; Kanehisa et al. 2014), Uniprot (UniProt 2015), and JGI databased

(Kanehisa et al. 2012; Nordberg et al. 2014; UniProt 2015) (Additional File 1). The

procedures required the following steps: (1) Identifier standardization - reaction

identifiers from the different databases were converted into KBASE rxn accessions and

added to the model; (2) Reaction reversibility calibration according to the KEGG

scheme; (3) Reaction elemental and redox balance verification; and (4) removal of

futile loops – active internal (non-exchange) reactions which remained active while all

external fluxes were blocked (upper and lower bounds set to zero). Remaining gaps

were then manually filled by an extensive literature survey and the use of other public

resources (Additional File 2).

All curation procedures were carried iteratively versus growth simulations to ensure

the production of all biomass components in Minimal Mineral Media (MMM: K^+, Mn^{2+},

CO2, Zn^{2+}, SO_4^-, Cu^{2+}, Ca^{2+}, HPO_4^{2-}, Mg^{2+}, Fe^{2+}, Cl^-), supplemented by alternative carbon

and nitrogen sources including atrazine, ethylamine, isopropylamine, ammonia,

glucose and selected amino acids (Additional File 3). As part of the curation process,

the hydrolytic pathway of atrazine degradation was *a priori* added to the model based

on previously published experimental data (Strong et al. 2002).

<u>Flux balance analyses simulations and dynamic, time dependent simulations</u>

Flux distributions in the metabolic model were determined using Flux Balance Analysis

(FBA) (Orth et al. 2010; Thiele and Palsson 2010; Varma and Palsson 1994). Briefly, a

metabolic model includes a network that is described as a stoichiometric matrix $\mathbf{S}_{m \times n}$,

where *m* represents the number of metabolites and *n* the number of reactions in the

model. In the assumed pseudo-steady-state, internal metabolites concentrations and

fluxes are assumed to be constant. The model can be represented as: $\mathbf{S} \times v = 0$, where

vector v signifies the fluxes through the internal reactions. As the number of reactions

(*n*) typically exceeds the number of metabolites (*m*), the system is under-determined.

In order to get a feasible solution space, linear programming (LP), subject to mass

balance preservation and flux constraints, was used by introducing an optimization

problem. Here, model growth was maximized by maximizing its biomass reaction,

which was used as objective function. Specific fluxes for metabolites consumption and

secretion were determined following fixation of the respective biomass reaction to its

maximal value and then using Flux Variability Analysis (FVA) (Mahadevan and Schilling

2003) to calculate possible ranges of these metabolites uptake and secretion. Dynamic

modeling was used for the prediction of the profile of consuming metabolites typical

to the biomass increase of *P. aurescens* TC1 and atrazine degradation across time. To

this end, we simulated the behavior of our metabolic model across time. This was

done as described in (Xu et al. 2019) and illustrated in Additional File 5, based on

concepts defined in (Harcombe et al. 2014). Briefly, the model works under the

following assumptions: (1) A finite start amount of media components is equally

available to all members of the population; (2) A maximal amount of uptake a single

cell can acquire from the media in a given time point is defined (the lower bound of

the exchange reaction value); (3) New substrate concentrations in time t, are

determined by the predicted substrate concentration for the previous step

augmented with any additional substrates provided or consumed in the current

iteration. The maximum uptake of a single organism was set to a ratio of up to 1 unit

of each metabolite available in the media to each organism; (4) After each time tick,

the biomass amount was updated according to the flux amount of the biomass

reaction in the model at this time tick. As biomass production rate serves as a proxy

for the size of the population in the simulated environment and substrate

uptake/secretion is mainly affected by to population size, the model was used to

evaluate the actual substrate uptake/secretion and growth rate given the supplied

media across time. Simulations were carried until reaching a state where time cycles

did not lead to an increase in biomass. Initial concentration values were set to a fixed

amount of 50 units (represented as the initial lower bounds, LB, of the exchange

reactions). Simulation parameters and defaults are available in Additional File 6.

Starting with one bacterium, the flux balance model was used to predict the uptake

of phosphate, sulfate, carbon and nitrogen sources (*i.e.* atrazine and glucose) by *P.*

aurescens TC1 across time.

All model simulations were done on an Intel i7 quad-core server with 32GB of

memory, running linux. The development programming language of our simulators

was JAVA, and our linear programming software was IBM CPLEX.

<u>*In vitro* bacterial growth conditions and chemical analysis</u>

P.aurescens TC1 (ATCC BAA-1386™) was grown on R medium (based on ATCC 2662 R)

plates with glucose (0.2%) and atrazine (30 mg/l, Agan Adama -Ashdod, Israel Israel)

as sole nitrogen. Stock cultures were maintained in glycerol at -80°C. Cultures were

resuscitated on R medium agar plates (1.5% Difco bacto™) and were used as starter

colonies for liquid medium growth experiments. The growth of *P.aurescens* TC1 was

tested in R media supplemented with different combinations of carbon and nitrogen

sources, including Glucose (0.2%), Atrazine (30 mg/l, Agan Adama -Ashdod, Israel),

ammonium chloride (37 mg/l), Ethylamine and Isopropylamine (31.1 and 40.8 mg/l,

Sigma-Aldrich -Rehovot, Israel) and the amino acids Histidine, Isoleucine and

Methionine (36, 92 and 103 mg/l respectively). Concentrations of amino acids,

ethylamine and Isopropylamine were calculated to provide the same nitrogen content

as the atrazine. In addition to the carbon and nitrogen sources, growth in media

supplemented by atrazine and atrazine together with glucose was tested under two

phosphate concentrations (K_2HPO_4 in 0.45 and 0.045 g/l). Each substrate was added

to sterile R medium and bacteria was grown in triplicate 250ml flasks incubated in

30°C with shaking in the dark. One flask contained only R medium and was used as

control. A 200 ul sample was taken from each flask every 8 hours for measurement of

OD in microplate reader (520nm, Infinite® 200 PRO Tecan, Männedorf Switzerland)

and 1 ml was taken to study atrazine degradation rate using HPLC. The HPLC analysis

was performed with Agilent 1100 HPLC (Waldbronn Germany) equipped with a DAD

detector. Samples were separated on Kinetex C18 column (Phenomenex Torrance, CA)

and the mobile phase consisted of 70% methanol flowing at 1 ml/min. Detection and

quantitation of atrazine was done at 240 nm with detection limit of 0.1 mg/l. Total

duration of each growth experiment was between 152 to 168 hours.

Visualizations

Venn diagrams were created using Venny (Oliveros 2007). Surface plots were created

using the MATLAB 2017b (MathWorks, Inc., Natick, Massachusetts, United States).

Pathway maps were created using Escher maps (King et al. 2015).

Results

P.aurescens TC1 model construction and validation

A genome-scale metabolic model for _P.aurescens_ TC1 (_i_RZ960) was reconstructed

and curated. Curation procedures were carried iteratively using growth simulations,

ensuring the production of all biomass components under the minimal feasible

conditions described in Strong et al. (2002). Reconstruction gaps in the initial draft

model were manually filled by the addition of _P. aurscens_ TC1 reactions retrieved

from alternative databases or supported by the literature (Additional File 2). The

curated model contains 960 genes, 2322 reactions, out of which 125 are exchange,

and 2409 metabolites. Model Reactions involved in atrazine degradation pathway

and its further conversion into cellular building blocks are illustrated in Figure 1. As

previously reported, _P.aurescens_ TC1 degrades atrazine and metabolizes

isopropylamine and ethylamine, both can provide sole carbon and nitrogen sources

(Strong et al. 2002). Growth simulations confirm that the model captures these

reported physiological performances and predictions were directly tested *in vitro*

under conditions equivalent to these used for simulations. Experimental growth

results verify that atrazine and its degradation products isopropylamine and

ethylamine can be used as sole carbon and nitrogen sources for *P. aurscens* TC1

(Figure 2A). Supplementing the media with glucose enhanced growth both *in silico*

and *in vitro* while supplementing it with ammonia had not shown an effect on

bacterial growth. This is in accordance with previous work stating that *P.aurescens*

TC1 growth is not inhibited by nitrogen sources such as ammonia (Silva et al.

2015).Overall, model simulations corresponded with experimental results and fully

repeat previous reports (Figure 2A).

Finally, growth simulations under a gradient of nitrogen *vs*. carbon   sources further

illustrate that atrazine serves as a good nitrogen source and as a poor carbon source

(Figure 2B). The growth promoting potential of combinations of atrazine with

ammonium (NH_4^+) and glucose as nitrogen and/or carbon sources respectively was

tested. The effect of nutrient ratio on growth rate was predicted by optimizing the

production of biomass across uptake gradients while constraining the fluxes through

exchange reactions for carbon and nitrogen sources. As expected, increasing glucose

uptake increased the growth rate (Figure 2B) and the maximal achievable growth

rate was much higher with glucose as a carbon source in comparison to atrazine as a

sole carbon source. However, atrazine is as equally good nitrogen source as

ammonia. This may be endorsed by ammonium as a byproduct of the atrazine

degradation process (Figure 1).

<u>Effect of media modifications on growth</u>

In addition to glucose, several carbon sources were previously reported to enhance

growth of several strains of *Arthrobacter* species in atrazine containing media (Guo et

al. 2014; Silva et al. 2015). To identify additional media modifications that can

potentially increase atrazine degradation, we simulated growth in 123 different media

combinations, each supplementing the atrazine containing minimal mineral media

with a single exchange metabolite. Several sugars, amino acids and dipeptides were

predicted to induce enhanced growth (Additional File 7). This is in accordance with the

previous characterization of *P.aurescens* TC1 as a metabolically versatile species

(Deutch et al. 2018; Mongodin et al. 2006). Out of a variety of potential nutritional

sources, *Paenarthrobacter* species have been shown to utilize different amino acids in

different capacities (Dsouza et al. 2015). The effect of amino-acids is media

dependent. For example, out of all amino acids tested, Isoleucine has the strongest

predicted effect on growth in atrazine containing medium whereas histidine has

stronger effect in the medium without atrazine; methionine has the weakest effect in

all media (Figure 3).   *P.aurescens* TC1 growth in *in vitro* media corresponding to the

media used for the *in silica* simulations were generally in agreement with model's

predictions supporting the superiority of isoleucine over histidine in atrazine

supplemented media but not in the atrazine deficient medium (Figure 3).

Supplementing media with methionine showed weaker effect on growth in

comparison to the other two amino acids tested, in accordance with its weak effect in

the predictions. <u>Prediction and evaluation of the effect of media modifications on</u>

<u>atrazine degradation</u>

Following the validation of the model predictions for growth capacities, we directly

evaluated atrazine degradation performances. To this end, we applied FVA, allowing

the identification of the ranges of flux variability that are possible within a given media

(Mahadevan and Schilling 2003). Once the biomass flux was maximized, we estimated

the rate of atrazine degradation according to the relevant flux. Atrazine degradation

over time was simulated in a medium that contains atrazine as a sole carbon and

nitrogen source *vs*. modified media sets. Atrazine degradation was estimated

according to the amount of atrazine left in the media following each time cycle.

Simulations showed that supplementing the media with glucose not only enhances

growth (Figure 2) but also induced an enhancement of atrazine consumption.

Predictions were confirmed by experimental validation (Figure 4).

Finally, the addition of phosphate - a main element in fertilizers, to soils has been

reported to have a significant effect on the degradation of herbicides. For instance, its

addition was shown to effect soil bacteria's degradation activity including a reported

effect on *Arthrobacter* sp. HB-5, a close relative of *P.aurescens* TC1 (Zhang et al. 2014).

Phosphate has been previously shown to accelerate atrazine degradation (Kanissery

and Sims 2011). We used the genome scale reconstruction to simulate the effect of

phosphate on growth and atrazine degradation. In accordance with these reports,

simulations showed similar trends, which were confirmed by experimental results,

indicating an effect of phosphate concentrations *on P. aurescens* TC1 growth (sup

Figure 4) and atrazine degradation (Figure 4).

Discussion

The accumulation and persistence of herbicides and their intermediates in soil and

drinking water has long been suggested to be disruptive, imposing a need for effective

strategies for their degradation and removal (Zhang et al. 2015). Bioremediation is a

widely applied approach for the degradation of herbicides including atrazine (Fan and

Song 2014; Megharaj et al. 2011; Nousiainen et al. 2015; Sagarkar et al. 2014; Singh

and Singh 2016). Here, we reconstructed a genome scale metabolic model, *i*RZ960, of

P. aurescnes TC1, a well-known atrazine degrader, in order to use the *in silica*

simulations for predicting optimal media to support its growth and atrazine

degradation activity. The *in silica* simulations were carried on an array of minimal

media using atrazine and its interim degradation products as alternative sole carbon

and nitrogen sources supplemented by potential enhancers. The simulations

suggested a group of metabolites promoting the increase in biomass yield including

sugars, amino acids and dipeptides. We hypothesize that the introduction of such an

efficient carbon source leads to an increase in biomass and a subsequent increase in

the degradation activity. Simulations were confirmed by *in vitro* experimental

validation. This genome scale reconstruction may be used for further experimenting

and the gain of new biological knowledge of this metabolically versatile bacteria.

While the metabolic modeling approach presented here was demonstrated useful for

generating predictions, its limitations should be acknowledged. First, model

construction relies on automatic protocols and publically free genomic information.

Biomass composition and energy maintenance requirements are hence

approximations (Henry et al. 2010; Oh et al. 2007). Second, data used to construct the

model is eclectic and annotations were gathered from numerous databases. The lack

of data standardization has led to many overlaps and misnomers resulting in draft

models containing massive gaps. The lion share of gap filling processes are still done

manually due to the differences in data description and cataloging in the different

databases. As technology and standardization advance, the process of constructing a

metabolic model able to produce testable predictions will be faster and have greater

accuracy. Despite relying on genomic data and a growth approximation, resulting

predictions are in agreement with experimental observations. We demonstrate that

through using metabolic time dependent simulations, an optimal range of nutrient

augmentation may be planned. Continuation of the manual curation process, also

based on added transcription information, is likely to improve the performances of

future versions.

Overall, we describe the construction of a mathematical stoichiometric metabolic

model of *Paenarthrobacter aurescens* TC1 (*i*RZ960), allowing simulating its

degradation activity in contaminated soil. The model successfully predicted both

bacterial growth and atrazine degradation in culture. Simulations were used to

predict growth and degradation following media modifications such as high/low

phosphate content and the addition of variety of amino acids to the standard

minimal R medium. Effects of such media were observed experimentally to effect

atrazine degradation. *i*RZ960 presented here is shown to be a useful tool for

prediction of the metabolic behavior of the bacteria in different media conditions

and may be used to further study this versatile organism. Natural degradation of

pollutants such as atrazine is of great importance worldwide. A predictive tool such

as the presented model is useful for achieving optimal degradation efficiencies.

Integration of such models in community models was demonstrated useful for the

development of ecologically friendly solutions for pollutant degradation in changing

environments (Xu et al. 2019). The integration of species-specific models together

with community approaches is likely to contribute to our understanding of

biodegradation processes in natural environments and promote the development of

bioaugmentation and biostimulation strategies.

LIST OF ABRAVATIONS

383 MMM - Minimal Mineral Media; FBA - Flux Balance Analysis; LP -Linear Programming;

384 FVA - Flux Variability Analysis; L/UB – lower/upper bounds.

DECLARATIONS

AVAILABILITY OF DATA AND MATERIALS

387 All data generated or analyzed during this study are included in this published article

388 and its supplementary information files.

FUNDING

391 This work was supported by grants from the NSFC-ISF joint program (31461143009),

392 Israel Science Foundation Grant no. 1416/14, and the Science Foundation of Jiangsu

393 Province, China (BK20150670). Atrazine was a gift of Agan Adama (Ashdod, Israel)

AUTHORS' CONTRIBUTIONS

RZ and SO equally contributed to model construction and predicted analysis. TL and

X.X., contributed to model curation, testing & development. ZR had designed the

experiments and SP and DG carried them. HE, JJ and SF and ZR X.X., R.Z. analyzed the

data and designed research. S.O, R.Z., ZR, S.F. and J.J. wrote the manuscript with

feedback from all authors.

All authors read and approved the final manuscript.

COMPETING INTERESTS

None of the authors have any competing interests in the manuscript.

REFERENCES

Abubucker S, Segata N, Goll J, Schubert AM, Izard J, Cantarel BL, Rodriguez-Mueller B,
Zucker J, Thiagarajan M, Henrissat B, White O, Kelley ST, Methe B, Schloss PD,
Gevers D, Mitreva M, Huttenhower C (2012) Metabolic reconstruction for
metagenomic data and its application to the human microbiome. PLoS
computational biology 8(6):e1002358 doi:10.1371/journal.pcbi.1002358
Adams GO, Fufeyin PT, Okoro SE, Ehinomen I (2015) Bioremediation, biostimulation
and bioaugmention: a review. International Journal of Environmental
Bioremediation & Biodegradation 3(1):28-39
Biswas K, Paul D, Sinha SN (2015) Biological Agents of Bioremediation: A Concise
Review. Microbiology 1(3):39-43
Burgard AP, Pharkya P, Maranas CD (2003) Optknock: a bilevel programming
framework for identifying gene knockout strategies for microbial strain
optimization. Biotechnol Bioeng 84(6):647-57 doi:10.1002/bit.10803
Chen Q, Wang H, Yang B, He F (2014) The combined effects of atrazine and lead (Pb):
relative microbial activities and herbicide dissipation. Ecotoxicol Environ Saf
102:93-9 doi:10.1016/j.ecoenv.2014.01.011
Contador CA, Rodriguez V, Andrews BA, Asenjo JA (2015) Genome-scale
reconstruction of Salinispora tropica CNB-440 metabolism to study strain-
specific adaptation. Antonie Van Leeuwenhoek 108(5):1075-90
doi:10.1007/s10482-015-0561-9
Cordova LT, Long CP, Venkataramanan KP, Antoniewicz MR (2015) Complete genome
sequence, metabolic model construction and phenotypic characterization of
Geobacillus LC300, an extremely thermophilic, fast growing, xylose-utilizing
bacterium. Metab Eng 32:74-81 doi:10.1016/j.ymben.2015.09.009
de la Torre A, Metivier A, Chu F, Laurens LM, Beck DA, Pienkos PT, Lidstrom ME,
Kalyuzhnaya MG (2015) Genome-scale metabolic reconstructions and
theoretical investigation of methane conversion in Methylomicrobium
buryatense strain 5G(B1). Microb Cell Fact 14(1):188 doi:10.1186/s12934-015-
0377-3

Deutch CE, Bui AP, Ho T (2018) Growth of Paenarthrobacter aurescens strain TC1 on
atrazine and isopropylamine during osmotic stress. Annals of Microbiology
68(9):569-577 doi:10.1007/s13213-018-1364-9
Dsouza M, Taylor MW, Turner SJ, Aislabie J (2015) Genomic and phenotypic insights
into the ecology of Arthrobacter from Antarctic soils. BMC Genomics 16:36
doi:10.1186/s12864-015-1220-2
Fan X, Song F (2014) Bioremediation of atrazine: recent advances and promises.
Journal of Soils and Sediments 14(10):1727-1737 doi:10.1007/s11368-014-
0921-5
Fang H, Lian J, Wang H, Cai L, Yu Y (2015) Exploring bacterial community structure and
function associated with atrazine biodegradation in repeatedly treated soils. J
Hazard Mater 286:457-65 doi:10.1016/j.jhazmat.2015.01.006
Fang Y, Scheibe TD, Mahadevan R, Garg S, Long PE, Lovley DR (2011) Direct coupling
of a genome-scale microbial in silico model and a groundwater reactive
transport model. J Contam Hydrol 122(1-4):96-103
doi:10.1016/j.jconhyd.2010.11.007
Guo Q, Zhang J, Wan R, Xie S (2014) Impacts of carbon sources on simazine
biodegradation by Arthrobacter strain SD3-25 in liquid culture and soil
microcosm. International Biodeterioration & Biodegradation 89:1-6
Harcombe WR, Riehl WJ, Dukovski I, Granger BR, Betts A, Lang AH, Bonilla G, Kar A,
Leiby N, Mehta P, Marx CJ, Segre D (2014) Metabolic resource allocation in
individual microbes determines ecosystem interactions and spatial dynamics.
Cell reports 7(4):1104-15 doi:10.1016/j.celrep.2014.03.070
Henry CS, DeJongh M, Best AA, Frybarger PM, Linsay B, Stevens RL (2010) High-
throughput generation, optimization and analysis of genome-scale metabolic
models. Nature biotechnology 28(9):977-82 doi:10.1038/nbt.1672
Henson MA (2015) Genome-scale modelling of microbial metabolism with temporal
and spatial resolution. Biochem Soc Trans 43(6):1164-71
doi:10.1042/BST20150146
Izallalen M, Mahadevan R, Burgard A, Postier B, Didonato R, Jr., Sun J, Schilling CH,
Lovley DR (2008) Geobacter sulfurreducens strain engineered for increased
rates of respiration. Metab Eng 10(5):267-75
doi:10.1016/j.ymben.2008.06.005
Kanehisa M, Goto S (2000) KEGG: kyoto encyclopedia of genes and genomes. Nucleic
Acids Res 28 doi:10.1093/nar/28.1.27
Kanehisa M, Goto S, Sato Y, Furumichi M, Tanabe M (2012) KEGG for integration and
interpretation of large-scale molecular data sets. Nucleic Acids Res
40(Database issue):D109-14 doi:10.1093/nar/gkr988
Kanehisa M, Goto S, Sato Y, Kawashima M, Furumichi M, Tanabe M (2014) Data,
information, knowledge and principle: back to metabolism in KEGG. Nucleic
Acids Res 42(Database issue):D199-205 doi:10.1093/nar/gkt1076
Kanissery RG, Sims GK (2011) Biostimulation for the Enhanced Degradation of
Herbicides in Soil. Applied and Environmental Soil Science 2011:10
doi:10.1155/2011/843450
Kastner M, Miltner A (2016) Application of compost for effective bioremediation of
organic contaminants and pollutants in soil. Appl Microbiol Biotechnol
100(8):3433-49 doi:10.1007/s00253-016-7378-y

King ZA, Drager A, Ebrahim A, Sonnenschein N, Lewis NE, Palsson BO (2015) Escher: A
Web Application for Building, Sharing, and Embedding Data-Rich Visualizations
of Biological Pathways. PLoS computational biology 11(8):e1004321
doi:10.1371/journal.pcbi.1004321
Lee J, Yun H, Feist AM, Palsson BØ, Lee SY (2008) Genome-scale reconstruction and in
silico analysis of the Clostridium acetobutylicum ATCC 824 metabolic network.
Appl Microbiol Biotechnol 80 doi:10.1007/s00253-008-1654-4
Mahadevan R, Bond DR, Butler JE, Esteve-Nunez A, Coppi MV, Palsson BO, Schilling
CH, Lovley DR (2006) Characterization of metabolism in the Fe(III)-reducing
organism Geobacter sulfurreducens by constraint-based modeling. Appl
Environ Microbiol 72(2):1558-68 doi:10.1128/AEM.72.2.1558-1568.2006
Mahadevan R, Lovley DR (2008) The degree of redundancy in metabolic genes is linked
to mode of metabolism. Biophys J 94(4):1216-20
doi:10.1529/biophysj.107.118414
Mahadevan R, Schilling CH (2003) The effects of alternate optimal solutions in
constraint-based genome-scale metabolic models. Metabolic engineering
5(4):264-76
Mani D, Kumar C (2013) Biotechnological advances in bioremediation of heavy metals
contaminated ecosystems: an overview with special reference to
phytoremediation. International Journal of Environmental Science and
Technology 11(3):843-872 doi:10.1007/s13762-013-0299-8
Megharaj M, Ramakrishnan B, Venkateswarlu K, Sethunathan N, Naidu R (2011)
Bioremediation approaches for organic pollutants: a critical perspective.
Environ Int 37(8):1362-75 doi:10.1016/j.envint.2011.06.003
Meyer F, Paarmann D, D'Souza M, Olson R, Glass E, Kubal M, Paczian T, Rodriguez A,
Stevens R, Wilke A, Wilkening J, Edwards R (2008) The metagenomics RAST
server – a public resource for the automatic phylogenetic and functional
analysis of metagenomes. BMC Bioinformatics 9(1):1-8 doi:10.1186/1471-
2105-9-386
Monard C, Vandenkoornhuyse P, Le Bot B, Binet F (2011) Relationship between
bacterial diversity and function under biotic control: the soil pesticide
degraders as a case study. The ISME journal 5(6):1048-56
doi:10.1038/ismej.2010.194
Mongodin EF, Shapir N, Daugherty SC, DeBoy RT, Emerson JB, Shvartzbeyn A, Radune
D, Vamathevan J, Riggs F, Grinberg V, Khouri H, Wackett LP, Nelson KE,
Sadowsky MJ (2006) Secrets of soil survival revealed by the genome sequence
of Arthrobacter aurescens TC1. PLoS Genet 2(12):e214
doi:10.1371/journal.pgen.0020214
Nilsson T, Rova M, Smedja Backlund A (2013) Microbial metabolism of oxochlorates:
a bioenergetic perspective. Biochim Biophys Acta 1827(2):189-97
doi:10.1016/j.bbabio.2012.06.010
Nordberg H, Cantor M, Dusheyko S, Hua S, Poliakov A, Shabalov I, Smirnova T,
Grigoriev IV, Dubchak I (2014) The genome portal of the Department of Energy
Joint Genome Institute: 2014 updates. Nucleic Acids Res 42(Database
issue):D26-31 doi:10.1093/nar/gkt1069
Nousiainen AO, Bjorklof K, Sagarkar S, Nielsen JL, Kapley A, Jorgensen KS (2015)
Bioremediation strategies for removal of residual atrazine in the boreal

groundwater zone. Appl Microbiol Biotechnol 99(23):10249-59 doi:10.1007/s00253-015-6828-2

Oberhardt MA, Zarecki R, Gronow S, Lang E, Klenk HP, Gophna U, Ruppin E (2015) Harnessing the landscape of microbial culture media to predict new organism-media pairings. Nat Commun 6:8493 doi:10.1038/ncomms9493

Oh YK, Palsson BO, Park SM, Schilling CH, Mahadevan R (2007) Genome-scale reconstruction of metabolic network in Bacillus subtilis based on high-throughput phenotyping and gene essentiality data. The Journal of biological chemistry 282(39):28791-9 doi:10.1074/jbc.M703759200

Oliveros JC (2007) VENNY. An interactive tool for comparing lists with Venn Diagrams.

Orth JD, Thiele I, Palsson BO (2010) What is flux balance analysis? Nat Biotech 28(3):245-248 doi:http://www.nature.com/nbt/journal/v28/n3/abs/nbt.1614.html#supplementary-information

Perez-Garcia O, Lear G, Singhal N (2016) Metabolic Network Modeling of Microbial Interactions in Natural and Engineered Environmental Systems. Front Microbiol 7:673 doi:10.3389/fmicb.2016.00673

Price ND, Papin JA, Schilling CH, Palsson BO (2003) Genome-scale microbial in silico models: the constraints-based approach. Trends Biotechnol 21(4):162-9 doi:10.1016/S0167-7799(03)00030-1

Reed JL, Vo TD, Schilling CH, Palsson BO (2003) An expanded genome-scale model of Escherichia coli K-12 (iJR904 GSM/GPR). Genome Biol 4(9):R54 doi:10.1186/gb-2003-4-9-r54

Sagarkar S, Mukherjee S, Nousiainen A, Bjorklof K, Purohit HJ, Jorgensen KS, Kapley A (2013) Monitoring bioremediation of atrazine in soil microcosms using molecular tools. Environ Pollut 172:108-15 doi:10.1016/j.envpol.2012.07.048

Sagarkar S, Nousiainen A, Shaligram S, Bjorklof K, Lindstrom K, Jorgensen KS, Kapley A (2014) Soil mesocosm studies on atrazine bioremediation. J Environ Manage 139:208-16 doi:10.1016/j.jenvman.2014.02.016

Shah J, Dahanukar N (2012) Bioremediation of organometallic compounds by bacterial degradation. Indian J Microbiol 52(2):300-4 doi:10.1007/s12088-011-0223-1

Silva VP, Moreira-Santos M, Mateus C, Teixeira T, Ribeiro R, Viegas CA (2015) Evaluation of Arthrobacter aurescens Strain TC1 as Bioaugmentation Bacterium in Soils Contaminated with the Herbicidal Substance Terbuthylazine. PloS one 10(12):e0144978

Singh B, Singh K (2016) Microbial degradation of herbicides. Crit Rev Microbiol 42(2):245-61 doi:10.3109/1040841X.2014.929564

Strong LC, Rosendahl C, Johnson G, Sadowsky MJ, Wackett LP (2002) Arthrobacter aurescens TC1 metabolizes diverse s-triazine ring compounds. Appl Environ Microbiol 68(12):5973-80

Thiele I, Palsson BO (2010) A protocol for generating a high-quality genome-scale metabolic reconstruction. Nat Protoc 5(1):93-121 doi:10.1038/nprot.2009.203

Thiele I, Vo TD, Price ND, Palsson BO (2005) Expanded metabolic reconstruction of Helicobacter pylori (iIT341 GSM/GPR): an in silico genome-scale characterization of single- and double-deletion mutants. J Bacteriol 187(16):5818-30 doi:10.1128/JB.187.16.5818-5830.2005

578   Udikovic-Kolic N, Scott C, Martin-Laurent F (2012) Evolution of atrazine-degrading
579   capabilities in the environment. Appl Microbiol Biotechnol 96(5):1175-89
580   doi:10.1007/s00253-012-4495-0
581   UniProt C (2015) UniProt: a hub for protein information. Nucleic Acids Res
582   43(Database issue):D204-12 doi:10.1093/nar/gku989
583   Varma A, Palsson BO (1994) Stoichiometric flux balance models quantitatively predict
584   growth and metabolic by-product secretion in wild-type Escherichia coli
585   W3110. Applied and environmental microbiology 60(10):3724-3731
586   Wang Q, Xie S (2012) Isolation and characterization of a high-efficiency soil atrazine-
587   degrading Arthrobacter sp. strain. International Biodeterioration &
588   Biodegradation 71:61-66 doi:10.1016/j.ibiod.2012.04.005
589   Xie S, Wan R, Wang Z, Wang Q (2013) Atrazine biodegradation by Arthrobacter strain
590   DAT1: effect of glucose supplementation and change of the soil microbial
591   community. Environ Sci Pollut Res Int 20(6):4078-84 doi:10.1007/s11356-012-
592   1356-4
593   Xu C, Liu L, Zhang Z, Jin D, Qiu J, Chen M (2013) Genome-scale metabolic model in
594   guiding metabolic engineering of microbial improvement. Appl Microbiol
595   Biotechnol 97(2):519-39 doi:10.1007/s00253-012-4543-9
596   Xu X, Zarecki R, Medina S, Ofaim S, Liu X, Chen C, Hu S, Brom D, Gat D, Porob S,
597   Eizenberg H, Ronen Z, Jiang J, Freilich S (2019) Modeling microbial
598   communities from atrazine contaminated soils promotes the development of
599   biostimulation solutions. The ISME journal 13(2):494-508 doi:10.1038/s41396-
600   018-0288-5
601   Zhang C, Li Y, Wang C, Niu L, Cai W (2015) Occurrence of endocrine disrupting
602   compounds in aqueous environment and their bacterial degradation: A review.
603   Critical Reviews in Environmental Science and Technology 46(1):1-59
604   doi:10.1080/10643389.2015.1061881
605   Zhang Q, Zhu L, Su J, Wang J, Xie H, Wang J, Wang F (2014) Impacts of nitrogen and
606   phosphorus on atrazine-contaminated soil remediation and detoxification by
607   Arthrobacter sp. strain HB-5. Environmental earth sciences 71(3):1465-1471
608   Zhang Y, Cao B, Jiang Z, Dong X, Hu M, Wang Z (2012) Metabolic ability and individual
609   characteristics of an atrazine-degrading consortium DNC5. J Hazard Mater 237-
610   238:376-81 doi:10.1016/j.jhazmat.2012.08.047
611   Zomorrodi AR, Segre D (2016) Synthetic Ecology of Microbes: Mathematical Models
612   and Applications. J Mol Biol 428(5 Pt B):837-61 doi:10.1016/j.jmb.2015.10.019
613   Zou W, Liu L, Zhang J, Yang H, Zhou M, Hua Q, Chen J (2012) Reconstruction and
614   analysis of a genome-scale metabolic model of the vitamin C producing
615   industrial strain Ketogulonicigenium vulgare WSH-001. J Biotechnol 161(1):42-
616   8 doi:10.1016/j.jbiotec.2012.05.015

FIGURE LEGENDS

Figure 1. Atrazine metabolism as implemented in the cytoplasm compartment of *P.*

aurescens TC1 genome scale metabolic model reconstruction (*i*RZ960). External

metabolite are notated with an e0;

Figure 2. Growth performances of *P.aurescens* TC1 in dependence on the availability

of different carbon and nitrogen sources. (A) Predicted and observed *A.aurescens* TC1

growth in mineral minimal media supplemented with Atrazine or its degradation

intermediates. ATZ-Atrazine, NH4, GLU-Glucose, ETH - Ethylamine, ISO –

Isopropylamine, CTRL - control. Values are normalized by dividing in maximal value.

(B) Surface plot rendering *P.aurescens* TC1 growth (biomass) in dependence on

atrazine, glucose and NH4 availability. Color scale is indicative of biomass (Z-axis).

Figure 3. Predicted and observed growth performances of *P. aurescens* TC1 growth in

mineral minimal media with glucose complemented by different amino acids and

atrazine. ATZ-Atrazine, HIS- Histdine, ILE-Isoluecine and MET- Metionine. Observed

values represents mean in triplicates; bars represent SD.

Figure 4. Predicted and observed atrazine degradation in modified media. High and

low phosphate concentrations: 50 and 5 mmol/gdw h for predicted and 0.081 and

0.081 g/l for observed, respectively. ATZ- atrazine, GLU- glucose. Observed values

represents mean in triplicates; bars represent SD.

ADDITIONAL FILES

Additional File 1. Supplementary Figure 1 (tiff). Venn diagram of *P.aurescens* TC1

annotations collected from four established databases.

Additional File 2. Table (xlsx). Reactions added to the model in the process of manual

curation and gap filling.

Additional File 3. Table (xlsx). Alternative carbon and nitrogen sources added to the

minimal mineral media used for *P.aurescens* TC1 growth.

Additional File 4. Metabolic model (sbml). *P.* aurescens TC1 metabolic model (iRZ960)

file.

Additional File 5. Supplementary Figure 2 (tiff). A schematic diagram describing a time

dependent simulation of nutrient dependent bacterial growth.

Additional File 6. Table (xlsx). Default values for exchange reactions as used for all

simulations.

Additional File 7. Table (xlsx). FVA analysis for different metabolites added to an

atrazine containing mineral media.

Additional File 8. Supplementary Figure 3 (tiff). Predicted and observed *P.aurescens*

TC1 growth in media supplemented with high and low phosphate concentrations (50

and 5 mmol/gdw h for predicted and 0.081 and 0.081 g/l for observed respectively).

Simulations and experiments were carried in MMM supplemented with atrazine and

glucose in ATZ-Atrazine, GLU-Glucose. Observed values represents mean in triplicates;

bars represent SD.

A

B