Title: Variation in space and time: a long-term examination of density-dependent dispersal in a woodland rodent

Authors: Denomme-Brown[1], Simon T.; Cottenie[1], Karl; Falls[2], J. Bruce; Falls[2], E. Ann; Brooks[1], Ronald J. and Andrew G. McAdam[1]

[1] University of Guelph, Dept. of Integrative Biology, 50 Stone Rd. E., Guelph, ON, Canada, N1G 2W1

[2] University of Toronto, Dept. of Zoology, Toronto, ON, Canada

Corresponding Author Information: Simon T. Denomme-Brown, denommes@uoguelph.ca, Phone Number: (519) 803-7993, Fax Number: NA

Keywords: Dispersal, Density-dependence, Small Mammals, Temporal Dynamics, Movement

Number of words in Body Text of Manuscript: 5287

Declaration of authorship: STDB, KC and AGM devised the main ideas of this work. STDB wrote the initial draft of the manuscript; all authors contributed substantially to its revision. STDB completed the analysis. JBF, EAF, RJB, STDB, and AGM all contributed to the collection and maintenance of the dataset.

Abstract

Dispersal is a fundamental ecological process that can be affected by population density, yet

studies report contrasting effects of density on propensity to disperse. Additionally, the

relationship between dispersal and density is seldom examined using densities measured at

different spatial scales or over extensive time-series. We used 51-years of trapping data to

examine how dispersal by wild deer mice (*Peromyscus maniculatus*) was affected by changes in

both local and regional population densities. We examined these patterns over both the entire

time-series and also in ten-year shifting windows to determine whether the nature and strength of

the relationship changed through time. Probability of dispersal decreased with increased local and

regional population density, and the negative effect of local density on dispersal was more

pronounced in years with low regional densities. Additionally, the strength of negative density-

dependent dispersal changed through time, ranging from very strong in some decades to absent in

other periods of the study. Finally, while females were less likely to disperse, female dispersal

was more density-dependent than male dispersal. Our study shows that the relationship between

density and dispersal is not temporally static and that investigations of density-dependent

dispersal should consider both local and regional population densities.

Introduction

Dispersal is defined as movement of an individual from a place of origin to one of

reproduction or from one reproductive locale to another (Greenwood 1980), and influences

fundamental ecological processes such as population growth rates and viability, genetic

connectivity, and range expansions (Greenwood 1980; Nathan 2006; Holyoak et al. 2008).

Dispersal is identified as one of four fundamental processes in community ecology (Vellend

2010), and a cornerstone of the metacommunity concept (Leibold et al. 2004). Yet the causes of

dispersal are often poorly understood and dispersal remains difficult to measure and include in

empirical studies (Koenig et al. 1996; Nathan et al. 2003; Logue et al. 2011).

Benefits of dispersal often stem from differences in conspecific densities between pre and

post-dispersal settlement sites. When conspecific densities at pre-dispersal sites are greater than

at post-dispersal sites (i.e. positive density-dependent dispersal), dispersing individuals may

benefit from reduced competition and agonistic interactions (Murray 1967; Gaines and

McClenaghan 1980; Matthysen 2005). Alternatively, negative density-dependent dispersal may

be beneficial when individuals disperse to gain access to a limiting resource, or when the benefits

of group living outweigh the costs of intraspecific competition (Bowler and Benton 2005).

Conflicting examples of positive and negative density-dependent dispersal (DDD) across

a variety of taxa (Bowler and Benton 2005; Matthysen 2005; Rodrigues and Johnstone 2014),

have left little consensus regarding general effects of density on dispersal. Matthysen (2005)

concluded that both birds and mammals predominantly exhibit positive DDD. The review,

however, excluded studies exhibiting temporal trends in density and seasonal variation which

Matthysen (2005) acknowledged could have changed their conclusions. Contradicting effects of

density on dispersal have also been found within species. DDD has been observed as both

negative and positive in separate studies of the same species (Garten and Smith 1974; Rehmeier

et al. 2004), or depending on when in the annual cycle density is measured (Betini et al. 2015), or

simultaneously negative and positive across a patchily distributed population (Kim et al. 2009).

44   In particular, variable effects of density on dispersal have been reported in deer mice,

Peromyscus maniculatus. *Peromyscus* spp. generally display male-biased dispersal (Fairbairn

1978b; Gaines and McClenaghan 1980; Rehmeier et al. 2004) and have been observed to exhibit

both positive (Garten and Smith 1974; Fairbairn 1978b; Anderson and Meikle 2010) and negative

DDD (Rehmeier et al. 2004; Wojan et al. 2017). These variable effects of density on dispersal

suggest that temporal and spatial variation in population densities could feasibly influence the

strength of density dependence in a system. Therefore, as others have proposed (e.g. Matthysen

2005), DDD must be examined using long time-series, while accounting for spatial and temporal

variation in population densities. In this study, we examined the frequency and extent of dispersal

by wild *P. maniculatus*, using monitoring data spanning 50 years from traplines distributed

across nearly 17km of temperate forest. These data allowed us to examine dispersal patterns at

large temporal and spatial scales and investigate how the strength of DDD was affected by both

temporal and spatial variation in population densities.

Materials and methods

Data Collection

Deer mice were live-trapped in Algonquin Provincial Park (APP), Ontario, Canada from 1960 -

2015. Trapping initially occurred on 10 traplines, with lines added and removed intermittently

throughout the study such that the same 17 lines were trapped consistently from 1995 – 2015.

Lines consisted of 10 trapping stations set along transects at 10m intervals. Early in the study,

most lines had one trap at each station, but by 1979 all lines had two traps per station in an

attempt to avoid trap saturation, for a total of 20 traps per line. Sherman-style live traps were used

exclusively until 2013, when one Longworth trap (Natural Heritage Book Society, Totnes,

Devon) replaced one Sherman at each station. Traps were baited with peanut butter and rolled

oats until 1991. After 1991, water-soaked sunflower seeds were used as bait. Freeze-killed

mealworms (*Tenebrio molitor*) were added to half the traps from 2013 - 2015 as a bait

supplement. Use of Longworth traps and mealworms were aimed at decreasing mortality in

shrew (*Soricidae*) species (Do et al. 2013; Shonfield et al. 2013). Distances between lines ranged

from 0.115 - 16.9km. Lines were located in forested habitats with different dominant trees

including sugar maple (*Acer saccharum*), white pine (*Pinus strobus*), white spruce (*Picea*

glauca), black spruce (*Picea mariana*), trembling aspen (*Populus tremuloides*), and red pine

(*Pinus resinosa*).

Each year, trapping was conducted on three consecutive nights either once or twice a

month from May through September, for a maximum of 10 three-night trapping sessions per

year. Traps were set just before dusk and checked after dawn. When captured, deer mice were

weighed, sexed, aged and assessed for reproductive condition. Before release, animals were fitted

with uniquely coded ear tags for individual identification upon recapture.

Dispersal Detection

Dispersal was initially detected when an individual deer mouse was captured on multiple

traplines within a single trapping season. Although these data were collected in a consistent

manner, any errors in data collection or transfer to digital records could mistakenly appear as

dispersal events (e.g. incorrect tag or trapline numbers were recorded). While the error rate is

likely low, the compounding of even a low rate of error over more than 50 years of data could

skew our estimates of dispersal. To minimize the frequency of such errors we used a strict

criterion to determine whether detected movements between traplines represented dispersal

events. To be considered a dispersing individual, an individual mouse must have been captured

on multiple, consecutive occasions on one line either preceded, or followed by at least one

capture on another line. We assumed that the mouse in question was either trapped while moving

towards a trapline where it subsequently settled, or was trapped while dispersing from a trapline

and either continued on or died. To meet this criterion, a mouse must have been captured on a

minimum of three occasions within a year. Accordingly, all model analyses and calculations of

dispersal frequencies discussed hereafter were performed on a dataset of mice caught three or

more times in a year. Additionally, we avoided including mis-read tags by ensuring that

dispersing animals had consistent records of individual traits such as sex and size. Instances

where an animal returned to their original line of capture following a capture on a different line

were not considered to be dispersal events. These could have represented nomadic wandering

between lines rather than dispersal or this could have been caused by a misread tag or data entry

error. Animals last caught on their initial line of capture were therefore not considered to have

dispersed (n = 136).

Movements occurring between years were not considered as few mice survive the winter

in APP, and thus there were relatively few potential dispersers over this period. Movements

between lines were considered to be potential dispersal events as the distance between the closest

traplines (115m) was greater than the diameter of home ranges for this species typically reported

in the literature (Stickel 1968; Bowers and Smith; 1979; Harestad and Bunnel 1979; Van Horne

1980; Abramson et al 2006; Wood 2010). Distances between traplines were calculated as

straight-line distance between the first trap station on each line. Finally, changes in methodology

and some loss of data necessitated the exclusion of five years of data from the dataset (1987-

1991).

Density Measures

We converted raw capture totals to the number of captures per hundred trap-nights per

year to account for variation in trapping effort both between years and among traplines. The first

measurement was raw local trapline density (hereafter termed raw local density. We converted

these raw local densities into two types of densities for the statistical analyses that determined the

independent effect of regional and local density on dispersal. The first density measure was the

average regional trapline density, calculated annually as the mean number of captures per

hundred trap-nights across all traplines in APP in that year (or regional density) Second, we

calculated relative local density (hereafter local density) for each trapline in each year by

subtracting the regional density in a year from the raw local density on each line that same year.

This local density measure thus captures the additional effect of local densities after accounting

for year-to-year fluctuations in regional desntiy associated with for instance seed mast events

(Falls et al. 2007). This relative local density thus allowed us to include uncorrelated local

densities and regional density estimates within the same models (*sensu* van de Pol and Wright

2009).

It should be noted that the density estimates we used do not account for spatial or

temporal variation in capture probability. While spatially explicit mark-recapture techniques

(SECR) represent a powerful set of statistical tools to account for these issues (Royle et al. 2013),

they were inappropriate for our long-term data for several reasons. First, Bayesian SECR models

are inappropriate for trapping data that are derived from single-catch traps such as those used in

our study (Gerber and Parmenter 2015). Second, when using frequentist SECR models the multi-

catch estimator can only be used if density is relatively constant over survey regions (Distiller

and Borchers 2015), and this was not the case in our data (see Results). Finally, the single-catch

estimator developed by Distiller and Borchers (2015), which allows for frequentist SECR

analysis using single catch traps, requires times for all captures which were not available for our

historical data. We discuss this issue further below.

Data Analysis

To assess how a deer mouse's propensity to disperse was affected by sex and population

density at different spatial scales, we fitted eight separate Generalized Linear Mixed Effects

Models (binomial family, logit link) associated with local and regional population densities as

well as sex of the individual mouse. They were 1) Biological Null Model, 2) Sex Model, 3) Local

Density Model, 4) Regional Density Model, 5) Raw Density Model (RWD), 6) Local and

Regional Density Model, 7) Density Interaction Model (DI), 8) Density and Sex Interaction

Model (DSI) (see Table 1). All eight models used the binary response variable of whether an

individual mouse dispersed or not and included the fixed effect of distance (in meters) from the

trapline where the mouse was initially caught to the next closest trapline. This effect was

included to account for the non-uniform spatial distribution of traplines and whether differences

in movement among lines were due primarily to spatial effects on the probability of detecting a

dispersal event (i.e. dispersal is more likely to be detected between nearby traplines). We used the

natural log of the raw distance in kilometers (km) to correct a significant non-linearity in the

relationship between this variable and the probability of dispersal. In all models, local density and

regional density were converted to z-scores ($z = x - \bar{x}/\sigma$) based on the mean and standard

deviation of all observations across years and lines.

We used the Akaike Information Criterion (AIC) to assess relative support for the eight

models (Burnham and Anderson 2007), which included various combinations of sex, and raw,

local and regional density as fixed effects (Table 1). We calculated AIC values for each model

and performed model comparisons using ΔAIC (Δ_i) values (Table 1). Akaike Weights (ω_i) were

calculated to examine the conditional probabilities of each model (Table 1). All models contained

the same random effects, with initial trapline of capture and year of capture included to capture

spatial and temporal variation in dispersal propensity. Likelihood ratio tests (LRT) were used to

assess the significance of the year and trapline random effects in each of the 8 candidate models.

Models were assessed for overdispersion and collinearity of variables and met model

assumptions. Statistical analyses were performed in R (Version 3.5; R Core Team 2018),

including the glmer function with a bobyqa optimizer in the lme4 package (Bates et al. 2015).

In addition to the model analysis described above, we performed a number of *post hoc*

analyses. When first visualizing the relationship between regional density and movement

frequency (Figure 1a) there appeared to be a marked change in the relationship between the two

variables for portions of the time-series, particularly over a 9-year period from 2002-2010. To

examine this, we ran model with raw local density as a fixed effect and *year* and *trapline* of

initial capture as random effects on a shifting ten-year window through the time-series. This

model was used because the RWD model was the simplest model with moderate support based

on AIC comparisons and contained only raw local density and sex as fixed effects. More

complex models would not fit these data given the smaller sample size provided in 10-year

windows of the time-series (Figure 1b).

The second set of *post hoc* analyses assessed how robust our results were to changes in

trapping protocols during this long-term study (e.g. number of traps per station, change in bait,

and trap types), that occurred at distinct points in our study (1979, 1991 and 2013 respectively).

For this we ran our best model on subsets of our dataset from before, between and after these

protocol changes occurred.

Results

Population Densities and Frequency of Movement

Over the 51 years included in this study, there were 3408 instances where a mouse was

caught at least three times in a year. Of these 3408 possible dispersers, 4.2% (n = 142) dispersed

between traplines. The proportion of the population that dispersed fluctuated between years ($\bar{x} \pm$

SE = 0.05 ± 0.008 mice dispersing in the population, n = 51 years), with no dispersal detected in

some years and a maximum of 28.6% of the population dispersing between lines in 1964 (Figure

1). The population sex ratio was slightly skewed, with males comprising 53.4% of individuals in

the dataset ($X^2 = 7.9$, df = 1, P < 0.01). This male skew was more pronounced among dispersing

individuals, with 64.8% of dispersing mice being male ($X^2 = 7.7$, df = 1, P < 0.01). Overall,

males were more likely to disperse (5.1%) than females (2.7%; Fisher Exact Test: P = 0.006).

Distances travelled by dispersing individuals ranged from 115m to 11.4 km ($\bar{x} \pm$ SE =

1.68 ± 0.20 km, n = 142) (Figure 2). There was no difference in dispersal distance between the

sexes (t = 0.98, df = 92, p-value = 0.3) (Figure 2). Only 51.4% (n = 73) of dispersals were to the

next most closely situated trapline (Figure 2). There were two traplines located only 115m apart,

but dispersal between these traplines represented only 16.9% (n = 24) of all movements.

The density of deer mice in APP fluctuated over an order of magnitude during this study,

with yearly regional population densities ranging from 1.9 to 27.8 captures per hundred trap

nights ($\bar{x} \pm$ SE = 11.15 ± 0.81 captures per hundred trap-nights, n = 51 years) (Figure 1a). Local

densities also varied, with the average range in local population densities among traplines within

years ($\bar{x} \pm$ SE = 19.37 ± 1.22 captures per hundred trap-nights, n = 51 years) approaching the

aforementioned range in regional densities among years. In particular, 9 of 51 years exhibited a

greater range in density among lines within a year than the range in regional density across the 51

years, with a maximum range of 47.2 captures per hundred trap-nights in 2001.

Dispersal Models

The model that had the lowest AIC score of the eight fitted models was the DSI model

which included the fixed effects of both regional and local density, sex, and interactions between

both local density and regional density, and local density and sex (Table 2). In the DSI model, the

probability of dispersal decreased with increasing distance to nearest trapline ($\beta \pm$ SE = -1.035 $\pm$

0.15, Z = -6.85, P < 0.0001), increased regional density ($\beta \pm$ SE = -0.63 ± 0.16, Z = -3.9, P <

0.0001), and increased local density ($\beta \pm SE$ = -0.091 ± 0.026, Z -3.52, P = 0.0004). The

significant positive interaction between local and regional densities ($\beta \pm SE$ = 0.036 ± 0.015, Z =

2.44, P = 0.015) indicated that in years of low regional density, local densities had a stronger

negative effect on the probability of dispersal (Figure 3a). For example, three of the five years

displaying the highest proportions of dispersers in the study were 1964 (0.286), 1962 (0.192) and

1976 (0.150), which represented three of the 11 lowest density years within the study.

The DSI model also included sex as fixed effect and while more males dispersed than

females, the effect of sex on dispersal was not significant ($\beta \pm SE$ = 0.32 ± 0.19, Z = 1.69, P =

0.09) (Figure 3b). However, the interaction between local density and sex was in fact significant

($\beta \pm SE$ = 0.064 ± 0.031, Z = 2.1, P = 0.036), indicating that while males disperse more than

females, females were more sensitive to local density than males and were more likely to disperse

when densities were low (Figure 3b). Initial *line of capture* was a significant random effect in this

model (LRT, Deviance = 1042, X^2_1= 11.2, P = 0.0008) whereas *year* was not (LRT, Deviance =

1042, X^2_1 = 0.2, P = 0.65). Overall, the DSI model explained 11.77% of the deviance.

There was nearly strong support ($\Delta_i < 2$) for the DI model($\Delta_i = 2.2$)   which was identical

to the DSI model except it did not include an interaction between local density and sex. All

effects in this model were in the same direction as in the DSI model and the DI model explained

10.05% of the overall deviance. Considerably less support was seen for models that did not

contain the interaction between local and regional densities, with all these models demonstrating

either moderate ($\Delta_i < 7$) or no support ($\Delta i > 10$) (Table 1).

Despite evidence of negative DDD overall, visual inspection of the time series of density

and dispersal probability (Figure 1a) suggested temporal changes in the strength of density-

dependent dispersal. When running the simplified model containing raw local density as a fixed

effect and *year and trapline* of initial capture as random effects on the shifting 10-year window

through the time series, the effect of raw density on dispersal probability never became positive.

The effect size did approach zero, however, and the negative relationship between raw density

and dispersal probability was significant for only 13 of 32 ten-year windows in the analysis

(Figure 1b). Finally, when running the DSI model on our data before and after each change in

trapping methodology, there were no changes in the direction of effects for all explanatory

variables save one. The exception was the effect of sex on dispersal, which was reversed prior to

the initial bait change, but the effect was not significant.

Discussion

Probability of Dispersal

Overall, deer mice in APP exhibited negative DDD, with a lower probability of dispersal

at high densities, regardless of the scale at which density was measured. Proposed mechanisms

for negative DDD include attraction to conspecifics (Danielson and Gaines 1987; Stamps 1991),

range constriction in the face of high densities (Stickel 1960; Taitt 1981) and exclusionary

behavior towards immigrants in periods of high density (i.e. "social fences"; Hestbeck 1982).

Had trap saturation occurred in APP it could mimic a natural social fence, denying immigrants

access to traps on densely populated lines and biasing our results towards the appearance of

negative DDD. This seems unlikely, however, as densities on even the most highly populated

traplines fell well short of saturation (i.e. average captures per 100 trap-nights across all lines

throughout this study was 22.62 with SE = 0.42 and SD = 10.66; the highest capture rate on any

line in any year was 73.81 captures per 100 trap-nights). Additionally, there was no obvious

change in the direction of density-dependence after the traps per station were doubled in 1979.

While the result of negative DDD runs counter to generalizations of positive DDD in

mammals (Matthysen 2005), previously reported relationships between dispersal and density in

Peromyscus spp. are inconsistent. Positive DDD has been observed in *P. maniculatus* (Fairbairn,

1978a), *P. polionotus* (Garten and Smith 1974), and genetic estimates show *P. leucopus* is more

likely to disperse from high density habitat patches (Anderson and Meikle 2010). Alternatively,

there is evidence of negative DDD in both *P. maniculatus* (Rehmeier et al. 2004) and *P. boylii*

(Wojan et al. 2015). The spatial and temporal breadth of our study may elucidate why studies on

Peromyscus spp., and other dispersing organisms, often yield divergent results. For while deer

mice in our study consistently exhibited negative DDD, the strength with which density affected

dispersal varied temporally, spatially and between sexes.

The relationship between density and dispersal varied throughout our time-series. This

variation is evident when examining the time-series of regional density and movement frequency

together (Figure 1a), particularly the period from 2002-2010 which visually suggests a possible

positive relationship. The effect of density on dispersal probability in the RWD model was never

positive in any 10-year window along the time-series, however, the effect size approached zero in

the 1990's before becoming increasingly negative (Figure 1b). Also, the majority of the 10-year

windows examined (n = 19) did not exhibit significant relationships between density and

dispersal. Few studies investigating DDD have used long-term data (Matthysen 2005), despite the

fact that inferences drawn from short-term studies can differ drastically from those generated by

long-term research (Kratz et al. 2003), and that effects of some ecological processes become

more pronounced over time (Cardinale et al. 2007). In an experiment using guppies (*Poecilia*

reticulata), De Bona et al. (2019) recently demonstrated that over the course of a colonization

event, density-dependent dispersal can change in strength and direction. Still, our findings are the

first to demonstrate that the density-dispersal relationship can vary through time in an

unmanipulated system and over large time-series. This highlights the value of long-term studies

for drawing accurate conclusions regarding the density-dispersal relationship, as an unfortunately

timed relatively long-term study (i.e. 10 years) would not detect the negative effect of density on

dispersal that exists in this system.

In addition to varying temporally, the strength with which local density affected dispersal

was mediated by regional densities. This interaction between the regional and local density

effects on dispersal indicates that the negative effect of local density on dispersal is greater in

years of low regional density and muted in years of high regional density (Figure 3a). This

suggests that local density can be an important driver of dispersal, but only when examined in

concert with regional density, as low local densities that increase the probability of dispersal are

only noteworthy in years when regional densities are also low. Local and regional densities acting

as interacting drivers of dispersal has been observed experimentally (De Bona et al. 2019) but

never before in natural, unmanipulated systems. Considering local densities within the context of

the accompanying regional densities is important moving forward, as many studies provide only

densities for localized sites of emigration (Matthysen 2005), which based on our findings may

lead to spurious results when not considered in the context of associated regional densities. As

well, the few large datasets that have been used to examine DDD fail to differentiate between

spatial and temporal variation in densities (Matthysen, 2005; but see Pasinelli and Walters 2002;

Catchpole et al. 2004) despite the fact both temporal variation and interactions between densities

at multiple spatial scales affect dispersal.

Several potential mechanisms could cause this interaction between the effects of local and

regional densities on dispersal. Deer mice populations in APP are closely tied to maple seed

crops, with mouse populations peaking after seed masting (Falls et al. 2007). *Peromyscus* spp.

also move based on habitat quality and population density (Morris and Diffendorfer 2004). It

may be that individuals select habitat and disperse to high quality, maple-rich areas when

possible; but in years of high regional densities, local densities in high-quality habitats rise

enough to prevent such immigration. This process would strike a balance between traditional

dispersal hypotheses predicting positive DDD due to increased competition (Greenwood 1980),

and ideas of conspecific attraction (Stamps 1991) and habitat selection. This balance between

competition and facilitation amongst conspecifics has been observed previously in blue-footed

booby (*Sula nebouxii*), where simultaneous positive and negative DDD results from the costs of

competition and the advantages of increased mating choice options (Kim et al. 2009).

311   Mate availability may also influence dispersal in APP as the strength of density-

dependence also differed between sexes. Males dispersed more often than females, which is

consistent with findings of male-biased dispersal in both *Peromyscus* spp. (Fairbairn 1978a;

Goundie and Vessey 1986; Rehmeier et al. 2004; Mabry 2014), and other mammals (Greenwood

1980; Dobson 1982; Mabry et al. 2013). However, we also found evidence that dispersal in

females was more sensitive to changes in local density than in males, with females dispersing

more frequently than males when local densities were very low. Examples of sex-biased dispersal

switching from male-bias to female-bias within a system are rare (but see Pérez-González and

Carranza 2009), but have implications for understanding how mating strategies influence

dispersal. The predominance of male-biased dispersal in mammals (and female-bias in birds) has

long been attributed to mating systems, although the simplicity of this relationship has been

questioned (Mabry et al. 2013). If males disperse to track females and females rely on available

resources for reproduction (Emlen and Oring 1977), then at exceedingly low densities the lack of

available mates may outweigh the benefits of familiar surroundings, resulting in increased female

dispersal. Alternatively, if low densities indicate low food availability, this would affect female

dispersal to a greater extent due to their reliance on resources for reproduction (Emlen and Oring

1977). Differences between male and female dispersal sensitivity to local densities mean that

spatial variation in density could potentially affect sex ratios, in turn affecting population density.

As mentioned previously, our density measures did not account for variation in capture

probabilities. While we described the practical reasons for not being able to utilize SECR

methods, we recognize that this is not ideal. However, the lack of trap saturation on any of our

lines throughout the study, the fact that frequency of recapture of individuals within this study ($\bar{x}$

$\pm$ SE = 2.63 $\pm$ 0.07) did not decrease with increased regional annual densities (linear regression; β

$\pm$ SE = 0.012 $\pm$ 0.01, t = 1.09, P = 0.28), and the fact that the density-dispersal relationships in

our models were not altered by the changes in trapping methodologies that occurred throughout

this study all speak to the robustness of our findings.

Frequency and Distance of Dispersal

Generally, deer mice dispersal occurred as often as expected based on existing literature,

but to greater distances. Rehmeier et al. (2004) reported that over a 10-year period 4% of deer

mice in a tallgrass prairie undertook long-distance dispersals (LDDs, >125m); Bowman et al.

(2001b) similarly reported 4.2% of deer mice demonstrating LDD (>100m) in temperate forests.

Based on these papers, our dispersals (>115m) can be classified as LDDs and the frequency of

LDD by deer mice in APP (4.2%) was remarkably similar to these frequencies reported in the

literature. The dispersal distances in our study exceeded previous reports for *P. maniculatus*, with

13% (n =20) of observed dispersals exceeding the previous reported maximum long-distance

movement (4287 $\pm$ 10m, in Wood and McKinney 2015). Our observed maximum dispersal

distance of 11.3km represents a 164% increase on that previous maximum. As well, we detected

more frequent movements of great distances than previous studies, with 3.1% of mice dispersing

over 200m compared to 1% of mice dispersing beyond 200m in previous studies (Brant 1962;

Groves and Keller 1986). We also observed 2.7% and 2.1% of mice dispersing beyond 300m and

500m respectively, while roughly 1% of populations travelled those distances in previous studies

(Diffendorfer et al. 1999; Rehmeier et al. 2004). This indicates that the tail of the dispersal

distance distribution of this species has previously been underestimated.

Dispersal distances tend to follow leptokurtic distributions; many individuals disperse

short distances with few LDDs. Understanding the rate and extent of LDDs, represented by the

tail of the distribution of dispersal distances, is important as they determine the spread of a

species in a landscape (Kot et al. 1996; Levin et al. 2003; Green and Figuerola 2005). Species

with "fat-tailed" dispersal distributions exhibit more rapid spread throughout landscapes (Kot et

al. 1996). This can influence persistence in the face of climate change (Schloss et al. 2012),

which small mammals in this region are already responding to (Myers et al. 2009), and the spread

of zoonotic diseases (Mundt et al. 2009), of which *Peromyscus* spp. are vectors (Meerburg et al.

2009; Gaitan and Millien 2016). These distribution tails are notoriously difficult to quantify

(Koenig et al. 1996), and LDD in deer mice is generally reported anecdotally or with little detail

(Jung et al. 2005; Wood and McKinney 2015). Our absolute frequencies of dispersal should be

interpreted cautiously as mark-recapture studies have limited ability to detect dispersal. The

majority of small mammal studies utilize trapping grids too small to detect LDDs (< 2ha, from

Bowman et al. 2001a), and comparisons with radio-tracking and genetic dispersal estimates show

that mark-recapture methods underestimate dispersal distances (Koenig et al. 1996).

Unfortunately, expanding trapping grids is often impractical, as increasing the maximum distance

across a grid exponentially increases area and monitoring effort. Our study takes an alternate

approach by instead using many smaller transects, reducing our ability to detect individual

dispersal events, but overcoming this via extensive temporal replication. Our results suggest that

deer mice LDDs are more common than the literature suggests, reinforcing claims that relatively

small-scale mark-recapture monitoring techniques fail to accurately capture LDD (Koenig et al.

1996).

We have demonstrated that deer mice exhibit negative DDD that is more pronounced

during periods of low regional densities and that female dispersal is more sensitive to these

changes in density than that of males. We also have demonstrated that the relationship between

density and dispersal is not static but rather varies conspicuously over large time scales. These

results show that long-term field studies are beneficial to the examination of ecological processes

such dispersal, as smaller scale projects are likely to miss or potentially misrepresent the broader

signal entirely. As well, our findings suggest that while local population densities are important,

their influence on dispersal can be affected by broader regional population densities. As such,

future studies should account for both local and regional densities when testing for DDD.

Additionally, we have shown that well-established phenomena such as mammalian male-biased

dispersal may also be density-dependent. These results together suggest that the conflicting

relationships between density and dispersal in the literature may result from broader context-

dependence in DDD. Finally, our results add to a previously modest body of empirical literature

examining LDD, by demonstrating that LDD by deer mice is not uncommon. This reinforces that

dispersal distributions estimated from short-term mark recapture studies may be underestimated,

which is potentially important for future conservation.

Acknowledgements

We would like to acknowledge countless individuals who collected these data. Thank you to the

Algonquin Wildlife Research Station for logistical support. This work was supported through

funding from the National Science and Engineering Research Council of Canada and the Ontario

Ministry of Natural Resources. We dedicate this work to the memory of our colleague Dr. Leslie

Rye who said it best: "I can't believe I am getting paid to work in Algonquin!"

References

Abramson G, Giuggioli L, Kenkre VM, Dragoo JW, Parmenter RR, Parmenter CR, Yates TL

(2006) Diffusion and home range parameters for rodents: Peromyscus maniculatus in

New Mexico. Ecol Complex 3: 64-70 doi: 10.1016/j.ecocom.2005.07.001

Anderson CS, Meikle DB (2010) Genetic estimates of immigration and emigration rates in

relation to population density and forest patch area in Peromyscus leucopus. Cons Gen

11:1593–1605 doi: 10.1007/s10592-009-0033-8

Bates D, Maechler M, Bolker B, Walker S (2015) Fitting linear mixed-effects models using lme4.

J Stat Soft 67:1–48 doi: 10.18637/jss.v067.i01

Betini GS, Pardy A, Griswold CK, Norris DR (2015) The role of seasonality and non-lethal

carry-over effects on density-dependent dispersal. Ecosphere 6:art272 doi: 10.1890/ES15-

00257.1

Bowers MA, Smith HD (1979) Differential habitat utilization by sexes of deermouse,

Peromyscus maniculatus. Ecology 60:869-875 doi: 10.2307/1936854

Bowler DE, Benton TG (2005) Causes and consequences of animal dispersal strategies: relating

individual behaviour to spatial dynamics. Biol Rev Camb Philos Soc 80:205–225 doi:

10.1017/S1464793104006645

Bowman J, Corkum CV, Forbes GJ (2001a) Spatial scales of trapping in small-mammal research.

Can Field-Nat 115:472–475

Bowman J, Forbes G, Dilworth T (2001b) Distances moved by small woodland rodents within

large trapping grids. Can Field-Nat 115:64-67

Brant DH (1962) Measures of the movements and population densities of small rodents. Univ

Calif publ zool 62:105–184

Burnham KP, Anderson DR (2007) Model selection and inference: A practical information-

theoretic approach, 2nd edn. Springer, New York, NY

Cardinale BJ, Wright JP, Cadotte MW, Carroll IT, Hector A, Srivastava DS, Loreau M, Weis JJ

(2007). Impacts of plant diversity on biomass production increase through time because of

species complementarity. Proc Natl Acad Sci 104:18123–18128

doi: 10.1073/pnas.0709069104

Catchpole EA, Fan Y, Morgan BJT, Coulson TN, Clutton-Brock TH (2004). Sexual dimorphism,

survival and dispersal in red deer. J Agric Biol Environ Stat 9:1–26

doi: 10.1198/1085711043172

Danielson BJ, Gaines MS (1987) The influences of conspecific and heterospecific residents on

colonization. Ecology 68:1778–1784 doi: 10.2307/1939869

De Bona S, Bruneaux M, Lee AEG, Reznick DN, Bentzen P, López-Sepulcre A

(2019) Spatio-temporal dynamics of density-dependent dispersal during a population

colonisation. Ecol Lett doi: 10.1111/ele.13205

Diffendorfer JE, Gaines MS, Holt RD (1999) Patterns and impacts of movements at different

scales in small mammals. In: Barrett GW, Peles JD (eds) Landscape ecology of small

mammals. Springer-Verlag, New York, pp 63-88

Distiller G, Borchers DL (2015) A spatially explicit capture-recapture estimator for single-catch

traps. Ecol Evol 21:5074-5087 doi: 10.1002/ece3.1748

Do R, Shonfield J, McAdam AG (2013) Reducing accidental shrew mortality associated with

small-mammal livetrapping II: a field experiment with bait supplementation. J Mammal

94:754-760 doi: 10.1644/12-MAMM-A-242.1

Dobson FS (1982) Competition for mates and predominant juvenile male dispersal in mammals.

Anim Behav 30:1183–1192 doi: 10.1016/S0003-3472(82)80209-1

Emlen ST, Oring LW (1977) Ecology, sexual selection and the evolution of mating systems.

Science 197:215–223 doi: 10.1126/science.327542

Fairbairn, DJ (1978a). Dispersal of deer mice, Peromyscus maniculatus - Proximal Causes and
Effects on Fitness. Oecologia 193:171–193 doi: 10.1007/BF00366070.

Fairbairn DJ (1978b) Behaviour of dispersing deer mice (Peromyscus maniculatus). Behav Ecol
Sociobiol 3:265–282 doi: 10.1007/BF00296313

Falls JB, Falls EA, Fryxell JM (2007) Fluctuations of deer mice in Ontario in relation to seed
crops. Ecol Monogr 77:19–32 doi: 10.1890/05-1485

Gaines MS, McClenaghan LR (1980) Dispersal in small mammals. Annu Rev Ecol Syst 11:163–
196 doi:10.1146/annurev.es.11.11080.001115

Gaitan J, Millien V (2016) Stress level, parasite load, and movement pattern in a small-mammal
reservoir host for Lyme disease. Can J Zool 94:565–573 doi: 10.1139/cjz-2015-0225

Garten CT, Smith MH (1974) Movement by oldfield mice and population regulation. Acta
Theriol 19:513–514

Gerber BD, Parmenter RR (2015) Spatial capture-recapture model performance with known
small-mammal densities. Ecol Appl 25:695-705 doi: 10.1890/14-0960.1

Goundie TR, Vessey SH (1986) Survival and dispersal of young white-footed mice born in nest
boxes. J Mammal 67:53–60 doi: 10.2307/1381001

Green AJ, Figuerola J (2005) Recent advances in the study of long-distance dispersal of aquatic
invertebrates via birds. Divers Distrib 11:149–156 doi: 10.1111/j.1366-
9516.2005.00147.x

Greenwood PJ (1980) Mating systems, philopatry and dispersal in birds and mammals. Anim
Behav 28:1140–1162 doi:10.1016/S0003-3472(80)80103-5

Groves CR, Keller BL (1986) Ecological characteristics of small mammals on a radioactive

waste disposal area in southeastern Idaho. Am Midl Nat 46:404–410 doi:

10.2307/2425405

Harestad AS, Bunnel FL (1979) Home range and body weight -- a reevaluation. Ecology 60:389–

402 doi: 10.2307/1937667

Hestbeck JB (1982) Population regulation of cyclic mammals: the Social Fence Hypothesis.

Oikos 39:157–163 doi: 10.2307/3544480

Holyoak MR, Casagrandi R, Nathan R, Revilla E, Spiegel O (2008) Trends and missing parts in

the study of movement ecology. Proc Natl Acad Sci USA 105:19060–19065 doi:

10.1073/pnas.0800483105

Jung T, O'Donovan K, Powell T (2005) Long-distance movement of a dispersing deer mouse,

Peromyscus maniculatus, in the boreal forest. Can Field-Nat 119:451–452 doi:

10.22621/cfn.v119i3.161

Kim SY, Torres R, Drummond H (2009) Simultaneous positive and negative density-dependent

dispersal in a colonial bird species. Ecology 90:230–239 doi: 10.1890/08-0133.1

Koenig W, Van Vuren D, Hooge PN (1996) Detectability, philopatry and the distribution of

dispersal distances in vertebrates. Trends Ecol Evol 11:514–517 doi: 10.1016/S0169-

5347(96)20074-6

Kot M, Lewis MA, Van Den Driessche P (1996) Dispersal data and the spread of invading

organisms. Ecology 77:2027–2042 doi: 10.2307/2265698

Kratz TK, Deegan LA, Harmon ME, Lauenroth WK (2003) Ecological variability in space and

time: insights gained from the US LTER program. BioScience 53:57–67 doi: 10.1641/0006-

3568(2003)053[0057:EVISAT]2.0.CO;2

Leibold MA, Holyoak M, Mouquet N, Amarasekare P, Chase JM, Hoopes MF, Holt RD, Shurin

JB, Law R, Tilman D, Loreau M, Gonzalez A (2004) The metacommunity concept: a

framework for multi-scale community ecology. Ecol Lett 7:601–613 doi: 10.1111/j.1461-

0248.2004.00608.x

Levin SA, Muller-Landau HC, Nathan R, Chave J (2003) The ecology and evolution of seed

dispersal: a theoretical perspective. Annu Rev Ecol Evol Syst 34:575–604

doi:10.1146/annurev.ecolsys.34.011802.132428

Logue JB, Mouquet N, Peter H, Hillebrand H (2011) Empirical approaches to metacommunities:

a review and comparison with theory. Trends Ecol Evol 26:482–91 doi:

10.1016/j.tree.2011.04.009

Mabry, KE (2014) Effects of sex and population density on dispersal and spatial genetic structure

in brush mice. J Mammal 95:981–991 doi: 10.1644/14-MAMM-A-008

Mabry KE, Shelley EL, Davis KE, Blumstein DT, van Vuren DH (2013) Social mating system

and sex-biased dispersal in mammals and birds: a phylogenetic analysis. PLoS ONE 8:1–

9 doi: 10.1371/journal.pone.0057980

Matthysen E (2005) Density-dependent dispersal in birds and mammals. Ecography 28:403–416

doi: 10.1111/j.0906-7590.2005.04073.x

Meerburg BG, Singleton GR, Kijlstra A (2009) Rodent-borne diseases and their risks for public

health. Crit Rev Microbiol 35:221-270 doi: 10.1080/1040841090298983

Morris DW, Diffendorfer JE (2004) Reciprocating dispersal by habitat-selecting white-footed

mice. Oikos 107:549–558 doi: 10.1111/j.0030-1299.2004.12895.x

Mundt CC, Sackett KE, Wallace LD, Cowger C, Dudley JP (2009) Long-distance dispersal and

accelerating waves of disease: empirical relationships. Am Nat 173:456–466 doi:

10.1086/597220.

Murray BGJ (1967) Dispersal in vertebrates. Ecology 48:975–978 doi: 10.2307/1934544

Myers P, Lundrigan BL, Hoffman SMG, Haraminac AP, Seto SH (2009) Climate-induced

changes in small mammal communities of the Northern Great Lakes Region. Glob

Change Biol 15:1434-1445 doi: 10.1111/j.1365-2486.2009.01846.x

Nathan R (2006) Long-distance dispersal of plants. Science 313:786–788 doi:

10.1126/science.1124975

Nathan R, Perry G, Cronin JT, Strand AE, Cain ML (2003) Methods for estimating long-distance

dispersal. Oikos 103:261–273 doi: 10.1034/j.1600-0706.2003.12146.x

Pasinelli G, Walters JR (2002) Social and environmental factors affect natal dispersal and

philopatry of male Red-cockaded Woodpeckers. Ecology 83:2229–2239 doi:

10.1890/0012-9658(2002)083[2229:SAEFAN]2.0.CO;2

Pérez-González J, Carranza J (2009) Female-biased dispersal under conditions of low male

mating competition in a polygynous mammal. Mol Ecol 18:4617–4630 doi:

10.1111/j.1365-294X.2009.04386.x.

Rehmeier RL, Kaufman GA, Kaufman DW (2004) Long-distance movements of the deer mouse

in tallgrass prairie. J Mammal 85:562–568 doi: 10.1644/1383956

Rodrigues AMM, Johnstone RA (2014) Evolution of positive and negative density-dependent

dispersal. Proc R Soc Lond B Biol Sci 281:20141226–20141226 doi:

10.1098/rspb.2014.1226

Royle JA, Chandler RB, Sollmann R, Gardner B (2013) Spatial capture-recapture. Academic

Press

Schloss CA, Nuñez TA, Lawler JJ (2012) Dispersal will limit ability of mammals to track climate

change in the Western Hemisphere. Proc Natl Acad Sci USA 109:8606–8611 doi:

10.1073/pnas.1116791109.

Shonfield J, Do R, Brooks RJ, McAdam AG (2013) Reducing accidental shrew mortality

associated with small-mammal livetrapping I: an inter- and intrastudy analysis. J Mammal

94: 745-753 doi: 10.1644/12-MAMM-A-271.1

Stamps JA (1991) The effects of conspecifics on habitat selection in teritorial species. Behav

Ecol Sociobiol 28:29–36 doi: 10.1007/BF00172136

Stickel LF (1960) Peromyscus ranges at high and low population densities. J Mammal 41:433–

441 doi: 10.2307/1377530

Stickel LF (1968) Home range and travels. In: King JA (eds) Biology of peromyscus (rodentia).

Special publication no. 2, Stillwater, OK pp: 373-411

Taitt MJ (1981) The effect of extra food on small rodent populations: I. deermice (Peromyscus

maniculatus). J Anim Ecol 50:111–124 doi: 10.2307/4035

van de Pol, M. and J. Wright (2009) A simple method for distinguishing within- versus between-

subject effects using mixed models. Anim Behav 77: 753-758 doi:

10.1016/j.anbehav.2008.11.006

Van Horne B (1980) Deomgraphy of Peromyscus maniculatus populatons in seral stages of

coastal coniferous forest in southeast Alaska. Can J Zool 59:1045-1061 doi: 10.1139/z81-

146

Vellend M (2010) Conceptual synthesis in community ecology. Q Rev Biol 85:183–206 doi:

10.1086/652373

Wojan CM, Knapp SM, Mabry KE (2015) Spatial variation in population density affects

dispersal behavior in brush mice. Ecology 96:1661-1669 doi: 10.1890/14-1661.1

Wood BA, Cao L, Dearing MD (2010) Deer mouse (Peromyscus maniculatus) home-range size

and fidelity in sage-steppe habitat. West N Am Nat 70:345-354 doi:

10.3398/064.070.0307

Wood CM, McKinney ST (2015) Record long-distance movement of a deer mouse, Peromyscus

569 maniculatus, in a New England montane boreal forest. Can Field-Nat 129:10–11 doi:

570 10.22621/cfn.v129i2.1699

571   Table 1: Eight generalized linear mixed effects models (binomial response, logit link) fitted to the
572   data for predicting the probability of an individual deer mouse dispersing

Model	K	AIC	ΔAIC	LogLik	AIC ω	Rank
Density and sex interaction model (DSI): ~ Local Density + Regional Density[+] + Sex[‡] + Closest Distance[¶] + Local Density:Sex + Local Density:Regional Density	9	1059.8	0	-520.8	0.686	1
Density interaction model (DI): ~ Local Density + Regional Density + Sex + Closest Distance + Local Density:Regional Density	8	1062.0	2.2	-523.0	0.228	2
Local and regional density model : ~ Local Density + Regional Density + Sex + Closest Distance	7	1065.3	5.5	-525.7	0.044	4
Raw density model (RWD): ~ Raw Density + Sex + Closest Distance	6	1065.6	5.7	-526.8	0.040	3
Regional density model: ~ Regional Density + Sex + Closest Distance	6	1071.4	11.6	-529.7	0.002	5
Local density model: ~ Local Density + Sex + Closest Distance	6	1074.7	14.9	-531.4	<0.001	6
Sex model: ~ Sex + Closest Distance	5	1081.0	21.2	-535.5	<0.001	7
Biological null model: ~ Closest Distance	4	1085.4	25.6	-538.7	<0.001	8

Note: K = number of parameters, AIC = Akaike Information Criterion, ΔAIC = Model's AIC -
the lowest AIC of any model, LogLik = log likelihood, AICω = Aikaike Weight.
Random effects in both models were year and trapline of emigration.
* density (captures/hundred trap-nights) of deer mice on the trapline an individual is initially
captured on minus the regional density in that year
[+] density (captures/hundred trap-nights) of deer mice averaged across all traplines in a year
‡ whether an animal is male or female, male used as reference category
¶ natural log of the distance in metres from the line of an individual's initial capture to the next
most closely situated line

Table 2: Summary of the statistical results from density and sex interaction model (DSI)

Variable – Variable Type	$\beta \pm SE$	Z	∂^2	X^2	P
Closest Distance - Fixed	-1.035 ± 0.15	-6.85	-	-	<0.0001
Sex (Male) - Fixed	0.32 ± 0.19	1.69	-	-	0.09
Local Density - Fixed	-0.091 ± 0.026	-3.52	-	-	0.0004
Regional Density - Fixed	-0.63 ± 0.16	-3.98	-	-	<0.0001
Local Density : Reg. Density - Fixed	0.036 ± 0.015	2.44	-	-	0.01
Local Density : Sex (male) - Fixed	0.064 ± 0.031	2.10	-	-	0.04
Year - Random	-	-	1042	0.2	0.65
Initial Trapline – Random	-	-	1042	11.2	0.0008

583   Note: This model best fitted the observed dispersal data and had strong support based on model comparisons using AIC.
584   Likelihood ratio tests were used to assess the significance of each random effect.

Fig 1: a) Proportion of possible dispersers that dispersed each year plotted alongside the density of the deer mouse (*Peromyscus maniculatus*) population in APP between 1960-2015. b) the effect size and significance of the relationship between raw density and dispersal probability as analyzed in 32 ten-year shifting windows using a model with raw local density as a fixed effect and *year* and *trapline* of initial capture as random effects. The magnitude of this effect of raw density on the probability of dispersing (effect size) is plotted against the midpoint of the 10-year window. Windows where the relationship is significant are represented with triangles and lack of significance is represented with squares. Significance determined at $\alpha = 0.05$.

Fig 2: Distances travelled by dispersing deer mice (N = 142). (F) is for females while (M) represents males. The number of individuals that dispersed to the closest trapline is depicted by the height of the open bars, whereas those that dispersed farther than the nearest trapline are depicted by the height of the shaded bar. There was no difference in dispersal distance between the sexes (t = 0.98, df = 92, p-value = 0.3)

Fig 3: a) Partial plot of the interaction between Regional Density and Local Density from the DSI model ($\beta \pm SE = 0.036 \pm 0.015$, Z = 2.44, P = 0.015). Regional Density is grouped by tertile. The positive interaction can be seen to make the negative density-dependent dispersal in response to local density more pronounced when regional density was low (green trend-line). When regional density was high (pink trend-line) there was no effect of local density. b) Partial plot of the interaction between Local Density and Sex (F=Female, M = Male) from the DSI model ($\beta \pm SE = 0.064 \pm 0.031$, Z = 2.10, P = 0.036). While males are more likely to disperse, this interaction is approaching significance and suggests that female dispersal may be more sensitive to changes in local density than male dispersal.

3