Running Head: Variable density-dependent dispersal

Title: Variation in space and time: a long-term examination of density-dependent dispersal in a

woodland rodent

Authors: Denomme-Brown[1], Simon T. (sdenommebrown@gmail.com); Cottenie[1], Karl

(cottenie@uoguelph.ca); Falls[2], J. Bruce (jbruce.falls@utoronto.ca); Falls[2], E. Ann

(fallseann@gmail.com); Brooks[1], Ronald J. (rjbrooks@uoguelph.ca) and Andrew G. McAdam[1]

(amcadam@uoguelph.ca)

[1] University of Guelph, Dept. of Integrative Biology, 50 Stone Rd. E., Guelph, ON, Canada,

N1G 2W1

[2] University of Toronto, Dept. of Zoology, Toronto, ON, Canada

Corresponding Author Information: Simon T. Denomme-Brown, sdenommebrown@gmail.com,

Dept. of Integrative Biology, University of Guelph, Ontario, Canada, N1G 2W1. Phone Number:

(519) 803-7993

Abstract

Dispersal is a fundamental ecological process, and population density has been observed as a

driver of dispersal in various taxa. Conflicting examples of positive and negative density-

dependent dispersal, however, leave little consensus regarding any general effects of density on

dispersal. Deer mice (*Peromyscus maniculatus*) have exhibited both negative and positive

density-dependent dispersal. Using 51-years of data on *P. maniculatus* live-trapping

abundances, we examined the spatial scale of density-dependent dispersal as well as its temporal

stability within this long time-series. We examined these patterns over both the entire time-series

and also in ten-year shifting windows to determine whether the nature and strength of the

relationship changed through time. Overall, the probability of dispersal decreased with increased

local and regional population density, and the negative effect of local density on dispersal was

more pronounced in years with low regional densities. Females were less likely to disperse, but

female dispersal was more density-dependent than male dispersal. Additionally, the strength of

negative density-dependent dispersal changed through time, from very strong in some decades to

absent in others. Our study shows that the relationship between density and dispersal is not

temporally static and that studies of density-dependent dispersal should consider both local and

regional population densities. As well, while male-biased dispersal is often considered

ubiquitous in mammals, we demonstrate that male and female dispersal may have differing

levels of density-dependence. Finally, our study highlights the importance of accounting for both

local and regional processes in natural systems as these types of long-term, spatially broad

examinations of dispersal are crucial if spatial processes are to be represented adequately and

accurately in population modelling and theory.

Keywords: Dispersal, Density-dependence, Peromyscus, Small Mammals, Spatial effects,

Temporal Dynamics, Movement

Introduction

Dispersal influences fundamental ecological processes such as population growth rates

and viability, genetic connectivity, and range expansions (Greenwood 1980; Nathan 2006;

Holyoak et al. 2008). Dispersal is identified as one of four fundamental processes in community

ecology (Vellend 2010), and a cornerstone of the metacommunity concept (Leibold et al. 2004).

Yet the causes of dispersal are often poorly understood and dispersal remains difficult to measure

and include in empirical studies (Koenig et al. 1996, Nathan et al. 2003, Logue et al. 2011).

Benefits of dispersal often stem from differences in conspecific densities between pre and

post-dispersal settlement sites. When conspecific densities at pre-dispersal sites are greater than

at post-dispersal sites (i.e. positive density-dependent dispersal), dispersing individuals may

benefit from reduced competition and agonistic interactions (Murray 1967; Gaines and

McClenaghan 1980; Matthysen 2005). Alternatively, negative density-dependent dispersal may

be beneficial when individuals disperse to gain access to a limiting resource, or when the benefits

of group living outweigh the costs of intraspecific competition (Bowler and Benton 2005).

Conflicting examples of positive and negative density-dependent dispersal (DDD) across

a variety of taxa (Bowler and Benton 2005; Matthysen 2005; Rodrigues and Johnstone 2014),

have left little consensus regarding general effects of density on dispersal. Matthysen (2005)

concluded that both birds and mammals predominantly exhibit positive DDD. The review,

however, excluded studies exhibiting temporal trends in density and seasonal variation which

Matthysen (2005) acknowledged could have changed their conclusions. Contradicting effects of

density on dispersal have also been found within species. DDD has been observed as both

negative and positive in separate studies of the same species (Garten and Smith 1974; Rehmeier

et al. 2004), or depending on when in the annual cycle density is measured (Betini et al. 2015), or

simultaneously negative and positive across a patchily distributed population (Kim et al. 2009).

In particular, variable effects of density on dispersal have been reported in deer mice,

Peromyscus maniculatus. *Peromyscus* spp. generally display male-biased dispersal (Fairbairn

1978b; Gaines and McClenaghan 1980; Rehmeier et al. 2004) and have been observed to exhibit

both positive (Garten and Smith 1974; Fairbairn 1978b; Anderson and Meikle 2010) and

negative DDD (Rehmeier et al. 2004; Wojan et al. 2017). These variable effects of density on

dispersal suggest that temporal and spatial variation in population densities could feasibly

influence the strength of density dependence in a system. Therefore, as others have proposed

(e.g. Matthysen 2005), DDD must be examined using long time-series, while accounting for

spatial and temporal variation in population densities. In this study, we examined the frequency

and extent of dispersal by wild *P. maniculatus*, using monitoring data spanning 50 years from

traplines distributed across nearly 17km of temperate forest. These data allowed us to examine

dispersal patterns at large temporal and spatial scales and investigate how the strength of DDD

was affected by both temporal and spatial variation in population densities.

Methods

Data Collection

Deer mice were live-trapped in Algonquin Provincial Park (APP), Ontario, Canada from 1960 -

2015. Trapping initially occurred on 10 traplines, with lines added and removed intermittently

throughout the study such that the same 17 lines were trapped consistently from 1995 – 2015.

Lines consisted of 10 trapping stations set along transects at 10m intervals. Early in the study,

most lines had one trap at each station, but by 1979 all lines had two traps per station in an

attempt to avoid trap saturation, for a total of 20 traps per line. Sherman-style live traps were

used exclusively until 2013, when one Longworth trap (Natural Heritage Book Society, Totnes,

Devon) replaced one Sherman at each station. Traps were baited with peanut butter and rolled

oats until 1991. After 1991, water-soaked sunflower seeds were used as bait. Freeze-killed

mealworms (*Tenebrio molitor*) were added to half the traps from 2013 - 2015 as a bait

supplement. Use of Longworth traps and mealworms were aimed at decreasing mortality in

shrew (*Soricidae*) species (Do et al. 2013; Shonfield et al. 2013). Distances between lines ranged

from 0.115 - 16.9km. Lines were located in forested habitats with different dominant trees

including sugar maple (*Acer saccharum*), white pine (*Pinus strobus*), white spruce (*Picea*

glauca), black spruce (*Picea mariana*), trembling aspen (*Populus tremuloides*), and red pine

(*Pinus resinosa*).

Each year, trapping was conducted on three consecutive nights either once or twice a

month from May through September, for a maximum of 10 three-night trapping sessions per

year. Traps were set just before dusk and checked after dawn. When captured, deer mice were

weighed, sexed, aged and assessed for reproductive condition. Before release, animals were

fitted with uniquely coded ear tags for individual identification upon recapture.

Dispersal Detection

Dispersal was defined to occur when an individual deer mouse was captured on multiple

traplines within a single trapping season. Movements occurring between years were excluded as

the biological drivers of inter-year dispersals likely differed from those occurring within a

trapping season. Additionally, few mice survive the winter in APP, and thus there were relatively

few potential dispersers over this period. Movements between lines were considered dispersal

events as the distance between the closest traplines (115m) was greater than the diameter of a

typical deer mouse home range (Harestad and Bunnel 1979). Distances between traplines were

calculated as straight-line distance between the first trap station on each line.

Although these data were collected in a consistent manner, any errors in data collection

or transfer to digital records could mistakenly appear as dispersal events (e.g. incorrect tag or

trapline numbers were recorded). While the error rate is likely low, the compounding of even a

low rate of error over more than 50 years of data could skew our estimates of dispersal. To

minimize the frequency of such errors we used a strict criterion to identify dispersal events. To

be considered a dispersing individual, an individual mouse must have been captured on multiple,

consecutive occasions on one line either preceded, or followed by at least one capture on another

line. We assumed that the mouse in question was either trapped while moving towards a trapline

where it subsequently settled, or was trapped while dispersing from a trapline and either

continued on or died. To meet this criterion, a mouse must have been captured on a minimum of

three occasions within a year. Accordingly, all model analyses and calculations of dispersal

frequencies discussed hereafter were performed on a dataset of mice caught three or more times

in a year. Additionally, we avoided including mis-read tags by ensuring that dispersing animals

had consistent records of individual traits such as sex and size. Individuals that moved to a new

line and then returned immediately to their initial line were also excluded as this likely indicated

a misread tag. Finally, changes in methodology and some loss of data necessitated the exclusion

of five years of data from the dataset (1987-1991).

Density Measures

We calculated three measures of density and converted each from raw capture totals to

captures per hundred trap-nights to account for variation in trapping effort both between years

and among traplines. The first measurement was raw local trapline density (hereafter termed raw

density), measured as the number of captures per hundred trap-nights at each trapline in each

year. The second density measure was an average regional trapline density, calculated annually

as the mean number of captures per hundred trap-nights across all traplines in APP in that year.

Finally, we calculated local density for each trapline in each year by subtracting the regional

density in a year from the raw density on each line that same year. This measure allowed us to

include local densities and regional densities within the same models.

It should be noted that the density estimates we used do not account for spatial or

temporal variation in capture probability. While spatially explicit mark-recapture techniques

(SECR) represent a powerful set of statistical tools to account for these issues (Royle et al.

2013), they were inappropriate for our long-term data for several reasons. First, Bayesian SECR

models are inappropriate for trapping data that are derived from single-catch traps such as those

used in our study (Gerber and Parmenter 2015). Second, when using frequentist SECR models

the multi-catch estimator can only be used if density is relatively constant over survey regions

(Distiller and Borchers 2015), and this was not the case in our data (see Results). Finally, the

single-catch estimator developed by Distiller and Borchers (2015), which allows for frequentist

SECR analysis using single catch traps, requires times for all captures which were not available

for our historical data. We discuss this issue further below.

Data Analysis

To assess how a deer mouse's propensity to disperse was affected by sex and population

density at different spatial scales, we fitted eight separate Generalized Linear Mixed Effects

Models (GLMM; binomial family, logit link) associated with local and regional population

densities as well as sex of the individual mouse. They were 1) Biological Null Model (BN), 2)

Sex Model (SX), 3) Local Density Model (LD), 4) Regional Density Model (RGD), 5) Raw

Density Model (RWD), 6) Local and Regional Density Model (LRD), 7) Density Interaction

Model (DI), 8) Density and Sex Interaction Model (DSI) (see Table 1). All eight models used the

binary response variable of whether an individual mouse dispersed or not and included the fixed

effect of distance (in meters) from the trapline where the mouse was initially caught to the next

closest trapline. This effect was included to account for the non-uniform spatial distribution of

traplines and whether differences in movement among lines were due primarily to spatial effects

on the probability of detecting a dispersal event (i.e. dispersal is more likely to be detected

between nearby traplines). We used the natural log of the raw distance in kilometers (km) to

correct a significant non-linearity in the relationship between this variable and the probability of

dispersal. In all models, local density and regional density were converted to z-scores ($z = x -$

$\bar{x}/\sigma$) based on the mean and standard deviation of all observations across years and lines.

We used the Akaike Information Criterion (AIC) to assess relative support for the eight

models (Burnham and Anderson 2007), which included various combinations of sex, and raw,

local and regional density as fixed effects (Table 1). We calculated AIC values for each model

and performed model comparisons using ΔAIC (Δ_i) values (Table 1). Akaike Weights (ω_i) were

calculated to examine the conditional probabilities of each model (Table 1). All models

contained the same random effects, with initial trapline of capture and year of capture included to

capture spatial and temporal variation in dispersal propensity. Likelihood ratio tests (LRT) were

used to assess the significance of the year and trapline random effects in each of the 8 candidate

models. Models were assessed for overdispersion and collinearity of variables and met model

assumptions. Statistical analyses were performed in R (Version 3.5; R Core Team 2018),

including the glmer function with a bobyqa optimizer in the lme4 package (Bates et al. 2015).

In addition to the model analysis described above, we performed a number of *post hoc*

analyses. When first visualizing the relationship between regional density and movement

frequency (Figure 1a) there appeared to be a marked change in the relationship between the two

variables for portions of the time-series, particularly over a 9-year period from 2002-2010. To

examine this, we ran the RWD model on a shifting ten-year window through the time-series. The

RWD model was used because it was the simplest model with moderate support based on AIC

comparisons. More complex models would not fit these data given the smaller sample size

provided in 10-year windows of the time-series (Figure 1b).

The second set of *post hoc* analyses assessed how robust our results were to changes in

trapping protocols during this long-term study (e.g. number of traps per station, change in bait,

and trap types), that occurred at distinct points in our study (1979, 1991 and 2013 respectively).

For this we ran our best model on subsets of our dataset from before, between and after these

protocol changes occurred.

Results

Population Densities and Frequency of Movement

Over the 51 years included in this study, there were 3408 instances where a mouse was

caught at least three times in a year. Of these 3408 possible dispersers, 4.5% (n = 154) dispersed

between traplines. The proportion of the population that dispersed fluctuated between years ($\bar{x} \pm$

SE = 0.053 ± 0.008 mice dispersing in the population, n = 51 years), with no dispersal detected

in some years and a maximum of 28% of the population dispersing between lines in 1964 (Figure

1). The population sex ratio was slightly skewed, with males comprising 53.4% of individuals in

the dataset ($X^2 = 7.9$, df = 1, P < 0.01). This male skew was more pronounced among dispersing

individuals, with 65.6% of dispersing mice being male ($X^2 = 7.7$, df = 1, P < 0.01). Overall,

males were more likely to disperse (5.5%) than females (3.3%; Fisher Exact Test: P = 0.002).

Distances travelled by dispersing individuals ranged from 115m to 11.4 km ($\bar{x} \pm$ SE =

1.57 ± 0.18 km, n = 154) (Figure 2). There was no difference in dispersal distance between the

sexes (t = 1.1, df = 95.7, p-value = 0.28) (Figure 2). Only 52.6% (n = 81) of dispersals were to

the next most closely situated trapline (Figure 2). There were two traplines located only 115m

apart, but dispersal between these traplines represented only 20.1% (n = 31) of all movements.

The density of deer mice in APP fluctuated over an order of magnitude during this study,

with yearly regional population densities ranging from 1.9 to 27.8 captures per hundred trap

nights ($\bar{x} \pm$ SE = 11.15 ± 0.81 captures per hundred trap-nights, n = 51 years) (Figure 1a). Local

densities also varied, with the average range in local population densities among traplines within

years ($\bar{x} \pm$ SE = 19.37 ± 1.22 captures per hundred trap-nights, n = 51 years) approaching the

aforementioned range in regional densities among years. In particular, 9 of 51 years exhibited a

greater range in density among lines within a year than the range in regional density across the

51 years, with a maximum range of 47.2 captures per hundred trap-nights in 2001.

Dispersal Models

The model that had the lowest AIC score of the eight fitted models was the DSI model

which included the fixed effects of both regional and local density, sex, and interactions between

both local density and regional density, and local density and sex (Table 2). In the DSI model,

the probability of dispersal decreased with increasing distance to nearest trapline ($\beta \pm$ SE = -1.16

± 0.17, Z = -7.0, P < 0.0001), increased regional density ($\beta \pm$ SE = -0.57 ± 0.15, Z = -3.9, P <

0.0001), and increased local density ($\beta \pm$ SE = -0.087 ± 0.025, Z -3.45, P = 0.0006). The

significant positive interaction between local and regional densities ($\beta \pm$ SE = 0.031 ± 0.014, Z =

2.23, P = 0.026) indicated that in years of low regional density, local densities had a stronger

negative effect on the probability of dispersal (Figure 3a). For example, three of the five years

displaying the highest proportions of dispersers in the study were 1964 (0.286), 1962 (0.192) and

1976 (0.150), which represented three of the 11 lowest density years within the study.

The DSI model also included sex as a significant predictor of dispersal, with males

dispersing more frequently than females ($\beta \pm$ SE = 0.38 ± 0.19, Z = 2.04, P = 0.04) (Figure 3b).

However, the interaction between local density and sex was marginally significant ($\beta \pm$ SE =

0.057 ± 0.029, Z = 2.0, P = 0.051), suggesting that while males disperse more than females,

females were more sensitive to local density than males and were more likely to disperse when

densities were low (Figure 3b). Year was a significant random effect in this model (LRT,

Deviance = 1098.3, X^2_1 = 9.8, P = 0.0017) whereas initial line of capture was not (LRT, Deviance

= 1106.7, X^2_1 = 1.5, P = 0.23). Overall, the DSI model explained 12.56% of the deviance.

There was also strong support for the density interaction model (DI) (Δ_i < 2), which was

identical to the DSI model except it did not include an interaction between local density and sex.

All effects in this model were in the same direction as in the DSI model and the DI model

explained 12.26% of the overall deviance. Considerably less support was seen for models that

did not contain the interaction between local and regional densities, with all these models

demonstrating either moderate (Δ_i < 7) or no support (Δi > 10) (Table 1).

Despite evidence of negative DDD overall, visual inspection of the time series of density

and dispersal probability (Figure 1a) suggested temporal changes in the strength of density-

dependent dispersal. When running the RWD model on the shifting 10-year window through the

time series, the effect of raw density on dispersal probability never became positive. The effect

size did approach zero, however, and the negative relationship between raw density and dispersal

probability was significant for only 14 of 32 ten-year windows in the analysis (Figure 1b).

Finally, when running the DSI model on our data before and after each change in trapping

methodology, there were no changes in the direction of effects for all explanatory variables save

one. The exception was the effect of sex on dispersal, which was reversed prior to the initial bait

change, but the effect was not significant.

Discussion

Probability of Dispersal

Overall, deer mice in APP exhibited negative DDD, with a lower probability of dispersal

under high-density conditions, regardless of the scale at which density was measured. Multiple

explanations have been proposed to explain negative DDD. Attraction to conspecifics

(Danielson and Gaines 1987; Stamps 1991), range constriction in the face of high densities

(Stickel 1960; Taitt 1981) and exclusionary behavior towards immigrants in periods of high

density (i.e. "social fences"; Hestbeck 1982) have all been proposed as possible mechanisms. If

trap saturation had occurred within our system it could mimic a natural social fence, denying

immigrants access to traps in densely populated lines and biasing our results towards the

appearance of negative DDD. This seems unlikely, however, as densities on even the most

highly populated traplines fell well short of saturation of available traps (i.e. average captures per

100 trap-nights in APP across all lines throughout this study was 22.62 with SE = 0.42 and a

standard deviation of 10.66, while the highest capture rate on any line in any year of the study

was 73.81 captures per 100 trap-nights). Additionally, there was no obvious change in the

direction of density-dependence before and after the traps per station were doubled in 1979.

While the result of negative DDD runs counter to generalizations of positive DDD in

mammals (Matthysen 2005), previously reported relationships between dispersal and density in

Peromyscus spp. have been inconsistent. Positive DDD has been observed in *P. maniculatus*

(Fairbairn, 1978a), *P. polionotus* (Garten and Smith 1974), and genetic estimates show that *P.*

leucopus is more likely to disperse from high density habitat patches (Anderson and Meikle

2010). In contrast, evidence of negative DDD has been found in both *P. maniculatus* (Rehmeier

et al. 2004) and *P. boylii* (Wojan et al. 2015). The spatial and temporal breadth of our study may

elucidate why studies on *Peromyscus* spp., and other dispersing organisms, often yield divergent

results. For while deer mice in our study consistently exhibited negative DDD, the strength with

which density affected dispersal varied temporally, spatially and between males and females.

The relationship between density and dispersal varied throughout our time-series. This

variation is evident upon examining the time-series of regional density and movement frequency

on the same axis (Figure 1a), particularly in the 9-year period from 2002-2010 which visually

suggests a possible positive relationship. The effect of raw density on dispersal probability in the

RWD model was never positive in any 10-year window along the time-series, however, the

effect size approached zero in the 1990's before becoming increasingly negative (Figure 1b).

Also, the majority of the 10-year windows examined (n = 18) did not exhibit a significant

relationship between density and dispersal. Few studies investigating DDD have been performed

using long-term data (Matthysen 2005), despite the fact that inferences drawn from short-term

studies can differ drastically from those generated by long-term research (Kratz et al. 2003), and

that effects of some ecological processes become more pronounced over time (Cardinale et al.

2007). Our data highlight the importance of long-term time-series to drawing accurate

conclusions regarding the density-dispersal relationship, as an unfortunately timed relatively

long-term study (i.e. 10 years) would not detect the negative effect of density on dispersal that

exists in this system.

In addition to varying temporally, the strength with which local density affected dispersal

was also mediated by regional densities. This positive interaction between the regional and local

density effects on dispersal indicates that the negative effect of local density on dispersal is

greater in years of low regional density and muted in years of high regional density (Figure 3b).

This suggests that local population density can be an important driver of dispersal in this system,

but only when examined in concert with regional density. Low local densities that increase the

probability of dispersal are only noteworthy in years when population densities across the APP

system are also low. This relationship between local and regional densities as interacting drivers

of dispersal has not been shown previously to our knowledge. It is important to highlight that

when examining dispersal, local population densities must be considered within the context of

their accompanying regional densities, as many existing studies provide only densities for

localized sites of emigration (Matthysen 2005), which based on our findings may lead to

spurious results if context is not provided in the form of associated regional densities. As well,

the few large datasets that have been used to examine DDD fail to differentiate between spatial

and temporal variation in densities (Matthysen, 2005, but see Pasinelli and Walters 2002;

Catchpole et al. 2004) despite the fact both temporal variation and interactions between densities

at multiple spatial scales affect dispersal.

Several potential mechanisms could cause this interaction between local and regional

densities and their effect on dispersal. Deer mice populations in APP are closely tied to maple

seed crops, with mouse populations peaking after seed masting events (Falls et al. 2007).

Peromyscus spp. will also move based on both habitat quality and population density (Morris

and Diffendorfer 2004). It may be that individuals select habitat and disperse to high quality,

maple-rich areas when possible; but in years of high regional densities, local densities in high-

quality habitats become high enough to prevent such immigration. Such a process would strike a

balance between traditional dispersal hypotheses predicting positive DDD in the face of

increased competition (Greenwood 1980), and ideas of conspecific attraction (Stamps 1991) and

habitat selection. This type of balance between competition and facilitation amongst conspecifics

has been observed previously in a colony of blue-footed booby (*Sula nebouxii*), where

simultaneous positive and negative DDD was thought to result from the costs of competition and

the advantages of increased mating choice options (Kim et al. 2009).

It may be that mate availability plays a role in dispersal in the APP system because the

strength of density-dependence also differed between males and females. Males dispersed more

often than females, which is consistent with findings of male-biased dispersal in both

Peromyscus spp. (Fairbairn 1978a; Goundie and Vessey 1986; Rehmeier et al. 2004; Mabry

2014), and other mammals (Greenwood 1980; Dobson 1982; Mabry et al. 2013). However, we

also found evidence that dispersal in females was more sensitive to changes in local density than

in males, such that females dispersed more frequently than males when local densities were very

low. Examples of sex-biased dispersal switching from male-bias to female-bias within a system

are rare (but see Pérez-González and Carranza 2009), but could have important implications for

understanding the role of mating strategies in dispersal. The predominance of male-biased

dispersal in mammals (and female-biased dispersal in birds) has long been attributed to mating

systems, although the simplicity of this relationship has been questioned (Mabry et al. 2013). If

males disperse to track females and females rely on available resources for reproduction (Emlen

and Oring 1977), then at exceedingly low densities the lack of available mates may outweigh the

benefits of familiar surroundings, resulting in a spike in female dispersal. Alternatively, if

densities are associated with food availability, then low densities may indicate low food

availability, which would affect dispersal in females more strongly than in males due to their

reliance on resources for reproduction (Emlen and Oring 1977). Such differences between male

and female dispersal sensitivity to local population density mean that spatial variation in density

could potentially affect sex ratios, which could in turn affect population density.

As mentioned previously, our density measures did not account for variation in capture

probabilities. While we have described the practical reasons for not being able to utilize SECR

methods, we recognize that this is not an ideal. However, the lack of trap saturation on any of our

lines throughout the study, the fact that frequency of recapture of individuals within this study ($\bar{x}$

$\pm$ SE = 2.63 $\pm$ 0.07) did not decrease with increased regional annual densities (linear regression;

$\beta \pm$ SE = 0.012 $\pm$ 0.01, t = 1.09, P = 0.28), and the fact that the density-dispersal relationships in

our models were not altered by the changes in trapping methodologies that occurred throughout

this study all speak to the robustness of our findings.

Frequency and Distance of Dispersal

In general, deer mice in APP dispersed as often as expected based on existing literature,

but over much greater distances than anticipated. Rehmeier et al. (2004) reported that over a 10-

year period 4% of deer mice in a tallgrass prairie undertook long-distance dispersals (LDDs,

>125m); Bowman et al. (2001b) similarly reported 4.2% of deer mice demonstrating LDD

(>100m) in mixed-wood forest. Based on these papers, our dispersals (>115m) can be classified

as LDDs and the yearly mean frequency of LDD by deer mice in APP of 4.5% was remarkably

similar to these other frequencies reported in the literature. However, these absolute frequencies

of dispersal should be interpreted with some caution as mark-recapture studies have limited

ability to detect dispersal. The majority of studies monitoring small mammal populations utilize

trapping grids of insufficient size to detect LDDs (< 2ha, from Bowman et al. 2001a).

Comparisons with radio tracking data and genetic estimates of dispersal have shown that such

trapping practices combined with mark-recapture methods underestimate dispersal distances

(Koenig et al. 1996). Unfortunately, expanding trapping grids to detect greater dispersal

distances is often impractical, as increasing the maximum straight-line distance across a grid

requires an exponential increase in area and monitoring effort. Our study provides an alternative

approach by instead monitoring dispersal using many smaller trapping transects. This reduced

our ability to detect individual dispersal events, however, this was overcome through extensive

temporal replication. So, while our raw rates of detected dispersals are surely downwardly

biased, our measurement of dispersal across 3408 individuals, 17 traplines and 51 years allowed

us to assess the effects of large-scale processes on the probability and extent of dispersal.

The spatial extent of dispersal within our system exceeded previous reports for this

species. Of 154 dispersal events we observed, 13% (n =20) exceeded the previous maximum

long-distance movement reported for *P. maniculatus* (4287 ± 10m, in Wood and McKinney

2015). The maximum dispersal distance of 11.3km in this study represents a 164% increase on

that previous reported maximum. As well, we detected more frequent movements of great

distances than previous studies. For instance, 3.3% of potentially dispersing mice in APP

dispersed over 200m compared to 1% of mice dispersing beyond this same distance in previous

studies (Brant 1962; Groves and Keller 1986). We also observed 2.8% and 2.1% of mice

dispersing beyond 300m and 500m respectively, while roughly 1% of populations travelled these

distances in previous studies (Diffendorfer et al. 1999; Rehmeier et al. 2004). This indicates that

the tail of the dispersal distance distribution of this species has previously been underestimated.

Dispersal distances tend to follow a leptokurtic distribution, with many individuals

dispersing short distances and few individuals exhibiting LDD. Understanding the rate and

extent of LDDs is particularly important as the spread of a species in a landscape is determined

largely by the frequency and extent of LDD, as represented in the tail of the species' distribution

of dispersal distances (Kot et al. 1996; Levin et al. 2003; Green and Figuerola 2005). However,

tails of dispersal distance distributions are notoriously difficult to quantify (Koenig et al. 1996),

which has led to LDD in deer mice being generally reported with little detail or as anecdotal

observations (Jung et al. 2005; Wood and McKinney 2015). Our results suggest that LDD by

deer mice is more common than reports in the literature would suggest, reinforcing claims that

relatively small scale mark-recapture techniques used to monitor many species fail to accurately

capture LDD (Koenig et al. 1996). Our results, however, indicate mark-recapture studies can

detect dispersal between distant locations, provided the temporal extent of the study is sufficient.

Our evidence of greater frequencies of LDD in this species has important ecological

implications as species with "fat-tailed" dispersal distributions are thought to exhibit more rapid

spread throughout landscapes (Kot et al. 1996). Dispersal distance distributions can influence

persistence in the face of climate change (Schloss et al. 2012) and the spread of zoonotic diseases

(Mundt et al. 2009), such as Hantavirus and Lyme disease for which *Peromyscus* spp. are vectors

(Meerburg et al. 2009; Gaitan and Millien 2016). This is particularly worthy of attention as small

mammal populations in this region are already responding to climate change (Myers et al. 2009).

We have demonstrated that deer mice exhibit negative DDD that is more pronounced

during periods of low regional densities and that female dispersal is more sensitive to these

changes in density than that of males. We also have demonstrated that the relationship between

density and dispersal is not static but rather varies conspicuously over large time scales. These

results show that long-term field studies are beneficial to the examination of ecological processes

such dispersal, as smaller scale projects are likely to miss or potentially misrepresent the broader

signal entirely. As well, our findings suggest that while local population densities are important,

their influence on dispersal can be affected by broader regional population densities. As such,

future studies should account for both local and regional densities when testing for DDD.

Additionally, we have shown that well-established phenomena such as mammalian male-biased

dispersal may also be density-dependent. These results together suggest that the conflicting

relationships between density and dispersal in the literature may result from broader context-

dependence in DDD. Finally, our results add to a previously modest body of empirical literature

examining LDD. We have demonstrated that deer mice dispersal of greater than 100m is not

uncommon and that LDD occurs more frequently, and to greater extents, than previously

thought. This reinforces that dispersal distributions estimated from short-term mark recapture

studies may be drastically underestimated, which is potentially important for future conservation.

Acknowledgements

We would like to acknowledge countless individuals who collected these data. Thank you to the

Algonquin Wildlife Research Station for logistical support. This work was supported through

funding from the National Science and Engineering Research Council of Canada and the Ontario

Ministry of Natural Resources. We dedicate this work to the memory of our colleague Dr. Leslie

Rye who said it best: "I can't believe I am getting paid to work in Algonquin!"

References

Anderson, C. S., and D. B. Meikle. 2010. Genetic estimates of immigration and emigration rates in relation to population density and forest patch area in Peromyscus leucopus. Conservation Genetics 11:1593–1605.

Bates, D., M. Maechler, B. Bolker, and S. Walker. 2015. Fitting Linear Mixed-Effects Models Using lme4. Journal of Statistical Software 67:1–48.

Betini, G. S., A. Pardy, C. K. Griswold, and D. R. Norris. 2015. The role of seasonality and non-lethal carry-over effects on density-dependent dispersal. Ecosphere 6:art272.

Bowler, D. E., and T. G. Benton. 2005. Causes and consequences of animal dispersal strategies: relating individual behaviour to spatial dynamics. Biological reviews of the Cambridge Philosophical Society 80:205–225.

Bowman, J., C. V. Corkum, and G. J. Forbes. 2001a. Spatial Scales of Trapping in Small-mammal Research. The Canadian Field-Naturalist 115:472–475.

Bowman, J., G. Forbes, and T. Dilworth. 2001b. Distances moved by small woodland rodents within large trapping grids. Canadian Field-Naturalist 115:64-67.

Burnham, K.P. and D.R. Anderson. 2007. Model Selection and Inference: A Practical Information-Theoretic Approach (2nd edition) New York, NY: Springer.

Brant, D. H. 1962. Measures of the Movements and Population Densities of Small Rodents. University of California Publications in Zoology 62:105–184.

Cardinale, B. J., J. P. Wright, M. W. Cadotte, I. T. Carroll, A. Hector, D. S. Srivastava, M. Loreau, and J. J. Weis. 2007. Impacts of plant diversity on biomass production increase through time because of species complementarity. Proceedings of the National Academy of

Sciences 104:18123–18128.

Catchpole, E. A., Y. Fan, B. J. T. Morgan, T. N. Coulson, and T. H. Clutton-Brock. 2004. Sexual

dimorphism, survival and dispersal in red deer. Journal of Agricultural, Biological and

Environmental Statistics 9:1–26.

Danielson, B. J., and M. S. Gaines. 1987. The influences of Conspecific and Heterospecific

Residents on Colonization. Ecology 68:1778–1784.

Diffendorfer, J. E., M. S. Gaines, and R. D. Holt. 1999. Patterns and impacts of movements at

different scales in small mammals. Pages 63–88 in G. W. Barrett and J. D. Peles, editors.

Landscape ecology of small mammals. Springer-Verlag, New York.

Distiller, G., and D. L. Borchers. 2015. A spatially explicit capture-recapture estimator for

single-catch traps. Ecology and Evolution 21:5074-5087.

Do, R., J. Shonfield, and A.G. McAdam. 2013. Reducing accidental shrew mortality associated

with small-mammal livetrapping II: a field experiment with bait supplementation. Journal of

Mammalogy 94:754-760.

Dobson. 1982. Competition for Mates and Predominant Juvenile Male Dispersal in Mammals.

Animal Behaviour 30:1183–1192.

Emlen, S. T., and L. W. Oring. 1977. Ecology, Sexual Selection and the Evolution of Mating

Systems. Science 197:215–223.

Fairbairn, D. J. 1978a. Dispersal of Deer Mice. Oecologia 193:171–193.

Fairbairn, D. J. 1978b. Behaviour of dispersing deer mice (Peromyscus maniculatus). Behavioral

Ecology and Sociobiology 3:265–282.

Falls, J. B., E. A. Falls, and J. M. Fryxell. 2007. Fluctuations of Deer Mice in Ontario in Relation

to Seed Crops. Ecological Monographs 77:19–32.

Gaines, M. S., and L. R. McClenaghan. 1980. Dispersal in Small Mammals. Annual Review of

Ecology and Systematics 11:163–196.

Gaitan, J., and V. Millien. 2016. Stress level, parasite load, and movement pattern in a small-

mammal reservoir host for Lyme disease. Canadian Journal of Zoology 94:565–573.

Garten, C. T., and M. H. Smith. 1974. Movement By Oldfield Mice and Population Regulation.

Acta Theriologica 19:513–514.

Gerber, B.D., and R.R. Parmenter. 2015. Spatial capture-recapture model performance with

known small-mammal densities. Ecological Applications 25:695-705.

Goundie, T. R., and S. H. Vessey. 1986. Survival and Dispersal of Young White-Footed Mice

Born in Nest Boxes. Journal of Mammalogy 67:53–60.

Green, A. J., and J. Figuerola. 2005. Recent advances in the study of long-distance dispersal of

aquatic invertebrates via birds. Diversity and Distributions 11:149–156.

Greenwood, P. J. 1980. Mating systems, philopatry and dispersal in birds and mammals. Animal

Behaviour 28:1140–1162.

Groves, C. R., and B. L. Keller. 1986. Ecological characteristics of small mammals on a

radioactive waste disposal area in southeastern Idaho. American Midland Naturalist

46:404–410.

Harestad, A. S., and F. L. Bunnel. 1979. Home Range and Body weight -- A Reevaluation.

Ecology 60:389–402.

Hestbeck, J. B. 1982. Population Regulation of Cyclic mammals: The Social Fence Hypothesis.

Oikos 39:157–163.

Holyoak, M., R. Casagrandi, R. Nathan, E. Revilla, and O. Spiegel. 2008. Trends and missing

parts in the study of movement ecology. Proceedings of the National Academy of Sciences

of the United States of America 105:19060–19065.

Jung, T., K. O'Donovan, and T. Powell. 2005. Long-distance movement of a dispersing Deer

Mouse, Peromyscus maniculatus, in the boreal forest. Canadian Field-Naturalist 119:451–

452.

Kim, S. Y., R. Torres, and H. Drummond. 2009. Simultaneous positive and negative density-

dependent dispersal in a colonial bird species. Ecology 90:230–239.

Koenig, W., D. Van Vuren, and P. N. Hooge. 1996. Detectability, philopatry and the distribution

of dispersal distances in vertebrates. Trends in Ecology & Evolution 11:514–517.

Kot, M., M. A. Lewis, and P. Van Den Driessche. 1996. Dispersal Data and the Spread of

Invading Organisms. Ecology 77:2027–2042.

Kratz, T. K., L. A. Deegan, M. E. Harmon, and W. K. Lauenroth. 2003. Ecological Variability in

Space and Time: Insights Gained from the US LTER Program. BioScience 53:57–67.

Leibold, M. a., M. Holyoak, N. Mouquet, P. Amarasekare, J. M. Chase, M. F. Hoopes, R. D.

Holt, J. B. Shurin, R. Law, D. Tilman, M. Loreau, and a. Gonzalez. 2004. The

metacommunity concept: a framework for multi-scale community ecology. Ecology Letters

7:601–613.

Levin, S. A., H. C. Muller-Landau, R. Nathan, and J. Chave. 2003. The ecology and evolution of

seed dispersal: a theoretical perspective. Annual review of Ecology, Evolution, and

Systematics 34:575–604.

Logue, J. B., N. Mouquet, H. Peter, and H. Hillebrand. 2011. Empirical approaches to

metacommunities: a review and comparison with theory. Trends in ecology & evolution

26:482–91.

Mabry, K. E. 2014. Effects of sex and population density on dispersal and spatial genetic

structure in brush mice. Journal of Mammalogy 95:981–991.

Mabry, K. E., E. L. Shelley, K. E. Davis, D. T. Blumstein, and D. H. van Vuren. 2013. Social

Mating System and Sex-Biased Dispersal in Mammals and Birds: A Phylogenetic Analysis.

PLoS ONE 8:1–9.

Matthysen, E. 2005. Density-dependent dispersal in birds and mammals. Ecography 28:403–416.

Meerburg, B. G., G. R. Singleton, and A. Kijlstra. 2009. Rodent-borne diseases and their risks

for public health. Page Critical Reviews in Microbiology.

Morris, D. W., and J. E. Diffendorfer. 2004. Reciprocating dispersal by habitat-selecting white-

footed mice. Oikos 107:549–558.

Mundt, C. C., K. E. Sackett, L. D. Wallace, C. Cowger, and J. P. Dudley. 2009. Long-distance

dispersal and accelerating waves of disease: empirical relationships. The American

Naturalist 173:456–466.

Murray, B. G. J. 1967. Dispersal in Vertebrates. Ecology 48:975–978.

Myers, P., B. L. Lundrigan, S. M. G. Hoffman, A. P. Haraminac, and S.H. Seto. 2009. Climate-

induced changes in small mammal communities of the Northern Great Lakes Region.

Global Change Biology 15:1434-1445.

Nathan, R. 2006. Long-Distance Dispersal of Plants. Science 313:786–788.

Nathan, R., G. Perry, J. T. Cronin, A. E. Strand, and M. L. Cain. 2003. Methods for estimating

long-distance dispersal. Oikos 103:261–273.

Pasinelli, G., and J. R. Walters. 2002. Social and environmental factors affect natal dispersal and

philopatry of male Red-cockaded Woodpeckers. Ecology 83:2229–2239.

Pérez-González, J., and J. Carranza. 2009. Female-biased dispersal under conditions of low male

mating competition in a polygynous mammal. Molecular Ecology 18:4617–4630.

Rehmeier, R. L., G. A. Kaufman, and D. W. Kaufman. 2004. Long-Distance Movements of the

Deer Mouse in Tallgrass Prairie. Journal of Mammalogy 85:562–568.

Rodrigues, A. M. M., and R. A. Johnstone. 2014. Evolution of positive and negative density-

dependent dispersal. Proceedings of the Royal Society B: Biological Sciences

281:20141226–20141226.

Royle, J. A., R. B. Chandler, R. Sollmann, and B. Gardner. 2013. Spatial capture-recapture.

Academic Press.

Schloss, C. A., T. A. Nuñez, and J. J. Lawler. 2012. Dispersal will limit ability of mammals to

track climate change in the Western Hemisphere. Proceedings of the National Academy of

Sciences of the United States of America 109:8606–8611.

Shonfield, J., R. Do, R. J. Brooks, and A.G. McAdam. 2013. Reduc King accidental shrew

mortality associated with small-mammal livetrapping I: an inter- and intrastudy analysis.

Journal of Mammalogy 94: 745-753.

Stamps, J. A. 1991. The effects of conspecifics on habitat selection in teritorial species.

Behavioral Ecology and Sociobiology 28:29–36.

Stickel, L. F. 1960. Perommyscus Ranges at High and Low Population Densities. Journal of

Mammalogy 41:433–441.

Taitt, M. J. 1981. The Effect of Extra Food on Small Rodent Populations: I. Deermice

(Peromyscus maniculatus). Journal of Animal Ecology 50:111–124.

Vellend, M. 2010. Conceptual synthesis in community ecology. The Quarterly review of biology

85:183–206.

Wojan, C.M., Knapp, S.M., and K.E. Mabry. 2015. Spatial variation in population density affects

dispersal behavior in brush mice. Ecology 96:1661-1669.

Wood, C. M., and S. T. McKinney. 2015. Record Long-distance Movement of a Deer Mouse,

Peromyscus maniculatus, in a New England Montane Boreal Forest. Canadian Field

Naturalist 129:10–11.

Table 1: Eight generalized linear mixed effects models (binomial response, logit link) fitted to
the data for predicting the probability of an individual deer mouse dispersing

Model	K	AIC	ΔAIC	LogLik	AIC ω	Rank
Density and sex interaction model (DSI): ~ Local Density + Regional Density[†] + Sex[‡] + Closest Distance[¶] + Local Density:Sex + Local Density:Regional Density	9	1115.16	0	-548.57	0.582	1
Density interaction model (DI): ~ Local Density + Regional Density + Sex + Closest Distance + Local Density:Regional Density	8	1116.92	1.764	-550.46	0.241	2
Raw density model (RWD): ~ Raw Density + Sex + Closest Distance	6	1118.64	3.482	-553.32	0.102	3
Local and regional density model (LRD) ~ Local Density + Regional Density + Sex + Closest Distance	7	1119.39	4.228	-552.69	0.070	4
Regional density model (RGD): ~ Regional Density + Sex + Closest Distance	6	1125.34	10.182	-556.67	0.004	5
Local density model (LD): ~ Local Density + Sex + Closest Distance	6	1128.20	13.044	-558.10	<0.001	6
Sex model (SX): ~ Sex + Closest Distance	5	1134.55	19.397	-562.28	<0.001	7
Biological null model (BN): ~ Closest Distance	4	1140.68	25.518	-566.34	<0.001	8

K = number of parameters, AIC = Akaike Information Criterion, ΔAIC = Model's AIC - the
lowest AIC of any model, LogLik = log likelihood, AICω = Aikaike Weight.
Random effects in both models were year and trapline of emigration.
* density (captures/hundred trap-nights) of deer mice on the trapline an individual is initially
captured on minus the regional density in that year
[†] density (captures/hundred trap-nights) of deer mice averaged across all traplines in a year
[‡] whether an animal is male or female, male used as reference category
[¶] natural log of the distance in metres from the line of an individual's initial capture to the next
most closely situated line

Table 2: Summary of the statistical results from density and sex interaction model (DSI)

Fixed	$\beta \pm SE$	Z	P
Closest Distance	-1.16 ± 0.17	-7.009	<0.0001
Sex (Male)	0.38 ± 0.19	2.037	0.04
Local Density	-0.087 ± 0.025	-3.449	0.0006
Regional Density	-0.57 ± 0.15	-3.858	0.0001
Local Density : Reg. Density	0.031 ± 0.014	2.232	0.03
Local Density: Sex (male)	0.057 ± 0.029	1.195	0.05

Random	∂^2	X^2_1	P
Year	0.316	9.818	0.002
Initial Trapline	0.0638	1.455	0.2

575
576   This model best fitted the observed dispersal data and had strong support based on model
577   comparisons using AIC.
578   Significant effects (P < 0.05) are in boldface
579   Likelihood ratio tests were used to assess the significance of each random effect.
580

Figure Legends
Figure 1: Proportion of possible dispersers that in fact dispersed each year plotted alongside the
density of the deer mouse (*Peromyscus maniculatus*) population in Algonquin Provincial Park
Ontario Canada between 1960-2015 (A). For much of the time series density and the proportion
of mice that dispersed were negatively associated. This is reflected in a negative effect of raw
density on the probability of dispersing (effect size in panel B), which was measured separately
for each of 32 ten-year moving windows. The magnitude of this effect of raw density on the
probability of dispersing (effect size) is plotted against the midpoint of the 10-year window (B).
Significant effects of raw density on the probability of dispersing are shown as triangles, whereas
non-significant effects are shown as circles. Significance determined at $\alpha = 0.05$.
591
Figure 2: Total number of deer mice that were observed to disperse (N = 154 mice) various
distances. The top panel (F) is for females while the bottom panel (M) is for males. The number
of individuals that dispersed to the closest trapline is depicted by the height of the open bars,
whereas those that dispersed farther than the nearest trapline are depicted by the height of the
shaded bar. There was no difference in dispersal distance between the sexes (t = 1.1, df = 95.73,
p-value = 0.28).
598
Figure 3:(A) Partial plot of the interaction between Local Density and Sex (F=Female, M =
Male) from the DSI model ($\beta \pm SE = 0.057 \pm 0.029$, Z = 1.951, P = 0.051). While males are
more likely to disperse, this interaction is approaching significance and suggests that female
dispersal may be more sensitive to changes in local density than male dispersal. (B) Partial plot
of the interaction between Regional Density and Local Density from the DSI model ($\beta \pm SE =$
0.031 ± 0.014, Z = 2.23, P = 0.026). Regional Density is grouped by tertile. The positive
interaction can be seen to make the negative density-dependent dispersal in response to local
density more pronounced when regional density was low (green trend-line). When regional
density was high (pink trend-line) there was no effect of local density.

610
611   Figure 1

612
613   Figure 2

615   Figure 3

616