The role of "spillover" in antibiotic resistance

Short title: Antibiotic resistance "spillover"

Scott W. Olesen[1], Marc Lipsitch[1,2], Yonatan H. Grad[1,3]

1 Department of Immunology and Infectious Diseases, Harvard T. H. Chan School of Public Health, Boston, Massachusetts 02115

2 Center for Communicable Disease Dynamics, Harvard T. H. Chan School of Public Health, Boston, Massachusetts 02115

3 Division of Infectious Diseases, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts 02115

Correspondence:
Yonatan Grad
Harvard T. H. Chan School of Public Health
665 Huntington Ave, Building 1, Room 715, Boston, Massachusetts 02115
617 432 2275
ygrad@hsph.harvard.edu

ABSTRACT (240 words)

Antibiotic use is a key driver of antibiotic resistance. Understanding the quantitative

association between antibiotic use and resulting resistance is important for predicting

future rates of antibiotic resistance and for designing antibiotic stewardship policy.

However, the use-resistance association is complicated by "spillover", in which one

population's level of antibiotic use affects another population's level of resistance via the

transmission of bacteria between those populations. Spillover is known to have effects

at the level of families and hospitals, but it is unclear if spillover is relevant at larger

scales. We used mathematical modeling and analysis of observational data to address

this question. First, we used dynamical models of antibiotic resistance to predict the

effects of spillover. Whereas populations completely isolated from one another do not

experience any spillover, we found that if even 1% of interactions are between

populations, then spillover may have large consequences: the effect of a change in

antibiotic use in one population on antibiotic resistance in that population could be

reduced by as much as 50%. Then, we quantified spillover in observational antibiotic

use and resistance data from US states and European countries for 3 pathogen-

antibiotic combinations, finding that increased interactions between populations were

associated with smaller differences in antibiotic resistance between those populations.

Thus, spillover may have an important impact at the level of states and countries, which

has ramifications for predicting the future of antibiotic resistance, designing antibiotic

resistance stewardship policy, and interpreting stewardship interventions.

INTRODUCTION

Antibiotic resistance is a major threat to public health (1). Outpatient antibiotic use, which accounts for approximately 80% of human antibiotic use (2,3), is considered a principal driver of antibiotic resistance in the community (4). Understanding the relationship between use and resistance is important because it allows accurate predictions of the future of antibiotic resistance and goal-oriented antibiotic stewardship policy. The use-resistance association has been previously characterized in many ecological studies at the level of US states (5–7) and European countries (8,9). However, antibiotic resistance is a complex, temporally dynamic phenomenon (10–13), and many factors complicate the use-resistance association, making what should be an "obvious" connection sometimes difficult to identify and quantify (11). Even when detected, observed use-resistance associations are sometimes weaker than might be expected (14). One factor that could account for the difficulty in detecting use-resistance associations in ecological studies and for the apparent weakness of such associations is "spillover".

"Spillover" is a consequence of the fact that antibiotic-resistant and -susceptible bacteria can be transmitted from person to person. Thus, one person's risk of an antibiotic resistant infection depends on their own antibiotic use (15,16) as well as the rates of antibiotic use among their contacts (17). For example, one person's use of antibiotics increases the risk of an antibiotic resistant infection among their family members (18–21). As another example, hospitalized patients with no recent antibiotic use can have a higher risk of resistance than people in the community with high antibiotic use (22)

because antibiotic use and resistance in other hospitalized patients are high.

Spillover is important for three reasons. First, it means antibiotic resistance is not merely

a localized problem. It is well-understood that new resistance determinants can emerge

in one geography and spread globally (23,24), but the role of spillover in determining the

levels of resistance in a given locale is not well-quantified. To what degree, for example,

can one US state expect that its antibiotic resistance levels are due to antibiotic use

within its borders, rather than in surrounding states? Second, spillover makes it difficult

to design antibiotic stewardship interventions and understand their results. For example,

if antibiotic use in one hospital changes, resistance might not change as expected

because of spillover, from the community or other hospitals into that hospital's patients.

Finally, spillover makes it difficult to interpret the results of controlled antibiotic

interventions, such as the effect of mass drug administration on antibiotic resistance

(25,26), when the intervention and control populations are not wholly epidemiologically

separate.

The effect of spillover should scale with the amount of interaction between populations.

If two populations do not interact at all, then antibiotic use in one population cannot

affect resistance in the other. However, if two populations liberally exchange bacteria,

then the rates of antibiotic resistance in the two populations will be very similar,

regardless of whether their rates of antibiotic use differ greatly.

The effects of spillover also depend on population sizes. For example, two large

populations will have most interactions within themselves, rather than between each

other. Spillover should therefore be most pronounced when considering small

populations and become less important for large populations. As mentioned above, a

single individual's risk of resistance is modulated by antibiotic use in their family or in

their healthcare facility. Spillover is also observed at the level of hospitals, as the level

of resistance in one hospital appears to be affected by resistance levels in nearby

hospitals as well as by antibiotic use rates in the surrounding communities (27–29).

Presumably, when examining ever larger populations, such as US Census tracts (30),

US states, or European countries, the effect of spillover will become less important.

However, the relationship between population size and spillover effects is not well

understood.

We hypothesized that US states and European countries, which are large populations

with relatively independent public health policies, may be subject to substantially lower

levels of antibiotic resistance spillover than family- or hospital-sized populations. This

hypothesis, if true, would mean that individual states or countries could act as

independent "laboratories" of antibiotic use and resistance. If not, it means that

outpatient antibiotic resistance policy must be national or international in order to

achieve its full effect. To evaluate this hypothesis, we first use mathematical models of

antibiotic use and resistance to make quantitative predictions about the effect of

spillover between populations as a function of their amount of mutual interaction. Then,

we search for signals of spillover in observational data of antibiotic use and resistance

in US states and European countries.

METHODS

Dynamical model of antibiotic resistance

To examine how interactions between populations could theoretically affect the

association between antibiotic use and resistance, we used the within-host neutrality

(WHN) mathematical model presented by Davies *et al*. (31) and described in the

Supplemental Methods. Briefly, the model predicts the prevalence ρ of antibiotic

resistance that results from an antibiotic use rate τ in a single, well-mixed population. To

verify that conclusions drawn from the WHN model are not specific to the model

structure, we also repeated all analyses with the "D-types" model of use and resistance

(32). We selected these two models because they demonstrate coexistence between

sensitive and resistant strains at equilibrium over a wide parameter space. Parameter

values and simulation methodology for both models are in the Supplemental Methods.

In the simulations, antibiotic use is measured as monthly treatments per capita and

resistance as the proportion of colonized hosts carrying resistant strains.

To conceptually frame and clarify the question of spillover, we simulated an antibiotic

stewardship intervention experiment using a structured host population approach

inspired by Blanquart *et al*. (33). We considered pairs of an intervention population with

antibiotic use rate τ_{int} and a control population with use rate τ_{cont}. To determine how

spillover affects the intervention's measured outcome, we modulated the proportion ε of

each population's contacts that are in the other population. For $\varepsilon = 0\%$, the populations

are completely separate. For ε = 50%, contacts across populations are just as likely as

contacts within populations (Supplemental Methods). We varied ε between 0% and

50%, and we varied the difference in use $\Delta\tau = \tau_{cont} - \tau_{int}$ between 0 and 0.15 treatments

per person per month while keeping the average use $0.5 \times (\tau_{cont} + \tau_{int})$ fixed at 0.125,

reflecting the range of antibiotic use rates in the original model presentations.

Observational data

We examined antibiotic use and resistance for 3 pathogen-antibiotic combinations: *S.*

pneumoniae and macrolides, *S. pneumoniae* and β-lactams, and *Escherichia coli* and

quinolones. We considered these 3 combinations because they are the subject of many

modeling (31,32) and empirical studies (5,15).

Observational data were drawn from 3 sources. First, we used MarketScan (34) and

ResistanceOpen (35) as previously described (7). The MarketScan data includes

outpatient pharmacy antibiotic prescription claims for 62 million unique people during

2011-2014. ResistanceOpen includes antibiotic resistance data collected during 2012-

2015 from 230 hospitals, laboratories, and surveillance units in 44 states. Second, we

used the QuintilesIMS Xponent database (36) and the US Centers for Disease Control

and Prevention's (CDC) National Healthcare Safety Network (NHSN) (37). The Xponent

data includes state-level data on US quinolone use during 2011-2014. NHSN includes

state-level data on quinolone resistance among *E. coli* catheter-associated urinary tract

infections during 2011-2014. Third, we used the European Center for Disease

Prevention and Control's (ECDC) ESAC-Net antimicrobial consumption database (38)

and EARS-Net Surveillance Atlas of Infectious Disease (39) for 2011-2015. The ESAC-

Net data includes country-level outpatient antibiotic use data provided by WHO and

Ministries of Health from member countries. The EARS-Net data includes country-level

resistance data. In the observational data, we quantified antibiotic use as yearly

treatments per capita and resistance as the proportion of collected isolates that were

non-susceptible. Further details about preparation of these data sources and their

availability are in the Supplemental Methods.

We excluded the *S. pneumoniae* resistance to β-lactams in US states from the analysis

because, in previous work using the same primary datasets, the point estimate for the

use-resistance relationship was negative (14).

Use-resistance relationships by populations' adjacency

To test the theoretical prediction that the same difference in antibiotic use will be

associated with smaller differences in antibiotic resistance when two populations (US

states or European countries) have stronger interactions, we tested whether the use-

resistance association is weaker in adjacent pairs of populations, which presumably

have more cross-population contacts, compared to non-adjacent populations. Two

populations were considered adjacent if they share a land or river border (40–42).

We quantified the use-resistance association as the percentage point difference (i.e.,

absolute risk difference) in resistance (proportion of non-susceptible isolates) divided by

the difference in antibiotic use. We summarized use-resistance associations among

adjacent pairs and non-adjacent pairs of populations using the median value. Because

use-resistance associations between pairs of populations are correlated, we used the

jackknife method to compute confidence intervals on the difference in medians between

groups. We used the Mann-Whitney U test to compute statistical significance. Because

our theoretical results suggested the ratio of percentage points difference in resistance

divided by difference in antibiotic use rates was a predictable function of the degree of

population mixing, we considered only this functional form for the use-resistance

association.

Use-resistance associations by interactions

Because adjacency might be too coarse measure of populations' interactions to detect

spillover, we performed a similar analysis as above, but predicting the use-resistance

association between populations using transportation data. For US states, we used

inter-county commuting statistics from the US Census (43). For European countries, we

used inter-country passenger flight data from Eurostat (44). Rather than trying to infer a

precise mathematical relationship between transportation statistics and epidemiological

contacts, we used a nonparametric approach: we assumed that pairs of populations

with relatively little inter-population transportation also have relatively few inter-

population contacts, but we infer only the rank ordering of inter-population contacts, not

their magnitudes. Specifically, we first converted the matrix of the number of counts

(workers in the commuting data and passengers in the flight data) from each population

to every other population into the proportion of counts moving from one population to

another (i.e., divided each row by its sum), then symmetrized the resulting matrix (taking

the elementwise average of the matrix and its transpose), and finally converted the

resulting values into ranks. We assumed that intra-population interactions outnumber

inter-population interactions and so set diagonal entries, which represent within-

population interactions, to the highest rank. We measured the association between

ranked interactions and use-resistance associations using the nonparametric

Spearman's correlation, computed confidence intervals using the jackknife method, and

tested for statistical significance using the Mantel test with 999 permutations. To

quantify the effect of spillover, we compared the median use-resistance associations

among the top and bottom decile of ranked interactions.

Simulations and observational analyses were made using R (version 3.6.0) (45). The

Mantel test used the *vegan* package (46). Multiple hypotheses were accounted for using

Benjamini-Hochberg false discovery rate.

RESULTS

In simulations of two populations, representing an intervention and control group,

interactions between the two populations attenuated the effect of the intervention

(Figure 1). With increasing interaction strength, the same intervention, that is, the same

difference in antibiotic use between the populations, was associated with a smaller

difference in antibiotic resistance. The difference in resistance between populations

increases with the difference in antibiotic use (Figure 1d), but the use-resistance

association, measured as the ratio of the difference in resistance to the difference in

use, depends strongly on the interaction strength (Figure 1e). Thus, spillover between

populations attenuates the measured use-resistance association.

The precise relationship between ε, the proportion of each population's contacts that are

in the other population, and the attenuation of the use-resistance association depended

on the choice of mathematical model (Supplemental Table 1, Supplemental Figure 1).

For $\varepsilon = 1\%$, the use-resistance declined by approximately 30% in the WHN model and

more than 60% in the "D-types" model. In other words, the models predict that as few as

1% of contacts need to be across populations, rather than within populations, to cause

the observed effect of an antibiotic stewardship intervention to shrink by one-third, or

even half.

To test whether spillover is important at the scale of US states or European countries,

we measured use-resistance associations between pairs of populations in 6

combinations of pathogen species, antibiotic class, and data source (Figure 2). We

reasoned that if spillover is relevant at these scales, then pairs of states or countries

with stronger interactions would have detectably weaker use-resistance associations.

We first tested whether pairs of physically adjacent populations (e.g., Massachusetts

and Connecticut) had weaker use-resistance associations than non-adjacent

populations (e.g., Massachusetts and Alaska). In 5 of 6 pathogen/antibiotic/dataset

combinations, the median use-resistance association was smaller among adjacent

populations than among non-adjacent populations (Figure 3), but in no case was the

difference statistically significant after multiple hypothesis correction (Supplemental

Table 2). Point estimates of the relative difference in median use-resistance

associations between adjacent populations were 18% to 50% weaker than between

non-adjacent populations (excepting *S. pneumoniae* with β-lactams, which was an

outlier), consistent with the theoretical modeling results showing a 50% reduction in the

use-resistance association for populations with approximately 1% of interactions across

populations (Supplemental Table 1).

Next, to account for the possibility that adjacency was too coarse a measure for

interactions between populations, we instead used a rank-ordered estimate of

interactions using US commuting and European airline passenger flows (Supplemental

Figure 2). In 4 of 6 dataset/pathogen/antibiotic combinations, the nonparametric

association between increased inter-population interactions and decreased use-

resistance associations was statistically significant, thus confirming the general trend

observed in the adjacency analysis (Figure 4, Supplemental Table 3). The weakest

significant result was for *S. pneumoniae* and macrolides in the

MarketScan/ResistanceOpen dataset (Spearman's ρ 0.07, 95% jackknife confidence

interval -0.03 to 0.18; p = 0.028, Mantel test), and the strongest was for *E. coli* and

quinolones in the Xponent/NHSN dataset (ρ = 0.13, 95% jackknife confidence interval

0.007 to 0.25; p = 0.001). The correlation for *S. pneumoniae* and macrolides in the

ECDC data has a point estimate suggesting spillover but was not statistically significant,

while the correlation for *S. pneumoniae* and β-lactams in the ECDC data had the

opposite point estimate, consistent with the adjacency analysis (Figure 3).

Finally, to quantify the effect of increased interactions on the observed use-resistance

associations, we compared the use-resistance associations in pairs of populations

within the lowest decile of interactions against those in the highest decile, using the

same approach as for the adjacency analysis above (Supplemental Table 4). In 5 of 6

dataset/pathogen/antibiotic combinations, the point estimate for the different in use-

resistance associations was consistent with spillover, with a weaker association among

pairs of populations with greater interactions. In those 5 cases, the point estimates

ranged from a 18% reduction up to a 75% reduction in use-resistance associations

among the highest-interacting pairs of populations, compared to the lowest-interacting

populations.

DISCUSSION

We used theoretical models to show that interactions between two populations can

attenuate the observed use-resistance association. In simulations, the quantitative

relationship between inter-population interactions and the attenuation of the use-

resistance association was dependent on the theoretical model used. However, we

found that, in two models of the use-resistance association, having on the order of 1%

of interactions between a control and intervention population was sufficient to attenuate

the observed effect of theoretical stewardship intervention by 50%, relative to a situation

where the two populations were completely isolated. These theoretical results suggest

that even small numbers of interactions could lead to substantial spillover.

When examining observational antibiotic use and resistance data from US states and

European countries, we did not detect a robust signal of spillover among pairs of

adjacent populations, as opposed to non-adjacent pairs, even across 3 pathogen-

antibiotic combinations in 3 separate datasets. However, when using more fine-grained

transportation data to estimate the relative ranking of epidemiological contacts between

those populations, we found a correlation between increased interactions and

attenuated use-resistance associations. Pairs of populations in the highest decile of

inter-population interactions, that is, those most subject to spillover, had use-resistance

associations on the order of 50% weaker than pairs in the lowest decile of interactions.

The 2 pathogen/antibiotic dataset combinations with data not indicative of spillover,

namely *S. pneumoniae* and β-lactams and macrolides in the ECDC data, may have not

shown the same signal as other cases because of the smaller number of populations in

those cases (27, versus 28 to 50 in the other cases) led to insufficient statistical power

or potentially because the biology or epidemiology of *S. pneumoniae* resistance in these

cases is somehow different and does not exhibit spillover.

These theoretical and empirical results suggest that spillover is relevant at the level of

US states and European countries. This finding has important ramifications. First,

attempts to attribute changes in a population's level of antibiotic resistance to changes

in that population's rates of antibiotic use may lead to inaccurate conclusions unless use

and resistance in surrounding populations is accounted for. Second, state- or country-

level antibiotic stewardship pilot studies may substantially underestimate the potential

reduction in antibiotic resistance that would follow from a reduction in antibiotic use if

that reduction were implemented at a larger scale. Third, mass drug administration trials

may lead to elevated levels of antibiotic resistance in the control populations if those

populations are not entirely separated from the intervention population. Finally, spillover

can at least partly explain why use-resistance associations at the level of US states or

European countries are sometimes difficult to detect and, when they are detected, are

sometimes weaker than expected (5,11,14). Furthermore, spillover means that

theoretical models of antibiotic use and resistance that treat US states or European

countries as epidemiologically independent populations will not accurately represent the

dynamics of resistance (33).

Our study has several limitations. First, we interpreted the theoretical results and

ecological data as if the association between antibiotic use and resistance were causal

and deterministic. However, decreases in the use of an antibiotic may not necessarily

lead to declines in resistance to that antibiotic in a target pathogen (12,47–49). We do

not address co-resistance and cross-selection (50,51), and we assumed that resistance

equilibrates on a timescale comparable to an intervention. Previous research has shown

that resistance among *E. coli*, *S. pneumoniae*, *N. gonorrhoeae* and other organisms can

respond to changes in antibiotic use on the timescale of months (52–55), but the

expected delay between a perturbation to antibiotic use and the resulting change in

resistance remains a subject of active study (13,52,56,57). Nevertheless, the use of

ecological data was essential to addressing our hypothesis, as data from multiple

controlled, state- or country-wide experiments are not available.

Second, our analyses attributed all differences in antibiotic resistance between

populations to differences in use across those populations and to interactions between

them. In fact, antibiotic resistance is associated with factors beyond antibiotic use

(6,58), and those factor are likely spatially correlated. In other words, closely interacting

populations might have more similar use-resistance associations because they tend to

be more similar with respect to other determinants of antibiotic use. Our estimates of the

correlation between inter-population interactions and the attenuation of use-resistance

relationships may therefore be overestimates. A more careful quantification of the

relative roles of spillover versus other spatially-correlated determinants of resistance is

required.

Third, our analysis only considered pairs of populations, when in fact spillover is

happening between all pairs of populations in our analysis simultaneously. We used the

pairs approach because it allowed for a simple theoretical model and a straightforward

comparison of theory with the observational data. However, more sophisticated

approaches that account for the network of spillover interactions will likely lead to more

refined characterizations of spillover.

Finally, analyses based on administrative entities like US states or European countries,

although logistically attractive "laboratories" of antibiotic stewardship, will always be

difficult to interpret because administrative entities average over important dimensions

of population structure like age (59), sexual networks (60), and race/ethnicity (61). Thus,

use-resistance associations measured across states and countries may be different

from those that appear among geographically-proximate populations with dissimilar

antibiotic use rates, such as the sexes (62) and racial/ethnic groups (63).

We suggest 3 lines of investigation that could refine our understanding about the role of

spillover at levels of US states and European countries. First, further mathematical

modeling studies with more realistic structuring of the host population might articulate

more detailed theoretical expectations about the relationship between intervention scale

and spillover. For example, models could be parameterized with epidemiological

information about individuals' contacts and travel patterns, as has been done for other

infectious diseases (64). Second, meta-analysis of existing studies of use-resistance

relationships (5,65,66), both experimental and observational, could potentially

determine the empirical relationship between intervention population size and the

importance of spillover. This kind of meta-analysis might reveal that populations other

than US states are feasible "laboratories" for resistance: it may be that cities, daycares,

schools, workplaces, or even families represent the optimal trade-off between

maximizing logistical feasibility and minimizing spillover. Finally, future experimental

outpatient antibiotic stewardship interventions should make careful and deliberate

decisions about the sizes and interconnectedness of the populations they target. We

hope that a better understanding of spillover will improve predictions about the future of

antibiotic resistance, the formulation of stewardship policy, the design of stewardship

interventions and antibiotic administration trials, and theoretical models of resistance.

References

1.   Review on Antimicrobial Resistance. Tackling drug-resistant infections globally:
final report and recommendations. 2016.

2.   Public Health England. English Surveillance Programme for Antimicrobial
Utilisation and Resistance (ESPAUR). 2014.

3.   Public Health Agency of Sweden, National Veterinary Institute. Consumption of
antibiotics and occurrence of antibiotic resistance in Sweden.

4.   US Centers for Disease Control and Prevention. Antibiotic Resistance Threats in
the United Staes, 2013. 2013.

5.   Bell BG, Schellevis F, Stobberingh E, Goossens H, Pringle M. A systematic review
and meta-analysis of the effects of antibiotic consumption on antibiotic resistance.
BMC Infect Dis. 2014 Jan 9;14:13.

6.   MacFadden DR, McGough SF, Fisman D, Santillana M, Brownstein JS. Antibiotic
resistance increases with local temperature. Nat Clim Change. 2018 Jun;8(6):510–
4.

7.   Olesen SW, Barnett ML, MacFadden DR, Brownstein JS, Hernández-Diaz S,
Lipsitch M, et al. The distribution of antibiotic use and its association with antibiotic
resistance. eLife [Internet]. 2018 Dec 18 [cited 2018 Dec 19];7. Available from:
https://elifesciences.org/articles/39435

8.   Goossens H, Ferech M, Vander Stichele R, Elseviers M. Outpatient antibiotic use
in Europe and association with resistance: a cross-national database study. The
Lancet. 2005 Feb 12;365(9459):579–87.

9.   van de Sande-Bruinsma N, Grundmann H, Verloo D, Tiemersma E, Monen J,
Goossens H, et al. Antimicrobial drug use and resistance in Europe. Emerg Infect
Dis. 2008 Nov;14(11):1722–30.

10. Livermore DM. The 2018 Garrod Lecture: Preparing for the Black Swans of
resistance. J Antimicrob Chemother. 2018 Nov 1;73(11):2907–15.

11. Turnidge J, Christiansen K. Antibiotic use and resistance--proving the obvious.
Lancet Lond Engl. 2005 Feb;365(9459):548–9.

12. Arason VA, Gunnlaugsson A, Sigurdsson JA, Erlendsdottir H, Gudmundsson S,
Kristinsson KG. Clonal Spread of Resistant Pneumococci Despite Diminished
Antimicrobial Use. Microb Drug Resist. 2002 Sep;8(3):187–92.

13. Lipsitch M. The rise and fall of antimicrobial resistance. Trends Microbiol. 2001
Sep;9(9):438–44.

14. Olesen SW, Barnett ML, MacFadden DR, Brownstein JS, Hernández-Díaz S,
Lipsitch M, et al. The distribution of antibiotic use and its association with antibiotic
resistance. eLife [Internet]. 2018 Dec 18 [cited 2019 Sep 23];7. Available from:
https://elifesciences.org/articles/39435

15. Costelloe C, Metcalfe C, Lovering A, Mant D, Hay AD. Effect of antibiotic
prescribing in primary care on antimicrobial resistance in individual patients:
systematic review and meta-analysis. BMJ. 2010 May 18;340:c2096.

16. Harbarth S, Harris AD, Carmeli Y, Samore MH. Parallel analysis of individual and
aggregated data on antibiotic exposure and resistance in gram-begative bacilli. Clin
Infect Dis. 2001;33:1462–8.

17. Lipsitch M. Measuring and Interpreting Associations between Antibiotic Use and
Penicillin Resistance in Streptococcus pneumoniae. Clin Infect Dis. 2001 Apr
1;32(7):1044–54.

18. Hannah EL, Angulo FJ, Johnson JR, Haddadin B, Williamson J, Samore MH. Drug-
resistant Escherichia coli, Rural Idaho. Emerg Infect Dis. 2005 Oct;11(10):1614–7.

19. Kalter HD, Gilman RH, Moulton LH, Cullotta AR, Cabrera L, Velapatiño B. Risk
factors for antibiotic-resistant Escherichia coli carriage in young children in Peru:
community-based cross-sectional prevalence study. Am J Trop Med Hyg. 2010
May;82(5):879–88.

20. Samore MH, Magill MK, Alder SC, Severina E, Morrison-De Boer L, Lyon JL, et al.
High rates of multiple antibiotic resistance in Streptococcus pneumoniae from
healthy children living in isolated rural communities: association with cephalosporin
use and intrafamilial transmission. Pediatrics. 2001 Oct;108(4):856–65.

21. Gottesman B-S, Low M, Almog R, Chowers M. Quinolone Consumption by Mothers
Increases Their Children's Risk of Acquiring Quinolone-Resistant Bacteriuria. Clin
Infect Dis [Internet]. 2019 Aug 29 [cited 2019 Dec 2]; Available from:
https://academic.oup.com/cid/advance-article/doi/10.1093/cid/ciz858/5556473

22. Lipsitch M, Samore MH. Antimicrobial use and antimicrobial resistance: a
population perspective. Emerg Infect Dis. 2002;8:347–54.

23. Moellering RC. NDM-1 — A Cause for Worldwide Concern. N Engl J Med. 2010
Dec 16;363(25):2377–9.

24. Wang R, van Dorp L, Shaw LP, Bradley P, Wang Q, Wang X, et al. The global
distribution and spread of the mobilized colistin resistance gene mcr-1. Nat
Commun [Internet]. 2018 Dec [cited 2020 Jun 26];9(1). Available from:
http://www.nature.com/articles/s41467-018-03205-z

25. Bogoch II, Utzinger J, Lo NC, Andrews JR. Antibacterial mass drug administration
for child mortality reduction: Opportunities, concerns, and possible next steps.
Bhutta ZA, editor. PLoS Negl Trop Dis. 2019 May 23;13(5):e0007315.

26. Doan T, Arzika AM, Hinterwirth A, Maliki R, Zhong L, Cummings S, et al. Macrolide
Resistance in MORDOR I — A Cluster-Randomized Trial in Niger. N Engl J Med.
2019 Jun 6;380(23):2271–3.

27. Cooper BS, Medley GF, Stone SP, Kibbler CC, Cookson BD, Roberts JA, et al.
Methicillin-resistant Staphylococcus aureus in hospitals and the community: Stealth
dynamics and control catastrophes. Proc Natl Acad Sci. 2004 Jul 6;101(27):10223–
8.

28. MacFadden DR, Fishman DN, Hanage WP, Lipsitch M. The Relative Impact of
Community and Hospital Antibiotic Use on the Selection of Extended-Spectrum
Beta-lactamase-Producing Escherichia coli. Clin Infect Dis [Internet]. 2018 [cited
2018 Nov 29]; Available from: https://www.ncbi.nlm.nih.gov/pubmed/30462185

29. Knight GM, Costelloe C, Deeny SR, Moore LSP, Hopkins S, Johnson AP, et al.
Quantifying where human acquisition of antibiotic resistance occurs: a
mathematical modelling study. BMC Med. 2018 23;16(1):137.

30. Klevens RM, Caten E, Olesen SW, DeMaria A, Troppy S, Grad YH. Outpatient
Antibiotic Prescribing in Massachusetts, 2011–2015. Open Forum Infect Dis
[Internet]. 2019 May 1 [cited 2019 Sep 15];6(5). Available from:
https://academic.oup.com/ofid/article/doi/10.1093/ofid/ofz169/5430941

31. Davies NG, Flasche S, Jit M, Atkins KE. Within-host dynamics shape antibiotic
resistance in commensal bacteria. Nat Ecol Evol. 2019 Mar;3(3):440–9.

32. Lehtinen S, Blanquart F, Croucher NJ, Turner P, Lipsitch M, Fraser C. Evolution of
antibiotic resistance is linked to any genetic mechanism affecting bacterial duration
of carriage. Proc Natl Acad Sci. 2017 Jan 31;114(5):1075–80.

33. Blanquart F, Lehtinen S, Lipsitch M, Fraser C. The evolution of antibiotic resistance
in a structured host population. J R Soc Interface. 2018 Jun;15(143).

34. Truven Health MarketScan Database. Commercial Claims and Encounters. Ann
Arbor, MI; 2015.

35. MacFadden DR, Fisman D, Andre J, Ara Y, Majumder MS, Bogoch II, et al. A
Platform for Monitoring Regional Antimicrobial Resistance, Using Online Data
Sources: ResistanceOpen. J Infect Dis. 2016 Dec 1;214(suppl_4):S393–8.

36. US Centers for Disease Control and Prevention. Patient Safety Atlas - Outpatient
Antibiotic Use [Internet]. [cited 2018 Oct 25]. Available from:
https://gis.cdc.gov/grasp/PSA/indexAU.html

37. US Centers for Disease Control and Prevention. Patient Safety Atlas - Antibiotic
Resistance [Internet]. [cited 2018 Oct 25]. Available from:
https://gis.cdc.gov/grasp/PSA/AboutTheData.html

38. European Centre for Disease Prevention and Control. Antimicrobial consumption
database [Internet]. European Centre for Disease Prevention and Control. [cited
2018 Oct 25]. Available from: http://ecdc.europa.eu/en/antimicrobial-
consumption/surveillance-and-disease-data/database

39. European Centre for Disease Prevention and Control. Surveillance Atlas of
Infectious Diseases [Internet]. [cited 2018 Oct 25]. Available from:
https://atlas.ecdc.europa.eu/public/index.aspx

40. US Census. County Adjacency File [Internet]. Available from:
https://www.census.gov/geo/reference/county-adjacency.html

41. Correlates of War Project. Direct Contiguity Data, 1816-2016 (v3.2) [Internet].
Available from: http://correlatesofwar.org/data-sets/direct-contiguity

42. Stinnett DM, Tir J, Diehl PF, Schafer P, Gochman C. The Correlates of War (Cow)
Project Direct Contiguity Data, Version 3.0. Confl Manag Peace Sci. 2002
Sep;19(2):59–67.

43. US Census. 2011-2015 5-Year ACS Commuting Flows [Internet]. Available from:
https://www.census.gov/data/tables/2015/demo/metro-micro/commuting-flows-
2015.html

44. Eurostat. Air transport measurement [Internet]. Available from:
https://ec.europa.eu/eurostat/cache/metadata/en/avia_pa_esms.htm

45. R Core Team. R: A language and environment for statistical computing [Internet].
Vienna; 2018. Available from: www.R-project.org

46. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, et al. vegan:
Community Ecology Package. R package version 2.5-6.

47. Hennessy TW, Petersen KM, Bruden D, Parkinson AJ, Hurlburt D, Getty M, et al.
Changes in Antibiotic-Prescribing Practices and Carriage of Penicillin-Resistant
Streptococcus pneumoniae: A Controlled Intervention Trial in Rural Alaska. Clin
Infect Dis. 2002 Jun 15;34(12):1543–50.

48. Sundqvist M, Geli P, Andersson DI, Sjölund-Karlsson M, Runehagen A, Cars H, et
al. Little evidence for reversibility of trimethoprim resistance after a drastic
reduction in trimethoprim use. J Antimicrob Chemother. 2010 Feb;65(2):350–60.

49. Enne VI, Livermore DM, Stephens P, Hall LM. Persistence of sulphonamide
resistance in Escherichia coli in the UK despite national prescribing restriction.
Lancet Lond Engl. 2001 Apr 28;357(9265):1325–8.

50. Pouwels KB, Freeman R, Muller-Pebody B, Rooney G, Henderson KL, Robotham
JV, et al. Association between use of different antibiotics and
trimethoprimresistance: going beyond the obvious crude association. J Antimicrob
Chemoth. 2018;

51. Tedijanto C, Olesen SW, Grad YH, Lipsitch M. Estimating the proportion of
bystander selection for antibiotic resistance among potentially pathogenic bacterial
flora. Proc Natl Acad Sci. 2018 Dec 18;115(51):E11988–95.

52. Olesen SW, Torrone EA, Papp JR, Kirkcaldy RD, Lipsitch M, Grad YH.
Azithromycin Susceptibility Among Neisseria gonorrhoeae Isolates and Seasonal
Macrolide Use. J Infect Dis [Internet]. 2018 [cited 2018 Nov 6]; Available from:
http://academic.oup.com/jid/advance-article/doi/10.1093/infdis/jiy551/5098400

53. Dagan R, Barkai G, Givon-Lavi N, Sharf AZ, Vardy D, Cohen T, et al. Seasonality
of Antibiotic-Resistant Streptococcus pneumoniae That Causes Acute Otitis Media:
A Clue for an Antibiotic-Restriction Policy? J Infect Dis. 2008 Apr 15;197(8):1094–
102.

54. Sun L, Klein EY, Laxminarayan R. Seasonality and Temporal Correlation between
Community Antibiotic Use and Resistance in the United States. Clin Infect Dis.
2012 Sep 1;55(5):687–94.

55. Blanquart F, Lehtinen S, Fraser C. An evolutionary model to predict the frequency
of antibiotic resistance under seasonal antibiotic use, and an application to
Streptococcus pneumoniae. Proc R Soc B. 2017 May 31;284(1855):20170679.

56. Dingle KE, Didelot X, Quan TP, Eyre DW, Stoesser N, Golubchik T, et al. Effects of
control interventions on Clostridium difficile infection in England: an observational
study. Lancet Infect Dis. 2017 Apr;17(4):411–21.

57. McCormick AW, Whitney CG, Farley MM, Lynfield R, Harrison LH, Bennett NM, et
al. Geographic diversity and temporal trends of antimicrobial resistance in
Streptococcus pneumoniae in the United States. Nat Med. 2003 Apr;9(4):424–30.

58. Collignon P, Beggs JJ, Walsh TR, Gandra S, Laxminarayan R. Anthropological and
socioeconomic factors contributing to global antimicrobial resistance: a univariate
and multivariable analysis. Lancet Planet Health. 2018 Sep;2(9):e398–405.

59. Mossong J, Hens N, Jit M, Beutels P, Auranen K, Mikolajczyk R, et al. Social
contacts and mixing patterns relevant to the spread of infectious diseases. PLoS
Med. 2008 Mar 25;5(3):e74.

60. Garett G, Hughes J, Anderson R, Stoner B, Aral S, Whittington W, et al. Sexual
mixing patterns of patients attending sexually transmitted diseases clinics. Sex
Transm Dis. 1996;23(3):248–57.

61.   Newman MEJ. Mixing patterns in networks. Phys Rev E Stat Nonlin Soft Matter
Phys. 2003 Feb;67(2 Pt 2):026126.

62.   Hicks LA, Bartoces MG, Roberts RM, Suda KJ, Hunkler RJ, Taylor TH, et al. US
Outpatient Antibiotic Prescribing Variation According to Geography, Patient
Population, and Provider Specialty in 2011. Clin Infect Dis. 2015 May
1;60(9):1308–16.

63.   Olesen SW, Grad YH. Racial/Ethnic Disparities in Antimicrobial Drug Use, United
States, 2014–2015. Emerg Infect Dis [Internet]. 2018 [cited 2018 Nov 6];24(11).
Available from: https://wwwnc.cdc.gov/eid/article/24/11/18-0762_article

64.   Charu V, Zeger S, Gog J, Bjørnstad ON, Kissler S, Simonsen L, et al. Human
mobility and the spatial transmission of influenza in the United States. PLoS
Comput Biol. 2017;13(2):e1005382.

65.   Schechner V, Temkin E, Harbarth S, Carmeli Y, Schwaber MJ. Epidemiological
interpretation of studies examining the effect of antibiotic usage on resistance. Clin
Microbiol Rev. 2013 Apr;26(2):289–307.

66.   O'Brien KS, Emerson P, Hooper P, Reingold AL, Dennis EG, Keenan JD, et al.
Antimicrobial resistance following mass azithromycin distribution for trachoma: a
systematic review. Lancet Infect Dis [Internet]. 2018 Oct 3 [cited 2018 Nov 7];
Available from:
http://www.sciencedirect.com/science/article/pii/S1473309918304444

Disclaimers

The views and opinions of the authors expressed herein do not necessarily state or

reflect those of the ECDC. The accuracy of the authors' statistical analysis and the

findings they report are not the responsibility of ECDC. ECDC is not responsible for

conclusions or opinions drawn from the data provided. ECDC is not responsible for the

correctness of the data and for data management, data merging and data collation after

provision of the data. ECDC shall not be held liable for improper or incorrect use of the

data.

Funding

This work was supported by the National Institutes of Health (grant number

U54GM088558 to ML). The funders had no role in study design, data collection and

interpretation, or the decision to submit the work for publication.

Acknowledgements

We thank Dr. Stephen M. Kissler for helpful comments on the manuscript.

Figure 1. **Interactions between populations attenuate the effect of interventions.**

(*a*) Schematic of the 2-population model. (*b*) Results of simulations of the 2-population

WHN model for a modest intervention (difference in antibiotic use between populations

$\Delta\tau$ = 0.05 monthly treatments per capita; average of control and intervention treatment

rates 0.125). As interaction strength (ε, horizontal axis) increases, the difference in

antibiotic resistance between the two populations decreases. Dotted line shows

resistance level in populations before the intervention. (*c*) The same pattern holds for a

stronger intervention ($\Delta\tau$ = 0.1, same average treatment rate). (*d*) In general, the

difference in resistance between populations ($\Delta\rho$, vertical axis) increases with the

difference in antibiotic use ($\Delta\tau$, horizontal axis). (*e*) However, in the WHN model, the

use-resistance relationship ($\Delta\rho/\Delta\tau$, vertical axis) depends mostly on the interaction

strength ε and is mostly independent of the difference in antibiotic use $\Delta\tau$.

Figure 2. **Use-resistance relationships across US states and European countries.**

Each point represents antibiotic use and resistance in a US state (top row) or European

country (bottom row). Lines show simple linear regression best fit. Gray areas show

95% confidence interval. Ec/q: *E. coli* and quinolones. Sp/m: *S. pneunomiae* and

macrolides. Sp/bl: *S. pneumoniae* and β-lactams. RO: ResistanceOpen. ECDC:

European CDC.

Figure 3. **Use-resistance relationships by adjacency.** Each point represents the use-

resistance association in populations (top row, US states; bottom row, European

countries), the same as those shown in Figure 2, arranged by whether the pair of

populations is physically adjacent. Physically adjacent populations tend to have weaker

use-resistance associations, but differences were not statistically significant. For visual

clarity, the vertical axes are truncated to show only the central 90% of data points. Ec/q:

E. coli and quinolones. Sp/m: *S. pneunomiae* and macrolides. Sp/bl: *S. pneumoniae*

and β-lactams. RO: ResistanceOpen. ECDC: European CDC.

Figure 4. **Use-resistance associations by ranked interaction.** Each point represents

the use-resistance association in a pair of US states (top row) or European countries

(bottom row), the same pairs as shown in Figure 3, rank ordered by increasing inter-

population interaction as inferred from transportation data. For visual clarity, the

horizontal axes are truncated to exclude outliers. The dashed red line is a visual

illustration of how increasing interaction is correlated with decreasing use-resistance

associations (robust regression; compare Supplemental Tables 3 and 4). The asterisk

(*) indicates a statistically significant association between increased interaction and

decreased use-resistance relationship (Supplemental Table 3). Ec/q: *E. coli* and

quinolones. Sp/m: *S. pneunomiae* and macrolides. Sp/bl: *S. pneumoniae* and β-

lactams. RO: ResistanceOpen. ECDC: European CDC.