1 **Outpatient antibiotic stewardship interventions: geographic scale and**
2 **associations between antibiotic use and resistance**
3
4 Short title: Geographical scales of outpatient stewardship interventions
5

6 Scott W. Olesen[1], Marc Lipsitch[1,2], Yonatan H. Grad[1,3]
7
8 1 Department of Immunology and Infectious Diseases, Harvard T. H. Chan School of
9 Public Health, Boston, Massachusetts 02115
10
11 2 Center for Communicable Disease Dynamics, Harvard T. H. Chan School of Public
12 Health, Boston, Massachusetts 02115
13
14 3 Division of Infectious Diseases, Department of Medicine, Brigham and Women's
15 Hospital, Harvard Medical School, Boston, Massachusetts 02115
16
Correspondence:
Yonatan Grad
Harvard T. H. Chan School of Public Health
665 Huntington Ave, Building 1, Room 715, Boston, Massachusetts 02115
617 432 2275
ygrad@hsph.harvard.edu
23

Abstract

Background: Antibiotic stewardship interventions aim to combat antibiotic resistance by reducing inappropriate antibiotic use. One obstacle to the rational design of outpatient stewardship programs is that small-scale pilot experiments that aim to reduce antibiotic resistance by reducing antibiotic use may produce results that are systematically different from results observed in larger-scale implementations. Here, we investigate the relationship between geographic scale and the effect of reductions in antibiotic use.

Methods and findings: First, we show that dynamical models of antibiotic resistance exhibit "spillover", such that resistance in an intervention population is partly due to antibiotic use in surrounding populations, which attenuates the intervention's effect size. Second, using observational antibiotic use and resistance data from US states and European countries for 3 pathogen-antibiotic combinations, we show that use-resistance associations are robust to aggregation above the level of US states or European countries. Finally, we did not detect differences in the strength of use-resistance associations measured between pairs of adjacent states or countries, which presumably have stronger spillover, compared to the associations among non-adjacent pairs.

Conclusions: These results imply that interventions at the level of US states will yield effect sizes that can be used to estimate the effects of regional or national interventions.

Introduction

Antibiotic resistance is a major threat to public health (1). Outpatient antibiotic use,

which accounts for approximately 80% of all human antibiotic use (2,3), is considered a

principal driver of antibiotic resistance in the community (4). Antibiotic stewardship

initiatives reduce antibiotic use with the goal of lowering healthcare costs (5), preventing

adverse drug events (6,7), and mitigating antibiotic resistance (8–10). Rational design of

stewardship initiatives requires quantitative models that predict the outcome of an

intervention. It is relatively simple to predict what reduction in antibiotic use is required

to achieve a target reduction in monetary costs or adverse events: each avoided

antibiotic prescription prevents the cost of that prescription and the risk of an adverse

event from that prescription. In contrast, quantitative predictions about how a reduction

in antibiotic use will affect antibiotic resistance are more challenging because resistance

is a complex, temporally dynamic phenomenon (11–14).

A critical feature of this complexity is that resistant bacteria can be transmitted, so that

the risk that an individual's infection is antibiotic resistant depends on that individual's

antibiotic use (15,16) as well as the rates of antibiotic use among that individual's

contacts (17), such as their family members (18–20). This effect of resistance "spilling

over" can be so strong that, for example, an individual in the hospital who has no recent

antibiotic use may have a higher risk of antibiotic resistance than an individual in the

community with a high antibiotic use rate (21). The same spillover phenomenon occurs

at the level of populations, such that resistance in a hospital can be affected by

resistance in nearby hospitals or by antibiotic use in the surrounding community (22–

24).

If resistance spills over into an outpatient stewardship intervention population from

surrounding populations not affected by the intervention, the effect on antibiotic

resistance in the intervention population may be smaller than it would be if the

intervention population were completely isolated, because spillover from the

surrounding population is not changed by the intervention. Conversely, an outpatient

stewardship intervention targeting a small population might underestimate the effect that

a certain reduction in antibiotic use would have when applied to a larger population.

As the population targeted by the intervention increases, the amount of bacterial

transmission and resistance spillover into the population presumably decreases relative

to the amount of transmission within the population, thus also mitigating the spillover

effect and providing ever more accurate predictions of an intervention's effect. It is

unclear if stewardship interventions at small scales can accurately inform the design of

interventions targeting larger populations. For example, an intervention at the level of a

city may not provide results that can be projected to predict the effects of that

intervention implemented at the level of a US state, which in turn may or may not be an

accurate prediction of a nationwide intervention's effect.

Ideally, one could determine what population size is sufficiently large to mitigate

spillover by consulting the results of randomized, controlled experiments that measure

how a reduction in antibiotic use affects resistance for the relevant pathogen and

antibiotic. In practice, interventions are often not controlled (25,26). Only a few

population-level, randomized experiments modulating antibiotic use have been

conducted (27,28), and many of those were intentional increases in antibiotic use as

part of mass drug administrations (29).

In contrast, the association between antibiotic use and antibiotic resistance has been

characterized in many observational studies, including ecological studies at the level of

US states (30–32), European countries (33,34), and smaller geographical areas (35–

37). However, even for observational data, it is not clear what kinds of populations

should be used to minimize the spillover problem (38,39). For example, larger

geographical areas would be expected to have less spillover. Aggregating smaller

geographical units into larger units for the purposes of analysis might therefore average

over the relevant scales of transmission, producing stronger correlations between use

and resistance (32,40). Conversely, it has been suggested that analyses at smaller

geographic scales may be more likely to detect relationships between use and

resistance (10), possibly because aggregating over larger areas obscures important

variations in use or resistance (41). In principle, multilevel models with individual-level

data can account for correlations between geographical units, but the selection of the

units will still affect the results (28,42,43) and few studies of antibiotic use and

resistance have assessed the sensitivity of the results to the choice of population used

in the analysis.

In this study, we aim to determine whether outpatient stewardship experiments at the

level of US states or European countries can be expected to provide accurate estimates

of the effect that the same reduction in antibiotic use would have if applied over a larger

area, indicating that interventions in smaller populations can be used to predict the

effect of an intervention in a larger population. First, we show that the spillover effect

does occur in mathematical models of antibiotic use and resistance, and we measure

how interactions between theoretical populations attenuate use-resistance associations.

Second, we look for empirical evidence that spillover has a measurably different effect

at scales above US states or European countries.

Methods

Dynamical model of antibiotic resistance

To examine how interactions between populations could theoretically affect the

association between antibiotic use and resistance, we use the within-host neutrality

(WHN) mathematical model presented by Davies *et al*. (44) and described in the

Supplemental Methods. Briefly, the model predicts the prevalence ρ of antibiotic

resistance that results from an antibiotic use rate τ in a single, well-mixed population. To

verify that conclusions drawn from the WHN model are not specific to the model

structure, we repeated all analyses with the "D-types" model of use and resistance (45).

Parameter values and simulation methodology for both models are in the Supplemental

Methods. In the simulations, antibiotic use as monthly treatments per capita and

resistance as the proportion of colonized hosts carrying resistant strains.

We adapted the WHN model to include multiple, interacting populations using a

structured host population approach inspired by Blanquart *et al.* (46). Interactions

between populations are modulated by the proportion ε of a population's contacts that

are in other populations. For $\varepsilon = 0$, each population is completely separate. For $\varepsilon = 1$,

contacts across populations are just as likely as contacts within populations

(Supplemental Methods).

We simulated a situation in which an intervention population has a lower antibiotic use

rate τ_{int} than a control population with use rate $\tau_{\mathrm{cont}} > \tau_{\mathrm{int}}$. To measure how contacts

between the two populations affect the intervention's effect size, we varied three

parameters, setting ε to each of the values 0.00, 0.01, 0.10, 0.25, and 0.50; and setting

$(\tau_{\mathrm{cont}}, \tau_{\mathrm{int}})$ to (0.15, 0.10) or (0.20, 0.05) treatments per person per month.

Mathematical models with nested population structure

To examine the effect of population structure on associations between antibiotic use

and resistance, we further adapted the multi-population model to include a nested

population structure with $n_{\mathrm{super}} \times n_{\mathrm{sub}}$ populations. The populations are grouped into

n_{super} "super-populations" representing geographic regions. Each super-population has

n_{sub} constituent subpopulations, each representing a smaller geographic area like a US

state or European country. Populations within a super-population interact according to

the parameter $\varepsilon_{\mathrm{sub}}$, while populations in different super-populations interact according to

$\varepsilon_{\mathrm{super}} \leq \varepsilon_{\mathrm{sub}}$ (Figure 1a, Supplemental Methods). The inequality encodes the idea that

US states within a region will interact more strongly with one another than with states in

other regions, for example.

To measure the effect of population structure on use-resistance associations, we set

$n_{\text{super}} = n_{\text{sub}} = 4$ and varied three parameters, setting ε_{sub} to 0.00, 0.01, 0.10, and 0.50;

$\varepsilon_{\text{super}}$ to the same values, subject to $\varepsilon_{\text{super}} \leq \varepsilon_{\text{sub}}$; and setting τ_i to a range of values

between 0.05 and 0.20 treatments per person per month (Supplemental Table 1).

Observational data

In this study, we examined antibiotic use and resistance for 3 pathogen-antibiotic

combinations: *S. pneumoniae* and macrolides, *S. pneumoniae* and β-lactams, and

Escherichia coli and quinolones. We considered these 3 combinations because they are

the subject of many modeling (44,45) and empirical studies (15,30).

Observational data were drawn from 3 sources. First, we used MarketScan (47) and

ResistanceOpen (48) as previously described (32). The MarketScan data includes

outpatient pharmacy antibiotic prescription claims for 62 million unique people during

2011-2014. ResistanceOpen includes antibiotic resistance data collected during 2012-

2015 from 230 hospitals, laboratories, and surveillance units in 44 states. Second, we

used the QuintilesIMS Xponent database (49) and the US Centers for Disease Control

and Prevention's (CDC) National Healthcare Safety Network (NHSN) (50). The Xponent

data includes state-level data on US quinolone use during 2011-2014. NHSN includes

state-level data on quinolone resistance among *E. coli* catheter-associated urinary tract

infections during 2011-2014. Third, we used the European Center for Disease

Prevention and Control's (ECDC) ESAC-Net antimicrobial consumption database (51)

and EARS-Net Surveillance Atlas of Infectious Disease (52) for 2011-2015. The ESAC-

Net data includes country-level outpatient antibiotic use data provided by WHO and

Ministries of Health from member countries. The EARS-Net data includes country-level

resistance data. In the observational data, we quantified antibiotic use as yearly

treatments per capita and resistance as the proportion of collected isolates that were

non-susceptible. Further details about preparation of these data sources and their

availability are in the Supplemental Methods.

Use-resistance associations by scale of aggregation

To test the idea that use-resistance associations are stronger when analyzing larger

populations, presumably by decreasing spillover, we measured use-resistance

associations when US states were aggregated into the 9 Census divisions or 4 Census

regions and when European countries were aggregated into the 4 United Nations

geoscheme sub-regions (53). Aggregate antibiotic use rates were computed as the

population-weighted mean antibiotic use (Supplemental Methods). Aggregate

resistance values were computed by summing the numerator number of resistant

isolates and the denominator number of total isolates. Use-resistance associations were

measured by logistic regression. Confidence intervals on the regression fits were

evaluated using 1,000 bootstrap replications.

Use-resistance relationships by adjacency

To test the idea that the same difference in antibiotic use will be associated with smaller

differences in antibiotic resistance when the two populations have stronger interactions,

we tested whether the use-resistance association is weaker for geographic units (US

states or European countries) that are physically adjacent to one another. Two units

were considered adjacent if they share a land or river border (Supplemental Methods).

We performed robust linear regressions (Tukey's bisquare) predicting the log odds ratio

of resistance between two units. Regressions were computed using the *rlm* function in

the MASS package (54) in R (version 3.5.1) (55). Predictors in the model were the

differences in antibiotic use, population density, per capita income, and mean

temperature (31) between the two units (Supplemental Methods). The model also

included an interaction term between antibiotic use and adjacency, which allows

adjacent pairs of geographic units to have a different use-resistance association from

non-adjacent pairs:

$$\Delta \mathrm{LO}(\rho)_i = \beta_\tau (\Delta \tau)_i + \beta_{\tau a} (\Delta \tau)_i a_i + \beta_{\mathrm{dens}} \Delta \mathrm{dens}_i + \beta_{\mathrm{income}} \Delta \mathrm{income}_i + \beta_{\mathrm{temp}} \Delta \mathrm{temp}_i + \varepsilon_i$$

where *i* indexes the pairs of units, $\Delta \mathrm{LO}(\rho)$ is the log odds ratio of resistance between the

two units $\Delta \tau$ is the difference in antibiotic use, *a* is a flag for whether the units in the pair

are adjacent, and ε is the error term. Confidence intervals on the regression fits were

evaluated using 1,000 bootstrap replications resampling the geographic units and

assembling new lists of pairs in each replication.

Results

In simulations of two populations, representing an intervention and control group,

interactions between the two groups attenuated the effect of the intervention (Figure 1).

With increasing interaction strength, the same difference in antibiotic use between the

populations was associated with a smaller difference in antibiotic resistance.

Similar results held for the D-types model (Supplemental Figure 1).

In simulations of nested populations, with state-like populations grouped into region-like

"super-populations", interactions within super-populations modulate use-resistance

associations within super-populations, while interactions across super-populations

modulate the use-resistance association across all populations (Figure 2a). When

aggregating the populations into super-populations, the use-resistance associations

across all super-populations are similar to the associations across all populations

(Figure 2b). However, analysis of pairs of populations can detect the within-super-

population interactions (Figure 2c) because pairs of populations from different super-

populations tend to have differences in resistance that scale with their differences in

antibiotic use, while pairs in the same super-population tend to have much smaller

differences in resistance for the same differences in antibiotic use. Similar results were

observed in the D-types model (Supplemental Figure 2).

In observational data of antibiotic use and resistance for 3 pathogen-antibiotic

combinations, we found that aggregating geographic units (US states or European

countries) into regional units (US Census division, US Census regions, or European

regions) produced similar use-resistance associations. Associations varied by

pathogen-antibiotic-dataset combination (Figure 3, Supplemental Figures 3). However,

similar to the theoretical prediction (Figure 2b), associations were similar when

measured across the original geographic units or across regional aggregations of those

units (Supplemental Figure 4).

Using the observational data, we evaluated whether use-resistance associations

between pairs of US states or European countries were weaker for adjacent pairs than

for non-adjacent pairs, as occurred for some parameterizations in theoretical

simulations (Figure 2c). We found no evidence for differences in the use-resistance

associations among adjacent pairs compared to non-adjacent pairs (Figure 4,

Supplemental Figure 5, Supplemental Table 2).

Discussion

We used theoretical models to show that interactions between a control and intervention

group can attenuate the reduction in antibiotic resistance expected from an antibiotic

stewardship intervention. However, consistent with at least one previous study (40),

empirical data did not provide robust evidence that aggregating US states or European

countries into regions yielded stronger use-resistance associations. Furthermore, the

same difference in antibiotic use between a pair of US states or European countries

was associated with similar differences in antibiotic resistance between the units in the

pair regardless of whether the units were physically adjacent or not. These results

suggest that spillover at the level of US states and European countries is not

substantially stronger than spillover at regional scales. Thus, outpatient stewardship

experiments at the level of US states may have effect sizes similar to those that would

be achieved in a national intervention. States may serve as accurate pilot populations

for designing national interventions.

Our study has multiple limitations. First, we used observational data to address

questions about the design of outpatient stewardship interventions, which requires

interpreting the theoretical results and ecological data as if the association between

antibiotic use and resistance were causal and deterministic. In fact, antibiotic resistance

is associated with factors beyond antibiotic use (31,56), and we used only a limited

number of determinants of resistance besides antibiotic use in our distance analysis.

Second, decreases in the use of an antibiotic may not necessarily lead to declines in

resistance to that antibiotic in a target pathogen (13,27,57,58). We do not address co-

resistance and cross-selection (59,60), and we assumed that resistance equilibrates on

a timescale comparable to the intervention. Previous research has shown that

resistance among *E. coli*, *S. pneumoniae*, *N. gonorrhoeae* and other organisms can

respond to changes in antibiotic use on the timescale of months (61–64), but the

expected delay between a perturbation to antibiotic use and the resulting change in

resistance remains a subject of active study (14,61,65,66).

Third, we only considered geographical populations. Although people within a US state

interact more often with other residents of that state than with residents of other states,

geography averages over important dimensions of population structure like age (67),

sexual networks (68), and race/ethnicity (69). Use-resistance relationships measured

across geographical units may be different from those that appear among

geographically-proximate populations with dissimilar antibiotic use rates, such as the

sexes (70) and racial/ethnic groups (71).

A final caveat is that our data sources limited us to analyzing geographical populations

at or larger than the scale of US states or European countries. Previous research has

shown that spillover is important for individuals (17–20), and the results of this study

suggest that US states and European countries do not have substantially stronger

spillover than larger regions, but the importance of spillover at smaller scales remains

unclear. Depending on the epidemiology of bacterial transmission and the distribution of

antibiotic use within the targeted populations, it may be that cities, daycares, schools,

workplaces, or even families represent the optimal trade-off between logistical feasibility

and the accuracy of measured effect size for a particular pathogen and antibiotic.

We suggest 3 lines of investigation that could help address the knowledge gap about

the important of spillover at levels between individuals and US states or European

countries. First, further mathematical modeling studies with more realistic structuring of

the host population might articulate more detailed theoretical expectations about the

relationship between intervention scale and spillover. For example, models could be

parameterized with epidemiological information about individuals' contacts and travel

patterns, as has been done for other infectious diseases (72). Second, meta-analysis of

existing studies of use-resistance relationships (28–30), both experimental and

observational, might determine how increasing population scales are associated with

increasing use-resistance associations. Finally, future experimental outpatient antibiotic

stewardship interventions should make careful and deliberate decisions about the sizes

and interconnectedness of the populations they target. The results of this study suggest

that outpatient interventions can be effective at scales smaller than US states. We hope

this means that outpatient stewardship can be effectively addressed by more

organizations, such as state and city health departments.

References

1.   Review on Antimicrobial Resistance. Tackling drug-resistant infections globally: final report and recommendations. 2016.

2.   Public Health England. English Surveillance Programme for Antimicrobial Utilisation and Resistance (ESPAUR). 2014.

3.   Public Health Agency of Sweden, National Veterinary Institute. Consumption of antibiotics and occurrence of antibiotic resistance in Sweden.

4.   US Centers for Disease Control and Prevention. Antibiotic Resistance Threats in the United Staes, 2013. 2013.

5.   US Centers for Medicare & Medicaid Services. A Field Guide to Antibiotic Stewardship in Outpatient Settings A Field Guide to Antibiotic Stewardship in Outpatient Settings July 2018 2 A Field Guide to Antibiotic Stewardship in Outpatient Settings. 2018.

6.   Shehab N, Lovegrove MC, Geller AI, Rose KO, Weidle NJ, Budnitz DS. US Emergency Department Visits for Outpatient Adverse Drug Events, 2013-2014. JAMA. 2016 Nov 22;316(20):2115.

7.   Dantes R, Mu Y, Hicks LA, Cohen J, Bamberg W, Beldavs ZG, et al. Association between Outpatient Antibiotic Prescribing Practices and Community-Associated Clostridium difficile Infection. Open Forum Infect Dis. 2015 Aug 11;ofv113.

8.   Sanchez G V., Fleming-Dutra KE, Roberts RM, Hicks LA. Core Elements of Outpatient Antibiotic Stewardship. MMWR Recomm Rep. 2016 Nov 11;65(6):1–12.

9.   Dellit TH, Owens RC, McGowan JE, Gerding DN, Weinstein RA, Burke JP, et al. Infectious Diseases Society of America and the Society for Healthcare Epidemiology of America guidelines for developing an institutional program to enhance antimicrobial stewardship. Clin Infect Dis Off Publ Infect Dis Soc Am. 2007 Jan 15;44(2):159–77.

10. Vellinga A, Murphy AW, Hanahoe B, Bennett K, Cormican M. A multilevel analysis of trimethoprim and ciprofloxacin prescribing and resistance of uropathogenic Escherichia coli in general practice. J Antimicrob Chemother. 2010 Jul;65(7):1514–20.

11. Livermore DM. The 2018 Garrod Lecture: Preparing for the Black Swans of resistance. J Antimicrob Chemother. 2018 Nov 1;73(11):2907–15.

12. Turnidge J, Christiansen K. Antibiotic use and resistance--proving the obvious. Lancet Lond Engl. 2005 Feb;365(9459):548–9.

365   13.   Arason VA, Gunnlaugsson A, Sigurdsson JA, Erlendsdottir H, Gudmundsson S,
366   Kristinsson KG. Clonal Spread of Resistant Pneumococci Despite Diminished
367   Antimicrobial Use. Microb Drug Resist. 2002 Sep;8(3):187–92.

368   14.   Lipsitch M. The rise and fall of antimicrobial resistance. Trends Microbiol. 2001
369   Sep;9(9):438–44.

370   15.   Costelloe C, Metcalfe C, Lovering A, Mant D, Hay AD. Effect of antibiotic
371   prescribing in primary care on antimicrobial resistance in individual patients:
372   systematic review and meta-analysis. BMJ. 2010 May 18;340:c2096.

373   16.   Harbarth S, Harris AD, Carmeli Y, Samore MH. Parallel analysis of individual and
374   aggregated data on antibiotic exposure and resistance in gram-begative bacilli. Clin
375   Infect Dis. 2001;33:1462–8.

376   17.   Lipsitch M. Measuring and Interpreting Associations between Antibiotic Use and
377   Penicillin Resistance in Streptococcus pneumoniae. Clin Infect Dis. 2001 Apr
378   1;32(7):1044–54.

379   18.   Hannah EL, Angulo FJ, Johnson JR, Haddadin B, Williamson J, Samore MH. Drug-
380   resistant Escherichia coli, Rural Idaho. Emerg Infect Dis. 2005 Oct;11(10):1614–7.

381   19.   Kalter HD, Gilman RH, Moulton LH, Cullotta AR, Cabrera L, Velapatiño B. Risk
382   factors for antibiotic-resistant Escherichia coli carriage in young children in Peru:
383   community-based cross-sectional prevalence study. Am J Trop Med Hyg. 2010
384   May;82(5):879–88.

385   20.   Samore MH, Magill MK, Alder SC, Severina E, Morrison-De Boer L, Lyon JL, et al.
386   High rates of multiple antibiotic resistance in Streptococcus pneumoniae from
387   healthy children living in isolated rural communities: association with cephalosporin
388   use and intrafamilial transmission. Pediatrics. 2001 Oct;108(4):856–65.

389   21.   Lipsitch M, Samore MH. Antimicrobial use and antimicrobial resistance: a
390   population perspective. Emerg Infect Dis. 2002;8:347–54.

391   22.   Cooper BS, Medley GF, Stone SP, Kibbler CC, Cookson BD, Roberts JA, et al.
392   Methicillin-resistant Staphylococcus aureus in hospitals and the community: Stealth
393   dynamics and control catastrophes. Proc Natl Acad Sci. 2004 Jul 6;101(27):10223–
394   8.

395   23.   MacFadden DR, Fishman DN, Hanage WP, Lipsitch M. The Relative Impact of
396   Community and Hospital Antibiotic Use on the Selection of Extended-Spectrum
397   Beta-lactamase-Producing Escherichia coli. Clin Infect Dis [Internet]. 2018 [cited
398   2018 Nov 29]; Available from: https://www.ncbi.nlm.nih.gov/pubmed/30462185

399   24.   Knight GM, Costelloe C, Deeny SR, Moore LSP, Hopkins S, Johnson AP, et al.
400   Quantifying where human acquisition of antibiotic resistance occurs: a
401   mathematical modelling study. BMC Med. 2018 23;16(1):137.

25. Seppälä H, Klaukka T, Vuopio-Varkila J, Muotiala A, Helenius H, Lager K, et al.
The Effect of Changes in the Consumption of Macrolide Antibiotics on
Erythromycin Resistance in Group A Streptococci in Finland. N Engl J Med. 1997
Aug 14;337(7):441–6.

26. Kristinsson KG. Effect of antimicrobial use and other risk factors on antimicrobial
resistance in pneumococci. Microb Drug Resist Larchmt N. 1997;3(2):117–23.

27. Hennessy TW, Petersen KM, Bruden D, Parkinson AJ, Hurlburt D, Getty M, et al.
Changes in Antibiotic-Prescribing Practices and Carriage of Penicillin-Resistant
Streptococcus pneumoniae: A Controlled Intervention Trial in Rural Alaska. Clin
Infect Dis. 2002 Jun 15;34(12):1543–50.

28. Schechner V, Temkin E, Harbarth S, Carmeli Y, Schwaber MJ. Epidemiological
interpretation of studies examining the effect of antibiotic usage on resistance. Clin
Microbiol Rev. 2013 Apr;26(2):289–307.

29. O'Brien KS, Emerson P, Hooper P, Reingold AL, Dennis EG, Keenan JD, et al.
Antimicrobial resistance following mass azithromycin distribution for trachoma: a
systematic review. Lancet Infect Dis [Internet]. 2018 Oct 3 [cited 2018 Nov 7];
Available from:
http://www.sciencedirect.com/science/article/pii/S1473309918304444

30. Bell BG, Schellevis F, Stobberingh E, Goossens H, Pringle M. A systematic review
and meta-analysis of the effects of antibiotic consumption on antibiotic resistance.
BMC Infect Dis. 2014 Jan 9;14:13.

31. MacFadden DR, McGough SF, Fisman D, Santillana M, Brownstein JS. Antibiotic
resistance increases with local temperature. Nat Clim Change. 2018 Jun;8(6):510–
4.

32. Olesen SW, Barnett ML, MacFadden DR, Brownstein JS, Hernández-Diaz S,
Lipsitch M, et al. The distribution of antibiotic use and its association with antibiotic
resistance. eLife [Internet]. 2018 Dec 18 [cited 2018 Dec 19];7. Available from:
https://elifesciences.org/articles/39435

33. Goossens H, Ferech M, Vander Stichele R, Elseviers M. Outpatient antibiotic use
in Europe and association with resistance: a cross-national database study. The
Lancet. 2005 Feb 12;365(9459):579–87.

34. van de Sande-Bruinsma N, Grundmann H, Verloo D, Tiemersma E, Monen J,
Goossens H, et al. Antimicrobial drug use and resistance in Europe. Emerg Infect
Dis. 2008 Nov;14(11):1722–30.

35. Garcia-Rey C, Aguilar L, Baquero F, Casal J, Dal-Re R. Importance of local
variations in antibiotic consumption and geographical differences of erythromycin
and penicillin resistance in Streptococcus pneumoniae. J Clin Microbiol. 2002
Jan;40:159–64.

36.   Bergman M, Nyberg ST, Huovinen P, Paakkari P, Hakanen AJ, and the Finnish
Study Group for Antimicrobial Resistance. Association between Antimicrobial
Consumption and Resistance in Escherichia coli. Antimicrob Agents Chemother.
2009 Mar 1;53(3):912–7.

37.   MacDougall C, Powell JP, Johnson CK, Edmond MB, Polk RE. Hospital and
community fluoroquinolone use and resistance in Staphylococcus aureus and
Escherichia coli in 17 US hospitals. Clin Infect Dis. 2005 Aug 15;41:435–40.

38.   Vellinga A, Tansey S, Hanahoe B, Bennett K, Murphy AW, Cormican M.
Trimethoprim and ciprofloxacin resistance and prescribing in urinary tract infection
associated with Escherichia coli: a multilevel model. J Antimicrob Chemother. 2012
Oct 1;67(10):2523–30.

39.   Donnan PT, Wei L, Steinke DT, Phillips G, Clarke R, Noone A, et al. Presence of
bacteriuria caused by trimethoprim resistant bacteria in patients prescribed
antibiotics: multilevel model with practice and individual patient data. BMJ. 2004
May 29;328(7451):1297.

40.   Priest P, Wise R, Yudkin P, McNulty C, Mant D. Antibacterial prescribing and
antibacterial resistance in English general practice: cross sectional study. BMJ.
2001 Nov 3;323(7320):1037–41.

41.   García-Rey C, Martín-Herrero JE, Baquero F. Antibiotic consumption and
generation of resistance in Streptococcus pneumoniae: the paradoxical impact of
quinolones in a complex selective landscape. Clin Microbiol Infect Off Publ Eur Soc
Clin Microbiol Infect Dis. 2006 May;12 Suppl 3:55–66.

42.   Openshaw S. The modifiable areal unit problem. Geo Books; 1984. (Concepts and
Techniques in Modern Geography).

43.   Vellinga A, Bennett K, Murphy AW, Cormican M. Principles of multilevel analysis
and its relevance to studies of antimicrobial resistance. J Antimicrob Chemother.
2012 Oct;67(10):2316–22.

44.   Davies NG, Flasche S, Jit M, Atkins KE. Within-host dynamics explain patterns of
antibiotic resistance in commensal bacteria. bioRxiv. 2018 Jan 3;217232.

45.   Lehtinen S, Blanquart F, Croucher NJ, Turner P, Lipsitch M, Fraser C. Evolution of
antibiotic resistance is linked to any genetic mechanism affecting bacterial duration
of carriage. Proc Natl Acad Sci. 2017 Jan 31;114(5):1075–80.

46.   Blanquart F, Lehtinen S, Lipsitch M, Fraser C. The evolution of antibiotic resistance
in a structured host population. J R Soc Interface. 2018 Jun;15(143).

47.   Truven Health MarketScan Database. Commercial Claims and Encounters. Ann
Arbor, MI; 2015.

48.   MacFadden DR, Fisman D, Andre J, Ara Y, Majumder MS, Bogoch II, et al. A
477   Platform for Monitoring Regional Antimicrobial Resistance, Using Online Data
478   Sources: ResistanceOpen. J Infect Dis. 2016 Dec 1;214(suppl_4):S393–8.

479   49.   US Centers for Disease Control and Prevention. Patient Safety Atlas - Outpatient
480   Antibiotic Use [Internet]. [cited 2018 Oct 25]. Available from:
481   https://gis.cdc.gov/grasp/PSA/indexAU.html

482   50.   US Centers for Disease Control and Prevention. Patient Safety Atlas - Antibiotic
483   Resistance [Internet]. [cited 2018 Oct 25]. Available from:
484   https://gis.cdc.gov/grasp/PSA/AboutTheData.html

485   51.   European Centre for Disease Prevention and Control. Antimicrobial consumption
486   database [Internet]. European Centre for Disease Prevention and Control. [cited
487   2018 Oct 25]. Available from: http://ecdc.europa.eu/en/antimicrobial-
488   consumption/surveillance-and-disease-data/database

489   52.   European Centre for Disease Prevention and Control. Surveillance Atlas of
490   Infectious Diseases [Internet]. [cited 2018 Oct 25]. Available from:
491   https://atlas.ecdc.europa.eu/public/index.aspx

492   53.   United Nations Statistics Division. Standard country or area codes or statistical use
493   [Internet]. New York: United Nations; 1999. Report No.: M No. 49. Available from:
494   https://unstats.un.org/unsd/methodology/m49/

495   54.   Venables WN, Ripley BD. Modern applied statistics with S [Internet]. 4th ed. New
496   York: Springer; 2002. Available from: www.stats.ox.ac.uk/pub/MASS4

497   55.   R Core Team. R: A language and environment for statistical computing [Internet].
498   Vienna; 2018. Available from: www.R-project.org

499   56.   Collignon P, Beggs JJ, Walsh TR, Gandra S, Laxminarayan R. Anthropological and
500   socioeconomic factors contributing to global antimicrobial resistance: a univariate
501   and multivariable analysis. Lancet Planet Health. 2018 Sep;2(9):e398–405.

502   57.   Sundqvist M, Geli P, Andersson DI, Sjölund-Karlsson M, Runehagen A, Cars H, et
503   al. Little evidence for reversibility of trimethoprim resistance after a drastic
504   reduction in trimethoprim use. J Antimicrob Chemother. 2010 Feb;65(2):350–60.

505   58.   Enne VI, Livermore DM, Stephens P, Hall LM. Persistence of sulphonamide
506   resistance in Escherichia coli in the UK despite national prescribing restriction.
507   Lancet Lond Engl. 2001 Apr 28;357(9265):1325–8.

508   59.   Pouwels KB, Freeman R, Muller-Pebody B, Rooney G, Henderson KL, Robotham
509   JV, et al. Association between use of different antibiotics and
510   trimethoprimresistance: going beyond the obvious crude association. J Antimicrob
511   Chemoth. 2018;

60. Tedijanto C, Olesen SW, Grad YH, Lipsitch M. Estimating the proportion of
bystander selection for antibiotic resistance among potentially pathogenic bacterial
flora. Proc Natl Acad Sci. 2018 Dec 18;115(51):E11988–95.

61. Olesen SW, Torrone EA, Papp JR, Kirkcaldy RD, Lipsitch M, Grad YH.
Azithromycin Susceptibility Among Neisseria gonorrhoeae Isolates and Seasonal
Macrolide Use. J Infect Dis [Internet]. 2018 [cited 2018 Nov 6]; Available from:
http://academic.oup.com/jid/advance-article/doi/10.1093/infdis/jiy551/5098400

62. Dagan R, Barkai G, Givon-Lavi N, Sharf AZ, Vardy D, Cohen T, et al. Seasonality
of Antibiotic-Resistant Streptococcus pneumoniae That Causes Acute Otitis Media:
A Clue for an Antibiotic-Restriction Policy? J Infect Dis. 2008 Apr 15;197(8):1094–
102.

63. Sun L, Klein EY, Laxminarayan R. Seasonality and Temporal Correlation between
Community Antibiotic Use and Resistance in the United States. Clin Infect Dis.
2012 Sep 1;55(5):687–94.

64. Blanquart F, Lehtinen S, Fraser C. An evolutionary model to predict the frequency
of antibiotic resistance under seasonal antibiotic use, and an application to
Streptococcus pneumoniae. Proc R Soc B. 2017 May 31;284(1855):20170679.

65. Dingle KE, Didelot X, Quan TP, Eyre DW, Stoesser N, Golubchik T, et al. Effects of
control interventions on Clostridium difficile infection in England: an observational
study. Lancet Infect Dis. 2017 Apr;17(4):411–21.

66. McCormick AW, Whitney CG, Farley MM, Lynfield R, Harrison LH, Bennett NM, et
al. Geographic diversity and temporal trends of antimicrobial resistance in
Streptococcus pneumoniae in the United States. Nat Med. 2003 Apr;9(4):424–30.

67. Mossong J, Hens N, Jit M, Beutels P, Auranen K, Mikolajczyk R, et al. Social
contacts and mixing patterns relevant to the spread of infectious diseases. PLoS
Med. 2008 Mar 25;5(3):e74.

68. Garett G, Hughes J, Anderson R, Stoner B, Aral S, Whittington W, et al. Sexual
mixing patterns of patients attending sexually transmitted diseases clinics. Sex
Transm Dis. 1996;23(3):248–57.

69. Newman MEJ. Mixing patterns in networks. Phys Rev E Stat Nonlin Soft Matter
Phys. 2003 Feb;67(2 Pt 2):026126.

70. Hicks LA, Bartoces MG, Roberts RM, Suda KJ, Hunkler RJ, Taylor TH, et al. US
Outpatient Antibiotic Prescribing Variation According to Geography, Patient
Population, and Provider Specialty in 2011. Clin Infect Dis. 2015 May
1;60(9):1308–16.

71. Olesen SW, Grad YH. Racial/Ethnic Disparities in Antimicrobial Drug Use, United
States, 2014–2015. Emerg Infect Dis [Internet]. 2018 [cited 2018 Nov 6];24(11).
Available from: https://wwwnc.cdc.gov/eid/article/24/11/18-0762_article

72. Charu V, Zeger S, Gog J, Bjørnstad ON, Kissler S, Simonsen L, et al. Human
mobility and the spatial transmission of influenza in the United States. PLoS
Comput Biol. 2017;13(2):e1005382.

Disclaimers

The views and opinions of the authors expressed herein do not necessarily state or

reflect those of the ECDC. The accuracy of the authors' statistical analysis and the

findings they report are not the responsibility of ECDC. ECDC is not responsible for

conclusions or opinions drawn from the data provided. ECDC is not responsible for the

correctness of the data and for data management, data merging and data collation after

provision of the data. ECDC shall not be held liable for improper or incorrect use of the

data.

Funding

This work was supported by the National Institutes of Health (grant number

U54GM088558 to ML). The funders had no role in study design, data collection and

interpretation, or the decision to submit the work for publication.

Acknowledgements

We thank Dr. Stephen M. Kissler for helpful comments on the manuscript.

Figures and figure legends

Figure 1. **Interactions between populations attenuate the effect of interventions.**

(*a*) Schematic of the 2-population WHN model. (*b*) Results of simulations of the 2-

population WHN model for a modest intervention (difference in antibiotic use between

populations $\Delta\tau$ = 0.05 monthly treatments per capita). As interaction strength (ε,

horizontal axis) increases, the difference in antibiotic resistance between the two

populations decreases. (*c*) The same pattern holds for a stronger intervention ($\Delta\tau$ =

0.15). Compare Supplemental Figure 1, which shows the analogous results for the D-

types model.

Figure 2. **Theoretical use-resistance associations with regional population**

structure. (*a*) Results of simulations of nested population simulations using the WHN

model for 3 parameter sets (panel columns). Populations (circles) in the same super-

population (color) interact more strongly ($\varepsilon_{\mathrm{sub}}$) with populations in the same super-

population than with other populations ($\varepsilon_{\mathrm{super}} \leq \varepsilon_{\mathrm{sub}}$). Lines show linear best fit within

each super-population. (*b*) Points show the populations in panel *a* but aggregated into

super-populations. Each super-population's use and resistance is the mean of its

constituent populations' values. Lines show linear best fit across super-populations. (*c*)

Each point represents a pair of populations from panel *a*. Points' positions represent the

differences in antibiotic use and resistance between the populations in the pair. Colors

indicate whether the two populations are in the same super-population. Lines show best

fit among the same-super-population and different-super-population pairs. Compare

Supplemental Figure 2, which shows the analogous results for the D-types model.

Figure 3. **Use-resistance associations when regionally aggregated.** Panels show

use-resistance relationships for 3 pathogen-antibiotic combinations in the

MarketScan/ResistanceOpen dataset. Points represent geographic units of analysis at

different aggregation levels (black, US states; green, US Census divisions; orange, US

Census regions). Curves show logistic regression fits. Shaded regions show 95%

bootstrap confidence intervals. Compare Figure 2b, which shows that theoretical

models predict the same use-resistance associations across aggregated and

unaggregated data. Compare also Supplemental Figure 3, which shows analogous

results using the other datasets.

Figure 4. **Use-resistance relationships by adjacency.** Each point represents a pair of

US states. The point's position represents the difference in use of macrolides between

the two states (horizontal axis) and the difference in macrolide resistance among *S.*

pneumoniae between the states (log odds ratio) using the MarketScan/ResistanceOpen

data, shown in one of the panels of Figure 3. The point's color indicates whether the

states are physically adjacent (red = adjacent, black = not adjacent). Lines show

predictions from robust linear regressions on the adjacent and non-adjacent pairs, using

the indicated difference in antibiotic use and mean values for the other model

predictors. Shaded areas indicate regressions' 95% bootstrap confidence intervals.

Compare Figure 2c, which shows that adjacency effects can be detected in theoretical

models. Compare also Supplemental Figure 5, which shows analogous results for other

pathogen-antibiotic combinations and other datasets.