Horizontally transmitted symbiont populations in deep-sea mussels are genetically isolated

Devani Romero Picazo[1*], Tal Dagan[1], Rebecca Ansorge[2], Jillian M. Petersen[3], Nicole Dubilier[2], Anne Kupczok[1*]

[1] Genomic Microbiology Group, Institute of General Microbiology, Christian-Albrechts University, Kiel
[2] Max Planck Institute for Marine Microbiology, Bremen
[3] Division of Microbiology and Ecosystem Science, University of Vienna
* Corresponding authors: dpicazo@ifam.uni-kiel.de, akupczok@ifam.uni-kiel.de

Abstract

Eukaryotes are habitats for bacterial organisms where the host colonization and dispersal among individual hosts have consequences for the bacterial ecology and evolution. Vertical symbiont transmission leads to geographic isolation of the microbial population and consequently to genetic isolation of microbiotas from individual hosts. In contrast, the extent of geographic and genetic isolation of horizontally transmitted microbiota is poorly characterized. Here we show that chemosynthetic symbionts of individual *Bathymodiolus brooksi* mussels constitute genetically isolated populations. The reconstruction of core genome-wide strain sequences from high-resolution metagenomes revealed distinct phylogenetic clades. Nucleotide diversity and strain composition vary along the mussel lifespan and individual hosts show a high degree of genetic isolation. Our results suggest that the uptake of environmental bacteria is a restricted process in *B. brooksi*, where self-infection of the gill tissue results in serial founder effects during symbiont evolution. We conclude that bacterial colonization dynamics over the host life-cycle is thus an important determinant of population structure and genome evolution of horizontally transmitted symbionts.

Bacteria inhabit most eukaryotes where their presence has consequences for key aspects of

their host biology[1], such as host development[2], nutrition[3], or behavior[4]. From the bacterial

perspective, animals constitute an ecological niche, where microbial communities utilize the

resources of their host habitat[5]. The microbiota biodiversity over the host life cycle is determined

by bacteria colonization dynamics and by host properties, including biotic and abiotic factors.

For example, the microbiota can be affected by the host diet[6] or the host physiological state

(e.g., hibernation[7] or pregnancy[8]). In addition, changes in the host environmental conditions

such as temperature[9] or the availability of reduced compounds[10] can have an effect on the

microbiota community composition.

Microbiota dispersal over the host life cycle depends on the level of fidelity between the

host and its microbiota; in faithful interactions, vertically transmitted bacteria are transferred

from adults to their progeny during early host developmental stages, while in less faithful

interactions, horizontally transmitted bacteria are acquired from the environment throughout the

host life cycle[11]. Strictly vertically transmitted bacteria are specialized in their host niche and

their association with the host imposes an extreme geographic isolation. Bacterial inheritance

over host generations imposes a strong bottleneck on the microbiota population and leads to

reduced intra-host genetic diversity[12]. Examples are monoclonal or biclonal populations

observed in symbiotic bacteria inhabiting grass sharpshooter[13] and pea aphids[14]. Furthermore,

the geographic isolation of vertically transmitted bacteria leads to genetic isolation and to

symbiont genome reduction over time as a consequence of genetic drift[15]. In contrast, dispersal

is expected to be higher for horizontally transmitted bacteria, where host-associated populations

are connected to one another through the environmental pool[16]. Nonetheless, the genetic

diversity of horizontally transmitted microbial populations may also be reduced due to

bottlenecks during symbiont transmission and host colonization. Stochastic effects in the

colonization of horizontally transmitted bacteria may manifest themselves in differences in

microbiota strain composition among hosts[17,18]. This would lead to subdivided symbiont

populations where the geographic isolation of the microbiota depends on the degree of

symbiont dispersal among individual hosts. Geographic isolation of individual hosts over the

host life span would then lead to genetic isolation of the symbiont populations and to symbiont

population structure. Genomic variation and genetic isolation have been observed for

horizontally transmitted symbionts of the human gut microbiome[19] and of the honey bee gut

microbiome[20]. Moreover, structured symbiont populations can also emerge within an individual

host, as observed for *Vibrio fischeri* colonizing the squid light organ, where different light organ

crypts are infected by a specific strain[21]. The degree of dispersal of horizontally transmitted

symbionts remains understudied; hence, whether populations from different microbiomes are

intermixing or are genetically isolated is generally unknown.

Here we study the microbiota strain composition of horizontally transmitted

endosymbionts across individual *Bathymodiolus brooksi* deep-sea mussels. *Bathymodiolus*

mussels live in a nutritional symbiosis with the chemosynthetic sulfur-oxidizing (SOX) and

methane-oxidizing (MOX) bacteria. The symbionts are acquired horizontally from the seawater

and are harbored in bacteriocytes within the gill epithelium[22,23]. Most Bathymodiolus species

harbor only a single 16S rRNA phylotype for each symbiont, including *B. brooksi*[24].

Nevertheless, a recent metagenomic analysis of *Bathymodiolus* species from hydrothermal

vents in the mid Atlantic ridge showed the presence of different SOX strains with differing

metabolic capacity[25]. Mussel gills constantly develop new filaments that are continuously

infected[26]. However, whether the new gill filaments in *Bathymodiolus brooksi* are colonized

predominantly by environmental bacteria or by symbionts from older filaments of the same host

remains unknown. These two alternative scenarios are expected to impose different degrees of

geographic isolation on the symbiont population: in continuous environmental acquisition, the

level of inter-host dispersal is high while self-infection limits the symbiont dispersal. Here we

studied the impact of tissue colonization dynamics of horizontally transmitted intracellular

symbionts on the degree of symbiont diversity. Furthermore, we quantified the level of genetic

isolation among communities across individual mussels and its impact on symbiont genome

evolution. For that, we implemented a high-resolution metagenomics approach that captures

genome-wide diversity for both symbionts in multiple *Bathymodiolus brooksi* individuals from a

single site.

Results

Gene-based metagenomics binning recovers SOX and MOX core genomes

To study the evolution of the SOX and MOX genomes in *Bathymodiolus* mussels we used a

high-resolution metagenomics approach. Twenty-three *B. brooksi* individuals of shell sizes

ranging between 4.8 cm and 24.3 cm were sampled from a single location at a cold seep site in

the northern Gulf of Mexico. Shell size correlates with mussel age[27]; thus, analyzing mussels

within a wide shell size range allowed us to study the symbiont population structure across host

ages. The mussels were sampled from three separate mussel 'clumps' (small mussel patches

residing on the sediment) that were at most 131m apart (Supplementary Fig. 1). Such a 'patchy'

distribution has often been observed in deep-sea mussels[28]. To obtain a comprehensive

representation of the bacterial population in individual mussels and to accurately infer strain-

specific genomes, homogenized gill tissue of each mussel was deeply sequenced (on average,

37.8 million paired-end reads of 250bp per sample, Supplementary Table 1). The resulting

metagenomic sequencing data was analyzed by a gene-based binning approach[29].

The prediction of protein-coding genes from the assembled metagenomes yielded a

non-redundant gene catalog of 4.4 million genes that potentially contains every gene present in

the samples. This includes genes from the microbial community and from the mussel host. In

the metagenomics binning step, genes that covary in their abundance across the different

samples were clustered into metagenomic species (MGSs). Our analysis revealed two MGSs

that comprise the SOX and MOX core genomes (Supplementary Fig. 2). The distribution of

gene coverage in individual samples shows that genes in each core genome have a similar

abundance within each mussel. This confirms the classification of the SOX and MOX MGSs as

core genomes. The MOX core genome is the largest MGS and it contains 2,518 genes with a

total length of 1.97 Mbp. A comparison to Gammaproteobacteria marker genes shows that it is

96.2% complete. Furthermore, it contains 1,568 genes (62.3%) that have homologs in MOX-

related genomes. The SOX core genome contains 1,439 genes, has a total length of 1.27 Mbp

and is considered as 80.2% complete. It contains 1,188 genes (82.6%) with homologs in SOX-

related genomes. In addition to the SOX and MOX core genomes, our analysis revealed a third

MGS of 1,449 genes (Supplementary Fig. 2) that was found in low abundance in a single

mussel and, in addition, 98,944 co-abundant gene groups (CAGs, 3-699 genes). Of the 23

metagenomes, four samples were discarded during the metagenomics binning. Two samples

were discarded prior to the binning due to high variance in symbiont marker gene coverages

and two samples were discarded after binning due to low coverage for both symbionts

(Supplementary Figs. 2,3). To gain insight into the SOX and MOX population structure between

hosts, we compared the characteristics of the core genomes across the remaining 19 samples.

The analysis of the core genome coverages shows that SOX is the dominant member of the

mussel microbiota. The differences in the SOX to MOX ratio among the mussel metagenomes

are likely explained by differences in the availability of H_2S and CH_4 among clumps, which is a

known determinant of SOX and MOX abundance in *Bathymodiolus*[30] (Supplementary

Information, Supplementary Fig. 4).

To study symbiont diversity below the species level, we analyzed single nucleotide

variants (SNVs) that were detected in the core genomes of the two symbionts. In this analysis,

we considered SNVs that are fixed in a metagenome as well as polymorphic SNVs, i.e., SNVs,
where both the reference and the alternative allele are observed in a single metagenome. We
found 18,070 SNVs in SOX (SNV density of 14 SNVs/kbp, 49 multi-state, 0.27%) and 4,652
SNVs in MOX (SNV density of 2.4 SNVs/kbp, 5 multi-state, 0.11%). The number of polymorphic
SNVs per sample ranges from 162 (0.9%) to 11,064 (61%) for SOX and from 27 (0.58%) to
3,026 (65%) for MOX (Supplementary Table 1), thus, most SNVs are polymorphic in at least
one sample. It is important to note that the observed difference in strain-level diversity between
SOX and MOX cannot be explained by the difference in sequencing depth (Supplementary
Information). These results are in agreement with previous reports of SOX genetic diversity in
other *Bathymodiolus* species[25]. We further revealed that there is genetic diversity in the MOX
symbiont.

Bathymodiolus microbiota is composed of SOX and MOX strains from several clades.

Diversity in natural populations of bacteria is characterized by cohesive associations among
genetic loci that contribute to lineage formation and generate distinguishable genetic clusters
beyond the species level[31]. The formation of niche-specific genotypes (i.e., ecotypes) has been
mainly studied in populations of free-living organisms such as the cyanobacterium
Prochlorococcus spp.[32]. Here we consider a strain to be a genetic entity that is present in
multiple hosts and is characterized by a set of clustered variants in the core genome. To study
lineage formation in symbiont populations associated with *Bathymodiolus* mussels, we
reconstructed the strain consensus core genomes from strain-specific variants that show similar
frequencies in a metagenomic sample.

The SNVs found in multiple samples and their covariation across samples were used for
strain deconvolution of the core genomes using DESMAN[33]. This revealed that SOX is
composed of eleven different strains with a mean strain core genome sequence identity of
99.52%. Phylogenetic reconstruction shows that the eleven strains cluster into four clades,
which are separated by relatively long internal branches (Fig. 1b). Notably, 849 of the SNVs on
the SOX core genome (4.7%) do not differentiate between strains. Thus, the resulting strain
alignment is invariant for each of these positions and they are termed invariant SNVs from here
on. For MOX, six strains with a mean core genome sequence identity of 99.88% were
reconstructed. The phylogenetic network shows that the six strains cluster into two clades
comprising three strains each (Fig. 1e). Of the total SNVs, 1,138 (24.4%) are invariant in the
strain alignment. The overall MOX branch lengths are shorter than those of SOX. We detected

no effect of sequencing coverage on the inference of the strain clades for SOX (Supplementary
Information, Supplementary Fig. 5).

To study the community assembly at the strain level, we examined the strain distribution
across individual mussels. Each SOX strain could be identified in between three and eight
samples (frequency ≥5%; Fig. 1a). Only one or two strains were detected with a frequency of at
least 5% in small mussels (≤7 cm), two to nine strains in medium-sized mussels (7.2 cm – 14.1
cm) and one to two strains in large mussels (14.6 cm – 24.1 cm). Notably, only strains from
clades S1 and S2 are present in large mussels (≥14.6 cm). One of the large mussels (S) is an
exception as it hosts three SOX strains and contains strains from both clades S1 and S2. Six
mussels have one dominant SOX strain (frequency ≥90%). Five of these are large mussels (M,
N, P, Q, R) and only one is a small mussel (C). The dominant strain is either S1.4, S2.1, or S2.2
(Fig. 1a; Supplementary Table 1). The MOX strain composition across mussels shows that each
MOX strain occurs (frequency ≥5%; Fig. 1d) in four to 17 mussels and each mussel contains
two to four MOX strains. Additionally, strains of clade M2 are dominant in ten of the mussels.

To investigate the degree of genetic cohesion within strain clades in the population, we
studied the allele frequency spectrum (AFS) of each mussel. A visual inspection of the derived
allele frequency spectra revealed multimodal distributions for both symbiont populations. The
modes reach high allele frequencies and are associated with the main phylogenetic clades; this
suggests that the clades constitute cohesive genetic units (Fig. 1c,f; Supplementary Fig. 6). The
presence of high-frequency modes is especially apparent for SOX in medium-sized mussels
that contain multiple strains. To identify sample-specific strain sequences, we reconstructed
dominant haplotypes (major allele frequency ≥90%) for the samples that contain a dominant
strain (strain frequency ≥90%). By comparing dominant haplotypes among samples containing
the same dominant strain, we found that these can contain between 42 and 74 differential SNVs
(Supplementary Table 1). This suggests that the fixation of variants within individual mussels
contributes to the observed population structure.

Overall, our results revealed that the symbiont populations are composed of strains that
cluster into few clades, which appear to be maintained by strong cohesive forces. In addition,
the strains are shared among multiple mussels and multiple strains are capable of dominating
different hosts. This suggests that stochastic processes are governing the symbiont community
assembly, as previously proposed for other *Bathymodiolus* species[34].

SOX strains evolve under purifying selection while MOX evolution is characterized by neutral processes

To study the evolution of SOX and MOX strains in *Bathymodiolus*, we examined the selection regimes that have been involved in the formation of cohesive genetic SOX and MOX units. The core genome-wide ratio of pN/pS is higher in MOX (pN/pS of 0.425) in comparison to SOX (pN/pS of 0.137), which indicates that the strength of purifying selection is higher for SOX. In addition, we estimated pN/pS for each of the symbiont core genes. This revealed that MOX genes are characterized by large pN/pS and small pS values, while SOX genes have small pN/pS and large pS values (Supplementary Fig. 7). The relative rate of nonsynonymous to synonymous substitutions has been shown to depend on the divergence of the analyzed species[35,36]. For populations of low divergence, SNVs comprise substitutions that have been fixed in the population and mutations that arose recently. The latter include slightly deleterious mutations that were not yet purged by selection, resulting in an elevated ratio of nonsynonymous to synonymous replacements. Thus, this ratio is not suitable for analyzing closely related genomes, which is usually the case when studying variation within bacterial species.

To circumvent the bias in pN/pS, we tested for differences in selection regimes in the evolution of SOX and MOX strains using the neutrality index (NI). NI is used to distinguish between divergent and polymorphic SNVs and to quantify the departure of a population from the neutral expectation. An excess of divergent nonsynonymous mutations (NI<1) indicates that the population underwent positive selection or an important demographic change in the past[37]. We estimated NI by considering two different levels of divergence and polymorphism. In the first level, all identified strains are considered as diverged taxonomic units; in the second level, we disregard the small-scale strain classification and consider only the clades as diverged taxonomic units (Table 1). Considering all strains as divergent, we observed a low NI^{MOX} (<1), which suggests that MOX evolved under a neutral (NI~1) or positive selection regime. NI^{MOX} increased when considering the clades as diverged, which suggests that the low NI^{MOX} observed at the strain level is the result of an excess of nonsynonymous SNVs within the strain clades that may constitute transient polymorphisms. Thus, the excess of nonsynonymous mutations observed for MOX is biased by the low level of divergence; hence, similar to the pN/pS ratio, it cannot serve as an indication for positive selection. On the other hand, we found that purifying selection is in action for SOX (NI^{SOX}>1). Similar to MOX, when using the clades as

divergent, NI^{SOX} slightly increases. This indicates that the SNVs that differ between clades are
more likely to be substitutions in comparison to those that differ among within-clade strains.

Altogether, these results suggest differences in the selection regimes during the
evolution of the SOX and MOX strains. While the SOX core genome is shaped by purifying
selection, we cannot detect deviation from the neutral expectation in the MOX core genome.
These differences likely stem from the different divergence levels among the strains of both
symbiont populations. The association of SOX with *Bathymodiolus* mussels is considered to be
ancient in chemosynthetic deep-sea mussels whereas the MOX association is thought to have
evolved secondarily during *Bathymodiolus* diversification[38]. This is in agreement with the larger
degree of divergence observed here for SOX. Since we observed no evidence for positive
selection on the symbiont core genomes, we suggest that the strains constitute cohesive
genetic units within one ecotype[39], where all strains are functionally equivalent at the core
genome level. Notwithstanding, the strains might be linked to differences in the accessory gene
content, as observed, for example, in the free-living cyanobacterium *Prochlorococcus* spp.[32] and
in SOX symbionts of other *Bathymodiolus* species[25].

Intra-sample diversity is higher for SOX than for MOX.

The association with the host limits the dispersal of bacterial populations where the association
across generations is likely maintained by symbiont dispersal between host individuals. If
symbionts are not continuously taken up from the environment, each individual host constitutes
an isolated habitat over its lifetime[5]. Geographic isolation between habitats results in genetic
isolation and contributes to the formation of cohesive associations of genetic loci[31]. Previous
studies showed that geographic isolation during vertical transmission can lead to the reduction
of intra-host genetic diversity in the bacterial populations[12], nonetheless, the degree of isolation
remains understudied for horizontally transmitted microbes. To characterize the contribution of
geographic isolation to strain formation in the *Bathymodiolous* symbiosis, we next studied the
degree of genetic isolation. Our sample collection of mussels covering a range of sizes (and
thus ages) enabled us to compare symbiont genome diversity among individual hosts of
different age within a single sampling site, thus minimizing the putative effect of biogeography
on population structure. The host species *B. brooksi* is ideal for such an analysis as it grows to
unusally large sizes and possibly lives longer than many other *Bathymodiolus* species. To study
differences in genome diversity of the two symbionts across individual mussels, we estimated

the intra-sample nucleotide diversity (π) and the ecological measure α-diversity at the resolution
of the SOX and MOX strains.

We found a high variability of π^{SOX} among different mussels (intra-sample π^{SOX} between
5.2×10^{-5} and 3.6×10^{-3}, Table 2, Fig. 2). Furthermore, π^{SOX} and the SOX α-diversity are
significantly positively correlated ($\rho^2 = 0.98$, $p < 10^{-6}$, Spearman correlation, Fig. 2a); hence, the
intra-sample strain diversity is well explained by the nucleotide diversity. The variability in π^{SOX}
agrees with the three age-related groups observed before for the number of SOX strains across
mussel size. Small mussels (≤7cm) and large mussels (14.6cm – 24.1cm) have a low π^{SOX} and
harbor one to two strains. Medium-sized mussels (7.2cm – 14.1cm) have a high π^{SOX} and harbor
two to nine strains. The community in the largest mussel is an exception, as it has a high π^{SOX},
similar to medium-sized mussels, which can be explained by the presence of three strains from
two clades.

The MOX nucleotide diversity is significantly lower in comparison to SOX (intra-sample
π^{MOX} between 5.6×10^{-6} and 7.0×10^{-4}, Table 2, Wilcoxon signed rank test, p=0.015, Fig. 2).
Similar to SOX, the MOX α-diversity is significantly positively correlated with π^{MOX} ($\rho^2 = 0.89$,
$p < 10^{-6}$, Spearman correlation) (Fig. 2b). One group of mussels harbors only MOX strains from
clade 2 and is characterized by low MOX nucleotide diversity (A, C, J, L, M, P, Q, R, S, π^{MOX}
between 5.6×10^{-6} and 2.1×10^{-5}), while the other group habors MOX strains from both clades and
is characterized by high MOX nucleotide diversity (B, D, E, F, G, H, I, K, N, O, π^{MOX} between
1.4×10^{-4} and 7.0×10^{-4}). These groups are not associated with mussel size. Taken together, we
observed a strong correlation between the nucleotide diversity π and α-diversity for both
symbionts. Notably, π is based on all the detected SNVs whereas the α-diversity is based only
on the strain composition and relatedness. Thus, the strong correlation demonstrates that the
strain diversity captures most of the core genome-wide nucleotide diversity.

A comparison of the π values estimated here to other microbiome studies shows that
higher π^{SOX} have been observed in other *Bathymodiolus* species (mean between 2.2 $\times 10^{-3}$ and
3.9×10^{-3})[25]. The average SOX and MOX nucleotide diversity estimated here is within the range
of π values observed in the clam *Solemya velum* microbiome where the symbiont transmission
mode is thought to be a mixture of vertical and horizontal transmission[40]. Furthermore, our π
estimates are lower than those observed for most bacterial species in the human gut
microbiome that are considered horizontally transmitted[19].

Geographic isolation of bacterial communities associated with individual mussels.

Symbiont transmission mode is an important determinant of the community assembly

dynamics[11]. For horizontally transmitted microbiota, similar community composition among

hosts may develop depending on factors that affect the community assembly such as the

environmental bacterial biodiversity or the order of colonization[41]. To study the degree of

geographic isolation between mussel hosts, we calculated genome-wide fixation index F_{ST} and

the ecological measure β-diversity at the strain resolution across the metagenomic samples for

the two symbionts. Small F_{ST} indicates that the samples stem from the same population

whereas large F_{ST} indicates that the samples constitute subpopulations.

Our results revealed generally high pairwise F_{ST} values, indicating a strong genetic

isolation between individual mussels (mean pairwise F_{ST}^{SOX} of 0.618, mean pairwise F_{ST}^{MOX} of

0.495, Fig. 2); hence, most mussels in our sample harbor an isolated symbiont subpopulation of

SOX and MOX. In addition, the SOX β-diversity is significantly positively correlated with

F_{ST}^{SOX}(ρ^2=0.7, p <10^{-6}, Spearman correlation). We observed subpopulations of mussels that are

characterized by a low pairwise F_{ST}^{SOX} within the subpopulation and a high pairwise F_{ST}^{SOX} with

other mussels. This subpopulation structure is also represented in the distribution of β-diversity

(Fig. 2). Thus, mussels from the same subpopulation harbor genetically similar SOX

communities and similar strain composition. Examples are one group of mussels including L, O,

P, and Q that contains only strains of clade S2 and another group including the mussels M, N,

and R that contains only strains of clade S1 (Fig. 2a). Notably, the two subpopulations contain

only large mussels that are characterized by a low π^{SOX}.

The distribution of pairwise F_{ST}^{MOX} revealed two main groups: one mussel group is

characterized by high pairwise F_{ST}^{MOX} and low π^{MOX} while the other group is characterized by

lower F_{ST}^{MOX} and high π^{MOX} (Fig. 2b). These correspond to the previously described groups,

where one contains mussels with a low π^{MOX} and strains from clade M2 and the other group

contains mussels with a high π^{MOX} and strains from both clades. We did not observe an

association between MOX β-diversity and F_{ST}^{MOX} (p>0.05, Spearman correlation), which can be

explained by the high proportion of invariant SNVs in MOX. Although the analysis of F_{ST}^{MOX} did

not reveal MOX subpopulations, the pattern of β-diversity uncovered subpopulations that show

a high pairwise F_{ST}^{MOX}. These subpopulations have a low β-diversity and a low nucleotide

diversity. One subpopulation consisting of large mussels (P, Q, S) is characterized by the

presence of strain M2.3 and the absence of clade M1. Another subpopulation (A, C, J, L, M, R) containing mussels of different sizes is characterized by the dominance of strains M2.1 and M2.2 and the absence of clade M1. Thus, the comparison of strain composition across mussels revealed that the population of MOX is substructured similarly to SOX. However, unlike SOX, the MOX subpopulations are not associated with specific mussel shell sizes.

The high F_{ST} values and the population structure we observed here reveal population stratification, that is especially pronounced for SOX. One possible factor that influences symbiont population structure is host genetics, whose impact on the composition of horizontally transmitted microbiota has been debated in the literature. Studies of the mammal gut microbiome showed that the host genotype had a contribution to the microbiome composition in mice[42], whereas the association with host genetics was reported to be weak in humans[43]. Analyzing 175 SNVs in 12 mitochondrial genes, we detected no association between mussel F_{ST} and symbiont F_{ST} for any of the two symbionts (Supplementary Information, Supplementary Fig. 8). Consequently, we conclude that the strong subpopulation structure observed for SOX and MOX cannot be explained by mussel relatedness (i.e., host genetics) or location.

Our results provide evidence for a strong genetic isolation between the symbiont populations associated with individual mussels. This finding is consistent with the observed individual-specific symbiont strain composition. In contrast, much lower F_{ST} values were found for SOX populations in other *Bathymodiolus* species sampled from hydrothermal vents (mean F_{ST} per site between 0.05 and 0.17), which implies a weaker genetic isolation in these vents[25]. Our analysis of cold seep *B. brooksi* data revealed SOX subpopulations with low genetic isolation that are observed using both F_{ST}, which takes all SNVs into account, and β-diversity at the level of strains. In contrast, only β-diversity disclosed subpopulations for MOX. Thus, strain-resolved metagenomics resolves similarities between individual mussel microbiomes below the species level.

Discussion

Our analysis revealed strong genetic isolation of symbiotic bacterial populations in individual mussel hosts, indicating geographic isolation between mussels. We hypothesize that this geographic isolation occurs through restricted uptake of SOX and MOX symbionts from the environment over time. The lack of evidence for strong adaptive selection in SOX and MOX strains suggests that the inter-host population structure is the result of neutral processes rather than host discrimination against different strains. Here, we propose a neutral model for symbiont

community assembly that explains how restricted symbiont uptake and colonization impose barriers to the symbiont dispersal, which can, over time, lead to inter-host population structure and contribute to the formation of cohesive genetic units within the symbiont population (Fig. 3). In our model, bacteria are acquired from the environmental symbiont pool in juveniles[44]. The presence of a symbiont environmental pool was suggested before based on the detection of symbiont genes in adjacent seawater[45,46]. Nevertheless, the loss of central metabolic enzymes suggests that bacteria disperse in a dormant state[47]. We hypothesize that the dormancy of free-living symbionts and the preservation of few symbiont cells inside bacteriocytes[23] contribute to the isolation of bacterial populations inside the host cells from the rest of the population, which can lead to recombination barriers. Our results support the self-infection hypothesis[26], according to which, once the gill is first colonized, bacteria present in ontogenically older tissue infect newly formed gill filaments; thus, the uptake of symbionts from the environment is limited. In addition, decreased growth rate in older mussels may also lead to decreased symbiont uptake. This model provides a plausible explanation for the observed pattern of strong symbiont genetic isolation between mussels and of reduced SOX strain diversity in large mussels. Possible later infections of the gill tissue from the environmental pool may occur due to symbiont loss and replacement driven by environmental changes or increased gill growth rate. Notably, our results are in contrast to a recent study on other *Bathymodiolus* species from hydrothermal vents, that concluded that SOX populations from individual mussels of the same site intermix[25]. This contrast may be explained by differences in the symbiont abundance in the seawater, which is expected to play a role in the colonization process. Our samples originate from a cold seep site with low mussel density (Supplementary Fig. 1); thus, the concentration of symbionts in the surrounding seawater may be correspondingly low. The low symbiont abundance would result in a low probability of later infections and a prevalence of self-infection. In contrast, the symbiont abundance in the seawater at large and densely populated mussel beds at hydrothermal vents is expected to be higher, resulting in a higher probability of later infections.

The colonization of new filaments over the mussel lifespan via self-infection entails serial founder events on the bacterial population. Throughout this process, new mutations arising in the symbiont population during the lifetime of the mussel can reach fixation due to genetic drift following population bottelnecks. This process is expected to lead to a reduction of symbiont genetic diversity over the mussel life time. Thus, individual mussels develop into independent habitats that harbor individual populations, which are genetically isolated from other mussel-associated symbiont populations and from the environmental pool. The evolution of vertically

transmitted endosymbiont populations is similarly affected by serial founder effects[48], as we
suggest here for horizontally transmitted bacteria. However, migration between host-associated
populations and the environmental pool results in an increased effective population size for
horizontally transmitted bacteria; thus, the population is not subject to the fate of genome
degradation as commonly observed in vertically transmitted symbionts[15]. Serial founder effects
and recombination barriers due to geographic isolation are important drivers of lineage
formation in bacteria[39]. Reduction of genetic diversity due to transmission bottlenecks is
considered a hallmark of pathogen genome evolution[49]; examples are *Yersinia pestis*[50] and
Listeria monocytogenes[51]. Our model demonstrates that, similar to pathogenic bacteria, genome
evolution of bacteria with a symbiotic lifestyle can be affected by serial founder effects due to
self-infection.

Methods

Collection and sequencing

Twenty-three individuals of *Bathymodiolus brooksi* mussels were collected during a research
cruise with the E/V *Nautilus* from the cold seep location GC853 at the northern Gulf of Mexico in
May 2015. The mussel distribution at the cold seep was patchy and mussel individuals were
collected from three distinct clumps within a radius of 131 meters (coordinates clump a:
28.1237, -89.1404 depth: -1073m, clump b: 28.1241, -89.1401 + depth: -1073m, clump c:
28.1237, -89.1404 + depth: -1073 to 1078m). The gills from each mussel individual were
dissected immediately after retrieval and homogenized with sterilized stainless steal beads, 3.2
mm in diameter (biostep, Germany). A subsample of the homogenate for sequencing analyses
was preserved in RNA later (Sigma, Germany) and stored at -80°C. DNA was extracted from
these subsamples as described by[52]. TruSeq library preparation and sequencing using Illumina
HiSeq2500 was performed by the Max Planck Genome Centre in Cologne, Germany, resulting
in 250 bp paired-end reads with a median insert size of 400 bp. The raw reads have been
deposited in NCBI under BioProject PRJNA508280.

Construction of the non-redundant gene catalog

Illumina paired-end raw reads from the samples were trimmed for adapters and filtered by
quality using BBMap tools[53]. Only reads with more than 30bp and quality above 10 were kept.
This results in 37.7 million paired-end reads per sample on average (Supplementary Table 1).

We assembled each of the metagenomic samples individually using metaSPAdes[54].
Genes were predicted *ab initio* on contigs with metaProdigal[55]. These predicted genes were
clustered by single-linkage according to sequence similarity using BLAT[56] (at least 95% of
sequence identity in at least 90% of the length of the shortest protein and e-value < 10^{-6}). To
reduce the potential inflation caused by the single-linkage clustering, we applied two additional
filters to discard hits: the maximum ratio allowed between the two compared sequence lengths
must be 4 and hits between partial and non-partial genes are discarded. These filters are meant
to remove spurious links between sequences due to the presence of commonly spread protein
domains. This clustering was performed in two successive steps; first, we obtained sample-
specific gene catalogs by performing intra-sample clustering. This is meant to reduce sequence
redundancy, resulting in an average of ~676,000 non-redundant genes per sample
(Supplementary Table 1). Second, one-sided similarity search was performed across all pairs of
sample catalogs. This resulted in 1,156,207 clusters (26.5%) and 3,207,869 (73.5%) singletons,
which make up a catalog of 4,364,076 million non-redundant genes. For each of the clusters,
we reconstructed a consensus sequence as cluster representative. To this end, we took the
majority nucleotide at each position (ties were resolved randomly).

Taxonomic annotation of gene catalog

Taxonomic annotation of the gene catalog was performed by aligning the translated genes to
the non-redundant protein NCBI database (date: 24/05/18) using diamond[57] (e-value<10^{-3},
sequence identity $\geq$ 30%) and obtaining the best hit. Genes were annotated as MOX-related if
their best hit is *Bathymodiolus platifrons* methanotrophic gill symbiont (NCBI Taxonomy ID
113268) or *Methyloprofundus sedimenti* (NCBI Taxonomy ID 1420851). For SOX, the genomes
of thioautotrophic symbionts belonging to four different Bathymodiolus species were used for
annotation (NCBI Taxonomy IDs: 2360, 174145, 113267 and 235205). In addition, the gene
catalog was screened for mitochondrial genes using best blastp hits against the *Bathymodiolus*
platifrons mitochondrial protein sequences (NC_035421.1)[58] (all e-values <10^{-40}). The gene
catalog was also screened for symbiont marker genes by best blastp hits to a published protein
database for *Bathymodiolus azoricus* symbionts[47] (80% of protein identity and 100% of query
coverage). This allowed to identify 86 SOX and 39 MOX marker genes. The marker gene
coverages are generally uniform across a sample, however a high variance in coverage is
present in two of the samples (Supplementary Fig. 3). Since the binning method relies on the
covariation of coverage across samples, the presence of a high variance in coverage can

interfere with the proper clustering of genes, thus, two samples were discarded from further
analysis (Dsc1, Dsc2).

Estimation of the gene catalog coverages

To estimate the gene abundances, we mapped the reads of each metagenomic sample to the
gene catalog using bwa mem[59]. Reads below 95% of sequence identity or mapping quality of
20, as well as not primary alignments were discarded. Coverage per position for each gene in
the catalog across samples was calculated using samtools depth[60] and the gene coverage is
given by the mean coverage across positions. We first downsampled the reads in each sample
to the minimum number of reads found (33M, Supplementary Table 1) and calculated mean
coverage per gene to perform the binning and the analyses of coverage variance across
symbiont marker genes (see above).

Genome binning and symbiont core genome identification

Next, we performed co-abundance gene segregation by using a canopy clustering algorithm[29],
which clusters genes into bins that covary in their abundances across the different samples.
This approach allows to recover from chimeric associations obtained in the assembly process
and to automatically separate core from accessory genes. Gene coverages across samples
were used as the abundance profiles for binning. First, genes with a Pearson correlation
coefficient (PCC) > 0.9 to the cluster abundance profile were clustered. Then, clusters with PCC
> 0.97 between their median abundance profiles were merged and outlier clusters for which the
coverage signal originates from less than three samples were removed. In addition, we removed
a gene from a cluster if Spearman correlation coefficient to the median canopy coverage profile
is lower than 0.7. Finally, overlaps among the clusters were removed by keeping a gene in the
largest of the clusters in which it has been found.

This allowed us to cluster 900,310 genes into 98,944 co-abundant gene groups (3 to 699
genes) and three MetaGenomic Species (MGSs, ≥700 genes). An additional filter was applied
to the MGSs to obtain final bins by removing outlier genes based on their coverage
(Supplementary Fig. 2). To this end, we used the Median Absolute Deviations (MAD) statistic as
a cutoff to discard highly or lowly covered genes. We removed genes that are at least 24 times
MAD far from the median in at least one of the samples. The bins after outlier gene removal
constitute the core genomes of the MGSs. We checked for the completeness of the symbiont

bins with CheckM, by screening for the presence of Gammaproteobacteria universal single copy
marker genes[61].

SNV discovery on the core genomes

To perform single nucleotide variant (SNV) discovery, we mapped the downsampled reads
individually for each sample to the gene catalog. Because sample size has been shown to have
an effect on variant detection[62], we normalized the data across samples. To this end, we
normalized each sample to the smallest median coverage found in a sample (482x coverage for
SOX, 36x coverage for MOX and 568x for mitochdonrial genes). LoFreq was used for
probabilistic realignment and variant calling of each sample independently[63]. SNVs detected
with LoFreq have been hard filtered using the parameters suggested by GATK best practices[64].
Briefly, SNVs with quality by depth below 2, Fisher's exact test Phred-scaled probability for
strand bias above 60, root mean square of mapping quality below 40, root mean square of base
quality above 30, mapping quality rank sum test below -12.5 and read position rank sum test
below -8 are kept for further analyses.

The resulting SNVs can be fixed or polymorphic in a sample. Polymorphic SNVs are
characterized by the allele frequency of the alternative allele whereas fixed SNVs have an allele
frequency of 1. Here, we define SNVs as polymorphic in a metagenomic sample if their
frequency is between 0.05 and 0.95 in the sample.

Population structure analyses

489 SNV data is used for calculating intra-sample and inter-sample nucleotide diversity (π) as
490 applied before to human gut microbiome species[19]. Intra-sample nucleotide diversity (π) is given
491 as:

$$\pi(H, G) = \frac{1}{|G|} \sum_{i=1}^{|G|} \sum_{B_1 \in \{ACTG\}} \sum_{B_2 \in \{ACTG\} \setminus B_1} \frac{X_{i,B_1}}{C_i} \frac{X_{i,B_2}}{C_i - 1}$$

where *H* corresponds to the sample, *G* to the bacterial genome, |*G*| is the length of the analyzed
genome and $X_{i,Bj}$ is the count of a specific nucleotide B_j at a specific locus *i* with coverage C_i.
Inter-sample nucleotide diversity (π) is then given as follows, where H_1 and H_2 correspond to the
two samples compared:

$$\pi(H_1, H_2, G) = \frac{1}{|G|} \sum_{i=1}^{|G|} \sum_{B_1 \in \{ACTG\}} \sum_{B_2 \in \{ACTG\} \setminus B_1} \frac{X_{i,B_1,S_1}}{C_{i,S_1}} \frac{X_{i,B_2,S_2}}{C_{i,S_2}}$$

Finally, these diversity measures are used to estimate the fixation index (F_{ST}), which measures

genetic differentiation based on the nucleotide diversity present within and between populations.

$$F_{ST}(S_1, S_2, G) = 1 - \frac{\pi_{within}}{\pi_{between}} = 1 - \frac{\pi(S_1, G) + \pi(S_2, G)}{2\,\pi(S_1, S_2, G)}$$

The scripts to calculate genome-wide inter and intra-sample nucleotide diversity (π) and fixation

index (F_{ST}) across all inter-sample comparisons from pooled SNV data have been deposited at

https://github.com/deropi/BathyBrooksiSymbionts.

Strain deconvolution

We reconstructed the strains for the core genomes with DESMAN[33]. The SNVs with two states

and their frequencies in each sample are used by DESMAN to identify strains in the core

genomes that are present over multiple samples. Thereby, the program uses the SNV

frequency covariation across samples to assign the SNV states to a specific genotype. For

SOX, we ran the strain deconvolution five times using different seed numbers and 500

iterations. Due to computational limitations, a subset of 5,000 SNVs was used and the

haplotypes considering the whole SNV dataset were inferred *a posteriori*. The five replicates

were run for an increasing number of strains from seven to twelve. The program uses posterior

mean deviance as a proxy for model fit. A posterior mean deviance lower than 5% was reached

in the transition from eleven to twelve strains, therefore the number of inferred SOX strains is

eleven. We did not run fewer numbers of strains due to the presence of large posterior mean

deviances between runs with a small strain number. Additionally, we ran DESMAN for the SOX

dataset that was subsampled to the MOX coverage with no replicates and eleven strains were

found using posterior mean deviance. For MOX, we ran four replicates using the whole SNV

dataset and 500 iterations. The runs were performed by using an increasing number of strains

from two to seven, reaching the optimal number of six strains. The consensus gene sequences

of each strain were concatenated to generate the strain core genomes, which were used for

further analyses. Splits network of the strain genome sequences were reconstructed using

SplitsTree[65] and uncorrected distances. The position of the root in the splits network was

estimated by the minimum ancestral deviation (MAD) method[66], which uses maximum likelihood

phylogenetic trees inferred with IQ-TREE[67].

α- and β-diversity

To study the microbial community composition, we estimated α- and β-diversity accounting for strain relatedeness in addition to species richness and eveness. α-diversity was estimated using phylogeny species eveness (PSE)[68] implemented in the R package 'Picante'[69]. β-diversity was estimated using the weighted Unifrac distance, which is implemented in the R package 'GUniFrac'[70]. This measure quantifies differences in strain community composition between two samples and accounts for phylogenetic relationships.

Allele frequency spectra estimation

The unfolded allele frequency spectra were calculated from biallelic SNVs for each of the bacterial species within individual samples. The unfolded allele frequency spectrum estimation relies on the presence of ancestral states in the population. Because we have no information about the ancestry relationship among the strains present in the samples, we made one main assumption in this regard: the ancestral SNV state in the population corresponds to the one which is present in the higher number of strains. Ties are resolved by arbitrarily assigning one tip of the tree as ancestral state: M2.2 for MOX and S4 for SOX.

pN/pS and Neutrality Index estimation

We estimated pN/pS for both bacterial populations, which is a variant of dN/dS that can be used based on intra-species SNVs. To this end we first calculated the expected ratio of nonsynonymous and synonymous mutations for each gene by accounting for each possible mutation occurring in each of the codons. Then, we estimated the observed nonsynonymous to synonymous ratio by using the biallelic SNVs. These two measures are later compared, resulting in the pN/pS ratio. pN/pS was estimated genome-wide as well as individually for each of the genes in the two symbiont species. The per-gene pN/pS calculation results into undefinded estimates for genes with no synonymous mutations. To circumvent this limitation, we added 1 to the number of observed synonymous mutations in each gene, which is a standard correction for dN/dS ratios[71].

The neutrality index (NI) accounts for differences in the ratio of nonsynonymous to synonymous variants between divergent and polymorphic SNVs in order to quantify the departure of a population from neutral evolution[37]. $NI = \frac{pN/pS}{dN/dS}$, where *pN*, and *pS* are the number of polymorphic synonymous and nonsynonymous sites, respectively, and *dN* and *dS*

are the number of divergent synonymous and nonsynonymous sites, respectively. For a coalescent population that evolves neutrally, the nature of fixed mutations that are involved in the divergence of the strains should not be different from that of the polymorphic mutations. An excess of divergent nonsynonymous mutations (NI<1) indicates that the population underwent positive selection or a large demographic change in the past[37].

Here we used the NI to analyze if differences in selection have been involved in the evolution of SOX and MOX strains. Different strains are typically found in more than one sample and this supports the notion that SNVs that characterize the strains constitute substitutions. We estimated NI by considering two different levels of divergence and polymorphism. First, we defined as divergent all those SNVs that have two possible states among the strains and as polymorphic all the invariant SNVs. Second, we used a more restrictive level of divergence. To this end, we excluded putative recently acquired SNVs from the set of divergent SNVs, by discarding those that have multiple states among strains from the same group. Polymorphic SNVs are all the remaining. The scripts to calculate the allele frequency spectra, pN/pS and NI have been deposited at https://github.com/deropi/BathyBrooksiSymbionts. Statistics and plotting were done in R[72].

Acknowledgements

We thank the captain and crew of the E/V *Nautilus* 2015 Expedition, the team of the ROV *Hercules*, the chief scientist on this expedition leg, Eric Cordes, and the expedition leader, Mike Brennan. Additionally, we thank Elie Jami, Robin Koch, Tanita Wein, and Christian Wöhle for critical comments on the manuscript. This work was supported by the CRC1182 *Origin and Function of Metaorganisms*, the Bioinformatics Network at Kiel University, and the North-German Supercomputing Alliance (HLRN, project shb00002). Additional funding support for ND, JMP and RA came from the Max Planck Society, the Gordon and Betty Moore Foundation Marine Microbiology Initiative Investigator Award through grant GBMF3811 to ND and a European Research Council Advanced Grant (BathyBiome, Grant 340535 to ND).

Author contributions

AK, TD, JMP, and ND designed the study, RA, JMP, and ND collected the data, DRP analyzed

586   the data, DRP, AK, and TD interpreted the results with contribution from RA, DRP, AK, and TD

587   wrote the manuscript with contributions from all authors.

589   **Figure Legends**

Figure 1: Symbiont strain abundances (a, d), symbiont strain relationships (b, e) and example allele frequency spectra (c, f). a, b, c, 11 strains reconstructed for SOX. These cluster into four clades, with two times four, three and one strain per clade, labelled by shades of green, red, blue, and yellow. The strains differ by between 669 SNVs (strains S2.2 and S2.3, sequence identity 99.95%) and 8,171 SNVs (strains S3.2 and S4 sequence identity 99.36%). Minimum number of SNVs between strains of different clades is 6,451 (strains S1.1 and S2.1, sequence identify 99.49%). **d, e, f,** 6 strains reconstructed for MOX. These cluster into two clades, labelled by shades of red and blue. Strains differ by between 105 (strain M2.2 and M2.3, sequence identity 99.99%) and 2,677 SNVs (strain M1.1 and M2.1, sequence identity 99.81). The minimum number of SNVs differentiating strains from different clades is 2,224 (strains M2.2 and M1.3, sequence identity 99.85%). **a, d,** Stacked barplot of strain relative abundances for each individual mussel. Mussel individuals are labeled with an assigned letter (A-S), followed by the sampling clump (a, b or c) and the shell size (cm). **b, e,** Splits network of the strain genome sequences. Scale bar shows the number of differences per site. The red dots indicate the position of the root. **c, f,** Example of derived allele frequency spectra (sample E). Different colors represent different strain clades (see also Supplementary Fig. 6).

 Figure 2: Symbiont population structure for a, SOX and b, MOX. Top left triangle: Intra-
sample π and symbiont fixation index (F_{ST}) based on SNVs. Lower right triangle: α- and β-
diversity based on reconstructed strains. Rows and columns are labelled by sample name,
sample location, and shell size. Heatmap hierarchical clustering is based on Euclidean distance
of F_{ST}. **a,** SOX: mean pairwise F_{ST} is 0.618. Two subpopulations show an extreme degree of
isolation: mean pairwise F_{ST} of subpopulation composed of M, N, R, is 0.313; mean pairwise F_{ST}
of subpopulation composed of L, O, P, Q is 0.308; mean F_{ST} of sample pairs where one sample
is M, N, or R and the other sample is L, O, P, or Q is 0.969. **b,** MOX: mean pairwise F_{ST} is
0.495. The clustering displays two groups: mean pairwise β-diversity of subpopulation
composed of A, B, C, D, E, F, G, H, I, J, L is 0.099; mean pairwise β-diversity of subpopulation
composed of K, M, N, O, P, Q, R, S is 0.383.

Figure 3. Symbiont colonization dynamics. a, The post larvae mussel gill does not take up endosymbionts until the gill presents several filaments and the gill epithelial cells reach a determined developmental stage[26]. At this time point, the filaments are simultaneously infected by different strains via endocytosis (1). This imposes the first bottleneck in the symbiont population, since most likely, not all the strains from the environmental pool can infect the tissue. **b,** Bacteria are released from the host tissue to the environmental pool. As the mussel grows, new filaments are continuously formed in the gill throughout the mussel life span (growing cells shaded in purple). The new tissue is colonized by a self-infection process[26], which involves infection of the newly formed filaments via endocytosis with bacteria that are released from old tissue via exocytosis (2). The spatial distribution of strains within the gill tissue also supports self-infection[45]. The continuous self-infection process imposes serial founder effects that lead to a reduction in strain diversity, which is mostly driven by drift. Additional sources of diversity loss are: tissue replacement (3) and regulated lysosomal digestion of symbionts[58] (4). **c,** In older mussels, a unique strain dominates the gill. In addition, *de novo* mutations occur in symbiont genomes (marked by x). Due to serial founder effects within the same mussel, those variants can be quickly fixed inside the mussel (5). **d,** As the mussel dies, bacteria are released from the gill, going back to the environmental pool (6).

Tables

Table 1. Neutrality index (NI) for the symbiont core genomes.

a

	SOX		MOX	
	divergent	polymorphic	divergent	polymorphic
nonsynonymous SNVs	5004	990	2115	704
synonymous SNVs	10577	1450	1313	515
nonsynonymous SNVs/synonymous SNVs	0.47	0.68	1.61	1.37
NI	1.44		0.85	

b

	SOX		MOX	
	divergent	polymorphic	divergent	polymorphic
nonsynonymous SNVs	2549	3455	1041	1778
synonymous SNVs	6370	5657	649	1179
nonsynonymous SNVs/synonymous SNVs	0.40	0.61	1.60	1.51
NI	1.52		0.94	

a, divergent SNVs are all those SNVs that differ between at least two strains, i.e., all identified strains are considered as diverged taxonomic units, and polymorphic SNVs are all the invariant SNVs. **b,** Divergent SNVs have the same state inside a strain clade and are not invariant and polymorphic SNVs are all the remaining, i.e., only the clades are considered as diverged taxonomic units.

Table 2. Nucleotide diversity (π), Fixation Index (F_{ST}), and pN/pS calculations for both symbiont populations.

	SOX	MOX
Intra-sample π range	5.2×10^{-5} - 3.6×10^{-3}	5.6×10^{-6} - 7.0×10^{-4}
intra-sample π mean	$1.4 \times 10^{-3} \pm 1.3 \times 10^{-3}$ (s.d)	$2.7 \times 10^{-4} \pm 2.8 \times 10^{-4}$ (s.d)
intra-sample π median	6.7×10^{-4}	1.4×10^{-4}
Pairwise F_{ST} range	0.151- 0.986	0.096- 0.898
Mean pairwise F_{ST}	0.618	0.495
pN/pS	0.137	0.425

References

1. McFall-Ngai, M. *et al.* Animals in a bacterial world, a new imperative for the life sciences. *Proc. Natl. Acad. Sci.* **110**, 3229–3236 (2013).
2. McFall-Ngai, M. J. The Importance of Microbes in Animal Development: Lessons from the Squid-Vibrio Symbiosis. *Annu. Rev. Microbiol.* **68**, 177–194 (2014).
3. Shabat, S. K. B. *et al.* Specific microbiome-dependent mechanisms underlie the energy harvest efficiency of ruminants. *Isme J.* **10**, 2958 (2016).
4. Schretter, C. E. *et al.* A gut microbial factor modulates locomotor behaviour in Drosophila. *Nature* **563**, 402–406 (2018).
5. Costello, E. K., Stagaman, K., Dethlefsen, L., Bohannan, B. J. M. & Relman, D. A. The Application of Ecological Theory Toward an Understanding of the Human Microbiome. *Science* **336**, 1255–1262 (2012).
6. David, L. A. *et al.* Diet rapidly and reproducibly alters the human gut microbiome. *Nature* **505**, 559–563 (2014).
7. Sommer, F. *et al.* The Gut Microbiota Modulates Energy Metabolism in the Hibernating Brown Bear *Ursus arctos. Cell Rep.* **14**, 1655–1661 (2016).
8. Koren, O. *et al.* Host Remodeling of the Gut Microbiome and Metabolic Changes during Pregnancy. *Cell* **150**, 470–480 (2012).
9. Jones, R. J., Hoegh-Guldberg, O., Larkum, A. W. D. & Schreiber, U. Temperature-induced bleaching of corals begins with impairment of the CO_2 fixation mechanism in zooxanthellae. *Plant Cell Environ.* **21**, 1219–1230 (1998).
10. Riou, V. *et al.* Influence of CH_4 and H_2S availability on symbiont distribution, carbon assimilation and transfer in the dual symbiotic vent mussel *Bathymodiolus azoricus. Biogeosciences* **5**, 1681–1691 (2008).
11. Bright, M. & Bulgheresi, S. A complex journey: transmission of microbial symbionts. *Nat. Rev. Microbiol.* **8**, 218–30 (2010).
12. Wernegreen, J. J. Endosymbiont evolution: predictions from theory and surprises from genomes: Endosymbiont genome evolution. *Ann. N. Y. Acad. Sci.* **1360**, 16–35 (2015).
13. Woyke, T. *et al.* One Bacterial Cell, One Complete Genome. *PLoS ONE* **5**, e10314 (2010).
14. Guyomar, C. *et al.* Multi-scale characterization of symbiont diversity in the pea aphid complex through metagenomic approaches. *Microbiome* **6**, 181 (2018).
15. Boscaro, V. *et al.* Parallel genome reduction in symbionts descended from closely related free-living bacteria. *Nat. Ecol. Evol.* **1**, 1160 (2017).
16. Klose, J. *et al.* Endosymbionts escape dead hydrothermal vent tubeworms to enrich the free-living population. **112**, 11300–11305 (2015).
17. Hagen, M. J. & Hamrick, J. L. Population level processes in *Rhizobium leguminosarum* bv. *trifolii*: the role of founder effects. *Mol. Ecol.* **5**, 707–714 (1996).
18. Vega, N. M. & Gore, J. Stochastic assembly produces heterogeneous communities in the *Caenorhabditis elegans* intestine. *PLOS Biol.* **15**, e2000633 (2017).
19. Schloissnig, S. *et al.* Genomic variation landscape of the human gut microbiome. *Nature* **493**, 45–50 (2013).
20. Ellegaard, K. M. & Engel, P. Genomic diversity landscape of the honey bee gut microbiota. *Nat. Commun.* **10**, 446 (2019).
21. Wollenberg, M. S. & Ruby, E. G. Population Structure of *Vibrio fischeri* within the Light Organs of *Euprymna scolopes* Squid from Two Oahu (Hawaii) Populations. *Appl. Environ. Microbiol.* **75**, 193–202 (2009).
22. Won, Y.-J. *et al.* Environmental Acquisition of Thiotrophic Endosymbionts by Deep-Sea Mussels of the Genus *Bathymodiolus. Appl. Environ. Microbiol.* **69**, 6785–6792 (2003).
23. Dubilier, N., Windoffer, R. & Giere, O. Ultrastructure and stable carbon isotope composition of the hydrothermal vent mussels *Bathymodiolus brevior* and *B.* sp. affinis *brevior* from the North Fiji Basin, western Pacific. *Mar. Ecol. Prog. Ser.* **165**, 187–193 (1998).
24. Duperron, S. *et al.* Diversity, relative abundance and metabolic potential of bacterial endosymbionts in three *Bathymodiolus* mussel species from cold seeps in the Gulf of Mexico. *Environ. Microbiol.* **9**, 1423–1438 (2007).

25. Ansorge, R. *et al.* Diversity matters: Deep-sea mussels harbor multiple symbiont strains. *bioRxiv*
531459 (2019).
26. Wentrup, C., Wendeberg, A., Schimak, M., Borowski, C. & Dubilier, N. Forever competent: Deep-sea
bivalves are colonized by their chemosynthetic symbionts throughout their lifetime. *Environ.*
Microbiol. **16**, 3699–3713 (2014).
27. Schöne, B. R. & Giere, O. Growth increments and stable isotope variation in shells of the deep-sea
hydrothermal vent bivalve mollusk *Bathymodiolus brevior* from the North Fiji Basin, Pacific Ocean.
Deep Sea Res. Part Oceanogr. Res. Pap. **52**, 1896–1910 (2005).
28. Van Dover, C. Community structure of mussel beds at deep-sea hydrothermal vents. *Mar. Ecol. Prog.*
Ser. **230**, 137–158 (2002).
29. Nielsen, H. B. *et al.* Identification and assembly of genomes and genetic elements in complex
metagenomic samples without using reference genomes. *Nat. Biotechnol.* **32**, 822–828 (2014).
30. Riou, V. *et al.* Influence of CH_4 and H_2S availability on symbiont distribution, carbon assimilation and
transfer in the dual symbiotic vent mussel *Bathymodiolus azoricus*. *Biogeosciences* **5**, 1681–1691
(2008).
31. Shapiro, B. J. & Polz, M. F. Ordering microbial diversity into ecologically and genetically cohesive
units. *Trends Microbiol.* **22**, 235–247 (2014).
32. Kashtan, N. *et al.* Single-cell genomics reveals hundreds of coexisting subpopulations in wild
Prochlorococcus. *Science* **344**, 416–20 (2014).
33. Quince, C. *et al.* DESMAN: a new tool for de novo extraction of strains from metagenomes. *Genome*
Biol. **18**, 181 (2017).
34. Ho, P.-T. *et al.* Geographical structure of endosymbiotic bacteria hosted by *Bathymodiolus* mussels
at eastern Pacific hydrothermal vents. *BMC Evol. Biol.* **17**, 121 (2017).
35. Rocha, E. P. C. *et al.* Comparisons of dN/dS are time dependent for closely related bacterial
genomes. *J. Theor. Biol.* **239**, 226–235 (2006).
36. Kryazhimskiy, S. & Plotkin, J. B. The population genetics of dN/dS. *PLoS Genet.* **4**, e1000304 (2008).
37. Rand, D. M. & Kann, L. M. Excess amino acid polymorphism in mitochondrial DNA: contrasts among
genes from *Drosophila*, mice, and humans. *Mol. Biol. Evol.* **13**, 735–748 (1996).
38. Lorion, J. *et al.* Adaptive radiation of chemosymbiotic deep-sea mussels. *Proc R Soc B* **280**,
20131243 (2013).
39. Achtman, M. & Wagner, M. Microbial diversity and the genetic nature of microbial species. *Nat. Rev.*
Microbiol. **6**, 431–440 (2008).
40. Russell, S. L., Corbett-Detig, R. B. & Cavanaugh, C. M. Mixed transmission modes and dynamic
genome evolution in an obligate animal–bacterial symbiosis. *ISME J.* **11**, 1359–1371 (2017).
41. Sprockett, D., Fukami, T. & Relman, D. A. Role of priority effects in the early-life assembly of the gut
microbiota. *Nat. Rev. Gastroenterol. Hepatol.* **15**, 197–205 (2018).
42. Benson, A. K. *et al.* Individuality in gut microbiota composition is a complex polygenic trait shaped by
multiple environmental and host genetic factors. *Proc. Natl. Acad. Sci.* **107**, 18933–18938 (2010).
43. Rothschild, D. *et al.* Environment dominates over host genetics in shaping human gut microbiota.
Nature **555**, 210–215 (2018).
44. Wentrup, C., Wendeberg, A., Huang, J. Y., Borowski, C. & Dubilier, N. Shift from widespread
symbiont infection of host tissues to specific colonization of gills in juvenile deep-sea mussels. *ISME*
J. **7**, 1244–7 (2013).
45. Ikuta, T. *et al.* Heterogeneous composition of key metabolic gene clusters in a vent mussel symbiont
population. *ISME J.* **10**, 990–1001 (2016).
46. Fontanez, K. M. & Cavanaugh, C. M. Evidence for horizontal transmission from multilocus phylogeny
of deep-sea mussel (Mytilidae) symbionts. *Environ. Microbiol.* **16**, 3608–3621 (2014).
47. Ponnudurai, R. *et al.* Metabolic and physiological interdependencies in the *Bathymodiolus azoricus*
symbiosis. *ISME J.* **11**, 463–477 (2017).
48. Reuter, M., Pedersen, J. S. & Keller, L. Loss of Wolbachia infection during colonisation in the
invasive Argentine ant Linepithema humile. *Heredity* **94**, 364–369 (2005).
49. Didelot, X., Walker, A. S., Peto, T. E., Crook, D. W. & Wilson, D. J. Within-host evolution of bacterial
pathogens. *Nat. Rev. Microbiol.* **14**, 150–162 (2016).
50. Gonzalez, R. J., Lane, M. C., Wagner, N. J., Weening, E. H. & Miller, V. L. Dissemination of a Highly
Virulent Pathogen: Tracking The Early Events That Define Infection. *PLOS Pathog.* **11**, e1004587
(2015).

51. Zhang, T. *et al.* Deciphering the landscape of host barriers to *Listeria monocytogenes* infection. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 6334–6339 (2017).

52. Zhou, J., Bruns, M. A. & Tiedje, J. M. DNA Recovery from Soils of Diverse Composition. *Appl. Environ. Microbiol.* **62**, 316–322 (1996).

53. Bushnell, Brian. BBMap. (2014). Available at: sourceforge.net/projects/bbmap/.

54. Nurk, S., Meleshko, D., Korobeynikov, A. & Pevzner, P. A. metaSPAdes: a new versatile metagenomic assembler. *Genome Res.* **27**, 824–834 (2017).

55. Hyatt, D., Locascio, P. F., Hauser, L. J. & Uberbacher, E. C. Gene and translation initiation site prediction in metagenomic sequences. *Bioinformatics* **28**, 2223–2230 (2012).

56. Kent, W. J. BLAT — The BLAST -Like Alignment Tool. *Genome Res.* **12**, 656–664 (2002).

57. Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using DIAMOND. *Nat. Methods* **12**, 59–60 (2015).

58. Sun, J. *et al.* Adaptation to deep-sea chemosynthetic environments as revealed by mussel genomes. *Nat. Ecol. Evol.* **1**, 0121 (2017).

59. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**, 1754–1760 (2009).

60. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).

61. Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res.* **25**, 1043–1055 (2015).

62. Subramanian, S. The effects of sample size on population genomic analyses – implications for the tests of neutrality. *BMC Genomics* **17**, 123 (2016).

63. Wilm, A. *et al.* LoFreq: A sequence-quality aware, ultra-sensitive variant caller for uncovering cell-population heterogeneity from high-throughput sequencing datasets. *Nucleic Acids Res.* **40**, 11189–11201 (2012).

64. Broad Institute. Best practices for variant calling with the GATK, https://www.broadinstitute.org/partnerships/education/broade/best-practices-variant-calling-gatk-1.

65. Huson, D. H. SplitsTree: analyzing and visualizing evolutionary data. *Bioinformatics* **14**, 68–73 (1998).

66. Tria, F. D. K., Landan, G. & Dagan, T. Phylogenetic rooting using minimal ancestor deviation. *Nat. Ecol. Amp Evol.* **1**, 0193 (2017).

67. Nguyen, L.-T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol. Biol. Evol.* **32**, 268–274 (2015).

68. Helmus, M. R., Bland, T. J., Williams, C. K. & Ives, A. R. Phylogenetic Measures of Biodiversity. *Am. Nat.* **169**, E68–E83 (2007).

69. Kembel, S. W. *et al.* Picante: R tools for integrating phylogenies and ecology. *Bioinformatics* **26**, 1463–1464 (2010).

70. Chen, J. GUniFrac: Generalized UniFrac Distances. (2018). Available at: https://CRAN.R-project.org/package=GUniFrac.

71. Stoletzki, N. & Eyre-Walker, A. The Positive Correlation between dN/dS and dS in Mammals Is Due to Runs of Adjacent Substitutions. *Mol. Biol. Evol.* **28**, 1371–1380 (2011).

72. R Core Team. *R: A Language and Environment for Statistical Computing.* (R Foundation for Statistical Computing, 2017).