Hif-1α stabilisation polarises macrophages via cyclooxygenase/prostaglandin E2 *in*

vivo

[1,2]Amy Lewis and [1,2]Philip M. Elks

[1]The Bateson Centre, University of Sheffield, Western Bank, Sheffield, UK. [2]Department of

Infection and Immunity and Cardiovascular Disease, University of Sheffield, Western Bank,

Sheffield, UK.

Corresponding author:

Dr Philip M. Elks

The Bateson Centre,

University of Sheffield,

Firth Court,

Western Bank,

Sheffield,

South Yorkshire,

S10 2TN.

UK

Tel: +44 (0) 1142 223609

p.elks@sheffield.ac.uk

Macrophage phenotype during inflammation and infectious disease are diverse and

are poorly understood, especially in the *in vivo* context where cues from the

microenvironment play a large role in cell behaviour. A crucial microenvironmental factor

associated with many disease pathologies is hypoxia, which has profound impacts upon

macrophage phenotypes through a variety of mechanisms. Further understanding of how

disease microenvironments impact macrophage polarisation may open up new therapeutic

avenues to fine-tune appropriate macrophage responses in disease situations. Here, we

show that hypoxia driven HIF-1α upregulates TNF production via cyclooxygenases in human

macrophages and in *in vivo* zebrafish systems, indicating a switch to an M1 phenotype via a

new mechanism. Our results suggest that macrophage phenotype could be altered

therapeutically via this axis in disease situations where hypoxia is an important feature of the

microenvironment.

Abstract

Macrophage subtypes are poorly characterised in disease systems *in vivo.* The initial

innate immune response to injury and infectious stimuli through M1 polarisation is important

for the outcome of disease. Appropriate macrophage polarisation requires complex

coordination of local microenvironmental cues and cytokine signalling to influence immune

cell phenotypes. If the molecular mechanisms behind macrophage polarisation were better

understood then macrophages could be pharmacologically tuned to better deal with bacterial

infections, for example tuberculosis. Here, using zebrafish *tnfa:GFP* transgenic lines as *in*

vivo readouts of M1 macrophages, we show that hypoxia and stabilisation of Hif-1α

polarises macrophages to a *tnfa* expressing phenotype. We demonstrate a novel

mechanism of Hif-1α mediated macrophage *tnfa* upregulation via a

cyclooxygenase/prostaglandin E2 axis, a mechanism that is conserved in human primary

macrophages. These findings uncover a novel macrophage HIF/COX/TNF axis that links

microenvironmental cues to macrophage phenotype that may have implications in

inflammation, infection and cancer, where hypoxia is a common microenvironmental feature

and where cyclooxygenase and Tnfa are major mechanistic players.

Introduction

The innate immune response to injury and pathogen invasion are complex and are

tightly regulated by coordination of microenvironmental cues and cytokine signalling.

Macrophages are important innate immune cells in disease and their activation states,

commonly termed polarisation states, are especially important during the early response to

injury and infection. Mammalian macrophages have been classified into two broad

polarisation/activation states; M1 (or classically activated) and M2 (alternatively activated)

(1). Pro-inflammatory, or M1, macrophages are highly antimicrobial and can phagocytose

and efficiently kill bacteria. Macrophages are also central players in tissue healing and

restoration of homeostasis post injury/infection, which requires a change in macrophage

phenotype to a wound healing, M2, state (2). A diverse variety in macrophage function has

been observed in many *in vitro* systems, yet control of macrophage subsets remains poorly

understood *in vivo*. Current classifications are based on *in vitro* mammalian datasets, using

monocytes that have been polarised to different macrophage phenotypes by addition of

cytokine-cocktails to the media (3). Specific subsets of macrophage polarisation are context

dependent *in vivo* and subtle changes in the microenvironment can lead to different

phenotypes of M1 and M2 macrophages, reflecting a spectrum rather than a binary

classification of behaviours (4). One important microenvironmental factor is hypoxia, which

has profound effects on macrophage phenotypes (5). To fully understand how

microenvironmental cues affect macrophage polarisation, *in vivo* systems are required to

study macrophage behaviour over time and space without disturbance from extrinsic factors.

Determining the molecular triggers and polarisation states of macrophages could open up

possibilities to manipulate macrophage phenotype during disease as novel therapeutic

avenues.

The tissue macrophage response to injury/bacterial infections is a rapid, pathogen

recognition receptor (PPR) mediated, switch to an M1 type phenotype, characterised by

production and release of pro-inflammatory mediators, including cytokines such as IL-1β, IL-

6, IL-12 and TNF (6, 7). This phenotypic switch is required for a successful response to the

invading pathogen and for clearance before infection can take hold (8, 9). However, in the

case of intracellular pathogens, such as *Mycobacterium tuberculosis* (Mtb), this macrophage

response is subverted, establishing a protective niche within innate immune cells (10–13).

This bacterial-mediated interference in macrophage polarisation highlights a gap for

potential therapeutic tuning of the macrophage response to allow for bacterial clearance (14,

15).

One potential macrophage tuning mechanism is via the Hypoxia Inducible Factor

(HIF) -1α transcription factor (16). HIF-1α is a master transcriptional regulator of the cellular

response to low levels of oxygen (hypoxia) (17). Leukocytes, including macrophages and

neutrophils, are exquisitely sensitive to tissue hypoxia, a signature of many disease

microenvironments (eg infections and cancer) due to lack of local blood supply and a high

turnover of local oxygen by pathogens (5, 18). Hypoxia and HIF-1α stabilisation have been

shown to have activating effects on macrophages *in vitro*, increasing their M1 profile and

bactericidal capabilities, yet the pro-inflammatory mechanisms behind this are not well

understood *in vivo* (5).

Hypoxia is a microenvironmental signal that increases epithelial

cyclooxygenase/prostaglandin E2 production in cancer models (19–21). Eicosanoids are

lipid signalling molecules and important pro-inflammatory mediators which are produced,

and released, by macrophages during early microbial infection (22, 23). The best

characterised of these are prostaglandins, strong inflammatory mediators that are known to

affect macrophage polarisation states during infection (24). A key family of enzymes for

prostaglandin biosynthesis is the cyclooxygenases (COX). COX-1 and -2 enzymes are

responsible for the production of prostaglandins as breakdown products of arachidonic acid.

Prostaglandins, in turn, are highly immunomodulatory, synergising with cytokines to amplify

pro-inflammatory responses (25). The potential effects of hypoxia-induced prostaglandins on

macrophage polarisation have not been investigated *in vivo*. Recent evidence indicates that

a HIF/COX/TNF axis may exist, with hypoxia-induced TNF expression in osteoblasts (bone

producing cells) being mediated via cyclooxygenase enzymes by an, as yet, unknown

mechanism (26). If observed in macrophages, this HIF/COX/TNF axis could have important

pro-inflammatory outcomes in disease and represent a novel therapeutic avenue.

Zebrafish have emerged as a useful model to understand innate immune cells in

their *in vivo* microenvironment (27). Development of multiple innate immune cell transgenic

lines, combined with non-invasive fluorescence imaging, has enabled mechanistic

investigation into macrophage biology (2, 28, 29). Well-characterised models of inflammation

and infection biology have emerged using zebrafish larvae. Mechanistic insights into

neutrophil and macrophage recruitment to an injury site have been gained through use of a

tailfin transection model (30), while host-pathogen interactions have been studied using the

natural fish pathogen *Mycobacterium marinum* (Mm) as a model of tuberculosis (27).

Recently, it has been demonstrated that *tnfa* promoter driven fluorescent transgenic lines

display an M1 type phenotype upon sterile injury immune challenge (31, 32). This advance

has enabled us to investigate macrophage polarisation within the zebrafish system during a

host-pathogen immune challenge. We have previously demonstrated that Hif-1α stabilisation

induces macrophage proinflammatory *il-1β* transcription and that this is host-protective

against Mm infection in the zebrafish model (33). We therefore hypothesised that Hif-1α

stabilisation would promote macrophage pro-inflammatory *tnfa* expression, via a

cyclooxygenase dependent mechanism, that could also be beneficial to the host during

infection.

Here, we investigated the regulation of expression of the important M1 pro-

inflammatory cytokine *tnfa* by Hif-1α stabilisation *in vivo*. We demonstrate that genetic,

pharmacological and hypoxia-mediated Hif-1α stabilisation upregulated macrophage *tnfa* in

zebrafish. We show that *tnfa* transcription is upregulated in macrophages after injury and

Mm infection (both at early- and granuloma- stage infection). We identify that Hif-1α

activates macrophage *tnfa* via cyclooxygenase/prostaglandin E2, while injury/Mm driven *tnfa*

production does not require active cyclooxygenase. Importantly, this novel macrophage

HIF/COX/TNF axis is conserved in primary human macrophages. These findings have

important implications in inflammation, infection and cancer biology where macrophage

polarisation states are influenced by microenvironmental hypoxia.

Materials and Methods

Zebrafish

Zebrafish were raised and maintained on a 14:10-hour light/dark cycle at 28ºC,

according to standard protocols (34), in UK Home Office approved facilities at The Bateson

Centre aquaria at the University of Sheffield. Strains used were Nacre (wildtype),

TgBAC(tnfa:GFP)pd1028,   *Tg(tnfa:eGFP-F)ump5Tg*,   *Tg(mpeg1:mCherry-F)ump2Tg,*

Tg(mpeg1:mCherryCAAX)sh378,   Tg(lyz:Ds-RED2)nz50   TgBAC(il-1β:eGFP)sh445 and

Tg(phd3:EGFP)i144 (31, 32, 35–37).

Tailfin injury

Inflammation was induced in zebrafish embryos by tail transection as previously

described (38). Embryos were anaesthetised at 2 days post fertilisation (dpf) by immersion

in 0.168mg/ml Tricaine, and transection of the tail was performed using a scalpel blade.

Larvae were imaged by confocal microscopy at 16 hour post wound (hpw) on a Leica TCS-

SPE confocal on an inverted Leica DMi8 base and imaged using 20x or 40x objective

lenses.

Mycobacterium marinum

Mm infection experiments were performed using M. marinum M (ATCC #BAA-535),

containing a psMT3-mCherry or psMT3 mCrimson vector (39). Injection inoculum was

prepared from an overnight liquid culture in the log-phase of growth resuspended in 2%

polyvinylpyrrolidone40 (PVP40) solution (CalBiochem) as previously described (40). 100 or

150 colony forming units (CFU) were injected into the caudal vein at 28-30hpf as previously

described (41).

Confocal microscopy of transgenic larvae

1dpi and 4dpi transgenic zebrafish larvae infected with fluorescent Mm strains were

mounted in 0.8-1% low melting point agarose (Sigma-Aldrich) and imaged on a Leica TCS-

SPE confocal on an inverted Leica DMi8 base and imaged using 20x or 40x objective

lenses.

For quantification purposes acquisition settings and area of imaging were kept the

same across groups. Corrected total cell fluorescence was calculated for each cell using

Image J as previously described (42, 43).

RNA injections

Embryos were injected with dominant *hif-1αb* (ZFIN: *hif1ab*) variant RNA at the one

cell stage as previously described (44). *hif-1α* variants used were dominant active (DA) and

dominant negative (DN) *hif-1α*. Phenol red (PR) (Sigma Aldrich) was used as a vehicle

control.

Pathway inhibitors

Unless otherwise stated, embryos were treated from 4 hours pre-Mm infection to 24

hours post infection (hpi) by addition to the embryo water and DMSO was used as a

negative solvent control. The pan hydroxylase inhibitor, DMOG (dimethyloxaloylglycine,

Enzo Life Sciences), was used at a 100µM concentration by incubation in E3 embryo media

as previously described (44). The selective PHD inhibitors JNJ-402041935 (Cayman

Chemicals) and FG-4592 (Selleckchem) were used at 100µM and 5µM respectively (45, 46).

The selective inhibitor of COX-1, SC-560 (Cayman Chemicals), was used at 30µM and the

selective inhibitor of COX-2, NS-398 (Cayman Chemicals), was used at 15µM by incubation

in E3 embryo media as previously described (47). The 15-LOX inhibitor PD146176 (Tocris)

was microinjected (1nl of 40nM) at 1 hour post infection (hpi). The leukotriene B4 receptor 1

(BLTR1) inhibitor, U-75302 (Abcam), was used at 30mM by incubation in E3 embryo media

as previously described (48) together with the BLTR2 inhibitor, LY255283 (Abcam), which

was used at 1mM. Exogenous prostaglandin E2 (Cayman Chemicals) and 15-keto-

prostaglandin E2 (Cayman Chemicals) were added by incubation in E3 at a concentration of

1µM (49).

Hypoxia

Embryos were incubated in 5% oxygen (with 5% carbon dioxide) in a hypoxia hood

(Baker-Ruskinn Sci-tive UM-027) from 32hpi for 6 hours and were imaged at 2dpf. Embryos

from the same clutch kept in incubated normoxic room air were used as controls (33).

Bacterial pixel count

Mm mCherry infected zebrafish larvae were imaged at 4dpi on an inverted Leica

DMi8 with a 2.5x objective lens. Brightfield and fluorescent images were captured using a

Hammamatsu OrcaV4 camera. Bacterial burden was assessed using dedicated pixel

counting software as previously described (50).

Human cells

Peripheral blood mononuclear cells (PBMC) were isolated by Ficoll-Paque PLUS

(GE Healthcare) density centrifugation of whole blood from healthy donors (National

Research Ethics Service reference 07/Q2305/7). PBMC were seeded at 2×10^6/ml in 24-well

tissue culture plates in RMPI 1640 media (Gibco) containing 2mmol/L L-glutamine (Lonza)

and 10% newborn calf serum (Gibco). Cells were cultured in 5% CO_2 at 37ºC and non-

adherent cells were removed after 24 hours. Remaining adherent cells were cultured for 14

days, with twice weekly media exchange, in RPMI 1640 supplemented with 2mmol/L L-

glutamine and 10% heat-inactivated fetal bovine serum (FBS) (PAN Biotech) (51).

For experiments RPMI 1640 + 25mM HEPES (Gibco) containing 2mmol/L L-

glutamine and 10% heat-inactivated FBS and drugs were pre-equilibrated in normoxia or

hypoxia (0.8% O_2, 5% CO_2, 70% humidity at 37ºC) for 24 hours prior to the experiment (52).

Cells were introduced to the hypoxic workstation (Baker-Ruskinn Sci-Tive UM-027)

where culture media was removed and replaced with hypoxia-equilibrated media as detailed

above. Cells were treated in duplicate or triplicate wells with 100µg/ml LPS (InvivoGen),

DMSO (Sigma-Aldrich), or 15mM COX-2 inhibitor (NS398) (Cayman Chemical) and

incubated for 18 hours. For normoxic controls, cells were treated identically within a class-2

tissue culture hood and transferred to a normoxic tissue-culture incubator.

Cell supernatants were collected following 18 hours incubation in normoxia or

hypoxia and were frozen for subsequent analysis. Culture supernatants were assayed in

duplicate or triplicate using a human TNF ELISA MAX (Biolegend). In samples where

undiluted supernatants fell outside the standard curve, samples were re-run following 1:10

dilution in assay buffer.

Statistical analysis

All data were analysed (Prism 7.0, GraphPad Software) using unpaired, two-tailed t-

tests for comparisons between two groups and one-way ANOVA (with Bonferonni post-test

adjustment) for other data. P values shown are: *$P < .05$, **$P < .01$, and ***$P < .001$.

Results

***tnfa:GFP* expression is upregulated by injury, infection and Hif-1α stabilisation**

TNF is a central regulator of the M1 inflammatory response to inflammation and

pathogen challenge. Here, we use two transgenic zebrafish lines that express GFP via a *tnfa*

promoter region; *TgBAC(tnfa:GFP)pd1028* and *Tg(tnfa:eGFP-F)ump5Tg* (31, 32). The

TgBAC(tnfa:GFP)pd1028 was made using BAC (bacterial artificial chromosome)

transgenesis and contains 50kb of promoter region (32). The *Tg(tnfa:eGFP-F)ump5Tg* line

has a smaller promoter region (3.8kb) than the BAC line (31) and has previously been

demonstrated to be upregulated in macrophages after tailfin injury in a zebrafish model (31).

Here, we demonstrate that this injury induced upregulation of macrophage *tnfa:GFP* is also

found in the BAC transgenic line (*TgBAC(tnfa:GFP)pd1028* crossed to

Tg(mpeg1:mCherryCAAX)sh378) shown 16 hours post wound (hpw) (Figure 1A). TNF has

been implicated in all stages of TB infection, however the cell types involved *in vivo* have

been difficult to observe (53). We used the *tnfa* BAC promoter GFP line to establish the

transcriptional regulation of *tnfa* during pre- and early- larval Mm granuloma formation

stages. *tnfa:GFP* expression was induced in larvae following Mm infection before granuloma

onset (at 1 day post infection, dpi), and after granuloma formation (at 4dpi) (Figure 1B).

tnfa:GFP expression was predominantly found in macrophages with internalised Mm, shown

by co-localisation of fluorescence with *mpeg-1:mCherry* expressing macrophages and

mCrimson expressing Mm (Figure 1C). Macrophage expression was confirmed using the

other *tnfa* promoter driven line, (*Tg(tnfa:eGFP-F)ump5Tg*, crossed to *Tg(mpeg1:mCherry-*

F)ump2Tg) (Figure S1). Our data demonstrate that injury and Mm induced *tnfa* expression

occurs in macrophages as part of an early M1 response.

Hif-1α can be stabilised in zebrafish both genetically and pharmacologically using

dominant active constructs and hydroxylase inhibitors respectively (29, 44). We tested

whether Hif-1α promotes M1 polarisation using the *tnfa:GFP* BAC transgenic line as a

readout of M1 macrophages (31, 32). Hif-1α was stabilised genetically in larvae using DA

Hif-1α RNA (44), which resulted in upregulation of *tnfa:GFP* expression (Figure 1D-F)

indicating a shift of macrophage phenotype towards an activated M1 response. Dominant

negative (DN) Hif-1α caused no upregulation of *tnfa:GFP* expression (Figure 1E-F). To

demonstrate that it was also possible to boost the Hif-1α M1 pro-inflammatory response

pharmacologically, the hypoxia mimetic DMOG (dimethyloxaloylglycine, a pan-hydroxylase

inhibitor that stabilises Hif-1α via inactivation of regulatory prolyl hydroxylase enzymes) was

used to stabilise endogenous levels of Hif-1α (57). DMOG treatment upregulated expression

of *tnfa*:*GFP* compared to DMSO solvent controls, phenocopying the DA-Hif-1α response

(Figure 1G). Recently described hypoxia mimetics with reportedly greater prolyl hydroxylase

selectively, JNJ-42041935 and FG4592, had similar effects to DMOG (Figure 1H and I) (33,

45, 46). During both homeostasis and disease physiology, Hif-1α stability is regulated by

microenvironmental hypoxia. To simulate this we subjected the *tnfa:GFP* line to 5% oxygen

for 6 hours at 32hpf and looked for GFP expression at 48hpf. This level of hypoxia was

sufficient to switch on GFP expression in the *Tg(phd3:GFP)i144* hypoxia reporter zebrafish

(Figure S2) (33). Six hours of 5% oxygen increased *tnfa:GFP* expression at 48hpf,

compared to normoxic controls, to a similar level as that observed with genetic or

pharmacological Hif-1α stabilisation (Figure 1J). Together, these data indicate that *tnfa*

expression is part of a pro-inflammatory M1 macrophage response to hypoxia and stabilised

Hif-1α, a response that is targetable by pharmacological agents and has the potential to aid

the host response to bacterial challenge.

Hif-1α dependent *tnfa:GFP* transcription requires cyclooxygenase

Eicosanoids are lipid signalling molecules produced by macrophages during early

microbial infection as part of an M1 response (22, 23). We tested whether cyclooxygenase

inhibition affected Hif-1α mediated *tnfa:GFP* production. At basal levels, *tnfa:GFP* was not

altered by the Cox-1 inhibitor SC560 or the Cox-2 inhibitor NS398 compared to DMSO

treated negative controls (Figure 2A-B). Strikingly, DA Hif-1α-induced *tnfa:GFP* expression

was significantly abrogated to basal levels by both SC560 and NS398 (Figure 2A-B). We

have previously shown that DA Hif-1α induces the expression of another important pro-

inflammatory cytokine, *il-1β,* shown by an *il-1β:GFP* transgenic line (Figure S3A) (33). Cox-1

inhibition by SC560 did not abrogate Hif-1α induced *il-1β:GFP*, while Cox-2 inhibition with

NS398 caused a small decrease that did not reach basal levels (Figure S3B-C). These data

indicate that Hif-1α-induced *tnfa* expression is caused by a product of the cyclooxygenase

arm of the arachidonic acid pathway via a novel macrophage HIF/COX/TNF axis (Figure

2C), while Hif-1α induced *il-1β* does not fully act via this pathway suggestive of complex

regulation of M1 pro-inflammatory cytokines by Hif-1α stabilisation.

We next tested whether injury- and infection-induced *tnfa* are cyclooxygenase

dependent processes. Macrophage *tnfa:GFP* expression induced after injury was not

abrogated by cyclooxygenase inhibition using NS398 (Figure 2D-E). Similarly,

cyclooxygenase inhibition using either SC560 or NS398 did not alter the expression of Mm-

induced *tnfa:GFP* (Figure 2F-G). Our data indicate that macrophage *tnfa* expression can be

driven by the presence of DAMPS (injury) or PAMPS (Mm) and that these are

cyclooxygenase independent mechanisms (Figure 2H).

We have previously shown that Hif-1α stabilisation is host-protective in the zebrafish

Mm model, an effect that is dependent on macrophage expression of proinflammatory *il-1β*

(33). We therefore hypothesised that priming macrophages with increased *tnfa* via Hif-1α

stabilisation may be protective against subsequent Mm infection. Neither genetic nor

pharmacological Hif-1α stabilisation were additive to the *tnfa:GFP* expression caused by

either injury or Mm infection (Figure 2C-F and S4A-D). To test whether DA Hif-1α priming of

macrophage *tnfa* had an effect on the outcome of Mm infection, bacterial burden was

quantified in DA Hif-1α-injected embryos with cyclooxygenase inhibition (having

demonstrated that Hif-1α-induced *tnfa* requires cyclooxygenase activity in Figure 2A-C).

Neither SC560 nor NS398 mediated inhibition of Hif-1α-induced *tnfa* expression diminished

the host-protective effect of DA Hif-1α (Figure S5A-C) or DMOG treated larvae (Figure S5D-

F). Taken together these data indicate that injury- and Mm- induced *tnfa* is not further

increased by Hif-1α stabilisation and that cyclooxygenase mediated early priming of *tnfa* in

macrophages is not required for the Hif-1α mediated reduction of bacterial burden.

Blocking cyclooxygenase independent arachidonic acid pathways does not abrogate

Hif-1α upregulation of *tnfa:GFP*

To investigate whether the effect of the cyclooxygenase inhibitors on *tnfa* was

specific to the prostaglandin path of the arachidonic acid pathway, we targeted the lipoxin

and leukotriene producing arms using the 15-Lipoxygenase inhibitor PD146176 and

leukotriene B4 receptor antagonists. The 15-Lipoxygenase inhibitor PD146176 did not block

the *tnfa:GFP* induced by DA Hif-1α (Figure 3A-B). Furthermore, PD147176 increased

tnfa:GFP expression on its own in the absence of infection, although not to the same extent

as DA Hif-α (Figure 3A-B). PD146176 also did not affect *tnfa:GFP* expression after Mm

infection (Figure 3A-B), nor did it block the protective effect of DA Hif-1α. Treatment with

PD146176 was sufficient to decrease Mm burden, but not to the same extent as DA Hif-1α

(Figure 3C). Leukotriene B4 inhibition, using the BLTR1/2 antagonists U75302 and

LY255283, did not increase *tnfa:GFP* levels and also did not block DA Hif-1α induced

tnfa:GFP (Figure 3D-E). These data indicate that blocking components of the lipoxygenase

dependent arms of the arachidonic acid pathway do not block the Hif-1α effect on *tnfa*

expression and do not replicate the effects observed by blocking the cyclooxygenase

dependent arm (Figure 3F).

Hif-1α-induced *tnfa:GFP* requires active prostaglandin E2

A key family of immune regulators downstream of arachidonic acid and cyclooxygenases are

prostaglandins. The best characterised of these as a regulator of macrophage function is

prostaglandin E2 (PGE2). We tested whether PGE2 was a mediator in the HIF/COX/TNF

pathway by addition of exogenous PGE2 to DA Hif-1α larvae. Exogenous PGE2 had no

effect on *tnfa:GFP* expression in the absence or presence of Mm infection (Figure 4A-D).

However, PGE2 was able to rescue the decrease in *tnfa:GFP* expression after Cox-1 (Figure

4A and C) or Cox-2 (Figure 4B and D) inhibition in the DA Hif-1α larvae. Furthermore,

addition of PGE2 alone led to a significant increase in the DA Hif-1α-induced *tnfa:GFP*

expression (Figure 4A-D). These rescuing effects were not observed by addition of

exogenous 15-keto prostaglandin E2, an immunologically inactive degradation product of

PGE2 (Figure 4E-F) (49, 58, 59). These data indicate that Hif-1α-induced *tnfa:GFP* requires

active prostaglandin E2 (Figure 4G).

The HIF-COX-TNF axis is conserved in human macrophages

To translate our findings from zebrafish to humans we tested whether HIF-1α

stabilisation in human macrophages induces TNF expression. We found that human

monocyte derived macrophages (hMDMs) produced higher levels of TNF protein in hypoxia

(0.8% oxygen) than those in normoxia, measured by an anti-human TNF ELISA (Figure 5A).

Furthermore, treatment with the COX-2 inhibitor, NS398, reduced this hypoxia-induced TNF

back to the equivalent levels found in normoxia (Figure 5A). This was replicated when HIF-

1α was stabilised in hHDMs using the hypoxia mimetic FG4592 (Figure 5B). These data

indicate that the HIF/COX/TNF axis is a conserved mechanism in human macrophages and

could be important in human disease.

Discussion

Control of macrophage function during homeostasis and infection are critical for

healthy tissues and must integrate changes in the local microenvironment with cytokine

signalling. Understanding of signalling pathways that link the microenvironment with

macrophage phenotypic outcomes will identify potential novel therapeutic avenues for

control of macrophages during disease. Currently, the mechanisms and molecular cues

leading to different macrophage phenotypes are not well-defined *in vivo*.

Here we show that a disease relevant microenvironmental cue, hypoxia signalling via

Hif-1α, upregulates macrophage *tnfa* expression in a cyclooxygenase dependent manner *in*

vivo. TNF regulation by hypoxia has been shown in a range of mammalian cells, and its

promoter region contains HIF responsive elements (HREs), resulting in some level of direct

regulation by HIF-1α (60–63). We observed an M1 pro-inflammatory *tnfa* response with

stabilised Hif-1α, demonstrating that hypoxia signalling alone can lead to a switch of

macrophage polarisation to an M1 phenotype, consistent with our previous observations

concerning *il-1β* activation (33). Not only could *tnfa* activation be achieved by genetic

stabilisation of Hif-1α, but also using hypoxia mimetics and physiological hypoxia. These

findings indicate that the Hif-1α pro-inflammatory switch is targetable by pharmaceuticals

and could be druggable during disease. Hypoxic regulation of TNF via COX has previously

been demonstrated in mammalian osteoblasts, however, little is known about this interaction

(26). Our data demonstrate that Hif-1α upregulation of macrophage *tnfa* is dependent on

cyclooxygenase and further shows that the mechanism is likely to be via the production of

PGE2. The degradation metabolite 15-keto-PGE2 did not rescue the loss of *tnfa* expression

after cyclooxygenase inhibition, consistent with previous reports that 15-keto-PGE2 is

unable to bind the prostaglandin E2 EP receptors, demonstrating a requirement for active

PGE2 (64).

Mycobacteria mediated macrophage polarisation is complex, with M1 and M2

phenotypes observed during pathogenesis, however, the mechanisms determining

macrophage phenotype are not fully characterised (56). The Mtb/Mm granuloma has been

widely studied in human, mice and zebrafish and is rich in pro-inflammatory cytokine

production (32, 33, 65–69). Here, we observed that an M1 pro-inflammatory *tnfa* response

was induced at pre-granuloma stages as well as during granuloma formation *in vivo*. A key

advantage of using fluorescent transgenic zebrafish lines, such as *tnfa:GFP*, is that they

enable identification of the cell type producing the transgene in an intact organism. This is

especially important when studying early innate immune cell interactions with pathogens, as

it eliminates the risk of activation during a sorting process. TNF is required for the control of

initial TB infection in mice and also of latent TB in humans (68, 70–73). Patients on anti-TNF

therapies for immune diseases such as Crohn's disease and rheumatoid arthritis, which

have proved effective treatments for these debilitating illnesses, have an associated

increased risk of TB reactivation and infectious disease (72, 74). In the zebrafish Mm model

it has been widely demonstrated that Tnfa is required for control of early infection, with

perturbation of Tnfa signalling leading to high infectious burdens (75). When Hif-1α driven

tnfa was downregulated by cyclooxygenase inhibition, there was no effect on the decrease

in bacterial burden resulting from stabilisation of Hif-1α. Furthermore, none of the

arachidonic acid pathway inhibitors used in our study had any effect on Tnfa production after

Mm infection. These findings are consistent with Mm induced macrophage Tnfa production

being independent of arachidonic acid, and is most likely TLR mediated as has been

reported elsewhere (43, 56). Our data indicate that there is no additional protective effect of

priming macrophages with increased levels of Tnfa before infection onset. Interestingly, this

is not the case for another important M1 cytokine, Il-1β, where Hif-1α stabilisation fails to

reduce bacterial burden if Il-1β is blocked early in infection, (33). Our findings indicate that

different M1 cytokines induced by Hif-1α stabilisation can have differential effects on the

outcome of infection, and highlight that, although TNF and IL-1β are often used as markers

of M1 polarisation, their roles in the pro-inflammatory macrophage are distinct.

Blocking cyclooxygenase independent arms of the arachidonic acid pathway did not

inhibit Hif-1α-induced macrophage *tnfa* transcription in the same way as blocking the

cyclooxygenase/PGE2 arm. Inhibition of the 15-lipoxygenase arm was sufficient to induce

macrophage *tnfa* transcription which was host protective and decreased Mm burden. These

findings indicate a potential shift towards the COX/TNF dependent axis when 15-

lipoxygenase is blocked and are consistent with previous observations suggesting that

inhibition of 15-Lipoxygenase can have immune activating effects (48).

Our data reveal a novel mechanism of TNF activation in macrophages via a Hif-1α/

cyclooxygenase/PGE2 axis. Importantly, this pathway is conserved in human macrophages,

meaning that it could be targetable for pharmaceutical intervention to promote macrophage

M1-type polarisation in human disease. Although the effect of the HIF/COX/TNF axis in early

Mm infection was negligible, this pathway is likely to have important roles in numerous

disease situations. Multiple disease pathologies have hypoxia as a features of the tissue

microenvironment, concurrent with macrophage influx. Hif-1α mediated M1 switching and

TNF activation may be pertinent in later TB infection situations where granulomas have

stabilised Hif-1α due to necrotic and hypoxic centres (76, 77). It is not only bacterial

infections such as TB that might be influenced by a HIF-1α mediated M1/TNF switch.

Hypoxia is a key hallmark of cancers with high levels of HIF-1α widely found in those that

produce large tumours where the centre is hypoxic (78). This is also true of

cyclooxygenase/PGE2 production, where these important immunomodulatory signals are

required for cellular adaption to the tumour microenvironment (79). Certain types of cancer

induce inflammatory microenvironments, where cytokines and other immunomodulators can

contribute to the progression of cancers (80). Macrophages play central roles in cancer-

inflammation that could be exploited utilising pharmaceutical control of this novel

HIF/COX/TNF mechanism. The zebrafish larvae is small enough to be fully oxygenated at

the 2-5dpf stages of this study (81). Further investigation of the HIF/COX/TNF axis in models

where hypoxia is a key hallmark of disease pathology is required to uncover the full

therapeutic relevance of this potentially important novel macrophage pathway.

In conclusion, we have identified a novel mechanism for macrophage *tnfa*

transcription via Hif-1α and cyclooxygenase that is conserved from zebrafish to humans,

highlighting a druggable mechanism to induce an M1-macrophage response. Importantly,

this axis links a microenvironmental cue, hypoxia, to a macrophage pro-inflammatory M1

cytokine, *tnfa* expression. We provide strong evidence to show that this response acts via

the cyclooxygenase/PGE2 arm of the arachidonic pathway. Due to the central roles of these

modulators in disease microenvironments we anticipate that this HIF/COX/TNF pathway

may have important implications in conditions such as inflammation, infection and cancer.

References

1.　　Benoit M, Desnues B, Mege J-L (2008) Infections Macrophage Polarization in

Bacterial Macrophage Polarization in Bacterial Infections. *J Immunol Ref J Immunol*

181:3733–3739.

2.   Wiegertjes GF, Wentzel AS, Spaink HP, Elks PM, Fink IR (2016) Polarization of

immune responses in fish: The 'macrophages first' point of view. *Mol Immunol* 69.

doi:10.1016/j.molimm.2015.09.026.

3.   Murray PJ, et al. (2014) Macrophage Activation and Polarization: Nomenclature and

Experimental Guidelines. *Immunity* 41(1):14–20.

4.   Shapouri-Moghaddam A, et al. (2018) Macrophage plasticity, polarization, and

function in health and disease. *J Cell Physiol* 233(9):6425–6440.

5.   Nizet V, Johnson RS (2009) Interdependence of hypoxic and innate immune

responses. *Nat Rev Immunol* 9(9):609–617.

6.   Weiss G, Schaible UE (2015) Macrophage defense mechanisms against intracellular

bacteria. *Immunol Rev* 264(1):182–203.

7.   Mortaz E, et al. (2015) Interaction of Pattern Recognition Receptors with

Mycobacterium Tuberculosis. *J Clin Immunol* 35(1). doi:10.1007/s10875-014-0103-7.

8.   Huang Z, et al. (2015) Mycobacterium tuberculosis-induced polarization of human

macrophage orchestrates the formation and development of tuberculous granulomas

in vitro. *PLoS One* 10(6). doi:10.1371/journal.pone.0129744.

9.   Qin Y, et al. (2015) Macrophage-Microglia Networks Drive M1 Microglia Polarization

After Mycobacterium Infection. *Inflammation* 38(4):1609–1616.

10.   Hussain Bhat K, Mukhopadhyay S (2015) Macrophage takeover and the host-bacilli

interplay during tuberculosis. *Future Microbiol* 10(5):853–872.

11.   Wang J, et al. (2015) Mycobacterium tuberculosis suppresses innate immunity by

coopting the host ubiquitin system. *Nat Immunol* 16(3):237–245.

12.   Ehrt S, et al. (2001) Reprogramming of the macrophage transcriptome in response to

interferon-gamma and Mycobacterium tuberculosis: signaling roles of nitric oxide

synthase-2 and phagocyte oxidase. *J Exp Med* 194(8):1123–1140.

13.   Roy S, et al. (2018) Transcriptional landscape of Mycobacterium tuberculosis infection

in macrophages. *Sci Rep* 8(1). doi:10.1038/s41598-018-24509-6.

14.   Dorhoi A, Kaufmann SHE (2015) Versatile myeloid cell subsets contribute to

tuberculosis-associated inflammation. *Eur J Immunol* 45(8):2191–2202.

15.   McClean CM, Tobin DM (2016) Macrophage form, function, and phenotype in

mycobacterial infection: Lessons from tuberculosis and other diseases. *Pathog Dis*

74(7). doi:10.1093/femspd/ftw068.

16.   Wang N, Liang H, Zen K (2014) Molecular mechanisms that influence the

macrophage M1-M2 polarization balance. *Front Immunol* 5(NOV).

doi:10.3389/fimmu.2014.00614.

17.   Cramer T, et al. (2003) HIF-1α is essential for myeloid cell-mediated inflammation.

Cell 112(5):645–657.

18.   Peyssonnaux C, et al. (2007) Cutting edge: Essential role of hypoxia inducible factor-

1alpha in development of lipopolysaccharide-induced sepsis. *J Immunol*

178(12):7516–7519.

19.   Kaidi A, Qualtrough D, Williams AC, Paraskeva C (2006) Direct transcriptional up-

regulation of cyclooxygenase-2 by hypoxia-inducible factor (HIF)-1 promotes

colorectal tumor cell survival and enhances HIF-1 transcriptional activity during

hypoxia. *Cancer Res* 66(13):6683–6691.

20.   Lee JJ, et al. (2010) Hypoxia activates the cyclooxygenase-2-prostaglandin E

synthase axis. *Carcinogenesis* 31(3):427–434.

21.   Zhao L, et al. (2012) Involvement of COX-2/PGE2signalling in hypoxia-induced

angiogenic response in endothelial cells. *J Cell Mol Med* 16(8):1840–1855.

22.   Norris PC, Reichart D, Dumlao DS, Glass CK, Dennis EA (2011) Specificity of

eicosanoid production depends on the TLR-4-stimulated macrophage phenotype. *J*

Leukoc Biol 90(3):563–574.

23.   Gupta S, Maurya MR, Stephens DL, Dennis EA, Subramaniam S (2009) An

integrated model of eicosanoid metabolism and signaling based on lipidomics flux

analysis. *Biophys J* 96(11):4542–4551.

24. Harizi H (2013) The immunobiology of prostanoid receptor signaling in connecting

innate and adaptive immunity. *Biomed Res Int* 2013. doi:10.1155/2013/683405.

25. Mancini AD, Di Battista JD (2011) The cardinal role of the phospholipase

A(2)/cyclooxygenase-2/prostaglandin E synthase/prostaglandin E(2) (PCPP) axis in

inflammostasis. *Inflamm Res* 60(12):1083–92.

26. Xing Y, et al. (2015) COX2 is involved in hypoxia-induced TNF-α expression in

osteoblast. *Sci Rep* 5(1):10020.

27. Meijer AH (2016) Protection and pathology in TB: learning from the zebrafish model.

Semin Immunopathol 38(2):261–273.

28. Henry KM, Loynes C a, Whyte MKB, Renshaw S a (2013) Zebrafish as a model for

the study of neutrophil biology. *J Leukoc Biol* 94(4):633–42.

29. Elks PM, Renshaw SA, Meijer AH, Walmsley SR, van Eeden FJ (2015) Exploring the

HIFs, buts and maybes of hypoxia signalling in disease: lessons from zebrafish

models. *Dis Model Mech* 8(11):1349–1360.

30. Loynes CA, et al. (2010) Pivotal Advance: Pharmacological manipulation of

inflammation resolution during spontaneously resolving tissue neutrophilia in the

zebrafish. *J Leukoc Biol* 87(2):203–212.

31. Nguyen-Chi M, et al. (2015) Identification of polarized macrophage subsets in

zebrafish. *Elife* 4(JULY 2015). doi:10.7554/eLife.07288.

32. Marjoram L, et al. (2015) Epigenetic control of intestinal barrier function and

inflammation in zebrafish. *Proc Natl Acad Sci U S A* 112(9):2770–5.

33. Ogryzko N, et al. (2018) Macrophage Il-1beta protects against mycobacterial infection

downstream of Hif-1alpha in zebrafish. *BioRxiv* 1(2):2000–2000.

34. Nusslein-Volhard C DR (2002) *Zebrafish: A Practical Approach* (Oxford University

Press, Oxford). 1st ed.

35. Bojarczuk A, et al. (2016) Cryptococcus neoformans Intracellular Proliferation and

Capsule Size Determines Early Macrophage Control of Infection. *Sci Rep* 6(1):21489.

36. Hall C, Flores M V, Storm T, Crosier K, Crosier P (2007) The zebrafish lysozyme C

promoter drives myeloid-specific expression in transgenic fish. *BMC Dev Biol* 7:42.

37.   Santhakumar K, et al. (2012) A zebrafish model to study and therapeutically

551   manipulate hypoxia signaling in tumorigenesis. *Cancer Res* 72(16):4017–4027.

552   38.   Elks PM, Loynes CA, Renshaw SA (2011) *Measuring inflammatory cell migration in*

553   *the zebrafish* doi:10.1007/978-1-61779-207-6_18.

554   39.   van der Sar AM, Spaink HP, Zakrzewska A, Bitter W, Meijer AH (2009) Specificity of

555   the zebrafish host transcriptome response to acute and chronic mycobacterial

556   infection and the role of innate and adaptive immune components. *Mol Immunol*

557   46(11–12):2317–2332.

558   40.   Cui C, et al. (2011) Infectious disease modeling and innate immune function in

559   zebrafish embryos. *Methods Cell Biol* 105:273–308.

560   41.   Benard EL, et al. (2012) Infection of zebrafish embryos with intracellular bacterial

561   pathogens. *J Vis Exp* (61). doi:10.3791/37813781 [pii].

562   42.   Elks PM, et al. (2013) PLOS Pathogens: Hypoxia Inducible Factor Signaling

563   Modulates Susceptibility to Mycobacterial Infection via a Nitric Oxide Dependent

564   Mechanism. *PLoS Pathog* 9(12):e1003789.

565   43.   Elks PM, et al. (2014) Mycobacteria counteract a TLR-mediated nitrosative defense

566   mechanism in a zebrafish infection model. *PLoS One* 9(6).

567   doi:10.1371/journal.pone.0100928.

568   44.   Elks PM, et al. (2011) Activation of hypoxia-inducible factor-1alpha (Hif-1alpha) delays

569   inflammation resolution by reducing neutrophil apoptosis and reverse migration in a

570   zebrafish inflammation model. *Blood* 118(3):712–722.

571   45.   Barrett TD, et al. (2011) Pharmacological Characterization of 1-(5-Chloro-6-

572   (trifluoromethoxy)-1H-benzoimidazol-2-yl)-1H-pyrazole-4-carboxylic Acid (JNJ-

573   42041935), a Potent and Selective Hypoxia-Inducible Factor Prolyl Hydroxylase

574   Inhibitor. *Mol Pharmacol* 79(6):910–920.

575   46.   Wu K, et al. (2016) Stabilization of HIF-1α by FG-4592 promotes functional recovery

576   and neural protection in experimental spinal cord injury. *Brain Res* 1632:19–26.

47. Feng Y, Renshaw S, Martin P (2012) Live imaging of tumor initiation in zebrafish

larvae reveals a trophic role for leukocyte-derived PGE2. *Curr Biol* 22(13):1253–1259.

48. Tobin DM, et al. (2010) The lta4h locus modulates susceptibility to mycobacterial

infection in zebrafish and humans. *Cell* 140(5):717–730.

49. Evans RJ, et al. (2017) Fungal derived 15-keto-prostaglandin E_2 activates host

peroxisome proliferator-activated receptor gamma (PPAR-γ) to promote C.

neoformans growth during infection. *bioRxiv*:113167.

50. Stoop EJ, et al. (2011) Zebrafish embryo screen for mycobacterial genes involved in

the initiation of granuloma formation reveals a newly identified ESX-1 component. *Dis*

Model Mech 4(4):526–536.

51. Dockrell DH, Lee M, Lynch DH, Read RC (2001) Immune-Mediated Phagocytosis and

Killing of *Streptococcus pneumoniae* Are Associated with Direct and Bystander

Macrophage Apoptosis. *J Infect Dis*. doi:10.1086/323084.

52. Ghezzi P, et al. (1991) Hypoxia increases production of interleukin-1 and tumor

necrosis factor by human mononuclear cells. *Cytokine*. doi:10.1016/1043-

4666(91)90015-6.

53. Tobin DM, et al. (2012) Host genotype-specific therapies can optimize the

inflammatory response to mycobacterial infections. *Cell* 148(3):434–446.

54. Van Der Vaart M, et al. (2014) The DNA damage-regulated autophagy modulator

DRAM1 links mycobacterial recognition via TLP-MYD88 to authophagic defense. *Cell*

Host Microbe 15(6):753–767.

55. Berod L, et al. (2014) MyD88 signalling in myeloid cells is sufficient to prevent chronic

mycobacterial infection. *Eur J Immunol* 44(5):1399–1409.

56. Basu J, Shin D-M, Jo E-K (2012) Mycobacterial signaling through toll-like receptors.

Front Cell Infect Microbiol 2. doi:10.3389/fcimb.2012.00145.

57. Elks PM, et al. (2011) Activation of hypoxia-inducible factor-1α (hif-1α) delays

inflammation resolution by reducing neutrophil apoptosis and reverse migration in a

zebrafish inflammation model. *Blood* 118(3). doi:10.1182/blood-2010-12-324186.

58.   Chen IJ, et al. (2018) Targeting the 15-keto-PGE2-PTGR2 axis modulates systemic

606   inflammation and survival in experimental sepsis. *Free Radic Biol Med* 115:113–126.

607   59.   Tai HH, Ensor CM, Tong M, Zhou H, Yan F (2002) Prostaglandin catabolizing

608   enzymes. *Prostaglandins Other Lipid Mediat* 68–69:483–493.

609   60.   Ghosh S, Paul A, Sen E (2013) Tumor Necrosis Factor Alpha-Induced Hypoxia-

610   Inducible Factor 1α-β-Catenin Axis Regulates Major Histocompatibility Complex Class

611   I Gene Activation through Chromatin Remodeling. *Mol Cell Biol* 33(14):2718–31.

612   61.   Charbonneau M, et al. (2007) Hypoxia-inducible factor mediates hypoxic and tumor

613   necrosis factor alpha-induced increases in tumor necrosis factor-alpha converting

614   enzyme/ADAM17 expression by synovial cells. *J Biol Chem* 282(46):33714–24.

615   62.   Imtiyaz HZ, Simon MC (2010) Hypoxia-inducible factors as essential regulators of

616   inflammation. *Curr Top Microbiol Immunol* 345(1):105–120.

617   63.   Sandau KB, Zhou J, Kietzmann T, Brüne B (2001) Regulation of the Hypoxia-

618   inducible Factor 1α by the Inflammatory Mediators Nitric Oxide and Tumor Necrosis

619   Factor-α in Contrast to Desferroxamine and Phenylarsine Oxide. *J Biol Chem*

620   276(43):39805–39811.

621   64.   Coggins KG, et al. (2002) Metabolism of PGE2by prostaglandin dehydrogenase is

622   essential for remodeling the ductus arteriosus. *Nat Med* 8(2):91–92.

623   65.   Ulrichs T, Kaufmann SH (2006) New insights into the function of granulomas in

624   human tuberculosis. *J Pathol* 208(2):261–269.

625   66.   Flynn JL, Chan J, Lin PL (2011) Macrophages and control of granulomatous

626   inflammation in tuberculosis. *Mucosal Immunol* 4(3):271–278.

627   67.   Via LE, et al. (2008) Tuberculous granulomas are hypoxic in guinea pigs, rabbits, and

628   nonhuman primates. *Infect Immun* 76(6):2333–2340.

629   68.   Orphanidou D, et al. (1996) Tumour necrosis factor, interleukin-1 and adenosine

630   deaminase in tuberculous pleural effusion. *Respir Med* 90(2):95–98.

631   69.   Benard EL, Rougeot J, Racz PI, Spaink HP, Meijer AH (2016) Transcriptomic

632   Approaches in the Zebrafish Model for Tuberculosis—Insights Into Host- and

Pathogen-specific Determinants of the Innate Immune Response. *Adv Genet* 95:217–

251.

70. Flynn JL, et al. (1995) Tumor necrosis factor-alpha is required in the protective

immune response against Mycobacterium tuberculosis in mice. *Immunity* 2(6):561–

72.

71. Kindler V, Sappino AP, Grau GE, Piguet PF, Vassalli P (1989) The inducing role of

tumor necrosis factor in the development of bactericidal granulomas during BCG

infection. *Cell* 56(5):731–740.

72. Ormerod LP (2004) Tuberculosis and anti-TNF-alpha treatment. *Thorax*

59(11):921,2004.

73. Wallis RS. d, Broder MS., Wong JY., Hanson ME., Beenhouwer DO. (2004)

Granulomatous infectious diseases associated with tumor necrosis factor antagonists.

Clin Infect Dis 38(9):1261–1265.

74. Xie X, Li F, Chen J-W, Wang J (2014) Risk of tuberculosis infection in anti-TNF-α

biological therapy: From bench to bedside. *J Microbiol Immunol Infect* 47(4):268–274.

75. Clay H, Volkman HE, Ramakrishnan L (2008) Tumor necrosis factor signaling

mediates resistance to mycobacteria by inhibiting bacterial growth and macrophage

death. *Immunity* 29(2):283–294.

76. Duque-Correa MA, et al. (2014) Macrophage arginase-1 controls bacterial growth and

pathology in hypoxic tuberculosis granulomas. *Proc Natl Acad Sci* 111(38):E4024–

E4032.

77. Qualls JE, Murray PJ (2016) Immunometabolism within the tuberculosis granuloma:

amino acids, hypoxia, and cellular respiration. *Semin Immunopathol* 38(2):139–152.

78. Ajdukovic J (2016) HIF-1 - A big chapter in the cancer tale. *Exp Oncol* 38(1):9–12.

79. Greenhough A, et al. (2009) The COX-2/PGE2 pathway: key roles in the hallmarks of

cancer and adaptation to the tumour microenvironment. *Carcinogenesis* 30(3):377–

386.

80. Landskron G, De La Fuente M, Thuwajit P, Thuwajit C, Hermoso MA (2014) Chronic

inflammation and cytokines in the tumor microenvironment. *J Immunol Res* 2014.

doi:10.1155/2014/149185.

81.   Oehlers SH, et al. (2015) Interception of host angiogenic signalling limits

mycobacterial growth. *Nature.* doi:10.1038/nature13967.

Acknowledgements

The authors would like to thank The Bateson Aquarium Team for fish care and the IICD

Technical Team for practical assistance (University of Sheffield). We gratefully thank, Michel

Bagnat (Duke University) for providing their *TgBAC(tnfa:GFP)pd1028* transgenic line,

Georges Lutfalla (Montpellier University) for providing the *Tg(tnfa:eGFP-F)ump5Tg* line,

Annemarie Meijer (University of Leiden) for M. marinum strains. We thank David Dockrell

(University of Edinburgh), Paul Collini (University of Sheffield) and Jon Kilby (University of

Sheffield) for help with hMDM experiments. Thanks to Alison Condliffe and Benjamin

Durham (University of Sheffield) for use of, and invaluable help with, their hypoxia hood. We

would like to thank Robert Evans (The Francis Crick, London) and Catherine Loynes

(University of Sheffield) for helpful discussions and advice on prostaglandin modulation.

Thanks to Simon Johnston, Hannah Isles and Stephen Renshaw (University of Sheffield) for

constructive comments on the manuscript.

Funding

AL and PME are funded by a Sir Henry Dale Fellowship jointly funded by the Wellcome

Trust and the Royal Society (Grant Number 105570/Z/14/Z) held by PME.

Author Contributions

Conceived and designed the experiments: PME, AL. Performed the experiments: PME, AL.

Analysed the data: PME, AL. Wrote the paper: PME.

Figure Legends

Figure 1. Macrophage *tnfa:GFP* is upregulated by injury, Mm infection and Hif-1α stabilisation

(A) Fluorescent confocal micrographs of 3 days post fertilisation larvae with or without tailfin wound (induced 16 hours previously) with example macrophages at the tailfin site. *tnfa* expression was detected by GFP levels, in green, using the *TgBAC(tnfa:GFP)pd1028* transgenic line. Macrophages are shown in red using a *Tg(mpeg1:mCherryCAAX)sh378* line. The asterisk in the not wounded control indicates a neuromast that is fluorescent in all channels as a marker of position.

 (B) Fluorescent confocal micrographs of 1dpi Mm infected larvae, prior to granuloma formation (left panels), and 4 dpi, early granuloma (right panels) stages of infection. *tnfa* expression was detected by GFP levels, in green, using the *TgBAC(tnfa:GFP)pd1028* transgenic line. Mm mCrimson is shown in the far-red (cyan) channel. Increased levels of *tnfa:GFP* expression were detectable in cells associated with infection.

(C) Fluorescent confocal micrographs of 1 day post infection (dpi) caudal vein region of infection. *tnfa* expression was detected by GFP levels, in green, using the *TgBAC(tnfa:GFP)pd1028* transgenic line. Macrophages are shown in red using a *Tg(mpeg1:mCherryCAAX)sh378* line. Mm Crimson is shown in the far-red channel (cyan, right panels) with a PVP control (left panels). Without infection there is little detectable *tnfa:GFP* expression, while infected and uninfected macrophages have higher levels of *tnfa:GFP* in the Mm group. Dotted lines indicate the yolk extension of the larvae where there is non-specific fluorescence.

(D) Fluorescent confocal micrographs of 2 days post fertilisation larvae in the caudal vein region. *tnfa* expression was detected by GFP levels, in green, using the *TgBAC(tnfa:GFP)pd1028* transgenic line. Macrophages are shown in red using a *Tg(mpeg1:mCherryCAAX)sh378* line. Larvae were injected at the 1 cell stage with dominant active (DA) Hif-1α or phenol red (PR) control.

(E) Fluorescent confocal micrographs of 2dpf zebrafish imaged around the caudal vein

region. *tnfa* expression was detected by GFP levels, in green, using the

TgBAC(tnfa:GFP)pd1028 transgenic line. Larvae were injected at the 1 cell stage with

dominant negative (DN) or dominant active (DA) Hif-1α or phenol red (PR) control. Dotted

lines indicate the yolk extension of the larvae where there is non-specific fluorescence.

(F) Graph shows corrected fluorescence intensity levels of *tnfa:GFP*. Dominant active Hif-1α

(DA1, filled bar) had significantly increased *tnfa:GFP* levels compared to phenol red (PR)

and dominant negative Hif-1α (DN1) injected controls (white bars). Data shown are mean ±

SEM, n=66 cells from 12 embryos accumulated from 3 independent experiments and

corresponds to images in (E).

(G) Fluorescent confocal micrographs of 2dpf *TgBAC(tnfa:GFP)pd1028* transgenic larvae

treated with DMOG or DMSO control. Dotted lines indicate the yolk extension of the larvae

where there is non-specific fluorescence. Graph shows corrected fluorescence intensity

levels of *tnfa:GFP* confocal z-stacks in larvae. DMOG treated larvae (filled bars) had

significantly increased *tnfa:GFP* levels compared to DMSO controls (white bars). Data

shown are mean ± SEM, n=24 cells from 6 embryos representative of 2 independent

experiments.

(G) Fluorescent confocal micrographs of 2dpf *TgBAC(tnfa:GFP)pd1028* transgenic larvae

treated with the PHD inhibitor JNJ-42041935 or a DMSO solvent control. Dotted lines

indicate the yolk extension of the larvae where there is non-specific fluorescence. Graph

shows corrected fluorescence intensity levels of *tnf-a:GFP* confocal z-stacks at 2dpf treated

with JNJ-42041935 (filled bars) or DMSO controls (white bars). Data shown are mean ±

SEM, n=48 cells from 8 embryos accumulated from 2 independent experiments.

(H) Fluorescent confocal micrographs of 2dpf *TgBAC(tnfa:GFP)pd1028* transgenic larvae

treated with the PHD inhibitor FG4592 or a DMSO solvent control. Dotted lines indicate the

yolk extension of the larvae where there is non-specific fluorescence. Graph shows

corrected fluorescence intensity levels of *tnfa:GFP* confocal z-stacks in larvae at 2dpf

treated with FG4592 (filled bars) or DMSO controls (white bars). Data shown are mean ±

SEM, n=54 cells from 9 embryos accumulated from 3 independent experiments.

(I) Fluorescent confocal micrographs of 2dpf *TgBAC(tnfa:GFP)pd1028* transgenic larvae

incubated in room normoxia or 5% oxygen (hypoxia) for 6 hours between 32-38hpf. Dotted

lines indicate the yolk extension of the larvae where there is non-specific fluorescence.

Graph shows corrected fluorescence intensity levels of *tnfa:GFP* confocal z-stacks in larvae

at 2dpf incubated in 5% oxygen (filled bars) or normoxia controls (white bars). Data shown

are mean ± SEM, n=72 cells from 12 embryos accumulated from 3 independent

experiments.

Figure 2. Hif-1α-activated *tnfa:GFP* is cyclooxygenase dependent while injury and

infection induced *tnfa:GFP* is cyclooxygenase independent

(A) Fluorescent confocal micrographs of 2dpf caudal vein region of larvae. *tnfa* expression

was detected by GFP levels, in green, using the *TgBAC(tnfa:GFP)pd1028* transgenic line.

Phenol red (PR) and dominant active Hif-1α (DA1) injected larvae were treated with DMSO,

SC560 (Cox-1 inhibitor) and NS398 (Cox-2 inhibitor). Dotted lines indicate the yolk extension

of the larvae where there is non-specific fluorescence.

(B) Corrected fluorescence intensity levels of *tnfa:GFP* confocal z-stacks in PR (white bars)

and DA1 (filled bars) injected larvae at 2dpf treated with Cox inhibitors. Data shown are

mean ± SEM, n=36 cells from 6 embryos representative of 3 independent experiments.

(C) Schematic of the arachidonic pathway indicating where stabilising Hif-1α upregulates

tnfa via cyclooxygenase, an effect that is blocked using the Cox1/2 inhibitors SC560/NS398.

(D) Fluorescent confocal micrographs of 3 days post fertilisation larvae with or without tailfin

wound (induced 16 hours previously) in phenol red (PR) or dominant active Hif-1α embryos.

tnfa expression was detected by GFP levels, in green, using the *TgBAC(tnfα:GFP)pd1028*

transgenic line. Embryos were treated with DMSO or NS398 (Cox-2 inhibitor).

(E) Corrected fluorescence intensity levels of *tnfa:GFP* confocal z-stacks in PR (white bars)

and DA1 (filled bars) injected larvae at 2dpf treated with Cox inhibitors. Data shown are

mean ± SEM, n=24 (in not wounded) or n=36 (in wounded) cells from 6 embryos

accumulated from 2 independent experiments. Not-wounded fish had fewer macrophages at

the tailfin site, hence the lower cell number in these groups.

(F) Fluorescent confocal micrographs of 1dpi caudal vein region of Mm Crimson infected

larvae. *tnfa* expression was detected by GFP levels, in green, using the

TgBAC(tnfa:GFP)pd1028 transgenic line. Phenol red (PR) and dominant active Hif-1α (DA1)

injected larvae were treated with DMSO, SC560 (Cox-1 inhibitor) and NS398 (Cox-2

inhibitor) and infected with Mm Crimson. Dotted lines indicate the yolk extension of the

larvae where there is non-specific fluorescence.

(G) Corrected fluorescence intensity levels of *tnfa:GFP* confocal z-stacks in PR (white bars)

and DA1 (filled bars) Mm Crimson infected larvae at 1dpi treated with Cox inhibitors. NB,

DMSO NoMm is a no Mm control taken from figure 2B for comparison with Mm infected data

(data is from the same experiment). Data shown are mean ± SEM, n=36 cells from 6

embryos representative of 3 independent experiments.

(H) Schematic showing that injury and infection upregulate *tnfa* independently of

cyclooxygenase, an effect that is not blocked by the Cox1/2 inhibitors SC560/NS398.

Figure 3. Blocking 15-lipoxygenase or leukotriene B4 receptors does not abrogate

DA-Hif-1α-upregulation of *tnfa:GFP*

(A) Fluorescent confocal micrographs of 1dpi caudal vein region of Mm and PVP infected

larvae. *tnfa* expression was detected by GFP levels, in green, using the

TgBAC(tnfa:GFP)pd1028 transgenic line. Phenol red (PR) and dominant active Hif-1α (DA1)

injected larvae were treated with DMSO or PD146176 (15-Lipoxygenase inhibitor). Non-

infected larvae are in the left panels (PVP) and Mm Crimson infected larvae are in the right

panels (Mm). Dotted lines indicate the yolk extension of the larvae where there is non-

specific fluorescence.

(B) Corrected fluorescence intensity levels of *tnfa:GFP* confocal z-stacks in uninfected

larvae (PVP, empty bars) and infected larvae (Mm, filled bars) at 1dpi of data shown in (A)

after treatment with DMSO or PD146176 (15-Lipoxygenase inhibitor). Data shown are mean

± SEM, n=42 cells from 7 embryos accumulated from 3 independent experiments.

(C) Bacterial burden at 4dpi after injection of DA Hif-1α (DA1) and 24 hours of the 15-

lipoxygenase inhibitor PD146176, using DMSO as a negative solvent control. Data shown

are mean ± SEM, n=48-50 as accumulated from 3 independent experiments.

(D) Fluorescent confocal micrographs of 1dpi caudal vein region of Mm and PVP infected

larvae shown in A. *tnfa* expression was detected by GFP levels, in green, using the

TgBAC(tnfa:GFP)pd1028 transgenic line. Phenol red (PR) and dominant active Hif-1α (DA1)

injected larvae were treated with DMSO or U75302/LY255283 (BLTR1/2 inhibitors). Non-

infected larvae are in the left panels (PVP) and Mm Crimson infected larvae are in the right

panels (Mm). Dotted lines indicate the yolk extension of the larvae where there is non-

specific fluorescence.

(E) Corrected fluorescence intensity levels of *tnfa:GFP* confocal z-stacks in uninfected

larvae (PVP, empty bars) and infected larvae (Mm, filled bars) at 1dpi of data shown in (D)

after treatment with DMSO or U75302/LY255283 (BLTR1/2 inhibitors). Data shown are

mean ± SEM, n=30 cells from 5 embryos accumulated from 2 independent experiments.

(F) Schematic of the arachidonic pathway indicating where stabilising Hif-1α upregulates

tnfa via cyclooxygenase, an effect that is not blocked using the 15-lipoxygenase inhibitor

PD146176 or BLTR1/2 inhibitors U75302/LY255283.

Figure 4. Hif-1α-induced *tnfa:GFP* requires active prostaglandin E2

(A) Fluorescent confocal micrographs of 1dpi caudal vein region of Mm. *tnfa* expression was

detected by GFP levels, in green, using the *TgBAC(tnfa:GFP)pd1028* transgenic line.

Phenol red (PR) and dominant active Hif-1α (DA1) injected larvae were treated with DMSO

or SC560 (Cox-1 inhibitor) in the presence or absence of endogenous prostaglandin E2

(PGE2). All larvae are PVP injected. Dotted lines indicate the yolk extension of the larvae

where there is non-specific fluorescence.

(B) Fluorescent confocal micrographs of 1dpi caudal vein region. *tnfa* expression was

detected by GFP levels, in green, using the *TgBAC(tnfa:GFP)pd1028* transgenic line.

Phenol red (PR) and dominant active Hif-1α (DA1) injected larvae were treated with DMSO

or NS398 (Cox-2 inhibitor) in the presence or absence of endogenous prostaglandin E2

(PGE2). All larvae are PVP injected. Dotted lines indicate the yolk extension of the larvae

where there is non-specific fluorescence.

(C) Corrected fluorescence intensity levels of *tnfa:GFP* confocal z-stacks in PVP injected

phenol red (PR, empty bars) and dominant active Hif-1α (DA1, filled bars) larvae at 1dpi of

data shown in (B) after treatment with DMSO/NS398/PGE2. Data shown are mean ± SEM,

n=54 cells from 9 embryos accumulated from 3 independent experiments.

(D) Corrected fluorescence intensity levels of *tnfa:GFP* confocal z-stacks in PVP injected

phenol red (PR, empty bars) and dominant active Hif-1α (DA1, filled bars) larvae at 1dpi of

data shown in (B) after treatment with DMSO/NS398/PGE2. Data shown are mean ± SEM,

n=54 cells from 9 embryos accumulated from 3 independent experiments.

(E) Fluorescent confocal micrographs of 1dpi caudal vein region. *tnf-a* expression was

detected by GFP levels, in green, using the *TgBAC(tnfa:GFP)pd1028* transgenic line.

Phenol red (PR) and dominant active Hif-1α (DA1) injected larvae were treated with DMSO

or NS398 (Cox-2 inhibitor) in the presence or absence of endogenous 15-keto prostaglandin

E2 (15K). All larvae are PVP injected. Dotted lines indicate the yolk extension of the larvae

where there is non-specific fluorescence.

(F) Corrected fluorescence intensity levels of *tnfa:GFP* confocal z-stacks in PVP injected

phenol red (PR, empty bars) and dominant active Hif-1α (DA1, filled bars) larvae at 1dpi of

data shown in (E) after treatment with DMSO/NS398/15-keto PGE2. Data shown are mean ±

SEM, n=24 cells from 4 embryos accumulated from 2 independent experiments.

(G) Schematic of the arachidonic pathway indicating where blocking cyclooxygenase

prevents Hif-1α upregulation of *tnfa*, an effect rescued by endogenous PGE2.

Figure 5. Hypoxia induces TNF in human MDMs in a COX-2 dependent manner

(A) TNF ELISA of human monocyte derived macrophages incubated in normoxia or 0.8%

hypoxia with or without treatment with NS398. Data shown are mean ± SEM, n=5-8

biological repeats from 3-4 donors.

(B) TNF ELISA of human monocyte derived macrophages incubated in normoxia treated

with FG4592 with or without treatment with NS398. Data shown are mean ± SEM, n=6

biological repeats from 3 donors.

Figure S1. Mm infection induced *tnfa:GFP* in macrophages in a second transgenic

line.

Fluorescent confocal micrographs of 1dpi caudal vein region of infection. *tnfa* expression

was detected by GFP levels, in green, using the *Tg(tnfa:eGFP-F)ump5Tg* transgenic line.

Macrophages are shown in red using a *Tg(mpeg1:mCherry-F)ump2Tg* line. Mm Crimson is

shown in the blue channel. Dotted lines indicate the yolk extension of the larvae where there

is non-specific fluorescence.

Figure S2. 5% oxygen induces expression of the *phd3:GFP* hypoxia transgene.

Fluorescent micrographs of 48hpf *Tg(phd3:eGFP)i144* hypoxia reporter zebrafish after

incubated in 6 hours of 5% oxygen from 32-38hpf, or normoxic controls.

Figure S3. DA-Hif-1α induced *il-1β:GFP* is not altered by Cox-1 inhibition and

decreased by Cox-2 inhibition.

(A) Fluorescent confocal micrographs of 2dpf caudal vein region. *il-1β* expression was

detected by GFP levels, in green, using the *TgBAC(il-1β:GFP)SH445* transgenic line.

Macrophages are shown in red using a *Tg(mpeg1:mCherryCAAX)sh378* line. Without

infection there is little detectable *il-1β:GFP* expression in phenol red (PR) controls, while DA-

Hif-1α macrophages have higher levels of *il-1β:GFP* in the Mm group.

(B) Fluorescent confocal micrographs of 1dpi caudal vein region of PVP injected larvae. *il-1β*

expression was detected by GFP levels. Phenol red (PR) and dominant active Hif-1α (DA1)

injected larvae were treated with DMSO, SC560 (COX-1 inhibitor) and NS398 (COX-2

inhibitor).

(C) Corrected fluorescence intensity levels of *il-1β:GFP* confocal z-stacks in PVP injected

larvae at 1dpi from (B). Data shown are mean ± SEM, n=36 cells from 6 embryos

representative of 3 independent experiments.

Figure S4. Mm infection induced *tnfa:GFP* is not altered by Hif-1α stabilisation

(A) Fluorescent confocal micrographs of Mm Crimson infected 1dpi zebrafish imaged around

the caudal vein region. *tnfa* expression was detected by GFP levels, in green, using the

TgBAC(tnfa:GFP)pd1028 transgenic line. Larvae were injected at the 1 cell stage with

dominant negative (DN) or dominant active (DA) Hif-1α or phenol red (PR) control. Dotted

lines indicate the yolk extension of the larvae where there is non-specific fluorescence.

(B) Graph shows corrected fluorescence intensity levels of *tnfa:GFP*. Dominant active Hif-1α

(DA1, filled bar) showed no change in *tnfa:GFP* levels in the presence of Mm bacterial

challenge compared to phenol red (PR) and dominant negative Hif-1α (DN1) injected

controls (white bars). NB, PR NoMm is a no Mm control taken from figure 1A for comparison

with Mm infected data, the data for which is from the same experiment as this one. Data

shown are mean ± SEM, n=66 cells from 12 embryos accumulated from 3 independent

experiments.

(C) Fluorescent confocal micrographs of 2dpf Mm infected *TgBAC(tnfa:GFP)pd1028*

transgenic larvae treated with DMOG or DMSO control. Dotted lines indicate the yolk

extension of the larvae where there is non-specific fluorescence.

(D) Graph shows corrected fluorescence intensity levels of *tnfa:GFP* confocal z-stacks in

larvae. DMOG treated larvae (filled bars) had significantly increased *tnfa:GFP* levels

compared to DMSO controls (white bars). Data shown are mean ± SEM, n=24 cells from 6

embryos representative of 2 independent experiments.

Figure S5. Cyclooxygenase inhibition does not block the protective effect of Hif-1α

stabilisation in Mm infection.

(A) Bacterial burden of SC560 treated larvae at 4dpi after injection with phenol red (PR) or

dominant active Hif-1α (DA1). Data shown are mean ± SEM, n=35 as accumulated from 3

independent experiments.

(B) Bacterial burden of NS398 treated larvae at 4dpi after injection with phenol red (PR) or

dominant active Hif-1α (DA1). Data shown are mean ± SEM, n=35 as accumulated from 3

independent experiments.

(C) Bacterial burden of DMSO treated larvae at 4dpi after co-treatment with DMSO or

DMOG. Data shown are mean ± SEM, n=33-36 as accumulated from 3 independent

experiments.

(D) Bacterial burden of SC560 treated larvae at 4dpi after co-treatment with DMSO or

DMOG. Data shown are mean ± SEM, n=33 as accumulated from 3 independent

experiments.

(E) Bacterial burden of NS398 treated larvae at 4dpi after co-treatment with DMSO or

DMOG. Data shown are mean ± SEM, n=33 as accumulated from 3 independent

experiments.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5