RESEARCH ARTICLE

Deep Brain Stimulation of the Ventromedial Prefrontal Cortex Disrupts Consolidation of Fear Memories

Shawn Zheng Kai **TAN**, Chi Him **POON**, Ying-Shing **CHAN**, Lee Wei **LIM**

School of Biomedical Sciences, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong SAR, China.

Corresponding author:

Lee Wei Lim MD, PhD, AM
School of Biomedical Sciences,
Li Ka Shing Faculty of Medicine,
The University of Hong Kong,
L4, 21 Sassoon Road,
Hong Kong SAR. China.
Tel: (852) 91572575
Email: limlw@hku.hk

Manuscript Information:
1. Number of words in the abstract: 150
2. Number of words in the manuscript (excluding references): 4258
3. Number of references:72
4. Number of figures: 5
5. Number of supplementary text: 1
6. Number of supplementary figures: 5
7. Number of supplementary tables: 1

Keywords: Neuromodulation, Deep Brain Stimulation; Anxiety; Fear; Memory; Dopamine

ABSTRACT

Anxiety disorders pose one of the biggest threats to mental health worldwide, yet

current therapeutics have been mostly ineffective due to issues with relapse, efficacy,

and toxicit. Deep brain stimulation (DBS) is a promising therapy for treatment-resistant

psychiatric disorders including anxiety, but very little is known about the effects of DBS

on fear memories. In this study, we used a modified plus-maze discriminative and

showed that DBS of the ventromedial prefrontal cortex (vmPFC) was able to disrupt

consolidation, but not acquisition or retrieval of fear memories. We validated these

results using a standard tone-footshock fear conditioning paradigm. We further

demonstrated short-term changes in dopaminergic receptor and c-Fos expression in the

ventral hippocampus (vHPC) and established a partial casual role of dopamine 2

receptors in this effect. Lastly, we showed changes in neurotransmitter levels in the

vHPC. This study highlights the potential therapeutic effect of vmPFC DBS to treat

anxiety disorders.

INTRODUCTION

Anxiety disorders are highly prevalent and pose one of the biggest threats to mental health worldwide (Ehlers, 1997; WHO, 2001). Anxiety disorders are commonly treated using exposure therapy, a form of cognitive-behavior therapy (CBT) that targets maladaptive associated fear memories. However, such exposure therapy involves new learning that attempts to inhibit or update the previous maladaptive learning rather than erase it (Khalaf et al., 2018), resulting in many patients unable to maintain the benefits of CBT and often relapsing (Barad, 2006; Baum, 1988; Bouton, 2002; Bouton and Bolles, 1979; McNally, 2007; Rescorla and Heth, 1975). Current attempts to enhance CBT using pharmacological or paradigm changes have met with complications such as low efficacy and drug toxicity (Farach et al., 2012; Klucken et al., 2016). Furthermore, improper administration of these techniques can lead to exacerbation of the condition (Eisenberg et al., 2003; Merlo et al., 2014; Pedreira and Maldonado, 2003).

Deep Brain Stimulation (DBS) is a surgically invasive technique that involves implanting electrodes in specific regions of the brain to modulate firing of neurons through electrical stimulation. It has been shown to be a promising non-pharmacological treatment for depression and anxiety disorders (Denys et al., 2010; Mayberg et al., 2005; Mian et al., 2010; Puigdemont et al., 2012). However, little work has been done to systematically investigate its effects on fear memory. In this study, we examined the hypothesis that DBS of the ventromedial prefrontal cortex (vmPFC), a structure we have previously argued to be an ideal target (Tan et al., 2019), is able to disrupt fear memories. Both innate and conditioned fear were investigated in rats using a modified avoidance plus maze in which bob cat urine (aversive odour) was placed in one closed arm of an elevated plus maze and rabbit urine (neutral odour) was placed in the opposite closed arm. The results were verified using a standard tone-footshock conditioning paradigm.

We then probed the role of the hippocampus in the effects seen, and showed that

ventral hippocampal dopaminergic transmission plays a central role in vmPFC DBS

ability to disrupt consolidation of fear memories.

METHODS AND MATERIALS

Subjects

The study was approved by the Committee on the Use of Live Animals in Teaching and

Research (CULATR) of The University of Hong Kong (Ref.: 4159-16). Male Sprague-

Dawley rats (n=173; 7-8 weeks old at the time of surgery) were individually housed in

standard cages with food and water available *ad libitum*. The environmental condition

was maintained at temperature (21±1 °C) and humidity (60-65%) under a reversed

12/12 h light/dark cycle.

Surgical and Deep Brain Stimulation Procedures

Surgery and DBS procedures were performed as previously described (Lim et al., 2015;

Liu et al., 2015). In brief, a construction of bilateral platinum-iridium electrodes (Synergy

Engineering Pte Ltd, Singapore) was implanted in the vmPFC (AP: +3.0 mm; ML: +/-0.6

mm; DV: -3.6 mm) based on the Paxinos & Watson Rat Brain Atlas (Paxinos and

Watson, 2006). Animals that received cannulation were also bilaterally implanted with

guide cannulas in the ventral hippocampus (AP: -5.3 mm; ML: +/-5.0 mm; DV: -5.6 mm).

Rats were connected to cables and stimulation was performed using a digital stimulator

(Model 3800 MultiStim: 8-Channel Stimulator; A-M Systems, Carlsborg, USA) and two

stimulus isolators (Model 3820; A-M Systems). Rats were stimulated according to the

experimental parameters (100 Hz, 200 µA and 100 µs pulse width) as previously

described (Lim et al., 2015; Liu et al., 2015). For verification of electrode/cannula

localization tip, hematoxylin-eosin (Merck, Darmstadt, Germany) was performed to

examine the implantation site. Detailed information about surgical and DBS procedures

are provided in the supplementary methods.

Administration of Drugs

Rats were infused with either Quinpirole–HCl (10 µg/side of salt, equivalent to 39.09

µmoles/side), Raclopride (1.67µg/side of salt, equivalent 3.36 µmoles/side) or artificial

cerebral spinal fluid (aCSF) at dosages previously shown to be effective in the vHPC

(Wilkerson and Levin, 1999). Drugs were infused into the vHPC through two Hamilton

syringes (10 µl) connected to an internal cannula via polyethylene tubing (Protech

International, Texas, USA). The infusion volume (2 µl) was delivered over approximately

3min and the internal cannula was left in for an additional 3min.

Modified Elevated Plus Maze

On day 1, a container with 5 ml of Bobcat urine (aversive odor; PredatorPee, Maine,

USA) was placed in one closed arm of the modified elevated plus maze (EPM) and a

container with 5 ml of rabbit urine (neutral odor) was placed in the opposite closed arm.

Urine was changed every 4-5 animals and testings were counter-balanced to ensure a

proper dispersion of odor across tests and subjects. On day 2, empty containers without

odor were placed in the two closed arms. Rats were stimulated (or sham stimulated)

accordingly (see protocol section) and placed in the central platform individually and

tested for 10min. More details can be found in the supplementary methods.

Real-time PCR

Immediately after experiments, animals were sacrificed, and brains were removed. The

dorsal hippocampus (dHPC), ventral hippocampus (vHPC), and medial prefrontal cortex

(mPFC) were micro-dissected (300 µm thickness) on a cryostat CM3050 (Leica

Microsystems, Wetzler, Germany). Total RNA was extracted, reverse transcribed, and

real-time PCR was performed. All primers used were previously published (Calabrese et

al., 2012; Covacu et al., 2009; Dick et al., 2015; Ermolinsky et al., 2008; Rogers et al.,

2004; Tan et al., 2015), and amplification efficiency was reassessed as previously

described (Livak and Schmittgen, 2001). Relative gene expression analysis was

performed using the $2^{-\Delta\Delta CT}$ method, and DBS animals were normalized to the sham

animals, as previously described (Livak and Schmittgen, 2001; Schmittgen and Livak,

2008). The list of primer sequences and the details of real-time PCR experiments can

be found in the Supplementary Table 1 and supplementary methods.

Mass Spectrometry

Metabolites were extracted from mPFC, dHPC, and vHPC slices, and derivatized using

a standard protocol. Gas chromatography–mass spectrometry (GC/MS) analysis was

performed on an Agilent 7890B GC - Agilent 7010 Triple Quadrapole Mass

Spectrometer system (Agilent, California, USA). Dopamine (DA), Serotonin (5-HT), γ-

Aminobutyric acid (GABA), Glutamic acid (Glu), 3,4-Dihydroxyphenylacetic acid

(DOPAC), Homovanillic acid (HVA), and 5-Hydroxyindole acetic acid (5-HIAA) were

measured with norvaline as the internal standard. More details can be found in the

supplementary methods.

Fear Conditioning

Fear conditioning was performed using a startle and fear conditioning system (Panlab

Harvard Apparatus, Massachusetts, USA). For acquisition, the conditioned stimulus

(CS) was a tone (Volume: 80 dB, Frequency: 5000 Hz) and the unconditioned stimulus

(US) was a 0.6 mA footshock. The protocol consisted of 2min of adaptation, followed by

three tone-footshock pairings, and then 2min of rest before being removed from the

chamber. For the context test, 24h after conditioning, rats were placed in the chamber

for 5min. Percentage freezing was reported based on the 5min test duration. For the

tone test, 24h after the context test, rats were placed in the chamber and tested in a

different context to the one received during conditioning. The tone tests consisted of

2min of adaptation, followed by five tone presentations (10s) with 10s inter-trial interval

(ITI) in the absence of footshock, and 2min of rest before being removed from the

chamber. Percentage freezing was reported as the average of the five CS

presentations.

To assess fear learning and memory, freezing was used as the dependent variable,

which is a species-specific defense response defined as the absence of all movement

except that required for respiration (Blanchard and Blanchard, 1969). Freezing values

were calculated using a high sensitivity Weight Transducer System (StartFear System,

Harvard Apparatus, Holliston, Massachusetts, USA). Open Field Test was conducted

24h after the tone test to control for locomotion differences. Animals were allowed to

explore the arena for 10min. The behavior of rats was recorded and analyzed using a

digital video camera and Any-maze video tracking system 5.0 (Stoelting Co).

181
Statistics

All statistical analyses were performed using GraphPad Prism 7.00. Statistical models

used are stated in the various results sections. Outliers were removed using the ROUT

method. Results were considered significant for $p<0.05$.

RESULTS

DBS of vmPFC disrupts consolidation, but not acquisition or retrieval of fear

memory

To investigate the effect of vmPFC DBS on fear memory, rats implanted with bilateral

electrodes in the vmPFC were tested in a modified EPM with aversive odor in one

closed arm, and then subjected to the retrieval task without odor 24h later. Rats were

either stimulated for 5min in the home cage before continual stimulation in the modified

EPM (Acquisition, Fig. 1A); stimulated for 5min in the home cage before continual

stimulation in the EPM during the retrieval task (Retrieval, Fig. 1F); or stimulated for a

further 15min at 15min after the acquisition task (Consolidation, Fig. 1K).

For the acquisition DBS group (n=20), unpaired t-test of the Discrimination Index (DI) on

day 1 showed no significant difference ($t_{(18)}$=0.24, p=0.81), indicating that vmPFC DBS

had no effect on acquisition of fear memory (Fig. 1B). Similarly, DI on day 2 showed no

significant difference ($t_{(18)}$=1.44, p=0.17), indicating that vmPFC DBS during acquisition

had no effect on retention of fear memory (Fig. 1C). There were no significant

differences in the time spent (t<0.82, all p>0.05) (Fig. 1D,E), or distance (t<0.62, all

p>0.05) (Supp. Fig. 1A,B), indicating that vmPFC DBS did not affect innate fear or

locomotion.

For the retrieval DBS group (n=27), unpaired t-tests showed no significant differences in

both acquisition ($t_{(23)}$=0.05, p=0.96) and retrieval ($t_{(24)}$=1.97, p=0.06) tasks (Fig. 1G,H),

indicating that vmPFC DBS had no effect on retrieval of fear memory. Similarly, no

significant differences were seen in the time spent (t<1.14, all p>0.05) (Fig. 1I,J) or

distance moved (t<0.95, all p>0.05) in the open arms on either day (Supp. Fig. 1D,E)

For the consolidation DBS group (n=18), unpaired t-test of the DI in the acquisition task

showed no significant difference ($t_{(14)}$=0.37, p=0.72), indicating the baseline fear

between the two groups was similar (Fig. 1L). Unpaired t-test of the DI on the day 2

retrieval task showed a significant difference ($t_{(15)}$=3.25, p=0.005), suggesting that

vmPFC DBS was able to disrupt consolidation of fear memory (Fig. 1M). There was no

significant difference in time spent in the open arms on either day (t<1.83, all p>0.05)

(Fig. 1N,O), suggesting the difference seen was not due to differences in innate fear.

Unpaired t-test of the distance moved on day 1 showed no significant difference

($t_{(14)}$=0.45, p=0.66) (Supp. Fig. 1H), whereas the distance moved on day 2 showed a

significant difference ($t_{(16)}$=2.17, p=0.045) (Supp. Fig. 1I). Taking into account all

previous experiments, these results seem to indicate that DBS did not cause any

locomotion differences, but rather vmPFC DBS animals had higher exploratory drive.

Single vmPFC stimulation during consolidation disrupts both tone and contextual

fear memory

To investigate if disruption of consolidation of memory affects both tone and contextual

fear and to validate the previous results, rats (n=19) implanted with bilateral electrodes

in the vmPFC were subject to the well-established tone-footshock fear conditioning

paradigm. 15min after the acquisition task, rats were stimulated (or sham stimulated) in

the home cage for 15min. 24h later, animals were placed back in the same context to

test contextual fear memory. 24h after the context test, animals were subjected to a new

context of five tones without footshock to test tone-footshock memory.

Two-way repeated measures ANOVA of the percentage of freezing in the conditioning

trial and the inter-trial intervals (ITIs) revealed an effect in the CS-US trial ($F_{(2,34)}$=80.9,

p<0.001; $F_{(3,51)}$=121.8, p<0.001), but not treatment or interaction (F<3.0, all p>0.05)

(Fig. 2A,B). This indicates proper acquisition of tone and context fear memory with no

difference between groups observed prior to DBS. Unpaired t-test of the context test

showed a significant difference between sham and DBS groups ($t_{(16)}$=2.38, p=0.03) (Fig.

2C). Unpaired t-test of the exploration period for tone test context showed no significant

differences between groups ($t_{(15)}$=0.32, p=0.75), indicating similar baseline levels for

new context. In the tone test, unpaired t-test of the average percentage of freezing for

all five tones showed a significant difference between sham and DBS groups ($t_{(13)}$=3.11,

p=0.008) (Fig. 2D). Animals were subsequently placed in an open field 24h after the

tone test to ensure the effects were not due to differences in locomotion. Unpaired t-test

of distance travelled showed no significant differences between sham and DBS groups

($t_{(17)}$=1.27, p=0.22), indicating no effects on locomotion (Fig. 2E).

Single stimulation during consolidation alters expressions of Drd2, Grm5,

GluN2A receptor, and c-Fos, expression in the vHPC

To understand the molecular mechanisms of vmPFC DBS on the hippocampus, rats

were sacrificed immediately after the trials on day 2. RT-qPCR was performed to detect

genes related to synaptic plasticity, learning and memory in dHPC and vHPC sections.

The t-tests showed no significant differences in any of the detected genes in the dHPC

and vHPC (t<1.74, all p>0.05) (Supp. Fig. 2A,B), indicating 15min of vmPFC DBS did

not cause long-term changes in gene expressions.

To examine the immediate changes in receptor expressions in the hippocampus upon

vmPFC DBS, rats (n=20) subjected to a similar trial as in the consolidation DBS group

were immediately sacrificed after DBS (Fig. 3A). RT-qPCR was performed to detect

genes in dHPC and vHPC sections (Fig. 3B). t-tests of the fold changes of genes in the

dHPC showed no significant differences (t<1.65, all p>0.05) (Fig. 3C,D). t-tests of fold

changes in the genes of the vHPC showed significant changes in the expressions of

Drd2 ($t_{(18)}$=2.37, p=0.029), Grm5 ($t_{(18)}$=2.12, p=0.048), and GluN2A ($t_{(18)}$=2.19, p=0.041),

but not Drd1, Grm2, Grm3, or GluN2B (t<0.17, all p>0.05) (Fig. 3D).

To detect the changes in neuronal activity of vmPFC stimulation in the hippocampus,

immediate early gene, c-Fos in mPFC, dHPC, and vHPC sections was examined by

RT-qPCR. The *t*-tests of the fold changes in c-Fos expression showed no significant

differences in the mPFC and dHPC ($t<1.12$, all $p>0.05$), but there was a significant

decrease in the vHPC ($t_{(17)}=2.16$, $p=0.045$) (Fig. 3E).

vHPC dopamine 2 receptors are involved in the effects of vmPFC DBS on

consolidation of memory

To establish the causal role of vHPC D2 receptor in the effects of vmPFC DBS on

consolidation, rats (n=69) were implanted with electrodes in the vmPFC and with guide

cannulas in the vHPC. Rats subjected to the modified EPM with an aversive odor in one

closed arm were then administered either aCSF, Quinpirole (a D2R agonist), or

Raclorpide (a D2R antagonist) via the guide cannula in the vHPC. At 15min after the

acquisition task, rats were stimulated (or sham stimulated) in the home cage for 15min.

Rats underwent the same EPM without odor 24h later to test retention of fear memory

(Fig. 4A).

Two-way ANOVA of the DI in the acquisition task showed no significant differences

($F<1.8$, all $p>0.05$) indicating the baseline fear between the two groups were similar

(Fig. 4B). Two-way ANOVA of the DI in the retrieval task showed an effect for

interaction ($F_{(2,58)}=3.95$, $p=0.025$), stimulation ($F_{(1,58)}=21.6$, $p<0.001$), and drugs

($F_{(2,58)}=7.84$, $p=0.001$). Tukey post-hoc test revealed significant differences in the aCSF

sham group compared to aCSF DBS ($p=0.002$) and Raclopride DBS ($p = 0.026$) groups,

but not Quinpirole DBS group. Within the DBS groups, aCSF and Raclopride groups

showed a significant difference compared with the Quinpirole group (aCSF: $p=0.001$;

Raclopride: $p=0.016$), but not with each other ($p=0.91$). All sham groups showed no

significant differences with each other (lowest p=0.93) (Fig. 4C). Two-way ANOVA of

the time spent in the open arms in the acquisition task showed an effect for drugs

($F_{(2,61)}$=5.07, p=0.01), but not interaction or stimulation (F<1.26, all p>0.05). Tukey post-

hoc test revealed a significant difference in only the aCSF sham group compared with

Raclopride sham group (p=0.01) (Fig. 4D). However, this effect disappeared in the

retrieval task, with the two-way ANOVA of the time spent in open arms showing no

effects (F<1.52, all p>0.05) (Fig. 4E). The differences seen in the acquisition task could

be attributed to either batch or random effects, although given the small differences in

the actual mean time (around 40s) and no differences in the retrieval task, we believe

the results are still valid. Lastly, no significant difference was seen in the distance

travelled in both the acquisition and retrieval tasks (F<2.50, all p>0.05) (Fig. 4F,G).

vmPFC DBS modulates neurotransmitters in the vHPC

To understand the effects of vmPFC DBS on neurotransmission, GC/MS was performed

for Glu, GABA, HVA, DOPAC, DA, 5-HIAA, and 5-HT in mPFC, dHPC, and vHPC

slices. All targets were within the linear ranges of the standard curves, except for DA,

which was excluded from the analysis as it was only detected in the vHPC of vmPFC

DBS group. t-test of the relative fold changes (target/average sham) in the vHPC

sections revealed significant differences in all targets (t>3.49, p<0.05) with decreases in

GABA, Glu, and 5-HIAA, and increases in HVA, DOPAC, and 5-HT (Fig. 5C). The t-test

of the relative fold changes in the dHPC sections revealed a significant decrease in only

5-HT with an increase in its metabolite 5-HIAA (t>4.22, all p<0.05) (Fig. 5B). The t-test

of the relative fold changes in mPFC sections revealed a significant decrease in only 5-

HIAA ($t_{(4)}$=3.68, p=0.021), although the difference in the mean of only 0.07 suggests no

clinically relevant significance (Fig. 5A). Chromatographs are shown in supplementary

materials (mPFC: Supp. Fig. 3; dHPC: Supp. Fig. 4; vHPC: Supp Fig. 5).

DISCUSSION

In this study, we systematically investigated the effects of acute vmPFC DBS on fear

memory. Using our experimental parameters, we showed vmPFC DBS was able to

disrupt consolidation, but not acquisition or retrieval of fear memory. The results from

the open arms and distance travelled further suggest that this is an effect of memory

rather than innate fear or locomotion. We found changes in the expression of

dopaminergic and glutamatergic receptors in the vHPC and established a partial causal

role of vHPC D2 receptors in this effect. Lastly, we found changes in neurotransmitters

in the vHPC.

Although using DBS to manipulate fear memory is not novel (Do-Monte et al., 2013;

Milad and Quirk, 2002; Reznikov et al., 2016; Rodriguez-Romaguera et al., 2012), few

studies have systematically studied its effects on the individual stages of the memory

process. Furthermore, most studies have focused on increasing extinction efficacy

rather than directly affecting the original memory (Do-Monte et al., 2013; Milad et al.,

2004; Milad and Quirk, 2002; Rodriguez-Romaguera et al., 2012). To the best of our

knowledge, this is the first study to isolate the effects on the consolidation of memory by

stimulating post acquisition. We also showed consistent results across several models

and repeat experiments. Our findings are of great importance as the effects of

neuromodulation on memories has been highly controversial, with some studies

showing improvements in memory (Fell et al., 2013; Miller et al., 2015; Suthana et al.,

2012; Titiz et al., 2017), and others showing disruption of memory (Coleshill, 2004;

Halgren et al., 1985; Halgren and Wilson, 1985; Jacobs et al., 2016; Lacruz et al., 2010;

Merkow et al., 2017).

We chose to target the vmPFC for its bidirectional connections in both the hippocampus

and the amygdala (Jin and Maren, 2015; Marek et al., 2013), structures which are

heavily implicated in fear memory (Kim et al., 1993; Ledoux, 1995; Maren, 2001).

Although microstimulation of the prelimbic cortex (PrL), and infralimbic cortex (IL) have

been shown to have opposing effects on the expression of conditioned fear (Vidal-

Gonzalez et al., 2006), we cannot comment on this specificity, as our results pertain to

the stimulation of the vmPFC as a whole. The proximity of PrL and IL means that

functional overlap of these structures cannot be ruled out (Giustino and Maren, 2015).

We investigated the downstream effects of vmPFC DBS on the hippocampus and

showed effects on the vHPC, but not the dHPC. This is perhaps not surprising due to

the direct monosynaptic projection from the vHPC to the vmPFC, which plays a crucial

role in anxiety (Padilla-Coreano et al., 2016). More surprising, our results suggest

effects on conditioned rather than innate fear - the functionality of the hippocampus is

thought to be split across the dorso-ventral axis, with the dHPC involved in spatial

encoding and the vHPC involved in innate fear (Bannerman et al., 1999; Fanselow and

Dong, 2010; Kheirbek et al., 2013; Kjelstrup et al., 2002; Moser et al., 1995). On the

other hand, the results could be related to how the dHPC and vHPC were

microdissected (Fig 3B), as vHPC slices were richer in CA3 compared to dHPC slices.

Obliteration of dopaminergic systems in CA3 has been shown to affect consolidation,

but not acquisition of memory (Wen et al., 2015). We cannot, however, entirely exclude

effects on innate fear, but rather suggest the effect is on memory. Furthermore, the

impairment of the "innate fear circuit" during consolidation might in turn lead to

disruption of the consolidation of the fear aspect of the memory, which could explain the

effects observed. Similarly, effects on the amygdala cannot be excluded, which

probably has a role in disrupting consolidation, as indicated by Klavir et al. (Klavir et al.,

2017), who found that high frequency optogenetic stimulation of amygdala inputs to the

PFC disrupted consolidation, but not acquisition of fear memory. Although optogenetic

stimulation is mechanistically different from DBS, it shares the same concept of axonal

activation (Abulseoud et al., 2012), that could suggest another mechanism by which

DBS can disrupt fear memories. Future work on the amygdala is needed to verify this.

Based on our gene expression results, we further studied the role of vHPC dopamine 2

receptor (D2R) on the effects of DBS. It has been shown vHPC D2R is involved in

spatial working memory, and there was a dose-related improvement with D2 agonist

Quinpirole and a dose-related deficit with D2 antagonist Raclopride (Wilkerson and

Levin, 1999). This is in line with our current results that showed a decrease in D2R gene

expression in the vHPC together with a deficit in memory. To establish a causal rather

than a correlative role of vHPC D2R, we infused Quinpirole and Raclopride separately

into the vHPC before DBS. Quinpirole was able to reverse the effects of DBS, whereas

Raclopride alone was not able to disrupt consolidation of memory. This suggests that

although vHPC D2Rs play a key role in the effects of vmPFC DBS on consolidation of

memory, it alone is not sufficient to disrupt consolidation, or fully explain the action of

vmPFC DBS.

One possible mechanism of how DBS exerts its action is through its modulatory effects

on neurotransmitters such as monoamines (Hamani et al., 2010; Van Dijk et al., 2012)

and glutamate (Agnesi et al., 2010; Jiménez-Sánchez et al., 2016; Tawfik et al., 2010).

We investigated how vmPFC stimulation during consolidation could be modulated by

various neurotransmitters. GC/MS results showed changes in all tested

neurotransmitters in vHPC, whereas dHPC showed only changes in the serotonin

system and mPFC showing no significant differences. As both glutamate and GABA

levels were lowered in vHPC, a "disruption hypothesis" might explain the observed

effects, in which information flow was disrupted (Chiken and Nambu, 2015). Lowered

glutamate level together with downregulated c-Fos might further suggest that vmPFC

DBS inhibits activity in the vHPC that in turn affects memory. However, the lower GABA

levels may complicate this theory and further work is needed to test this hypothesis.

Similar to the study by Volle et al. (Volle et al., 2018), we showed that vmPFC DBS

increased 5-HT levels in the vHPC. Contrary to another study (Hamani et al., 2010), we

showed 5-HT was lower in the dHPC, but this could be due to differences in length of

the stimulation (15min in our study compared to 4h in other study). The increase in 5-HT

in the vHPC was accompanied by lower 5-HIAA, which appears to be similar to the

actions of a MAO-inhibitor (Kaehler and Philippu, 1999) and hints at increased 5-HT

availability, although this requires further study. More pertinently and consistent with our

current study, we found significant changes in dopaminergic metabolites in the vHPC,

but not in dHPC or mPFC. Unfortunately, dopamine levels were too low to detect

reliably, making it difficult to accurately measure the dopamine turn-over. Interestingly,

we found increased dopamine metabolites in the vHPC of vmPFC DBS groups, and

dopamine was detectable only in the vmPFC DBS group, but not in the sham group.

This initially seems to contradict our results (in that dopamine agonist reversed the

effects of vmPFC DBS), however, it should be noted that Quinpirole has been shown to

lower dopamine and DOPAC (Santiago et al., 1993), which could explain the reversing

effects on vmPFC DBS. Further studies are needed to fully understand how vmPFC

DBS affects the complex play of dopamine in modulating memory processes.

To the best of our knowledge, this is the first study to show vmPFC DBS can disrupt

consolidation of fear memories. Our results were reproducible over multiple experiments

and paradigms, consistently showing the involvement of the vHPC and dopaminergic

system (particularly D2 receptors). However, the effects of DBS are complex and the

precise mechanisms, including the involvement of serotonergic and glutamatergic

systems, are still very much unknown and need to be further explored. Overall, we

provide strong evidence and a strong basis for further research into the use of

neuromodulation to disrupt fear memory processes as a possible treatment for anxiety

disorders.

Acknowledgements & Disclosures

The scientific work was funded by grants from the Hong Kong Research Grant Council

(RGC-ECS 27104616), and The University of Hong Kong URC Supplementary Funding

(102009728) that was awarded to LWL. We would like to thank the Proteomics and

Metabolomics Core Facility (HKU), the Laboratory Animal Unit (HKU), as well as Anna

Tse, Junhao Koh, and Smaranda Badea for their valuable input. All authors declared no

biomedical financial interests or potential conflicts of interests.

Figure Legends

Figure 1. vmPFC DBS affects consolidation but not acquisition or retrieval of fear

memory. Rats were trained in a modified elevated plus maze with aversive odor in one

closed arm, and neutral odor in the opposite closed arm. Rats were tested for arm

preference 24h after acquisition. Rats were stimulated either during acquisition (A),

retrieval (F), or consolidation (K). Both acquisition stimulation (A, B) and retrieval

stimulation (G, H) showed no significant differences in the discrimination index of the

closed arm with aversive odor vs neutral odor during the acquisition and retrieval tests,

indicating no effect on memory. For the consolidation stimulation, L shows there was no

significant difference in the discrimination index during acquisition, whereas B shows

significant differences in the retrieval test, indicating disruption of consolidation of

memory. No significant differences were seen in the time spent in the open arms in both

the acquisition and retrieval tasks for the acquisition DBS group (D, E), and retrieval

DBS group (I, J). For the consolidation stimulation, N shows there was no significant

difference in the time spent in the open arms, whereas O shows a significant difference

in the time spent in the open arms during the retrieval task. * $p<0.05$, ** $p<0.01$.

Figure 2. vmPFC DBS disrupts consolidation of both contextual and tone-

footshock memory. Rats were fear conditioned in a tone-footshock paradigm. vmPFC

DBS was carried out 15min after acquisition. Rats were then tested for contextual fear

memory and tone-footshock memory. A & B shows there was effective learning of both

CS-US and context-US fear memory, with no differences between groups at baseline.

Figure C shows there was significantly less freezing in the context test, whereas D

shows significantly less freezing in the tone test. E shows there was no significant

difference in the distance travelled in the open field test, indicating no difference in

locomotion. F shows electrode implantation sites. * $p<0.05$, ** $p<0.01$.

Figure 3. qPCR of various synaptic plasticity- and neuronal activity-related genes

immediately after stimulation. Rats were trained in a modified elevated plus maze

with aversive odor in one closed arm and neutral odor in the opposite closed arm.

vmPFC DBS was carried out 15 min after acquisition and rats were immediately

sacrificed (A). mPFC, dHPC, and vHPC sections were micro-dissected for qPCR (B). C

shows there were no significant differences in the dHPC. D shows downregulation of

Drd2, Grm5, and Grin2A in the vHPC. C-Fos gene expression was not significantly

changed in the mPFC or dHPC (E and F) but was significant downregulated in the

vHPC (G). * $p<0.05$.

Figure 4. vHPC Dopamine 2 receptor is involved in the effects of DBS on

consolidation of memory. Rats were trained in a modified elevated plus maze with

aversive odor in one closed arm and neutral odor in the opposite closes arm. aCSF,

Quinpirole, or Raclopride were immediately administered to the vHPC. vmPFC DBS

was carried out 15min after acquisition (A). B shows there was no significant difference

in the discrimination index during acquisition, whereas C shows significant difference

between DBS and sham groups for aCSF and Raclopride, but not Quinpirole group.

None of the sham groups showed significant differences with each other. Quinpirole

DBS group was significantly different from the aCSF and Raclorpide groups (C). There

was a significant difference between the aCSF and Raclopride sham pretreatment

groups (D), but no significant difference was observed in the retrieval task (E), which

suggests minor batch differences, but overall no effect on innate fear. No significant

difference was seen in the distance travelled indicating no effects on locomotion (F, G).

Figure 5. Mass spectrometry analysis of various neurotransmitters and

metabolites. mPFC, dHPC, and vHPC slices were analyzed by GC/MS for Glutamate,

GABA, HVA, DOPCA, 5-HIAA, and 5-HT. No Significant differences were observed in

the mPFC (A), while dHPC showed significant differences with an increase in 5-HIAA

and decrease in 5-HT. Significant differences were also observed in the vHPC with

decreases in Glutamate, GABA, and 5-HIAA, and increases in HVA, DOPAC, and 5-HT.

* $p<0.05$.

Supplementary Table 1. List of primers used for qPCR. Table of primer sequences

for qPCR with references. All primers were tested for efficiency before use.

Supplementary Figure 1. vmPFC DBS does not affect locomotion. No significant

differences were seen in the distance travelled for both the acquisition and retrieval

tasks in the acquisition DBS group (A, B) and retrieval DBS group (D, E). For

consolidation stimulation, H shows there was no significant difference in the distance

travelled, whereas I shows a significant difference in distance travelled, suggesting

higher exploratory drive rather than a difference in locomotion. C, F, & J show electrode

implantation sites. * $p<0.05$.

Supplementary Figure 2. qPCR of various synaptic plasticity- and neuronal

activity-related genes 24h after stimulation. Rats from the consolidation experiment

(Fig. 1K) were sacrificed, and dHPC and vHPC sections were micro-dissected for

qPCR. No significant differences were seen in the detected gene expressions in both

dHPC and vHPC (A, B).

Supplementary Figure 3. Mass spectrometry chromatographs of various

neurotransmitters and metabolites in the mPFC. mPFC slices were analyzed by

GC/MS for Glutamate, GABA, HVA, DOPCA, 5-HIAA, and 5-HT. A shows

chromatographs of Sham group, while B shows chromatographs of vmPFC DBS group.

Supplementary Figure 4. Mass spectrometry chromatographs of various

neurotransmitters and metabolites in the dHPC. dHPC slices were analyzed by

GC/MS for Glutamate, GABA, HVA, DOPCA, 5-HIAA, and 5-HT. A shows

chromatographs of Sham group, while B shows chromatographs of vmPFC DBS group.

Supplementary Figure 5. Mass spectrometry chromatographs of various

neurotransmitters and metabolites in the vHPC. vHPC slices were analyzed by

GC/MS for Glutamate, GABA, HVA, DOPCA, 5-HIAA, and 5-HT. A shows

chromatographs of Sham group, while B shows chromatographs of vmPFC DBS group.

References

Abulseoud OA, Knight EJ, Lee KH. 2012. Chapter 18: Neurotransmitter Release During Deep Brain Stimulation In: Denys D, Feenstra MGP, Schuurman R, editors. Deep Brain Stimulation: A New Frontier in Psychiatry. Springer. pp. 193–204.

Agnesi F, Blaha CD, Lin J, Lee KH. 2010. Local glutamate release in the rat ventral lateral thalamus evoked by high-frequency stimulation. *J Neural Eng* **7**:26009. doi:10.1088/1741-2560/7/2/026009

Bannerman DM, Yee BK, Good M a, Heupel MJ, Iversen SD, Rawlins JN. 1999. Double dissociation of function within the hippocampus: a comparison of dorsal, ventral, and complete hippocampal cytotoxic lesions. *Behav Neurosci* **113**:1170–1188. doi:10.1037/0735-7044.113.6.1170

Barad M. 2006. Is extinction of fear erasure or inhibition? Why both, of course. *Learn Mem* **13**:108–109. doi:10.1101/lm.211306

Baum M. 1988. Spontaneous recovery from the effects of flooding (exposure) in animals. *Behav Res Ther* **26**:185–186. doi:10.1016/0005-7967(88)90118-0

Blanchard RJ, Blanchard DC. 1969. Crouching as an index of fear. *J Comp Physiol Psychol* **67**:370. doi:10.1037/h0026779

Bouton ME. 2002. Context, ambiguity, and unlearning: Sources of relapse after behavioral extinction. *Biol Psychiatry* **52**:976–986. doi:10.1016/S0006-3223(02)01546-9

Bouton ME, Bolles RC. 1979. Contextual control of the extinction of conditioned fear. *Learn Motiv* **10**:445–466. doi:10.1016/0023-9690(79)90057-2

Calabrese F, Guidotti G, Molteni R, Racagni G, Mancini M, Riva MA. 2012. Stress-induced changes of hippocampal NMDA receptors: Modulation by duloxetine treatment. *PLoS One* **7**:3–10. doi:10.1371/journal.pone.0037916

Chiken S, Nambu A. 2015. Mechanism of Deep Brain Stimulation: Inhibition, Excitation, or Disruption? *Neuroscientist* **22**:313–322. doi:10.1177/1073858415581986

Coleshill SG. 2004. Material-Specific Recognition Memory Deficits Elicited by Unilateral Hippocampal Electrical Stimulation. *J Neurosci* **24**:1612–1616. doi:10.1523/JNEUROSCI.4352-03.2004

Covacu R, Arvidsson L, Andersson A, Khademi M, Erlandsson-Harris H, Harris RA, Svensson MA, Olsson T, Brundin L. 2009. TLR activation induces TNF-alpha production from adult neural stem/progenitor cells. *J Immunol* **182**:6889–6895. doi:10.4049/jimmunol.0802907

Denys D, Mantione M, Figee M, Munckhof P van den, Koerselman F, Westenberg H, Bosch A, Schuurman R. 2010. Deep Brain Stimulation of the Nucleus Accumbens for Treatment-Refractory Obsessive-Compulsive Disorder. *Arch Gen Psychiatry* **67**:1061–1068.

Dick ALW, Pooters T, Gibbs S, Giles E, Qama A, Lawrence AJ, Duncan JR. 2015.

NMDA receptor binding is reduced within mesocorticolimbic regions following chronic inhalation of toluene in adolescent rats. *Brain Res* **1624**:239–252. doi:10.1016/j.brainres.2015.07.037

Do-Monte FH, Rodriguez-Romaguera J, Rosas-Vidal LE, Quirk GJ. 2013. Deep brain stimulation of the ventral striatum increases BDNF in the fear extinction circuit. *Front Behav Neurosci* **7**:102. doi:10.3389/fnbeh.2013.00102

Ehlers A. 1997. 'Anxiety disorders: Challenging negative thinking.' Quoted in the Wellcome Trust Reveiws.

Eisenberg M, Kobilo T, Berman DE, Dudai Y. 2003. Stability of Retrieved Memory: Inverse Correlation with Trace Dominance. *Science (80-)* **301**:1102–1104. doi:10.1126/science.1086881

Ermolinsky B, Pacheco Otalora LF, Arshadmansab MF, Zarei MM, Garrido-Sanabria ER. 2008. Differential changes in mGlu2 and mGlu3 gene expression following pilocarpine-induced status epilepticus: A comparative real-time PCR analysis. *Brain Res* **1226**:173–180. doi:10.1016/j.brainres.2008.05.073

Fanselow M, Dong H-W. 2010. Are the Dorsal and Ventral Hippocampus functionally distinct structures. *Neuron* **65**:1–25. doi:10.1016/j.neuron.2009.11.031.Are

Farach FJ, Pruitt LD, Jun JJ, Jerud AB, Zoellner LA, Roy-Byrne PP. 2012. Pharmacological treatment of anxiety disorders: Current treatments and future directions. *J Anxiety Disord* **26**:833–843. doi:10.1016/j.janxdis.2012.07.009

Fell J, Staresina BP, Do Lam ATA, Widman G, Helmstaedter C, Elger CE, Axmacher N. 2013. Memory modulation by weak synchronous deep brain stimulation: A pilot study. *Brain Stimul* **6**:270–273. doi:10.1016/j.brs.2012.08.001

Giustino TF, Maren S. 2015. The Role of the Medial Prefrontal Cortex in the Conditioning and Extinction of Fear. *Front Behav Neurosci* **9**:298. doi:10.3389/fnbeh.2015.00298

Halgren E, Wilson CL. 1985. Recall deficits produced by afterdischarges in the human hippocampal formation and amygdala. *Electroencephalogr Clin Neurophysiol* **61**:375–380. doi:10.1016/0013-4694(85)91028-4

Halgren E, Wilson CL, Stapleton JM. 1985. Human medial temporal-lobe stimulation disrupts both formation and retrieval of recent memories. *Brain Cogn* **4**:287–295. doi:10.1016/0278-2626(85)90022-3

Hamani C, Diwan M, Macedo CE, Brandão ML, Shumake J, Gonzalez-Lima F, Raymond R, Lozano AM, Fletcher PJ, Nobrega JN. 2010. Antidepressant-Like Effects of Medial Prefrontal Cortex Deep Brain Stimulation in Rats. *Biol Psychiatry* **67**:117–124. doi:10.1016/j.biopsych.2009.08.025

Jacobs J, Miller J, Lee SA, Coffey T, Watrous AJ, Sperling MR, Sharan A, Worrell G, Berry B, Lega B, Jobst BC, Davis K, Gross RE, Sheth SA, Ezzyat Y, Das SR, Stein J, Gorniak R, Kahana MJ, Rizzuto DS. 2016. Direct Electrical Stimulation of the Human Entorhinal Region and Hippocampus Impairs Memory. *Neuron* **92**:983–990. doi:10.1016/j.neuron.2016.10.062

Jiménez-Sánchez L, Castañé A, Pérez-Caballero L, Grifoll-Escoda M, López-Gil X, Campa L, Galofré M, Berrocoso E, Adell A. 2016. Activation of AMPA Receptors Mediates the Antidepressant Action of Deep Brain Stimulation of the Infralimbic Prefrontal Cortex. *Cereb Cortex* **26**:2778–2789. doi:10.1093/cercor/bhv133

Jin J, Maren S. 2015. Prefrontal-Hippocampal Interactions in Memory and Emotion. *Front Syst Neurosci* **9**:170. doi:10.3389/fnsys.2015.00170

Kaehler ST, Philippu A. 1999. Effects of local MAO inhibition in the locus coeruleus on extracellular serotonin and 5-HIAA during exposure to sensory and cardiovascular stimuli. *Naunyn-Schmiedeberg's Arch Pharmacol* 187–193.

Khalaf O, Resch S, Dixsaut L, Gorden V, Glauser L, Gräff J. 2018. Fear Memory Attenuation. *Science (80-)* **1242**:1239–1242. doi:10.1126/science.aas9875

Downloaded

Kheirbek MA, Drew LJ, Burghardt NS, Costantini DO, Tannenholz L, Ahmari SE, Zeng
H, Fenton AA. 2013. Differential Control of Learning and Anxiety Along the
Dorsoventral Axis of the Dentate Gyrus. *Neuron* **77**:955–968.
doi:10.1016/j.neuron.2012.12.038

Kim JJ, Rison R a, Fanselow MS. 1993. Effects of amygdala, hippocampus, and
periaqueductal gray lesions on short- and long-term contextual fear. *Behav*
Neurosci **107**:1093–1098. doi:10.1037/0735-7044.107.6.1093

Kjelstrup KG, Tuvnes FA, Steffenach H-A, Murison R, Moser EI, Moser M-B. 2002.
Reduced fear expression after lesions of the ventral hippocampus. *Proc Natl Acad*
Sci U S A **99**:10825–30. doi:10.1073/pnas.152112399

Klavir O, Prigge M, Sarel A, Paz R, Yizhar O. 2017. Manipulating fear associations via
optogenetic modulation of amygdala inputs to prefrontal cortex. *Nat Neurosci* **20**.
doi:10.1038/nn.4523

Klucken T, Kruse O, Schweckendiek J, Kuepper Y, Mueller EM, Hennig J, Stark R.
2016. No evidence for blocking the return of fear by disrupting reconsolidation prior
to extinction learning. *Cortex* **79**:112–122. doi:10.1016/j.cortex.2016.03.015

Lacruz ME, Valentín A, Seoane JJG, Morris RG, Selway RP, Alarcón G. 2010. Single
pulse electrical stimulation of the hippocampus is sufficient to impair human
episodic memory. *Neuroscience* **170**:623–632.
doi:10.1016/j.neuroscience.2010.06.042

Ledoux JE. 1995. EMOTION□: Clues from the Brain. *Annu Rev Psychol* **46**:209–235.

Lim LW, Prickaerts J, Huguet G, Kadar E, Hartung H, Sharp T, Temel Y. 2015.
Electrical stimulation alleviates depressive-like behaviors of rats: investigation of
brain targets and potential mechanisms. *Transl Psychiatry* **5**:e535.
doi:10.1038/tp.2015.24

Liu A, Jain N, Vyas A, Lim LW. 2015. Ventromedial prefrontal cortex stimulation
enhances memory and hippocampal neurogenesis in the middle-aged rats. *Elife*
2015:1–21. doi:10.7554/eLife.04803

Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-
time quantitative PCR and. *Methods* **25**:402–408. doi:10.1006/meth.2001.1262

Marek R, Strobel C, Bredy TW, Sah P. 2013. The amygdala and medial prefrontal
cortex: partners in the fear circuit. *J Physiol* **591**:2381–91.
doi:10.1113/jphysiol.2012.248575

Maren S. 2001. Neurobiology of Pavlovian Fear Conditioning. *Annu Rev Nuerosci*
24:897–931.

Mayberg HS, Lozano AM, Voon V, McNeely HE, Seminowicz D, Hamani C, Schwalb
JM, Kennedy SH. 2005. Deep brain stimulation for treatment-resistant depression.
Neuron **45**:651–660. doi:10.1016/j.neuron.2005.02.014

McNally RJ. 2007. Mechanisms of exposure therapy: How neuroscience can improve
psychological treatments for anxiety disorders. *Clin Psychol Rev* **27**:750–759.
doi:10.1016/j.cpr.2007.01.003

Merkow MB, Burke JF, Ramayya AG, Sharan AD, Sperling MR, Kahana MJ. 2017.
Stimulation of the human medial temporal lobe between learning and recall
selectively enhances forgetting. *Brain Stimul* **10**:645–650.
doi:10.1016/j.brs.2016.12.011

Merlo E, Milton AL, Goozee ZY, Theobald DE, Everitt BJ. 2014. Reconsolidation and
extinction are dissociable and mutually exclusive processes: behavioral and
molecular evidence. *J Neurosci* **34**:2422–2431. doi:10.1523/JNEUROSCI.4001-
13.2014

Mian MK, Campos M, Sheth SA, Eskandar EN. 2010. Deep brain stimulation for
obsessive-compulsive disorder: past, present, and future. *Neurosurg Focus*

676   **29**:E10. doi:10.3171/2010.4.FOCUS10107

677 Milad MR, Vidal-Gonzalez I, Quirk GJ. 2004. Electrical stimulation of medial prefrontal
678   cortex reduces conditioned fear in a temporally specific manner. *Behav Neurosci*
679   **118**:389–394. doi:10.1037/0735-7044.118.2.389

680 Milad MRR, Quirk GJJ. 2002. Neurons in medial prefrontal cortex signal memory for
681   fear extinction. *Nature* **420**:70–74. doi:10.1038/nature01144.1.

682 Miller JP, Sweet JA, Bailey CM, Munyon CN, Luders HO, Fastenau PS. 2015. Visual-
683   spatial memory may be enhanced with theta burst deep brain stimulation of the
684   fornix: A preliminary investigation with four cases. *Brain* **138**:1833–1842.
685   doi:10.1093/brain/awv095

686 Moser MB, Moser EI, Forrest E, Andersen P, Morris RG. 1995. Spatial learning with a
687   minislab in the dorsal hippocampus. *Proc Natl Acad Sci U S A* **92**:9697–701.
688   doi:10.1073/pnas.92.21.9697

689 Padilla-Coreano N, Bolkan SS, Pierce GM, Blackman DR, Hardin WD, Garcia-Garcia
690   AL, Spellman TJ, Gordon JA. 2016. Direct ventral hippocampal-prefrontal input is
691   required for anxiety-related neural activity and behavior. *Neuron* **17**:857–866.
692   doi:10.1016/j.neuron.2016.01.011.Direct

693 Paxinos G, Watson C. 2006. The Rat Brain in Stereotaxic Coordinates Sixth Edition by.
694   *Acad Press* **170**:547612. doi:10.1016/0143-4179(83)90049-5

695 Pedreira ME, Maldonado H. 2003. Protein synthesis subserves reconsolidation or
696   extinction depending on reminder duration. *Neuron* **38**:863–869.
697   doi:10.1016/S0896-6273(03)00352-0

698 Puigdemont D, Pérez-Egea R, Portella MJ, Molet J, de Diego-Adeliño J, Gironell A,
699   Radua J, Gómez-Anson B, Rodríguez R, Serra M, de Quintana C, Artigas F,
700   Álvarez E, Pérez V. 2012. Deep brain stimulation of the subcallosal cingulate gyrus:
701   further evidence in treatment-resistant major depression. *Int J*
702   *Neuropsychopharmacol* **15**:121–133. doi:10.1017/S1461145711001088

703 Rescorla RA, Heth CD. 1975. Reinstatement of Fear to an Extinguished Conditioned
704   Stimulus **104**:88–96.

705 Reznikov R, Binko M, Nobrega JN, Hamani C. 2016. Deep Brain Stimulation in Animal
706   Models of Fear, Anxiety and Post-Traumatic Stress Disorder.
707   *Neuropsychopharmacology* 1–8. doi:10.1038/npp.2016.34

708 Rodriguez-Romaguera J, Monte FHM Do, Quirk GJ. 2012. Deep brain stimulation of the
709   ventral striatum enhances extinction of conditioned fear. *Proc Natl Acad Sci*
710   **109**:8764–8769. doi:10.1073/pnas.1200782109/-
711   /DCSupplemental.www.pnas.org/cgi/doi/10.1073/pnas.1200782109

712 Rogers A, Schmuck G, Scholz G, Williams DC. 2004. c-fos mRNA Expression in Rat
713   Cortical Neurons During Glutamate-Mediated Excitotoxicity. *Toxicol Sci* **82**:562–
714   569. doi:10.1093/toxsci/kfh279

715 Santiago M, Machado A, Cano J. 1993. Effects of Age and Dopamine Agonists and
716   Antagonists on Striatal Dopamine Release in the Rat: An In Vivo Microdialysis
717   Study. *Mech Ageing Dev* **67**:261–267.

718 Schmittgen TD, Livak KJ. 2008. Analyzing real-time PCR data by the comparative CT
719   method. *Nat Protoc* **3**:1101–1108. doi:10.1038/nprot.2008.73

720 Suthana N, Haneef Z, Stern J, Mukamel R, Behnke E, Knowlton B, Fried I. 2012.
721   Memory Enhancement and Deep-Brain Stimulation of the Entorhinal Area. *N Engl J*
722   *Med* **366**:502–10. doi:10.1056/NEJMoa1107212

723 Tan SZK, Ganella DE, Dick ALW, Duncan JR, Ong-Palsson E, Bathgate RAD, Kim JH,
724   Lawrence AJ. 2015. Spatial Learning Requires mGlu5 Signalling in the Dorsal
725   Hippocampus. *Neurochem Res* **40**:1303–1310. doi:10.1007/s11064-015-1595-0

726 Tan SZK, Sheng V, Chan Y-S, Lim LW. 2019. Eternal Sunshine of the Neuromodulated
727   Mind: Altering Fear Memories Through Neuromodulation. *Exp Neurol* **314**:9–19.

Tawfik VL, Chang SY, Hitti FL, Roberts DW, Leiter JC, Jovanovic S, Lee KH. 2010.
Deep brain stimulation results in local glutamate and adenosine release:
Investigation into the role of astrocytes. *Neurosurgery* **67**:367–375.
doi:10.1227/01.NEU.0000371988.73620.4C
Titiz AS, Hill MRH, Mankin EA, Aghajan ZM, Eliashiv D, Tchemodanov N, Maoz U,
Stern J, Tran ME, Schuette P, Behnke E, Suthana NA, Fried I. 2017. Theta-burst
microstimulation in the human entorhinal area improves memory specificity. *Elife*
6:1–18. doi:10.7554/eLife.29515
Van Dijk A, Klompmakers AA, Feenstra MGP, Denys D. 2012. Deep brain stimulation of
the accumbens increases dopamine, serotonin, and noradrenaline in the prefrontal
cortex. *J Neurochem* **123**:897–903. doi:10.1111/jnc.12054
Vidal-Gonzalez I, Vidal-Gonzalez B, Rauch SL, Quirk GJ. 2006. Microstimulation
reveals opposing influences of prelimbic and infralimbic cortex on the expression of
conditioned fear. *Learn Mem* **13**:728–733. doi:10.1101/lm.306106
Volle J, Bregman T, Scott B, Diwan M, Raymond R, Fletcher PJ, Nobrega JN, Hamani
C. 2018. Deep brain stimulation and fluoxetine exert different long-term changes in
the serotonergic system. *Neuropharmacology* **135**:63–72.
doi:10.1016/j.neuropharm.2018.03.005
Wen J, Xue L, Wang R, Chen Z, Shi Y, Zhao H. 2015. Involvement of the dopaminergic
system in the consolidation of fear conditioning in hippocampal CA3 subregion.
Behav Brain Res **278**:527–534. doi:10.1016/j.bbr.2014.10.049
WHO. 2001. World health report 2001: mental health: new understand, new hope.
World Heal Organ Rep no 9240681701.
Wilkerson A, Levin ED. 1999. Ventral hippocampal dopamine D1 and D2 systems and
spatial working memory in rats. *Neuroscience* **89**:743–749. doi:10.1016/S0306-
4522(98)00346-7