Time series experimental design under one-shot sampling: The importance of condition diversity

Xiaohan Kang[1], Bruce Hajek[1], Faqiang Wu[2] Yoshie Hanzawa[2],

1 Coordinated Science Laboratory and Department of Electrical and Computer Engineering, University of Illinois at Urbana–Champaign, Urbana, Illinois, United States of America
2 Department of Biology, California State University, Northridge, Northridge, California, United States of America

1   Abstract

Many biological data sets are prepared using one-shot sampling, in which each individual organism provides only one sample. Time series therefore do not follow trajectories of individuals over time. However, samples collected at different times from individuals grown under the same conditions share the same perturbations of the biological processes, and hence behave as surrogates for multiple samples from a single individual at different times. This implies the importance of growing individuals under multiple conditions if one-shot sampling is used. This paper models the condition effect explicitly by correlated perturbations in the variations driving the expression level dynamics, quantifies the performance of network structure estimators both analytically and numerically, and illustrates the difficulty in network reconstruction under one-shot sampling when the condition effect is absent.

2   Introduction

Time series data is important for studying biological systems because of their dynamic nature. Ideally it is desirable to use time series with *multi-shot sampling*, where each individual[1] is sampled multiple times to produce the trajectory of the biological process, as in Fig 1. Then the natural biological variations in different individuals can be leveraged for statistical inference, and thus inference can be made even if the samples are prepared under the same experimental condition.

Fig 1. Multi-shot sampling. Each plant is observed four times.

However, in many experiments multi-shot sampling is not possible. Due to stress response of the organisms and/or the large amount of cell tissue required for accurate measurements, the dynamics of the relevant biological process in an individual of the organism cannot be observed at multiple times without interference. For example, in an

---

[1]We sometimes call the individual organism a "plant" in this paper, while the model and analysis apply to animals and microorganisms as well.

RNA-seq experiment an individual plant is often only sampled once in its entire life, 12
leaving the dynamics unobserved at other times. See Appendix A.1 for a review of 13
literature on this subject. We call the resulting time series data, as illustrated in Fig 2, 14
a time series with *one-shot sampling*. Because the time series with one-shot sampling do

Fig 2. One-shot sampling. Each plant is observed once.

not follow the trajectories of the same individuals, they do not capture all the 16
correlations in the biological processes. For example, the trajectory of observations on 17
plants 1–2–3–4 and that on 1–6–7–4 in Fig. 2 are statistically identical. The resulting 18
partial observation renders some common models for the biological system dynamics 19
inaccurate or even irrelevant. 20

To address this problem, instead of getting multi-shot time series of single 21
individuals, one can grow multiple individuals under each condition with a variety of 22
conditions, and get one-shot time series of the single conditions. The one-shot samples 23
from the same condition then become a surrogate for multi-shot samples for a single 24
individual, as illustrated in Fig 3. In essence, if we view the *condition effect* on the

Fig 3. One-shot sampling with two different conditions.

samples as being random, then there should be a positive correlation among samples 26
grown under the same condition. It is similar to the effect of biological variation of a 27
single individual sampled at different times (if such sampling were possible). 28

For each condition, the one-shot samples at different times are also complemented by 29
biological replicates, which are samples from independent individuals taken at the *same* 30
time used to reduce measurement errors and/or biological variations. See Appendix A.2 31
for a review on how replicates are used for biological inference. With a budget over the 32
number of samples, a balance must be kept between the number of conditions, the 33
number of sampling times and the number of replicates. 34

To illustrate and quantify the effect of one-shot sampling in biological inference, we 35
introduce a simple dynamic gene expression model with a condition effect. We consider 36
a hypothesis testing setting and model the dynamics of the gene expression levels at 37
different sampling times by a dynamic Bayesian network (DBN), where the randomness 38

comes from biological and environmental variations. The condition-dependent variation of gene j is the same for all plants in that condition and makes up a fraction γ_j of all the variation, the remaining of which is biological variation and is independent for the individuals. The fraction γ_j is called the *condition correlation coefficient* of gene j for the varying conditions and it indicates the strength of the condition effect on the expression of gene j. For example, for varying temperature conditions a differentially expressed gene may have a γ of 0.4 while a non-differentially expressed gene may have a γ of 0.1. In contrast to GeneNetWeaver [1], where the condition effect is modeled by the same random perturbation to the network coefficients, in our model the condition effect is characterized by correlation in the variation terms of the dynamics. Note in both models samples from different individuals under the same condition are statistically independent given the randomness associated with the condition.

The contributions of this paper are threefold.

1. A composite hypothesis testing problem on the gene regulatory network is formulated and a gene expression dynamic model that explicitly capture the per-gene condition effect and the gene regulatory interactions is proposed.

2. The performance of gene regulatory network structure estimators is analyzed for both multi-shot and one-shot sampling, with focus on two algorithms. Furthermore, single-gene and multi-gene simulation results indicate that multiple-condition experiments can somewhat mitigate the shortcomings of one-shot sampling.

3. The difficulty of network reconstruction under one-shot sampling with no condition effect is illustrated. This difficulty connects network analysis and differential expression analysis, two common tasks in large-scale genomics studies, in the sense that the part of network involving non-differentially expressed genes may be harder to reconstruct.

3 Materials and Methods

3.1 Stochastic model of time-series samples

This section formulates the hypothesis testing problem of learning the structure of the gene regulatory network (GRN) from gene expression data with one-shot or multi-shot sampling. The GRN is characterized by an unknown adjacency matrix. Given the GRN, a dynamic Bayesian network model is used for the gene expression evolution with time. A parameter γ_j is used for each gene j to explicitly capture the correlation between the variations in the gene expression levels of the individuals under the same experimental condition.

3.1.1 Notation

For any positive integer n, let $[n] = \{1, 2, \ldots, n\}$. We use $(f(x))_{x \in \mathcal{I}}$ to denote the family of elements in the set $\{f(x) \colon x \in \mathcal{I}\}$ indexed by the index set $\mathcal{I}$. The indicator function on a statement or a set P is denoted by $\mathbb{1}_P$. The n-by-n identity matrix is denoted by I_n. The transpose of matrix A is denoted by A^*.

3.1.2 Model for gene regulatory network topology

Let n be the number of genes and let $A \in \mathcal{A} \subseteq \mathbb{R}^{n \times n}$ be the unknown adjacency matrix of the GRN. The sign of the entry a_{ij} of A for $i \neq j$ indicates the type of regulation of j by i, and the absolute value the strength of the regulation. A zero entry $a_{ij} = 0$ with

$i \neq j$ indicates no regulation of j by i. The diagonal of A characterizes gene self-regulation. Let $\mathcal{G} = \{G_1, G_2, \ldots, G_{|\mathcal{G}|}\}$ be a finite set of network structures and let D be a mapping from $\mathcal{A}$ to $\mathcal{G}$. Then $\mathcal{A}$ is partitioned by the associated network structures. Fix a loss function $l \colon \mathcal{G}^2 \to \mathbb{R}$. Let $Y \in \mathcal{Y}$ be the random observation and let $\delta \colon \mathcal{Y} \to \mathcal{G}$ be an estimator for the structure. The performance of an estimator is evaluated by the expected loss $\mathbb{E}\, l(D(A), \delta(Y))$. This is a hypothesis testing problem with composite hypotheses $\{D^{-1}(G) \colon G \in \mathcal{G}\}$. This paper considers network reconstruction up to regulation type with $D(A) = (\mathrm{sgn}(A_{ij}))_{(i,j)\in([n]\times[n])}$, where $\mathrm{sgn}(s) = \mathbb{1}_{\{s>0\}} - \mathbb{1}_{\{s<0\}}$. In other words, the ternary value of the edge signs (positive, negative, or no edge) are to be recovered. Some examples of loss functions are as follows.

- Ternary false discovery rate (FDR)

$$l_{\mathrm{FDR}}(S, S') = 1 - \frac{\sum_{i=1}^{n}\sum_{j=1}^{n} \mathbb{1}_{\{S_{ij}=S'_{ij}\neq 0\}}}{\sum_{i=1}^{n}\sum_{j=1}^{n} \mathbb{1}_{\{S'_{ij}\neq 0\}}}.$$

- Ternary false negative rate (FNR)

$$l_{\mathrm{FNR}}(S, S') = 1 - \frac{\sum_{i=1}^{n}\sum_{j=1}^{n} \mathbb{1}_{\{S_{ij}=S'_{ij}\neq 0\}}}{\sum_{i=1}^{n}\sum_{j=1}^{n} \mathbb{1}_{\{S_{ij}\neq 0\}}}.$$

- Ternary false positive rate (FPR)

$$l_{\mathrm{FPR}}(S, S') = 1 - \frac{\sum_{i=1}^{n}\sum_{j=1}^{n} \mathbb{1}_{\{S_{ij}=S'_{ij}=0\}}}{\sum_{i=1}^{n}\sum_{j=1}^{n} \mathbb{1}_{\{S_{ij}=0\}}}.$$

- Ternary error rate

$$l_{\mathrm{E}}(S, S') = \frac{1}{n^2} \sum_{i=1}^{n}\sum_{j=1}^{n} \mathbb{1}_{\{S_{ij}\neq S'_{ij}\}}.$$

Note $l_{\mathrm{FDR}}(S, S') = l_{\mathrm{FNR}}(S', S)$. Also if S does not contain zeros then $l_{\mathrm{FNR}}(S, S') = l_{\mathrm{E}}(S, S')$. Similarly if S' does not contain zeros then $l_{\mathrm{FDR}}(S, S') = l_{\mathrm{E}}(S, S')$. As an example, for a random guessing algorithm with probabilities of $S'_{ij} = 0, 1, -1$ being $1-q, q/2, q/2$ and a network prior with probabilities of $S_{ij} = 0, 1, -1$ being $1-p, p/2, p/2$, $l_{\mathrm{FDR}} = 1 - p/2$, $l_{\mathrm{FNR}} = 1 - q/2$, and $l_{\mathrm{FPR}} = q$.

3.1.3 Model for gene expression dynamics

This section models the gene expression dynamics of individuals by dynamic Bayesian networks with a parameter γ_j for gene j, which is the condition correlation coefficient of variation in the expression level dynamics of gene j.

Let K, T and C be the number of individuals, sampling times, and conditions, respectively. Let $X_j^k(t) \in \mathbb{R}$ be the expression level of gene $j \in [n]$ in individual $k \in [K]$ at time $t \in [T]$, and let $c_k \in [C]$ be the label that indicates the condition for individual k. The gene expression levels evolve according to the *Gaussian linear model*[2] (GLM) with initial condition $X_j^k(0) = 0$ for any $j \in [n]$, $k \in [K]$ and recursion

$$
\begin{aligned}
X_j^k&(t+1) \\
&= \sum_{i=1}^{n} X_i^k(t) A_{ij} \\
&\quad + \sigma_j \sqrt{\gamma_j} W_{\mathrm{cond},j}^{c_k}(t+1) + \sigma_j \sqrt{1-\gamma_j} W_{\mathrm{biol},j}^{k}(t+1),
\end{aligned}
\tag{1}
$$

---

[2]A logarithm transform is often applied to the estimated concentrations of proteins such that lowly expressed genes have negative expression levels.

for $j \in [n]$, $k \in [K]$, and $t \in \{0, 1, \dots, T-1\}$, where $\hspace{1cm}$
$\left(W^c_{\text{cond},j}(t)\right)_{(c,j,t) \in [C] \times [n] \times [T]}$, $\left(W^j_{\text{biol},k}(t)\right)_{(j,k,t) \in [n] \times [K] \times [T]}$ are collections of independent standard Gaussian random variables that are used to drive the dynamics, $\sigma_j \geq 0$ the driving noise level of gene j, and $\gamma_j \in [0, 1]$ the fraction of condition-dependent variation for gene j. Here the last two terms in (1) denotes the condition-dependent variation and biological variation respectively. To write (1) in matrix form, we let $X(t) = \left(X^k_j(t)\right)_{(k,j) \in [K] \times [n]}$ and $W(t) = \left(W^k_j(t)\right)_{(k,j) \in [K] \times [n]}$ be K-by-n matrices, where $W^k_j(t) = \sigma_j \sqrt{\gamma_j} W^{c_k}_{\text{cond},j}(t) + \sigma_j \sqrt{1 - \gamma_j} W^k_{\text{biol},j}(t)$. Then

$$X(t+1) = X(t)A + W(t+1) \tag{2}$$

and hence

$$X(t) = \sum_{\tau=1}^{t} W(\tau) A^{t-\tau}. \tag{3}$$

3.1.4 Model for sampling method

This paper considers a *full factorial design* with CRT samples obtained under C
conditions, R replicates and T sampling times, denoted by $Y = (Y^{c,r,t})_{(c,r,t) \in [C] \times [R] \times [T]}$.
Here the triple index for each sample indicates the condition, replicate and time.
Consider two sampling methods that generate Y: multi-shot sampling and one-shot
sampling.

Multi-shot sampling Assume an individual can be sampled multiple times. This sampling model requires $K = CR$ and $c_k = \lceil \frac{k}{R} \rceil \in [C]$ for all $k \in [K]$. Equivalently, multi-index (c, r) can be used to determine the individual instead of k for X and W with c denoting the condition and r the replicate. Then (1) for multi-shot sampling can be rewritten as

$$X^{c,r}_j(t+1)$$
$$= \sum_{i=1}^{n} X^{c,r}_i(t) A_{ij}$$
$$+ \sigma_j \sqrt{\gamma_j} W^c_{\text{cond},j}(t+1) + \sigma_j \sqrt{1 - \gamma_j} W^{c,r}_{\text{biol},j}(t+1),$$

and the observation for condition c, replicate r and time t is

$$Y^{c,r,t}_j = X^{c,r}_j(t) + \sigma_{Z,j} Z^{c,r,t}, \tag{4}$$

with $\left(Z^{c,r,t}_j\right)_{(j,c,r,t) \in [n] \times [C] \times [R] \times [T]}$ being a collection of independent standard Gaussian
random variables modeling the observation noise, and $\sigma_{Z,j}$ is the observation noise level
of gene j. We see that for fixed c and r the observations of different times are from the
same individual with the multi-index (c, r).

One-shot sampling Assume an individual can be sampled only once. This model requires $K = CRT$ and $c_k = \lceil \frac{k}{RT} \rceil \in [C]$ for all $k \in [K]$. Equivalently, with multi-index (c, r, s) denoting the condition, the replicate, and the target sampling time, the evolution (1) for one-shot sampling can be rewritten as

$$X^{c,r,s}_j(t+1)$$
$$= \sum_{i=1}^{n} X^{c,r,s}_i(t) A_{ij}$$
$$+ \sigma_j \sqrt{\gamma_j} W^c_{\text{cond},j}(t+1) + \sigma_j \sqrt{1 - \gamma_j} W^{c,r,s}_{\text{biol},j}(t+1),$$

and the observation is
$$Y^{c,r,t} = X^{c,r,t}(t) + \sigma_{Z,j} Z^{c,r,t}. \tag{5}$$

Again $\sigma_{Z,j}$ is the observation noise level of gene j and the Z's are independent standard Gaussian random variables. Note that for fixed c and r the observations of different times are from *different* individuals because the triple indices are different.

4 Results

4.1 Performance evaluation of network structure estimators

This section studies the performance of network structure estimators with multi-shot and one-shot sampling data. First, general properties of the two sampling methods are obtained. Then two algorithms, the generalized likelihood-ratio test (GLRT) and the basic sparse linear regression (BSLR) are studied. The former is a standard decision rule for composite hypothesis testing problems, and is shown to have some properties in Section 4.1.2 but is computationally infeasible for even a small number of genes. The latter, described in Section 4.1.2, is an algorithm based on linear regression, and is feasible for a moderate number of genes. Finally simulation results for a single-gene network with GLRT and for a multi-gene network with BSLR are shown in Sections 4.1.2 and 4.1.4.

4.1.1 General properties

By (3), (4) and (5), the samples Y are jointly Gaussian with zero mean. The covariance of the random tensor Y is derived under the two sampling methods in the following.

Under multi-shot sampling, the samples under different conditions are independent and hence uncorrelated. Consider $Y^{c,r,t}$ and $Y^{c,r',t'}$, which are two samples under the same condition and collected at times t and t'. The covariance matrix between $Y^{c,r,t}$ and $Y^{c,r',t'}$ is the sum of the covariance matrices of their common variations at times τ for $1 \le \tau \le t \wedge t'$ multiplied by $(A^*)^{t-\tau}$ on the left and $A^{t'-\tau}$ on the right, plus covariance for the observation noise. The covariance matrices of the variations are $\Sigma_W = \mathrm{diag}(\sigma_1^2, \sigma_2^2, \ldots, \sigma_n^2)$ if the two samples are from the same individual (i.e., $r = r'$), and $\Sigma_W \Gamma$ otherwise, where $\Gamma = \mathrm{diag}(\gamma_1, \gamma_2, \ldots, \gamma_n)$. Let $\Sigma_Z = \mathrm{diag}(\sigma_{Z,1}^2, \sigma_{Z,2}^2, \ldots, \sigma_{Z,n}^2)$. This yields:

$$\mathbb{E}((Y^{c,r,t})^* Y^{c',r',t'})$$
$$= \begin{cases} \sum_{\tau=1}^{t} (A^*)^{t-\tau} \Sigma_W A^{t-\tau} + \Sigma_Z & \text{if } (c,r,t) = (c',r',t'), \\ \sum_{\tau=1}^{t \wedge t'} (A^*)^{t-\tau} \Sigma_W A^{t'-\tau} & \text{if } (c,r) = (c',r') \text{ and } t \ne t', \\ \sum_{\tau=1}^{t \wedge t'} (A^*)^{t-\tau} \Sigma_W \Gamma A^{t'-\tau} & \text{if } c = c' \text{ and } r \ne r', \\ 0 & \text{if } c \ne c'. \end{cases}$$

Under one-shot sampling the only difference compared with multi-shot sampling is that two samples indexed by (c,r,t) and (c,r,t') are from different individuals if $t \ne t'$. So

$$\mathbb{E}((Y^{c,r,t})^* Y^{c',r',t'}) \tag{6}$$
$$= \begin{cases} \sum_{\tau=1}^{t} (A^*)^{t-\tau} \Sigma_W A^{t-\tau} + \Sigma_Z & \text{if } (c,r,t) = (c',r',t'), \\ \sum_{\tau=1}^{t \wedge t'} (A^*)^{t-\tau} \Sigma_W \Gamma A^{t'-\tau} & \text{if } c = c' \text{ and } (r,t) \ne (r',t'), \\ 0 & \text{if } c \ne c'. \end{cases}$$

For any fixed network structure estimator:

1. If $\gamma_j = 1$ for all j and C, R and T are fixed, the joint distribution of the data is the same for both types of sampling. (So the performance of the estimator would be the same for multi-shot and one-shot sampling.)

2. If $\gamma_j = 1$ for all gene j, $\sigma_Z^2 = 0$ (no observation noise) and C, T are fixed, the joint distribution of the data is the same for both types of sampling (as noted in item 1) and any replicates beyond the first are identical to the first. (So the performance of the estimator can be obtained even if all replicates beyond the first are discarded.)

3. Under multi-shot sampling, when C, R, T are fixed with $R = 1$, the joint distribution of the data does not depend on γ_j for all $\gamma_j \in [0, 1]$. (So the performance of the estimator would be the same for all $\gamma_j \in [0, 1]$, $j \in [n]$.)

4. In the homogeneous gene case with $\sigma_j = \sigma$, $\gamma_j = \gamma$, $\sigma_{Z,j} = \sigma_Z$ for all j, if the estimator δ is based on replicate averages $y = (y^{c,t})_{(c,t) \in [C] \times [T]}$ with $y^{c,t} = \frac{1}{R} \sum_{r=1}^{R} Y^{c,r,t}$, and if δ is scale-invariant (i.e., $\delta(Y) = \delta(cY)$ for any $c \neq 0$ and Y), then its performance under multi-shot sampling depends on σ^2, σ_Z^2, γ and R only through

$$\widetilde{\sigma}^2 = \frac{\sigma^2}{\sigma_Z^2}(\gamma R + 1 - \gamma).$$

Under one-shot sampling, the estimator's performance depends on σ^2, σ_Z^2, γ and R only through $\widetilde{\sigma}^2$ and $\widetilde{\gamma}$, where $\widetilde{\sigma}^2$ is as defined above and

$$\widetilde{\gamma} = \frac{\gamma}{\gamma + (1 - \gamma)/R}.$$

To see 4), note that averaging reduces the variance of the biological variation and that of the observation noise by a factor of R due to independence, but preserves the condition-dependent variation because it is identical across replicates. Hence the variance of the variation in the dynamics of the averages is $\sigma^2(\gamma + \frac{1-\gamma}{R})$ and the variance of the observation noise of the averages is σ_Z^2/R. Then the averages are essentially single-replicate data, and the performance under multi-shot sampling depends only on the ratio of the new total variance to the new observation noise variance. For one-shot sampling the new condition correlation coefficient also matters for the single-replicate data, so the performance depends both on $\widetilde{\sigma}^2$ and $\widetilde{\gamma}$.

4.1.2 Network reconstruction algorithms

In this section we introduce GLRT and BSLR. GLRT is a standard choice in composite hypothesis testing setting. We observe some properties for GLRT under one-shot and multi-shot sampling. However, GLRT involves optimizing the likelihood over the entire parameter space, which grows exponentially with the square of the number of genes. Hence GLRT is hard to compute for multiple-gene network reconstruction. In contrast, BSLR is an intuitive algorithm based on linear regression, and will be shown in simulations to perform reasonably well for multi-gene scenarios.

GLRT The GLRT (see, e.g., page 38, Chapter II.E in [15]) is given by $\delta(y) = D(\hat{A}_{\mathrm{ML}}(y))$, where $\hat{A}_{\mathrm{ML}}(y)$ is the maximum-likelihood estimate for A based on the covariance of Y in Section 4.1.1 given the observation $Y = y$.

Proposition 1. *Suppose all genes have the same condition effect; i.e., $\gamma_j = \gamma$ for all $j \in [n]$. Then GLRT (with the knowledge of γ) has the following properties.*

1. *Under multi-shot sampling with $\sigma_{Z,j}^2 = 0$ (no observation noise) and $\sigma_j^2 = \sigma^2$, the performance of GLRT (with the knowledge of σ) for sign estimation is the same for all (R, γ) excluding $(R \geq 2, \gamma = 1)$.* **165 166 167**

2. *Under one-shot sampling and $\gamma = 0$, the log likelihood of the data as a function of A (i.e. the log likelihood function) is invariant with respect to replacing A by $-A$. (So, for the single-gene $n = 1$ case, the log likelihood function is an even function of A, and thus the GLRT will do no better than random guessing.)* **168 169 170 171**

For 2 it suffices to notice in (6) the covariance is invariant with respect to changing A to $-A$. A proof of 1 is given below. **172 173**

Proof of 1). We first prove it for the case of a single gene with constant T and a constant number of individuals, CR. To do that we need to look at the likelihood function closely. **174 175 176**

Because the trajectories for different conditions are independent (for given parameters (A, γ)), we shall first consider the case with a single condition; i.e., $C = 1$. There are hence R trajectories of length T. We may assume $\sigma^2 = 1$. Then the covariance matrix of the length-R driving vector used at time t for the trajectories is

$$\mathrm{Cov}(W(t)) = (1 - \gamma)I_R + \gamma J_R =: \Sigma.$$

When $\gamma > 0$, Σ is not the identity matrix multiplied by some constant; i.e., the noise vector $W(t)$ is colored across replicates. It can be checked when $\gamma < 1$ the matrix Σ is positive definite. Then there exists an orthogonal matrix U and a diagonal matrix Λ with positive diagonal elements such that $\Sigma = U\Lambda U^*$. Let $\Sigma^{-1/2} = U\Lambda^{-1/2}U^*$ and let

$$\widetilde{X}(t) = \Sigma^{-1/2}X(t),$$

$$\widetilde{W}(t) = \Sigma^{-1/2}W(t)$$

for all $t \in [T]$. Then the trajectories for the R replicates in a single condition become:

$$\widetilde{X}(t+1) = \widetilde{X}(t)A + \widetilde{W}(t+1).$$

It can be checked that after the linear transformation by $\Sigma^{-1/2}$, which does not depend on A, the new driving vectors are white (i.e., $\mathrm{Cov}(\widetilde{W}(t)) = I_R$). It follows that the distribution of $\widetilde{X}|(A, \gamma)$ is the same as the distribution of $X|(A, 0)$ (i.e. $\gamma = 0$). Therefore, for $x = (x^r(t))_{(r,t)\in[R]\times[T]}$, if we let $L_X(x|A, \gamma)$ denote the likelihood of $X = x$ for parameters A, γ, then

$$L_X(x|A, \gamma) = L_{\widetilde{X}}(\widetilde{x}|A, \gamma)d(R, T, \gamma) = L_X(\widetilde{x}|A, 0)d(R, T, \gamma),$$

where $d(R, T, \gamma) = (\det \Sigma)^{-T/2}$ is a function of R, T and γ, and $\widetilde{x}(t) = \Sigma^{-1/2}x(t)$. **177**

Now consider the likelihood function for all CRT samples with general C. It is the product of C likelihood functions for the samples prepared under the C different conditions. It is thus equal to $d(R, T, \gamma)^C$ times the likelihood of the transformed expression levels $\widetilde{x}$, which is the likelihood function for $\gamma = 0$ and a total of CRT samples. The form of the product depends on C and R only through CR, because under the transformation, all CR trajectories are independent. Hence, for fixed A, γ, C, R, T the distribution of the maximum likelihood estimate of A, when the samples are generated using a given $\gamma > 0$ (so the R individuals under each condition are correlated) and the likelihood function also uses γ, is the same as the distribution of

the maximum likelihood estimate of A when $\gamma = 0$ (in which case the CR individual trajectories are independent and identically distributed). Formally,

$$\mathbb{E}_\gamma \, l(D(A), \delta(Y)) = \mathbb{E}_\gamma \, l(D(A), D(\arg\max_A L_X(X|A, \gamma)))$$

$$= \mathbb{E}_\gamma \, l(D(A), D(\arg\max_A L_X(\widetilde{X}|A, 0)))$$

$$= \mathbb{E}_0 \, l(D(A), D(\arg\max_A L_X(X|A, 0)))$$

$$= \mathbb{E}_0 \, l(D(A), \delta(Y)),$$

where $\mathbb{E}_\gamma$ denotes that the condition effect of the random elements X and Y is γ. The above fails if $\gamma = 1$ and $R \geq 2$ because then Σ is singular. It also fails if γ is unknown to the GLRT.

For the general model with multiple genes, if γ is the same for each gene, 1) holds as before – for the proof, apply left multiplication by $\Sigma^{-\frac{1}{2}}$ for each gene, time, and condition to all R samples in the condition. View (2) as an update equation for an $R \times n$ matrix for each group of R samples in one condition. One column of length R per gene, and one row per sample. $\qquad\square$

BSLR In BSLR, replicates are averaged and the average gene expression levels at different times under different conditions are fitted in a linear regression model with best-subset sparse model selection, followed by a Granger causality test to eliminate the false discoveries. BSLR is similar to other two-stage linear regression–based network reconstruction algorithms, notably oCSE [16] and CaSPIAN [17]. Both oCSE and CaSPIAN use greedy algorithms in the first build-up stage, making them more suitable for large-scale problems. In contrast, BSLR uses best subset selection, which is conceptually simpler but computationally expensive for large n. For the tear-down stage both BSLR and CaSPIAN use the Granger causality test, while oCSE uses a permutation test.

Build-up stage In the first stage BSLR finds potential regulatory interactions using a linear regression model. Let $Y_j^c(t) = \frac{1}{R} \sum_{r=1}^R Y^{c,r,t}$ and let $Y(t) = \left(Y_j^c(t)\right)_{(c,j) \in [C] \times [n]}$ denote the C-by-n matrix. Let

$$\Psi(1) = \begin{pmatrix} Y(2) \\ Y(3) \\ \vdots \\ Y(T) \end{pmatrix}$$

and

$$\Psi(0) = \begin{pmatrix} Y(1) \\ Y(2) \\ \vdots \\ Y(T-1) \end{pmatrix}.$$

For each target gene $j \in [n]$, BSLR solves the following best subset selection problem with a subset size $k < n$:

$$\underset{A_{\cdot j}, b_j, d_j}{\text{minimize}} \quad \|\Psi_j(1) - \Psi(0)A_{\cdot j} - d_j \Psi_j(0) - b_j \mathbf{1}\|_2^2$$

$$\text{subject to} \quad \|A_{\cdot j}\|_0 \leq k \text{ and } A_{jj} = 0.$$

Denote the solution by (A^*, b^*, d^*). The output of the first stage is then A^*.

A naive algorithm to solve the above optimization has a computational complexity of $O(n^{k+1})$ for fixed k as $n \to \infty$. Faster near-optimal alternatives exist [18].

Tear-down stage   The second stage is the same as that of CaSPIAN. For each $j \in [n]$ and each $i \in \text{supp}(A_{\cdot j})$, let the unrestricted residual sum of squares be

$$\text{RSS}_{\text{u}} = \|\Psi_j(1) - \Psi(0)A_{\cdot j}^* - d_j^*\Psi_j(0) - b_j^*\mathbf{1}\|_2^2$$

and the restricted residual sum of squares

$$\text{RSS}_{\text{r}} = \inf\{\|\Psi_j(1) - \Psi(0)A_{\cdot j} - d_j\Psi_j(0) - b_j\mathbf{1}\|_2^2:$$
$$\text{supp}(A_{\cdot j}) = \text{supp}(A_{\cdot j}^*)\backslash\{i\}\}.$$

The F-statistic is given by

$$F = \frac{\text{RSS}_r - \text{RSS}_u}{\text{RSS}_u/(C(T-1) - k - 2)}.$$

The potential parent i of j is removed in the tear-down stage if the p-value of the F-statistic with degrees of freedom $(1, C(T-1) - k - 2)$ is above the preset significance level (e.g., 0.05). Note the tests are done for all parents in $A_{\cdot j}$ simultaneously; both the restricted and the unrestricted models contain the other potential parents regardless of the results of the tests on them. secnum

4.1.3   Simulations on single-gene network reconstruction

Simulations on single-gene network reconstruction

The Gaussian linear model is used to simulate one-shot sampling data with a single gene. In order to compare one-shot and multi-shot sampling, we view the main expense to be proportional to the number of samples to prepare as opposed to the number of individuals to grow. We thus fix a total budget of $CRT = 180$ samples and consider full factorial design with C and R varying with $CR = 30$, and $T = 6$ with 10 000 simulations. Results are plotted in Fig. 4. The four plots on the left are for one-shot sampling and the four on the right are for multi-shot sampling. For each plot the observed probability of sign (of A) error is shown for $\gamma \in \{0, 0.2, 0.4, 0.6, 0.8, 1.0\}$ and for R ranging over the divisors of 30 from smallest to largest. Plots 4a–4d show the performance for the GLRT algorithm assuming no observation noise ($\sigma_Z^2 = 0$) and known γ. Plots 4e–4h show the performance for the GLRT algorithm assuming observation noise with known variance $\sigma_Z^2 = 1$, and with both γ and A unknown to the algorithm.
  The numerical simulations reflect the following observations implied by the analytical model.

1. Under one-shot sampling, when $\gamma = 0$, the GLRT is equivalent to random guessing.

2. The GLRT performs the same under one-shot and multi-shot sampling when $\gamma = 1$.

3. Under no observation noise, the performance for multi-shot sampling is the same for all $\gamma < 1$.

Some empirical observations are in order.

1. Multi-shot sampling outperforms one-shot sampling (unless $\gamma = 1$, where they have the same error probability).

2. For one-shot sampling, the performance improves as γ increases. Regarding the number of replicates R per condition, if $\gamma = 0.2$ (small condition effect), a medium number of replicates (2 to 5) outperforms the single replicate strategy. For larger γ, one replicate per condition is the best.

3. For multi-shot sampling, performance worsens as γ increases. One replicate per condition ($R = 1$) is best.

4. Comparing 4a–4d vs. 4e–4h, we observe that the performance degrades with the addition of observation noise, though for moderate noise ($\sigma_Z = 1.0$) the effect of observation noise on the sign error is not large. Also, the effect of the algorithm not knowing γ is not large.

4.1.4 Simulations on multi-gene network reconstruction

This subsection studies the case when multiple genes interact through the GRN. The goal is to compare one-shot vs. multi-shot sampling for BSLR under a variety of scenarios, including different homogeneous γ values, varying number of replicates, varying observation noise level, and heterogeneous γ values.

The performance evaluation for multi-gene network reconstruction is trickier than the single-gene case because of the many degrees of freedom introduced by the number of genes. First, the network adjacency matrix A is now an n-by-n matrix. While some notion of "size" of A (like the spectral radius or the matrix norm) may be important, potentially every entry of A may affect the reconstruction result. So instead of fixing a ground truth A as in Fig. 4, we fix a prior distribution of A with split Gaussian prior described in Appendix C. Second, because the prior of A can be subject to sparsity constraints and thus far from a uniform distribution, multiple loss functions that are more meaningful than the ternary error rate can be considered for performance. So we consider ternary FDR, ternary FNR and ternary FPR for the multi-gene case. In the simulations we have 20 genes and $d_{\max} = 3$ with in-degree uniformly distributed over $\{0, 1, \ldots, d_{\max}\}$, so the average in-degree is 1.5. The number of sampling times is $T = 6$ and $CR = 30$.

Varying γ, R and σ_Z   In this set of simulations we fix the observation noise level and vary the number of replicates R and the condition correlation coefficient γ. The performance of BSLR under one-shot and multi-shot sampling is shown in Fig 5 ($\sigma_Z = 0$) and Fig 6 ($\sigma_Z = 1$). Note BSLR does not apply to a single condition with 30 replicates due to the constraint that the degrees of freedom $C(T - 1) - k - 2$ in the second stage must be at least 1.

For one-shot sampling, when $\gamma = 0$, we see in both Fig. 5 and Fig. 6 that BSLR is no different from random guessing, with an FDR close to $1 - \frac{1}{2}\frac{1.5}{19} \approx 0.96$ and an FNR and an FPR such that $l_{\mathrm{FNR}} + \frac{1}{2}l_{\mathrm{FPR}} \approx 1$ (recall the example of random guessing at the end of Section 3.1.2). When $\gamma = 1$, BSLR performs similarly with one-shot or multi-shot sampling, which is consistent with the property 1 in Section 4.1.1. As γ increases from 0 to 1, under one-shot sampling for a fixed number of replicates, the FDR and FNR reduce greatly. For example, as γ increases from 0.2 to 1, the FDR for single replicate under one-shot sampling decreases from 0.74 to 0.31 with noiseless data (Fig. 5), and from 0.79 to 0.36 with noisy data (Fig. 6), while the FNR decreases from 0.70 to 0.00 with noiseless data, and from 0.78 to 0.04 with noisy data. This decrease is more pronounced for smaller number of replicates. Note the trend of the performance of BSLR under one-shot sampling as a function of R and γ is very similar to that of GLRT in Figs. 4e and 4g.

For multi-shot sampling, in the noiseless case, we see all three losses are invariant with respect to different γ for fixed R, which is consistent with Property 4 in Section 4.1.1 because BSLR is an average-based scale-invariant algorithm. In the noisy case, the FDR and FNR slightly decrease as γ increases, which is an opposite trend compared with Figs. 4f and 4h.

In summary, the main conclusions from Figs. 5 and 6 are the following.

(a) One-shot sampling, known γ, $A = 0.1$, $\sigma_Z = 0$.

(b) Multi-shot sampling, known γ, $A = 0.1$, $\sigma_Z = 0$.

(c) One-shot sampling, known γ, $A = 0.5$, $\sigma_Z = 0$.

(d) Multi-shot sampling, known γ, $A = 0.5$, $\sigma_Z = 0$.

(e) One-shot sampling, unknown γ, $A = 0.1$, $\sigma_Z = 1$.

(f) Multi-shot sampling, unknown γ, $A = 0.1$, $\sigma_Z = 1$.

(g) One-shot sampling, unknown γ, $A = 0.5$, $\sigma_Z = 1$.

(h) Multi-shot sampling, unknown γ, $A = 0.5$, $\sigma_Z = 1$.

Fig 4. Performance of the GLRT in single-gene sign recovery with different numbers of replicates.

(a) FDR of one-shot sampling.

(b) FDR of multi-shot sampling.

(c) FNR of one-shot sampling.

(d) FNR of multi-shot sampling.

(e) FPR of one-shot sampling.

(f) FPR of multi-shot sampling.

Fig 5. Performance of the BSLR in multi-gene network reconstruction with different numbers of replicates, $\sigma_Z = 0$.

(a) FDR of one-shot sampling.

(b) FDR of Multi-shot sampling.

(c) FNR of one-shot sampling.

(d) FNR of multi-shot sampling.

(e) FPR of one-shot sampling.

(f) FPR of multi-shot sampling.

Fig 6. Performance of the BSLR in multi-gene network reconstruction with different numbers of replicates, $\sigma_Z = 1$.

- The performance of BSLR under multi-shot sampling is consistently better than that under one-shot sampling.

- The performance of BSLR under one-shot sampling varies with γ, from random guessing performance at $\gamma = 0$ to the same performance as multi-shot sampling at $\gamma = 1$.

- By comparing Fig. 5 with Fig. 6, we see the observation noise of $\sigma_Z = 1$ has only a small effect on the performance with the two sampling methods.

Reduced number of directly differentially expressed genes In the above simulations we have assumed all genes are equally directly differentially expressed. In other words, we took γ_j to be the same for all j. To test what happens more generally, we conducted simulations such that only half of the genes are directly differentially expressed genes (DEGs), while the other half are non-DEGs. To do so, we assign $\gamma_j = 1$ for $1 \leq j \leq 10$, and $\gamma_j = 0$ for $11 \leq j \leq 20$. The results for $R = 3$ are pictured in Fig. 7. We see that with one-shot sampling the edges coming out of the DEGs are reconstructed with lower FDR and FNR compared to those coming out of non-DEGs. However, under one-shot sampling, even the edges from the non-DEGs in Fig. 7 are recovered with much lower FDR and FNR, as compared to one-shot sampling in Fig. 6 with $\gamma = 0$ and $R = 3$. The results indicate that the performance of BSLR under one-shot sampling benefits from diversity in conditions even when not all genes are directly differentially expressed.

5 Discussion

5.1 Information limitation for reconstruction under one shot sampling without condition effect

In the previous section it is shown that both GLRT and BSLR are close to random guessing under one-shot sampling when $\gamma_j = 0$ for all j. This leads us to the following question: is the network reconstruction with no condition effect ($\gamma_j = 0$ for all j) information theoretically possible? In this section we examine this question under general estimator-independent settings. Note in this case the trajectories of all individuals are independent given A regardless of $(c_k)_{k \in [K]}$.

As we have seen in Proposition 1 part 2, when $\gamma_j = 0$ for all j, the distribution of the observed data Y is invariant under adjacency matrix A or $-A$, implying any estimator will have a sign error probability no better than random guessing for the average or worst case over A and $-A$. Here, instead of sign error probability, we consider the estimation for A itself.

The extreme case with infinite number of samples available for network reconstruction is considered to give a lower bound on the accuracy for the finite data case. Note that with infinite number of samples a sufficient statistic for the estimation of the parameter A is the marginal distributions of $X^1(t)$; no information on the correlation of $\left(X^1(t)\right)_{t \in [T]}$ across time t can be obtained. A similar observation is made in [19] for sampling stochastic differential equations.

We first consider the transient case with $X(0) = 0$ as stated in Section 3.1.3. With infinite data the covariance matrix $\Sigma_t \triangleq \mathrm{Cov}(X(t)) = \sum_{\tau=1}^{t} (A^*)^{t-\tau} A^{t-\tau}$ can be recovered for $t \in [T]$. Now we want to solve A from $(\Sigma_t)_{t \in [T]}$. As a special case, if $A^*A = \rho I_n$ (i.e., $\rho^{-1/2}A$ is orthogonal) then we will have $\Sigma_t = \sum_{\tau=0}^{t-1} \rho^\tau I_n$. As a result, given $(\Sigma_t)_{t \in [T]}$ in the above form, no more information of A can be obtained other than $\rho^{-1/2}A$ being orthogonal, with $\frac{n(n-1)}{2}$ degrees of freedom remaining. In general case it is not clear if A can be recovered from $(\Sigma_t)_{t \in [T]}$.

(a) FDR.

(b) FNR.

(c) FPR.

Fig 7. Performance of BSLR for heterogeneous γ_i with $\sigma_Z = 1$.

Now consider the case where X^k is in steady state; i.e., $X(0)$ is random such that $\mathrm{Cov}(X(t))$ is invariant with t. With infinite amount of data we can get the covariance matrix Σ, which satisfies $\Sigma = A^*\Sigma A + I$. Since covariance matrices are symmetric, we have $\frac{n(n+1)}{2}$ equations for n^2 variables in A. Thus A is in general not determined by the equation uniquely. In fact, note that Σ is positive definite. Then by eigendecomposition $\Sigma = Q\Lambda Q^*$, where Q is an orthogonal matrix and Λ the diagonal matrix of the eigenvalues of Σ. Then $\Lambda = (Q^*AQ)^*\Lambda(Q^*AQ) + I$. Let $B = QAQ^*$. Then $\Lambda = B^*\Lambda B$. By the Gram–Schmidt process, B can be determined with $\frac{n(n-1)}{2}$ degrees of freedom. So the network cannot be recovered from the stationary covariance matrix.

In summary, the recovery of the matrix A is generally not possible in the stationary case, and also not possible in the transient case at least when A is orthogonal. To reconstruct A, further constraints (like sparsity) may be required.

6 Conclusion

One-shot sampling can miss a lot of potentially useful correlation information. Often gene expression data collected from plants is prepared under one-shot sampling. One factor that can partially mitigate the shortcomings of one-shot sampling is to prepare samples under a variety of conditions or perturbations. Samples grown under the same condition can then be thought of as a surrogate for an individual plant.

To clarify issues and take a step towards quantifying them, we proposed a model with a parameter $\gamma_j \in [0, 1]$ which is the correlation coefficient for the expression level of gene j induced by preparation of samples under the same condition. The model includes the joint expression level dynamics of multiple genes.

There is little agreement across the biology literature about how to model the impact of condition on the gene regulatory network. In some cases, it is not even clear that we are observing the same network structure as conditions vary. Nevertheless, our results suggest that the preparation of samples under different conditions can partially compensate for the shortcomings of one-shot sampling.

Appendices

A   Related work

A.1   One-shot sampling in the literature

This section reviews the sampling procedures reported in several papers measuring gene expression levels in biological organisms with samples collected at different times to form time series data. In all cases, the sampling is one-shot, in the sense that a single plant or cell is only sampled at one time.

A.1.1   Microorganisms

In the transcriptional network inference challenge from DREAM5 [2], three compendia of biological data sets were provided based on microorganisms (*E. coli*, *S. aureus*, and *S. cerevisiae*), and some of the data corresponded to different sampling times in a time series. Being based on microorganisms, the expression level measurements involved multiple individuals per sample, a form of one-shot sampling.

A.1.2 Plants

In [3], the plants are exposed to nitrate, which serves as a synchronizing event, and samples are taken from three to twenty minutes after the synchronizing event. The statement ". . . each replicate is independent of all microarrays preceding and following in time" suggests the experiments are based on one-shot sampling. In contrast, the state-space model with correlation between transcription factors in an earlier time and the regulated genes in a later time fits multi-shot sampling. [4] studied the gene expression difference between leaves at different developmental stages in rice. The 12th, 11th and 10th leaf blades were collected every 3 days for 15 days starting from the day of the emergence of the 12th leaves. While a single plant could provide multiple samples, namely three different leaves at a given sampling time, no plant was sampled at two different times. Thus, from the standpoint of producing time series data, the sampling in this paper was one-shot sampling. [5] devised the phenol-sodium dodecyl sulfate (SDS) method for isolating total RNA from *Arabidopsis*. It reports the relative level of mRNA of several genes for five time points ranging up to six hours after exposure to a synchronizing event, namely being sprayed by a hormone *trans*-zeatin. The samples were taken from the leaves of plants. It is not clear from the paper whether the samples were collected from different leaves of the same plant, or from leaves of different plants.

A.1.3 Animals

 [6] likely used one-shot sampling for their $-24, 60, 120, 168$ hour time series data from mouse skeletal muscle C2C12 cells without specifying whether the samples are all taken from different individuals. [7] produced time series data by extracting cells from a human, seeding the cells on plates, and producing samples in triplicate, at a series of six times, for each of five conditions. Multiple cells are used for each sample with different sets of cells being used for different samples, so this is an example of one-shot sampling of *in vitro* experiment in the sense that each plate of cells is one individual. The use of (five) multiple conditions can serve as a surrogate for a single individual set of cells to gain the effect of multi-shot sampling. Similarly, the data sets produced by [8] involving the plating of HeLa S3 cells can be classified as one-shot samples because different samples are made from different sets of individual cells. Interestingly, the samples are prepared under one set of conditions, so the use of different conditions is not adopted as a surrogate for multi-shot sampling. However, a particular line of cells was selected (HeLa S3) for which cells can be highly synchronized. Also, the paper does not attempt to determine causal interactions.

A.1.4 In silico

The three *in silico* benchmark suites described in the GeneNetWeaver paper on performance profiling of network inference methods [1] consisted of steady state, and therefore one-shot, samples from dynamical models. However, the GeneNetWeaver software can be used to generate multi-shot time series data, and some of that was included in the network inference challenges, DREAM3, DREAM4, and DREAM5 [1, 2].

A.2 On biological replicates

This section summarizes how data from biological replicates is used in the analysis of experimental data for the purpose of differential expression analysis of genes, or inference of gene regulatory networks.

In many biological experiments, independent biological replicates are used to reduce the variation in the measurements and to consequently increase the power of the statistical tests. [9] studies the effect of biological replication together with dye

switching in microarray experiments and recommends biological replication when precision in the measurements is desired. [10] studies the tradeoff between biological replication and sequencing depth under a sequencing budget limit in RNA-seq differential expression (DE) analysis. They propose a metric for cost effectiveness that suggests a sequencing depth of 10 million reads per library of human breast cells and 2–6 biological replicates for optimal RNA-seq DE design. [11] studies the number of necessary biological replicates in RNA-seq differential expression experiments on $S.$ $cerevisiae$ quantitatively with various statistical tools and concludes with the usage of a minimum of six biological replicates. A recent survey [12] suggests a minimum of three replicates for RNA-seq experiments whenever sample availability allows.

The choice of replication strategy depends on how the statistical algorithm uses the replicate data. In many differential analysis software packages replicates are treated as independent samples with identical experimental conditions. For example, in edgeR [13] and sleuth [14] the logarithm of the abundance of gene i in sample m is assumed to be $x_m^* \beta_i$, where x_m is the column vector of design characteristics with respect to p variates for sample m and β_i the column vector of the associated effects of the p variates to gene i. Replicate samples can then be used to expand the design matrix x with identical columns. Note that, as a result, replicates are not necessary for edgeR and sleuth because samples with different design characteristics can all contribute to the estimation of β. It is then not clear whether replicates bring more benefit than the sheer additional amount of data compared to samples under different conditions.

For regulatory network reconstruction there is even less consensus on how replicates should be used. One straightforward way is to reduce the replicates into a single set of data by averaging either directly or after a random resampling of the original replicated data. In this case the mean of the replicates are used as a better estimate of the population than each single replicate, while higher moments of the empirical distribution of the replicates are practically ignored. Another way adopted in [3] is to account for all four potential transitions between two replicates in two adjacent sampling times in their machine learning algorithm due to the one-shot nature of the replicates.

B   Joint A, γ and observation noise estimation

The simulation results in Fig. 4 are for the GLRT estimator that is assumed to know the level of the driving noise random variables (i.e. the $W_j^k(t)$ random variables), which for brevity we normalized to one. To account for not knowing the variance we could multiply the W's by a positive parameter σ, such that σ that is not known to the algorithm. In addition, it was assumed that the GLRT knows the variance of the observation noise, σ_Z^2. This section presents additional simulations, for a GLRT estimator that needs to estimate all four parameters, A, γ, σ, and σ_Z, again for recovery of the sign of regulation for a single gene. The sign error rates with the true $A = 0.1, 0.5$ and one-shot and multi-shot sampling are shown in Fig. 8. It can be seen in comparison with Fig. 4 that the performance of GLRT is not significantly worse without the knowledge of σ and σ_Z.

The scatter plots of the estimate $\hat{A}$ against the actual A and that of the estimate $\hat{\gamma}$ against the actual γ with different joint prior on A and γ and $C = 10$, $R = 3$, $T = 6$ are shown in Figs. 9–13. Each plot is based on 1000 simulations. For all these figures, the plots on the left are for one-shot sampling and the plots on the right are for multi-shot sampling. Also, for all these figures the GLRT algorithm did not have access to σ and σ_Z. Fig. 9 shows scatterplots that indicate how well A and γ can be estimated when the true value of (A, γ) is uniformly distributed over $[-1, 1] \times [0, 1]$. Figs 10–13 show how well A and γ can be estimated when only one of these parameters is uniformly distributed and the other is fixed at 0.1 or 0.5.

(a) One-shot sampling, $A = 0.1$, $\sigma_Z = 1$.

(b) Multi-shot sampling, $A = 0.1$, $\sigma_Z = 1$.

(c) One-shot sampling, $A = 0.5$, $\sigma_Z = 1$.

(d) Multi-shot sampling, $A = 0.5$, $\sigma_Z = 1$.

Fig 8. Error rates of the GLRT for the sign of a single gene with confidence intervals based on 10 000 simulations. All four parameters (A, γ, σ and σ_Z) are unknown to the algorithm.

We can see that for all cases shown, the estimates of A and γ are more accurate for multi-shot sampling vs. one-shot sampling. Comparing Figs. 10 and 11, we see that increasing A slightly helps the estimation of γ. Comparing Figs. 12 and 13, we see that increasing γ greatly improves the estimation of A for one-shot sampling, while it slightly hurts the estimation of A for multi-shot sampling. These observations are consistent with the performance of GLRT for sign recovery seen in Figs. 4 and 8. Namely, for one-shot sampling the performance significantly improves as γ increases, while for multi-shot sampling the performance slightly degrades as γ increases (but it is much worse for $\gamma \approx 1$).

C   Split Gaussian network prior

Let the prior of the adjacency matrix be such that the in-degree of each vertex is uniform over $\{0, 1, \ldots, d_{\max}\}$ for some $d_{\max} \geq 1$, and the set of parents is selected uniformly at random from the rest of vertices given the in-degree. For each edge from vertex i to vertex j, let the corresponding element in the adjacency matrix be split Gaussian distributed with unit gap and unit variance (i.e., $a_{ij} = (0.5 + (0.554553)|Z|)\operatorname{sgn}(Z)$ for $Z \sim \mathcal{N}(0,1)$.). The degradation and the self-regulation are ignored; i.e., $a_{ii} = 0$ for all i. We call this prior the *split Gaussian prior* with max in-degree $d_{\max}$.

References

1. Schaffter T, Marbach D, Floreano D. GeneNetWeaver: *in silico* benchmark generation and performance profiling of network inference methods. Bioinform. 2011;27(16):2263–2270. doi:10.1093/bioinformatics/btr373.

(a) $\hat{A}$ against A with one-shot sampling. $\rho = 0.76$.

(b) $\hat{A}$ against A with multi-shot sampling. $\rho = 0.96$.

(c) $\hat{\gamma}$ against γ with one-shot sampling. $\rho = 0.87$.

(d) $\hat{\gamma}$ against γ with multi-shot sampling. $\rho = 0.89$.

Fig 9. Joint estimation scatter plots with $\sigma = 1$, $\sigma_Z = 1.0$ and $(A, \gamma) \sim \mathsf{Unif}([-1, 1] \times [0, 1])$.

(a) Histogram of $\hat{A}$ for fixed $A = 0.1$ with one-shot sampling.

(b) Histogram of $\hat{A}$ for fixed $A = 0.1$ with multi-shot sampling.

(c) $\hat{\gamma}$ against γ with one-shot sampling and fixed $A = 0.1$. $\rho = 0.87$.

(d) $\hat{\gamma}$ against γ with multi-shot sampling and fixed $A = 0.1$. $\rho = 0.88$.

Fig 10. Joint estimation plots with $\sigma = 1$, $\sigma_Z = 1$, fixed $A = 0.1$ and $\gamma \sim \mathsf{Unif}([0, 1])$.

(a) Histogram of $\hat{A}$ for fixed $A = 0.5$ with one-shot sampling.

(b) Histogram of $\hat{A}$ for fixed $A = 0.5$ with multi-shot sampling.

(c) $\hat{\gamma}$ against γ with one-shot sampling and fixed $A = 0.5$. $\rho = 0.88$.

(d) $\hat{\gamma}$ against γ with multi-shot sampling and fixed $A = 0.5$. $\rho = 0.89$.

Fig 11. Joint estimation plots with $\sigma = 1$, $\sigma_Z = 1$, fixed $A = 0.5$ and $\gamma \sim \mathsf{Unif}([0,1])$.

(a) $\hat{A}$ against A with one-shot sampling and fixed $\gamma = 0.1$. $\rho = 0.41$.

(b) $\hat{A}$ against A with multi-shot sampling and fixed $\gamma = 0.1$. $\rho = 0.98$.

(c) Histogram of $\hat{\gamma}$ for fixed $\gamma = 0.1$ with one-shot sampling.

(d) Histogram of $\hat{\gamma}$ for fixed $\gamma = 0.1$ with multi-shot sampling.

Fig 12. Joint estimation scatter plots with $\sigma = 1$, $\sigma_Z = 1$, fixed $\gamma = 0.1$ and $A \sim \mathsf{Unif}([-1, 1])$.

(a) $\hat{A}$ against A with one-shot sampling and fixed $\gamma = 0.5$. $\rho = 0.82$.

(b) $\hat{A}$ against A with multi-shot sampling and fixed $\gamma = 0.5$. $\rho = 0.97$

(c) Histogram of $\hat{\gamma}$ for fixed $\gamma = 0.5$ with one-shot sampling.

(d) Histogram of $\hat{\gamma}$ for fixed $\gamma = 0.5$ with multi-shot sampling.

Fig 13. Joint estimation plots with $\sigma = 1$, $\sigma_Z = 1$, fixed $\gamma = 0.5$ and $A \sim \mathsf{Unif}([-1, 1])$.

2. Marbach D, Costello JC, Küffner R, Vega NM, Prill RJ, Camacho DM, et al. 487
   Wisdom of crowds for robust gene network inference. Nature methods. 488
   2012;9(8):796–804. doi:10.1038/nmeth.2016. 489

3. Krouk G, Mirowski P, LeCun Y, Shasha DE, Coruzzi GM. Predictive network 490
   modeling of the high-resolution dynamic plant transcriptome in response to 491
   nitrate. Genome Biology. 2010;11(12):R123. doi:10.1186/gb-2010-11-12-r123. 492

4. Yamaoka C, Suzuki Y, Makino A. Differential Expression of Genes of the 493
   Calvin–Benson Cycle and its Related Genes During Leaf Development in Rice. 494
   Plant and Cell Physiology. 2016;57(1):115–124. doi:10.1093/pcp/pcv183. 495

5. Taniguchi M, Kiba T, Sakakibara H, Ueguchi C, Mizuno T, Sugiyama T. 496
   Expression of *Arabidopsis* response regulator homologs is induced by cytokinins 497
   and nitrate. FEBS Letters. 1998;429(3):259–262. 498
   doi:10.1016/s0014-5793(98)00611-5. 499

6. Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, et al. 500
   Transcript assembly and quantification by RNA-Seq reveals unannotated 501
   transcripts and isoform switching during cell differentiation. Nature 502
   Biotechnology. 2010;28(5):511–515. doi:10.1038/nbt.1621. 503

7. Kiselev VY, Juvin V, Malek M, Luscombe N, Hawkins P, Novère NL, et al. 504
   Perturbations of PIP3 signalling trigger a global remodelling of mRNA landscape 505
   and reveal a transcriptional feedback loop. Nucleic Acids Research. 506
   2015;43(20):9663–9679. doi:10.1093/nar/gkv1015. 507

8. Whitfield ML, Sherlock G, Saldanha AJ, Murray JI, Ball CA, Alexander KE, 508
   et al. Identification of genes periodically expressed in the human cell cycle and 509
   their expression in tumors. Molecular Biology of the Cell. 2002;13(6):1977–2000. 510
   doi:10.1091/mbc.02-02-0030. 511

9. Liang M, Briggs AG, Rute E, Greene AS, Cowley AW. Quantitative assessment 512
   of the importance of dye switching and biological replication in cDNA microarray 513
   studies. Physiological Genomics. 2003;14(3):199–207. 514
   doi:10.1152/physiolgenomics.00143.2002. 515

10. Liu Y, Zhou J, White KP. RNA-seq differential expression studies: more 516
    sequence or more replication? Bioinformatics. 2014;30(3):301–304. 517
    doi:10.1093/bioinformatics/btt688. 518

11. Schurch NJ, Schofield P, Gierliński M, Cole C, Sherstnev A, Singh V, et al. How 519
    many biological replicates are needed in an RNA-seq experiment and which 520
    differential expression tool should you use? RNA. 521
    2016;doi:10.1261/rna.053959.115. 522

12. Conesa A, Madrigal P, Tarazona S, Gomez-Cabrero D, Cervera A, McPherson A, 523
    et al. A survey of best practices for RNA-seq data analysis. Genome Biology. 524
    2016;17(1):13. doi:10.1186/s13059-016-0881-8. 525

13. McCarthy DJ, Chen Y, Smyth GK. Differential expression analysis of multifactor 526
    RNA-Seq experiments with respect to biological variation. Nucleic Acids 527
    Research. 2012;40(10):4288–4297. doi:10.1093/nar/gks042. 528

14. Pimentel H, Bray NL, Puente S, Melsted P, Pachter L. Differential analysis of 529
    RNA-seq incorporating quantification uncertainty. Nature Methods. 530
    2017;14:687–690. doi:10.1038/nmeth.4324. 531

15. Poor HV. An Introduction to Signal Detection and Estimation. Springer-Verlag   532
    New York; 1994.   533

16. Sun J, Taylor D, Bollt EM. Causal network inference by optimal causation   534
    entropy. SIAM Journal on Applied Dynamical Systems. 2015;14(1):73–106.   535
    doi:10.1137/140956166.   536

17. Emad A, Milenkovic O. CaSPIAN: A causal compressive sensing algorithm for   537
    discovering directed interactions in gene networks. PloS one. 2014;9(3):e90781. 538

18. Bertsimas D, King A, Mazumder R. Best subset selection via a modern   539
    optimization lens. Ann Statist. 2016;44(2):813–852. doi:10.1214/15-AOS1388.   540

19. Bento J, Ibrahimi M, Montanari A. Learning Networks of Stochastic Differential 541
    Equations. In: Advances in Neural Information Processing Systems (NIPS); 2010.   542
    p. 172–180. Available from: `http://papers.nips.cc/paper/`   543
    `4055-learning-networks-of-stochastic-differential-equations.pdf`.   544