Sonic hedgehog signaling in astrocytes mediates cell type-specific synaptic organization

AUTHORS: Steven Hill[1,§], Andrew Blaeser[1,§], Austin A. Coley[2], Yajun Xie[3], Katherine A. Shepard[1], Corey C. Harwell[3], Wen-Jun Gao[2], A. Denise R. Garcia[1,2*]

[1]Department of Biology, Drexel University, Philadelphia, PA 19104, USA

[2]Department of Neurobiology and Anatomy, Drexel University College of Medicine, Philadelphia, PA 19129, USA

[3]Department of Neurobiology, Harvard Medical School, Boston, MA 02115, USA

[§]Authors contributed equally to this work.

Correspondence:

A. Denise R. Garcia

Department of Biology

Department Neurobiology and Anatomy

Drexel University

3245 Chestnut St. PISB 422

Philadelphia, PA 19104

adg82@drexel.edu

Astrocytes have emerged as integral partners with neurons in regulating synapse formation and function, but the mechanisms that mediate these interactions are not well understood. Here, we show that Sonic hedgehog (Shh) signaling between neurons and astrocytes is required for establishing structural organization and remodeling of cortical synapses in a cell type-specific manner. In the postnatal cortex, Shh signaling is active in a subpopulation of mature astrocytes localized primarily in deep cortical layers. Selective disruption of Shh signaling in astrocytes produces a dramatic increase in synapse number specifically on layer V apical dendrites that emerges during adolescence and persists into adulthood. Dynamic turnover of dendritic spines is impaired in mutant mice and is accompanied by an increase in neuronal excitability and a reduction of the glial-specific, inward-rectifying K^+ channel Kir4.1. These data identify a critical role for Shh signaling in astrocyte-mediated modulation of neuronal activity required for sculpting synapses.

INTRODUCTION

The organization of synapses into the appropriate number and distribution occurs through a process of robust synapse addition followed by a period of refinement during which excess synapses are eliminated. Failure to establish or maintain appropriate synaptic organization is a hallmark of many neurodevelopmental disorders [1]. Considerable evidence now shows that, together with neurons, astrocytes are critical regulators of synaptic connectivity and function [2–4]. Astrocytes interact intimately with synapses to regulate their formation, maturation, and function, and a growing number of astrocyte-secreted proteins that directly mediate synapse formation and elimination have been identified [5–8]. Nevertheless, despite the remarkable progress in our understanding of the molecular interactions between astrocytes and synapses that facilitate synaptogenesis, the underlying signaling programs mediating astrocyte-dependent regulation of synapse organization remain poorly understood.

The molecular signaling pathway Sonic hedgehog (Shh) governs a broad array of neurodevelopmental processes in the vertebrate embryo, including morphogenesis, cell proliferation and specification, and axon pathfinding [9,10]. However, Shh activity persists in multiple cell populations in the postnatal and adult CNS, including progenitor cells, as well as in differentiated neurons and astrocytes [11–15], where novel and unexpected roles for Shh activity are emerging [16]. Following injury, Shh has been shown to mitigate inflammation [17,18], and in the cerebellum, Shh derived from Purkinje neurons instructs phenotypic properties of mature Bergmann glia [19]. In the postnatal cortex, Shh is required for establishing local circuits between two distinct projection neuron populations [14]. Shh produced by layer V neurons guides the formation of synaptic connections to its layer II/III presynaptic partners, which transduce the Shh signal through non-canonical, Gli-independent mechanisms. We have previously shown that Shh signaling is also active in a discrete subpopulation of cortical astrocytes [15], suggesting that Shh signaling mediates both homotypic and heterotypic cellular interactions. Astrocytes engaging in Shh activity are identified by expression of Gli1, a transcriptional effector of canonical Shh signaling [9]. Whether Shh signaling in cortical astrocytes plays a role in synaptic organization of neurons is not known.

In this study, we examined the organization and dynamics of dendritic spines on cortical neurons following selective disruption of Shh signaling in astrocytes. Dendritic spines are the structural hosts of most excitatory synapses and play an important role in the organization and function of neural circuits. We show that deep layer neurons exhibit long-lasting aberrations in the density and turnover of dendritic spines that emerge during postnatal development when astrocytic Shh signaling is disrupted. Notably, these perturbations in synaptic organization are not observed in upper layer neurons where Gli1 astrocytes are relatively sparse. Chronic *in vivo* imaging of dendritic spines reveals that mutant mice exhibit lower rates of spine turnover, suggesting impaired structural plasticity. In addition, these mice show a pronounced deficit in expression of the glial-specific inward-rectifying K^+ channel, Kir4.1, and concomitant hyperexcitability of cortical neurons. Taken together, these data demonstrate that astrocytes act as key modulators of neural activity and structure during postnatal development, and further establish Shh signaling as a fundamental mediator of synaptic connectivity.

RESULTS

Gli1 astrocytes exhibit a distinct laminar distribution in the adult cortex

In the mature cortex, a subpopulation of astrocytes express the transcription factor Gli1, indicating active Shh signaling [15]. Notably, the distribution of Gli1 astrocytes throughout the cortex is non-uniform, showing a laminar-specific pattern (**Figure 1**). We analyzed the laminar distribution of Gli1 astrocytes in adult *Gli1^{CreER/+};*Ai14 mice, in which tamoxifen administration promotes Cre-mediated recombination of the fluorescent tdTomato reporter protein, permanently marking Gli1-expressing cells. The vast majority of marked cells were observed within layers IV and V, with few marked cells observed in layers II/III or VI (**Figure 1**). We performed immunostaining with the pan-astrocytic marker S100β and quantified the fraction of marked astrocytes in cortical layers (**Figure 1**). This analysis showed that 44% of astrocytes in layer IV express Gli1 (**Figure 1**), while 36% of astrocytes in layer V express Gli1. Interestingly, the distribution of Gli1 astrocytes in layer V was not homogenous, showing an enrichment of marked cells in layer Vb, and a distinctive paucity of marked cells in layer Va (**Figure 1**), consistent with the localization of Shh producing neurons in layer Vb [14,15]. We analyzed the fraction of Gli1 astrocytes in each sublayer of layer V and found that 75% of marked cells were found in layer Vb whereas only 25% were found in layer Va. Layers VI and II/III showed the lowest fractions of marked astrocytes, with only 11% and 22% of S100β cells expressing tomato, respectively. These data demonstrate that Gli1 astrocytes are preferentially localized in deep cortical layers and suggest their interactions with local neurons may differ from that of Gli1-negative astrocytes in superficial layers.

Figure 1: Gli1 astrocytes are distributed in a laminar-specific manner. (A) Tamoxifen-induced reporter expression in *Gli1^{CreER};*Ai14 mice reveals a laminar distribution of cells undergoing active Shh signaling (red). Scale bar, 250 µm. **(B)** Gli1+ cells (red) are a subset of all astrocytes (S100β, green, see inset; scale bar, 12.5 µm). **(C)** Layer analysis of Gli1 astrocytes (red) with DAPI and the deep layer marker Ctip2 (gray) illustrate their layer-specific organization. Scale bar, 125 µm. **(D)** The fraction of astrocytes expressing Gli1 in each cortical layer. Data points indicate individual animals, bars represent mean ± SEM. Statistical signficance was assessed by one-way ANOVA with Tukey's post hoc test for multiple comparisons. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.01$. **(E)** The fraction of Gli1 astrocytes in sublayers Va and Vb. Bar graphs represent mean value ± SEM. n = 4 animals.

Shh signaling is required for cell type-specific synaptic connectivity

Transduction of canonical Shh signaling begins with binding of Shh to its transmembrane receptor, Patched1 (Ptch), relieving inhibition of a second, obligatory co-receptor Smoothened (Smo). To

investigate the role of Shh signaling in astrocyte function, we performed conditional knockout of Smo selectively in astrocytes using GFAP-Cre transgenic mice in which Cre expression is regulated by the full-length mouse GFAP promoter[20] (mGFAP Smo CKO). Because mGFAP-mediated Cre recombination targets astrocyte progenitors expressing GFAP which are present at birth, we can effectively delete Shh activity in all cortical astrocytes [21]. Nevertheless, because Shh signaling occurs in postnatal neurons, and because recombination in cortical neurons has been observed in some GFAP-Cre transgenic lines[22], we performed further validation of the Cre-mediated recombination pattern by crossing mGFAP-Cre mice with Ai14 reporter mice (*mGFAP-Cre;*Ai14). We analyzed the identity of tdTomato-expressing cells (tomato) in the somatosensory cortex of *mGFAP-Cre;*Ai14 reporter mice by immunohistochemistry with cell type-specific markers (**Figure 2 – figure supplement 1**). Although individual floxed alleles possess distinct recombination efficiencies, analysis of recombination in a reporter line provides a useful approximation of the expression pattern of a given Cre driver. The vast majority of tomato-positive cells showed a bushy morphology, typical of protoplasmic astrocytes (**Figure 2 – figure supplement 1**). Double staining with S100β showed that nearly 70% of tomato-expressing cells were co-labeled, identifying them as astrocytes. A small fraction (9%) of tomato cells were co-labeled with NeuN, identifying them as neurons. Conversely, only 1.3% of cortical neurons were labeled with tomato, and these were localized primarily to upper layers (**Figure 2 – figure supplement 1**). We also identified a small fraction (6%) of tomato cells as oligodendrocytes. Importantly, although a minor fraction of recombined cells was identified as neurons or oligodendrocytes, nearly all cortical astrocytes analyzed expressed the tdTomato reporter protein (95%), suggesting that this mGFAP-Cre driver is an effective tool for selective deletion of Smo in all cortical astrocytes.

Figure 2 – Figure supplement 1: Gfap-Cre recombination occurs primarily in astrocytes. (A) Expression of the astrocyte marker S100β and tdTomato in somatosensory cortex. Scale bar, 25 μm. **(B)** Expression of the neuronal marker NeuN and tdTomato in somatosensory cortex. **(C)** Expression of the oligodendrocyte marker CAII and tdTomato in somatosensory cortex. White arrows show cells that express the indicated marker and are derived from a GFAP+ cell. Blue arrows show cells that express the indicated marker but are not derived from a GFAP+ cell. Yellow arrows show cells derived from a GFAP+ cell that does not express the given markers. **(D-F)** Fraction of recombined cells identified as astrocytes (D), neurons (E), or oligodendrocytes (F) in the cortex of adult *Gfap-Cre;*Ai14 mice. Data points represent individual brain sections analyzed. Bar graph represent mean ± SEM. *n* = 2 animals.

During postnatal development, Shh is required for establishing synaptic connectivity between layer V neurons and their presynaptic partners in layer II/III, which is mediated by Gli-independent, non-canonical signaling [14]. To examine whether canonical, Gli-mediated Shh signaling in astrocytes plays a role in establishing synaptic connectivity of cortical neurons, we evaluated spine density of apical dendrites on layer V neurons in the somatosensory cortex of mGFAP Smo CKO mice across postnatal

development. Dendritic spines receive
the vast majority of excitatory input,
thereby serving as a useful readout of
synaptic connectivity[23]. To visualize
neurons for reliable identification and
tracing, mGFAP Smo CKO mice were
crossed with Thy1-GFPm transgenic mice,
which sparsely express GFP in a subset of
layer V cortical neurons [24]. Dendritic
spines undergo a period of dynamic
reorganization during postnatal
development during which there is an
initial overproduction of spines during
the first 2-3 weeks of postnatal
development, followed by a period of
synaptic pruning that refines the precise
connectivity of cortical circuits [25]. At P14,
spine density was comparable between
mGFAP Smo CKO mice and wild type (WT)
littermate controls, suggesting that early
synaptogenesis does not require
astrocytic Shh signaling (0.58 ± 0.03
spines/µm and 0.56 ± 0.01 spines/µm,
WT [*n* = 7 cells] and mGFAP Smo CKO [*n* =

Figure 2: Loss of Shh signaling in astrocytes results in increased spine density on layer V neurons. (**A-B**) GFP immunostaining of representative apical dendritic segments from layer V neurons in a WT (A) and mGfap Smo CKO (B) mice at P21. Arrows indicate individual spines. Scale bar, 5 µm. (**C**) Spine density of apical dendrites from layer V neurons across various postnatal ages and in adult mice. Data points represent individual cells, bars represent mean ± SEM. Statistical analysis performed by two-way ANOVA with Tukey's post-hoc test for multiple comparisons. *, $p < 0.05$; **, $p < 0.01$; ****, $p < 0.0001$. Significance is stated as mGFAP Smo CKO compared to WT at a given age. (**D-E**) Spine density of apical dendrites from CA1 pyramidal neurons (D) and layer II/III (E) in WT and mGfap Smo CKO mice. (**F**) Spine density of apical dendrites from layer V neurons in adult CamKIIa Smo CKO mice. Data points represent individual cells, bars represent mean ± SEM. Statistical signfiiance assessed by Student's t-test. *n* ≥ 3 animals per genotype for all ages but P14, where *n* = 2 animals per genotype.

6 cells], respectively, *p* = 0.64, 2 animals per genotype). Between P14 and P21, spine density remained
stable in WT, but showed a dramatic increase at P28 (**Figure 2**). This period of synapse addition was
followed by a steady reduction in spine density at P42 and into
adulthood (≥P90), reflecting the developmental elimination of
spines. In contrast, mGFAP Smo CKO mice showed an accelerated
timeline of spine addition in which spine density increased
dramatically between P14 and P21 (**Figure 2**). Spine density was
significantly higher at P21 in mGFAP Smo CKO mice compared to
controls (0.53 ± 0.05 spines/µm and 0.76 ± 0.06 spines/µm, WT and
mGFAP Smo CKO, respectively, *p* = 0.008, *n* = 9 cells from 3 animals
per genotype). Although there was a modest reduction in spine
density as animals matured, spine density remained elevated in
adult mGFAP Smo CKO mice compared to WT controls (0.36 ± 0.02
spines/um and 0.60 ± 0.02 spines/µm in WT [*n* = 24 cells] and
mGFAP Smo CKO [*n* = 23 cells], respectively, *p* < 0.0001, 4 animals
per genotype; **Figure 2**). Thus, whereas WT mice experience a 49%
reduction in spine density from the peak of spine density at P28 to
adulthood, mGFAP Smo CKO mice exhibit a 22% reduction from

Figure 2 – figure supplement 2: Spine density on layer V basal dendrites are not affected in adult mGFAP Smo CKO mice. (**A-B**) Representative images of basal dendrites analyzed from layer V neurons expressing *Thy1GFPm* in both WT (*Smo^fl/fl^*) and mGFAP Smo CKO (*Gfap-Smo^fl/fl^*) animals. Scale bar, 5 µm. (**C**) Graph comparing the density of spines on basal dendrites from WT and mGFAP Smo CKO animals. Data points represent individual cells, bars represent mean ± SEM. Statistical significance was assessed using unpaired Student's t-test. *n* = 3 animals per genotype.

their peak at P21 to adulthood. Interestingly, we find no difference in basal dendrite spine density in
mGFAP Smo CKO mice (0.60 ± 0.04 spines/um and 0.62 ± 0.04 spines/μm in WT and mGFAP Smo CKO,
respectively, *p* = 0.71, *n* = 10 cells from 3 animals per genotype; **Figure 2 – figure supplement 1**). This is
in contrast to embryonic deletion of Shh from neurons using the EmxCre driver, which produces a
reduction in spine density of basal dendrites on layer V neurons [14]. These data suggest that while the
early stages of synapse formation proceed independently of astrocytic Shh signaling, the maturing
cortical circuit requires intact Shh activity in astrocytes for the developmental pruning of excess
synapses necessary to achieve its mature organization.

To confirm that elevated spine density was due specifically to a loss of astrocytic Shh signaling, we
interrogated the spine density of pyramidal cells in two different regions where Gli1 astrocytes are
relatively sparse (see **Figure 1**). Pyramidal neurons in layer II/III of WT mice showed a higher spine
density than layer V neurons (0.79 ± 0.06 spines/μm, *n* = 7 cells, 4 animals) consistent with previous
studies [26]. However, this was not significantly different from the spine density observed in mGFAP Smo
CKO mice (0.76 ± 0.06 spines/μm, *p* = 0.76, *n* = 9 cells, 5 animals; **Figure 2**). We also analyzed pyramidal
neurons in the hippocampus. Although the dentate gyrus of the hippocampus harbors a population of
Gli1-expressing adult neural progenitor cells [12,27], mature, differentiated astrocytes expressing Gli1 are
relatively sparse (**Figure 1**). The spine density of CA1 pyramidal neurons in adult mice was higher than in
cortical neurons, consistent with previous studies [28,29]. However, there was no significant difference in
spine density between mGFAP Smo CKO mice and WT controls (0.97 ± 0.05 spines/μm and 0.90 ± 0.05
spines/μm in WT and mGFAP Smo CKO, respectively, *p* = 0.32, *n* = 11 cells from 3 animals per genotype;
Figure 2). Our analysis of the recombination pattern of the mGFAP-Cre driver showed that some cortical
neurons undergo recombination (**Figure 2 – figure supplement 1**). To rule out the possibility that Smo
deletion in a small population of neurons is responsible for the elevated spine density of layer V
neurons, we deleted Smo in excitatory neurons using the CamKIIa Cre driver (CK2 Smo CKO), and
evaluated spine density of layer V neurons in adult mice. We found no significant difference in spine
density between CK2 Smo CKO mice and littermate controls (0.77 ± 0.07 spines/um and 0.59 ± 0.06
spines/um in WT [*n* = 10 cells] and CK2 Smo CKO [*n* = 9 cells], respectively, *p* = 0.07, 3 animals per
genotype; **Figure 2**). Taken together, these data suggest that Shh activity in astrocytes is necessary for
the developmental reorganization of dendritic spines specifically on layer V cortical neurons. Notably,
we observed elevated spine density along the apical dendrites of layer V neurons, which traverse layer
II/III, where few Gli1 astrocytes are found. This suggests that the modification of synapse organization
mediated by Gli1 astrocytes does not occur through direct astrocyte-synapse interactions.

Synapses are more stable in mGFAP Smo CKO mice

Dendritic spines are dynamic structures that undergo rapid formation and elimination during postnatal
cortical development. As cortical circuits mature, the fraction of spines undergoing dynamic turnover
declines concomitant with an increase in spine stability [25,26]. The dynamic turnover of spines has long
been considered a structural correlate of synaptic plasticity and can be regulated in an activity-
dependent manner [30–33]. To evaluate the role of astrocytic Shh signaling in mediating spine dynamics,
we performed repeated *in vivo* imaging of the apical tufts of layer V neurons in the somatosensory

cortex through a cranial window [34]. We confirmed the layer V identity of imaged neurons by following individual dendritic segments to their soma and created 3D reconstructions of imaged neurons. Only dendrites with soma in layer V, typically 500 μm – 600 μm from the surface of the brain, were analyzed. In Thy1-GFPm mice, expression of GFP is developmentally regulated such that at P14, the intensity of GFP expression and the density of fluorescently labeled neurons within the 3 mm cranial window is insufficient for reliable imaging, as has been previously reported [26]. However, by P17, GFP expression was sufficiently dense and bright to enable reliable imaging. We analyzed the fraction of spines undergoing dynamic turnover over 2 days in young mice at P17-P21 and P28-P32. In WT mice, the turnover ratio was 0.14 at P17-P21 and 0.09 in P28-P32 mice, though this was not statistically significant (p = 0.12, n = 5 mice; **Figure 3**). In mGFAP Smo CKO mice, however, the turnover ratio was lower at P17-P21 than in WT mice, and remained constant at P28-P32 (0.11 and 0.10, P17-P21 and P28-P32, respectively, n = 4 mice; **Figure 3**). This suggests that dendritic spines in mGFAP Smo CKO mice stabilize earlier than those in WT mice. Consistent with this, we found that the fraction of filopodia was lower in juvenile mGFAP Smo CKO mice compared to WT controls (**Figure 3**). Filopodia are structural precursors of dendritic spines and exhibit higher rates of dynamic turnover than mature spines [35,36]. Accordingly, the density of filopodia declines as the cortex matures [25,37]. Consistent with this, the fraction of all protrusions in juvenile WT mice identified as filopodia was 24%, but that fraction declined significantly to 15% in adults (p = 0.0016; **Figure 3**). In contrast, we did not

Figure 3. Loss of astrocytic Shh signaling impairs structural plasticity. (**A-C**) Example dendrite segment imaged repeatedly over 6 weeks. Day 0 indicates the first day of imaging, subsequent imaging days indicated. White arrows, stable spines; cyan arrows, transient spines; orange arrows, filopodia. (**D**) Turnover ratios in WT (n = 5 animals) and mGFAP Smo CKO (n = 4 animals) juvenile mice analyzed over 2 days. (**E**) Turnover ratios in WT (n = 6 animals) and mGFAP Smo CKO (n = 9 animals) adult mice analyzed over 7 days. Data points represent individual cells, bars represent mean ± SEM. Statistical analysis by two-way ANOVA (D) and unpaired Student's t-test (E). (**F**) Fraction of protrusions identified as filopodia in in juvenile (WT, n = 11 animals; mGFAP Smo CKO, n = 10 animals) and adult mice (WT, n = 16: mGFAP Smo CKO, n = 7 animals). Statistics by two-way ANOVA, Tukey's post hoc test. (**G**) Comparison of spine survival curves for WT and mGFAP Smo CKO neurons. Each dashed curve represents the curve from an individual mouse (WT, n = 7 animals; mGFAP Smo CKO, n = 5 animals). Solid curves represent best-fits to exponential decay models. (**H**) The fraction of spines belonging to the stable, recurrent, and transient populations. Statistical significance was assessed by Student's t-test for each class (stable, recurrent, or transient; n = 7 and 8 animals, WT and mGFAP Smo CKO, respectively). Stable, p = 0.0485; Recurrent, p = 0.1023; Transient, p = 0.7709.

observe this age-dependent decline in mGFAP Smo CKO mice (16% and 13% in juvenile and adult mGFAP
Smo CKO mice, respectively). The filopodial fraction in juvenile mGFAP Smo CKO mice was already
significantly below WT levels (16% vs 24% in mGFAP Smo CKO and WT mice, respectively) and similar to
adult WT levels (15% and 13% in WT and mGFAP Smo CKO adult mice, respectively). It is well established
that the rate of synaptic turnover declines considerably as animals mature, reflecting an increase in
stability of synaptic connections [25,26]. These results indicate that disruption of Shh signaling selectively in
astrocytes accelerates the stabilization of dendritic spines on layer V cortical neurons.

We next sought to investigate the long-term stability of individual spines in adult mice by imaging
dendrites weekly for up to 6 weeks. We investigated the fraction of spines identified on the first day of
imaging that persisted in subsequent imaging sessions and calculated the survival fraction by fitting to
an exponential decay model (**Figure 3;** see Methods). This analysis revealed a larger proportion of long-
lived stable spines in mGFAP Smo CKO mice compared to WT control (71% and 79% in WT [n = 7 mice]
and mGFAP Smo CKO [n = 5 mice], respectively, p = 0.0068, Extra sum-of-squares F test; **Figure 3**),
suggesting an increase in stability of dendritic spines. Interestingly, among the population of dynamic
spines, we found two distinct populations consisting of transient spines, which disappeared and did not
reappear for the duration of the study, and recurrent spines, which disappeared and then subsequently
reappeared. Our data showed that the shift towards increased spine stability in the mGFAP Smo CKO
neurons was entirely due to a reduction in the proportion of recurrent spines, with nearly identical
fractions of transient spines observed across both genotypes (**Figure 3**). Taken together, these data
suggest that structural plasticity is impaired in mGFAP Smo CKO mice. This deficit in structural plasticity
emerges during the third week of postnatal development and persists into adulthood. This suggests
that Shh signaling between neurons and astrocytes is required for refining synaptic connections in the
cortex.

Shh signaling regulates Kir4.1 expression in cortical astrocytes

In both the developing and adult CNS, astrocytes directly eliminate synapses in an activity-dependent
manner through MEGF10 and MERTK, two phagocytic receptors enriched in astrocytes [38]. To investigate
whether these genes are negatively
regulated in mGFAP Smo CKO mice, we
examined their expression in the cortex
of adult mice by quantitative PCR (qPCR).
There was no difference in expression
level of *Mertk* between WT and mGFAP
Smo CKO mice (dCq values: 5.99 ± 0.16
and 5.83 ± 0.06 in WT [n = 4 mice] and
mGFAP Smo CKO [n = 5 mice]
respectively, p = 0.35; **Figure 4 – figure**
supplement 1). Expression of *Megf10*
showed a trend towards increased
expression (dCq values: 8.13 ± 0.29 and

Figure 4 – figure supplement 1: Quantitive PCR analysis of known spine and synapse modulators show no change in expression in mGFAP Smo CKO mice. (A) Quantitive PCR shows a slight, but not significant, increase in *Megf10* and no change in *Mertk* in mGFAP Smo CKO animals compared to controls (n = 4 and 5 samples of cortical RNA from WT and mGFAP Smo CKO animals, respectively). Statistical significance assessed by unpaired Student's t-test of the quantification cycles for each sample. Error bars represent standard error of the mean. **(B)** Droplet digital PCR shows no difference in the number of trancripts of either *GLT-1* or *GLAST* in mGFAP Smo CKO animals (n = 3 cortical RNA samples per genotype). Statistical significance assessed by unpaired Student's t-test of the copy number for each target in each genotype. Error bars represent standard error of the mean.

7.53 ± 0.11 in WT [*n* = 4 mice] and mGFAP Smo CKO [*n* = 5 mice], respectively; **Figure 4 – figure**

supplement 1), though this difference was not statistically significant (*p* = 0.075). These data suggest

that direct engulfment of synapses by astrocytes through these pathways is not regulated by Shh

signaling and is not responsible for the increased spine density seen in mGFAP Smo CKO animals.

In the mature

cerebellum, Shh

signaling between

Purkinje neurons

and Bergmann glia

regulates

expression of the

glial specific,

inward-rectifying

K^+ channel, Kir4.1

[19]. In the cortex,

the distribution of

Kir4.1 exhibits a

laminar pattern

similar to that of

Gli1 astrocytes,

such that

expression is

enriched in layers

IV and Vb, and

reduced in layer Va

(**Figure 4**). High

resolution,

confocal analysis

showed that Kir4.1

is localized in the

Figure 4: Shh signaling regulates Kir4.1 expression in the cortex. **(A)** Fluorescence micrographs showing the distribution of Gli1 astrocytes (A, red) and Kir4.1 (A, green) in the cortex of an adult *Gli1Cre^{ER}*;Ai14 mouse counterstained with DAPI (A, blue). **(B-E)** Immunofluorescence for Kir4.1 in the cortex of adult WT (B), mGfap Smo CKO (C), CK2 Smo CKO (D) and GLAST Ptch CKO mice (E). Scale bar, 125 μm. **(F-G)** Confocal images of Gli1 (F, red) Kir4.1 (green) and NeuN (white) from adult WT (F), mGfap SmoCKO (G), or CK2 SmoCKO (H) mice. Scale bar, 25 μm. **(I)** Gene expression levels in the cortex of Kir4.1 from WT (*n* = 6 animals) and mGfap Smo CKO or CK2 Smo CKO mice (*n* = 4 each). Data points represent individual animals and bar graphs represent mean value ± SEM. Statistical significance was assessed by one-way ANOVA with Tukey's test for multiple comparisons. Significance stated on the graph is compared to WT.

processes of Gli1 astrocytes (**Figure 4**). Kir4.1 expression is localized to astrocytic endfeet and is found

surrounding neuronal somata in the spinal cord and brain [39–41]. In WT mice, expression of Kir4.1

surrounded many NeuN-positive neuronal somata in layer V (**Figure 4**). In mGFAP Smo CKO mice,

however, there was a pronounced reduction in the expression of Kir4.1 throughout the cortex, and peri-

somal Kir4.1 expression was severely diminished (**Figure 4**). In order to quantify expression levels of

Kir4.1, we measured transcript abundance in the cortex of mGFAP Smo CKO mice and WT controls by

droplet digital PCR (ddPCR). In mGFAP Smo CKO mice, there was a 50% reduction in *Kir4.1* transcripts

compared to WT controls ($1.99 \times 10^6 \pm 1.93 \times 10^5$ copies/μl and $9.92 \times 10^5 \pm 8.49 \times 10^4$ copies/μl in WT [*n*

= 6 mice] and mGFAP Smo CKO [*n* = 4 mice], respectively, *p* = 0.02). There was no difference in *Kir4.1*

expression in CK2 Smo CKO mice compared to WT ($1.88 \times 10^6 \pm 3.77 \times 10^5$ copies/μl, *p* = 0.95, *n* = 4 mice;

Figure 4), suggesting that downregulation of *Kir4.1* is due to astrocyte-specific, and not neuronal,

disruption of Shh signaling. Conversely, GLAST-CreER-mediated deletion of the Shh receptor *Ptch1*, a

negative regulator of Shh signaling, dramatically upregulated Kir4.1 expression in the cortex (Ptch CKO;

Figure 4). Kir4.1 expression is associated with glutamate uptake both *in vitro* and *in vivo* [42–44]. We

therefore measured expression levels of the astrocyte-specific glutamate transporters *GLAST* and *GLT1*.

There was no difference in expression of these genes in mGFAP Smo CKO mice compared to WT (<u>*GLT1*</u>:

$1.08 \times 10^7 \pm 6.6 \times 10^5$ copies/µl and $9.58 \times 10^6 \pm 2.1 \times 10^5$ copies/µl in WT and mGFAP Smo CKO,

respectively, p = 0.15, *n* = 3 animals per genotype; <u>*GLAST*</u>: $7.4 \times 10^6 \pm 4.4 \times 10^5$ copies/µl and $6.7 \times 10^6 \pm$

6.5×10^5 copies/µl in WT and mGFAP Smo CKO, respectively, *p* = 0.41, *n* = 3 animals per genotype; **Figure**

4 – figure supplement 1). It should be noted that Kir4.1 was not restricted to Gli1 astrocytes in layers IV

and Vb. Nevertheless, single transcript quantification of Kir4.1 transcripts in mGFAP Smo CKO show a

pronounced reduction, suggesting that Kir4.1 expression in cortical astrocytes is regulated, in part, by

Shh signaling.

Neurons exhibit

hyperexcitability in

mGFAP Smo CKO mice

Astrocytes play critical

roles in buffering K⁺ in the

extracellular space

through Kir4.1, and loss of

Kir4.1 produces

disturbances in neuronal

firing properties [39,40,43–47].

To examine whether

neurons in mGFAP Smo

CKO mice exhibit

disruptions in

physiological activity, we

performed whole-cell

patch clamp recordings

and first examined action

potential firing and

membrane properties of

layer V pyramidal neurons

from coronal brain slices

at P21. Using current

clamp, we recorded action

potential spikes per

injected current at

multiple step currents

Figure 5. mGFAP Smo CKO neurons exhibit a significant increase in excitability and synaptic transmission. (A) Example traces of action potentials from layer V pyramidal neurons in WT and mGFAP Smo CKO mice at P21. Samples of action potential spikes (dashed lines, lower panel) describe AP threshold, AP peak amplitude, and AP ½ width of pyramidal neuron. **(B)** Line graph displays relationship between action potential spike numbers (y-axis) versus current injection; mGFAP Smo CKO neurons exhibit an increase in AP spikes (>500 pA) compared to WT neurons. Statistical signifcance was assessed by two-way ANOVA with Sidak's test for multiple comparisons and stated as mGFAP Smo CKO compared to WT at a given current (n = 11 and 7 cells from WT and mGFAP Smo CKO, respectively). **(C-F)** Bar graphs describe an increase in total AP spike numbers (C), reduction in AP threshold (mV) (D), reduction in AP 1/2 width (E), and increase in AP peak amplitude (pA) (F) in mGFAP Smo CKO neurons. **(E)** Example traces of sEPSCs from layer V pyramidal neurons recorded in the presence of picrotoxin from WT and mGFAP Smo CKO mice at P21. **(F)** Summary graphs of sEPSC frequency and amplitude (n = 14 and 11 cells from WT and mGFAP Smo CKO animals, respectively). **(G)** Example traces of mEPSCs from layer V pyramidal neurons recorded in the presence of picrotoxin and TTX. **(H)** Summary graphs of sEPSC frequency and amplitude (n = 16 cells per genotype). mGFAP Smo CKO neurons exhibit increases in both frequency and amplitude in sEPSCs and mEPSCs recordings. Statistical significance was assessed by unpaired Student's t-test. All data are from at least 3 animals per genotype.

from -300 pA to +650 pA. Our
results revealed a significant
increase in spike numbers
with high current injections
(>500 pA) and total overall
spikes in mGFAP Smo CKO
animals compared to WT
control (326 ± 54 and 574 ± 71
spikes in WT [n = 11 cells from
4 animals] and mGFAP Smo

	WT (n = 11 cells)	mGFAP Smo CKO (n = 7 cells)	*P* value
Input resistance (MΩ)	130.6 ± 21.52	97.93 ± 11.49	0.15
Resting membrane potential (mV)	-59.37 ± 2.059	-63.73 ± 3.248	0.25
Tau (ms)	11.76 ± 0.71	11.75 ± 1.398	0.79

Figure 5 – figure supplement 1. Membrane properties of mGFAP Smo CKO layer V pyramidal neurons. There is a significant decrease in input resistance (MΩ) in mGFAP Smo CKO neurons compared to WT neurons. No significant differences were observed in resting membrane potential (mV), rheobase (pA), and tau (ms) in mGFAP Smo CKO neurons. Statistical significance was assessed by Mann-Whitney U test (input resistance and tau) and unpaired Student's t-test (resting membrane potential).

CKO [n = 7 cells cells from 3 animals], respectively, p = 0.01; **Figure 5**). We also measured other
membrane properties, including input resistance, resting membrane potential and tau, and found no
significant difference between mGFAP Smo CKO and WT neurons (**Figure 5 – figure supplement 1**).
However, we observed a significant decrease in action potential threshold (-30.6 ± 2.0 and -37.9 ± 2.0
mV in WT and mGFAP Smo CKO, respectively, p = 0.0441), accompanied by an increase in action
potential amplitude (59.6 ± 2.6 and 73.3 ± 3.0 pA in WT and mGFAP Smo CKO, respectively, p = 0.0083),
and a trending decrease in action potential ½ width (1.0 ± 0.92 and 0.76 ± 0.02 ms in WT and mGFAP
Smo CKO, respectively, p = 0.05) in mGFAP Smo CKO neurons, consistent with an increase in neuron
excitability (**Figure 5**). Together, these results suggest mGFAP Smo CKO neurons exhibit an increase in
neuronal excitability consistent with excess extracellular K^+.

To examine excitatory synaptic transmission, we recorded spontaneous and miniature excitatory
postsynaptic currents (sEPSCs and mEPSCs) in layer V pyramidal neurons. We found an increase in both
the frequency (0.74 ± 0.08 and 1.18 ± 0.19 Hz in WT and mGFAP Smo CKO, respectively, p = 0.042) and
amplitude (8.54 ± 0.38 and 11.33 ± 0.91 pA in WT and mGFAP Smo CKO, respectively, p = 0.008) of
sEPSCs in slices from mGFAP Smo CKO mice compared to WT control slices (n = 16 cells from 4 animals
per genotype; **Figure 5**). Furthermore, we observed an increase in the frequency (0.40 ± 0.05 and 1.10 ±
0.23 Hz in WT and mGFAP Smo CKO, respectively, p = 0.003) and amplitude (7.54 ± 0.57 and 10.34 ± 0.74
pA in WT and mGFAP Smo CKO, respectively, p = 0.005) in mEPSCs in the presence of tetrodotoxin (TTX),
indicating an increase in postsynaptic response of neurons in mGFAP Smo CKO mice (n = 14 and 11 cells
from WT and mGFAP Smo CKO, respectively, 4 animals per genotype; **Figure 5**). These data suggest that
loss of Shh activity in astrocytes produces disturbances in both pre and postsynaptic activity. Moreover,
these data suggest that astrocytes modulate both neuronal excitability and excitatory synaptic
transmission in a Shh-dependent manner.

Selective disruption of Shh signaling in astrocytes produces mild reactive gliosis

We previously demonstrated that cortical astrocytes in mGFAP Smo CKO mice upregulate GFAP
expression and exhibit cellular hypertrophy, two classic hallmarks of reactive astrogliosis [15]. Astrocytes
exhibiting these features were broadly distributed across cortical layers (**Figure 6**), in contrast to the
distribution of Gli1 astrocytes which are found predominantly in layers IV and V. In addition to
upregulation of GFAP, astrocytes in mGFAP Smo CKO mice show dramatic changes in morphological
structure. Sholl analysis revealed no difference in the number of primary branches in mGFAP Smo CKO

mice compared to WT
controls (7.2 and 7.5
branches, WT and mGFAP
Smo CKO, respectively; *n* = 9
cells from 3 animals per
genotype). However,
astrocytes from mGFAP Smo
CKO animals did show an
increase in the number of
higher order branches, the
length of the longest tree
(136.6 ± 17 and 212.1 ± 28
µm for WT and mGFAP Smo
CKO, respectively, *p* = 0.035),
and the total length of all
processes (397 ± 66 and 674
± 91 µm for WT and mGFAP
Smo CKO, respectively, *p* =
0.025) compared to WT
controls (**Figure 6**).
Consistent with this, there

Figure 6: Disruption of Shh signaling in astrocytes results in widespread reactive changes in morphology. (A & B) Brightfield immunohistochemistry reveals a reactive upregulation of GFAP across cortical layers in mGFAP Smo CKO astrocytes. Scale bar, 125 µm. **(C & E)** Representative high-magnification images of both WT (C) and mGFAP Smo CKO (E) astrocytes are shown. Scale bar, 10 µm. **(D & F)** Representative traces of GFAP expression from WT (D) and mGFAP Smo CKO (F) cortical astrocytes. **(G & H)** Sholl analysis shows significant increases in complexity of mGFAP Smo CKO astrocytes compared to WT controls. Statistical significance was assessed by one-way ANOVA with Bonferroni's for multiple comparisons. **(I-K)** Quantification of various morphological features of traced astrocytes. Statistical analysis performed by one-way ANOVA (G & H) and unpaired Student's t-test (I-K). Data points represent individual cells. Graphs represent mean value ± SEM. *n* = 3 animals per genotype.

was a significant increase in the number of branches intersecting
concentric shells at various distances from the soma (**Figure 6**). Notably,
in CK2 Smo CKO mice, GFAP staining in the cortex was indistinguishable
from WT controls (**Figure 6 – figure supplement 1**), suggesting that
astrocytes exhibit dramatic changes in morphological structure following
astrocytic, but not neuronal, disruption of Shh signaling. Such changes in
morphology are consistent with mild reactive gliosis, a complex cellular
response of astrocytes to disturbances in physiological homeostasis [48].
Notably, the distribution of reactive astrocytes in mGFAP Smo CKO mice
is broader than that defined by Gli1 expression, arguing against the idea
that Shh signaling regulates the intrinsic state of astrocytes. Rather,
reactive gliosis may instead reflect a cellular response to the mild, but
persistent, aberrations in neuronal activity observed in mGFAP Smo CKO
mice.
473

Figure 6 – figure supplement 1: Reactive gliosis does not occur when neuronal Shh signaling is disrupted. Brightfield immunohistochemistry for GFAP in *CamKIIa;Smo^fl/fl* animals (B) and littermate controls (A) does not reveal mild reactive gliosis. Scale bar, 250 µm.

DISCUSSION

In this study, we demonstrate that refinement of neural circuits during
postnatal development requires neuron-astrocyte signaling mediated by Shh signaling. Our data show
that Gli1 astrocytes are enriched in deep cortical layers with a relative paucity in upper cortical layers.

Selective disruption of Shh signaling in astrocytes leads to an overabundance of spines on the apical
dendrites of layer V, but not layer II/III, cortical neurons. The overabundance of spines emerges during
postnatal development, continues into adulthood and is accompanied by a reduction in dynamic
turnover. mGFAP Smo CKO mice exhibit a pronounced reduction in Kir4.1 and a concomitant increase in
intrinsic neuronal excitability, consistent with a deficit in astrocyte-mediated K$^+$ buffering. Finally, we
show that cortical astrocytes exhibit dramatic changes in morphology and GFAP expression, a
phenotype consistent with reactive gliosis. Taken together, these data suggest that local neuron-
astrocyte interactions mediated by Shh signaling facilitate modulation of neuronal activity that regulates
synaptic remodeling during postnatal development.

In the postnatal cortex, Shh signaling from layer V neurons mediates synaptic connectivity with their
layer II/III presynaptic partners [14]. Here, we show that, in addition to neuronal Shh signaling, Shh
signaling in astrocytes is required for the establishment and maintenance of cortical circuits. This is
supported by two key observations in our study. First, we find that in mGFAP Smo CKO mice, apical
dendrites of cortical neurons exhibit an increase in spine density that emerges during postnatal
development and persists into adulthood. Importantly, genetic deletion of *Smo* in mature neurons
instead of astrocytes failed to show any differences in spine density, pointing to a specific role for Shh
signaling in astrocyte regulation of synaptic connectivity. Interestingly, deletion of Shh ligand using an
EmxCre driver produces a reduction in spine density of basal dendrites [14], whereas we found no
difference in basal dendrite spine density, suggesting that Shh acts in distinct, cell type-specific ways to
regulate synaptic connectivity of local cortical circuits. Indeed, whereas astrocytes express Gli1, neurons
do not [15], indicating differential transduction of Shh through canonical and non-canonical, Gli-
independent pathways, respectively, in different cell types. Second, our data show fewer filopodia in
juvenile mGFAP Smo CKO mice compared to WT controls along with a concomitant increase in spine
stability. In the adult, spines continue to show deficits in dynamic turnover, demonstrating long-term
impairments in structural plasticity. This suggests that Shh signaling is required not only for identifying
appropriate synaptic partners during development, but it also plays an important role in mediating
synaptic plasticity.

We previously demonstrated that Gli1 expression is restricted to a subpopulation of cortical astrocytes
[15]. Here, we extend these findings and demonstrate that the distribution of Gli1 astrocytes in the cortex
is not uniform across cortical layers, but rather, occurs in a laminar-specific fashion. Cortical layers have
been historically defined by neuronal populations and their specific functional properties and
connectivity. However, emerging evidence suggests that astrocytes also exhibit cortical lamination
patterns based on distinct gene expression profiles [49]. Our data show that Gli1 astrocytes are enriched
in layers IV and V with a relatively sparse distribution in layer II/III. Interestingly, in the visual cortex, the
BMP antagonist, chordin-like 1 (Chrdl1) is preferentially enriched in upper layer cortical astrocytes [8],
suggesting that Gli1 and Chrdl1 may identify two distinct, but regionally complementary, astrocyte
populations. Alternatively, Shh signaling may actively repress Chrdl1 expression in deep layer
astrocytes. Notably, our observation that Gli1 astrocytes only comprise a fraction of astrocytes in layers
IV and V suggests that, in addition to the laminar-specific distribution of these cells, there may be
additional heterogeneity of astrocytes even with a given cortical layer. Whether and how the laminar

distribution of astrocytes with distinct gene expression profiles confers functional specialization is not
well understood and requires further study.

Interestingly, we observed that the disturbances in synaptic organization of cortical neurons in mGFAP
Smo CKO mice occurs selectively in apical dendrites of layer V, but not layer II/III, neurons. Since apical
dendrites of layer V neurons traverse through layer II/III, this suggests that Shh-dependent regulation of
synaptic refinement is not mediated by direct interaction between Gli1 astrocytes and synapses. One
possibility is that Gli1 astrocytes interact with neuronal soma in layer V and effectively modulate their
excitability by providing peri-somatic buffering of extracellular K^+. Indeed, a recent study demonstrated
that Kir4.1 surrounds neuronal soma in the lateral habenula and regulates their firing properties [40].
Astrocytes play essential roles in buffering extracellular K^+ through Kir4.1, and several studies
demonstrate that loss of function of Kir4.1 inhibits K^+ uptake and increases neuronal excitability [43,44,50].
The establishment of appropriate synapse number and connectivity are tightly regulated processes that
are profoundly shaped by experience or activity [33,51]. Our observation that mGFAP Smo CKO mice show
a pronounced reduction in Kir4.1 expression accompanied by increases in spontaneous neuronal activity
and excitability suggests that the activity-dependent processes that govern the refinement of excess
synapses are impaired.

Aberrant synaptic number and organization characterize a number of neurodevelopmental disorders [1].
In addition, remodeling of spines following insult or disease is essential for preserving or restoring
function. Here, we demonstrate that Shh signaling mediates neuron-astrocyte interactions that are
necessary for refining developing neural circuits. The deficits in synaptic organization observed in
mGFAP Smo CKO mice persist for the life of the animal. These deficits are not limited to synapse
number but also include lower rates of spine turnover, suggesting long-lasting deficits in structural
plasticity. Together with the observation that Shh guides local synaptic connectivity of cortical neurons
[14], these studies reveal that Shh signaling exerts considerable influence on the establishment and
maintenance of cortical circuits. Further, these results reveal a specific role for astrocytes in mediating
synaptic connectivity of local cortical circuits. These results point to Shh signaling in astrocytes as a
potential molecular and cellular target for developing novel strategies to overcome aberrant synaptic
connectivity.

METHODS

Animals

All experiments were approved by Drexel University's Institutional Animal Care and Use Committee and
were conducted according to approved protocols. We used the following strains of transgenic mice on
the C57BL/6 background: *Gli1$^{CreER/+}$* [12], Ai14 [52], *Smo$^{fl/fl}$* [53], *mGFAP-Cre* [20], *CaMKIIa-Cre* [54] and *Thy1-GFPm* [24].
Mice of either sex were included in these studies.

Immunohistochemistry

Mice were deeply anesthetized by intraperitoneal injection of ketamine/xylazine/acepromazine.
Animals were then transcardially injected with 100 units heparin and perfused with phosphate-buffered
saline (PBS) followed by 4% paraformaldehyde (PFA, Sigma-Aldrich, St. Louis, MO). Brains were fixed

overnight at 4°C and then transferred to 30% sucrose for at least 48 hours or until sectioned. Brains
were cryosectioned (Leica CM3050S, Wetzlar, Germany) and collected in 40 μm sections. Sections were
stored in 0.1M Tris-buffered saline (TBS) with 0.05% sodium azide at 4°C. Immunohistochemistry was
performed using the following primary antibodies: rabbit anti-GFP (AB3080; Millipore, Burlington, MA),
rabbit anti-Kir4.1 (AB5818; Millipore), rabbit anti-GFAP (Z033429-2; Agilent, Santa Clara, CA), mouse
anti-NeuN (MAB377; Millipore), rabbit anti-RFP (PM005; MBL International, Woburn, MA), rat anti-Ctip2
(ab18465; Abcam, Cambridge, UK), sheep anti-CAII (AHP206; Bio-Rad), rabbit anti-S100β (Z0311,
Agilent), and rabbit anti-Olig2 (AB9610; Millipore). Sections were washed three times in 0.1M TBS at
room temperature (10 minutes/wash). Sections were then blocked in 10% Normal Serum with 0.5%
Triton-X (Sigma-Aldrich) for one hour at room temperature and incubated with primary antibody in 0.5%
Triton-X at 4°C overnight. For fluorescent immunohistochemistry, sections were rinsed the following day
in 0.1M TBS and incubated in Alexafluor-conjugated secondary antibodies with 10% Normal Serum and
0.1M TBS and incubated at room temperature for two hours. The sections were then rinsed in 0.1 TBS
and incubated in DAPI (1:50,000; Life Technologies, Carlsbad, CA) for 15 minutes. Sections were rinsed
again in 0.1M TBS and mounted onto microscope slides (Fisherbrand, Waltham, MA) and coverslipped
using ProLong Gold Antifade Mountant (Invitrogen, Carlsbad, CA) and Fisherfinest Premium Cover Glass.
For brightfield immunohistochemistry, sections were rinsed as previously described but were incubated
with biotinylated secondary antibodies against the primary antibody species (Vector Laboratories,
Burlingame, CA) for one hour. Sections were then placed in Avitin-Biotin Complex solution (Vector) for
one hour and visualized using 3,3'-diaminobenzidine (DAB Peroxidase Substrate Kit, Vector). Sections
were then mounted onto slides and coverslipped with DPX Mountant (Fisher).

Quantification of dendritic spine density

Neurons were traced using Neurolucida 11 (MicroBrightField Biosciences, Williston, VT) and individual
spines were marked. Apical dendrites of layer V neurons were counted from ~100 μm below the primary
bifurcation through the apical tuft. Apical dendrites of layer II/III cells were analyzed just below the
primary bifurcation through the apical tuft. Basal dendrites were analyzed beginning ~50 μm from the
soma through the end of the processes. CA1 hippocampal neurons were analyzed from just below the
primary bifurcation through the apical tuft. All analysis was performed on an upright Zeiss microscope
using a 63x oil objective.

Cranial window

Cranial window surgeries were adapted from a published protocol[34]. Immediately prior to each surgery,
mice received a subcutaneous injection of carprofen (Rimadyl, 5 mg/kg), and a follow-up injection was
given 24 hours after surgery. Mice were deeply anesthetized with isoflurane (5%) in an induction
chamber, and then transferred to a stereotaxic apparatus where they inhaled 1.5% isoflurane
continuously for the duration of the surgery. The scalp was then removed and a metal bar (~1 cm long)
with threadings for screws was affixed to the skull, using superglue and dental acrylic to form a head
post that sealed off the skin while leaving the right parietal bone exposed. After allowing 24 hours for
the glue and acrylic to set, the mouse was returned to the stereotaxic apparatus and head-fixed using
the headpost. A circular craniotomy ~3 mm in diameter and centered 2.5 mm lateral and posterior to

bregma was performed. The exposed dura was treated with saline-soaked gel foam until any minor
bleeding ceased. Finally, a 3 mm glass coverslip was gently pressed onto the dura and sealed in place
using superglue and dental acrylic. Adult mice were allowed at least 1 week for recovery. To mitigate the
tendency of rapid bone growth to destabilize the windows at younger ages, juvenile mice were allowed
24 hours to recover.

Two-photon imaging

A commercial two-photon microscope (Bruker, Billerica, MA) with a tunable Ti:Sapphire laser (Coherent,
Santa Clara, CA) was used for all experiments. The laser was set to 920 nm and power was tuned on the
sample as necessary to obtain consistently high-quality images without damaging the tissue. Mice were
head-fixed and anesthetized with continuous isoflurane (1-1.5%) during imaging. Under a 20X water
immersion objective, GFP-expressing neurons were identified visually and then imaged down to the cell
body to determine layer. From the apical dendrite tuft, individual dendrite segments that projected
primarily along the xy-plane were selected for analysis, and z-stacks (0.5 μm step size) were collected.
To track individual dendrites, repeated imaging was generally performed on the following schedule (in
days, relative to first imaging session): 0, 1, 2, 7, 14, 21, 28, 42. Some data from studies using other
schedules were also included. For juvenile mice, this schedule was: 0, 1, 2.

Analysis of structural plasticity

Initial measures of structural plasticity were determined by side-by-side manual comparison of two z-
stacks across a fixed time-interval (7 days for adults, 2 days for juvenile). These comparisons were
conducted by trained observers blinded to mouse genotype. Long, thin protrusions lacking a bulbous
head were identified as filopodia, and any other prominent dendritic protrusion that extended > 0.4 μm
from the shaft was counted as a spine. The filopodial fraction was calculated as the proportion of all
analyzed features present at the first imaging timepoint that were identified as filopodia. The position
and dynamic status (stable/eliminated/formed) of each spine was recorded using ImageJ's CellCounter
plugin. Each neuron's turnover ratio was calculated as:

$$TO = (N_{elim} + N_{form})/(N_{elim} + N_{form} + 2N_{stable})$$

For longitudinal studies, the dendritic features identified in our initial ImageJ-based analysis were
imported into MATLAB (Mathworks, Natick, MA) and manually tracked across all imaged time-points
using custom-written software. Spines that were observed consistently across all time point were
classified as stable. Spines that disappeared at some time point and were observed again in a
subsequent time point were classified as recurrent. Spines that disappeared and never reappeared were
classified as transient. To compare long-term dynamics between WT and mGFAP Smo CKO mice, we
examined the survival curves, S(t), for each mouse, where:
S(t) = # of spines consistently present to time point t / # of spines initially observed
Survival curves for each mouse were pooled by genotype and fit to a single-phase exponential decay
model:
$$S(t) = S_p + S_i * exp(-t/\tau)$$

where S_p and S_i represent the relative fractions of permanent and impermanent spines respectively and
τ is the characteristic lifespan of impermanent spines. Fitting and statistical comparison of survival data
was performed with PRISM 6 (GraphPad, San Diego, CA) software.

Transcript Quantification

Mice were deeply anesthetized by intraperitoneal injection of ketamine/xylazine/acepromezine before
rapid decapitation. Cortices were dissected into TRIzol reagent (Thermo-Fisher, Waltham, MA) and RNA
was extracted according to the standard protocol. RNA was then purified with an RNeasy Micro Kit
(Qiagen, Hilden, Germany) and subsequently reverse transcribed to cDNA using the High-Capacity cDNA
Reverse Transcription Kit (Thermo-Fisher). Droplet digital PCR (ddPCR) was performed with the QX200
Droplet Digital PCR System using Evagreen Supermix (Bio-Rad, Hercules, CA). Alternatively, for *Megf10*
and *Mertk*, qPCR was performed with the CFX96 Touch Real-Time PCR Detection System (Bio-Rad) using
PowerUp SYBR Green Master Mix (Thermo-Fisher). PrimePCR ddPCR Expression EvaGreen Assay (Bio-
Rad) was used for *Gapdh* primers; all other primers were designed using NCBI Primer-BLAST as follows:
Kir4.1 F, GCTGCCCCGCGATTTATCAG; *Kir4.1* R, AGCGACCGACGTCATCTTGG; *GLAST* F,
TCCTCTACTTCCTGGTAACCC; *GLAST* R, TCCACACCATTGTTCTCTTCC; *Glt1* F, CATCAACAGAGGGTGCCAAC;
Glt1 R, CACACTGCTCCCAGGATGAC; *Megf10* F, CTCACTGCTCTGTCACTGGGTG; *Megf10* R,
GGTAGCTGATTCTGTGCCGTGT; *Mertk* F, AAACTGCATGTTGCGGGATGAC; *Mertk* R,
TCCCACATGGTCACGCCAAA. ddPCR absolute quantification of targets was calculated using Quantasoft
Version 1.7 software (Bio-Rad). Samples were run in triplicate, and *Gapdh* was quantified for each
sample as a loading control with no difference detected between sample groups.

Slice electrophysiology

Mice aged P21 were anesthetized with intraperitoneal injection of euthasol (0.2 ml/kg) and brains were
rapidly removed. 300 μm coronal slices were cut using a vibratome tissue slicer (Leica) and transferred
to a holding chamber, submerged in oxygenated artificial cerebrospinal fluid (ACSF, in mM: 124 NaCl, 2.5
KCl, 1.25 NaH_2PO_4, 2 $CaCl_2$, 1 $MgSO_4$, 26 $NaHCO_3$, and 10 dextrose, pH 7.4) at 36°C for 1 hour and then
maintained at room temperature. Individual slices containing the barrel cortex were placed into a
recording chamber immersed in oxygenated ACSF mounted on an Olympus upright microscope (BX51).
Neurons were visualized with infrared differential interference video microscopy.

Whole-cell current clamp was used to record action potentials from layer V pyramidal cells using patch
electrodes with an open tip resistance of 7–10 MΩ. Patch electrodes were filled with potassium
gluconate internal solution (in mM): 120 potassium gluconate, 20 KCl, 4 ATP-Na, 0.3 Na_2GTP, 5 Na-
phosphocreatine, 0.1 EGTA, 10 HEPES, pH 7.3, 305 mosmol/l) and action potential responses were
measured in response to various step currents from -300 pA to +650 pA with 50 pA increments. Action
potentials were described as spike numbers per depolarized current injections. The resting membrane
potential, input resistance, action potential (AP) threshold, AP half-width, and peak AP amplitude were
also measured.

Whole-cell voltage-clamp recordings were obtained from layer V pyramidal cells using patch electrodes
with Cs^+-containing solution (in mM): 120 Cs-gluconate, 5 lidocaine, 6 $CaCl_2$, 1 Na_2ATP, 0.3 Na_2GTP, and

10 HEPES (pH 7.3 adjusted by CsOH). To record spontaneous excitatory postsynaptic currents (sEPSCs),

cells were held at a membrane potential of -70mV in the presence of $GABA_A$ receptor antagonist

picrotoxin (PTX; 100 µM, Sigma-Aldrich) and recorded for 5 minutes. Miniature EPSCs (mEPSCs) were

then recorded for an additional 5 minutes in the presence of both picrotoxin and tetrodotoxin (TTX, 0.5

µM, Hello Bio, Princeton, NJ). All recordings were conducted with Axon MultiClamp 700B amplifier

(Molecular Devices, San Jose, CA). The sEPSCs and mEPSCs frequency and amplitudes were measured by

averaging 5 sweeps from the onset of recording with Clampfit 9.2 software (Molecular Devices). The

s/mEPSCs were detected and characterized using a sample template for the 5-minute data period. Based

upon the selected sample template, the frequency (number of event detections) and amplitude of

events were measured using a threshold set in Clampfit.

Statistical Analyses

Statistical analyses used for various datasets are indicated both in the figure legends and in the text.

Prism 6 software (GraphPad) was used for all analyses and to generate graphs.

REFERENCES

1. Penzes, P., Cahill, M. E., Jones, K. A., VanLeeuwen, J.-E. & Woolfrey, K. M. Dendritic spine pathology in neuropsychiatric disorders. *Nat. Neurosci.* **14**, 285–293 (2011). DOI: 10.1038/nn.2741

2. Allen, N. J. & Eroglu, C. Cell Biology of Astrocyte-Synapse Interactions. *Neuron* **96**, 697–708 (2017). DOI: 10.1016/j.neuron.2017.09.056

3. Adamsky, A., Kol, A., Groysman, M., London, M., Kreisel, T., Doron, A., Ozeri-Engelhard, N., Melcer, T., Refaeli, R., Horn, H., Regev, L. & Goshen, I. Astrocytic Activation Generates De Novo Neuronal Potentiation and Memory Enhancement. *Cell* **174**, 59–71 (2018). DOI: 10.1016/j.cell.2018.05.002

4. Risher, W. C., Patel, S., Kim, I. H. wan, Uezu, A., Bhagat, S., Wilton, D. K., Pilaz, L. J., Singh Alvarado, J., Calhan, O. Y., Silver, D. L., Stevens, B., Calakos, N., Soderling, S. H. & Eroglu, C. Astrocytes refine cortical connectivity at dendritic spines. *Elife* **3**, 166–175 (2014). DOI: 10.7554/eLife.04047

5. Allen, N. J., Bennett, M. L., Foo, L. C., Wang, G. X., Chakraborty, C., Smith, S. J. & Barres, B. A. Astrocyte glypicans 4 and 6 promote formation of excitatory synapses via GluA1 AMPA receptors. *Nature* **486**, 410–414 (2012). DOI: 10.1038/nature11059

6. Kucukdereli, H., Allen, N. J., Lee, A. T., Feng, A., Ozlu, M. I., Conatser, L. M., Chakraborty, C., Workman, G., Weaver, M., Sage, E. H., Barres, B. A. & Eroglu, C. Control of excitatory CNS synaptogenesis by astrocyte-secreted proteins Hevin and SPARC. *Proc. Natl. Acad. Sci. U. S. A.* **108**, E440-9 (2011). DOI: 10.1073/pnas.1104977108

7. Christopherson, K. S., Ullian, E. M., Stokes, C., Mullowney, C., Hell, J., Agah, A., Lawler, J., Mosher, D., Bornstein, P. & Barres, B. A. Thrombospondins are astrocyte-secreted proteins that promote CNS synaptogenesis. *Cell* **120**, 421–433 (2005).

8. Blanco-Suarez, E., Liu, T.-F., Kopelevich, A. & Allen, N. J. Astrocyte-Secreted Chordin-like 1 Drives Synapse Maturation and Limits Plasticity by Increasing Synaptic GluA2 AMPA Receptors. *Neuron* 1–17 (2018). DOI: 10.1016/j.neuron.2018.09.043

9. Fuccillo, M., Joyner, A. L. & Fishell, G. Morphogen to mitogen: the multiple roles of hedgehog signalling in vertebrate neural development. *Nat. Rev. Neurosci.* **7**, 772–83 (2006). DOI: 10.1038/nrn1990

10. Ruiz i Altaba, A., Palma, V. & Dahmane, N. Hedgehog–Gli Signalling and the Growth of the Brain. *Nat. Rev. Neurosci.* **3**, 24–33 (2002). DOI: 10.1038/nrn704

11. Traiffort, E., Angot, E. & Ruat, M. Sonic Hedgehog signaling in the mammalian brain. *Journal of Neurochemistry* **113**, 576–590 (2010). DOI: 10.1111/j.1471-4159.2010.06642.x

12. Ahn, S. & Joyner, A. L. In vivo analysis of quiescent adult neural stem cells responding to Sonic hedgehog. *Nature* **437**, 894–897 (2005). DOI: 10.1038/nature03994

13. Ihrie, R. A., Shah, J. K., Harwell, C. C., Levine, J. H., Guinto, C. D., Lezameta, M., Kriegstein, A. R. & Alvarez-Buylla, A. Persistent Sonic Hedgehog Signaling in Adult Brain Determines Neural Stem Cell Positional Identity. *Neuron* **71**, 250–262 (2011). DOI: 10.1016/j.neuron.2011.05.018

14. Harwell, C. C., Parker, P. R. L., Gee, S. M., Okada, A., McConnell, S. K., Kreitzer, A. C. & Kriegstein, A. R. Sonic Hedgehog Expression in Corticofugal Projection Neurons Directs Cortical Microcircuit

728   Formation. *Neuron* **73**, 1116–1126 (2012). DOI: 10.1016/j.neuron.2012.02.009

729 15. Garcia, A. D. R., Petrova, R., Eng, L. & Joyner, A. L. Sonic hedgehog regulates discrete populations
730   of astrocytes in the adult mouse forebrain. *J. Neurosci.* **30**, 13597–608 (2010). DOI:
731   10.1523/JNEUROSCI.0830-10.2010

732 16. Garcia, A. D. R., Han, Y.-G., Triplett, J. W., Farmer, W. T., Harwell, C. C. & Ihrie, R. A. The Elegance
733   of Sonic Hedgehog: Emerging Novel Functions for a Classic Morphogen. *J. Neurosci.* **38**, 9338–
734   9345 (2018). DOI: 10.1523/JNEUROSCI.1662-18.2018

735 17. Alvarez, J. I., Dodelet-Devillers, A., Kebir, H., Ifergan, I., Fabre, P. J., Terouz, S., Sabbagh, M.,
736   Wosik, K., Bourbonnière, L., Bernard, M., Van Horssen, J., De Vries, H. E., Charron, F. & Prat, A.
737   The hedgehog pathway promotes blood-brain barrier integrity and CNS immune quiescence.
738   *Science (80-.).* **334**, 1727–1731 (2011). DOI: 10.1126/science.1206936

739 18. Allahyari, R. V., Clark, K. L., Shepard, K. A. & Garcia, A. D. R. Sonic hedgehog signaling is negatively
740   regulated in reactive astrocytes after forebrain stab injury. *Sci. Rep.* **9**, 565 (2019). DOI:
741   10.1038/s41598-018-37555-x

742 19. Farmer, W. T., Abrahamsson, T., Chierzi, S., Lui, C., Zaelzer, C., Jones, E. V, Bally, B. P., Chen, G. G.,
743   Théroux, J.-F., Peng, J., Bourque, C. W., Charron, F., Ernst, C., Sjöström, P. J. & Murai, K. K.
744   Neurons diversify astrocytes in the adult brain through sonic hedgehog signaling. *Science (80-.).*
745   **351**, 849–854 (2016). DOI: 10.1126/science.aab3103

746 20. Garcia, A. D. R., Doan, N. B., Imura, T., Bush, T. G. & Sofroniew, M. V. GFAP-expressing
747   progenitors are the principal source of constitutive neurogenesis in adult mouse forebrain. *Nat.*
748   *Neurosci.* **7**, 1233–1241 (2004). DOI: 10.1038/nn1340

749 21. Petrova, R., Garcia, A. D. R. & Joyner, A. L. Titration of GLI3 Repressor Activity by Sonic Hedgehog
750   Signaling Is Critical for Maintaining Multiple Adult Neural Stem Cell and Astrocyte Functions. *J.*
751   *Neurosci.* **33**, 17490–17505 (2013). DOI: 10.1523/JNEUROSCI.2042-13.2013

752 22. Zhuo, L., Theis, M., Alvarez-Maya, I., Brenner, M., Willecke, K. & Messing, A. hGFAP-cre
753   Transgenic Mice for Manipulation of Glial and Neuronal Function In Vivo With the goal of
754   performing astrocyte-specific modifica-tion of genes in the mouse. (2001). DOI:
755   10.1002/gene.10008

756 23. Nimchinsky, E. A., Sabatini, B. L. & Svoboda, K. Structure and Function of Dendritic Spines. *Annu.*
757   *Rev. Physiol.* **64**, 313–353 (2002). DOI: 10.1146/annurev.physiol.64.081501.160008

758 24. Feng, G., Mellor, R. H., Bernstein, M., Keller-Peck, C., Nguyen, Q. T., Wallace, M., Nerbonne, J. M.,
759   Lichtman, J. W. & Sanes, J. R. Imaging Neuronal Subsets in Transgenic Mice Expressing Multiple
760   Spectral Variants of GFP. *Neuron* **28**, 41–51 (2000). DOI: 10.1016/S0896-6273(00)00084-2

761 25. Zuo, Y., Lin, A., Chang, P. & Gan, W.-B. Development of Long-Term Dendritic Spine Stability in
762   Diverse Regions of Cerebral Cortex. *Neuron* **46**, 181–189 (2005). DOI:
763   10.1016/j.neuron.2005.04.001

764 26. Holtmaat, A., Trachtenberg, J. T., Wilbrecht, L., Shepherd, G. M., Zhang, X., Knott, G. W. &
765   Svoboda, K. Transient and persistent dendritic spines in the neocortex in vivo. *Neuron* **45**, 279–
766   291 (2005). DOI: 10.1016/j.neuron.2005.01.003

767 27. Han, Y. G., Spassky, N., Romaguera-Ros, M., Garcia-Verdugo, J. M., Aguilar, A., Schneider-
768   Maunoury, S. & Alvarez-Buylla, A. Hedgehog signaling and primary cilia are required for the
769   formation of adult neural stem cells. *Nat. Neurosci.* **11**, 277–284 (2008). DOI: 10.1038/nn2059

28.   Attardo, A., Fitzgerald, J. E. & Schnitzer, M. J. Impermanence of dendritic spines in live adult CA1
hippocampus. *Nature* **523**, 592–596 (2015). DOI: 10.1038/nature14467

29.   Perez-Cruz, C., Nolte, M. W., Van Gaalen, M. M., Rustay, N. R., Termont, A., Tanghe, A., Kirchhoff,
F. & Ebert, U. Reduced Spine Density in Specific Regions of CA1 Pyramidal Neurons in Two
Transgenic Mouse Models of Alzheimer's Disease. *J. Neurosci.* **31**, 3926–3934 (2011). DOI:
10.1523/JNEUROSCI.6142-10.2011

30.   Lendvai, B., Stern, E. a, Chen, B. & Svoboda, K. Experience-dependent plasticity of dendritic
spines in the developing rat barrel cortex in vivo. *Nature* **404**, 876–881 (2000). DOI:
10.1038/35009107

31.   Trachtenberg, J. T., Chen, B. E., Knott, G. W., Feng, G., Sanes, J. R., Welker, E. & Svoboda, K. Long-
term in vivo imaging of experience-dependent synaptic plasticity in adult cortex. *Nature* **420**,
788–794 (2002). DOI: 10.1038/nature01273

32.   Holtmaat, A. & Svoboda, K. Experience-dependent structural synaptic plasticity in the
mammalian brain. *Nat. Rev. Neurosci.* **10**, 647–658 (2009). DOI: 10.1038/nrn2721

33.   Zuo, Y., Yang, G., Kwon, E. & Gan, W.-B. Long-term sensory deprivation prevents dendritic spine
loss in primary somatosensory cortex. *Nature* **436**, 261–265 (2005). DOI: 10.1038/nature03715

34.   Holtmaat, A., Bonhoeffer, T., Chow, D. K., Chuckowree, J., De Paola, V., B, H. S., Hübener, M.,
Keck, T., Knott, G., Lee, W.-C. A., Mostany, R., Mrsic-Flogel, T. D., Nedivi, E., Portera-Cailliau, C.,
Svoboda, K., Trachtenberg, J. T. & Wilbrecht, L. Long-term, high-resolution imaging in the mouse
neocortex through a chronic cranial window. *Nat. Protoc.* **4**, 1128–44 (2009). DOI:
10.1038/nprot.2009.89

35.   Ziv, N. E. & Smith, S. J. Evidence for a role of dendritic filopodia in synaptogenesis and spine
formation. *Neuron* **17**, 91–102 (1996). DOI: 10.1016/S0896-6273(00)80283-4

36.   Berry, K. P. & Nedivi, E. Spine Dynamics: Are They All the Same? *Neuron* **96**, 43–55 (2017). DOI:
10.1016/j.neuron.2017.08.008

37.   Grutzendler, J., Kasthuri, N. & Gan, W. B. W.-B. Long-term dendritic spine stability in the adult
cortex. *Nature* **420**, 812–816 (2002). DOI: 10.1038/nature01151.1.

38.   Chung, W.-S., Clarke, L. E., Wang, G. X., Stafford, B. K., Sher, A., Chakraborty, C., Joung, J., Foo, L.
C., Thompson, A., Chen, C., Smith, S. J. & Barres, B. a. Astrocytes mediate synapse elimination
through MEGF10 and MERTK pathways. *Nature* **504**, 394–400 (2013). DOI: 10.1038/nature12776

39.   Kelley, K. W., Ben Haim, L., Schirmer, L., Tyzack, G. E., Tolman, M., Miller, J. G., Tsai, H. H., Chang,
S. M., Molofsky, A. V., Yang, Y., Patani, R., Lakatos, A., Ullian, E. M. & Rowitch, D. H. Kir4.1-
Dependent Astrocyte-Fast Motor Neuron Interactions Are Required for Peak Strength. *Neuron* 1–
14 (2018). DOI: 10.1016/j.neuron.2018.03.010

40.   Cui, Y., Yang, Y., Ni, Z., Dong, Y., Cai, G., Foncelle, A., Ma, S., Sang, K., Tang, S., Li, Y., Shen, Y.,
Berry, H., Wu, S. & Hu, H. Astroglial Kir4.1 in the lateral habenula drives neuronal bursts in
depression. *Nature* **554**, 323–327 (2018). DOI: 10.1038/nature25752

41.   Higashi, K., Fujita, A., Inanobe, A., Tanemoto, M., Doi, K., Kubo, T. & Kurachi, Y. An inwardly
rectifying K(+) channel, Kir4.1, expressed in astrocytes surrounds synapses and blood vessels in
brain. *Am J Physiol Cell Physiol* **281**, C922-31 (2001). DOI: 10.1002/cne.903280405

42.   Kucheryavykh, Y. V., Kucheryavykh, L. Y., Nichols, C. G., Maldonado, H. M., Baksi, K., Reichenbach,
A., Skatchkov, S. N. & Eaton, M. J. Downregulation of Kir4.1 inward rectifying potassium channel

subunits by RNAi impairs potassium transfer and glutamate uptake by cultured cortical
astrocytes. *Glia* **55**, 274–281 (2007). DOI: 10.1002/glia.20455

43. Djukic, B., Casper, K. B., Philpot, B. D., Chin, L. S. & McCarthy, K. D. Conditional knock-out of
Kir4.1 leads to glial membrane depolarization, inhibition of potassium and glutamate uptake, and
enhanced short-term synaptic potentiation. *J. Neurosci.* **27**, 11354–11365 (2007). DOI:
10.1523/jneurosci.0723-07.2007

44. Tong, X., Ao, Y., Faas, G. C., Nwaobi, S. E., Xu, J., Haustein, M. D., Anderson, M. A., Mody, I.,
Olsen, M. L., Sofroniew, M. V. & Khakh, B. S. Astrocyte Kir4.1 ion channel deficits contribute to
neuronal dysfunction in Huntington's disease model mice. *Nat. Neurosci.* **17**, 694–703 (2014).
DOI: 10.1038/nn.3691

45. Nwaobi, S. E., Cuddapah, V. A., Patterson, K. C., Randolph, A. C. & Olsen, M. L. The role of glial-
specific Kir4.1 in normal and pathological states of the CNS. *Acta Neuropathol.* **132**, 1–21 (2016).
DOI: 10.1007/s00401-016-1553-1

46. Olsen, M. L. & Sontheimer, H. Functional implications for Kir4.1 channels in glial biology: From K +
buffering to cell differentiation. *Journal of Neurochemistry* **107**, 589–601 (2008). DOI:
10.1111/j.1471-4159.2008.05615.x

47. Chever, O., Djukic, B., McCarthy, K. D. & Amzica, F. Implication of Kir4.1 Channel in Excess
Potassium Clearance: An In Vivo Study on Anesthetized Glial-Conditional Kir4.1 Knock-Out Mice.
J. Neurosci. **30**, 15769–15777 (2010). DOI: 10.1523/JNEUROSCI.2078-10.2010

48. Sofroniew, M. V. Astrocyte barriers to neurotoxic inflammation. *Nat. Rev. Neurosci.* **16**, 249–263
(2015). DOI: 10.1038/nrn3898

49. Bayraktar, O. A. et al. Single-cell in situ transcriptomic map of astrocyte cortical layer diversity.
bioRxiv 432104 (2018). DOI: 10.1101/432104

50. Sibille, J., Pannasch, U. & Rouach, N. Astroglial potassium clearance contributes to short-term
plasticity of synaptically evoked currents at the tripartite synapse. *J. Physiol.* **592**, 87–102 (2014).
DOI: 10.1113/jphysiol.2013.261735

51. Yang, G., Pan, F. & Gan, W.-B. Stably maintained dendritic spines are associated with lifelong
memories. *Nature* **462**, 920–924 (2009). DOI: 10.1038/nature08577

52. Madisen, L., Zwingman, T. A., Sunkin, S. M., Oh, S. W., Zariwala, H. A., Gu, H., Ng, L. L., Palmiter,
R. D., Hawrylycz, M. J., Jones, A. R., Lein, E. S. & Zeng, H. A robust and high-throughput Cre
reporting and characterization system for the whole mouse brain. *Nat. Neurosci.* **13**, 133–40
(2010). DOI: 10.1038/nn.2467

53. Long, F., Zhang, X. M., Karp, S., Yang, Y. & McMahon, A. P. Genetic manipulation of hedgehog
signaling in the endochondral skeleton reveals a direct role in the regulation of chondrocyte
proliferation. *Development* **128**, 5099–108 (2001). DOI: 10.1016/s0092-8674(00)80094-x

54. Tsien, J. Z., Chen, D. F., Gerber, D., Tom, C., Mercer, E. H., Anderson, D. J., Mayford, M., Kandel, E.
R. & Tonegawa, S. Subregion- and cell type-restricted gene knockout in mouse brain. *Cell* **87**,
1317–1326 (1996). DOI: 10.1016/S0092-8674(00)81826-7