Evidence accumulation during perceptual decision-making is sensitive to the dynamics of

attentional selection

Running title: Decision making depends on attentional selection

Authors: Dragan Rangelov[a] and Jason B. Mattingley[a,b]

Affiliations:

[a]Queensland Brain Institute, The University of Queensland, St Lucia, QLD 4072, Australia

[b]School of Psychology, The University of Queensland, St Lucia, QLD 4072, Australia

Corresponding Author: Dragan Rangelov, d.rangelov@uq.edu.au, +61 401 281 181

Conflict of interest: The authors declare no competing financial interests.

11   **Abstract**

The ability to select and combine multiple sensory inputs in support of accurate

decisions is a hallmark of adaptive behaviour. Attentional selection is often needed to

prioritize stimuli that are task-relevant and to attenuate potentially distracting sources of

sensory information. As most studies of perceptual decision-making to date have made use

of task-relevant stimuli only, relatively little is known about how attention modulates

decision making. To address this issue, we developed a novel 'integrated' decision-making

task, in which participants judged the average direction of successive target motion signals

while ignoring concurrent and spatially overlapping distractor motion signals. In two

experiments that varied the role of attentional selection, we used linear regression to

quantify the influence of target and distractor stimuli on behaviour. Using

electroencephalography, we characterised the neural correlates of decision making,

attentional selection and feature-specific responses to target and distractor signals. While

targets strongly influenced perceptual decisions and associated neural activity, we also

found that concurrent and spatially coincident distractors exerted a measurable bias on

both behaviour and brain activity. Our findings suggest that attention operates as a real-

time but imperfect filter during perceptual decision-making by dynamically modulating the

contributions of task-relevant and irrelevant sensory inputs.

Keywords: decision making, visual attention, encephalography, mixture distribution

modelling, frequency tagging, multivariate analyses

Introduction

Cognition can be conceived of as a cascade of processes, beginning with the

encoding of sensory input and ending with appropriate behavioural choices. Sequential-

sampling decision-making models have had considerable success in describing choice

behaviour in terms of both behavioural performance and neural activity (Forstmann,

Ratcliff, & Wagenmakers, 2016; O'Connell, Shadlen, Wong-Lin, & Kelly, 2018; Ratcliff, Smith,

Brown, & McKoon, 2016). According to these models, decision-making relies on a central

decision variable, an abstract quantity that accumulates in time toward one of several

possible choice alternatives. On these accounts, a choice is made once the decision variable

reaches threshold. In one widely employed decision-making task – the random-dot motion

discrimination task (Ratcliff et al., 2016; Shadlen & Kiani, 2013) – participants are shown

fields of randomly moving dots and are asked to judge the motion direction of a fraction of

the dots that move coherently. Studies in both animals (Shadlen & Kiani, 2013) and humans

(Heekeren, Marrett, & Ungerleider, 2008; Kelly & O'Connell, 2013; Liu & Pleskac, 2011;

O'Connell, Dockree, & Kelly, 2012; Ratcliff, Philiastides, & Sajda, 2009; Twomey, Kelly, &

O'Connell, 2016) have shown that neural activity recorded during random-dot motion

discrimination follows a trajectory that resembles the accumulation of a decision variable,

thereby lending neurobiological support to sequential-sampling decision-making models.

To date, much has been learned about simple perceptual decision-making in humans

and other animal species (Hanks & Summerfield, 2017; O'Connell et al., 2018; Summerfield

& Tsetsos, 2015). In the real world, however, decisions are often more complex, requiring

judgements on stimuli that are separated in time and space, and which may or may not be

task relevant. For example, deciding to cross a busy road requires us to integrate the motion

directions of vehicles coming from both the left and right sides, while disregarding any

movements made by people or objects on the sidewalk. In recent years there has been

increased interest in studying behaviour and neural activity in more complex decision-

making tasks (Churchland & Ditterich, 2012; Churchland, Kiani, & Shadlen, 2008;

Itthipuripat, Cha, Deering, Salazar, & Serences, 2018; Kohl, Spieser, Forster, Bestmann, &

Yarrow, 2018; Leite & Ratcliff, 2010; Spitzer, Blankenburg, & Summerfield, 2016; Spitzer,

Waschke, & Summerfield, 2017; Wyart, Myers, & Summerfield, 2015). To increase decision

complexity, many of these studies have varied the number of response alternatives (e.g.,

two vs. four), and have found significant costs associated with a larger number of potential

responses (e.g., Churchland et al., 2008; Leite and Ratcliff, 2010; Itthipuripat et al., 2018;

Kohl et al., 2018). To date, however, the question about how the brain regulates decision-

making when it receives competing – task-relevant and irrelevant – sensory inputs has

received little attention in the literature. The central aim of the present study was to

characterise the effects of selective attention on perceptual decision-making. Specifically,

we were interested in measuring the degree to which relevant and irrelevant signals are

integrated into a decision variable that is used as the basis for behavioural choices.

According to sequential-sampling decision-making models, selective attention

should mediate sensory input and evidence accumulation by filtering out task-irrelevant

inputs and preventing them from being accumulated into the decision variable (Leite &

Ratcliff, 2010; Smith & Ratcliff, 2009; White, Ratcliff, & Starns, 2011; Wyart et al., 2015). This

framework raises at least two inter-related questions. First, what is the relationship

between the time-course of attentional selection and evidence accumulation, and second,

how effective is attention in mediating sensory encoding and evidence accumulation?

Conceptually, selection and accumulation could take place sequentially so that only

prioritised, task-relevant input is accumulated. Alternatively, the two processes could take

place in parallel, so that task-irrelevant inputs might also impact upon evidence

accumulation to some extent, particularly in the early stages of attentional selection.

Studies on visual attention have repeatedly shown robust behavioural and neural

costs when task-irrelevant, distracting stimuli are presented together with task-relevant

target stimuli (e.g., S. K. Andersen & Müller, 2010; Eriksen, 1995; Theeuwes, 1992). While

the literature agrees that attentional selectivity is imperfect, relatively little is known about

the effects of attention on the dynamics of evidence accumulation. When both target and

distractor signals are present, attention might influence decision making in at least two

ways. On the one hand, target and distractor stimuli could each trigger a separate decision-

making process, in which case attention would modulate the probability with which the two

processes drive behaviour. On the other hand, if attention operates as a filter between

sensory input and evidence accumulation, it would be expected to modulate the degree to

which target and distractor signals are accumulated into a single decision variable. The

critical distinction between these two accounts relates to the content of decision variable. In

the former case, the decision variable could reflect either the target or distractor signal,

whereas in the latter it would be a mixture of *both* target and distractor signals.

In the present study, across two experiments, we used random-dot displays

comprising **two** simultaneous, intermingled fields of dots, one in a cued (target) colour, the

other in an uncued (distractor) colour (see Fig. 1). Past research has suggested that the

number of targets may influence the effectiveness of attentional selection (Frătescu, Van

Moorselaar, & Mathôt, 2019; Hollingworth & Beck, 2016; van Moorselaar, Theeuwes, &

Olivers, 2014). Motivated by these findings, we wanted test whether and to which extent

the complexity of attentional set influences evidence accumulation. To that purpose, in

different blocks of trials, we cued our participants to monitor for either a single target

colour (e.g., blue) or two target colours (e.g., blue and yellow). In each trial, participants

were presented with successive epochs of dot motion (each containing the two overlapping

dot fields, one in the target colour and the other in a distractor colour). At the end of the

trial, participants' were instructed to reproduce the ***average target motion direction***,

ignoring the directions of the concurrent distractor-dot fields. Thus, for example, if target

motion direction was 90° for the first display and 110° for the second, the *average* motion

direction would be 100°. Participants made their responses on a continuous scale by

rotating a dial to match their judgement of the average motion direction of the two targets.

This approach enabled us to characterise the precision with which participants integrated

the two target signals and to separate noisy target responses from random guesses using

mixture distribution modelling (Schneegans & Bays, 2016). Finally, a continuous nature of

our response variable enabled us to use linear regression to quantify the degree to which

presented target and distractor motion signals influenced the responses.

In addition to these behavioural measures, we recorded participants' neural activity

using electroencephalography (EEG). We focused in particular on the temporal dynamics of

decision processes as reflected in the central-parietal positivity (CPP). Previous research has

shown that only task-relevant stimuli evoke the CPP (Loughnane et al., 2016) and that the

CPP is observed even when no response is required (Twomey et al., 2016), suggesting that

this component is neither a passive sensory response nor a correlate of motor planning.

Further, the CPP time-course closely resembles the time-course of evidence accumulation:

specifically, its amplitude builds up gradually (O'Connell et al., 2012) and the build-up slope

is proportional to stimulus quality (Kelly & O'Connell, 2013) suggesting that the CPP is

intimately related to decision making.

127   **Experiment 1**

Materials and Methods

Participants. 36 healthy volunteers (19 females, average age 21.2 ± 2.5 years)

participated in Experiment 1. All had normal or corrected-to-normal visual acuity and

normal colour vision. Three participants were identified as outliers based on their

behavioural performance and/or EEG signals (see below for details), leaving 33 participants

in the final sample. The study was approved by the Human Research Ethics Committee of

The University of Queensland (Approval 2016001247), and was conducted in accordance

with the Human Subjects Guidelines of the Declaration of Helsinki. Participants provided

written informed consent prior to taking part in the study.

Stimuli, task and procedure. At the beginning of each trial of Experiment 1, a central

disk, shown for 1 s, cued participants to the target colour (Fig. 1a). In separate blocks of

trials, participants were cued to look for either one target colour (e.g., blue), or two

different target colours (e.g., blue and yellow). The cue was followed by a .5 s buffer period

showing a field of 80 grey, randomly-moving dots. Then, over a 1.5 s period, the dots' colour

saturation gradually increased before returning to grey. The increase in saturation revealed

two overlapping, coloured-dot patches moving in different directions. Each trial comprised

three epochs of colour modulation – in two of these epochs, one dot patch was rendered in

the target colour, and the other appeared in a distractor colour. In the remaining ("no

target") epoch, neither of the patches had a target colour. Target patches were displayed

equally often in the first, second and third epochs of a trial. Participants were asked to

monitor successively presented epochs and to indicate the ***average direction*** of target-

motion events at the end of the trial using a response dial. A response feedback symbol

showing the correct answer was presented after every trial. Participants were instructed to

prioritise response accuracy over response speed, and the response times were not

analysed further. Participants completed 30 practice trials followed by six blocks of 44 trials

(264 trials in total) which took approximately one hour.

In Experiment 1, dot-motion was generated at a suprathreshold level (40%

coherence), with a speed of 19°/s, 100 ms dot-life, and with different signal-dots across

frames and random-walk noise-dots (Scase, Braddick, & Raymond, 1996). Although

coherent motion was present throughout the epoch, different motion directions were

discernible only at higher colour saturations. The motion directions were uniformly sampled

from a 0°–359° range in 1° steps, with the target-distractor and target-target difference

remaining within ±30°–150°. Five easily distinguishable colours were selected in the HSL

colour-space with a minimum hue difference of 72° and medium saturation and brightness

(both 50). Two randomly-selected colours served as targets and two as distractors. In two

target-colour trials, there was an equal number of trials in which the target colour repeated

or changed between two target-present epochs within the trial.

Apparatus. The experiments were conducted in a dark, acoustically and

electromagnetically shielded room. The stimuli were presented on a 24" monitor with

1920×1080 resolution and a refresh rate of 144 Hz. The experimental software was custom-

coded in Python using the PsychoPy toolbox (Peirce, 2007, 2009). EEG signals were recorded

using 64 Ag-AgCl electrodes (BioSemi ActiveTwo) arranged in the 10-20 layout, and sampled

at 1,024 Hz.

Behavioural analyses. To identify outlier participants, the distributions of error

magnitudes (i.e., the angular difference between the response and the correct answer)

were compared to a uniform distribution (i.e., pure guessing) using the Kolmogorov-Smirnov

test. Participants for whom the probability of the null hypothesis (i.e., a uniform distribution

of error magnitudes) exceeded .001 were removed from further analyses.

The remaining distributions per experimental condition and per participant were

fitted to a theoretical model (Schneegans & Bays, 2016), and responses were separated into

noisy target responses and random guesses. We estimated two parameters per participant

and experimental condition: (i) the response precision of the target responses (K parameter)

and (ii) the guessing rates (α).

Finally, we quantified the degree to which presented motion signals influenced the

response. To that purpose, a multiple linear regression (OLS) model with a term for each of

the presented motion directions, expressed as complex numbers, was fitted to the

responses, separately per participant and experimental condition. Transforming the angles

into complex numbers was necessary, as both our independent and the dependent

variables were circular variables. We could not use real values, because real-valued linear

regression would treat the 0° and 360° angles as different even though the two are in fact

alternative representations of the same angle. The absolute value of the resulting regression

coefficients reflects the influence of each of the presented coherent motion signals on the

response, i.e., its *decision weight*. As the decision weights were always positive, we used

permutation testing (10,000 permutations) to generate the empirical null distribution

against which the empirical decision weights were tested.

EEG analyses. EEG signals were analysed using the MNE-Python toolbox (Gramfort et

al., 2013). The data were re-referenced offline to the average electrode, low-pass filtered at

99 Hz and notch-filtered at 50 Hz to eliminate line noise. The recorded signal was pre-

processed using the FASTER algorithm for automated artefact rejection (Nolan, Whelan, &

Reilly, 2010). The pre-processed signal was down-sampled to 256 Hz, segmented into

epochs between the cue-onset and the response-display onset, baseline-corrected relative

to -.1–0 s pre-trial and linearly de-trended. Outlier epochs and participants were identified

using the FASTER algorithm and removed from further analyses.

The CPP was computed by averaging Cz/CPz electrodes across trials. Each epoch

within a trial was extracted from the onset of the increase in colour saturation and baseline-

corrected relative to a common baseline during -.1–0 s, averaged across all epochs. We

quantified the CPP onset latency, slope and peak amplitude. The peak CPP amplitudes were

identified using the jackknife-method (Ulrich & Miller, 2001). Using the same method, the

CPP onset latencies were computed at 50% peak amplitude. The CPP slopes were computed

as the amplitude difference between -.05 s and .05 s relative to the ERP onset, and divided

by .1 s so as to yield μV/s.

To characterise feature-specific neural responses to coherent motion, we used a

multivariate analysis approach described previously (Ede, Chekroud, Stokes, & Nobre, 2018,

2019; Wolff, Jochim, Akyürek, & Stokes, 2017). This analysis splits the whole data set into

training and test epochs and uses the results derived from the training set to characterise

the brain's response in the test epochs. We used an iterative, leave-one-out algorithm in

which each epoch of coherent motion served as a test epoch and the remaining epochs

served as the training epochs. Two separate analyses were conducted, one for the target

motion direction, the other for the distractor motion direction. The training epochs were

first sorted into 16 equidistant bins (β centres range = -180–180°, width = 90°) based on the

angular distance between motion directions in the test and the training epochs. The epochs

within a bin were then averaged, yielding a bin-specific ERP. For each time sample, a

multidimensional Mahalanobis distance between the test and each bin ERP was computed

using temporally smoothed (Gaussian window, SD = 16 ms) and spatially filtered (surface-

Laplacian) signals from all 64 electrodes. This analysis yielded a distance matrix

($D_{M\ bins \times N\ samples}$) containing the distance (d_{mn}) between a test epoch and a given bin m

for time sample n. The distances were then normalised by subtracting the mean bin

distance per time sample ($\hat{d}_{mn} = d_{mn} - \frac{1}{M\ bins}\sum_{i=1}^{M\ bins} d_{in}$), temporally smoothed using a

Gaussian window (SD = 16 ms), and expressed as a similarity matrix by inverting the sign

($S_{M \times N} = -\hat{D}_{M \times N}$).

If the test epoch contains feature-specific signals, the similarity matrix should

contain the largest values for bins with the smallest angular difference relative to the test

direction (e.g., ±15°), and the smallest values for bins with the largest difference (e.g.,

±180°). In a final step, we computed an aggregate measure of motion-specific response (i.e.,

motion tuning) by convolving the similarity matrix with a vector of cosine-transformed bin

centres ($Tuning_N = \cos(\beta_M) \cdot S_{M \times N}$). This non-parametric measure provides a robust

index of feature-specific brain responses. It returns a value of zero when the similarity is the

same across all bins and increases gradually with sharpening of the similarity profile across

bins.

Results

To quantify how well participants could perform the task, we used mixture

distribution modelling to estimate the response precision (K) and guessing rates (α) per

participant and experimental condition. Initial analyses showed no differences between

repetitions and changes of target colour within a trial (all ts < 1), so these two trial types

were averaged for further analyses. The average response precision was high both in the

one- and two-target colour conditions ($K_{M/SEM}$ = 10.52/1.70 and 11.79/2.08), and did not

differ significantly between conditions (F < 1). Overall, the full-width-at-half-maximum

(FWHM) of the fitted distribution of target responses was around 40°. For reference, the

average tuning width of individual neurons in macaque area MT is about 60° FWHM (Treue,

Hol, & Rauber, 2000). The average guessing rates (α) were around 26%, and did not change

across experimental conditions (F < 1).

To quantify how much each of the presented target and distractor signals influenced

observers' average-motion decisions, we used linear regression modelling with complex-

valued data (see Methods). The decision weights for all motion signals in all experimental

conditions were significantly different from zero (all p < .001), indicating that both task-

relevant and task-irrelevant signals contributed to the decision. As with the mixture

distribution modelling, initial analyses of decision weights revealed no significant

differences between target colour repetitions and changes (all Fs < 1.97, all ps > .170), so

the two conditions were averaged for further analyses. Overall, the weights for targets

(M/SEM = .44/.01, Fig. 1c) were much higher ($F_{2,64}$ = 326.80, p < .001, η_p^2 = .91) than the

weights for either concurrently presented distractors (.19/.01) or distractors presented on

their own (.13/.01). Unexpectedly, there was also a reliable order effect, with higher target

weights for the first epoch than for the second (.50/.01 vs. .38/.02, t_{32} = 4.74, p < .001);

analogously, distractor weights were lower for the first epoch than for the second (.14/.01

vs. .23/.01, t_{32} = 4.68, p < .001). Overall, the decision weight analyses revealed that

distractors, presented both with and without the target, contributed significantly to the

response, albeit to a lower degree than target motions.

Figure 1. Schematic of the stimuli, task and the behavioural results in Experiment 1. a, Example stimulus

sequence from a typical trial (above) together with the timing of colour saturation modulation and coherent-

motion onsets (below). Participants had to reproduce the *average* motion direction of target-coloured dots by

adjusting the orientation of a response dial (black pointer). After indicating their response they received

feedback on accuracy within the same display (white pointer). **b**, Group-level histograms (bars) of the error

magnitude (reproduced motion direction minus expected). The pink lines show fitted distributions of target

responses and guesses using mixture-distribution modelling of each individual participant's data. **c**, Average

decision weights showing the contributions of each presented motion direction to the decision. Since there

were no significant differences (t < 1) between the two distractor motions in the "No target" condition, only

the average of the two is shown. Grey dots denote individual participants.

We next characterised the temporal dynamics the CPP ERP. We were particularly

interested in the CPP in no-target epochs: if attentional selection and evidence

accumulation operate sequentially, then attention should suppress motion signals,

predicting no CPP in this epoch. By contrast, if selection and accumulation take place

concurrently, there should be a significant CPP even in the no-target epoch. Inspection of

the ERP topographies (Fig. 2a) revealed a central-medial positivity consistent with the usual

CPP component (O'Connell et al., 2012). The CPP (Fig. 2b) closely followed the time-course

of colour modulation, with three distinct peaks coinciding with the three epochs within each

trial. Between epochs, as the colour saturation of the dots returned to grey, the CPP

amplitude also returned close to baseline.

The whole-trial epochs were then divided into three segments relative to the onset

of colour saturation modulation, and baseline-corrected relative to the average of all three

epochs over a window of 100 ms prior to onset. Thus, all three epochs had the same

baseline, permitting comparisons between them. The epochs were then sorted according to

experimental conditions (first target, second target, no target; one- and two-target colours)

and averaged across trials (Fig. 2c). As we observed no effects of target colour repetitions

vs. changes (all $F_c < 1$), all subsequent analyses were conducted using the average of these

two trial types.

Overall, there was a statistically significant positive deflection in all experimental

conditions (Fig. 2c). The onset latency (522 ms) and the CPP slope (3.31 μV/s) were similar

across experimental conditions (all $F_c < 1$), suggesting that the rate of evidence

accumulation did not differ between conditions. The peak CPP amplitude, however,

decreased between epochs ($F_{c(2,64)} = 46.41$, $p < .001$), from the first target epoch to the

second and then to the no-target epoch (2.74, 2.43 and 1.06 µV, respectively, all $t_{c(32)} \geq$

1.93, p ≤ .031). Finally, the number of target colours (one vs two) had no significant effect

on the peak CPP amplitude, either as a main effect ($F_c < 1$) or in combination with epoch

($F_{c(2,64)}$ = 1.44, p = .244).

Time after onset of colour saturation modulation

Figure 2. Topography and time course of the CPP component in Experiment 1. a, Surface-Laplacian-filtered

EEG topography averaged across all conditions at times of peak colour saturation for all three epochs (1, 3, and

5 s after stimulus onset). The markers denote the electrode pair [CPz and Cz] used for the ERP analyses. **b**,

Time-course of the CPP averaged across all experimental conditions (M ± 1 within-participants SEM; Morey,

2008) for the total trial duration. **c**, CPP time-course during the gradual increase in colour saturation, shown

separately per epoch and experimental condition. Symbols above the x-axis denote time samples, averaged

across experimental conditions, for which the voltage was significantly different from 0 at $p_{FDR} < .05$. Full-line

and dashed-line arrows indicate onset latencies in the one and two target colours conditions, respectively. For

the purpose of data presentation, all traces were low-pass filtered at 10 Hz using a 4th order Butterworth IIR

filter. Data analyses were conducted using unfiltered data.

Taken together, the results of Experiment 1 revealed strong effects of selective

attention on decision-making. Behaviourally, target signals were associated with

significantly larger decision weights than distractors. At the neural level, attention had a

robust effect on the peak CPP amplitude, with higher peaks for target-present epochs than

for no-target epochs. Importantly, however, neither the CPP onset nor the slope differed

between target-present and no-target epochs, suggesting that both task-relevant and task-

irrelevant inputs triggered a process related to decision making. There was a robust and

significant influence of distractor events on behaviour, suggesting that attention attenuates

but does not fully suppress irrelevant inputs during integrated decision making. Significant

distractor weights and the CPP ERP in the no-target epoch are consistent with the

hypothesis that both target and distractor signals are integrated into a single decision

variable.

In Experiment 1, we used an increase in colour saturation to gradually reveal to

participants the task-relevance of the overlapping motion stimuli, and to render the

coherent motion signals discernible. While this design was clearly effective for measuring

the impact of distractors on decision making, it is possible that attentional filtering of

distractors might have been more successful if participants had been given an opportunity

to fully select the task-relevant stimuli (i.e., dot fields in the cued target colour) prior to the

onset of coherent motion events. To test this hypothesis, in Experiment 2 the trial epochs

were structured so that the colours of the dot stimuli reached full saturation prior to the

onset of coherent motion events, thus effectively separating selection-related and

response-related signals in time.

 Experiment 2

Materials and Methods

Participants. 36 healthy volunteers (20 females, average age 22.4 ± 4.4 years)

participated in the study. All had normal or corrected-to-normal visual acuity and normal

colour vision. Based on the same criteria as in Experiment 1, five participants were

identified as outliers and removed from further analyses, leaving 31 participants in total. All

other details were the same as in Experiment 1.

Stimuli, task and procedure. In Experiment 2 (Fig. 3a), the colour saturation

increased rapidly (.25 s) from grey to full saturation, and remained at maximum saturation

for 1 s. Coherent motion events were presented for .5 s during the maximum saturation

period, and at variable times (.50–.75 s asynchrony relative to the onset of colour

modulation).

To measure responses to targets and distractors independently, the dot fields

corresponding to target and distractor signals flickered (ON-OFF) concurrently at different

frequencies (12 and 18 Hz). The pairing between a given frequency and stimulus type (target

vs. distractor) was counterbalanced across trials. The introduction of these flickering

displays yielded distinct, time-resolved steady-state evoked potentials (SSVEPs, see Norcia

et al., 2015), which enabled us to read out the strength of target and distractor

representations on each trial. Previous research has shown that attention modulates the

SSVEP amplitude with stronger responses to task-relevant than irrelevant stimuli (Andersen

& Müller, 2010; O'Connell et al., 2012; Painter, Dux, Travis, & Mattingley, 2014) suggesting

that the SSVEP amplitude is closely related to attentional selection dynamics.

To optimise the display for recording SSVEPs, the field size, dot size and dot number

were doubled in Experiment 2 relative to Experiment 1. The dot-motion was generated at

2.5°/s, with an infinite dot life. Three distinctive colours (maximum saturation 75, minimum

colour separation 90°) were selected; two colours served as targets, and the remaining one

served as a distractor. The distractor and target colours were counterbalanced across

participants. As the SSVEP analyses permitted independent tracking of target and distractor

signals, only two epochs of dots were shown in Experiment 2, and both epochs within a trial

contained a target-motion patch (i.e., there was no longer a "no target" epoch). Participants

completed 64 practice trials followed by six blocks of 64 trials (384 trials in total). All other

details were the same as in Experiment 1.

EEG analyses. In Experiment 2, time-resolved SSVEP power at tagged frequencies

was computed using Morlet wavelets with fixed-width .5 s Gaussian taper, and normalized

relative to the neighbouring frequencies. The mean power per epoch was computed during

the 1 s of maximal colour saturation. For analyses of the target and distractor SSVEPs, the

power per epoch was baseline-corrected relative to a .1 s interval around the onset of

maximal colour saturation (.5 and 2.5 s, for the first and the second epochs, respectively).

All other analyses were the same as in Experiment 1.

Results

The behavioural results of Experiment 2 were qualitatively similar to those of

Experiment 1, with a continuous unimodal distribution of error magnitudes centred at 0

(Fig. 3b). Initial analyses revealed no effect of repetitions vs. changes in the target colour (all

$ts < 1$), so all subsequent analyses used the average of the two. Response precision (Fig. 3b)

was high and comparable between one- and two-target colour trials (K/FWHM = 7.96/48°

and 7.37/50°, respectively, $F_{1,30}$ = 2.37, p = .134). The guessing rates were low and virtually

identical for one- and two-target colour trials (ca. 12%, F < 1).

Figure 3. Schematic of the stimuli, task and the behavioural results in Experiment 2. a, Example stimulus

sequence from a typical trial (above) together with the timing of colour saturation modulation and coherent-

motion onsets (below). Only target-present epochs were presented, and the increase in colour saturation and

onset of coherent motion were separated in time. Dot patches were frequency-tagged at 12 and 18 Hz

(counterbalanced across target and distractor stimuli). **b**, Group-level histograms (bars) of the error magnitude

together with the best-fitting mixture-distribution models (pink lines). **c**, Average decision weights showing the

contributions of each presented motion direction to the decision. Grey dots denote individual participants.

Conventions as in Figure 1.

Analyses of decision weights revealed that both target and distractor signals

contributed to the average motion-direction response in all conditions (all p < .001). The

weights (Fig. 3c) for targets, however, were seven times higher than those for distractors

(.56/.01 and .08/.01, respectively, $F_{1,30}$ = 1,845, p < .001, η_p^2 = .98). The serial order effect

from Experiment 1 was replicated, with higher decision weights for the first epoch than for

the second, both for targets (.58/.01 vs. .54/.01, t_{30} = 3.06, p = .004) and distractors (.09/.01

vs. .07/.01, t_{30} = 3.32, p = .002). Finally, there was no significant difference between one- and

two-target colour trials, either as a main effect or in combination with the other factors (all

F ≤ 1.35, all p ≥ .254).

We next characterised the time-course of the CPP. The ERP topography (Fig. 4a) in

Experiment 2 was broadly consistent with that of Experiment 1, showing a central-medial

positivity. The time-course of the CPP (Fig. 4b) throughout the trial closely followed the

modulation of colour saturation, peaking sharply around .8 and 2.8 s and decaying quickly

thereafter. There was also a second modulation, broadly coinciding with the coherent

motion onsets (at around 1.5 and 3.5 s), which, due to the jittered timing of the motion

onsets, rose and fell more slowly than the colour-evoked components.

Analyses of the CPP (Fig. 4c) time-locked to the onset of colour modulation and

coherent motion, respectively, revealed a significant positive deflection in both cases.

Analyses of **colour-locked CPPs** for the first epoch relative to the second revealed a shorter

onset latency (220 vs. 231 ms, $F_{c(1,30)}$ = 5.04, p = .032), a marginally steeper slope (17.57 vs.

13.34 μV/s, $F_{c(1,30)}$ = 3.23, p = .083) and a higher peak amplitude (1.95 and 1.42 μV, $F_{c(1,30)}$ =

6.40, p = .017). In the second epoch, dividing attention between two target colours yielded a

significantly higher peak amplitude for two- versus one-target colour trials (1.56 and 1.29

μV, respectively; $t_{c(1,30)}$ = 2.30, p = .029) . Analyses of **motion-locked CPPs**, by contrast,

revealed no significant differences in onset latency (270 ms), slope (9.44 μV/s) or peak

amplitude (1.27 μV) between experimental conditions (all F_c <= 1.44, all ps >= .240). Overall,

the ERP analyses of Experiment 2 suggest that discerning the task-relevant colour patch (i.e.,

selection-relevant feature) and the coherent motion direction (i.e., response-relevant

feature) both engaged decision-making processes to some degree. Further, as only the

colour-locked CPP differed between the first and second epoch, it appears that the serial

order effect observed in both Experiments 1 and 2 was more closely related to processing of

colours rather than processing of the motion signal itself.

Figure 4. Topography and time course of the CPP component in Experiment 2. a, Surface-Laplacian-filtered

EEG topography averaged across all conditions at the middle of peak colour saturation for both epochs. **b**, The

CPP component averaged across all experimental conditions for the total trial duration. **c**, Time-course of the

CPP time-locked to the onset of colour saturation (left panels), and to the onset of coherent motion (right

panels), shown separately per epoch and experimental condition. Note the difference in scale between panels.

Conventions as in Figure 2.

To independently characterise the brain's response to target and distractor signals,

we next analysed the amplitude of SSVEP responses elicited by the frequency-tagged

motion stimuli. There was a strong (≈2 dB) frequency-specific response which peaked at the

Oz electrode over the occipital pole (Fig. 5a,b). Time-frequency analyses revealed that the

power of both target and distractor signals was significantly different from zero throughout

the trial (all $p_{FDR-corrected} < .05$). Moreover, the power was similar for target and distractor

patches prior to the increase in colour saturation in both the first and second epochs (Fig.

5c).

Figure 5. Time-course of target and distractor processing measured as SSVEP power in Experiment 2. a,

Topography of the total power at 12 Hz and 18 Hz normalized relative to the neighbouring frequencies,

computed using a fast-Fourier transform during periods of maximal colour saturation (.5–1.5 s, and 2.5–3.5 s).

b, Normalized total SSVEP power (M ± 95% CI) at Oz electrode during the period of maximal colour saturation

averaged across epochs and all experimental conditions. **c**, Time-resolved target- and distractor-evoked

SSVEPs for the whole trial duration averaged across all experimental conditions (M $\pm$ 1 within-participants

SEM). Black symbols above the x-axis denote times at which the difference between target and distractor

power was significant at $p_{FDR\text{-corrected}}$ < .05. **d**, Target-evoked (upper panels) and distractor-evoked (lower

panels) SSVEPs per epoch and per experimental condition baseline-corrected relative to .1 s interval around

the onset of maximum colour saturation (.5 and 2.5 s for the first and second epoch, respectively). Black

symbols above the x-axis denote times at which the baseline-corrected power was significantly different from

zero at p_{FDR} < .05.

As the colour saturation increased, there was a transient *decrease* in power for the

two patches in both the first epoch (around .5 s) and the second (around 2.5 s). Whereas

the power for target patches in both epochs recovered quickly (Fig. 5d, upper panels), the

power for distractor patches remained unchanged throughout the epochs (Fig. 5d, lower

panels), yielding a robust effect of task-relevance (average power: .24 and .10 dB for target

and distractor patches, respectively, $F_{1,30}$ = 11.64, p = .002, η_p^2 = .28). Finally, the SSVEP power

was comparable across the first and second epochs, and across one- and two-target colours

(all F $\leq$ 1.25, all p $\geq$ .272).

460   **Analyses of feature-specific brain responses**

In a final analysis, we characterised the strength of feature-specific brain responses

to coherent motion events in both Experiments 1 and 2 (Fig. 6a,b). Inspection of the motion

tuning in Experiment 1 revealed a phasic response to both target and distractor motion 0–.5

s after the onset of coherent motion. During the following 1 s period, a feature-specific

response developed gradually, peaking at around 1.5 s. Interestingly, throughout the epoch,

motion tuning was comparable for the target and distractor motion directions. To test the

effect of stimulus type (target vs distractor) on tuning strength, we conducted stepwise

468 linear mixed effects modelling with participant as a random effect, separately for each time

469 sample. In the first step, the model included only the average tuning strength per sample

470 (the intercept). In the second step, the stimulus type (target vs distractor) was added as a

471 fixed effect, and the increase in the goodness-of-fit between different models (Intercept

472 only vs Intercept + Stimulus) was assessed using a log-likelihood ratio test. The resulting p-

473 values for all time samples were corrected for multiple comparisons (FDR-correction). The

474 modelling revealed overall significant tuning at around .25 s and 1.5 s (Fig. 3a, lower panel,

475 the intercept). The effect of stimulus did not reach significance at any time point.

Figure 6. Time-course of feature-specific responses to coherent motion direction (i.e., motion tuning). a,

Upper panel shows tuning strength (M ± 1 within-participants SEM) averaged across one- and two-target

colours, and first and second target epochs, in Experiment 1. Higher values indicate stronger tuning. Lower

panel shows FDR-corrected p-values (-log10-transformed) for the overall tuning strength (Intercept) and the

effect of stimulus (Target vs Distractor) per time-sample. The horizontal line denotes .05 cut-off. **b**, Tuning

strength in Experiment 2. Conventions as in panel a. Note the difference in x and y scales between panels a

and b.

Inspection of motion tuning in Experiment 2 revealed strong tuning to target motion,

starting from around .2 s after the onset of coherent motion. By contrast, tuning to

distractor motion was negligible throughout the epoch. These observations were confirmed

by a stepwise LME model which revealed both strong motion tuning overall (the intercept)

and a strong effect of stimulus (target vs distractor). Relative to the tuning analyses

conducted in Experiment 1, tuning to target motion direction in Experiment 2 was roughly

four times stronger, consistent with the paradigm differences between experiments. More

specifically, whereas in Experiment 1 participants had to concurrently select the task-

relevant dot field and discern its motion direction, in Experiment 2 they could first select the

target field and then dedicate themselves to processing motion direction.

Overall, the results of the motion tuning analyses in Experiments 1 and 2 are broadly

consistent with the behavioural decision weights in the two experiments. In Experiment 1,

the behavioural target weights were four times stronger than the distractor weights,

whereas in Experiment 2 target weights were around seven times stronger. Similarly, the

tuning strength was overall numerically lower in Experiment 1 than in Experiment 2. These

results suggest that providing additional time for attentional selection (as in Experiment 2)

increases the precision with which task-relevant features are represented and decreases the

influence of concurrently presented task-irrelevant features.

502   **Discussion**

The goal of the present study was to characterise the behavioural and neural

processes associated with making integrated perceptual decisions. Unlike simple perceptual

decision-making (Forstmann et al., 2016; Gold & Shadlen, 2007; Ratcliff et al., 2016), which

requires judgments on a single, task-relevant stimulus, our decision-making task involved

selective processing of target signals in the presence of distractors, and required integration

of sensory evidence over an extended time period within a trial. The two main questions we

sought to address in this study were: (i) whether attentional selection and decision making

take place in parallel or sequentially, and (ii) whether task-relevant and task-irrelevant

sensory inputs are accumulated into a single decision variable.

To characterise the role of selective attention in decision making, we used mixture

distribution modelling and regression analyses. To characterise the neural correlates of

decision-making, we used a widely accepted correlate of decision making in humans, the

centro-parietal positivity (CPP) component (Hanks & Summerfield, 2017; O'Connell et al.,

2018; Spitzer et al., 2017). To monitor processing of different, task-relevant and task-

irrelevant sensory inputs, in Experiment 2 we flickered stimuli at different frequencies and

measured SSVEPs to targets and distractors. Finally, to characterise feature-specific brain

responses to coherent motion direction, we used a multivariate analysis approach that was

successfully used in the past to characterise feature-specific responses in a variety of tasks

(Ede et al., 2018; Wolff et al., 2017).

In Experiment 1, there was a measurable CPP in no-target epochs, coupled with a

robust and significant effect of distractor motion signals on behaviour. Taken together,

these results suggest that participants were unable to suppress distractor signals from being

accumulated into the decision variable. Importantly, both the peak CPP amplitude and the

decision weights in the no-target epochs were lower than the CPP amplitude and target

weights in target-present epochs, suggesting that selective attention modulated the

contribution of distractor motion to decision making. Consistent with Experiment 1, the

neural responses to distractors in Experiment 2, while significantly smaller than those to

targets, were not completely suppressed and remained stable over the period during which

target and distractor patches were discernible within the trial. There was also a statistically

significant distractor effect on responses, albeit smaller than the distractor effect in

Experiment 1. Finally, we also observed qualitative differences between Experiments 1 and

2. Specifically, the behavioural target weights were numerically higher and the distractor

weights numerically lower in Experiment 2 than Experiment 1. Similarly, the motion tuning

analyses revealed a significant difference in tuning to target and distractor motion

directions in Experiment 2, whereas there was no such difference in Experiment 1.

Taken together, the results of the two experiments suggest that: (i) selective

attention unfolds in parallel with evidence accumulation, and that (ii) both target and

distractor signals are accumulated into a single decision variable. As attentional selectivity

increases, distractor inputs are attenuated. Importantly, our findings demonstrate that

attention does not completely suppress the accumulation of task-irrelevant sensory input.

The differences between Experiments 1 and 2 can be parsimoniously explained by

differences in the experimental paradigm: in Experiment 1, coherent motion onset within

each trial coincided with modulation of colour saturation, whereas in Experiment 2 motion

onset followed modulation of colour saturation. Assuming that attentional selectivity at the

onset of coherent motion differed between experiments accounts for the discrepancies

between decision weights and motion tuning in a straightforward way.

Overall, the results of the present study are consistent with the notion that sensory

evidence accumulates in parallel (rather than sequentially) with an increase in selectivity for

task-relevant features. Computational modelling of response speed and accuracy in the

classic flanker task – in which target and distractor stimuli are presented concurrently at

different spatial locations (e.g., Eriksen, 1995) – has shown that similar principles apply to

selectivity for task-relevant locations (White et al., 2011). Of note, in a recent study Wyart et

al. (2015) had participants report the average orientation of one of two spatially separated

streams of achromatic gratings. Unlike the present study, decision weights for the spatially

separated distractors in their study were statistically indistinguishable from zero. While

differences in experimental tasks and analytical approaches prevent direct comparisons

between our study and that of Wyart et al. (2015), the two sets of findings suggest that

space- and feature-based filtering of sensory inputs during evidence accumulation may

exhibit different properties. Specifically, sensory evidence accumulation might be restricted

to a finite region of space so that under conditions in which signals are sufficiently

separated they engage independent accumulators.

Interestingly, across both the current experiments, having participants divide

attention between two potential target colours yielded no behavioural cost relative to

having them focus on a single target colour. In addition, there was no evidence of a cost for

switching between target colours across two epochs, compared with trials in which the two

colours repeated. Since in the present study the target colour patches were rendered

increasingly visible over a prolonged time interval and the responses were not speeded, it is

possible that our paradigm was insensitive to potential behavioural costs of dividing

attention (but see, e.g., Eimer and Grubert, 2014; Grubert et al., 2017). On the other hand,

the neural correlates of decision making, as characterised by the onset latency, slope, and

peak CPP amplitude, were consistent with the behavioural findings: the CPP time-traces

were virtually identical for one- and two-target colour trials in both experiments. As the

specifics of our experimental design prevent any strong conclusions, these findings call for

further investigation.

In both experiments, we also found a significant serial order bias with higher

decision weights for the first target epoch relative to the second. In Experiment 1, higher

target weights were accompanied by lower distractor weights, whereas in Experiment 2 the

weights for both target and distractor signals were higher in the first than the second epoch.

This primacy effect was not expected and, in fact, studies using similar paradigms have

typically reported either no order effects or even the opposite, recency effects (Cheadle et

al., 2014; Wyart, de Gardelle, Scholl, & Summerfield, 2012; Wyart et al., 2015). Mirroring

the behavioural primacy effects, the colour-locked peak CPP amplitude in Experiment 2 was

higher for the first than the second epoch. By contrast, analyses of the motion-locked CPP

revealed no order effects. These results suggest that the primacy effects might be related to

processing colours rather than motion. Since previous studies did not modulate attention-

related signals in the way we did here, the primacy effects we observed are not necessarily

inconsistent with the literature. Further studies will be necessary to characterise this effect

in more detail.

The motion averaging task we have introduced here has potential as a tool for

investigating the mechanisms underlying more complex, integrated decision-making.

Dynamic weighting of individual decisions to inform a single, integrated decision is

ubiquitous in daily life: for example, when deciding whether or not to cross a busy road. In

our task, individual motion signals are uncorrelated and, considered in isolation, are not

predictive of the final decision. Hence, the averaging task permits independent

parameterisation of both *simple* and *integrative* decision-making processes in a

straightforward and well characterised motion discrimination task. Using neuroimaging

methods with high spatial resolution such as fMRI, future studies could identify the brain

areas involved in simple decision-making, on the one hand, and integrative decision-making,

on the other. Using this task to characterise decision-making across the lifespan or in clinical

groups could also shed light on the development of integrative decision-making abilities

which are a hallmark of adaptive, intelligent behaviour.

Acknowledgments

This work was supported by the Australian Research Council (ARC) Centre of

Excellence for Integrative Brain Function (ARC Centre Grant CE140100007). JBM was

supported by an ARC Australian Laureate Fellowship (FL110100103). The funders had no

role in study design, data collection and analysis, decision to publish, or preparation of the

manuscript.

References

Andersen, S. K., & Müller, M. M. (2010). Behavioral performance follows the time course of neural facilitation and suppression during cued shifts of feature-selective attention. *Proceedings of the National Academy of Sciences*, *107*(31), 13878–13882. https://doi.org/10.1073/pnas.1002436107

Cheadle, S., Wyart, V., Tsetsos, K., Myers, N., de Gardelle, V., Herce Castañón, S., & Summerfield, C. (2014). Adaptive Gain Control during Human Perceptual Choice. *Neuron*, *81*(6), 1429–1441. https://doi.org/10.1016/j.neuron.2014.01.020

Churchland, A. K., & Ditterich, J. (2012). New advances in understanding decisions among multiple alternatives. *Current Opinion in Neurobiology*, *22*(6), 920–926. https://doi.org/10.1016/j.conb.2012.04.009

Churchland, A. K., Kiani, R., & Shadlen, M. N. (2008). Decision-making with multiple alternatives. *Nature Neuroscience*, *11*(6), 693–702. https://doi.org/10.1038/nn.2123

Ede, F. van, Chekroud, S. R., Stokes, M. G., & Nobre, A. C. (2018). Decoding the influence of anticipatory states on visual perception in the presence of temporal distractors. *Nature Communications*, *9*(1), 1449. https://doi.org/10.1038/s41467-018-03960-z

Ede, F. van, Chekroud, S. R., Stokes, M. G., & Nobre, A. C. (2019). Concurrent visual and motor selection during visual working memory guided action. *Nature Neuroscience*, *22*(3), 477. https://doi.org/10.1038/s41593-018-0335-6

Eimer, M., & Grubert, A. (2014). Spatial Attention Can Be Allocated Rapidly and in Parallel to New Visual Objects. *Current Biology*, *24*(2), 193–198. https://doi.org/10.1016/j.cub.2013.12.001

Eriksen, C. W. (1995). The flankers task and response competition: A useful tool for investigating a variety of cognitive problems. *Visual Cognition*, *2*(2).

Forstmann, B. U., Ratcliff, R., & Wagenmakers, E.-J. (2016). Sequential Sampling Models in Cognitive Neuroscience: Advantages, Applications, and Extensions. *Annual Review of Psychology*, *67*(1), 641–666. https://doi.org/10.1146/annurev-psych-122414-033645

Frătescu, M., Van Moorselaar, D., & Mathôt, S. (2019). Can you have multiple attentional templates? Large-scale replications of Van Moorselaar, Theeuwes, and Olivers (2014) and Hollingworth and Beck (2016). *Attention, Perception, & Psychophysics*, *81*(8), 2700–2709. https://doi.org/10.3758/s13414-019-01791-8

Gold, J. I., & Shadlen, M. N. (2007). The Neural Basis of Decision Making. *Annual Review of Neuroscience*, *30*(1), 535–574. https://doi.org/10.1146/annurev.neuro.29.051605.113038

Gramfort, A., Luessi, M., Larson, E., Engemann, D. A., Strohmeier, D., Brodbeck, C., … Hämäläinen, M. (2013). MEG and EEG data analysis with MNE-Python. *Frontiers in Neuroscience*, *7*. https://doi.org/10.3389/fnins.2013.00267

Grubert, A., Fahrenfort, J., Olivers, C. N. L., & Eimer, M. (2017). Rapid top-down control over template-guided attention shifts to multiple objects. *NeuroImage*, *146*(Supplement C), 843–858. https://doi.org/10.1016/j.neuroimage.2016.08.039

Hanks, T. D., & Summerfield, C. (2017). Perceptual Decision Making in Rodents, Monkeys, and Humans. *Neuron*, *93*(1), 15–31. https://doi.org/10.1016/j.neuron.2016.12.003

Heekeren, H. R., Marrett, S., & Ungerleider, L. G. (2008). The neural systems that mediate human perceptual decision making. *Nature Reviews. Neuroscience*, *9*(6), 467–479. https://doi.org/10.1038/nrn2374

Hollingworth, A., & Beck, V. M. (2016). Memory-based attention capture when multiple items are maintained in visual working memory. *Journal of Experimental Psychology. Human Perception and Performance*, *42*(7), 911–917. https://doi.org/10.1037/xhp0000230

Itthipuripat, S., Cha, K., Deering, S., Salazar, A. M., & Serences, J. T. (2018). Having More Choices Changes How Human Observers Weight Stable Sensory Evidence. *Journal of Neuroscience*, *38*(40), 8635–8649. https://doi.org/10.1523/JNEUROSCI.0440-18.2018

Kelly, S. P., & O'Connell, R. G. (2013). Internal and External Influences on the Rate of Sensory Evidence
Accumulation in the Human Brain. *The Journal of Neuroscience*, *33*(50), 19434–19441.
https://doi.org/10.1523/JNEUROSCI.3355-13.2013

Kohl, C., Spieser, L., Forster, B., Bestmann, S., & Yarrow, K. (2018). The Neurodynamic Decision Variable in
Human Multi-alternative Perceptual Choice. *Journal of Cognitive Neuroscience*, *31*(2), 262–277.
https://doi.org/10.1162/jocn_a_01347

Leite, F. P., & Ratcliff, R. (2010). Modeling reaction time and accuracy of multiple-alternative decisions.
Attention, Perception, & Psychophysics, *72*(1), 246–273. https://doi.org/10.3758/APP.72.1.246

Liu, T., & Pleskac, T. J. (2011). Neural correlates of evidence accumulation in a perceptual decision task. *Journal*
of Neurophysiology, *106*(5), 2383–2398. https://doi.org/10.1152/jn.00413.2011

Loughnane, G. M., Newman, D. P., Bellgrove, M. A., Lalor, E. C., Kelly, S. P., & O'Connell, R. G. (2016). Target
Selection Signals Influence Perceptual Decisions by Modulating the Onset and Rate of Evidence
Accumulation. *Current Biology*, *26*(4), 496–502. https://doi.org/10.1016/j.cub.2015.12.049

Morey, R. D. (2008). Confidence intervals from normalized data: A correction to Cousineau. *Tutorials in*
Quantitative Methods for Psychology , *4*, 61–64.

Nolan, H., Whelan, R., & Reilly, R. B. (2010). FASTER: Fully Automated Statistical Thresholding for EEG artifact
Rejection. *Journal of Neuroscience Methods*, *192*(1), 152–162.
https://doi.org/10.1016/j.jneumeth.2010.07.015

O'Connell, R. G., Dockree, P. M., & Kelly, S. P. (2012). A supramodal accumulation-to-bound signal that
determines perceptual decisions in humans. *Nature Neuroscience*, *15*(12), 1729–1735.
https://doi.org/10.1038/nn.3248

O'Connell, R. G., Shadlen, M. N., Wong-Lin, K., & Kelly, S. P. (2018). Bridging Neural and Computational
Viewpoints on Perceptual Decision-Making. *Trends in Neurosciences*, *41*(11), 838–852.
https://doi.org/10.1016/j.tins.2018.06.005

Peirce, J. W. (2007). PsychoPy—Psychophysics software in Python. *Journal of Neuroscience Methods*, *162*(1–2),
8–13. https://doi.org/10.1016/j.jneumeth.2006.11.017

Peirce, J. W. (2009). Generating Stimuli for Neuroscience Using PsychoPy. *Frontiers in Neuroinformatics*, *2*.
https://doi.org/10.3389/neuro.11.010.2008

Ratcliff, R., Philiastides, M. G., & Sajda, P. (2009). Quality of evidence for perceptual decision making is indexed
by trial-to-trial variability of the EEG. *Proceedings of the National Academy of Sciences*, *106*(16), 6539–
6544. https://doi.org/10.1073/pnas.0812589106

Ratcliff, R., Smith, P. L., Brown, S. D., & McKoon, G. (2016). Diffusion Decision Model: Current Issues and
History. *Trends in Cognitive Sciences*, *20*(4), 260–281. https://doi.org/10.1016/j.tics.2016.01.007

Scase, M. O., Braddick, O. J., & Raymond, J. E. (1996). What is Noise for the Motion System? *Vision Research*,
36(16), 2579–2586. https://doi.org/10.1016/0042-6989(95)00325-8

Schneegans, S., & Bays, P. M. (2016). No fixed item limit in visuospatial working memory. *Cortex*,
83(Supplement C), 181–193. https://doi.org/10.1016/j.cortex.2016.07.021

Shadlen, M. N., & Kiani, R. (2013). Decision Making as a Window on Cognition. *Neuron*, *80*(3), 791–806.
https://doi.org/10.1016/j.neuron.2013.10.047

Smith, P. L., & Ratcliff, R. (2009). An integrated theory of attention and decision making in visual signal
detection. *Psychological Review*, *116*(2), 283–317. https://doi.org/10.1037/a0015156

Spitzer, B., Blankenburg, F., & Summerfield, C. (2016). Rhythmic gain control during supramodal integration of
approximate number. *NeuroImage*, *129*, 470–479. https://doi.org/10.1016/j.neuroimage.2015.12.024

Spitzer, B., Waschke, L., & Summerfield, C. (2017). Selective overweighting of larger magnitudes during noisy
numerical comparison. *Nature Human Behaviour*, *1*(8), 0145. https://doi.org/10.1038/s41562-017-0145

Summerfield, C., & Tsetsos, K. (2015). Do humans make good decisions? *Trends in Cognitive Sciences*, *19*(1),
27–34. https://doi.org/10.1016/j.tics.2014.11.005

Theeuwes, J. (1992). Perceptual selectivity for color and form. *Perception & Psychophysics*, *51*(6), 599–606.

Treue, S., Hol, K., & Rauber, H.-J. (2000). Seeing multiple directions of motion—Physiology and psychophysics.
Nature Neuroscience, *3*(3), 270–276. https://doi.org/10.1038/72985
Twomey, D. M., Kelly, S. P., & O'Connell, R. G. (2016). Abstract and Effector-Selective Decision Signals Exhibit
Qualitatively Distinct Dynamics before Delayed Perceptual Reports. *Journal of Neuroscience*, *36*(28), 7346–
7352. https://doi.org/10.1523/JNEUROSCI.4162-15.2016
Ulrich, R., & Miller, J. (2001). Using the jackknife-based scoring method for measuring LRP onset effects in
factorial designs. *Psychophysiology*, *38*(5), 816–827. https://doi.org/10.1111/1469-8986.3850816
van Moorselaar, D., Theeuwes, J., & Olivers, C. N. L. (2014). In competition for the attentional template: Can
multiple items within visual working memory guide attention? *Journal of Experimental Psychology. Human*
Perception and Performance, *40*(4), 1450–1464. https://doi.org/10.1037/a0036229
White, C. N., Ratcliff, R., & Starns, J. J. (2011). Diffusion models of the flanker task: Discrete versus gradual
attentional selection. *Cognitive Psychology*, *63*(4), 210–238.
https://doi.org/10.1016/j.cogpsych.2011.08.001
Wolff, M. J., Jochim, J., Akyürek, E. G., & Stokes, M. G. (2017). Dynamic hidden states underlying working-
memory-guided behavior. *Nature Neuroscience*, *20*(6), nn.4546. https://doi.org/10.1038/nn.4546
Wyart, V., de Gardelle, V., Scholl, J., & Summerfield, C. (2012). Rhythmic Fluctuations in Evidence Accumulation
during Decision Making in the Human Brain. *Neuron*, *76*(4), 847–858.
https://doi.org/10.1016/j.neuron.2012.09.015
Wyart, V., Myers, N. E., & Summerfield, C. (2015). Neural Mechanisms of Human Perceptual Choice Under
Focused and Divided Attention. *Journal of Neuroscience*, *35*(8), 3485–3498.
https://doi.org/10.1523/JNEUROSCI.3276-14.2015
726