Selective attention modifies weights of distinct visual signals in human decision making

Authors: Dragan Rangelov[a] and Jason B. Mattingley[a,b,c]

Affiliations:

[a]Queensland Brain Institute, The University of Queensland, St Lucia, QLD 4072, Australia

[b]School of Psychology, The University of Queensland, St Lucia, QLD 4072, Australia

[c]Canadian Institute for Advanced Research (CIFAR)

Corresponding Author:

Dragan Rangelov

ORCID: 0000-0001-7648-6447

Post: Queensland Brain Institute

13   The University of Queensland

14   St Lucia, QLD 4072, Australia

Phone: +61 401 281 181

Email: d.rangelov@uq.edu.au

Keywords: selective attention, decision-making, electroencephalography, mixture

distribution modelling

Abstract

The ability to decide on the basis of multiple goal-relevant signals while ignoring

distractors is a hallmark of adaptive, intelligent behaviour. While attention and decision-

making processes have been well characterised in isolation, little is known about how the

two interact. To independently manipulate task relevance and information integration, we

presented moving dots in both a cued (target) colour and an uncued (distractor) colour.

Using electroencephalography, we recorded brain activity in human observers as they

reproduced the average motion direction of two consecutively presented targets. Although

completely spatially intermingled with the targets, the distractors had remarkably little

influence on either behavioural or electrophysiological correlates of decision-making.

Consistent with this, we found only target-related processing in a follow-up experiment

which used frequency-tagging to individually monitor target and distractor processing. Our

findings suggest that two task-relevant stimuli can be efficiently combined to inform

decision-making, implying a previously undocumented coupling between attention and

decision-making.

      Introduction

Cognition can be conceived of as a cascade of processes, beginning with the

encoding of sensory input and ending with appropriate behavioural choices. Sequential-

sampling decision-making models (Forstmann, Ratcliff and Wagenmakers, 2016; Ratcliff *et*

al., 2016) have had considerable success in describing choice behaviour in terms of both

behavioural performance and neural activity. According to these models, decision-making

relies on a central decision variable, an abstract quantity that accumulates in time toward

one of several possible choice alternatives. On these accounts, a choice is made once the

decision variable reaches threshold. In one widely employed decision-making task – the

random-dot motion discrimination task (Shadlen and Kiani, 2013; Ratcliff *et al.*, 2016) (see

Fig. 1a) – participants are shown fields of randomly moving dots and are asked to judge the

motion direction of a fraction of the dots that move coherently. Studies in both animals

(Shadlen and Kiani, 2013) and humans (Heekeren, Marrett and Ungerleider, 2008; Ratcliff,

Philiastides and Sajda, 2009; O'Connell, Dockree and Kelly, 2012; Kelly and O'Connell, 2013;

Twomey, Kelly and O'Connell, 2016) have shown that neural activity recorded during

random-dot motion discrimination follows a trajectory that resembles the accumulation of a

decision variable, thereby lending neurobiological support to sequential-sampling decision-

making models.

To date, much has been learned (Summerfield and Tsetsos, 2015; Spitzer, Waschke

and Summerfield, 2017) about simple perceptual decision-making in humans and other

animal species. In the real world, however, decisions are often more complex, requiring

judgements on stimuli that are separated in time and space, and which may or may not be

task relevant. For example, deciding to cross a busy road requires us to integrate the motion

directions of vehicles coming from both the left and right sides, while disregarding any

movements made by people or objects on the sidewalk. Little is known about how the brain

regulates decision-making when it receives competing – task-relevant and irrelevant –

sensory inputs. With a single, task-relevant dot-motion patch, integrating all dot trajectories

is clearly the optimal approach as it would yield strongest noise suppression. It remains

unclear, however, how efficiently observers can integrate only a task-relevant *subset* of dots

from amongst a spatially overlapping field of task-irrelevant stimuli. In this scenario,

integrating all dot trajectories is clearly suboptimal as the net motion direction would be the

average of both the relevant and irrelevant signals. To discern the task-relevant motion

direction, selectively attending to task-relevant visual input would be necessary. Despite the

critical role of attention for accurate decisions, most work on decision-making to date has

used only task-relevant dot patches. In the current study we characterised the behavioural

and neural mechanisms via which attention regulates complex perceptual decision-making.

To investigate whether and to what extent task-irrelevant, distractor signals

influence decision-making, we used random-dot displays (Fig. 1a) comprising *two*

intermingled dot patches moving in different directions at supra-threshold coherence. The

two patches had different colours, with one having a cued (target) colour and the other an

uncued (distractor) colour. As recent studies have suggested that task demands may

influence decision-making (Churchland, Kiani and Shadlen, 2008; Leite and Ratcliff, 2010;

Tsetsos, Usher and McClelland, 2011; Churchland and Ditterich, 2012; Loughnane *et al.*,

2016; Li *et al.*, 2017; Spitzer, Waschke and Summerfield, 2017), we cued participants, in

separate blocks of trials, to monitor either for a single target colour across epochs, or for

each of two different target colours.

To investigate integration of individual decisions over time, we measured

behavioural and neural responses while participants performed a novel, complex decision-

making task. We presented *two* epochs of target motion per trial and observers had to

reproduce the average target motion direction. This averaging task enabled us to

characterise the integration of two simple decisions into one, complex decision.

Participants' reproduction errors provided a continuous index of decision accuracy, which

we analysed using computational modelling, as described below, to quantify the unique

contributions of target and distractor events to the final decision. In addition, the averaging

task allowed us to dissociate decision-making from response selection and execution

processes, as the stimuli were orthogonal to the final response in this task.

Results

In Experiment 1, a group of 36 participants judged average motion direction across

three successive epochs within each trial (Fig. 1a). Participants were instructed to monitor

the motion direction of the target-coloured patches, and ignore the distractor-coloured

patches. Two, randomly selected epochs contained concurrent target and distractor motion

patches, while a third epoch contained two distractor patches. To quantify task

performance, we computed error magnitudes (Fig. 1b) by subtracting the true average

motion direction from the reproduced direction. The distributions of error magnitudes per

experimental condition were used to estimate the response precision (K) and guessing rates

(α), which reflect how well participants performed the task and how often they guessed,

respectively. The average guessing rates (α) decreased significantly ($F_{1,32} = 5.56$, p = .025, η_p^2 =

.15) as attentional demands increased: $\alpha_{M/SEM}$ =.34/.05 and .28/.05 for one- and two-target

colour conditions, respectively. The average response precision was high (one target colour:

$K_{M/SEM}$ = 10.90/1.77, 41° full-width-at-half-maximum (FWHM); two colours: $K_{M/SEM}$ =

12.72/1.92, 38° FWHM) and did not differ significantly between conditions ($F_{1,32}$ = 1.39, p =

.248). For reference, the average tuning width (Treue, Hol and Rauber, 2000) of individual

neurons in macaque area MT is about 60° FWHM. The excellent behavioural performance of

our participants demonstrates highly effective target selection from the concurrent and

spatially interspersed distractors.

Figure 1. Schematic of the stimuli, task and the behavioural results in Experiment 1. a, Example stimulus

sequence from a typical trial (above) together with the timing of colour saturation modulation and coherent-

motion onsets (below). The cue indicated task-relevant (target) colours in that trial. Two, randomly selected,

epochs of coloured dots comprised both target- and distractor-coloured dots (First and Second target),

whereas the remaining epoch comprised only distractor-coloured dots (No Target). Participants had to

reproduce the *average* motion direction of target-coloured dots by adjusting the orientation of a response dial

(black pointer) followed by a correct feedback within the same display (white pointer). **b**, Group-level

histograms (bars) of the error magnitude (reproduced motion direction – expected) in Experiment 1, shown

separately for one- and two-target colour trials. The lines show fitted distributions of target responses (full

lines) and guesses (dashed lines) using mixture-distribution modelling of each individual participant's data. **c**,

Average decision weights showing the contributions of each presented motion direction to the decision. The

yellow and green bars denote decision weights for target and distractor motion, respectively. Since there were

no significant differences (t < 1) between the two non-target motions in the "No target" condition (blue bars),

only the average of the two is shown. Grey dots denote individual participants.

To quantify how much each of the presented target and distractor signals influenced

observers' average-motion decisions, we used linear regression modelling with complex-

valued data (see Methods). The decision weights (Fig. 1c) were much higher ($F_{2,64} = 326.80$,

$p < .001$, $\eta_p^2 = .91$) for targets (.44/.01) than for either concurrently presented distractors

(.19/.01) or distractors presented on their own (.13/.01), indicating efficient attentional

selection of task-relevant signals. There was a reliable order effect with higher target

weights for the first epoch than for the second (.50/.01 vs. .38/.02, $t_{32} = 4.74$, $p < .001$);

analogously, distractor weights were lower for the first epoch than for the second (.14/.01

vs. .23/.01, $t_{32} = 4.68$, $p < .001$). Finally, the target weights were marginally higher for two

target colours relative to one (.46/.01 vs. .42/.01, , $t_{32} = 2.14$, $p = .047$), while attentional

demands had no effect on distractor weights ($t_{32} = 1.63$, $p = .113$).

We next analysed the dynamics of sensory and attentional processes feeding into

decision-making stages. If attention mediates perception and decision-making, then neural

correlates of attentional selection should be stronger for target-present than target-absent

epochs. To characterise attentional selection, we measured the onset latency, peak

amplitude and slope of a recently described (Loughnane *et al.*, 2016) event-related potential

(ERP) component, the N2b, which manifests as a bilateral negativity over posterior scalp

sites.

Figure 2. Time course of ERP correlates of attentional selection and decision making in Experiment 1. a,

Surface-Laplacian-filtered EEG topography averaged across all conditions at times of peak colour saturation for

all three epochs (1 s, 3 s, and 5 s after stimulus onset). The markers denote electrode pairs [O1 and O2] and

[CPz and Cz] which were used for ERP analyses. **b**, Time-course of target selection (the N2b, blue trace) and

decision-making (the CPP, red) ERPs averaged across all experimental conditions (M $\pm$ 1 within-participants

SEM (Morey, 2008)) for the total trial duration. **c**, Time-course of the N2b ERP during the gradual increase in

colour saturation, separately per epoch and experimental condition. **d**, Time-course of the CPP ERP during the

same period. For the purpose of data presentation, all traces were low-pass filtered at 10 Hz using a 4th order

Butterworth IIR filter. Data analyses were conducted using unfiltered data.

Inspection of the ERP topographies (Fig. 2a) revealed a bilateral posterior negativity

consistent with the N2b topography. The N2b time-traces (Fig. 2b) peaked at .7, 2.7, and 4.7

s, closely mirroring the modulation of colour saturation within each trial. The N2b onset

latency (403 ms, Fig. 2c) was comparable across experimental conditions ($F_c < 1$), indicating

that the modulation of colour saturation was registered irrespective of the colour's task-

relevance or attentional demands. Task-relevance, by contrast, had a significant effect on

the *slope* of the N2b ($F_{c(1,32)} = 3.58$, p = .034), with significantly steeper slopes for the target

epochs (-11.82 μV/s and -11.55 μV/s for the first and second epoch, respectively, $t_{c(32)} < 1$)

than for the no-target epoch (-4.02 μV/s). Mirroring the behavioural order effect, the N2b

peak amplitude was higher for the first target epoch than for the second (-3.59 μV and -2.84

μV, respectively), and both were higher than the no-target epoch (-2.04 μV, all $t_{c(32)} \geq 3.46$,

all p < .001). Also in line with the behavioural results, the robust differences between target-

and no-target epochs suggest that only task-relevant sensory input is passed on to decision-

making stages. Interestingly, increasing the attentional demands from one to two target

colours did not affect N2b slopes ($F_c < 1$). This finding nicely complements recent reports

(Eimer and Grubert, 2014; Grubert *et al.*, 2017) that spatial attention can be guided by two

attentional templates (i.e., two target colours) just as efficiently as by one template (i.e.,

one colour). Here we go beyond this work by showing that similar mechanisms control not

only spatial attention, but feature-based attention too. Critically, attentional demands

influenced the peak N2b amplitude which was significantly lower for one target colour than

for two (-2.69 and -2.96 μV, respectively, $F_{c(1,32)} = 4.96$, p = .033). Using a sequential-

sampling decision-making framework to interpret the N2b results, it appears that higher

attentional demands increased the decision threshold (as measured by the peak N2b

amplitude) for selecting the target colour, but did not affect the speed of evidence

accumulation (as measured by the N2b slope).

Next, we characterised the time-course of decision-making. If evidence accumulation

relies primarily on task-relevant signals, then neural correlates of evidence accumulation

should be stronger for target epochs relative to the no-target epoch. To characterise the

temporal dynamics of evidence accumulation, we quantified the onset latency, slope and

peak amplitude of an ERP known as the central-parietal positivity (CPP). Previous research

has shown that the time-course of the CPP closely resembles the time-course of evidence

accumulation: specifically, its amplitude builds up gradually (O'Connell, Dockree and Kelly,

2012), the build-up slope is proportional to stimulus quality (Kelly and O'Connell, 2013), and

it is observed even in the absence of overt responses (Twomey, Kelly and O'Connell, 2016).

Inspection of the ERP topographies (Fig. 2a) revealed a central-medial positivity consistent

with the usual CPP (O'Connell, Dockree and Kelly, 2012). Throughout the trial, the CPP (Fig.

2b) closely followed the time-course of colour modulation, with three distinct peaks

coinciding with the three episodes of maximum colour saturation. The onset latency (522

ms) and the CPP slope (3.31 μV/s) were similar across experimental conditions (all $F_c < 1$),

suggesting that the rate of evidence accumulation did not differ between epochs or

attentional demands. This is not entirely surprising, as the rate of evidence accumulation for

simple visual decisions has been shown to depend on motion coherence (Twomey, Kelly and

O'Connell, 2016), which was held constant in our paradigm. The peak CPP amplitude,

however, *decreased* between epochs ($F_{c(2,64)}$ = 46.41, p < .001), from the first target epoch

to the second epoch and then to the no target epoch (2.72, 2.44 and 1.10 µV, respectively,

all $t_{c(32)}$ ≥ 1.79, p ≤ .041).

Finally, attentional demands (one vs. two target colours) had no significant effect on

the peak CPP amplitude, either as a main effect (F_c < 1) or in combination with epoch ($F_{c(2,64)}$

= 1.44, p = .244). Overall, the CPP analyses revealed that the decision threshold (as indexed

by the peak CPP amplitude) was sensitive to the presence of a target motion patch,

suggesting that more sensory evidence was needed to reach a decision in target-present

than in no-target epochs.

Taken together, the ERP results of Experiment 1 revealed reliable differences

between target-present and no-target epochs for both the N2b and CPP components,

suggesting that complex visual decision-making – here involving the averaging of two

successive motion targets – relies on attentional selection of task-relevant stimuli. It

remains unclear, however, in what way does visual attention influence the processing of

concurrently presented target- and distractor-coloured signals: attention could boost target

encoding, suppress distractor processing, or potentially influence both. The goal of

Experiment 2 was to characterise how attention affects target and distractor processing in

our novel decision-making task. To measure responses to targets and distractors

independently, in Experiment 2 the dot fields corresponding to these two stimulus types

flickered concurrently at different frequencies (12 and 18 Hz; see Methods). The

introduction of these flickering displays yielded distinct, time-resolved steady-state evoked

potentials (SSVEPs, see Norcia *et al.*, 2015). Using a similar approach, studies on visual

attention have shown (Andersen and Müller, 2010; Andersen, Müller and Hillyard, 2015)

that when two differently-coloured, spatially overlapping random-dot patches are

presented, electrophysiological responses to the task-relevant, target patch are stronger

than those to concurrent distractor stimuli. Unlike previous studies, the modulation of

colour saturation enabled us to track attentional deployment within a trial as a function of

stimulus discriminability.

Experiment 2 differed from Experiment 1 in two further important ways (Fig. 3a).

First, since we wished to focus on the processing of concurrently presented target and

distractor colours, we removed no-target epochs and presented only target-present epochs.

Second, to separate in time neural processes associated with selection of the target versus

distractor colour from processes associated with motion processing, we first increased

colour saturation within each epoch, and then presented the critical coherent motion

events over a jittered interval during the period of maximum saturation. A new group of 36

participants took part in Experiment 2.

The behavioural results of Experiment 2 were strikingly similar to those of

Experiment 1. Response precision (Fig. 3b) was high and comparable between the one- and

two-target colour conditions (K/FWHM = 7.96/48° and 7.37/50°, $F_{1,30}$ = 2.37, p = .134).

Guessing rates were low and virtually identical for one and two target colours (.12 for both

conditions, F < 1). Decision weights (Fig. 3c) for targets were seven times higher than those

for distractors (.56/.01 and .08/.01, respectively, $F_{1,30}$ = 1,845, p < .001, η_p^2 = .98). We also

replicated the serial order effect, with higher decision weights for the first epoch than the

second, both for targets (.58/.01 vs. .54/.01, t_{30} = 3.06, p = .004) and distractors (.09/.01 vs.

.07/.01, t_{30} = 3.32, p = .002). Finally, increasing attentional demands from one to two target

colours had no effect on decision weights, either as a main effect or in combination with the

other factors (all F ≤ 1.35, all p ≥ .254).

Figure 3. Schematic of the stimuli, task and the behavioural results in Experiment 2. a, Example stimulus

sequence from a typical trial (above) together with the timing of colour saturation modulation and coherent-

motion onsets (below). Only target-present epochs were presented, and the increase in colour saturation and

onset of coherent motion were separated in time. Dot patches were frequency-tagged at 12 and 18 Hz

(counterbalanced across target and distractor stimuli). **b**, Group-level histograms (bars) of the error magnitude

together with the best-fitting mixture-distribution models (lines). **c**, Average decision weights showing the

contributions of each presented motion direction to the decision.

As in Experiment 1, we next characterised the time-course of attentional selection

and decision-making by analysing the N2b and CPP, respectively. The ERP topography (Fig.

4a) in Experiment 2 was broadly consistent with that of Experiment 1, showing a bilateral

posterior negativity and a centro-medial positivity. The time-course of the N2b and CPP (Fig.

4b) followed closely the modulation of colour saturation, peaking sharply around 1.8 and

3.8 s and decaying quickly thereafter. There was a second modulation of both components,

broadly coinciding with the coherent motion onsets (ca. 2.5 and 4.5 s), which rose and fell

more slowly than the colour-evoked components, most likely due to jittered timing of the

motion onsets.

Figure 4. Time course of ERP correlates of attentional selection and decision making in Experiment 2. a, Surface-Laplacian-filtered EEG topography averaged across all conditions at the middle of peak colour saturation for both epochs (2 s and 4 s after dot onset). The markers denote electrode pairs [O1 and O2] and [CPz and Cz] which were used for computing the N2b and CPP, respectively. **b**, The N2b (blue trace) and the CPP (red) averaged across all experimental conditions for the total trial duration. Notice the difference between the voltage scale for the N2b and CPP time-traces. **c**, Time-course of the N2b during the gradual increase in colour saturation, shown separately per epoch and experimental condition. **d**, Time-course of the CPP during the same period. Conventions as in Figure 2.

Mimicking the behavioural serial order effect, the N2b onset latency (Fig. 4c) was

somewhat shorter for the first epoch than for the second (235 and 240 ms, $F_{c(1,30)}$ = 4.10, p =

.052). Similarly, the N2b slopes were steeper and the peak N2b amplitude was higher in the

first epoch than in the second (slopes: -61.50 and -45.47 μV/s, $F_{c(1,30)}$ = 21.13, p < .001,

amplitudes: 1.95 and 1.41 μV, $F_{c(1,30)}$ = 88.31, p < .001). While for the first epoch, there were

no differences in the peak N2b amplitude for the one- and two-target conditions (t_c < 1), for

the second epoch, the peak N2b amplitude was higher for two-target than for one-target

colour trials (-5.17 and -4.28 μV, $t_{c(1,30)}$ = 3.30, p = .002). Similar to the N2b findings, analyses

of the CPP (Fig. 4d) revealed a shorter onset latency (220 and 231 ms, $F_{c(1,30)}$ = 5.04, p =

.032), a marginally steeper slope (17.57 and 13.34 μV/s, $F_{c(1,30)}$ = 3.23, p = .083) and a higher

peak amplitude (1.95 and 1.42 μV, $F_{c(1,30)}$ = 6.40, p = .017) for the first epoch relative to the

second. Increasing attentional demands had a marginally significant effect on the peak CPP

amplitude (1.62 and 1.57 μV, for one- and two target colours, respectively, $F_{c(1,30)}$ = 4.01, p =

.054) which was due to a higher peak amplitude for two target colours relative to one in the

second epoch ($t_{c(1,30)}$ = 2.30, p = .029). Overall, the ERP analyses of Experiment 2 replicated

the most salient findings of Experiment 1. First, the ERP time-course mirrored the

behavioural order effect and, second, an increase in attentional demands increased the

peak amplitudes, albeit only in the second epoch.

To independently characterise target and distractor processing, we next analysed

the amplitude of SSVEP responses elicited by the frequency-tagged motion stimuli. There

was a strong (≈2 dB, Fig. 5b), frequency-specific response which peaked at the Oz electrode

over the occipital pole (Fig. 5a).

Figure 5. Time-course of target and distractor processing measured as SSVEP power in Experiment 2. a,

Topography of the total power at 12 Hz and 18 Hz normalized relative to the neighbouring frequencies,

computed using a fast-Fourier transform during periods of maximal colour saturation (1.5–2.5 s, and 3.5–4.5

s). **b,** Normalized total SSVEP power (M ± 95% CI) at Oz electrode during the period of maximal colour

saturation averaged across epochs and all experimental conditions. **c,** Time-resolved target- and distractor-

evoked SSVEP for the whole trial duration averaged across all experimental conditions (M ± 1 within-

participants SEM). Black dots denote times at which the difference between target and distractor power was

significant at $p_{FDR-corrected} < .05$. **d,** Target-evoked SSVEPs per epoch and per experimental condition baseline-

corrected relative to the trough power in the -50–50 ms window around the peak colour saturation (1.5 s and

3.5 s for the first and the second epochs, respectively). **e**, Distractor-evoked baseline-corrected SSVEPs per

epoch and experimental condition.

Time-frequency analyses revealed that the power was similar for target and

distractor patches (Fig. 5c) prior to the increase in colour saturation in both the first epoch

and the second. This suggests that participants divided attention across both patches during

periods in which the two patches were indistinguishable. As the colour saturation increased,

there was a transient *decrease* in power for the two patches in both the first epoch (ca. 1.5

s) and the second (ca. 3.5 s). Whereas the power for target patches in both epochs (Fig. 5d,

grey dots) recovered quickly, the respective power for distractor patches remained at the

same (decreased) level throughout the epochs (Fig. 5e), yielding a robust effect of task-

relevance (average power: .24 and .10 dB for target and distractor patches, respectively, $F_{1,30}$

= 11.64, p = .002, η_p^2 = .28). Finally, the SSVEP power was comparable for the first and second

epoch, and for the one- and two-target colour conditions (all $F \leq 1.25$, all $p \geq .272$).

The transient decrease in power at the epoch onset (Fig. 5c) may reflect a gating

mechanism described in computational models of neuronal firing rates (Purcell *et al.*, 2010,

2012). In the absence of a stimulus, such gating prevents neuronal noise from reaching a

decision threshold by accident. Presenting a stimulus (in this case, increasing colour

saturation) would initially strengthen the gating, which might take time to overcome and

result in a transient decrease in power. A comparable decrease in power for both target and

distractor patches suggests that attentional selection does not modulate the gating. The

degree to which different patches were able to escape the gating, however, was strongly

influenced by task-relevance.

Taken together, the results of Experiment 2 suggest that, following initial

suppression of both target and distractor signals, attention selectively boosts task-relevant

target signals, but leaves task-irrelevant, distractor signals relatively unaffected. Consistent

with the behavioural decision weights, the SSVEP analyses revealed that participants

successfully selected targets and disregarded distractors, despite the fact that the two dot

patches were spatially overlapped at fixation.

Discussion

The central goal of the present study was to characterise the behavioural and neural

processes associated with complex decision-making. Unlike simple decision-making (Gold

and Shadlen, 2007; Forstmann, Ratcliff and Wagenmakers, 2016; Ratcliff *et al.*, 2016), our

complex decision-making task involves selective processing of target signals in the presence

of distractors and requires integration of several, independent decisions into a single,

'complex' decision. Although one could think of many other attributes of complex decision-

making, here we focused on these two specifically as we could characterise them readily

using the standard dot-motion assay of decision-making. The behavioural results of

Experiments 1 and 2 demonstrated that complex decision-making relies almost solely on

task-relevant sensory input, even when a similar but task-irrelevant signal is concurrently

presented. Since most previous studies on decision-making used only task-relevant stimuli,

the present study adds to the literature by demonstrating a profound influence of selective

attention on visual decision-making.

The averaging task that we developed permitted us to characterise integration of

independent decisions into one, complex decision. We found that complex decisions are

biased for sequential visual presentations, such that participants weight more heavily the

first of two successive target stimuli. As the two signals had identical, supra-threshold

motion coherence, differences in sensory processing of the first and the second epoch

cannot account for the order effect. Neither can a simple process of memory decay explain

the order effect, as this would predict stronger decision weights for the *second*, more recent

epoch, than the first. The sequential order bias we observed here echoes the "robust

averaging" effect reported in studies that require averaging of more than two signals

(Gardelle and Summerfield, 2011; Li *et al.*, 2017; Spitzer, Waschke and Summerfield, 2017).

In these tasks, stimuli that are similar to the grand average of all presented stimuli carry

greater weight than dissimilar stimuli. It is unlikely, however, that the order effect we

observed in the present study reflects robust averaging as the two target signals in our task

were, by design, equally informative. Instead, the order effect we observed suggests the

existence of a distinct, higher-level mechanism specifically involved in complex decision-

making.

In Experiment 1, analyses of a neural correlate of evidence accumulation, the CPP

ERP, revealed higher peak amplitude in epochs with target-coloured signals relative to no-

target epochs. As colour was always task-relevant, our findings suggest that *the degree* to

which task-relevant stimulus properties are processed depends on the match with the

relevant target features. In combination with previous studies (Twomey, Kelly and

O'Connell, 2016) which have found no CPP to completely task-irrelevant stimulus attributes

(e.g., orientation), our findings demonstrate that evidence accumulation evolves from

coarse stimulus representation (e.g., are the stimuli coloured or not?) to fine (e.g., are the

stimuli target- or distractor-coloured?). The fact that, in target-absent epochs, participants

were able to stop processing relatively quickly suggests that a decision can be made at

several levels of accumulated evidence, rather than after reaching a single, fixed decision-

threshold.

Finally, deciding about the motion direction of one or two possible target colours

mattered little for the CPP time-course in both Experiments 1 and 2. These results are of

particular importance for the decision-making literature. A critical distinction between

different evidence-accumulation models is whether they assume mutual inhibition between

different choices (Leite and Ratcliff, 2010) or independent evidence accumulation (Usher

and McClelland, 2001) across choice alternatives. The mutual inhibition architecture

predicts changes in decision-making dynamics as the number of choices increases. By cueing

our participants to attend to just one target colour or two colours we varied the number of

choice alternatives. By showing that the neural correlates of decision-making do not co-vary

with the number of choices – at least in the range of choice alternatives used in the present

study – our findings are consistent with the notion of independent evidence accumulation.

In Experiment 1, our index of the neural correlate of attentional selection (the N2b)

revealed a stronger response to task-relevant versus irrelevant stimuli. Likewise, in

Experiment 2 only target SSVEPs were modulated by visual attention, whereas the distractor

SSVEPs remained unaffected. These findings imply that the reason for participants' accurate

responses was effective attentional selection of task-relevant sensory input. Remarkably,

we also found that having participants maintain two attentional sets, as opposed to just

one, had no reliable influence on either behaviour or electrophysiological correlates of

attentional selection. Since target and distractor stimuli were always presented

concurrently within a foveated and attended region of space, these findings go beyond

recent work (Eimer and Grubert, 2014; Grubert *et al.*, 2017) which found that the time-

course of spatial attention changes little as attentional demands increase. Whereas previous

studies (Andersen, Hillyard and Müller, 2008; Andersen, Müller and Hillyard, 2015) using

similar methods found that multiple stimulus features *across* different dimensions (e.g.,

colour and orientation) can be attended in parallel, the present study shows that multiple

features *within* a single feature dimension (e.g., colour) can also be attended with little, if

any, cost.

The motion averaging task we have introduced here has great potential as a tool for

investigating the mechanisms underlying complex decision-making. Dynamic weighting of

individual decisions is ubiquitous in daily life: for example, when deciding whether or not to

cross a busy road. Our averaging paradigm captures both individual decisions and weighting

components of complex decision-making in a straightforward and well characterised motion

discrimination task. In our task, individual motion signals are uncorrelated and, considered

in isolation, are not predictive of the final decision. Hence, the averaging task permits

independent parameterisation of both *simple* and *complex* decision-making processes.

Using neuroimaging methods with high spatial resolution such as fMRI, future studies could

identify the brain areas involved in simple decision-making, on the one hand, and complex

decision-making, on the other.

The integrative aspect of complex decision-making is particularly interesting. In the

present study, we found that participants were biased such that they relied more strongly

on the first target epoch than the second. This finding suggests that integrating two simple

decisions into one, complex decision involves higher-level cognitive control processes.

Future studies could investigate determinants of decision biases in the averaging task: for

example, whether or not the order effect persists even when the two signals have different

coherence levels. As the averaging task can capture the integration process, using this task

to characterise complex decision-making across the lifespan or in clinical groups with

difficulties in cognitive control will shed light on the development of complex decision-

making abilities which are a hallmark of adaptive, intelligent behaviour.

Methods

Participants. Seventy-two healthy volunteers (39 females, average age 21.5 ± 2.5

years, 36 per experiment) participated in the study after providing written informed

consent. The sample size per experiment was estimated so that the statistical tests (t-test)

would have sufficient power (>= .80) to detect medium to strong effects (d >= .5). All

participants had normal or corrected-to-normal visual acuity and normal colour vision. In

Experiment 1, three participants were identified as outliers based on their behavioural

performance and/or EEG signals (see below for details). Data from five participants were

removed from Experiment 2, based on the same criteria, leaving 33 and 31 participants per

experiment, respectively. The study was approved by the Human Research Ethics

Committee of The University of Queensland, and was conducted in accordance with the

Human Subjects Guidelines of the Declaration of Helsinki.

Stimuli, task and procedure. In Experiment 1, participants were asked to monitor

the direction of successively presented epochs of coherent motion, and to indicate the

average direction of two target-motion events at the end of the trial using a response dial. A

response feedback symbol showing the correct answer was presented after every trial.

At the beginning of each trial, a central disk cued participants to look for one or two

target colours in the subsequent displays (Fig. 1a). This was followed by a buffer period

showing a field of 40 grey, randomly-moving dots. The dots' colour saturation gradually

increased (Fig. 1b) to reveal two overlapping, coloured-dot patches, before decreasing in

saturation and returning to grey. Each trial consisted of three such epochs – in two of these

epochs one of the two overlapping dot patches was rendered in the target colour, and the

other appeared in a distractor colour. Each trial contained two epochs in which a target was

present, and one epoch in which both dot patches were rendered in different distractor

colours (no-target epochs). Target patches were displayed equally often in the first, second

and third epochs of a trial. A single colour was cued in the first two blocks with a different

colour per block. In the remaining four blocks, two colours were cued. Between target-

present epochs, the target colours either repeated or changed equally often.

Dot-motion was generated at 40% coherence, with a speed of 19 dva/s, 100 ms dot-

life, different signal-dots across frames and random-walk noise-dots (Scase, Braddick and

Raymond, 1996). Note that we deliberately used a relatively high motion coherence value as

we wanted to measure participants' *average* decision to the two target events, rather than

their success in discriminating individual targets. Although coherent motion was present

throughout the epoch (1.5 s, Fig. 1b), different motion-directions were discernible only at

higher colour saturations. The motion directions were uniformly sampled from 0°–359°,

with the target-distractor and target-target difference remaining within 30°–150°. Five

easily distinguishable colours (Fig. 1b) were selected in the HSB colour-space with a

minimum hue difference of 72° and medium saturation and brightness (50). Two randomly-

selected colours served as targets and two as distractors. Participants completed 30 practice

trials followed by six blocks of 44 trials (= 264 trials). Each experimental session lasted

approximately one hour.

In Experiment 2, the field size, dot size and dot number were doubled relative to

Experiment 1, to optimise the display for recording SSVEPs. The dot-motion was generated

at 2.5 dva/s, with an infinite dot life. Three distinctive colours (maximum saturation 75,

minimum colour separation 90°, Fig. 3a) were selected; two colours served as targets, and

the remaining one served as a distractor. The distractor and target colours were

counterbalanced across participants. Different dot patches were frequency-tagged (Norcia

et al., 2015) by flickering (ON-OFF) the patches at 12 Hz and 18 Hz. Different frequencies

were randomly and equally often paired with target and distractor colours. The colour

saturation increased rapidly (.25 s) from grey to full saturation, and remained at maximum

saturation for 1 s. Only two epochs of coloured dots were shown, and both epochs within a

trial contained a target-motion patch. Coherent motion events were presented for .5 s

during the maximum saturation period, and at variable times (.25–.50 s after maximum

colour saturation was reached). Participants completed 64 practice trials followed by six

blocks of 64 trials (= 384 trials). All other details were the same as in Experiment 1.

Apparatus. The experiments were conducted in a dark, acoustically and

electromagnetically shielded room. The stimuli were presented on a 24" monitor with

1920×1080 resolution and a refresh rate of 144 Hz. The experimental software was custom-

coded in Python using the PsychoPy (Peirce, 2007, 2009) toolbox. EEG signals were recorded

using 64 Ag-AgCl electrodes (BioSemi ActiveTwo) arranged in the 10-20 layout and sampled

at 1,024 Hz.

Behavioural analyses. To identify outlier participants, the distributions of error

magnitudes (i.e., the angular difference between the response and the correct answer)

were compared to a uniform distribution (i.e., pure guessing) using the Kolmogorov-Smirnov

test. Participants for whom the probability of the null hypothesis (i.e., a uniform distribution

of error magnitudes) exceeded .001 were removed from further analyses. The remaining

distributions per experimental condition and per participant were fitted to a theoretical

model (Schneegans and Bays, 2016), and responses were separated into noisy target

responses and random guesses. To quantify decision weights, a multiple-regression (OLS)

model with a term for each of the presented motion directions, expressed as complex

numbers, was fitted to the responses, separately per participant and experimental

condition. The absolute value of the resulting regression coefficients reflects the influence

of each of the presented coherent motion signals on the response, i.e., its *decision weight*.

EEG analyses. EEG signals were analysed using the MNE-Python (Gramfort *et al.*,

2013) toolbox. The data were offline re-referenced to the average electrode, low-pass

filtered at 99 Hz and notch-filtered at 50 Hz to eliminate line noise. The recorded signal was

pre-processed using the FASTER (Nolan, Whelan and Reilly, 2010) algorithm for automated

artefact rejection. The pre-processed signal was down-sampled to 256 Hz, segmented into

epochs between the cue-onset and the response-display onset, baseline-corrected relative

to -.1–0 s pre-trial and linearly de-trended. Outlier epochs and participants were identified

using the FASTER algorithm and removed from further analyses. The N2b and CPP ERPs

were computed by averaging O1/O2 and Cz/CPz electrodes, respectively, across trials. Each

motion epoch within a trial was extracted from the onset of the increase in colour

saturation and baseline-corrected relative to a common baseline during -.1–0 s, averaged

across all epochs. The peak N2b and CPP amplitudes were identified using the jackknife-

method (Ulrich and Miller, 2001). Using the same method, ERP onset latencies were

computed at 50% peak amplitude. The ERP slopes were computed as the amplitude

difference between -.05 s and .05 s relative to the ERP onset, and divided by .1 s so as to

yield $\mu V/s$. Time-resolved SSVEP power at tagged frequencies was computed using Morlet

wavelets with fixed-width .5 s Gaussian taper, and normalized relative to the neighbouring

frequencies. For analyses of the target (Fig. 5d) and distractor SSVEPs (Fig. 5e), the power

per epoch was baseline-corrected relative to the mean power in the -.05–.05 s interval

around the onset of maximal colour saturation (1.5 and 3.5 s, for the first and the second

epoch, respectively). The mean power per epoch was computed during the 1 s of maximal

colour saturation.

Acknowledgments

This work was supported by the Australian Research Council (ARC) Centre of

Excellence for Integrative Brain Function (ARC Centre Grant CE140100007). JBM was

supported by ARC Australian Laureate Fellowship (FL110100103).

References

Andersen, S. K., Hillyard, S. A. and Müller, M. M. (2008) 'Attention Facilitates Multiple Stimulus Features in
Parallel in Human Visual Cortex', *Current Biology*, 18(13), pp. 1006–1009. doi: 10.1016/j.cub.2008.06.030.

Andersen, S. K. and Müller, M. M. (2010) 'Behavioral performance follows the time course of neural facilitation
and suppression during cued shifts of feature-selective attention', *Proceedings of the National Academy of*
Sciences, 107(31), pp. 13878–13882. doi: 10.1073/pnas.1002436107.

Andersen, S. K., Müller, M. M. and Hillyard, S. A. (2015) 'Attentional Selection of Feature Conjunctions Is
Accomplished by Parallel and Independent Selection of Single Features', *Journal of Neuroscience*, 35(27),
pp. 9912–9919. doi: 10.1523/JNEUROSCI.5268-14.2015.

Churchland, A. K. and Ditterich, J. (2012) 'New advances in understanding decisions among multiple
alternatives', *Current Opinion in Neurobiology*. (Decision making), 22(6), pp. 920–926. doi:
10.1016/j.conb.2012.04.009.

Churchland, A. K., Kiani, R. and Shadlen, M. N. (2008) 'Decision-making with multiple alternatives', *Nature*
Neuroscience, 11(6), pp. 693–702. doi: 10.1038/nn.2123.

Eimer, M. and Grubert, A. (2014) 'Spatial Attention Can Be Allocated Rapidly and in Parallel to New Visual
Objects', *Current Biology*, 24(2), pp. 193–198. doi: 10.1016/j.cub.2013.12.001.

Forstmann, B. U., Ratcliff, R. and Wagenmakers, E.-J. (2016) 'Sequential Sampling Models in Cognitive
Neuroscience: Advantages, Applications, and Extensions', *Annual Review of Psychology*, 67(1), pp. 641–
666. doi: 10.1146/annurev-psych-122414-033645.

Gardelle, V. de and Summerfield, C. (2011) 'Robust averaging during perceptual judgment', *Proceedings of the*
National Academy of Sciences, 108(32), pp. 13341–13346. doi: 10.1073/pnas.1104517108.

Gold, J. I. and Shadlen, M. N. (2007) 'The Neural Basis of Decision Making', *Annual Review of Neuroscience*,
30(1), pp. 535–574. doi: 10.1146/annurev.neuro.29.051605.113038.

Gramfort, A. *et al.* (2013) 'MEG and EEG data analysis with MNE-Python', *Frontiers in Neuroscience*, 7. doi:
10.3389/fnins.2013.00267.

Grubert, A. *et al.* (2017) 'Rapid top-down control over template-guided attention shifts to multiple objects',
NeuroImage, 146(Supplement C), pp. 843–858. doi: 10.1016/j.neuroimage.2016.08.039.

Heekeren, H. R., Marrett, S. and Ungerleider, L. G. (2008) 'The neural systems that mediate human perceptual
decision making', *Nature Reviews. Neuroscience*, 9(6), pp. 467–479. doi: 10.1038/nrn2374.

Kelly, S. P. and O'Connell, R. G. (2013) 'Internal and External Influences on the Rate of Sensory Evidence
Accumulation in the Human Brain', *The Journal of Neuroscience*, 33(50), pp. 19434–19441. doi:
10.1523/JNEUROSCI.3355-13.2013.

Leite, F. P. and Ratcliff, R. (2010) 'Modeling reaction time and accuracy of multiple-alternative decisions',
Attention, Perception, & Psychophysics, 72(1), pp. 246–273. doi: 10.3758/APP.72.1.246.

Li, V. *et al.* (2017) 'Robust averaging protects decisions from noise in neural computations', *PLOS*
Computational Biology, 13(8), p. e1005723. doi: 10.1371/journal.pcbi.1005723.

Loughnane, G. M. *et al.* (2016) 'Target Selection Signals Influence Perceptual Decisions by Modulating the
Onset and Rate of Evidence Accumulation', *Current Biology*, 26(4), pp. 496–502. doi:
10.1016/j.cub.2015.12.049.

Morey, R. D. (2008) 'Confidence intervals from normalized data: A correction to Cousineau', in *Tutorials in*
Quantitative Methods for Psychology , 4, pp. 61–64.

Nolan, H., Whelan, R. and Reilly, R. B. (2010) 'FASTER: Fully Automated Statistical Thresholding for EEG artifact
Rejection', *Journal of Neuroscience Methods*, 192(1), pp. 152–162. doi: 10.1016/j.jneumeth.2010.07.015.

Norcia, A. M. *et al.* (2015) 'The steady-state visual evoked potential in vision research: A review', *Journal of*
Vision, 15(6), pp. 4–4. doi: 10.1167/15.6.4.

O'Connell, R. G., Dockree, P. M. and Kelly, S. P. (2012) 'A supramodal accumulation-to-bound signal that
determines perceptual decisions in humans', *Nature Neuroscience*, 15(12), pp. 1729–1735. doi:
10.1038/nn.3248.

Peirce, J. W. (2007) 'PsychoPy—Psychophysics software in Python', *Journal of Neuroscience Methods*, 162(1–
2), pp. 8–13. doi: 10.1016/j.jneumeth.2006.11.017.

Peirce, J. W. (2009) 'Generating Stimuli for Neuroscience Using PsychoPy', *Frontiers in Neuroinformatics*, 2.
doi: 10.3389/neuro.11.010.2008.

Purcell, B. A. *et al.* (2010) 'Neurally Constrained Modeling of Perceptual Decision Making', *Psychological*
Review, 117(4), pp. 1113–1143. doi: 10.1037/a0020311.

Purcell, B. A. *et al.* (2012) 'From Salience to Saccades: Multiple-Alternative Gated Stochastic Accumulator
Model of Visual Search', *The Journal of Neuroscience*, 32(10), pp. 3433–3446. doi:
10.1523/JNEUROSCI.4622-11.2012.

Ratcliff, R. *et al.* (2016) 'Diffusion Decision Model: Current Issues and History', *Trends in Cognitive Sciences*,
20(4), pp. 260–281. doi: 10.1016/j.tics.2016.01.007.

Ratcliff, R., Philiastides, M. G. and Sajda, P. (2009) 'Quality of evidence for perceptual decision making is
indexed by trial-to-trial variability of the EEG', *Proceedings of the National Academy of Sciences*, 106(16),
pp. 6539–6544. doi: 10.1073/pnas.0812589106.

Scase, M. O., Braddick, O. J. and Raymond, J. E. (1996) 'What is Noise for the Motion System?', *Vision*
Research, 36(16), pp. 2579–2586. doi: 10.1016/0042-6989(95)00325-8.

Schneegans, S. and Bays, P. M. (2016) 'No fixed item limit in visuospatial working memory', *Cortex*,
83(Supplement C), pp. 181–193. doi: 10.1016/j.cortex.2016.07.021.

Shadlen, M. N. and Kiani, R. (2013) 'Decision Making as a Window on Cognition', *Neuron*, 80(3), pp. 791–806.
doi: 10.1016/j.neuron.2013.10.047.

Spitzer, B., Waschke, L. and Summerfield, C. (2017) 'Selective overweighting of larger magnitudes during noisy
numerical comparison', *Nature Human Behaviour*, 1(8), p. 0145. doi: 10.1038/s41562-017-0145.

Summerfield, C. and Tsetsos, K. (2015) 'Do humans make good decisions?', *Trends in Cognitive Sciences*, 19(1),
pp. 27–34. doi: 10.1016/j.tics.2014.11.005.

Treue, S., Hol, K. and Rauber, H.-J. (2000) 'Seeing multiple directions of motion—physiology and
psychophysics', *Nature Neuroscience*, 3(3), pp. 270–276. doi: 10.1038/72985.

Tsetsos, K., Usher, M. and McClelland, J. L. (2011) 'Testing Multi-Alternative Decision Models with Non-
Stationary Evidence', *Frontiers in Neuroscience*, 5. doi: 10.3389/fnins.2011.00063.

Twomey, D. M., Kelly, S. P. and O'Connell, R. G. (2016) 'Abstract and Effector-Selective Decision Signals Exhibit
Qualitatively Distinct Dynamics before Delayed Perceptual Reports', *Journal of Neuroscience*, 36(28), pp.
7346–7352. doi: 10.1523/JNEUROSCI.4162-15.2016.

Ulrich, R. and Miller, J. (2001) 'Using the jackknife-based scoring method for measuring LRP onset effects in
factorial designs', *Psychophysiology*, 38(5), pp. 816–827. doi: 10.1111/1469-8986.3850816.

Usher, M. and McClelland, J. L. (2001) 'The time course of perceptual choice: the leaky, competing
accumulator model', *Psychological Review*, 108(3), pp. 550–592.

a

Cue

Buffer

1st Epoch

2nd Epoch

3rd Epoch

Response

2 dva

7.8 dva

.3 dva

1st Target

No Target

2nd Target

Coherent motion

Saturation

.5 s

b

One target colour

Two target colours

Fitted target responses — — Fitted guesses

Proportion of trials

0.3

0.2

0.1

0.0

$-\pi$ 0 π

$-\pi$ 0 π

c

First target

Second target

No target

Decision weights [a.u.]

0.6

0.4

0.2

0.0

Target

Distractor

One target colour   Two target colours

a

Cue

Buffer

1st Epoch

2nd Epoch

Response

2 dva

15.6 dva

.6 dva

Coherent motion — Onset jitter

Saturation

.5 s

b

One target colour   Two target colours

—— Fitted target responses   - - - Fitted guesses

Proportion of trials

0.3

0.2

0.1

0.0

-π   0   π   -π   0   π

c

First target   Second target

Decision weights [a.u.]

0.6

0.4

0.2

0.0

Target

Distractor

One target colour   Two target colours

a

12 Hz   18 Hz

2
1 dB
0

b

1 dB
-1 dB

12   18 Hz

c

First target   Second target

.5 dB

—— Target
—— Distractor

1.5   2.0   2.5   3.0   3.5   4.0   4.5 s   5.0 s

—— One target colour   - - - Two target colours   ● p_{FDR} < .05

d

.25 dB

1.5   1.75   2.0   2.25   2.5 s   3.5   3.75   4.0   4.25   4.5 s

e

.25 dB

1.5   1.75   2.0   2.25   2.5 s   3.5   3.75   4.0   4.25   4.5 s