Causes and consequences of cellulose production by *Pseudomonas fluorescens* SBW25 at the air-liquid interface

Maxime Ardré[1,*] and Paul B Rainey[1,2,*]

[1]Laboratoire de Génétique de l'Evolution, Ecole Superieure de Physique et de Chimie Industrielles de la Ville de Paris (ESPCI Paris Tech), CNRS UMR 8231, PSL Research University, 75231 Paris, France.

[2]Department of Microbial Population Biology, Max Planck Institute for Evolutionary Biology, Plön 24306, Germany.

[*]For correspondence: maxime.ardre@espci.fr and rainey@evolbio.mpg.de

Keywords: diguanylate cyclase — bioconvection — pyoverdin — Raleigh-Taylor instability — continuum field models — ecological complexity

Running title: Cellulose and colonisation of the air-liquid interface

ABSTRACT

Cellulose over-producing wrinkly spreader mutants of *Pseudomonas fluorescens* SBW25 have been the focus of much investigation, but conditions promoting the production of cellulose in ancestral SBW25, its effects and consequences have escaped investigation through lack of *in vitro* phenotype. Here, using a custom built device, we reveal that in static broth microcosms ancestral SBW25 encounters environmental signals at the air-liquid interface that activate, via three diguanylate cyclase-encoding pathways (Wsp, Aws and Mws), production of cellulose. Secretion of the polymer at the meniscus leads to modification of the environment and growth of numerous micro-colonies that extend out from the surface. Accumulation of cellulose and associated microbial growth leads to Rayleigh-Taylor instability resulting in bioconvection and rapid transport of water-soluble products over tens of millimetres. Drawing upon data we build a mathematical model that recapitulates experimental results and captures the interactions between biological, chemical and physical processes.

IMPORTANCE This work reveals a new bacterial behaviour that manifests at the air-liquid interface, which depends on production of cellulose, and hints to hitherto unrecognised dimensions to bacterial life at surfaces. Additionally, the study links activation of known diguanylate cyclase-encoding pathways to cellulose expression and to signals encountered at the air-liquid interface. Further significance

48   stems from recognition of the consequences of fluid instabilities

49   arising from surface production of cellulose for transport of water-

50   soluble products over large distances.

Introduction

Surfaces are frequently colonised by microbes. Surface-associated microbes grow as dense populations / communities termed "biofilms" (1-3). Growth at surfaces provides microbes with nutrients and opportunities for cross-feeding (4,5). For the pathogens surface colonisation is often a prelude to invasion (6,7). Microbes in high-density populations can find protection against external factors such as antibiotics and toxic agents (8). At the same time, microbes in biofilms experience intense competition for resources and can be negatively impacted by costs associated with exposure to metabolic waste products (9). For long-term survival, escape from surfaces and dispersal is crucial (10).

Much attention has been given to colonisation of solid-liquid surfaces (11,12). This owes as much to the importance of these surfaces as it does the ease with which they can be studied. For example, colonisation of abiotic surfaces can be measured by simple histochemical assay or by microscopic observation using flow cells (13,14). Decades of study have revealed much about adhesive factors including polymers and proteinaceous adhesions involved in surface attachment and the regulatory pathways controlling their expression (15). A particular focus has been pathways for synthesis and degradation of the secondary signalling molecule cyclic-di-GMP (16). For the most part, the precise signals activating these regulatory pathways are unclear. Moreover, the frequent use of mutants —

sometimes intentionally but often inadvertently — that constitutively

over-produce adhesive factors has stymied progress in understanding

many subtleties surrounding surface colonisation.

Surfaces are also a feature of the interface between gas and liquid,

but colonisation of such surfaces has been little studied (17,18).

Air-liquid interfaces (ALIs) are of special relevance for aerobic

organisms because colonisation of the meniscus provides access to

oxygen that is typically absent from the broth phase. While many

motile aerobic bacteria display taxis toward oxygen, this alone may

be insufficient to allow cells to contend with the forces required to

overcome the effects of surface tension and colonise the meniscus.

Once achieved, in the absence of mechanisms to achieve buoyancy,

cells growing at the interface must contend with the effects of

gravity.

The interface between air and liquid has further significance in that

it often marks the divide between aerobic and anaerobic conditions.

This has implications for surface chemistry with ensuing

physiological effects for bacteria. For example, iron, an essential

element, exists in the insoluble and biologically unavailable ferric

form in the presence of oxygen, but is water soluble and freely

available in the absence of oxygen (19). Bacteria growing within an

initially resource-rich and oxygen replete broth phase consume oxygen

and thus further growth requires access to the ALI (20). Bacteria

that achieve colonisation of this surface must then contend with iron

deplete conditions requiring the synthesis of siderophores (21).

To date, studies of colonisation of the ALI have been largely centred

on genotypes that constitutively produce polymers such as cellulose

(22). Often these have arisen as a consequence of selection

experiments in static broth microcosms where mutants with

constitutively active diguanylate cyclases (and ensuing constitutive

production of the respective polymers) have a selective advantage

that arises from capacity to form a dense microbial mat at the air-

liquid interface (23-25). While such mutants have made clear the

central importance of cellulose and related polymers the generality

of conclusions arising from the use of constitutively active mutants

need to be treated with caution (24). Desirable would be analysis of

the biophysics of ALI colonisation in wild type bacteria where

regulation of polymer production is unaffected by mutation.

Almost two decades ago it was reported that in well-mixed culture the

fitness of a cellulose-defective mutant of *Pseudomonas fluorescens*

SBW25 was equivalent to that of the wild type (ancestral) bacterium

(22). Also reported in that study was a significant reduction in

fitness of a cellulose defective mutant in static broth culture, but

the reasons were not determined. Recent observations of the growth of

a cellulose-defective mutant of wild type (ancestral) SBW25 made

during the course of analyses of evolutionary convergence in polymer

production by SBW25 (26) led to the realisation of a subtle phenotype

associated with absence of growth in the cellulose-defective mutant

at the air-liquid interface. Unlike ancestral SBW25, it grows

exclusively within the broth phase with ensuing negative effects of

oxygen limitation responsible for its previously noted low fitness

(22).

Here we seek to understand the biological role of cellulose and do so

via a device that combines spectrophotometry with multi-perspective

time-lapse imaging. Aided by the device we monitor surface growth,

reveal the contribution made by cellulose and show that it involves

regulatory contributions from three known diguanylate cyclase-

encoding regulatory pathways. The production of cellulose allows

formation of a lawn of micro-colonies at the meniscus that eventually

coalesce into a thin film of bacteria. The mass of bacteria and

cellulose generates a gravitational force that leads to Rayleigh

Taylor instability and causes bioconvection (27). One consequence of

bioconvection is the rapid transport of the water-soluble iron-

binding siderophore, pyoverdin. A mathematical model based on

partial differential equations with fluid dynamics described by the

Navier-Stokes (NS) equation with Boussinesq approximation accounts

for the diffusion-reaction and convection processes occurring in the

microcosm.

RESULTS

Describing microbial colonisation of the air-liquid interface (hereafter ALI), although in principle straightforward, is fraught with difficulty. While advanced microscopic techniques offer possibilities to observe colonisation at the single cell level, much stands to be gained from more macroscopic perspectives, aided by low power microscopy in conjunction with time-lapse photography.

Device. To understand and quantitate growth of ancestral SBW25 and the cellulose-defective mutant SBW25 Δ*wssA-J* a device was constructed that allows growth at the ALI and in the broth phase to be monitored from multiple perspectives (Fig. 1). It comprises three cameras: one placed perpendicular to the microcosm to record growth within the microcosm and on the under surface of the meniscus, one mounted at a 45° angle above the ALI to capture surface growth and one to detect the light emitted from excitation of the fluorescent signal arising from production of the iron-chelating siderophore, pyoverdin. Additionally, the device incorporates a laser and corresponding photodiode to vertically scans the flask at regular (5 min) time intervals.

Cellulose is required for colonisation of the ALI. Figure 2 shows the growth dynamics of ancestral SBW25 and SBW25 Δ*wssA-J* determined by the scanning laser and calibrated using direct plate counts. SBW25 Δ*wssA-J* is slower to enter exponential growth than SBW25, it grows at

approximately the same rate (SBW25, $0.53+/-0.02$ h^{-1}; SBW25 ΔwssA-J

$0.57+/-0.03$ h^{-1}), but density in stationary phase is consistently

lower. Notable in SBW25 at ~24 hours is a reproducible plateau of

growth followed by a further increase and a widening of difference in

cell density compared to SBW25 $\Delta wssA-J$ (Fig. 2). No such

intermediate plateau occurs in the cellulose mutant.

Time-lapse observation of the ALI from a 45° angle in flasks

inoculated with SBW25 reveal presence of a thin film at 19 h that is

more prominent at 26 h and still evident albeit weakly at 40 h (Fig.

3a and supplementary movie file 1). In contrast, no evidence of

colonisation of the ALI is evident in SBW25 $\Delta wssA-J$ (Fig. 3b and

supplementary movie file 2). Observations from the camera

perpendicular to the flasks confirmed presence of surface growth in

SBW25 (Fig. 3a), but not in the cellulose mutant (Fig. 3b).

Additionally rapid streaming was observed in the broth phase for the

ancestral genotype but not for SBW25 $\Delta wssA-J$ (Fig. 3a and 3b,

supplementary movie files 3 and 4 respectively). The significance of

this streaming dynamic is considered in detail below.

Curious as to the nature of the previously unseen surface growth we

obtained high-resolution photos at hourly intervals (between 15h and

20 h) from directly above the surface using a light source for

illumination positioned at an oblique angle to the surface. No

surface growth was evident for SBW25 $\Delta wssA-J$ (Fig. 4b), but

remarkably, from the ancestral genotype, numerous micro-colonies

emerged from the surface of the meniscus and grew outward as if on an

agar plate (Fig. 4a). By 19 hours the micro-colonies coalesced into a

single structure (Fig. 4a) that upon coalescence collapsed

(supplementary movie file 5).

Regulation of cellulose and ALI colonisation by multiple diguanylate cyclase-encoding regulatory pathways. Numerous studies of

constitutive cellulose over-producing mutants — the so named wrinkly

spreader (WS) types (28) — have shown the phenotype to arise

primarily by mutations in the Wsp, Aws and Mws pathways (25-30).

Mutations in the negative regulators of these diguanylate cyclase-

encoding pathways (DGCs) result in over-production of cyclic-di-GMP,

over-production of cellulose and formation of substantive and

enduring mats at the ALI. While these findings have connected *over-expression* of DGC-encoding pathways to the WS phenotype, the

relationship between known DGC-encoding pathways and cellulose

expression in the absence of DGC over-activating mutations has been a

mystery. Recognition that ancestral SBW25 activates cellulose

production at the ALI leading to micro-colony formation and a frail

film of cells, allowed investigation of the role of Wsp, Aws and Mws

in expression of this phenotype.

A reduction in the formation of micro-colonies in SBW25 Δ*wspABCDEFR*,

SBW25 Δ*awsXRO*, and SBW25 Δ*mwsR* demonstrates for the first time a

connection between the Wsp, Aws and Mws pathways, the production of

cellulose and colonisation / micro-colony formation at the ALI (Fig.

4c). Surprisingly, no single pathway mutant resulted in a cellulose

defective phenotype that matched that of the cellulose defective

wssA-J deletion mutant (Fig. 4c). Equally surprising was that all

three pathways make some contribution to colonisation of the ALI

(Fig. 4c). The most pronounced phenotype was associated with SBW25

Δ*mwsR*, followed by SBW25 Δ*awsXRO* and SBW25 Δ*wspABCDEFR*. A mutant

lacking all three pathways was indistinguishable from SBW25 Δ*wssA-J*

(Fig. 4c).

Cellulose causes bioconvection. As noted above, in microcosms

inoculated with cellulose-producing ancestral SBW25, material falls

in finger-like plumes that stream from the ALI (Fig. 3a). Analysis of

time-lapse movies (supplementary movie file 6) shows plumes to be

characteristic of long-range convection (Fig. 5), which arises as a

consequence of instability of the interface between the cellulose-

rich meniscus and the less dense broth phase beneath. The phenomenon

is known as Rayleigh-Taylor instability. That cellulose is the

critical component stems from the fact that the streaming plumes are

evident in ancestral SBW25, but not in cultures of the cellulose

negative mutant (SBW25 Δ*wssA-J*).

Quantification of the streaming plumes shows instability at ~25 h and

continues until ~40 h at which point streaming ceases and the medium

becomes homogeneous. The velocity of the falling plumes ranges from

500 to 2000 µm·min^{-1} (Fig. 5). From this it is possible to calculate

the Péclet number that defines the contribution of diffusion relative

to bioconvection on the transport of water-soluble products. In this

instance the Péclet number (*Pe*) is ~1,000 (calculated by multiplying

the typical plume length (1 cm) by its velocity (~$1·10^{-3}$ cm·s^{-1}) and

then dividing by the diffusion coefficient of pyoverdin ~$1·10^{-6}$ cm^2·s^{-}

^{1}). The Péclet number, being greater than 1 (*Pe* is a dimensionless

number), means that bioconvection is a more significant contributor

to the transport of soluble products than diffusion.

Bioconvection affects spatial distribution of extracellular products.

A soluble product of relevance to *P. fluorescens* SBW25 in static

culture is the water-soluble iron binding siderophore, pyoverdin

(31). That it is fluorescent means that it is readily monitored.

Figure 6a shows the average concentration of pyoverdin at the ALI as

imaged via camera 2 equipped with suitable optical filters (see Fig.

1). The first indication of pyoverdin production occurs at the ALI at

~19 h and coincides precisely with the first visible stages of

surface colonisation where micro-colonies begin to form at the

meniscus (Fig. 4).

The first signs of pyoverdin production are restricted to the ALI

despite the fact that at 19 h and thereafter, the broth phase is

turbid with growth (supplementary movie file 3 shows turbidity in the

flask and supplementary movie file 7 shows pyoverdin in the flask).

This is consistent with oxygen being available at the broth surface

(and absent in the bulk phase due to metabolic activity) causing iron

at the ALI to exist in the insoluble ferric form, leading to

activation of pyoverdin synthesis solely at the ALI. The kinetics of

pyoverdin production were quantified by fitting data to a simple

logistic model (Fig. 6a) whose fit indicates that the underlying

chemical reaction is autocatalytic and characteristic of positive

feedback regulation that controls pyoverdin synthesis (31).

Visible plumes of pyoverdin (Fig. 6b) were quantified by measuring

pixel intensity across a single horizontal profile (inset Fig. 6c) as

indicated by the red line in Figure 6b. To determine the

characteristic plume width (Fig. 6c), the data were analysed by Fast

Fourier transformation (FFT). The transformation shows that pyoverdin

is concentrated in plumes with a horizontal width of 3mm.

Modelling microcosm dynamics. Surface colonisation by *P. fluorescens*

SBW25, interaction of cells with oxygen and ensuing effects,

including bioconvection and transport of pyoverdin, draw attention to

striking ecological complexity in this simplest of microcosms. To

determine the match between current understanding of the interplay

between biological, chemical and physical processes and the extent to

which simple biophysical mechanisms explain the observed dynamics, we

constructed a model based on diffusion-reaction processes and

hydrodynamics. The degree of fit between model and data stands to

show how well the system is understood.

The model is based on experimental quantification of bacterial

density, pyoverdin concentration, and fluid flow. It uses partial

differential equations to account for the diffusion-reaction-

convection processes within in the flask. The local concentration of

bacteria, oxygen, pyoverdin and cellulose are described as continuous

fields. The liquid environment is modelled as an incompressible

Newtonian fluid with a mass density that depends on the

concentrations of bacteria and cellulose. Its dynamic is described by

the Navier-Stokes (NS) equation using the Boussinesq approximation,

in which the variations of density are neglected except in the

buoyancy force (32). The coupled equations allow for inclusion of

different physical interactions between the components. Details are

provided in the Materials and Methods section.

The model was solved numerically as a means of validation.

Simulations were performed on a two-dimensional grid representing a

physical domain of size 1 cm^2. The top of the domain corresponds to

the ALI with free fluid slip (liquid can move along the ALI) and no

penetration boundary conditions (the meniscus cannot be deformed).

The sides correspond to the lateral walls of the microcosm and the

bottom of the flask. The boundary conditions on the wall allow no

fluid slip (liquid cannot move along the wall) and no penetration.

The results of the simulation are showing in Figure 7 (and

supplementary movie files 8-11) and closely reproduce the dynamics

observed in microcosms. Bacteria replicate and consume oxygen until

growth saturates at ~$3 \cdot 10^8$ cfu·ml^{-1}. At 16 h oxygen is available at

the meniscus and in a single millimetre layer immediately below the

ALI. Also at 16 h pyoverdin production begins; at 19 h the first

indication of cellulose production become visible resulting in an

increase in density of the surface layer. Soon after, cellulose-

laden regions begin to form descending plumes marking the onset of

Rayleigh-Taylor instability. Plumes flow from the ALI to the bottom

of the flask at a speed of ~1000 µm·min^{-1}. This is in accord with

experimental observations. Additionally, plumes serve to transport

pyoverdin (over a millimetre scales) and oxygen, which penetrates

several millimetres into the liquid phase.

DISCUSSION

The interface between liquid and air defines a niche of significance for many bacteria (4). For aerobic organisms it is an environment replete with oxygen, it offers opportunities for unfettered surface spreading that may aid dispersal and indirectly, may allow rapid colonisation of solid surfaces; colonisation of the ALI may also allow bacteria to escape grazing by solid-surface associated predators. Despite its ecological relevance knowledge of mechanisms and consequences of surface colonisation are poorly understood.

For more than two decades studies of evolution in experimental microcosms have drawn attention to adaptive mutants of *P. fluorescens* SBW25 that specialise in colonisation of the ALI (22,23,28,33). These mutants, which constitutively overproduce cellulose as a consequence of DGC-activating mutations (24,25,26,29,30,34), reap a significant adaptive advantage in static broth microcosms because of ability to grow at the ALI and thus access to oxygen. Largely unknown however has been the ecological significance of cellulose in the ancestral type and more generally, the role of cellulose in the natural environment. Impeding progress has been the fact that cellulose production is not evident on standard agar plate culture and neither is it produced in shaken broth culture. In the absence of a phenotype *in vitro* it is difficult to make progress.

Nonetheless, several previous studies have indicated environmental

relevance: Gal et al (35) showed a cellulose defective mutant to be

significantly less fit than the ancestral type in assays of plant

colonisation and Giddens et al (36) showed the cellulose-encoding *wss*

operon to be specifically activated on plant root surfaces. Perhaps

the most significant finding, but at the time overlooked, was from

competitive fitness assays between ancestral SBW25 and a *wss*

defective mutant performed in shaken and unshaken microcosms (22): in

shaken culture the fitness of the cellulose defective mutant was no

different to the ancestral type, but in unshaken culture the mutant

was significantly less fit. Here, prompted by recent observation of

the poor growth in unshaken culture of SBW25 Δ*wssA-J* (26), combined

with new tools of observation, we have come a step closer to

understanding the biological significance of bacterial cellulose

production.

Apparent from use of the device shown in Figure 1 is that ancestral

SBW25 activates cellulose production in static broth culture and that

polymer production allows cells to break through the meniscus and

remarkably, grow as micro-colonies on the surface. In the absence of

cellulose production, cells are unable to penetrate the ALI and fail

to reap the growth advantage that comes from a plentiful supply of

oxygen (Fig. 2). Just how cellulose enables bacteria to break through

the ALI is unclear. One possibility is that the polymer changes

viscosity and this alone is sufficient to propel bacteria through the

meniscus, another possibility is that the polymer alters surface

charge and that altered electrostatic properties of the cells affects

interactions with the surface.

Also unknown is the signal(s) that lead(s) to activation of cellulose

production. What is clear is that known DGC-encoding pathways are

necessary to transduce effects through to cellulose production. The

fact that three pathways all contribute to differing extents points

to complexity in the mapping between DGCs and the cellulosic target

(37). It is tempting to suggest that the signal is oxygen, but this

seems unlikely because it is incompatible with the previous finding

that SBW25 and a cellulose defective mutant are equally fit in an

oxygen-replete environment (22). Our suspicion is that the signal

stems from some physical attribute of the ALI, possibly surface

tension and Marangoni forces arising as a consequence of evaporation

or production of surfactant – a subject that received momentary

attention almost a century ago (38-40).

The ecological significance of the behaviour is unknown. Assuming

our observations are relevant to the natural environment and not just

to laboratory culture, then one possibility is that cells use

cellulose to colonise the ALI of water films on plant roots / leaves

(the natural environment of SBW25 (41)) and use this environment to

aid rapid and unimpeded dispersal. An additional benefit may then

accrue on drying when the widely dispersed bacteria are bought back

in contact with a solid substrate. Suggestive though that the

remarkable growth extending up and out of the liquid surface may hint

at a more complex and as yet unrecognised behaviour is the

involvement of three DGC-encoding pathways. Why involve three

pathways to regulate cellulose production when one would seem to

suffice?

As colonisation of the surface begins to saturate, the heavier

material on top becomes unstable and collapses in plumes typical of

Rayleigh-Taylor instability. That such behaviour occurs is consistent

with the thesis that cellulose is produced just at the meniscus and

is not evenly distributed throughout the broth phase. Numerous

consequences arise from the ensuing bioconvection, one of which is

the rapid transport of water-soluble products. Our particular

attention has been the fluorescent molecule pyoverdin, which by

virtue of association with cells, is rapidly mixed from the point of

production (the ALI) through the entire broth phase of the microcosm.

Bioconvection additionally alters the chemical status of the

environment, not only through mixing of extant products, but also

through effects wrought by enhanced transport of oxygen.

Transport of pyoverdin has particular significance in light of a

previous analysis of SBW25 populations propagated in static KB

culture of an extended period (21,42). Common mutant types that rose

to prominence harboured mutations that abolished pyoverdin

production. The evolutionary advantage of these mutants stemmed not

from scavenging of pyoverdin (akin to "cheating"), but from avoidance

of the cost of producing pyoverdin when it was not required (21).

That pyoverdin is not required in the broth phase (because lack of

oxygen means iron exists in the soluble ferrous state) is evident

from the time-lapse movies (supplementary movie 7) where pyoverdin

production is initiated exclusively at the ALI. However upon reaching

the point of Rayleigh-Taylor instability, bioconvection due to

cellulose rapidly transports pyoverdin into the broth phase where, in

complex with iron, it serves to positively activate transcription of

pyoverdin synthetic genes (43)- even though pyoverdin is not required

by broth-colonising cells.

Imaging of cultures as reported here draws attention to the

complexity and interdependence of biological, chemical and physical

processes. A primary goal of the modelling exercise was to see just

how far physical descriptions of measured phenomena such as plume

velocity, bacterial density and pyoverdin concentration could account

for observed dynamics. Similar approaches have been taken previously

in analysis of microbial systems (44,45). Specifically, the model

shows how dynamical processes occurring in the liquid can be affected

by biofilm formation at the ALI. It also reveals how proliferation

of biomass affects the production and transport of pyoverdin.

Additionally it accounts for physical transport of water-soluble

products and the relative contributions of diffusion versus

bioconvection to this process.

The model generates results consistent with cellulose production at

the ALI being sufficient to generate Rayleigh-Taylor instability and

initiate fluid movement. The specific mechanisms in the model

involve the imbalance between the force of mass repartition in the

fluid and the damping force of viscosity. The model also supports

hypotheses concerning the critical role of cellulose in

bioconvection: numerical resolution of the model showed plumes to

have a velocity of ~1000 $\mu m.min^{-1}$ as observed in the experiment.

Integrity of plumes – often tens of millimetres in length – is also

explained by the model, and arises from the fact diffusion is a minor

contributor to fluid dynamics relative to the effects of

bioconvection. A further insight concerns ability of bioconvection to

mix oxygen into the top few millimetres of the broth phase at a rate

that is greater than its consumption. All these effects follow from

the Rayleigh-Taylor instability wrought by the production of

cellulose at the meniscus.

Together this study has shed new light on colonisation of the ALI and

particularly the role of cellulose — a widespread microbial product

(46). Previous work has drawn attention to cellulose as an adhesive

substance affecting the relationship between bacteria and solid

surfaces (47-50), but these findings stem from study systems that do

not provide opportunity for ALI colonisation and perhaps by design

even select mutants that over-express cellulose and thus mislead as

to ecological significance. This stated, cellulose may play different

ecological roles in different organisms and under different

conditions. Nonetheless, recognition that production of a polymer

can modify an environment thus significantly changing the

relationship between the organism and its environment – and the

environment in a more general sense – has implications for

understanding environments and processes affected by ALI biofilms,

such as those encountered in sewage treatment plants, marine and

fresh water systems, and in terrestrial environments where transient

films of moisture exist in soil pores and on plant surfaces. It also

raises intriguing possibilities for future research on the importance

of surface tension as a cue eliciting phenotypic responses in

bacteria.

MATERIALS AND METHODS

Bacterial strain and growth conditions. The ancestral strain of *P. fluorescens* SBW25 was isolated from the leaf of a sugar beet plant at the University of Oxford farm (Wytham, Oxford, U.K.; Silby et al. 2009). The Δ*wssA-J* strain is deleted of the entire *wssA-J* operon (PFLU0300-PFLU0309) in the ancestral background and comes from (26). The Δ*wsp*, Δ*aws* and Δ*mws* were previously constructed by a two-step allelic exchange strategy (25).

Strains are cultured in King's Medium B (KB) (51) at 28°C. KB contains (per litre) 20 g bacto™ proteose peptone No. 3 (BD ref211693), 10 g glycerol, 1.5 g K2HPO4 and 1.5 g MgSO4·7H2O.

To follow bacterial dynamics in experimental flasks bacteria were pre-cultured in KB overnight, centrifuged (6000rpm, 4min) and resuspended in fresh KB. The OD of suspended cultures was adjusted to an OD600nm of 0.8 and stored in 20µl aliquots containing 10µL of cultures of OD 0.8 and 10µL of 60w% autoclaved glycerol. The aliquots were conserved at -80°C.

To establish each experiment, a rectangular flask (Easy Flask 25cm² Nunc) was filled with 20ml of KB medium. A 20µl -80°C cultlure was then defrosted and inoculated in the KB at a final dilution of 10^4 cfu.ml^{-1}. The flask was positioned in the setup shown in Figure 1 and incubated at 28°C in an IGS60 HERATHERM static incubator.

Experimental setup to measure the dynamics of unshaken bacterial culture. The setup was designed and built to perform custom measurements and details are available from the authors upon request. The device comprises a laser-photodiode alignment to measure optical density in the flask and three cameras to observe the ALI as well as the biomass and the pyoverdin in the liquid phase.

To measure the optical density of the liquid phase a vertical profile was obtained by scanning with a laser-photodiode detector mounted on a lifter. A plastic piece that was produced by a 3D-printer on a stratasys fortus250 in ABS (yellow on Fig. 1) joined the laser-photodiode to a carriage that was free to slide on a vertical rail (ingus TS-01-15/TW-01-15) driven by a M10 cage bolt coupled to a M10 threaded rod. This ensured a precise vertical and horizontal positioning of the laser-photodiode alignment. The thread rode was smoothly rotated using a 7,2Vcc motor. A L293D power switch controlled by an Arduino board MEGA 2560 directed rotation of the motor. The thread rod rotation angle was measured with an optical encoder HEDS5500 500CPR. Ultimately, this allowed measurement of the vertical position of the laser beam with a resolution of ~3µm.

The photodiode was from Thorlabs (FDS1010), the laser a HLM1230 of wavelength 650 nm and power 5mW. To ensure that the laser did not harm bacteria the light was attenuated by a NE520B (Thorlabs) neutral

density filter of OD=2. The optical density of the culture was

evaluated by measuring the photo-current produced by the laser

hitting the photodiode after it went through the flask. To ensure

linearity between the intensity of light hitting the photodiode and

its conversion in photo-current, the photodiode was polarised in

inverse with 5V provided by a LM4040 electronic component. The photo-

current was estimated by measuring the voltage of a 437 ± 5% kΩ

resistor mounted in serial with the photodiode. The voltage was

monitored by the Arduino MEGA 2560 board encoding a 0-5V analogic

input on 10 bits giving a resolution of 5mV. After subsequent

calibration, the signal acquired by the system allowed estimation of

the bacterial concentration in the flask within a range of 10^7-$5\cdot10^9$

bact$\cdot$ml^{-1}.

Synchronized with the laser-photodiode, were three cameras: a uEyeLE

USB2.0 Camera, a 1/2" CMOS Monochrome Sensor, and a 1280x1024 Pixel

equipped with a CMFA0420ND 4 mm ½ inch lens. The first camera (Fig.

1) records a bright field image of the vertical side view of the

culture medium. The second camera is equipped with a band pass

optical filter 470 ± 10nm (Thorlabs FB470-10). It takes a side view

of the culture medium. During its acquisition a 405nm laser (405MD-5-

10-1235) illuminates the flask to excite pyoverdin fluorescence. The

third camera takes a bright field image of the ALI with an angle of

~45°. Acquisition of optical density data and photos were

synchronized using a master script written in Python that recorded

the data produced by the Arduino board and saved the photos taken by

the cameras.

Colonization of the air-liquid interface ALI (Fig. 4). To observe the

effect of Δ*wsp*, Δ*aws* and Δ*mws* mutations on ALI colonization, we use

6-well plates Greiner bio-one 657160 filled with 8ml of KB. Each well

was inoculated from glycerol stocks. The 6-well plates ere incubated

at 28°C without shaking. Pictures were taken the ALI with a Nikon

D7000 camera equipped with an AF-S DX NIKKOR 18-105mm f/3.5-5.6G ED

VR objective.

The Model. The model uses six fields to describe the system: the

vector field of the fluid vorticity (ω), the scalar field of the

fluid stream function (Ψ) and the scalar fields of bacterial (b),

oxygen (o), cellulose (c) and pyoverdin (p) concentrations. We also

use a derived vector field that represents the velocity of the fluid

(u). The model is valid for a three dimensional space but we estimate

its validity in a two dimensional space in order to reduce the time

of numerical computation. That is why we choose a fluid description

in term of vorticity (ω) and the stream function (Ψ). This

description gives two advantages for the numerical resolution of the

model. First, the equation of the fluid incompressibility is solved

by construction; second, the calculation of the vorticity vector can

be reduced to the calculation of a simple scalar field (for more

detail see (35)).

With the six fields given above come six partial differential

equations that describe their dynamics.

The first equations related to the stream function. This scalar field

is calculated by a Poisson equation:

Equation 1

$$\Delta\Psi = -\omega$$

611 Where, Δ is the Laplace operator.

612 The second equation deals with the vorticity reduced to a simple

613 scalar field. Its dynamics can be derived from the Navier-Stokes (NS)

614 equation:

615 **Equation 2**

$$\frac{\partial\omega}{\partial t} + \left(\vec{u}\bullet\vec{V}\right)\omega = v\Delta\omega - g\frac{\partial}{\partial x}\left(\frac{\rho}{\rho_0}\right)$$

The left side of NS is the Lagrangian derivative of the vorticity.

The right side contains damping of the vorticity by the viscosity v,

and a gravity term traduces the generation of vorticity due to the

uneven spatial repartition of the mass density ρ relative to the

density of the fluid medium ρ_0. The operator $\left(\vec{u}\bullet\vec{V}\right)$ stem for the

convective derivative. In the equation, the local mass density ρ

takes into account the mass density of the liquid medium ρ_0, bacteria

ρ_b and cellulose ρ_c. Hence we consider the local mass density (ρ) as

the sum of the mass contribution of the liquid medium, the bacteria

and the cellulose. Explicitly the notation ρ *stem for*:

$$\rho = \rho_0 + \Phi_b(\rho_b - \rho_0) + \Phi_c(\rho_c - \rho_0)$$

where Φ_b is the local volume fraction of the bacteria and Φ_b the

local volume fraction of the cellulose.

To calculate the dynamics of the concentration of bacteria (*b*),

cellulose (*c*) and pyoverdin (*p*), we write a diffusion-reaction-

convection equation.

To write the third equation dealing with bacteria we make several

assumptions:

Equation 3

$$\frac{\partial b}{\partial t} + (\vec{u} \bullet \vec{\nabla})b = D_b \Delta b + \delta b \left(1 - \frac{b}{b_{sat}}\right)$$

bacteria grow exponentially until they reach the saturation b_{sat}

measured experimentally (Fig. 2) and, bacteria consume oxygen that is

dissolved into the liquid.

The left-hand side of the bacterial equation is the Lagrangian

derivative applied to b. The right-hand side contains a diffusive

term that takes into account the random motility of bacteria with a

diffusion coefficient D_b, and an exponential growth term with a rate

δ that goes to zero when the concentration reach the b_{sat} value.

The fourth equation describes the dynamics of the oxygen (o) field.

Bacteria consume the oxygen at a rate γ.

Equation 4

$$\frac{\partial o}{\partial t} + \left(\vec{u} \bullet \vec{\nabla}\right)o = D_o \Delta o - \gamma b\, \Theta(o)$$

The coefficient of diffusion is D_o. Oxygen consumption goes to zero

when there is no more oxygen. This is ensure that multiplying the

consumption term by the Heaviside function ($\Theta(o)$) is 1 when o is

above zero, but zero otherwise.

The fifth equation assumes that the cellulose is produced with an

exponential rate (α) as long as the oxygen concentration is higher

than o^* and the concentration of bacteria is higher than $b^\star$.

Additionally, cellulose production saturates when c tends to 1, its

maximal value. The equation is:

Equation 5

$$\frac{\partial c}{\partial t} + \left(\vec{u} \bullet \vec{\nabla}\right)c = D_c \Delta c + \alpha c(1-c)\Theta(b-b^*)\,\Theta(o-o^*)$$

661   The sixth equation describes the dynamics of pyoverdin production.

662   Provided that the local concentration of oxygen is sufficiently high

663   bacteria produce pyoverdin according to the autocatalytic synthesis

664   measured experimentally (Fig. 5D) with a rate β. The equation is:

665   **Equation 6**

$$\frac{\partial p}{\partial t} + \left(\vec{u} \bullet \vec{\nabla}\right)p = D_p \Delta p + \beta p(1-p)\,\Theta(o-o^*)$$

Here, pyoverdin production goes to zero when oxygen concentration is

below $o*$ by multiplying the production term by a Heaviside function

$\Theta(o-o*)$.

Finally, to calculate the fluid velocity we use the derivative of the

stream function where u_x and u_y stand for the horizontal and vertical

components of the fluid velocity (u):

$$u_x = \frac{\partial \Psi}{\partial y}$$

$$u_y = -\frac{\partial \Psi}{\partial x}$$

Numerical Simulations. We used a finite difference method to solve

the coupled reaction-diffusion-convection equations (35). The

simulation was performed on a Linux system: Debian 4.9.51-1, gcc

6.3.0. The hardware CPU was an Intel(R)Core(TM) i7-7700K @ 4.2Ghz

with 16GB RAM. The parameters used in the simulation displayed Figure

7 are listed in Table 1.

ACKNOWLEDGEMENTS

We thank Nicolas Desprat, Clara Moreno Fenoll, Steven Quistad and

Guilhem Doulcier for discussion and comment. MA was supported by HFSP

grant RGP0010/2015.

REFERENCES

1. Costerton JW, Stewart PS, Greenberg EP. 1999. Bacterial

biofilms: a common cause of persistent infections. Science

284:1318-22.

2. Davey ME, O'Toole G A. 2000. Microbial biofilms: from ecology

to molecular genetics. Microbiol Mol Biol Rev 64:847-67

3. Flemming HC, Wingender J, Szewzyk U, Steinberg P, Rice SA,

Kjelleberg S. 2016. Biofilms: an emergent form of bacterial

life. Nat Rev Microbiol 14:563-75.

4. Marshall, K. C. 1996. *Adhesion as a strategy for access to*

nutrients (pp. 59-87). Wiley: New York.

5. Watnick P, Kolter R. 2000. Biofilm, city of microbes. J

Bacteriol 182:2675-9.

6. Donlan RM. 2002. Biofilms: microbial life on surfaces.

Emerging infectious diseases 8:881-890.

7. Alsharif G, Ahmad S, Islam MS, Shah R, Busby SJ, Krachler AM.

2015. Host attachment and fluid shear are integrated into a

mechanical signal regulating virulence in *Escherichia coli*

O157: H7. Proc Natl Acad Sci U.S.A. 102:5503-5508.

8. Stewart PS. 2002. Mechanisms of antibiotic resistance in

bacterial biofilms. Int J Med Microbiol 292:107-13.

9. Dang H, Lovell CR. 2016. Microbial surface colonization and

biofilm development in marine environments. Microbiol Mol

Biol Rev 80:91-138.

10.Hamilton WD, May RM. 1977. Dispersal in stable habitats.

711   Nature 269:578-581.

712 11. Monds RD, O'Toole GA. 2009. The developmental model of

713   microbial biofilms: ten years of a paradigm up for review.

714   Trends Microbiol 17:73-87.

715 12. O'Toole GA, Wong GC. 2016. Sensational biofilms: surface

716   sensing in bacteria. Curr Opin Microbiol 30:139-146.

717 13. Heydorn A, Nielsen AT, Hentzer M, Sternberg C, Givskov M,

718   Ersboll BK, Molin S. 2000. Quantification of biofilm

719   structures by the novel computer program COMSTAT. Microbiol

720   146:2395-407.

721 14. Azeredo J, Azevedo NF, Briandet R, Cerca N, Coenye T, Costa

722   AR, Desvaux M, Di Bonaventura G, Hébraud M, Jaglic Z. 2017.

723   Critical review on biofilm methods. Crit Rev Microbiol

724   43:313-351.

725 15. Petrova OE, Sauer K. 2012. Sticky situations: key components

726   that control bacterial surface attachment. J Bacteriol

727   194:2413-2425.

728 16. Valentini M, Filloux A. 2016. Biofilms and cyclic di-GMP (c-

729   di-GMP) signaling: Lessons from *Pseudomonas aeruginosa* and

730   other bacteria. J Biol Chem 291:12547-55.

731 17. Kjelleberg, S. (1985). Mechanisms of bacterial adhesion at

732   gas-liquid interfaces. In *Bacterial Adhesion* (pp. 163-194).

733   Springer, Boston, MA.

734 18. Wotton RS, Preston TM. 2005. Surface films: Areas of water

735   bodies that are often overlooked. AIBS Bull. 55:137-145.

19.Neilands JB. 1981. Microbial iron compounds. Annu Rev

Biochem 50:715-31.

20.Koza A, Moshynets O, Otten W, Spiers AJ. 2011. Environmental

modification and niche construction: developing O2 gradients

drive the evolution of the Wrinkly Spreader. ISME J 5:665-73.

21.Zhang XX, Rainey PB. 2013. Exploring the sociobiology of

pyoverdin-producing *Pseudomonas*. Evolution 67:3161-74.

22.Spiers AJ, Kahn SG, Bohannon J, Travisano M, Rainey PB.

2002. Adaptive divergence in experimental populations of

Pseudomonas fluorescens. I. Genetic and phenotypic bases of

wrinkly spreader fitness. Genetics 161:33-46.

23.Rainey PB, Rainey K. 2003. Evolution of cooperation and

conflict in experimental bacterial populations. Nature

425:72.

24.Bantinaki E, Kassen R, Knight C, Robinson Z, Spiers A,

Rainey P. 2007. Adaptive divergence in experimental

populations of Pseudomonas fluorescens. III. Mutational

origins of wrinkly spreader diversity. Genetics 176:441-453.

25.McDonald MJ, Gehrig SM, Meintjes PL, Zhang X-X, Rainey PB.

2009. Adaptive divergence in experimental populations of

Pseudomonas fluorescens. IV. Genetic constraints guide

evolutionary trajectories in a parallel adaptive radiation.

Genetics. 161: 33-46.

26.Lind PA, Farr AD, Rainey PB. 2017. Evolutionary convergence

in experimental *Pseudomonas* populations. ISME J 11:589.

27. Plesset MS, Whipple CG, Winet H. 1976. Rayleigh-Taylor

instability of surface layers as the mechanism for

bioconvection in cell cultures. J Theor Biol 59:331-351.

28. Rainey PB, Travisano M. 1998. Adaptive radiation in a

heterogeneous environment. Nature 394:69-72.

29. Lind PA, Farr AD, Rainey PB. 2015. Experimental evolution

reveals hidden diversity in evolutionary pathways. eLife

4:e07074.

30. Lind PA, Libby E, Herzog J, Rainey PB. 2019. Predicting

mutational routes to new adaptive phenotypes. eLife 8:e38822.

31. Visca P, Imperi F, Lamont IL. 2007. Pyoverdine siderophores:

from biogenesis to biosignificance. Trends Microbiol 15:22-

30.

32. Pozrikidis C. 2016. *Fluid Dynamics: Theory,Ccomputation, And*

Numerical Simulation. Springer.

33. Spiers AJ, Bohannon J, Gehrig SM, Rainey PB. 2003. Biofilm

formation at the air-liquid interface by the *Pseudomonas*

fluorescens SBW25 wrinkly spreader requires an acetylated

form of cellulose. Mol Microbiol 50:15-27.

34. Goymer P, Kahn S, Malone J, Gehrig S, Spiers A, Rainey P.

2006. Adaptive divergence in experimental populations of

Pseudomonas fluorescens. II. The role of the GGDEF regulator

WspR in evolution and development of the wrinkly spreader

phenotype. Genetics. 161:33-46.

35. Gal M, Preston GM, Massey RC, Spiers AJ, Rainey PB. 2003.

Genes encoding a cellulosic polymer contribute toward the

ecological success of *Pseudomonas fluorescens* SBW25 on plant

surfaces. Mol Ecol 12:3109-3121.

36.Giddens SR, Jackson RW, Moon CD, Jacobs MA, Zhang X-X,

Gehrig SM, Rainey PB. 2007. Mutational activation of niche-

specific genes provides insight into regulatory networks and

bacterial function in a complex environment. Proc Natl Acad

Sci U.S.A. 104(46):18247-18252.

37.Yan J, Deforet M, Boyle KE, Rahman R, Liang R, Okegbe C,

Dietrich LE, Qiu W, Xavier JB. 2017. Bow-tie signaling in c-

di-GMP: Machine learning in a simple biochemical network.

PLoS Comp Biol 13:e1005677.

38.Frobisher Jr M. 1926. Relations of surface tension to

bacterial phenomena. J Infect Dis 38:66-91.

39.Gibbs WM, Batchelor H, Sickels T. 1926. Surface tension and

bacterial growth. J Bacteriol 11:393.

40.Pizarro OR. 1927. The relation of surface tension to

bacterial development. J Bacteriol 13:387.

41.Rainey PB, Bailey MJ. 1996. Physical and genetic map of the

Pseudomonas fluorescens SBW25 chromosome. Mol Microbiol

19:521-533.

42.Rainey PB, Desprat N, Driscoll WW, Zhang XX. 2014. Microbes

are not bound by sociobiology: Response to Kümmerli and Ross-

Gillespie (2013). Evolution 68:3344-3355.

43.Lamont IL, Beare PA, Ochsner U, Vasil AI, Vasil ML. 2002.

Siderophore-mediated signaling regulates virulence factor

production in *Pseudomonas aeruginosa*. Proc Natl Acad Sci

U.S.A. 99:7072-7077.

44. Ardré M, Henry H, Douarche C, Plapp M. 2015. An individual-

based model for biofilm formation at liquid surfaces. Phys

Biol 12:066015.

45. Tuval I, Cisneros L, Dombrowski C, Wolgemuth CW, Kessler JO,

Goldstein RE. 2005. Bacterial swimming and oxygen transport

near contact lines. Proc Natl Acad Sci U.S.A. 102:2277-2282.

46. Ross P, Mayer R, Benziman M. 1991. Cellulose biosynthesis

and function in bacteria. Microbiol Mol Biol Rev 55:35-58.

47. Matthysse A. 1983. Role of bacterial cellulose fibrils in

Agrobacterium tumefaciens infection. J Bacteriol 154:906-915.

48. Da Re S, Ghigo J-M. 2006. A CsgD-independent pathway for

cellulose production and biofilm formation in *Escherichia*

coli. J Bacteriol 188:3073-3087.

49. Serra DO, Richter AM, Hengge R. 2013. Cellulose as an

architectural element in spatially structured *Escherichia*

coli biofilms. J Bacteriol 195:5540-5554.

50. Zogaj X, Nimtz M, Rohde M, Bokranz W, Römling U. 2001. The

multicellular morphotypes of *Salmonella typhimurium* and

Escherichia coli produce cellulose as the second component of

the extracellular matrix. Mol Microbiol 39:1452-1463.

51. King EO, Ward MK, Raney DE. 1954. Two simple media for the

demonstration of pyocyanin and fluorescin. Transl Res 44:301-

307.

52.Flemming HC, Wingender J. 2010. The biofilm matrix. Nat Rev

Microbiol 8:623.

53.Bratbak G, Dundas I. 1984. Bacterial dry matter content and

biomass estimations. App Env Microbiol 48:755-757.

54.Berg HC. 1993. *Random Walks in Biology*. Princeton University

Press.

55.Krieger IM, Mulholland GW, Dickey CS. 1967. Diffusion

coefficients for gases in liquids from the rates of solution

of small gas bubbles. J Phys Chem 71:1123-1129.

56.Martí S, Rodríguez-Baño J, Catel-Ferreira M, Jouenne T, Vila

J, Seifert H, Dé E. 2011. Biofilm formation at the solid-

liquid and air-liquid interfaces by *Acinetobacter* species.

BMC Res Notes 4:5.

FIGURES

FIG 1 Experimental device. A polycarbonate cell-culture bottle

filled with 20 ml of King's Medium B and inoculated with bacteria is

placed on a fixed vertical stand. The device and associated cameras

are maintained within a 28°C incubator. The flask is scanned

vertically every 5min with a 600 nm laser beam with 1mm section.

Light passing through the flask is collected by a photodiode. To

obtain a measure of the optical density in the flask along a vertical

profile, the alignment laser-photodiode is coupled to a motorised

device that ensures smooth vertical translation. Three cameras are

located around the flask. The first (camera 1) obtains a side-view

image of the liquid phase of the medium using bright-field

illumination. The second (camera 2), also fixed perpendicular to the

flask, monitors fluorescence associated with pyoverdin (excitation

405/emission 450nm). The third camera (camera 3) is oriented with a

45° angle and captures growth at the ALI using bright-field

illumination.

FIG 2 Production of cellulose maximises growth in static broth culture. Dynamics of growth of *P. fluorescens* SBW25 (black lines) and *P. fluorescens* SBW25 Δ*wssA-J* (cellulose negative mutant) (grey lines) in unshaken King's Medium B as determined by the scanning laser device and associated photodiode depicted in Figure 1. Data are average optical density at 600 nm (OD_{600}) obtained from scanning the vertical section of three independent flasks at 5 min intervals. OD_{600} measures are calibrated using direct plate counts of colony forming units (equivalent cfu·ml^{-1}). The arrow denotes the onset of bioconvection caused by production of cellulose that marks a secondary increase in growth. This second growth phase is absent in the cellulose negative mutant.

FIG 3 Cellulose is necessary for growth at the air-liquid interface

(ALI) and results in bioconvection. Bright-field images of ancestral

P. fluorescens SBW25 (a) and *P. fluorescens* SBW25 Δ*wssA-J* (cellulose

negative mutant) (b) taken at four time intervals. Complete movies

are available as SI movie files 1,2,6 and 7. Images above the solid

line show growth at the ALI captured using camera 3; images below the

line are from camera 1 (see Figure 1). At time 0 h the medium is

inoculated with ~10^4 cells·ml^{-1}. By 19h the ancestral cellulose-

producing genotype has formed a thin white pellicle at the ALI

(visible by both camera 1 and 3). No pellicle formation is seen in

the cellulose negative mutant, but growth is evident in the broth

phase. By 26 h, in cultures of the cellulose-producing ancestral

type, plumes characteristic of bioconvection stream from the ALI. No

evidence of mat formation or streaming is seen in SBW25 Δ*wssA-J*. By

40h streaming has largely ceased in the ancestral type, although

growth is still apparent at the ALI. Scale bars are 5mm. Contrast

has been adjusted to highlight salient features.

FIG 4 Multiple diguanylate cyclase are required for colonisation of

the ALI. Micro-colony formation at the ALI for ancestral *P.*

fluorescens SBW25 and a range of mutants captured from a camera

mounted directly above individual wells of a six-well tissue culture

plate containing 5 ml King's Medium B. Time course of micro-colony

formation for ancestral *P. fluorescens* SBW25 (a). Comparison with

SBW25 Δ*wssA-J* (cellulose negative mutant) at 19 h (b). Patterns of

micro-colony formation at 19 h in mutants devoid of Wsp (Δ*wsp*), Aws

(Δ*aws*) and Mws (Δ*mws*) diguanylate cyclase-encoding pathways and

combinations there-of. Scale bar is 1 mm, except for the entire well

in (a) which is 5 mm.

FIG 5 Bioconvection caused by cellulose. Time lapse images via bright

field camera 1 (Fig. 1) capture biomass dynamics in the liquid

medium. By 25 h Rayleigh-Taylor instability generates plumes of

biomass that fall from the ALI to the bottom of the flask (inset).

The velocity of movement is obtained by tracking trajectories of the

plumes. The frequency distribution of plume velocity reveals a mean

speed of 983 ± (SD) 373 μm·min^{-1}.

a

p=K/[a·exp(-β· t)+K]
K =5.86e-02
a =6.34e+03
β=4.77e-01
K =5.74e-01
a =1.21e+04
β=3.99e-01

b

c

FIG 6 Bioconvection. Camera 2 (see Figure 1) monitors pyoverdin
concentration in the flask by measuring fluorescence. Pyoverdin is
produced primarily at the ALI. The fluorescence at the position of
the ALI increases with time like a sigmoid curve (a). The adhoc
logistic function of the inset gives the normalized intensity (*p*) as
a function of the time (*t*) and the parameters of the fit (*K*, *a* and

β). The fitted curves (dotted line) adjust the experimental curves
(plain line) for the estimated values of the parameters given in
inset. Plumes due to Rayleigh-Taylor instability transport pyoverdin
from the ALI to the liquid phase. Pyoverdin concentration is
transiently higher along vertical columns that correspond to the
plumes flowing from the ALI. The white scale bar is 5mm. The
fluorescence intensity profile along the red horizontal line (b)
shows that pyoverdin is distributed with a fluctuating spatial
structure (inset). Fast Fourier transformation (FFT) of the intensity
profile reveals these fluctuations to have a characteristic
wavelength of 3mm.
949

FIG 7 Numerical simulation of the mathematical model. Images display

the dynamics of the simulated microcosm from inoculation at 0h to

28h. Time resolved movies are available in supplementary movies 8-11.

The first row above shows the dynamics of the biomass in the bulk

(bacteria and cellulose) as if observed with bright field

illumination 1 (Fig. 1). In experiments, at 24h, plumes concentrated

in biomass flow are evident in the liquid phase. The second row shows

the concentration of pyoverdin in the liquid phase. The plumes

transport pyoverdin into the bulk phase. The third row shows the

dynamics of liquid velocity. When bioconvection is activated fluid

flow is of the order of 1000 μm·min^{-1}, which is consistent with the

measurements shown (Fig. 5). The fourth row shows the dynamics of

oxygen concentration. Soon after inoculation oxygen in the bulk phase

is eliminated due to metabolic (oxygen consuming) activities of

bacteria. The supply of oxygen at the ALI combined with growth of

bacteria and production of cellulose means a gradient of oxygen 2-3

mm into the liquid. Images at 24h and 28h show that oxygen transport

from the ALI before consumption by bacteria in the liquid phase.

The square images are 1 cm^2 and contrast is identical across each

row.

 TABLE 1. Parameters, values and sources of data used for the model.

Name	Symbol	Value	Unit	Source
time step	Δt	$1 \cdot 10^{-3}$	s	
spatial step	Δx	$1 \cdot 10^{-4}$	m	
grid size NxN	N	100	pixel	
fluid dynamics viscosity	η	$8.9 \cdot 10^{-4}$	$kg \cdot m \cdot s$	handbook
bacterial volume	V_b	$3 \cdot 10^{-18}$	m^{-3}	experimental evidence
water mass density	ρ_0	$0.995 \cdot 10^3$	$kg \cdot m^{-3}$	(52)
bacterial mass density	ρ_b	$1.193 \cdot 10^3$	$kg \cdot m^{-3}$	(56)
cellulose mass density	ρ_c	$1.5 \cdot 10^3$	$kg \cdot m^{-3}$	
maximal bacterial concentration	b_{sat}	$3 \cdot 10^{14}$	bacteria $\cdot m^{-3}$	experimental fit
initial concentration of bacteria	b_0	$2 \cdot 10^{10}$	bacteria $\cdot m^{-3}$	experimental settings
diffusion coefficient of bacteria	D_b	$1 \cdot 10^{-10}$	$m^2 \cdot s^{-1}$	(57)
diffusion coefficient of cellulose	D_c	$5 \cdot 10^{-11}$	$m^2 \cdot s^{-1}$	adhoc
diffusion coefficient of	D_o	$1 \cdot 10^{-9}$	$m^2 \cdot s^{-1}$	(58)

Name	Symbol	Value	Unit	Source
oxygen				
diffusion coefficient of pyoverdin	D_p	$3 \cdot 10^{-10}$	$m^2 \cdot s^{-1}$	measured
initial oxygen concentration	O_0	$1.5 \cdot 10^{23}$	molecules $\cdot m^{-3}$	(45)
initial pyoverdin concentration	p_0	$3.36 \cdot 10^{-5}$	au	experimental fit
initial cellulose concentration	c_0	$1 \cdot 10^{-8}$	au	adhoc
production of pyoverdin	β	$1.22 \cdot 10^{-4}$	s	experimental fit
production of cellulose	α	$1 \cdot 10^{-3}$	s	adhoc
growth rate bacteria	δ	0.53	h^{-1}	experimental fit
oxygen consumption per bacteria per second	γ	$1 \cdot 10^6$	molecules $\cdot bacteria^{-1} \cdot s^{-1}$	(45)
volume of cellulose produced per bacteria	$10 \cdot V_b$	$3 \cdot 10^{-17}$	m^3	(55)

Name	Symbol	Value	Unit	Source
minimal bacterial concentration for cellulose production	$b^* = b_{sat}/3$	$1 \cdot 10^{14}$	bacteria $\cdot m^{-3}$	experimental evidence
Boundary condition of oxygen at the top	O_0	$1.5 \cdot 10^{23}$	molecules $\cdot m^{-3}$	(45)
acceleration of gravity	g	9.81	$m \cdot s^{-2}$	
minimal oxygen concentration for cellulose and pyoverdin production	o^*	$O_0 \cdot 10^{-1}$	molecules $\cdot m^{-3}$	adhoc

SUPPLEMENTARY FILES

Supplementary movie file 1.

Name of file: supplementaryMovie1_ancestor_Surface.avi

Growth of ancestral SBW25 imaged from camera 2.

Supplementary movie file 2.

Name of file: supplementaryMovie2_Wss_Surface.avi

Growth of SBW25 Δ*wssA-J* imaged from camera 2.

Supplementary movie file 3.

Name of file: supplementaryMovie3_ancestor_Turbidity.avi

Growth of ancestral SBW25 imaged from camera 1.

Supplementary movie file 4.

Name of file: supplementaryMovie4_Wss_Turbidity.avi

Growth of Δ*wssA-J* SBW25 imaged from camera 1.

Supplementary movie file 5.

Name of file: supplementaryMovie5_ancestor_ALI.avi

Growth of ancestral SBW25 imaged by SLR camera mounted directly above

the well.

Supplementary movie file 6.

Name of file: supplementaryMovie6_ancestor_Bioconvection.avi

1000   Replicate of ancestral SBW25 growth imaged by camera 1 with a high

1001   frequency frame rate.

1002

1003   Supplementary movie file 7.

1004   Name of file: supplementaryMovie7_ancestor_Pvd.avi

1005   Growth of ancestral SBW25 imaged from camera 3.

1006

1007   Supplementary movie file 8.

1008   Name of file: supplementaryMovie8_pvd.gif

1009   Simulation of the pyoverdin dynamics.

1010

1011   Supplementary movie file 9.

1012   Name of file: supplementaryMovie9_velocity.gif

1013   Simulation of the velocity dynamics.

1014

1015   Supplementary movie file 10.

1016   Name of file: supplementaryMovie10_biomass.gif

1017   Simulation of the biomass dynamics.

1018

1019

1020   Supplementary movie file 11.

1021   Name of file: supplementaryMovie10_O2.gif

1022   Simulation of the oxygen dynamics.

1023